1 **Genomic insights into adaptive divergence and speciation among malaria**
2 **vectors of the *Anopheles nili* group**
3

Caroline Fouet[1]*, Colince Kamdem[1], Stephanie Gamez[1], Bradley J. White[1,2]*

[1]Department of Entomology, University of California, Riverside, CA 92521

[2]Center for Disease Vector Research, Institute for Integrative Genome Biology, University of

California, Riverside, CA 92521

*Corresponding authors: caroline.fouet@ucr.edu; bwhite@ucr.edu

Abstract

Ongoing speciation in most African malaria vectors gives rise to cryptic populations, which

differ remarkably in their behaviour, ecology and capacity to vector malaria parasites.

Understanding the population structure and the drivers of genetic differentiation among

mosquitoes is critical for effective disease control because heterogeneity within species

contribute to variability in malaria cases and allow fractions of vector populations to escape

control efforts. To examine the population structure and the potential impacts of recent

large-scale control interventions, we have investigated the genomic patterns of

differentiation in mosquitoes belonging to the *Anopheles nili* group, a large taxonomic group

that diverged ~3-Myr ago. Using 4343 single nucleotide polymorphisms (SNPs), we

detected strong population structure characterized by high F_{ST} values between multiple

divergent populations adapted to different habitats within the Central African rainforest.

Delineating the cryptic species within the *Anopheles nili* group is challenging due to

incongruence between morphology, ribosomal DNA and SNP markers consistent with

incomplete lineage sorting and/or interspecific gene flow. A very high proportion of loci are

fixed (F_{ST} = 1) within the genome of putative species, which suggests that ecological and/or

reproductive barriers are maintained by strong selection on a substantial number of genes.

Key words: *Anopheles nili*, divergent selection, high-F_{ST} regions, speciation.

1. Introduction

One of the principal goals of population genetics is to summarize the genetic similarities and differences between populations (Wright, 1984). This task can be relatively straightforward for some taxa, but the genetic relationship among populations can also be difficult to summarize, especially for species whose evolutionary history is complex and reticulate. The best known mosquito species of the genus *Anopheles* — which includes all vectors of *Plasmodium*, the parasite of human malaria — exhibit very complex range-wide population structure due to the combined effects of cryptic speciation, adaptive flexibility and ongoing gene flow (Harbach, 2013; Krzywinski and Besansky, 2003). For example, almost all major malaria vectors of the Afrotropical region belong to large taxonomic groups encompassing multiple incipient species relatively isolated reproductively and geographically from one another (reviewed by Sinka et al. 2010; Antonio-Nkondjio and Simard 2013; Coetzee and Koekemoer 2013; Dia et al. 2013; Lanzaro and Lee 2013). These characteristics make them promising model systems to study speciation and the processes which contribute to reproductive barriers (e.g. Turner et al. 2005; Lawniczak et al. 2010; Neafsey et al. 2010; Fontaine et al. 2015; Weng et al. 2016), but can also have far-reaching practical consequences. Both spatial and temporal variability in malaria cases and the effectiveness of vector control measures are greatly impacted by heterogeneity within vector species (Molineaux and Gramiccia 1980; Van Bortel et al. 2001). For these reasons, research on the genetic structure among the major African malaria vector mosquitoes has intensified over the last few decades (Antonio-Nkondjio and Simard, 2013; Coetzee and Koekemoer, 2013; Dia et al., 2013; Lanzaro and Lee, 2013).

The recent scaling up of insecticide-treated nets usage and indoor insecticide

spraying to a lesser extent have led to a dramatic reduction in malaria morbidity across the

continent (WHO, 2016). However, other implications of these large-scale interventions

include increased insecticide resistance (reviewed by Hemingway et al. 2016; Ranson and

Lissenden 2016), range shift (e.g. Bøgh et al. 1998; Derua et al. 2012; Mwangangi et al.

2013) and profound evolutionary changes among vector populations. In contrast to

insecticide resistance and range shift, which has been extensively studied, the recent

adaptive changes among mosquito populations have just started to be addressed. These

changes — which involve local adaptation and genetic differentiation, introgressive

hybridization and selective sweeps across loci conferring resistance to xenobiotics — are

particularly evident in the most anthropophilic species (Barnes et al., 2017; Clarkson et al.,

2014; Kamdem et al. , 2017; Norris et al., 2015).

The ecology, taxonomic complexity, geographic distribution, role in transmission

and evolutionary potential of each major African malaria vector is unique. Consequently,

further research is needed to specifically resolve the fine-scale population structure and the

genomic targets of natural selection in all of the important taxa including currently

understudied species. The present work focused on a group of malaria vector species

representing a large taxonomic unit named *Anopheles nili* group. Despite the significant role

of some of its species in sustaining high malaria transmission, this group has received little

attention. To date, four species that occur in forested areas of Central and West Africa and

are distinguishable by slight morphological variations are known within the *An. nili* group:

An. nili sensu stricto (hereafter *An. nili*), *An. ovengensis*, *An. carnevalei*, and *An. somalicus*

(Awono-Ambene et al., 2004; Gillies and Coetzee, 1987; Gillies and De Meillon, 1968). These

species are characterized by reticulate evolution and complex phylogenies that have been

challenging to resolve so far (Awono-Ambene et al., 2004; Awono-Ambene et al., 2006;

Kengne et al., 2003; Ndo et al., 2010, 2013; Peery et al., 2011; Sharakhova et al., 2013).

Populations of *An. nili* and *An. ovengensis* are very anthropophilic and efficient vectors of

Plasmodium in rural areas where malaria prevalence is particularly high (Antonio-Nkondjio

et al., 2006).

To delineate genomic patterns of differentiation, we sampled mosquito populations

throughout the range of species of the *An. nili* group in Cameroon and used reduced

representation sequencing to develop genome-wide SNP markers that we genotyped in 145

individuals. We discovered previously unknown subpopulations characterized by high

pairwise differentiation within *An. ovengensis* and *An. nili*. We further explored the genetic

differentiation across the genome and revealed the presence of a very high number of

outlier loci that are targets of selection among locally adapted subpopulations. These

findings provide significant baseline data on the genetic underpinnings of adaptive

divergence and pave the way for further genomic studies in this important group of

mosquitoes. Notably, a complete reference genome will allow in-depth studies to decipher

the functional and phenotypic characteristics of the numerous differentiated loci as well as

the contribution of recent selective events in ongoing adaptation.

2. Materials and methods

(a) Mosquito species

We surveyed 28 locations within the geographic ranges of species of the *An. nili*

group previously described in Cameroon (Figure 1) (Antonio-Nkondjio et al., 2009; Awono-

Ambene et al., 2004; Awono-Ambene et al., 2006; Ndo et al., 2010, 2013). The genetic

structure of *Anopheles* species is most often based on macrogeographic or regional

subdivisions of gene pools, but can also involve more subtle divergence between larvae and

adults, or between adult populations found in or around human dwellings (e.g. Riehle et al.,

2011). To effectively estimate the genetic diversity and identify potential cryptic

populations within species, we collected larvae and adult mosquitoes within and around

human dwellings using several sampling techniques (Service, 1993) in September-October

2013 (Table S1). To identify the four currently known members of the *An. nili.* group, we

used morphological keys and a diagnostic PCR, which discriminates species on the basis of

point mutations of the ribosomal DNA (Awono-Ambene et al., 2004; Gillies and Coetzee,

1987; Gillies and De Meillon, 1968; Kengne et al., 2003).

(b) Library preparation, sequencing and SNP discovery

We created double-digest Restriction site Associated DNA (ddRAD) libraries as

described in Kamdem et al. 2017 using a modified version of the protocol designed by

Peterson et al*.* 2012. Briefly, genomic DNA of mosquitoes was extracted using the DNeasy

Blood and Tissue kit (Qiagen) and the Zymo Research MinPrep kit for larvae and adult

samples respectively. Approximately 50ng (10μl) of DNA of each mosquito was digested

simultaneously with *MluC1* and *NlaIII* restriction enzymes. Digested products were ligated

to adapter and barcode sequences enabling identification of individuals. Samples were

pooled, purified, and 400-bp fragments selected. The resulting libraries were amplified via

PCR and purified, and fragment size distribution was checked using the BioAnalyzer. PCR

products were quantified, diluted and single-end sequenced to 100 base reads on Illumina

HiSeq2000.

(b) SNP discovery and genotyping

The *process_radtags* program of the Stacks v 1.35 pipeline (Catchen et al., 2013; Catchen et al., 2011) was used to demultiplex and clean Illumina sequences. Reads that passed quality filters were aligned to the *An. nili* Dinderesso draft genome assembly (Giraldo-Calderon et al., 2015) made up of 51048 short contigs (~200-30512bp long) using Gsnap (Wu and Nacu, 2010). To identity and call SNPs within consensus RAD loci, we utilized the *ref_map.pl* program of Stacks. We set the minimum number of reads required to form a stack to three and allowed two mismatches during catalogue creation. We generated SNP files in different formats for further downstream analyses using the *populations* program of Stacks and Plink v1.09 (Purcell et al., 2007).

(d) Population genomics analyses

We analyzed the genetic structure of *An. nili* sensu lato (s.l.) populations using a Principal Component Analysis (PCA) and an unrooted Neighbor-Joining tree (NJ). We also examined ancestry proportions and admixtures between populations in Admixture v1.23 (Alexander et al., 2009) and Structure v2.3.4 (Pritchard et al., 2000). We used the package *adegenet* (Jombart, 2008) to implement the PCA in R (R Development Core Team 2016). The NJ tree was generated from SNP allele frequencies via a matrix of Euclidian distance using the R package *ape* (Paradis et al., 2004). We ran Admixture with 10-fold cross-validation for values of k from 1 through 8. Similarly, we analyzed patterns of ancestry from k ancestral populations in Structure, testing five replicates of k = 1-8. We used 200000 iterations and discarded the first 50000 iterations as burn-in for each Structure run. Clumpp v1.1.2 (Jakobsson and Rosenberg, 2007) was used to summarize assignment

results across independent runs. To identify the optimal number of genetic clusters in our

sample, we applied simultaneously the lowest cross-validation error in Admixture, the ad-

hoc statistic deltaK (Earl and VonHoldt, 2012; Evanno et al., 2005) and the Discriminant

Analysis of Principal Component (DAPC) method implemented in *adegenet.* To examine the

level of genomic divergence among populations, we assessed genetic differentiation (F_{ST})

across SNPs using the *populations* program of the Stacks pipeline. Mean F_{ST} values were also

used to quantify pairwise divergence between populations. To infer the demographic

history of different populations, we used the diffusion approximation method implemented

in the package ∂a∂i v 1.6.3 (Gutenkunst et al., 2009). Single-population models were fitted

to allele frequency spectra, and the best model was selected using the lowest likelihood and

Akaike Information Criterion as well as visual inspections of residuals.

3. Results

(a) SNP genotyping

We collected mosquitoes from four locations out of 28 sampling sites (Figure 1,

Table S1) and sequenced 145 individuals belonging, according to morphological criteria

and diagnostic PCRs, to two species (*An. nili* (n = 24) and *An. ovengensis* (n = 121)). We

assembled 197724 RAD loci that mapped to unique positions throughout the reference

genome. After applying stringent filtering rules, we retained 408 loci present in all

populations and in at least 50% of individuals in each population. Within these loci, we

identified 4343 high-quality biallelic markers that were used to analyze population

structure and genetic differentiation.

(b) Morphologically defined species do not correspond to genetic clusters

PCA and a NJ tree show that the genetic variation across 4343 SNPs is best

explained by more than two clusters, implying subdivisions within *An. nili* and *An.*

ovengensis (Figure 2). Three subgroups are apparent within *An. nili* while two distinct

clusters segregate in *An. ovengensis*. These five subpopulations are associated with the

different sampling sites suggesting local adaptation of divergent populations. Importantly,

Structure and Admixture analyses reveal that, at k = 2, one population identified by

morphology and the diagnostic PCR as *An. nili* has almost the same ancestry pattern as the

largest *An. ovengensis* cluster (Figure 3). Such discrepancies between morphology-based

and molecular taxonomies can be due to a variety of processes including phenotypic

plasticity, introgressive hybrydization or incomplete lineage sorting (i.e., when independent

loci have different genealogies by chance) (Arnold, 1997; Combosch and Vollmer, 2015;

Fontaine et al., 2015; Weng et al., 2016). At k = 2 and k = 3, some populations also depict

half ancestry from each morphological species suggestive of gene flow. We found a

conflicting number of genetic clusters in our samples likely reflecting the complex history of

subdivisions and admixtures among populations (Figure 4). The Evanno et al. method,

which highlights the early states of divergence between *An. nili* and *An. ovengensis* indicates

two probable ancestors. DAPC and the Admixture cross-validation error, which are more

sensitive to recent hierarchical population subdivisions, show five or more distinct clusters

as revealed by PCA and the NJ tree (Figure 4).

As suggested by the long internal branches, which connect subpopulations on the NJ

tree, there is strong differentiation between and within morphological species

characterized by globally high F_{ST} values (Table 1). Relatively lower F_{ST} values observed

between certain clusters may be due to greater inter-population migration and intermixing

or more recent divergence. The F_{ST} values do not reflect the morphological delimitation of

species, and the level of genetic differentiation is even higher between some subpopulations

within the same morphological species. Overall, patterns of genetic structure and

differentiation reveal a group of populations whose phylogenies and species status are

likely confounded by hybridization and/or incomplete lineage sorting. We argue that the

current taxonomy based on morphology and ribosomal DNA does not capture the optimal

reproductive units among populations of this group of mosquitoes.

(c) Genomic signatures of divergent selection and demographic history

We analyzed patterns of genetic differentiation across SNP loci throughout the

genome. Pairwise comparisons are based on filtered variants that satisfy all criteria to be

present in both populations, which explain the discrepancy in the number of SNPs observed

between specific paired comparisons (Figure 5). The distribution of locus-specific F_{ST}

values between the five subpopulations revealed a bimodal shape characterized by two

peaks around 0 and 1. The great majority of SNPs have low to moderate divergence, but a

substantial number of variants are extremely differentiated between populations. The

maximum F_{ST} among SNPs is 1 and the proportion of loci with $F_{ST} = 1$ varies from 6.52%

between the populations we termed *An. nili* group 1 and *An. ovengensis* group 1 to 44.74%

between the subgroups called *An. nili* group 2 and *An. nili* group 3 (Figure 5). This pattern of

genome-wide divergence suggests that a very high number of sites with abrupt

differentiation — which likely contain genes that contribute to divergent selection and/or

reproductive isolation — coexist with regions of weak divergence that can be freely

exchanged between species. As is the case with the overall genetic differentiation,

morphology is not a reliable predictor of locus-specific divergence. Precisely, the lowest

percentage of fixed SNPs is found between *An. ovengensis* from Nyabessan and *An. nili*
collected from Mbébé and Ebebda (Figure 1, Figure 5). In contrast, the greatest proportion
of fixed loci occurs between locally adapted subgroups within the same morphological
species, *An. nili*. The draft reference genome made up of short contigs didn't allow us to test
hypotheses about the genomic distribution of differentiated loci. For example, it remains
unknown whether the numerous SNPs that are fixed among populations are spread
throughout the entire genome or clustered in genomic regions of low recombination
including chromosomal inversions and chromosome centers (Nosil and Feder, 2012; Roesti
et al., 2012).

Models of population demography indicate that all subgroups have experienced an
increase in effective size in a more or less recent past (Table 2). Nevertheless, confidence
intervals of population parameters are high in some populations, and our results should be
interpreted with the necessary precautions. The population growth is less significant in *An.*
nili group 1.

4. Discussion

(a) Genetic differentiation

Advances in sequencing and analytical approaches have opened new avenues for the
study of genomes of disease vectors. We have focused on malaria mosquitoes of the *An. nili*
group, whose taxonomy and population structure have been challenging to resolve with
low-resolution markers. We analysed the genetic structure using genome-wide SNPs and
found strong differentiation and local adaption among populations belonging to the two
morphologically defined species *An. nili* and *An. ovengensis*. The exact number of

subpopulations remains contentious, with the suggested number of divergent clusters

varying from two to five within both species. Significant population structure at eight

microsatellite loci has been described among *An. nili* populations from Cameroon, with F_{ST}

values as high as 0.48 between samples from the rainforest area (Ndo et al., 2013). By

contrast, *An. ovengensis* was discovered recently and the genetic structure of this vector

remains understudied. This species was initially considered as a sibling of *An. nili* (Awono-

Ambene et al., 2004; Awono-Ambene et al., 2006; Kengne et al., 2003), but more recent

studies have started to challenge the assumed relatedness between the two species due to

the high divergence of their polytene chromosomes (Sharakhova et al., 2013). Our findings

call for careful review of the current taxonomy within this group of species, which is a

necessary first step for accurately delineating the role played by the different

subpopulations in malaria transmission.

Our samples were collected from locations characterized by a more or less degraded

forest in the rainforest area of Cameroon. In these habitats, larvae of *An. nili* s.l. exploit

relatively similar breeding sites consisting of slow-moving rivers (Antonio-Nkondjio et al.,

2009). The ecological drivers of genetic differentiation remain unknown, and will be

difficult to infer from our data given the apparent similarity of habitats among populations.

Further study is needed to clearly address the environmental variables that may be

correlated with ongoing adaptive divergence at adult and larval stages. One of the most

expected outcomes of current large-scale malaria control measures that are underway in

Sub-Saharan African countries concerns the effects of increased insecticide exposure on the

genetic diversity and population demography of vectors. A substantial population decline

that may considerably affect the adaptive potential of vector species has been occasionally

reported following the distribution of insecticide-treated bed nets and/or indoor residual

house spraying (e.g. Athrey et al., 2012). The inferred demographic history of the different

subpopulations within the *An. nili* group does not reveal signatures of bottlenecks that can

be potentially correlated with increased usage of insecticides and insecticide-treated nets.

This result is consistent with the demography of several other important malaria vectors of

the Afrotropical region, including *An. gambiae*, *An. coluzzii*, *An. funestus* and *An. moucheti*,

which depict a substantial population increase suggesting that recent intense insecticide

exposure has yet to leave deep or detectable impacts on patterns of genetic variation among

mosquito populations (Fouet et al., 2017; Kamdem et al., 2017; O'Loughlin et al., 2014).

(b) Genomic architecture of geographic and reproductive isolation

Understanding the genomic architecture of reproductive isolation may reveal crucial

information on the sequence of events that occur between the initial stages of divergence

among populations and the onset of strong reproductive barriers (e.g. Turner et al., 2005;

Harr 2006; Nadeau et al., 2012; Ellegren et al., 2012; Carneiro et al., 2014; Burri et al.,

2015). One influential concept of speciation coined the "genic view of species" proposes

that boundaries between species are properties of individual genes or genome regions and

not of whole organisms or lineages (Barton and Hewitt, 1985; Harrison and Larson, 2014;

Harrison, 1990; Key, 1968; Nosil and Feder, 2012; Rieseberg et al., 1999; Wu, 2001). We

have discovered a dramatically high number of SNPs that are strongly differentiated

between populations and often fixed within subgroups of *An. nili* s.l. Interpreting this

intriguing pattern of genomic differentiation is not straightforward due to the complex

interactions between numerous forces — including positive or negative selection,

recombination, introgressive hybridization and incomplete lineage sorting — that can affect

the level of divergence among SNPs (Begun and Aquadro, 1992; Cutter and Payseur, 2013;

Harrison and Larson, 2016; Nachman and Payseur, 2012; Roesti et al., 2012). Some of these

variants exhibiting high divergence among populations certainly contain markers of

ecological and/or reproductive isolation. However, as far as reproductive barriers are

concerned, recent studies have indicated a complex relationship between genetic

differentiation and gene flow at the genome level (e.g. Gompert et al., 2012; Hamilton et al.,

2013a, b; Larson et al., 2013, 2014; Parchman et al., 2013; Taylor et al., 2014). Highly

divergent genomic regions do not necessarily coincide with regions of reduced gene flow.

Several alternative interpretations exist for the numerous high-F_{ST} regions we detected in

all pairwise comparisons (Cruickshank and Hahn, 2014; Delmore et al., 2015; Nachman and

Payseur, 2012; Noor and Bennett, 2009). Nevertheless, careful examination of these outliers

of differentiation may reveal significant insights into the wide range of genes and traits that

contribute to ecological divergence and/or reproductive isolation between subgroups of *An.*

nili s.l. A complete genome assembly will be necessary to better delineate specific regions of

the genome under natural selection, and thereby clarify the genomic basis of phenotypic

fitness differences between divergent populations. This will also help understand the extent

to which recent selection associated with human interventions contribute to local

adaptation and genetic differentiation as observed in *An. gambiae* and *An. coluzzii* (Kamdem

et al., 2017).

Signals consistent with gene flow between *An. nili* and *An. ovengensis* are apparent

in our data despite significant time since divergence (~3M-yr) (Ndo et al., 2013). Some

individuals display almost half ancestry from each morphological species. The

disagreement between morphology/PCR and molecular taxonomies observed in Structure

and Admixture analyses also suggests that incongruent genealogies may be widespread

along chromosomes due to hybridization. However, hybridization can be difficult to detect

because other factors such as incomplete lineage sorting or technical artefacts can leave

signatures that are similar to those of interspecific gene flow (Liu et al., 2014; Patterson et

al., 2012). A complete reference genome is also needed to analyze the detailed distribution

of genealogies across small genomic windows and to disentangle the relative contribution

of processes that generate the putative admixtures and species confusion observed among

divergent populations (Fontaine et al., 2015; Martin et al., 2013; Weng et al., 2016).

5. Conclusions and implications

Delineating the fine-scale population structure of mosquito populations is crucial for

understanding their epidemiological significance and their potential response to vector

control measures. Moreover, recent malaria control efforts are affecting interspecific gene

flow, genetic differentiation, population demography and natural selection in mosquitoes

(Athrey et al., 2012; Barnes et al., 2017; Clarkson et al., 2014; Kamdem et al., 2017; Norris et

al., 2015). Deciphering the signatures of all these processes across mosquito genomes is

important to minimize their negative impacts on vector control. Our findings shed some

light on the complex evolutionary history and provide a framework for future

investigations into the genetic basis of ecological and reproductive barriers among species

of the *An. nili* group.

Acknowledgements

Funding for this project was provided by the University of California Riverside and NIH
grants 1R01AI113248 and 1R21AI115271 to BJW. We thank inhabitants and administrative
authorities of the sampling sites included in this study for their collaboration.
329
330   **Data Archiving Statement**
331
332   Data for this study are available at: to be completed after manuscript is accepted for
333   publication
334

335   **References**
336

Alexander, D. H., Novembre, J., & Lange, K. (2009). Fast model-based estimation of ancestry
in unrelated individuals. *Genome Research*, *19*, 1655–1664.
Antonio-Nkondjio, C., Kerah, C. H., Simard, F., Awono-Ambene, P., Chouaibou, M.,
Tchuinkam, T., & Fontenille, D. (2006). Complexity of the malaria vectorial system in
Cameroon: contribution of secondary vectors to malaria transmission. *Journal of*
Medical Entomology, *43*, 1215–1221. https://doi.org/10.1603/0022-
2585(2006)43[1215:COTMVS]2.0.CO;2
Antonio-Nkondjio, C., Ndo, C., Costantini, C., Awono-Ambene, P., Fontenille, D., & Simard, F.
(2009). Distribution and larval habitat characterization of Anopheles moucheti,
Anopheles nili, and other malaria vectors in river networks of southern Cameroon.
Acta Tropica, *112*(3), 270–276. https://doi.org/10.1016/j.actatropica.2009.08.009
Antonio-Nkondjio, C., & Simard, F. (2013). Highlights on Anopheles nili and Anopheles
moucheti, Malaria Vectors in Africa. In S. Manguin (Ed.), *Anopheles mosquitoes - New*
insights into malaria vectors (INTECH).
Arnold, M. L. (1997). *Natural Hybridization and Evolution. Oxford University Press, Oxford*.
Athrey, G., Hodges, T. K., Reddy, M. R., Overgaard, H. J., Matias, A., Ridl, F. C., … Slotman, M. a.
(2012). The effective population size of malaria mosquitoes: large impact of vector
control. *PLoS Genetics*, *8*(12), e1003097.
https://doi.org/10.1371/journal.pgen.1003097
Awono-Ambene, H. P., Kengne, P., Simard, F., Antonio-Nkondjio, C., & Fontenille, D. (2004).
Description and bionomics of Anopheles (Cellia) ovengensis (Diptera: Culicidae), a new
malaria vector species of the Anopheles nili group from south Cameroon. *Journal of*
Medical Entomology, *41*, 561–568. https://doi.org/10.1603/0022-2585-41.4.561
Awono-Ambene, H. P., Simard, F., Antonio-Nkondjio, C., Cohuet, A., Kengne, P., & Fontenille,
D. (2006). Multilocus enzyme electrophoresis supports speciation within the
Anopheles nili group of malaria vectors in Cameroon. *The American Journal of Tropical*
Medicine and Hygiene, *75*, 656–658. https://doi.org/75/4/656 [pii]
Barnes, K. G., Weedall, G. D., Ndula, M., Irving, H., Mzihalowa, T., Hemingway, J., & Wondji, C.
S. (2017). Genomic Footprints of Selective Sweeps from Metabolic Resistance to
Pyrethroids in African Malaria Vectors Are Driven by Scale up of Insecticide-Based
Vector Control. *PLoS Genetics*, *13(2): e10*, 1–22.
https://doi.org/10.1371/journal.pgen.1006539
Barton, N., & Hewitt, G. . (1985). Analysis of hybrid zones. *Annual Review of Ecology and*
Systematics, *16*, 113–148.
Begun, D., & Aquadro, C. (1992). Levels of naturally occurring DNA polymorphism correlate

with recombination rates in D. melanogaster. *Nature*, *356*, 519–520.

Bøgh, C., Pedersen, E. M., Mukoko, D. a, & Ouma, J. H. (1998). Permethrin-impregnated
bednet effects on resting and feeding behaviour of lymphatic filariasis vector
mosquitoes in Kenya. *Medical and Veterinary Entomology*, *12*(1), 52–9.

Burri, R., Nater, A., Kawakami, T., Mugal, C. F., Olason, P. I., Smeds, L., … Ellegren, H. (2015).
Linked selection and recombination rate variation drive the evolution of the genomic
landscape of differentiation across the speciation continuum of Ficedula flycatchers.
Genome Research, *25*(11), 1656–1665. https://doi.org/10.1101/gr.196485.115

Carneiro, M., Albert, F. W., Afonso, S., Pereira, R. J., Burbano, H., Campos, R., … Ferrand, N.
(2014). The genomic architecture of population divergence between subspecies of the
European Rabbit. *PLoS Genetics*, *10*(8). https://doi.org/10.1371/journal.pgen.1003519

Catchen, J., Hohenlohe, P. a., Bassham, S., Amores, A., & Cresko, W. . (2013). Stacks: An
analysis tool set for population genomics. *Molecular Ecology*, *22*, 3124–3140.
https://doi.org/10.1111/mec.12354

Catchen, J. M., Amores, A., Hohenlohe, P., Cresko, W., & Postlethwait, J. H. (2011). Stacks:
building and genotyping Loci de novo from short-read sequences. *G3 (Bethesda, Md.)*,
1(3), 171–82. https://doi.org/10.1534/g3.111.000240

Clarkson, C. S., Weetman, D., Essandoh, J., Yawson, A. E., Maslen, G., Manske, M., … Donnelly,
M. J. (2014). Adaptive introgression between Anopheles sibling species eliminates a
major genomic island but not reproductive isolation. *Nature Communications*, *5*(May),
4248. https://doi.org/10.1038/ncomms5248

Coetzee, M., & Koekemoer, L. (2013). Molecular Systematics and Insecticide Resistance in
the Major African Malaria Vector Anopheles funestus. *Annual Review of Entomology*.
https://doi.org/10.1146/annurev-ento-120811-153628

Combosch, D. J., & Vollmer, S. V. (2015). Trans-Pacific RAD-Seq population genomics
confirms introgressive hybridization in Eastern Pacific Pocillopora corals. *Molecular
Phylogenetics and Evolution*, *88*, 154–162.
https://doi.org/10.1016/j.ympev.2015.03.022

Cruickshank, T. E., & Hahn, M. W. (2014). Reanalysis suggests that genomic islands of
speciation are due to reduced diversity, not reduced gene flow. *Molecular Ecology*,
23(13), 3133–57. https://doi.org/10.1111/mec.12796

Cutter, A. D., & Payseur, B. A. (2013). Genomic signatures of selection at linked sites:
unifying the disparity among species. *Nature Reviews. Genetics*, *14*(4), 262–74.
https://doi.org/10.1038/nrg3425

Delmore, K. E., Hübner, S., Kane, N. C., Schuster, R., Andrew, R. L., Câmara, F., … Irwin, D. E.
(2015). Genomic analysis of a migratory divide reveals candidate genes for migration
and implicates selective sweeps in generating islands of differentiation. *Molecular
Ecology*, (24), 1873–1888. https://doi.org/10.1111/mec.13150

Derua, Y. a, Alifrangis, M., Hosea, K. M., Meyrowitsch, D. W., Magesa, S. M., Pedersen, E. M., &
Simonsen, P. E. (2012). Change in composition of the Anopheles gambiae complex and
its possible implications for the transmission of malaria and lymphatic filariasis in
north-eastern Tanzania. *Malaria Journal*, *11*(1), 188. https://doi.org/10.1186/1475-
2875-11-188

Dia, I., Guelbeogo, M., & Ayala, D. (2013). Advances and Perspectives in the Study of the
Malaria Mosquito Anopheles funestus. In S. Manguin (Ed.), *Anopheles mosquitoes - New
insights into malaria vectors* (INTECH).

Earl, D. A., & VonHoldt, B. M. (2012). STRUCTURE HARVESTER: a website and program for
visualizing STRUCTURE output and implementing the Evanno method. *Conservation*
Genetics Resources, *4*(359– 361).
Ellegren, H., Smeds, L., Burri, R., Olason, P. I., Backström, N., Kawakami, T., … Wolf, J. B. W.
(2012). The genomic landscape of species divergence in Ficedula flycatchers. *Nature*,
491(7426), 756–60. https://doi.org/10.1038/nature11584
Evanno, G., Goudet, J., & Regnaut, S. (2005). Detecting the number of clusters of individuals
using the software structure: a simulation study. *Molecular Ecology*, *14*, 2611–2620.
Fontaine, M. C., Pease, J. B., Steele, a., Waterhouse, R. M., Neafsey, D. E., Sharakhov, I. V., …
Besansky, N. J. (2015). Extensive introgression in a malaria vector species complex
revealed by phylogenomics. *Science*, *347*(6217), 1258524–1258524.
https://doi.org/10.1126/science.1258524
Fouet, C., Kamdem, C., Gamez, S., & White, B. J. (2017). Extensive genetic diversity among
populations of the malaria mosquito Anopheles moucheti revealed by population
genomics. *Infection, Genetics and Evolution*, *48*, 27–33.
https://doi.org/10.1016/j.meegid.2016.12.006
Gillies, M. T., & Coetzee, M. (1987). *A supplement to the Anophelinae of Africa south of the*
Sahara. Johannesburg: The South African Institute for Medical Research.
Gillies, M. T., & De Meillon, B. (1968). *The Anophelinae of Africa South of the Sahara* (Second
Edi). Johannesburg: Publications of the South African Institute for Medical Research.
Giraldo-Calderon, G. I., Emrich, S. J., MacCallum, R. M., Maslen, G., Dialynas, E., Topalis, P., …
Lawson, D. (2015). VectorBase: an updated bioinformatics resource for invertebrate
vectors and other organisms related with human diseases. *Nucleic Acids Research*,
43(D1), D707–D713. https://doi.org/10.1093/nar/gku1117
Gompert, Z., Lucas, L. K., Nice, C. C., Fordyce, J. A., Forister, M. L., & Buerkle, C. A. (2012).
Genomic regions with a history of divergent selection affect fitness of hybrids between
two butterfly species. *Evolution*, *66*(7), 2167–2181.
https://doi.org/10.5061/dryad.f0b2f083
Gutenkunst, R. N., Hernandez, R. D., Williamson, S. H., & Bustamante, C. D. (2009). Inferring
the joint demographic history of multiple populations from multidimensional SNP
frequency data. *PLoS Genetics*, *5*(10), e1000695.
https://doi.org/10.1371/journal.pgen.1000695
Hamilton, J. a., Lexer, C., & Aitken, S. N. (2013). Differential introgression reveals candidate
genes for selection across a spruce (Picea sitchensis × P. glauca) hybrid zone. *New*
Phytologist, *197*(3), 927–938. https://doi.org/10.1111/nph.12055
Hamilton, J. A., Lexer, C., & Aitken, S. N. (2013). Genomic and phenotypic architecture of a
spruce hybrid zone (Picea sitchensis × P. glauca). *Molecular Ecology*, *22*(3), 827–841.
https://doi.org/10.1111/mec.12007
Harbach, R. E. (2013). The Phylogeny and Classification of Anopheles. In S. Manguin (Ed.),
Anopheles mosquitoes - New insights into malaria vectors (INTECH).
Harr, B. (2006). Genomic islands of differentiation between house mouse subspecies, 730–
737. https://doi.org/10.1101/gr.5045006.entiation
Harrison, R. G. (1990). Hybrid zones: windows on evolutionary process. In D. Futuyma & J.
Antonovics (Eds.) (Oxford Uni, pp. 69–128). New York.
Harrison, R. G., & Larson, E. L. (2014). Hybridization, introgression, and the nature of
species boundaries. *Journal of Heredity*, *105*(S1), 795–809.

https://doi.org/10.1093/jhered/esu033

Harrison, R. G., & Larson, E. L. (2016). Heterogeneous genome divergence, differential
introgression, and the origin and structure of hybrid zones. *Molecular Ecology*, *25*,
2454–2466. https://doi.org/10.1111/mec.13582

Hemingway, J., Ranson, H., Magill, A., Kolaczinski, J., Fornadel, C., Gimnig, J., … Hamon, N.
(2016). Averting a malaria disaster: Will insecticide resistance derail malaria control?
The Lancet, *387*(10029), 1785–1788. https://doi.org/10.1016/S0140-6736(15)00417-
1

Jakobsson, M., & Rosenberg, N. (2007). CLUMPP: a cluster matching and permutation
program for dealing with multimodality in analysis of population structure.
Bioinformatics, *23*, 1801– 1806.

Jombart, T. (2008). adegenet: a R package for the multivariate analysis of genetic markers.
Bioinformatics, *24*, 1403–1405.

Kamdem, C., Fouet, C., Gamez, S., & White, B. J. (2017). Pollutants and insecticides drive local
adaptation in African malaria mosquitoes. *Molecular Biology and Evolution*.
https://doi.org/DOI: 10.1093/molbev/msx087 [Epub ahead of print]

Kengne, P., Awono-Ambene, H. P. Antonio-Nkondjio, C., Simard, F., & Fontenille, D. (2003).
Molecular identification of the Anopheles nili group African malaria vectors. *Med Vet*
Entomol, *17*, 67–74.

Key, K. H. L. (1968). The concept of stasipatric speciation. *Syst. Zool.*, *17*, 14–22.

Krzywinski, J., & Besansky, N. J. (2003). Molecular systematics of Anopheles : from
subgenera to subpopulations. *Annual Review of Entomology*, *48*(1), 111–139.
https://doi.org/10.1146/annurev.ento.48.091801.112647

Lanzaro, G. C., & Lee, Y. (2013). Speciation in Anopheles gambiae — The Distribution of
Genetic Polymorphism and Patterns of Reproductive Isolation Among Natural
Populations. In S. Manguin (Ed.), *Anopheles mosquitoes - New insights into malaria*
vectors (INTECH).

Larson, E. L., Andrés, J. A., Bogdanowicz, S. M., & Harrison, R. G. (2013). Differential
introgression in a mosaic hybrid zone reveals candidate barrier genes. *Evolution*,
67(12), 3653–3661. https://doi.org/10.1111/evo.12205

Larson, E. L., White, T. A., Ross, C. L., & Harrison, R. G. (2014). Gene flow and the
maintenance of species boundaries. *Molecular Ecology*, *23*(7), 1668–1678.
https://doi.org/10.1111/mec.12601

Lawniczak, M. K. N., Emrich, S. J., Holloway, a K., Regier, a P., Olson, M., White, B., …
Besansky, N. J. (2010). Widespread divergence between incipient Anopheles gambiae
species revealed by whole genome sequences. *Science (New York, N.Y.)*, *330*(6003),
512–4. https://doi.org/10.1126/science.1195755

Liu, K. J., Dai, J., Truong, K., Song, Y., Kohn, M. H., & Nakhleh, L. (2014). An HMM-Based
Comparative Genomic Framework for Detecting Introgression in Eukaryotes. *PLoS*
Computational Biology, *10*(6). https://doi.org/10.1371/journal.pcbi.1003649

Martin, S. H., Dasmahapatra, K. K., Nadeau, N. J., Salazar, C., Walters, J. R., Simpson, F., …
Jiggins, C. D. (2013). Genome-wide evidence for speciation with gene flow in Heliconius
butterflies. *Genome Research*, *23*(11), 1817–28.
https://doi.org/10.1101/gr.159426.113

Molineaux, L., & Gramiccia, G. (1980). *The Garki Project. Research on the epidemiology and*
control of malaria in the sudan savanna of West Africa. World Health Organization.

Mwangangi, J. M., Mbogo, C. M., Orindi, B. O., Muturi, E. J., Midega, J. T., Nzovu, J., ... Beier, J. C.
(2013). Shifts in malaria vector species composition and transmission dynamics along
the Kenyan coast over the past 20 years. *Malaria Journal*, *12*, 13.
https://doi.org/10.1186/1475-2875-12-13
Nachman, M. W., & Payseur, B. A. (2012). Recombination rate variation and speciation :
theoretical predictions and empirical results from rabbits and mice. *Philosophical*
Transactions of the Royal Society B: Biological Sciences, *367*, 409–421.
https://doi.org/10.1098/rstb.2011.0249
Nadeau, N. J., Whibley, A., Jones, R. T., Davey, J. W., Dasmahapatra, K. K., Baxter, S. W., ...
Jiggins, C. D. (2012). Genomic islands of divergence in hybridizing Heliconius
butterflies identified by large-scale targeted sequencing, 343–353.
https://doi.org/10.1098/rstb.2011.0198
Ndo, C., Antonio-Nkondjio, C., Cohuet, A., Ayala, D., Kengne, P., Morlais, I., ... Simard, F.
(2010). Population genetic structure of the malaria vector Anopheles nili in sub-
Saharan Africa. *Malaria Journal*, *9*, 161. https://doi.org/10.1186/1475-2875-9-161
Ndo, C., Simard, F., Kengne, P., Awono-Ambene, P., Morlais, I., Sharakhov, I., ... Antonio-
Nkondjio, C. (2013). Cryptic Genetic Diversity within the Anopheles nili group of
Malaria Vectors in the Equatorial Forest Area of Cameroon (Central Africa). *PLoS ONE*,
8(3), 1–12. https://doi.org/10.1371/journal.pone.0058862
Neafsey, D. E., Lawniczak, M. K. N., Park, D. J., Redmond, S. N., Coulibaly, M. B., Traoré, S. F., ...
Muskavitch, M. a T. (2010). SNP genotyping defines complex gene-flow boundaries
among African malaria vector mosquitoes. *Science (New York, N.Y.)*, *330*(6003), 514–7.
https://doi.org/10.1126/science.1193036
Noor, M. a F., & Bennett, S. M. (2009). Islands of speciation or mirages in the desert?
Examining the role of restricted recombination in maintaining species. *Heredity*,
103(6), 439–444. https://doi.org/10.1038/hdy.2010.13
Norris, L. C., Main, B. J., Lee, Y., Collier, T. C., Fofana, A., Cornel, A. J., & Lanzaro, G. C. (2015).
Adaptive introgression in an African malaria mosquito coincident with the increased
usage of insecticide-treated bed nets. *Proceedings of the National Academy of Sciences*,
201418892. https://doi.org/10.1073/pnas.1418892112
Nosil, P., & Feder, J. L. (2012). Widespread yet heterogeneous genomic divergence.
Molecular Ecology, *21*(12), 2829–2832. https://doi.org/10.1111/j.1365-
294X.2012.05580.x
O'Loughlin, S. M., Magesa, S., Mbogo, C., Mosha, F., Midega, J., Lomas, S., & Burt, A. (2014).
Genomic Analyses of Three Malaria Vectors Reveals Extensive Shared Polymorphism
but Contrasting Population Histories. *Molecular Biology and Evolution*, 1–14.
https://doi.org/10.1093/molbev/msu040
Paradis, E., Claude, J., & Strimmer, K. (2004). Analyses of Phylogenetics and Evolution in R
language. *Bioinformatics*, *20*(2), 289–290.
Parchman, T. L., Gompert, Z., Braun, M. J., Brumfield, R. T., McDonald, D. B., Uy, J. a C., ...
Buerkle, C. a. (2013). The genomic consequences of adaptive divergence and
reproductive isolation between species of manakins. *Molecular Ecology*, *22*(12), 3304–
17. https://doi.org/10.1111/mec.12201
Patterson, N., Moorjani, P., Luo, Y., Mallick, S., Rohland, N., Zhan, Y., ... Reich, D. (2012).
Ancient admixture in human history. *Genetics*, *192*(3), 1065–1093.
https://doi.org/10.1534/genetics.112.145037

556 Peery, A., Sharakhova, M. V, Antonio-Nkondjio, C., Ndo, C., Weill, M., Simard, F., & Sharakhov,
557   I. V. (2011). Improving the population genetics toolbox for the study of the African
558   malaria vector Anopheles nili: microsatellite mapping to chromosomes. *Parasites &*
559   *Vectors*, *4*(1), 202. https://doi.org/10.1186/1756-3305-4-202
560 Peterson, B. K., Weber, J. N., Kay, E. H., Fisher, H. S., & Hoekstra, H. E. (2012). Double Digest
561   RADseq: An Inexpensive Method for De Novo SNP Discovery and Genotyping in Model
562   and Non-Model Species. *PLoS ONE*, *7*(5), e37135.
563   https://doi.org/10.1371/journal.pone.0037135
564 Pritchard, J. K., Stephens, M., & Donnelly, P. (2000). Inference of population structure using
565   multilocus genotype data. *Genetics*, *155*, 945–959.
566 Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M., Bender, D., … Sham, P. (2007).
567   PLINK: a toolset for whole-genome association and population-based linkage analysis.
568   *American Journal of Human Genetics*, *81*.
569 Ranson, H., & Lissenden, N. (2016). Insecticide Resistance in African Anopheles Mosquitoes:
570   A Worsening Situation that Needs Urgent Action to Maintain Malaria Control. *Trends in*
571   *Parasitology*, *32*(3), 187–196. https://doi.org/10.1016/j.pt.2015.11.010
572 R Development Core Team (2016). *R: A language and environment for statistical computing*.
573   R Foundation for Statistical Computing, Vienna, Austria.
574 Riehle, M. M., Guelbeogo, W. M., Gneme, A., Eiglmeier, K., Holm, I., Bischoff, E., … Vernick, K.
575   D. (2011). A cryptic subgroup of Anopheles gambiae is highly susceptible to human
576   malaria parasites. *Science (New York, N.Y.)*, *331*(6017), 596–8.
577   https://doi.org/10.1126/science.1196759
578 Rieseberg, L. H., Whitton, J., & Gardner, K. (1999). Hybrid Zones and the Genetic
579   Architecture of a Barrier to Gene Flow Between Two Sunflower Species.
580 Roesti, M., Hendry, A. P., Salzburger, W., & Berner, D. (2012). Genome divergence during
581   evolutionary diversification as revealed in replicate lake-stream stickleback population
582   pairs. *Molecular Ecology*, *21*(12), 2852–62. https://doi.org/10.1111/j.1365-
583   294X.2012.05509.x
584 Service, M. W. (1993). *Mosquito ecology: field sampling methods*. (U. K. Elsevier Applied
585   Science, London, Ed.).
586 Sharakhova, M. V., Peery, A., Antonio-Nkondjio, C., Xia, A., Ndo, C., Awono-Ambene, P., …
587   Sharakhov, I. V. (2013). Cytogenetic analysis of Anopheles ovengensis revealed high
588   structural divergence of chromosomes in the Anopheles nili group. *Infection, Genetics*
589   *and Evolution*, *16*, 341–348. https://doi.org/10.1016/j.meegid.2013.03.010
590 Sinka, M. E., Bangs, M. J., Manguin, S., Coetzee, M., Mbogo, C. M., Hemingway, J., … Hay, S. I.
591   (2010). The dominant Anopheles vectors of human malaria in Africa, Europe and the
592   Middle East: occurrence data, distribution maps and bionomic précis. *Parasites &*
593   *Vectors*, *3*(1), 117. https://doi.org/10.1186/1756-3305-3-117
594 Taylor, S. A., Curry, R. L., White, T. A., Ferretti, V., & Lovette, I. (2014). Spatiotemporally
595   consistent genomic signatures of reproductive isolation in a moving hybrid zone.
596   *Evolution*, *68*(11), 3066–3081. https://doi.org/10.1111/evo.12510
597 Turner, T. L., Hahn, M. W., & Nuzhdin, S. V. (2005). Genomic islands of speciation in
598   Anopheles gambiae. *PLoS Biology*, *3*(9), 1572–1578.
599   https://doi.org/10.1371/journal.pbio.0030285
600 Van Bortel, W., Harbach, R. E., Trung, H. D., Roelants, P., Backeljau, T., & Coosemans, M.
601   (2001). Confirmation of Anopheles varuna in vietnam, previously misidentified and

mistargeted as the malaria vector Anopheles minimus, *65*(April 1999), 729–732.
Weng, D., Yu, Y., Hahn, M. W., & Nakhleh, L. (2016). Reticulate evolutionary history and
extensive introgression in mosquito species revealed by phylogenetic network
analysis. *Molecular Ecology*, *Special is.* https://doi.org/10.1111/mec.13544
WHO. (2016). *World malaria report 2016*.
Wright, S. (1984). *Evolution and the Genetics of Populations: Genetics and Biometric*
Foundations v. 1 (Genetic & Biometric Foundations); New Edition. University of Chicago
press.
Wu, C. (2001). The genic view of the process of speciation. *Journal of Evolutionary Biology*,
14, 851–865.
Wu, T. D., & Nacu, S. (2010). Fast and SNP-tolerant detection of complex variants and
splicing in short reads. *Bioinformatics*, *26*(7), 873–881.

Author contributions

Conceived and designed the experiments: CF CK BJW. Performed the experiments: CF CK SG
BJW. Analysed the data: CF CK. Wrote the paper: CF CK BJW.

618
619
620

Tables

Table 1: Pairwise F_{ST} between divergent subpopulations of *An. Nili* s.l.

F_{ST}	*An. nili* group 1	*An. nili* group 2	*An. nili* group 3	*An. ovengensis* group 1	*An. ovengensis* group 2
An. nili **group 1**	-				
An. nili **group 2**	0.374	-			
An. nili **group 3**	0.506	0.552	-		
An. ovengensis **group 1**	0.135	0.275	0.364	-	
An. ovengensis **group 2**	0.432	0.458	0.492	0.349	-

Table 2: Demographic models of different subgroups of *An. nili* s.l.

Population	Best Model	Log Likelihood	Final Population Size[a] (95% CI)	Time[b] (95% CI)
An. nili group 1	*Growth*	-18.42	6.41 (5.326 - 20.71)	3.70 (1.11 - 13.31)
An. nili group 2	*Two-epoch*	-19.97	17.87 (9.33 - 35.50)	11.27 (4.93 - 19.64)
An. ovengensis group 1	*Growth*	-112.18	13.04 (12.15 - 17.26)	0.70 (0.58 - 1.08)
An. Ovengensis group 2	*Growth*	-22.98	19.95 (14.45 - 45.70)	5.11 (2.33 - 15.13)

[a] Relative to ancestral population size.

[b] Expressed in units 2Ne generations from start of growth to present.

 Figures

627 **Figure 1**: Map showing the sampling locations and relative frequencies of the
628 morphological species *An. nili* and *An. ovengensis*. Small and large black dots indicate
629 respectively the 28 locations surveyed and the four sampling sites where mosquitoes were
630 collected.
631

632
633

Figure 2: Population genetic structure inferred from 4343 SNPs using PCA (A) and a neighbor-joining tree (B). The percentage of variance explained is indicated on each PCA axis.

641 **Figure 3:** Ancestry proportions inferred in ADMIXTURE with k = 2 – 8.
642

643
644
645

646 **Figure 4:** Identification of the optimal number of genetic clusters using the delta k method
647 of Evanno et al (A), DAPC (B) and 10-fold cross-validation in ADMIXTURE (C). The lowest
648 BIC and CV error and the highest delta k indicate the most probable number of clusters.
649
650

651
652
653

Figure 5: Distribution of F_{ST} values throughout the genome between *An. nili* group 1 and *An.*
nili group 2 (A); *An. nili* group 1 and *An. nili* group 3 (B); *An. nili* group 1 and *An. ovengensis*
group 1 (C); *An. nili* group 1 and *An. ovengensis* group 2 (D); *An. nili* group 2 and *An. nili*
group 3 (E); *An. nili* group 2 and *An. ovengensis* group 1 (F); *An. nili* group 2 and *An.*
ovengensis group 2 (G); *An. nili* group 3 and *An. ovengensis* group 1 (H); *An. nili* group 3 and
An. ovengensis group 2 (I); *An. ovengensis* group 1 and *An. ovengensis* group 2 (J). The
number of SNPs with $F_{ST} = 1$ is indicated in each pairwise comparison as well as the total
number of SNPs in parenthesis.
662

663
664

Supplemental information

Table S1: Information on mosquito samples included in this study.

Sampling locations	Geographic coordinates	Sampling methods			Total
		HLC-OUT	**HLC-IN**	**LC**	
Ebebda	4°20'00"N, 11°17'00"E			6	6
Nkoteng	4°31'00"N, 12°02'00"E			8	8
Nyabessan	2°24'00"N, 10°24'00"E	63	44		107
Mbébé	4°10'00"N, 11°04'00"E	13	3	8	24
Total		76	47	22	145

HLC-OUT, human landing catches performed outdoor; HLC-IN, human landing catches performed indoor; LC, larval collection