1 # Genomic signatures of introgression at late stages of speciation in a
2 # malaria mosquito
3

Colince Kamdem[1]*, Caroline Fouet[1], Stephanie Gamez[1], Bradley J. White[1,2]*

[1]Department of Entomology, University of California, Riverside, CA 92521

[2]Center for Disease Vector Research, Institute for Integrative Genome Biology, University of

California, Riverside, CA 92521

*Corresponding authors: colincek@ucr.edu; bwhite@ucr.edu

Key words: Introgression, speciation, linkage disequilibrium, divergent selection,

Anopheles nili

Abstract

Hybridization plays a central role during the evolution of species boundaries, but the relative impact of gene flow on genomic divergence and vice versa remains largely unknown. The genome architecture of populations and emerging species exhibiting various levels of divergence along the speciation continuum should provide insights into the events that promote or prevent speciation. In this work, we have used a combination of population genomic approaches to examine the genomic signatures of hybridization between *Anopheles nili sensu stricto* and *An. ovengensis*, two malaria mosquitoes that have split ~3-Myr ago. Despite this substantial time since divergence, the two species hybridize extensively in nature, giving rise to a unique population of differentiated hybrids in contact zones. Using genomic clines and Bayesian models, we showed that signatures of introgression are widespread across the genome suggesting that recent hybridization between *Anopheles nili sensu stricto* and *An. ovengensis* involves multiple fitness traits and functional classes. Linkage Disequilibrium analyses allowed us to identified a block of 39 linked loci that segregated between hybrids and parental species and may harbour genes responsible for reproductive isolation. Our results demonstrate that genome-wide admixtures can persist in the face of species divergence over long periods of time during speciation due to increased gene flow at loci providing selective advantage.

Introduction

Most cases of speciation occur in a gradual manner so that emerging species continue to

mate and exchange genes before the onset of complete reproductive isolation (Nosil 2012;

Nadeau *et al.* 2013). As a result, secondary contacts between diverging lineages are

pervasive in nature and can lead to several scenarios including extensive hybridization and

the creation of a new hybrid species or genetic homogenization (James 2007; Ozerov *et al.*

2016). The role of hybridization either in the creation of biodiversity through speciation or

in its reduction through genetic homogenization has been under increasing scrutiny over

the last decades and has been recognized as one major force driving the evolution of plant

and animal species (James 2007; Abbott *et al.* 2013; Mallet *et al.* 2015). However, although

there is consistent evidence that hybridization can provide the raw material for

evolutionary diversification (Grant 2015), examples of homoploid hybrid species (where

the hybrid offspring has the same ploidy level as the two parental species) are rather rare in

nature (James 2007; Abbott *et al.* 2013). Most often, hybrids have reduced fitness relative to

parental populations and barely persist as a reproductively isolated unit in the wild.

It is particularly difficult to predict the outcome of secondary contacts between diverging

lineages and specifically the rate of hybridization (Payseur & Rieseberg 2016). One main

issue is that the level of divergence between species supposed to encourage or prevent

hybridization remains obscure. Until a recent past, estimates of the population

differentiation parameter F_{ST} have been used to predict the rate of hybridization between

incomplete species. Variations in F_{ST} among groups of related species were often

interpreted as indicative of differences in rates of gene flow across populations, high F_{ST}

values being considered as evidence of limited gene flow. However, instances of

hybridization and introgression between highly divergent taxa are accumulating, thereby

indicating that estimates of population differentiation are particularly poor predictors of

hybridization between two species (Nydam & Harrison 2011; Roux *et al.* 2013; Parchman *et*

al. 2013; Martin *et al.* 2013; Canestrelli *et al.* 2014).

Likewise, genomic regions showing high differentiation among populations have been

thought to harbor hybrid incompatibility or other reproductive isolation (RI) factors. In

support of this view, genome scans in many couples of incipient species have identified

genomic regions of extreme F_{ST} values qualified as "speciation islands", considered to be

enriched in RI factors and resistant to gene flow (Turner *et al.* 2005; Hohenlohe *et al.* 2010;

Lawniczak *et al.* 2010; Ellegren *et al.* 2012). Yet, not only cases where genes involved in RI

have been explicitly identified in F_{ST} outliers regions of the genome are extremely rare, but

also extensive introgression has been found to sometimes coincide with regions of high

differentiation. For example, (Parchman *et al.* 2013) used genomic clines and Bayesian

models to study the correlation between regions of exceptional introgression and genetic

differentiation manakins birds (*Manacus*). Contrary to expectations, they found that loci of

strong introgression were relatively positively correlated to the genetic differentiation. The

most plausible interpretation of this intriguing result is that divergent selection may

promote introgression of select genes or genomic regions, resulting in differential patterns

of introgression across the genome. A relative concordance between locus-specific

divergence and locus-specific measures of introgression has also been described in the

Lycaeides butterfly (Gompert *et al.* 2012, 2013) and in *Drosophila melanogaster* (Pool *et al.*

2012). Overall, the relationship between introgressive hybridization and genetic divergence

at the species or the genome level is more complex than previously thought. Whether

exceptionally differentiated regions of the genome harbor genes that can, paradoxically,

introgress easily is an empirical question that remains opened and should be answered by

studying closely related species at hybrid zones.

Mosquitoes of the genus *Anopheles* are ideal system in which to address the evolutionary

processes at hybrid zones due to the prevalence of adaptive speciation (Ndo *et al.* 2013;

Neafsey *et al.* 2015; Kamdem *et al.* 2016). In this paper, we focused on *An. nili*, a group of

African malaria vectors characterized by a reticulate evolution leading to complex

phylogenies that have been challenging to clarify (Kengne *et al.* 2003; Awono-Ambene *et al.*

2004, 2006; Ndo *et al.* 2010, 2013; Peery *et al.* 2011; Sharakhova *et al.* 2013). The group

harbour four known species identified by slight morphological differences: *An. nili sensu*

stricto, *An. ovengensis*, *An. carnevalei*, and *An. somalicus. An. nili sensu stricto* (thereafter *An.*

nili) is the most widespread across the continent while the three other species are more

patchily distributed primarily in the equatorial rainforest. *An. nili* and *An. ovengensis* are

important vectors of human malaria parasites (Antonio-Nkondjio *et al.* 2006). We sampled

populations across the ranges of the four species of the *An. nili* group in Cameroon and we

used a population resequencing approach to develop genome-wide SNP markers that we

genotyped in 145 individuals. Our first aim was to clarify the evolutionary relationships

between populations. We discovered new cryptic species within *An. ovengensis* and *An. nili*

as well as a hybrid subgroup resulting from massive hybridization in sympatric areas.

Finally, we took advantage of the recent implementation of genomic cline models that

enables the investigation of footprints of introgression across genomes of non-model

organisms without the need of a high-quality reference genome (Gompert & Alex 2011;

Gompert & Buerkle 2012), to critically evaluate the interplays between the locus-specific

divergent selection and introgression.

Materials and methods

Mosquito sample

We surveyed a total of 28 geographic locations that were representative of the main habitats of species of the *An. nili* group previously described in Cameroon (Fig 1A) (Awono-Ambene *et al.* 2004, 2006; Antonio-Nkondjio *et al.* 2009; Ndo *et al.* 2010, 2013). The different species were identified using reference morphological identification keys and a diagnostic PCR that discriminates the four currently known members of the *An. nili.* group on the basis of a point mutation of the ribosomal DNA (Gillies & De Meillon 1968; Gillies & Coetzee 1987; Kengne *et al.* 2003).

Library preparation, sequencing and SNP discovery

We created double-digest RAD (ddRAD) libraries using a modified version of the protocol described by Peterson et al, 2012 (Peterson *et al.* 2012). Genomic DNA of mosquitoes was extracted using the DNeasy Blood and Tissue kit (Qiagen) and the Zymo Research MinPrep kit on larvae and adult samples respectively. Approximately 50 ng (10µl) of DNA of each mosquito sample was digested simultaneously with *MluC1* and *NlaIII* restriction enzymes. Digested products were then ligated to adapter and barcode sequences to enable the unique identification of the individual associated with each sequencing read. The samples were pooled, purified, and 400-bp fragments selected. The resulting libraries were then amplified via PCR and purified. The distribution of library fragment-size was checked on a BioAnalyzer (Agilent Technologies, Inc., USA). The PCR products were quantified and diluted for sequencing on Illumina HiSeq2000 (Illumina Inc., USA) to yield single-end reads of 101 bp.

Bioinformatics pipeline

The *An. nili* Dinderesso draft genome assembly comprises 51,048 contigs, varying between 100 and 26775kb in length, to which short reads can be aligned and SNP called. However, members of the *An. nili* group found in Cameroon diverged ~0.2-6 million years ago (Ndo *et al.* 2013). Thus, alignments of our samples to *An. nili* Dinderesso reference genome might be subjected to an important bias associated with the inconsistent mapping of reads from highly divergent populations. To make sure that this potential reference sequence bias didn't undermined our analyses, we compared our results across two SNP sets that were identified within RAD loci created using two distinct approaches: a *de novo* assembly and an assembly of reads aligned onto the reference genome. Recent studies using RAD sequencing on *Heliconius* butterflies showed that combining *de novo* assemblies and reference alignments provided a robust approach to perform rigorous test on introgression and phylogenetic relationships in distantly related species (The Heliconius Genome Consortium 2012). The *process_radtags* program of the Stacks v 1.35 pipeline was used to demultiplex and clean raw reads. Reads without the *NlaIII* restriction site and those bearing ambiguous barcode sequences or having low-quality score (average Phred score < 33) were discarded. Reads were trimmed to 96-bp by removing index and barcode sequences. We aligned the short reads to the *An. nili* Dinderesso draft genome assembly using Gsnap (Wu & Nacu 2010) with a maximum of five nucleotide mismatches allowed. The *ref_map.pl* and *denovo_map.pl* programs in Stacks were used to identify consensus RAD loci and to call SNPs within these loci across our populations using respectively the Gsnap-aligned SAM files or the individual fastq files as input. For both analyses, we set the minimum number of

reads required to form a stack to 3. In the *denovo* assembly, we allowed a maximum of three

mismatches when creating loci in every individual (M parameter in *denovo_map.pl*) and two

mismatches when building the catalogue of loci across individuals (n parameter). In

reference-based assembly, we specified n = 2 in *ref_map.pl* to allow two mismatches during

catalogue creation. We generated SNP files in different formats for further downstream

analyses using the *populations* program of Stacks and PLINK v1.09 (Purcell *et al.* 2007).

Population genetic structure

We analyzed the genetic structure of *An. nili s.l.* populations using Principal Component

Analysis (PCA) and Neighbor-Joining trees (NJ). We also examined ancestry proportions

and admixtures between populations in ADMIXTURE v1.23 (Alexander *et al.* 2009) and

STRUCTURE v2.3.4 (Pritchard *et al.* 2000). We performed these tests using filtered SNPs

identified in RAD loci present in every population and in at least 50% of individuals by the

populations program in Stacks. We used the R package *adegenet* (Jombart 2008) to

implement the PCA. Neighbor-Joining trees were generated from matrixes of Euclidian

distance computed from allele frequencies at genome-wide SNPs using the R package *ape*

(Paradis *et al.* 2004). We ran ADMIXTURE with 10-fold cross-validation for values of k from

1 through 20. We analyzed patterns of ancestry from k ancestral populations in

STRUCTURE, testing five replicates of k = 1-10. We used 200000 iterations and discarded

the first 50000 iterations as burn-in for each STRUCTURE run. CLUMPP v1.1.2 (Jakobsson &

Rosenberg 2007) was used to summarize assignment results across independent runs and

DISTRUCT v1.1 (Rosenberg 2004) to plot STRUCTURE results. To identify the optimal

number of genetic clusters in our samples, we used simultaneously the lowest cross-

validation error in ADMIXTURE, the ad-hoc statistic deltaK (Evanno *et al.* 2005; Earl &

VonHoldt 2012) and the Discriminant Analysis of Principal Component (DAPC) method in

adegenet.

Population genomics analyses

To quantify the genetic differentiation between species and putative subgroups within

species, we used a subset of 1000 high-quality SNPs to calculate the overall pairwise

differentiation index F_{ST} (Weir & Cockerham 1984) in Genodive v 1.06 (Meirmans & Van

Tienderen 2004). Statistical significance was assessed with 10000 permutations. To further

examine the genomic footprints of selection and introgression, we used ANGSD v 0.612

(Korneliussen *et al.* 2014) and ngsTools (Fumagalli *et al.* 2014) to derive locus-specific

estimates of the nucleotide diversity (measured as θ_w and θ_π), Tajima's *D*, absolute

sequence divergence (d_{xy}) and F_{ST} across 9622 sites using Gnap alignments without SNP

calling. To identify statistical outliers of F_{ST}, we used the software LOSITAN which applies

the coalescent simulation method (FDIST2) (Beaumont & Nichols 1996) to identify loci with

exceptionally high F_{ST} values relative to a "neutral" genome-wide F_{ST} value expected under

neutral evolution. We ran LOSITAN using the neutral F_{ST} option, 50,000 simulations and a

false discovery rate of 0.05%. Finally, we used the R package LDna (Kemppainen *et al.*

2015) for identifying clusters of linkage disequilibrium in our samples. This analysis was

conducted primarily to assess the possible influence of inversion polymorphism on the

genomic architecture of divergence and hybridization. Precisely, we wished to test if loci

with great significance in the genetic divergence and/or introgression were independent

loci scattered throughout the genome or clusters of linked loci encapsulated in low

recombination regions like chromosomal inversions. We used PLINK to calculate the

linkage disequilibrium (estimated as the r^2 correlation coefficient) between all pairs of SNPs

and we applied the graph-based method implemented in LDna to search for clusters of

strongly correlated SNPs.

Tests of introgression

In addition to the interpretation of patterns of ancestry provided by clustering analyses in

STRUCTURE and ADMIXTURE, we conducted formal tests to infer the history of population

splits and admixtures. We used the three-population (f_3) and the four-population (f_4)

statistics, introduced in (Reich *et al.* 2009) as methods to estimate mixture proportions in

an admixed group, to test for introgression among species. The f_3 and f_4 statistics exploit the

idea that the genetic drift, defined as a function of allele frequency, should be uncorrelated

in unadmixed populations. As a result, correlations detected between allele frequencies of

three (f_3) or four (f_4) populations joined by an unrooted tree indicate episodes of gene flow.

Specifically, the f_3 (X; Y, W) tests for admixture between a test population X and two

reference populations Y and W. The expected value of f_3 is positive in case of no mixture and

negative if X is admixed with Y or W, or both. Similarly, the f_4 for an unrooted tree (A,B;C,D),

tests whether allele frequency differences between A and B are correlated with differences

between C and D. f_4 is equal to zero if there are no correlations and no admixtures across

branches of the tree. In contrast, f_4 is significantly positive if admixture occurs between A

and C, or between B and D, or both, and significantly negative if admixture occurs between

A and D, or between B and C, or both. The statistical significance of f_3 and f_4 values is

assessed using a Z-score: an f_3 or f_4 value divided by its standard deviation. We used a

threshold Z-score of 2.5 corresponding to a p-value of 0.05 as suggested in (Reich *et al.*

2009). The *threepop* and *fourpop* programs of the software *TreeMix* (Pickrell & Pritchard

2012) were used to calculate f_3 and f_4 statistics and Z-scores for all possible triples and

quadruples of populations we described within *An. nili s.l.*

We next applied the graph-based method developed in *TreeMix* to determine the

directionality and quantify the extent of gene flow among species. The *TreeMix* approach

uses allele frequencies at genome-wide polymorphisms and a Gaussian approximation of

the genetic drift among populations to first construct a Maximum Likelihood (ML)

phylogeny connecting sampled populations by simple bifurcations. The model then

compares the covariance structure modeled by this dendrogram and adds edges to the

phylogeny to account for admixtures. We first conducted one *TreeMix* run without

migration. We noted the percentage of explained variance of the models and visually

inspected the residuals of covariance matrixes among populations. We next ran *TreeMix*

100 times without migration using different numbers of random seed and we built a ML

consensus tree from the 100 trees in *SumTrees* (Sukumaran & Holder 2010) using the 95%

majority rule. We finally added 1 and 4 migration edges to the ML trees and examined the

changes in the percentage of explained variance and the residual fit. For all f_3, f_4 and

TreeMix analyses, we used 4343 SNPs that were present in all populations and in at least

50% of individuals in every population.

Genomic clines

Loci of exceptional introgression compared with genome-wide average admixture may be

important for local adaptation or the maintenance of species barriers. To identify sites with

the greatest contribution to introgression between *An. nili* and *An. ovengensis*, we used the

Bayesian model of genomic clines as implemented in *bgc* (Gompert & Buerkle 2012). The

model first estimates a genome-wide average of a hybrid index (h) that varies between zero

and one for every potentially admixed individual, using allele frequencies of "pure" parental

populations (Gompert & Alex 2011). The value zero corresponds to pure individuals of the

alternative species (*An. ovengensis* in our case) and the value one to pure individuals of the

reference species (*An. nili* in our case). Two genomic clines parameters α and β are then

used to evaluate the deviation of individual loci in admixed individuals from the expected

genome-wide hybrid index (h). The cline parameter α reflects the probability of *An. nili*

ancestry relative to the base expectation (h), whereas the genomic cline parameter β

denotes the rate of transition from low to high probability of *An. ovengensis* ancestry as a

function of hybrid index (Gompert & Alex 2011). Deviations of loci from the genome-wide

average of α and β were examined on the basis of departures from the 95% confidence

envelope.

The "pure" parental populations we used in genomic clines analyses were apparently

allopatric *An. nili* and *An. ovengensis* populations collected in Nyabessan and Nkoteng,

respectively (Fig. 1A). The two locations are separated by ~420 km, which presumably

reduces the rate of gene flow between *An. nili* and *An. ovengensis*. Hybrids were admixed

individuals collected from the sympatric area in Mbébé (Fig. 1A). These individuals had

almost equivalent ancestry proportions from *An. nili* and *An. ovengensis* as revealed by both

STRUCTURE and ADMIXTURE clustering analyses. In *bgc*, we estimated genomic cline

parameters using 9622 SNPs deriving from aligned RAD-loci in a dataset consisting of 11

admixed individuals, 8 *An. nili* parents and 15 *An. ovengensis* parents. We calculated average

values of α and β across five runs of *bgc*, each run including 50000 steps with samples from

the posterior distribution recorded every 25th step following a 25000 step burn-in. We

visually inspected the MCMC output to assess convergence to the stationary distribution.

To examine the relationship between the strength of selection and introgression, we tested

whether loci with extreme or outlier genomic clines were enriched in genomic regions that

were targets of selection. To do so, we studied the correlation between locus-specific cline

parameters (α and β) and several divergence and diversity statistics (F_{ST}, d_{xy}, θ_w, θ_π,

Tajima's D) across 9622 SNPs in the genome. The strength of correlation was assessed

using Pearson's product moment correlation coefficient.

Results

SNP genotyping

We collected mosquitoes from four locations (Fig. 1A, Table S1) and sequenced 145

individuals belonging, according to morphological identifications and PCR, to two species

(*An. nili* (n = 24) and *An. ovengensis* (n = 121)). We aligned reads from all 145 individuals to

the reference genome and we assembled 197724 96-bp RAD loci that mapped to unique

positions throughout the genome. We retained 408 loci present in all populations and in at

least 50% of individuals in each population and identified 4343 high-quality biallelic

markers within these loci. We also identified another set of 704408 unique loci by building

consensus RAD loci *de novo* without aligning reads to the reference genome. We applied the

same stringent filtration as with aligned reads to identify 3071 high-quality SNPs.

Genetic structure of populations

PCA and NJ trees showed that the genotype variation at 4343 genome-wide SNPs among the

145 sequenced individuals is best explained by more than two clusters, implying cryptic

subdivisions among *An. nili* and *An. ovengensis* populations (Fig. 1C-D). The first three PCA

axes and NJ trees clearly distinguished three subgroups within *An. nili* and two clusters in

An. ovengensis. The five different clusters were associated with the sampling locations

suggesting a strong correlation between the genetic structure and a local distinctness of

populations. This marked geographic structure of *An. nili* and *An. ovengensis* populations

can be explained by the ongoing adaptive divergence and ecological speciation within the

two species. Importantly, both STRUCTURE and ADMIXTURE analyses revealed that, at k =

3, a subgroup containing 11 individuals corresponded to a cluster of hybrids with almost

half ancestry from *An. nili* and half from *An. ovengensis* (Fig. 1B). This result is surprising

given the substantial time since divergence. We applied three different methods to identify

the optimal number of genetic clusters in our samples. The DAPC suggested the presence of

five clusters as indicated by PCA and NJ trees (Fig. 1E). However, the method of Evanno et

al. indicated two probable ancestors while the distribution of the cross-validation error as a

function of the number of putative populations in ADMIXTURE failed to unambiguously

reveal an optimal number of genetic clusters (Fig. 1F,G). These conflicting results between

methods are sometimes observed when the history of subdivisions and admixtures events

is very complex as it is the case in *An. nili s.l* (Decker *et al.* 2014). The Evanno et al. method

is overwhelmed by early divergence between *An. nili* and *An. ovengensis* while the results of

DAPC and the ADMITURE cross-validation error reflect recent hierarchical population

subdivisions within the two species. The inferred genetic structure was consistent when we

used respectively 4343 and 3071 SNPs identified from reference-based and *de novo*

assemblies (Fig. 1 and Fig. S1).

Genetic differentiation

We estimated the level of population differentiation between the five genetic clusters

identified in our samples using the overall pairwise F_{ST} (Table 1). We found strong pairwise

differentiation characterized by extreme F_{ST} values, including between populations

classified as belonging to the same species. The level of differentiation is even higher

between some pairs of populations within the same species than between *An. nili* and *An.*

ovengensis. The highest level of within-species genetic differentiation was recorded

between "allopatric" populations of *An. ovengensis* collected from locations separated by

~<u>350</u>km (F_{ST} = 0.896, p < 0.005). These findings strongly suggest that, in addition to

hybridization, local differentiation associated with late stages of adaptive divergence within

species are overwhelming current taxonomic descriptions in *An. nili s.l.* populations. The

results obtained with reference alignments perfectly mirrored those of *de novo* assemblies

(Table 1, S2). Taken as a whole, the population genetic structure and divergence of *An. nili*

and *An. ovengensis* depicts a radiating group involving a collection of species whose

phylogenetic relationships are blurred by ongoing hybridizations. As a result, current

taxonomic classifications based on morphological characters and point mutations on the

ribosomal DNA cannot effectively describe the actual reproductive units and the vectorial

capacity within the *An. nili* species group.

Evidence for recent gene flow between *An. nili* and *An. ovengensis*

We used *TreeMix* to construct a tree connecting the five species and to effectively describe

and visualize the mixture(s) event(s). The consensus of 100 ML *TreeMix* tree inferred from

4343 SNPs without migration resumes population-level relationships described by

neighbor-joining analysis, PCA and clustering analyses (Fig. 2A). The long terminal

branches leading to *An. nili* group 1, *An. nili* group 2 and *An. ovengensis* group 1 in the

inferred tree reflect the signatures of strong bottlenecks in the history of these species.

Interestingly, the residual fit from the ML model without migration suggested some

correlations between species that were consistent with the admixture events inferred from

ADMIXTURE and STRUCTURE analyses, particularly between the two species *An. nili* group

2 and *An. ovengensis* group 2 (Fig. 1B). Additional support to admixture was provided by

the four-population test. Across all possible combinations of populations, there was a highly

334 significant allele frequency correlation between *An. nili* group 2 and *An. ovengensis* group 1

335 confirming that *An. nili* group 1 is the result of an admixture event between *An. nili* group 2

336 and *An. ovengensis* group 1 (Table 2 and S3). Nevertheless, five independent runs of *TreeMix*

337 including one or two migrations edges captured none of these migration events. Also, the

338 explained variance of the *TreeMix* ML model without migration edge was very high (99.8%)

339 indicating that a bifurcation tree can also match the phylogenetic relationships among

340 sampled populations. Finally, we found no significant f_3 statistics in all triplets of

341 populations, but this test may be underpowered because of the low-level and the precocity

342 of admixture or because of the complex demographic history of our populations (Decker *et*

343 *al.* 2014). As we showed previously with the population structure and genetic divergence,

344 the admixture events we described between *An. nili* and *An. ovengensis* also hold across

345 assembly methods (*de novo* or reference alignments)(Fig. 2 and S2).

346 **Genomic signatures of divergent selection and differential introgression**

347 We retained 9622 SNP loci identified from a reference-based assembly that we used to

348 examine the genomic footprints of divergent selection and introgression in a dataset

349 containing 8 "pure" *An. nili s.s.* individuals, 15 "pure" *An. ovengesis* and 11 admixed

350 individuals (see Materials and methods). The values of F_{ST} among SNP loci between parent

351 species *An. nili* and *An. ovengensis* were heterogeneous across the genome (Fig. 3). The

352 empirical distribution of locus-specific F_{ST} also revealed an unusually bimodal shape

353 featuring F_{ST} peaks centered on values around 0 and 1. The great majority of sites have low

354 to moderate differentiation, but a substantial number of loci are extremely differentiated

355 between the two species (Fig. 3A). We thinned our SNP set to 4003 using more stringent

filtration criteria and identified 42 statistical outliers of F_{ST} in LOSITAN, which were all

above the 99th percentile of the empirical distribution of F_{ST}. The bimodality of the genetic

divergence between *An. nili* and *An. ovengensis* is also evident in the empirical distribution

of d_{xy} (Fig. 3B). As exemplified by some empirical cases like the introgressive hybridization

observed between two closely related species of monkeyflowers (*Mimulus*) (Brandvain *et*

al. 2014), the bimodal divergence is due to the coexistence between sites of abrupt

differentiation reflecting the real level of divergence between the two species and sites of

low divergence resulting from recent gene flow. We conducted Linkage Disequilibrium (LD)

analyses to understand if highly differentiated sites are clustered into blocks of linked loci

or dispersed throughout the genome. Particularly, polymorphic chromosomal inversions

are important sources of genetic variation in *Anopheles* species and play a key role in local

adaptation and speciation. High F_{ST} values can aggregate in genomic regions containing

chromosomal inversions polymorphisms that are under strong divergent selection and

drive adaptive segregation as often observed in other African anopheline mosquitoes

(Ayala *et al.* 2011; Cheng *et al.* 2012; Kamdem *et al.* 2016). Cytogenetic studies have

detected no polymorphic chromosomal inversion among *An. ovengensis* samples. Two

polymorphic inversions (2*Rb*, 2*Rc*) have been identified in *An. nili*, but samples from

Cameroon bore only the 2*Rc* at a low frequency (Sharakhova *et al.* 2013). Rearranged

regions of the genome are characterized by reduced recombination and increased LD

relative to the genome-wide average. To effectively examine the extent of LD in the

genomes of *An. nili* and *An. ovengensis* and to test for the effect of chromosomal inversion

polymorphisms in adaptive evolution, we used the package LDna to identify clusters of LD.

To prevent spurious clusters due to LD between SNPs located on the same RAD locus, we

used a dataset in which only one SNP was randomly selected within each RAD locus (1330

SNPs in total). We optimized LDna clustering parameters, which indicated the presence of 6

LD blocks (Single Outlier Clusters (SOCs)) among our samples (Fig. 4). These clusters

represent signals of independent or compound events in the evolutionary history that left

imprints on LD across the genome (Kemppainen *et al.* 2015). To further understand the role

of the 6 LD clusters in the evolutionary history of *An. nili* and *An. ovengensis*, we conducted

downstream analysis using PCA to examine the population structure of SNPs within each

SOC. We found that one SOC (containing 39 SNPs) clearly discriminated a cluster

encompassing all hybrid individuals from the two parent species (Fig. 4B). Two other SOCs

(containing respectively 383 and 112 SNPs) consistently separated one parent species from

a group comprising the other parent species and all hybrid individuals. The last three SOCs

revealed no clear clustering patterns. None of the SOCs differentiated clusters that could be

associated with the three alternative karyotypes (inverted homozygotes, heterozygotes and

uninverted homozygotes) expected in case of polymorphic inversion. As suggested by

cytogenetic observations, polymorphic inversions likely play only a moderate role in the

genomic divergence in *An. nili s.l,* but a more intensive sampling is needed to make any

definitive conclusion. Interestingly, the three most important blocks of LD found in the

genomes of *An. nili* and *An. ovengensis* are instead associated with hybridization, which

emphasizes the central role played by interspecific gene flow in the genomic architecture of

adaptive speciation in the two species. Our results also suggest that loci resistant to

introgression and genes responsible for reproductive isolation between *An. nili* and *An.*

ovengensis are in strong linkage disequilibrium. This finding is consistent with recent

introgression whereby long-range haplotypes that are generated by recent gene flow

between genetically distinct populations have not have sufficient time to be broken down

by recombination (admixture-induced LD) (Martin *et al.* 2013). However, LDna estimates

the LD irrespective of the physical linkage between markers and LD blocks can also result

from a strong correlation between allelic frequencies of SNPs scattered throughout the

genome. A detailed characterization of long-range haplotypes across genomes of *An. nili* and

An. ovengensis will provide a more powerful examination of the genomic architecture of

reproductive isolation between the two species.

Here we have used another innovative approach (genomic cline) to further understand the

relationship between divergence, selection and differential introgression at the genomic

level. First, consistent with STRUCTURE and ADMIXTURE results, the empirical distribution

of hybrid index among admixed individuals (average hybrid index ~0.3) shows a slight

predominance of *An. ovengensis* ancestry (Fig. 5A, Fig. 1B). Using estimates of genomic cline

parameters, we noted that introgression was very heterogeneous across loci (Fig. 5B).

Indeed, the level of introgression at 1297 SNP loci differed significantly from the genome-

wide average (outliers) and we detected an excess *An. ovengensis* ancestry for 2506 loci

(lower bound of 95% CI for $\alpha > 0$) and excess *An. nili* ancestry for 2635 loci (upper bound of

95% CI for $\alpha < 0$). Estimates of the genomic cline parameter α range from -5 to 5.25 while

our values of β are low overall (β varies from -0.13 to 0.09). Moreover, we identified no loci

with significantly elevated estimates of genomic cline rate. The β parameter assesses the

rate of transition from one ancestry to the other and thereby scores the steepness of the

genomic cline at each locus. Extreme values of β are expected when there is population

structure in the hybrid zone, selection against hybrids or gene flow among parent species

(Parchman *et al.* 2013). Our hybrid population is highly differentiated ($F_{ST} > 0.8$) from both

parent species and is well adapted in hybrid zone, which certainly explain the low β values

observed. One key aspect in the Bayesian implementation of genomic clines is the

relationship between α and β, which is crucial to understand the rate of transition of sites of

exceptional introgression from one side to the other of the genome. As we have shown

previously with F_{ST} and d_{xy}, recent introgression in divergent genomes of *An. nili* and *An.*

ovengensis has resulted in bimodal genomic divergence featuring two blocks of sites with

extreme divergence values. This pattern presumes that most variant sites will have very

steep genomic clines because the transition from one ancestry to the other is very abrupt.

In agreement with this prediction, the scatterplot of β as a function of α indicates the

presence of two blocks of SNPs with either high probability of *An. nili* or of *An. ovengensis*

ancestry (Fig. 5B). The genomes of both species are split into two compartments of ancestry

due to recent introgression resulting in steep genomic clines at hybrid zone (Fig. 5B). To

better understand the biological significance of these outlier loci of introgression in

admixed individuals, we assessed the correlation between α and β and locus-specific

estimates of divergence and selection parameters. We found a detectable negative

correlation between F_{ST} and values of α (r = -0.15, p < 0.005). Among the 32 LOSITAN F_{ST}

outliers, 13 loci had extreme α estimates, and 11 were outlier of β, but the average values of

both α and β were not significantly different between F_{ST} outliers and the 9622 genome-

wide SNP loci (Mann-Whitney U-test, P < 0.001). Erroneous correlation between F_{ST} and

exceptional introgression can inferred when the F_{ST} between parental populations is < 0.1

(Parchman *et al.* 2013). The high overall F_{ST} (~0.8) between *An. nili* and *An. ovengensis*

minimizes such errors in our study. We also detected a negative correlation between α and

d_{xy} (r = -0.13, p < 0.005), which confirms previous results with F_{ST}, suggesting that sites that

diverge strongly resist to differential introgression because they likely contain reproductive

isolation factors. The negative correlation between d_{xy} and estimates of locus-specific

admixture indicates that perhaps gene flow has had sufficient time to reduce sequence

divergence at admixed loci despite the relatively recent hybridization process. We next

correlated cline parameters (α and β) to the estimates of locus-specific diversity (θ_w and θ_π

) and the allele frequency spectrum (Tajima's *D*). The results provided another clear

illustration of the steep genomic clines described previously with alpha and beta values.

Notably, there is a strong positive correlation between θ_w and α (r = 0.53, p< 0.005) in *An.*

ovengensis and a negative correlation of the same magnitude with the diversity of *An. nili* (r

= -0.49, p< 0.005), which translate the segregation of genomes of admixed individuals in

two sources of ancestry that are at opposite ends of the genomic cline. Interestingly, the

high correlation between alpha and diversity indicate that sites that are favored by

introgression are not constrained by divergent selection that should have likely resulted in

depression in nucleotide diversity. In contrast to what has been shown in manakin birds

(Parchman *et al.* 2013) and in *Lycaeides* butterflies (Gompert *et al.* 2012, 2013), there is no

evidence of increased introgression at loci under divergent selection between *An. nili* and

An. ovengensis.

Discussion

Hybridization at late stages of speciation

We have described a complex case of speciation and hybridization in a group of related *Anopheles* mosquitoes endemic to Sub-Saharan Africa. Both incomplete speciation and pervasive hybridization are leading to a very dynamic pattern of genetic structure between and within species. We first analysed the population genetic structure and revealed cryptic subdivisions between the two malaria vectors *An. nili* and *An. ovengensis*. The exact number of demes or cryptic subgroups within each species in our samples was difficult to determine. We found conflicting results between estimates from three different genetic clustering methods. However, three different methods suggested the existence of 2 to 5 clusters in our sample. *An. nili* is subdivided into three clusters while ongoing adaptive speciation in *An. ovengensis* results so far in two cryptic species. The extremely high genetic differentiation between populations indicates that all these cryptic subgroups are almost complete species. Our results therefore suggest that *An. nili* and *An. ovengensis,* contrary to the current taxonomy, represent probably two different complexes of cryptic species. The geographic origin of samples explains a great part of the genetic variance among individuals consistent with strong local differentiation and adaptive speciation. Significant population structure has been described among *An. nili* populations from Cameroon with 8 microsatellite loci (Ndo *et al.* 2013). Ndo *et al*. 2013 also found F_{ST} values as high as 0.48 between two populations from the forest area. Using genome-wide SNPs, we have identified new subdivisions and revealed that the population genetic structure of *An. nili* is more complex. Our work highlights the strength of Next Generation Sequencing (NGS)

approaches and the necessity of fine-scale genomic examinations in the resolution of

intricate patterns of ancestry in this group of mosquito. By contrast, the ongoing speciation

we have observed within *An. ovengensis* has never been described in the past, and this

species has been sometimes considered as a sibling of *An. nili* (Kengne *et al.* 2003; Awono-

Ambene *et al.* 2004, 2006; Ndo *et al.* 2013). Nevertheless, more recent studies have started

to challenge the assumed relatedness between the two species (Ndo *et al.* 2013; Sharakhova

et al. 2013). Precisely, analyses of polytene chromosomes revealed high karyotypic

divergence of *An. ovengensis* from *An. nili* (Sharakhova *et al.* 2013) and estimates of time

since divergence indicated that the two species split from one another 3 to 6-Myr ago (Ndo

et al. 2013). Intriguingly, our work provides multiple lines of evidence supporting the

existence of extensive ongoing gene flow between the two species despite this strong

divergence. First, clustering methods and Bayesian implementation of genomic clines

identified individuals with almost half ancestry from both species and formal tests of

population admixture corroborated these ongoing admixture events. Second, perhaps the

most compelling evidence for hybridization is the presence of admixture-induced linkage

disequilibrium (ALD) characterized by blocks of linked SNPs that discriminate hybrid

populations from parental species. In general, linkage disequilibrium (LD) or the

correlation between allele frequencies of different loci across the genome, which can have

multiple origins including selection, genetic drift, or population structure, is normally

eroded by recombination in the course of time. ALD is caused by associations between

nearby loci co-inherited on an intact chromosomal block from one of the ancestral mixing

populations (Loh *et al.* 2013). Signatures of ALD are frequent in genomes of recently

admixed populations for which recombination has not yet broken the large introgressed

chromosomal segments into smaller portions. ALD is a well-known feature in evolutionary

history of humans and estimates of long-range LD have been proposed as an approach to

measure the extent and the timing of admixture events that have shaped the genetic

polymorphism across genomes of extant populations (Loh *et al.* 2013). In insects,

signatures of ALD have been found for example in the *Heliconius* butterfly genome,

especially around loci involved in mimicry of color patterns that circulate between species

(Martin *et al.* 2013). *An. nili* provides another rare example of ALD across the genome of an

insect species. Further studies using a more comprehensive genomic sequencing and a

reference genome of better quality will help us to improve our knowledge of functional and

sequential characteristics of admixed LD blocks found in the *An. nili* genome.

Although the concept of speciation-with-gene-flow has become the dominant paradigm in

speciation studies, we remain ignorant about the conditions that prevent or motivate gene

flow between divergent lineages before the onset of complete reproductive isolation.

Moreover, even the notion of "complete reproductive isolation" is now challenged because

an increasing number of examples from diverse taxa showed rampant gene exchange across

strong reproductive barriers, sometimes between established species that diverged several

million years ago (Nydam & Harrison 2011; Roux *et al.* 2013; Parchman *et al.* 2013; Martin

et al. 2013; Canestrelli *et al.* 2014). These studies and ours suggest that processes

underlying hybridization and introgression in the presence of clear genetic differentiation

will be best addressed within speciation continuum rather than across couples of

occasionally mating species. A continuum of speciation featuring a collection of taxa

occupying a gradient of genetic/ecological divergence provides ideal conditions where the

relation between divergence and hybridization can be inferred. Contrary to what we

initially thought, the *An. nili* group does not represent a speciation continuum but instead a

collection of complete species sharing a common ancestry. Moreover, despite extensive

efforts, we couldn't sample populations of the two remaining species of the group: *An.*

somalicus and *An. carnevalei*. Therefore, patterns of admixtures and ancestry among

populations of this group of mosquitoes are probably more complex than what we have

shown. Nevertheless, instead of a comprehensive description of splits and admixtures in *An.*

nili s.l., we have focused our efforts on the examination of the genomic architecture of

divergence and introgression between *An. nili* and *An. ovengensis.* The results we discuss

below are among the rare cases that address the genomic signatures of gene flow and the

relationship between divergence, selection and introgression at late stages of the speciation

process.

Genomic architecture of adaptive introgression

Owing to the increasing availability of high-throughput sequencing information, genomes of

multiple taxa have been scanned and analyzed in comparative frameworks to search for

genomic signatures of divergence and speciation. The prevailing idea behind these

approaches is that regions of extreme differentiation between incipient or complete species

contain factors that maintain reproductive isolation (RI) among their populations. In

reality, the genomic distribution of highly differentiated regions has been more contentious

(Nosil & Feder 2012, 2013). Overall, two models are well documented: the "speciation

island" model whereby RI loci are thought to be caught up in in a few regions of the genome

where outliers of genetic differentiation cluster (e.g. (Andrew & Rieseberg 2013)) and the

"heterogeneous" distribution model, which posits that genomic divergence is instead lead

by numerous small genomic regions scattered throughout all chromosomes (e.g.

(Lawniczak *et al.* 2010; Roesti *et al.* 2012)). A shift in this paradigm is envisioned as

evidence is accumulating indicating that highly differentiated loci can paradoxically

coincide with regions of elevated introgression between two species (Gompert *et al.* 2012,

2013; Pool *et al.* 2012; Parchman *et al.* 2013). The genomic distribution of the genetic

divergence between *An. ovengensis* and *An. nili* shows that the compound effects of strong

divergence and recent introgression generates a bimodal pattern of divergence which

assigns most sites into two main categories: a majority of low divergence sites and a small

cluster of high divergence loci with F_{ST} values centered around 1. In most of the widespread

Anopheles species, signatures of high divergence can be found in large chromosomal

segments corresponding to rearranged regions of chromosomes where recombination is

rare (Neafsey *et al.* 2015). This is the case for example in the 2L*a* inversion locus, which

depicts signatures of strong divergent selection along a latitudinal cline (Cheng *et al.* 2012).

In agreement with cytogenetic studies, which found roughly no polymorphic inversions

among *An. nili* samples from Cameroon, we observed no tendency for high F_{ST} or d_{xy} to

cluster within regions that can be assimilated to chromosomal rearrangements. In contrast

to what has been recently demonstrated in manakin birds (Parchman *et al.* 2013) and in

Lycaeides butterflies (Gompert *et al.* 2012, 2013), there is a negative correlation between

the introgression parameter α and genetic divergence (estimated both as F_{ST} an d_{xy})

between *An. nili* and *An. ovengensis*. As expected, gene flow is favored across neutral loci

and those that provide selective advantage, but are presumably not under strong divergent

selection. Moreover, the relationship between α and β suggests a prevalence of steep clines

in genomes of admixed individuals at contact zones. This also translates into a positive

correlation between α and the nucleotide diversity in *An. ovengensis* and a negative

correlation of α to the diversity of *An. nili*. In theory, SNPs with steep cline are hypothesized

to be near genes involved in reproductive isolation and as such are possibly under selection

in hybrids (Janoušek *et al.* 2015). Therefore, the abundance of steep clines provides another

clear illustration of the mosaic genome characterized by coexistence between high

divergence and consistent gene flow observed in the hybrid species.

A substantial body of evidence indicates that genomic material coming from related species

can confer an advantage to populations (adaptive introgression). Adaptive introgression of

one or two loci has been widely studied over the last two decades and excellent examples

have been described. The most prominent cases include the transfer of genes involved in

mimicry of color patterns in *Heliconius* butterflies (Consortium 2012), the circulation of

resistant alleles of insecticides in mosquitoes (Clarkson *et al.* 2014; Norris *et al.* 2015) and

rodenticide in mice (Song *et al.* 2011). However, although convincing signatures of adaptive

introgression around one gene or a few linked loci can now be described in an increasing

number of species, the extent and the magnitude of introgressive hybridization across the

genome remained unknown. Further, due in part to the fact that most species in which the

concomitance of introgressive hybridization and high divergence has been observed are

nonmodel species with little genomic resources, the knowledge of genomic characteristics

of adaptive introgression in these species remains relatively modest. Meanwhile, a

consistent pattern has started to emerge from the few cases that have been studies with

substantial genomic details. In general, genomic regions exhibiting non-random

introgression are widely dispersed across the genome, rather than co-localized in a few

discrete blocks. This heterogeneity of genome-wide introgression patterns has been

observed for example between manakin birds (*Manacus candei* and *M. vitellinus*)

(Parchman *et al.* 2013) in different mice subspecies (Liu *et al.* 2015), and among

mosquitoes of the *Anopheles gambiae* species complex (Fontaine *et al.* 2014; Kamdem *et al.*

2016). Our results also show that selective introgression can be widespread across the

genome of two highly divergent species. However, in contrast to most of the reference cited,

LD analyses in *An. nili* and *An. ovengensis* have revealed a LD cluster separating hybrids

from the two parental species, which suggests that at least some of the recently

introgression loci consist of relatively large chromosome segments that have yet to be

further characterized sequentially and functionally with a high-quality reference genome.

In addition to the lack of knowledge about the genomic architecture, the functional and

phenotypic aspects of introgressive hybridization between established species remain

obscures. In mice for example, introgression is associated with a polarization of GO terms,

regions of elevated introgression exhibiting a disproportionate number of genes involved in

signal transduction and olfactory receptor genes (Janoušek *et al.* 2015). Hybrids between

An. nili and *An. ovengensis* were collected from an area of the equatorial forest whose

environmental features were not apparently very divergent from those of the locations

where the parent species were sampled (Fig. 1A). As a result, it is hard to pinpoint

environmental gradients and the life history traits that can be considered as the main

drivers of speciation and introgression among species of *An. nili* and *An. ovengensis*.

Conclusions and implications

Although hybridization has been recognized as one of the major forces that affect the

evolution of living species, the detailed study of its fundamental and applied implications

has been hampered by methodological limitations. Advances in high-throughput DNA

sequencing and statistical genomics are revolutionizing experimental and conceptual

approaches, allowing a very sensitive examination of the heterogeneity of hybridization

across species and genomes. We have used a combination of tests tailored to infer patterns

of introgression across genomes of nonmodel species. Although our results still need to be

replicated in other contact zones across the distribution range of *An. nili* in Africa, they

highlight the complex relationships between divergence, selection and introgression during

the split of taxa. Our work has methodological, conceptual and applied implications. Most

genome scans assume a negative correlation between genetic divergence and introgression.

It has been suggested that that the opposite is possible (Parchman *et al.* 2013; Gompert *et*

al. 2013), but our findings do no support this hypothesis. Climate change and anthropogenic

disturbance are contributing to expand geographic ranges of mosquito species worldwide

thereby increasing contact between previously isolated species that are capable of

exchanging gene flow. This interspecific gene flow in mosquitoes often leads to the spread

of insecticide resistance alleles and other epidemiologically significant genes. Our work

provided a methodological validation of a cost-effective population genomic approach that

can be applied to investigate the bases of introgressive hybridization in other mosquito

species.

Acknowledgements

Funding for this project was provided by the University of California Riverside and NIH
grants 1R01AI113248 and 1R21AI115271 to BJW.

References

Abbott R, Albach D, Ansell S *et al.* (2013) Hybridization and speciation. *Journal of Evolutionary Biology*, **26**, 229–246.

Alexander DH, Novembre J, Lange K (2009) Fast model-based estimation of ancestry in unrelated individuals. *Genome Research*, **19**, 1655–1664.

Andrew RL, Rieseberg LH (2013) Divergence is focused on few genomic regions early in speciation: incipient speciation of sunflower ecotypes. *Evolution; international journal of organic evolution*, **67**, 2468–82.

Antonio-Nkondjio C, Kerah CH, Simard F *et al.* (2006) Complexity of the malaria vectorial system in Cameroon: contribution of secondary vectors to malaria transmission. *Journal of medical entomology*, **43**, 1215–1221.

Antonio-Nkondjio C, Ndo C, Costantini C *et al.* (2009) Distribution and larval habitat characterization of Anopheles moucheti, Anopheles nili, and other malaria vectors in river networks of southern Cameroon. *Acta Tropica*, **112**, 270–276.

Awono-Ambene HP, Kengne P, Simard F, Antonio-Nkondjio C, Fontenille D (2004) Description and bionomics of Anopheles (Cellia) ovengensis (Diptera: Culicidae), a new malaria vector species of the Anopheles nili group from south Cameroon. *Journal of medical entomology*, **41**, 561–568.

Awono-Ambene HP, Simard F, Antonio-Nkondjio C *et al.* (2006) Multilocus enzyme electrophoresis supports speciation within the Anopheles nili group of malaria vectors in Cameroon. *The American journal of tropical medicine and hygiene*, **75**, 656–658.

Ayala D, Fontaine MC, Cohuet A *et al.* (2011) Chromosomal inversions, natural selection and adaptation in the malaria vector Anopheles funestus. *Molecular biology and evolution*, **28**, 745–758.

Beaumont MA, Nichols RA (1996) Evaluating loci for use in the genetic analysis of population structure. *Proceedings of the Royal Society B: Biological Sciences*, **263**, 1619–1626.

Brandvain Y, Kenney AM, Flagel L, Coop G, Sweigart AL (2014) Speciation and Introgression between Mimulus nasutus and Mimulus guttatus. *PLoS Genetics*, **10**.

Canestrelli D, Bisconti R, Nascetti G (2014) Extensive unidirectional introgression between two salamander lineages of ancient divergence and its evolutionary implications. *Scientific reports*, **4**, 6516.

Cheng C, White BJ, Kamdem C *et al.* (2012) Ecological genomics of Anopheles gambiae along a latitudinal cline: a population-resequencing approach. *Genetics*, **190**, 1417–32.

Clarkson CS, Weetman D, Essandoh J *et al.* (2014) Adaptive introgression between Anopheles sibling species eliminates a major genomic island but not reproductive isolation. *Nature communications*, **5**, 4248.

Consortium THG (2012) Butterfly genome reveals promiscuous exchange of mimicry adaptations among species. *Nature*, **487**, 94–8.

Decker JE, McKay SD, Rolf MM *et al.* (2014) Worldwide Patterns of Ancestry, Divergence, and Admixture in Domesticated Cattle. *PLoS Genetics*, **10**.

Earl DA, VonHoldt BM (2012) STRUCTURE HARVESTER: a website and program for visualizing STRUCTURE output and implementing the Evanno method. *Conservation*

692  *Genetics Resources*, **4**.

693 Ellegren H, Smeds L, Burri R *et al.* (2012) The genomic landscape of species divergence in
694  Ficedula flycatchers. *Nature*, **491**, 756–60.

695 Evanno G, Goudet J, Regnaut S (2005) Detecting the number of clusters of individuals using
696  the software structure: a simulation study. *Molecular Ecology*, **14**, 2611–2620.

697 Fontaine MC, Pease JB, Steele A *et al.* (2014) Extensive introgression in a malaria vector
698  species complex revealed by phylogenomics. *Science*, **347**, DOI:
699  10.1126/science.1258524.

700 Fumagalli M, Vieira FG, Linderoth T (2014) ngsTools : methods for population genetics
701  analyses from Next-Generation Sequencing data. *Bioinformatics*.

702 Gillies MT, Coetzee M (1987) *A supplement to the Anophelinae of Africa south of the Sahara*.
703  The South African Institute for Medical Research, Johannesburg.

704 Gillies MT, De Meillon B (1968) *The Anophelinae of Africa South of the Sahara*. Publications
705  of the South African Institute for Medical Research, Johannesburg.

706 Gompert Z, Alex BC (2011) Bayesian estimation of genomic clines. *Molecular ecology*, **20**,
707  2111–2127.

708 Gompert Z, Buerkle C a. (2012) <tt>bgc</tt> : Software for Bayesian estimation of genomic
709  clines. *Molecular Ecology Resources*, **12**, 1168–1176.

710 Gompert Z, Lucas L, Nice C (2012) Genomic regions with a history of divergent selection
711  affect fitness of hybrids between two butterfly species. …, 2167–2181.

712 Gompert Z, Lucas LK, Nice CC, Buerkle CA (2013) Genome divergence and the genetic
713  architecture of barriers to gene flow between Lycaeides idas and L. Melissa. *Evolution;*
714  *international journal of organic evolution*, **67**, 2498–514.

715 Grant BR (2015) Introgressive hybridization and natural selection in Darwin's finches.
716  *Biological Journal of the Linnean SocietyBiolog*, 812–822.

717 Hohenlohe P a, Bassham S, Etter PD *et al.* (2010) Population genomics of parallel adaptation
718  in threespine stickleback using sequenced RAD tags. *PLoS genetics*, **6**, e1000862.

719 Jakobsson M, Rosenberg N (2007) CLUMPP: a cluster matching and permutation program
720  for dealing with multimodality in analysis of population structure. *Bioinformatics*, **23**,
721  1801– 1806.

722 James M (2007) Hybrid speciation. *Nature*, **446**, 279–283.

723 Janoušek V, Munclinger P, Wang L, Teeter KC, Tucker PK (2015) Functional organization of
724  the genome may shape the species boundary in the house mouse. *Molecular Biology*
725  *and Evolution*, **32**, 1208–1220.

726 Jombart T (2008) adegenet: a R package for the multivariate analysis of genetic markers.
727  *Bioinformatics*, **24**, 1403–1405.

728 Kamdem C, Fouet C, Gamez S, White BJ (2016) Pollutants and insecticides drive local
729  adaptation in African malaria mosquitoes. *bioRxiv*.

730 Kemppainen P, Knight CG, Sarma DK *et al.* (2015) Linkage disequilibrium network analysis
731  (LDna) gives a global view of chromosomal inversions, local adaptation and geographic
732  structure. *Molecular Ecology Resources*, n/a–n/a.

733 Kengne P, Awono-Ambene, H. P. Antonio-Nkondjio C, Simard F, Fontenille D (2003)
734  Molecular identification of the Anopheles nili group African malaria vectors. *Med Vet*
735  *Entomol*, **17**, 67–74.

Korneliussen T, Albrechtsen A, Nielsen R (2014) ANGSD: Analysis of Next Generation
Sequencing Data. *BMC Bioinformatics*, **15**, 356.
Lawniczak MKN, Emrich SJ, Holloway a K *et al.* (2010) Widespread divergence between
incipient Anopheles gambiae species revealed by whole genome sequences. *Science*
(New York, N.Y.), **330**, 512–4.
Liu KJ, Steinberg E, Yozzo A *et al.* (2015) Interspecific introgressive origin of genomic
diversity in the house mouse. *Proceedings of the National Academy of Sciences*, **112**,
196–201.
Loh PR, Lipson M, Patterson N *et al.* (2013) Inferring admixture histories of human
populations using linkage disequilibrium. *Genetics*, **193**, 1233–1254.
Mallet J, Besansky N, Hahn MW (2015) How reticulated are species? *BioEssays*, 140–149.
Martin SH, Dasmahapatra KK, Nadeau NJ *et al.* (2013) Genome-wide evidence for speciation
with gene flow in Heliconius butterflies. *Genome research*, **23**, 1817–28.
Meirmans P, Van Tienderen P (2004) GENOTYPE and GENODIVE: two programs for the
analysis of genetic diversity of asexual organisms. *Molecular Ecology Notes*, **4**, 792–794.
Nadeau NJ, Martin SH, Kozak KM *et al.* (2013) Genome-wide patterns of divergence and
gene flow across a butterfly radiation. *Molecular ecology*, **22**, 814–26.
Ndo C, Antonio-Nkondjio C, Cohuet A *et al.* (2010) Population genetic structure of the
malaria vector Anopheles nili in sub-Saharan Africa. *Malaria journal*, **9**, 161.
Ndo C, Simard F, Kengne P *et al.* (2013) Cryptic Genetic Diversity within the Anopheles nili
group of Malaria Vectors in the Equatorial Forest Area of Cameroon (Central Africa).
PLoS ONE, **8**, 1–12.
Neafsey DE, Waterhouse RM, Abai MR *et al.* (2015) Highly evolvable malaria vectors: The
genomes of 16 Anopheles mosquitoes. *Science*, **347**, 1258522–1258522.
Norris LC, Main BJ, Lee Y *et al.* (2015) Adaptive introgression in an African malaria
mosquito coincident with the increased usage of insecticide-treated bed nets.
Proceedings of the National Academy of Sciences, 201418892.
Nosil P (2012) *Ecological Speciation*.
Nosil P, Feder JL (2012) Widespread yet heterogeneous genomic divergence. *Molecular*
Ecology, **21**, 2829–2832.
Nosil P, Feder JL (2013) Genome evolution and speciation: toward quantitative descriptions
of pattern and process. *Evolution; international journal of organic evolution*, **67**, 2461–
7.
Nydam ML, Harrison RG (2011) Introgression Despite Substantial Divergence in a
Broadcast Spawning Marine Invertebrate. *Evolution*, **65**, 429–442.
Ozerov MY, Gross R, Bruneaux M *et al.* (2016) Genome-wide introgressive hybridization
patterns in wild Atlantic salmon influenced by inadvertent gene flow from hatchery
releases. *Molecular Ecology*, n/a–n/a.
Paradis E, Claude J, Strimmer K (2004) Analyses of Phylogenetics and Evolution in R
language. *Bioinformatics*, **20**, 289–290.
Parchman TL, Gompert Z, Braun MJ *et al.* (2013) The genomic consequences of adaptive
divergence and reproductive isolation between species of manakins. *Molecular ecology*,
22, 3304–17.
Payseur BA, Rieseberg LH (2016) A Genomic Perspective on Hybridization and Speciation.

780  *Molecular Ecology*, n/a–n/a.

781 Peery A, Sharakhova M V, Antonio-Nkondjio C *et al.* (2011) Improving the population
782  genetics toolbox for the study of the African malaria vector Anopheles nili:
783  microsatellite mapping to chromosomes. *Parasites & Vectors*, **4**, 202.

784 Peterson BK, Weber JN, Kay EH, Fisher HS, Hoekstra HE (2012) Double Digest RADseq: An
785  Inexpensive Method for De Novo SNP Discovery and Genotyping in Model and Non-
786  Model Species. *PLoS ONE*, **7**, e37135.

787 Pickrell JK, Pritchard JK (2012) Inference of Population Splits and Mixtures from Genome-
788  Wide Allele Frequency Data. *PLoS Genetics*, **8**.

789 Pool JE, Corbett-Detig RB, Sugino RP *et al.* (2012) Population Genomics of Sub-Saharan
790  Drosophila melanogaster: African Diversity and Non-African Admixture. *PLoS Genetics*,
791  **8**.

792 Pritchard JK, Stephens M, Donnelly P (2000) Inference of population structure using
793  multilocus genotype data. *Genetics*, **155**, 945–959.

794 Purcell S, Neale B, Todd-Brown K *et al.* (2007) PLINK: a toolset for whole-genome
795  association and population-based linkage analysis. *American Journal of Human*
796  *Genetics*, **81**.

797 Reich D, Thangaraj K, Patterson N, Price AL, Singh L (2009) Reconstructing Indian
798  population history. *Nature*, **461**, 489–494.

799 Roesti M, Hendry AP, Salzburger W, Berner D (2012) Genome divergence during
800  evolutionary diversification as revealed in replicate lake-stream stickleback population
801  pairs. *Molecular ecology*, **21**, 2852–62.

802 Rosenberg N (2004) DISTRUCT: a program for the graphical display of population
803  structure. *Molecular Ecology Resources*, **4**, 137–138.

804 Roux C, Tsagkogeorga G, Bierne N, Galtier N (2013) Crossing the species barrier: Genomic
805  hotspots of introgression between two highly divergent ciona intestinalis species.
806  *Molecular Biology and Evolution*, **30**, 1574–1587.

807 Sharakhova M V., Peery A, Antonio-Nkondjio C *et al.* (2013) Cytogenetic analysis of
808  Anopheles ovengensis revealed high structural divergence of chromosomes in the
809  Anopheles nili group. *Infection, Genetics and Evolution*, **16**, 341–348.

810 Song Y, Endepols S, Klemann N *et al.* (2011) Adaptive introgression of anticoagulant rodent
811  poison resistance by hybridization between old world mice. *Current Biology*, **21**, 1296–
812  1301.

813 Sukumaran J, Holder MT (2010) DendroPy: A Python library for phylogenetic computing.
814  *Bioinformatics*, **26**, 1569–1571.

815 Turner TL, Hahn MW, Nuzhdin S V. (2005) Genomic islands of speciation in Anopheles
816  gambiae. *PLoS Biology*, **3**, 1572–1578.

817 Weir BS, Cockerham CC (1984) Estimating F-statistics for the analysis of population
818  structure. *Evolution*, **38**, 1358–1370.

819 Wu TD, Nacu S (2010) Fast and SNP-tolerant detection of complex variants and splicing in
820  short reads. *Bioinformatics*, **26**, 873–881.

822 **Author contributions**
823 Conceived and designed the experiments: CK CF BJW. Performed the experiments: CK CF SG
824 BJW. Analyzed the data: CK CF BJW. Wrote the paper: CK CF BJW.
825

826 **Tables**

828 **Table 1**: Pairwise F_{ST} between *An. nili* and *An. ovengensis* populations. p < 0.005 for all
829 values.

F_{ST}	*An. nili* group 1	*An. nili* group 2	*An. nili* group 3	*An. ovengensis* group 1	*An. ovengensis* group 2
An. nili group 1	-				
An. nili group 2	0.794	-			
An. nili group 3	0.838	0.863	-		
An. ovengensis group 1	0.655	0.834	0.873	-	
An. ovengensis group 2	0.857	0.863	0.858	0.896	-

831 **Table 2**: Results of the most significant f_4 tests for gene flow.

Test	$f_4 \pm$ std err			Z-score	P-value
f_4(Oveng 1 ,Nili 3 ; Nili 2, Nili 1)	-0.03506	±	0.00287	-12.20	< 0.00001
f_4(Oveng 1, Nili 3 ; Nili 2, Oveng 2)	0.02241	±	0.00263	8.52	< 0.00001
f_4(Oveng 1, Oveng 2 ; Nili 3, Nili 1)	-0.05775	±	0.00355	-16.25	< 0.00001
f_4(Oveng 1, Oveng 2 ; Nili 2, Nili 1)	-0.03718	±	0.00292	-12.72	< 0.00001
f_4(Nili 3, Nili 1 ; Nili 2, Oveng 2)	-0.02272	±	0.00263	-8.63	< 0.00001

Oveng 1: *An. ovengensis* group 1 ; Oveng 2: *An. ovengensis* group 2 ; Nili 1: *An. nili* group 1 ; Nili 2: *An. nili* group 2 ; Nili 3: *An. nili* group 3

834   **Table 3**: Pearson's correlation coefficient assessing the relationship between cline
835   parameter (α and β) and locus-specific estimates of five population genomic parameters
836   (pairwise genetic divergence (F_{ST} and d_{xy}), nucleotide diversity (θ_w and θ_π) and allele
837   frequency spectrum (Tajima's D)).
838

	Hybrids						An. nili			An. ovengensis			
	F_{ST}	d_{xy}	θw	$\theta\pi$	TD		θw	$\theta\pi$	TD	θw	$\theta\pi$	TD	
α	-0.151	-0.130	0.057	0.052	0.016	*	-0.498	-0.414	0.088	0.537	0.432	-0.033	*
β	0.060	0.048	-0.016	* -0.017	* -0.011	*	0.230	0.188	-0.047	-0.266	-0.212	0.018	*

* not significant (p>0.005)

TD: Tajima's D

Figures

Figure 1: Population genetic structure of *An. nili sensu lato* inferred from 4343 SNPs
identified with a reference-based assembly. (A) Map showing the sampling locations and
relative frequencies of the two *An. nili* and *An. ovengensis*. (B) ADMIXTURE plots with k
from 2 trough 8. (C) and (D) neighbor-joining tree and PCA. Each PCA axis is labeled with
the percentage of variance explained. (E), (F) and (G) Identification of the optimal number
of genetic clusters using the delta k method of Evanno et al, DAPC, and 10-fold cross-
validation in ADMIXTURE. The lowest BIC and CV error and the highest delta k indicate the
most probable number of genetic clusters.

Figure 2. *TreeMix* Maximum Likelihood (ML) trees depicting the signals of gene flow
between *An. nili* and *An. ovengensis*. ML tree and residual fit from the ML model inferred
with (A) no migration edge, (B) a single migration edge and (C) two migration edges. The
small arrow on each indicates the directionality of gene flow migration edge and the color
of the edge reflect the intensity of admixture. Heat colors depict the residual covariance
between each pair of populations. Darker colors indicate populations more closely related
to each other than expected under a bifurcating maximum likelihood tree, suggestive of
gene flow.

Figure 3: Frequency distribution of F_{ST} (A) and d_{xy} (B) based on 9622 variant sit

Figure 4: Results of Linkage disequilibrium analyses in LDna. (A) LDna graph suggesting
the presence of 6 LD clusters (Single Outlier Clusters (SOCs)) based on 1330 SNPs in a
dataset containing *An. nili*, *An. ovengensis,* and hydrids of the two species. Values of the two
parameters: φ (which controls when clusters are defined as outliers) and $|E|_{min}$, the
minimum number of edges required for a LD cluster to be considered as an outlier, are
indicated on top of the graph. Corresponding LD thresholds are shown on the x-axis. (B)
Population genetic structure of the six SOCs identified.

Figure 5: Genomic cline analysis. (A) Frequency distribution of the hybrid index in admixed
individuals (hybrid index of pure *An. nili* = 1.0 and pure *An. ovengensis* = 0.0). (B)
Prevalence of steep genomic clines illustrated by a scatterplot between the cline
parameters α and β.

872
873

874   **Supplemental Material**
875
876   **Table S1:** Information on *An. nili sensu lato* mosquitoes included in this study.

Sampling locations	Geographic coordinates	Sampling methods			Total
		HLC-OUT	**HLC-IN**	**LC**	
Ebebda	4°20'00"N, 11°17'00"E			6	6
Nkoteng	4°31'00"N, 12°02'00"E			8	8
Nyabessan	2°24'00"N, 10°24'00"E	63	44		107
Mbébé	4°10'00"N, 11°04'00"E	13	3	8	24
Total		76	47	22	145

HLC-OUT, human landing catches performed outdoor; HLC-IN, human landing catches performed indoor; LC, larval collection

878   **Table S2**: Pairwise F_{ST} estimated from a *de novo* assembly. p < 0.005 for all values.

F_{ST}	*An. nili* group 1	*An. nili* group 2	*An. nili* group 3	*An. ovengensis* group 1	*An. ovengensis* group 2
An. nili group 1	-				
An. nili group 2	0.791	-			
An. nili group 3	0.844	0.862	-		
An. ovengensis group 1	0.705	0.861	0.902	-	
An. ovengensis group 2	0.854	0.862	0.867	0.907	-

880 **Table S3**: Results of the most significant f_4 tests for gene flow (*de no*vo assembly).

Test	$f_4 \pm$ std err			Z-score	P-value
f_4(Oveng 1 ,Nili 3 ; Nili 2, Nili 1)	-0.03692	±	0.00360	-10.25	<0.00001
f_4(Oveng 1, Nili 3 ; Nili 2, Oveng 2)	0.02787	±	0.00351	7.94	<0.00001
f_4(Oveng 1, Oveng 2 ; Nili 3, Nili 1)	-0.06610	±	0.00451	-14.64	<0.00001
f_4(Oveng 1, Oveng 2 ; Nili 2, Nili 1)	-0.04053	±	0.00368	-11.01	<0.00001
f_4(Nili 3, Nili 1 ; Nili 2, Oveng 2)	-0.02776	±	0.00349	-7.96	<0.00001

Oveng 1: *An. ovengensis* group 1 ; Oveng 2: *An. ovengensis* group 2 ; Nili 1: *An. nili* group 1 ; Nili 2: *An. nili* group 2 ; Nili 3: *An. nili* group 3

 Figure S1: Population genetic structure of *An. nili sensu lato* inferred from 3071 SNPs
882 identified with a *de novo* assembly. (A) ADMIXTURE plots with k from 2 trough 8. (B) and
883 (C) neighbor-joining tree and PCA. (D), (E) and (F) Identification of the optimal number of
884 genetic clusters using the delta k method of Evanno et al, DAPC, and a 10-fold cross-
885 validation in ADMIXTURE.

Figure S2: *TreeMix* Maximum Likelihood (ML) trees estimated from 3071 SNPs identified
with a *de novo* assembly. ML tree and residual fit from the ML model inferred with (A) no
migration, (B) a single migration and (C) two migration edges. See Fig. 2. in the main text
for additional description.