Extensive genetic diversity among populations of the malaria

mosquito *Anopheles moucheti* revealed by population genomics

Caroline Fouet[1]*, Colince Kamdem[1], Stephanie Gamez[1], Bradley J. White[1,2]*

[1]Department of Entomology, University of California, Riverside, CA 92521

[2]Center for Disease Vector Research, Institute for Integrative Genome Biology,

University of California, Riverside, CA 92521

*Corresponding authors

Address: Department of Entomology, University of California Riverside. 900

University Ave. Riverside, CA 92521. Tel: +1 951 827 2626

Email address: caroline.fouet@ucr.edu; bwhite@ucr.edu

Abstract

Malaria vectors are exposed to intense selective pressures due to large-scale intervention programs that are underway in most African countries. One of the current priorities is therefore to clearly assess the adaptive potential of Anopheline populations, which is critical to understand and anticipate the response mosquitoes can elicit against such adaptive challenges. The development of genomic resources that will empower robust examinations of evolutionary changes in all vectors including currently understudied species is an inevitable step toward this goal. Here we constructed double-digest Restriction Associated DNA (ddRAD) libraries and generated 6461 Single Nucleotide Polymorphisms (SNPs) that we used to explore the population structure and demographic history of wild-caught *Anopheles moucheti* from Cameroon. The genome-wide distribution of allelic frequencies among samples best fitted that of an old population at equilibrium, characterized by a weak genetic structure and extensive genetic diversity, presumably due to a large long term effective population size. Estimates of F_{ST} and Linkage Disequilibrium (LD) across SNPs reveal a very low genetic differentiation throughout the genome and the absence of segregating LD blocks among populations, suggesting an overall lack of local adaptation. Our study provides the first investigation of the genetic structure and diversity in *An. moucheti* at the genomic scale. We conclude that, despite a weak genetic structure, this species has the potential to challenge current vector control measures and other rapid anthropogenic and environmental changes thanks to its great genetic diversity.

38   **Key words:** *Anopheles moucheti*, population genomics, RADseq, *de novo* assembly

1. Introduction

Despite having a widely acknowledged epidemiological significance, most African malaria mosquitoes are so-called "neglected vectors" because the efforts devoted to their study and control are clearly insufficient. *Anopheles moucheti sensu lato* is one of the best examples. This mosquito vector is a group of three related species (*An. moucheti moucheti*, *An. moucheti nigeriensis*, and *An. moucheti bervoetsi*) distributed across the equatorial forest and distinguishable from each other by slight morphological differences (Kengne et al., 2007). The nominal species of the group, *An. moucheti moucheti* (hereafter *An. moucheti*), is a very efficient and anthropophilic vector especially in rural areas where the highest malaria burden due to *Plasmodium falciparum* infections are recorded (Antonio-Nkondjio et al., 2009, 2008, 2002). In such settings, abundant populations of *An. moucheti* breed year-round in slow moving streams and rivers and often outcompete other main malaria mosquitoes. Despite this epidemiological significance, the evolutionary history and the adaptive potential of this vector remain understudied. Early investigations of the genetic structure based on allozymes and microsatellites showed a significant genetic differentiation among samples from three different countries (Antonio-Nkondjio et al., 2008), but detected little divergence within populations from the same country (Antonio-Nkondjio et al., 2007, 2002). Precisely, very low levels of genetic differentiation were found between populations from Cameroon across eight microsatellite loci, suggesting extensive gene flow at such geographic scales, but detailed studies in other countries are still lacking to fully support this hypothesis. On the other hand, African anopheline populations are

increasingly exposed to strong selective pressures associated with insecticide-based

malaria control campaigns that have been recently intensified (World Health

Organization, 2013). Such pressures represent particularly efficient driving forces

that often contribute to the rapid diversification of vector populations in a few

decades (Clarkson et al., 2014; Kamdem et al., Unpublished data a; Norris et al.,

2015). As a result, a detailed characterization of the genomic architecture of all

vectors is important for a critical appraisal of the impacts of malaria control efforts.

In this framework, we set out to perform the first genome-wide investigation of

natural polymorphism in *An. moucheti*. One of our main goals was to know to what

extent assessing the genetic diversity could provide clues about the spatial

distribution and help predict the environmental resilience of this species. In

principle, evolutionary responses of species to human-induced or natural changes

rely largely on available heritable variation, which reflects the evolutionary

potential and adaptability to novel environments (Orr and Unckless, 2008).

Therefore, the screening of genome-wide variation is supposed to be a sensible

approach that may provide a generalized measure of evolutionary potential in

species like *An. moucheti* for which direct ecological, evolutionary or functional tests

are impossible (Harrisson et al., 2014).

Thanks to recent progresses in sequencing technology, high-resolution sequence

information can be generated for virtually any living organism. These technological

advances are extraordinary helpful for non-model species with limited genomic

resources like mosquitoes (Ellegren, 2014). However, at the exception of *Anopheles*

gambiae for which significant genomic studies have been carried out using high-

quality sequencing data (Fontaine et al., 2015; Kamdem et al., Unpublished data a;

O'Loughlin et al., 2014), the other African malaria vectors have yet to fully benefit

from the explosive growth of methods for assessing genetic variation at a fine scale.

These neglected vectors face a vicious cycle whereby the lack of basic genomic

resources that are critical to generate high-quality sequencing information and to

enable robust interpretations of natural polymorphisms greatly contributes to their

marginalization. One typical example is *An. moucheti*, which lacks all the vital

resources ranging from a laboratory strain, a reference genome assembly, and a

physical or linkage map.

To start filling this gap and to shed some light on the evolutionary history and

adaptive potential of this vector, we have performed a high-throughput sequencing

of reduced representation libraries in 98 wild-caught individuals from Cameroon

and identify thousands of RAD loci scattered throughout the genome. Using high-

quality Single Nucleotide Polymorphisms (SNPs) identified within these loci, we

have investigated the genetic structure of populations and scan genomes of our

samples to detect footprints of local adaptation and natural selection. We found that,

in our study zone, populations of *An. moucheti* are characterized by a great genetic

diversity and extensive gene flow. We argue that this vector is particularly adapted

to challenge the selective pressures imposed by vector controls and rapid

environmental modifications.

2. Material and methods

2.1. Mosquito sampling and sequencing

This study included two *An. moucheti* populations from the Cameroonian equatorial forest. A total of 98 mosquitoes (97 adults and 1 larva) were collected in August and November 2013 from Olama and Nyabessan, respectively (Table 1). The two locations are separated by ~200 km (Fig. 1A) and are crossed respectively by the Nyong and the Ntem rivers that provide the breeding sites for *An. moucheti* larvae. Specimens were identified as *An. moucheti moucheti* using morphological identification keys (Gillies and Coetzee, 1987; Gillies and De Meillon, 1968) and a diagnostic PCR, which targets mutations on the ribosomal DNA (Kengne et al., 2007). We extracted genomic DNA using the DNeasy Blood and Tissue kit (Qiagen) for larvae and the Zymo Research MinPrep kit for adult mosquitoes. We used 10µl (~50ng) of genomic DNA to prepare double-digest Restriction-site Associated DNA libraries following a modified protocol of Peterson et al., 2012. *MluC1* and *NlaIII* restriction enzymes were used to digest DNA of individual mosquitoes, yielding RAD-tags of different sizes to which short unique DNA sequences (barcodes and adaptors) were ligated to enable the identification of reads belonging to each specimen. The digestion products were purified and pooled. DNA fragments of around 400bp were selected and amplified via PCR. The distribution of fragment sizes was checked on a BioAnalyzer (Agilent Technologies, Inc., USA) before sequencing. The sequencing was performed on an Illumina HiSeq2000 platform (Illumina Inc., USA) (Genomic Core Facility, University of California, Riverside) to yield single-end reads of 101bp.

2.2 SNP discovery and genotyping

We used the bioinformatics pipeline Stacks v1.35 (Catchen et al., 2013) to process

Illumina short reads. The program *process_radtags* was first used to sort the reads

according to the barcodes and to trim all reads to 96bp in length by removing index

and barcode sequences from the ends of the reads. Reads with ambiguous barcodes,

those that did not contain the *NlaIII* recognition site and those with low-quality

scores (average Phred score < 33) were excluded. The program *ustacks* was then

utilized to perform a *de novo* assembly (i.e., the assembly of reads in "stacks"

enabling the creation of consensus RAD loci without prior alignment onto a

reference genome sequence) (Catchen et al., 2013, 2011) in each individual in our

populations. We allowed a maximum of 2 nucleotide mismatches between stacks (M

parameter in *ustacks*) and we required a minimum of three reads to create a stack

(m parameter in *ustacks*). Using the *cstacks* program, a catalogue of loci was built to

synchronize variations across all individuals in our populations. Finally, we utilized

the *populations* program to calculate population genetic parameters and output

SNPs in different formats. To avoid bias associated with less informative SNPs or

possible false positive SNPs (due to sequencing or pipeline errors), only RAD loci

scored in at least 70-75% of individuals were retained for further analyses.

2.3. Population genomic analyses

SNP files outputted by the *populations* program were used to assess the population

genetic structure with a Principal Component Analysis (PCA) and a Neighbor-Joining

(NJ) tree analysis using respectively the R packages *adegenet* and *ape* (Jombart,

2008; Paradis et al., 2004; R Development Core Team, 2008). We also explored

patterns of ancestry and admixture among individuals in ADMIXTURE v1.23

(Alexander et al., 2009) with 10-fold cross-validation for k assumed ancestral

populations (k= 1 through 6). The optimal number of clusters was confirmed using

the Discriminant Analysis of Principal Component (DAPC) method, which explores

the number of genetically distinct groups by running a k-means clustering

sequentially with increasing numbers and by comparing different clustering

solutions using Bayesian Information Criterion (BIC) (Jombart, 2008). We examined

the population genetic diversity, conformity to Hardy-Weinberg equilibrium and

demographic background using several statistics calculated with the *populations*

program. Precisely, to assess the global genetic diversity per population, we

calculated the overall nucleotide diversity (π) and the frequency of polymorphic

sites within population. To make inferences on the demographic history and to test

for departures from Hardy–Weinberg equilibrium, we used the allele frequency

spectrum and the Wright's inbreeding coefficient (F_{IS}). To quantify the geographic

and genetic differentiation between allopatric populations, we estimated the

genome-wide average F_{ST} (Weir and Cockerham, 1984) on 2000 randomly selected

SNPs in Genodive v1.06 (Meirmans and Van Tienderen, 2004). We also conducted

an hierarchical Analysis of Molecular Variance (AMOVA) (Excoffier et al., 1992) on

the same SNP set to quantify the effects of the geographic origin on the genetic

variance among individuals. The statistical significance of F_{ST} and AMOVA was

assessed with 10000 permutations. Finally, to have a detailed picture of the genomic

architecture of divergence, we inspected the genome-wide distribution of locus-

specific estimates of F_{ST}.

2.4. Identification of segregating polymorphic chromosomal inversions

In structured *Anopheles* populations whose ecological/genetic divergence is due to polymorphic chromosomal inversions, high values of F_{ST} are expected between divergent populations within inversion loci, a pattern consistent with local adaptation of alternative karyotypes (Ayala and Coluzzi, 2005). This is the case for most populations of the main African malaria vectors *An. funestus* and *An. gambiae*, which depict multiple inversion clines in nature (Ayala et al., 2011; Fouet et al., Unpublished data; Kamdem et al., Unpublished data a; O'Loughlin et al., 2014). In addition to scanning genomes of our individuals to identify outlier values of F_{ST} that are indicators of selection and local adaptation, we also used Linkage Disequilibrium (LD) analysis to search for the presence of LD blocks corresponding to putative inversion polymorphisms. LD (the nonrandom association of alleles at different loci) provides information about past events and is affected by local adaptation and geographical structure, the demographic history, or the magnitude of selection and recombination across the genome (Lewontin and Kojima, 1960). Notably, high LD is expected in regions bearing inversions relative to the rest of the genome because the neutral recombination rate is notoriously reduced within inversions (Kirkpatrick and Barton, 2006). Thus, assessing genome-wide patterns of LD can reveal clusters of strongly correlated SNPs (LD blocks) corresponding potentially to chromosomal inversions. The R package LDna (Kemppainen et al., 2015) allows the examination of the distinct LD network clusters within the genome of non-model species without the need of a linkage map or reference genome. We have calculated LD, estimated as the r^2 correlation coefficient between all pairs of SNPs, in PLINK

200   v1.09 (Purcell et al., 2007). To avoid spurious LD due to the strong correlation

201   between SNPs located on the same RAD locus, we randomly selected only one SNP

202   within each RAD locus resulting in a dataset of 2569 variants containing less than

203   15% missing data. LDna was then used to identify LD blocks whose population

204   genetic structure was examined with a PCA.

3. Results

3.1. *De novo* assembly

In total, 518,218 unique 96-bp RAD loci were identified from *de novo* assembly of reads in 98 individuals. We retained 946 loci that were present in all sampled populations and in at least 75% of individuals in every population, and we identified 3027 high-quality biallelic SNPs from these loci.

3.2. Population genetic structure

First, we tested for the presence of cryptic genetic subdivision within *An. moucheti* with PCA, NJ trees and the ADMIXTURE ancestry model. A NJ tree constructed from a matrix of Euclidian distance using allele frequencies at 3027 genome-wide SNPs showed a putative subdivision of *An. moucheti* populations in two genetic clusters (Fig. S1A). The first three axes of PCA also revealed a number of outlier individuals separated from a main cluster (Fig. S1B). However, when we ranked our sequenced individuals based on the number of sequencing reads, we noticed that one of the putative genetic clusters corresponded to a group of individuals having the lowest sequencing coverage (Fig. S1 and Table S1). We excluded all these individuals and reduced our dataset to 78 individuals. We conducted a new *de novo* assembly and analyzed the relationship between the 78 remaining individuals at 6461 SNPs present in at least 70% of individuals using PCA, NJ trees and ADMIXTURE. Both the k-means clustering (DAPC) and the variation of the cross-validation error as a function of the number of ancestral populations in ADMIXTURE revealed that the polymorphism of *An. moucheti* resulted from only one ancestral population (k = 1) (Fig. 1B and 1C). PCA and NJ depicted a homogeneous cluster comprising all 78

individuals providing additional evidence of the lack of genetic or geographic

structuring among populations (Fig. 1D and 1E). Unsurprisingly, the overall F_{ST} was

remarkably low between populations from the two sampling locations Olama and

Nyabessan (F_{ST} = 0.008, p < 0.005). Similarly, the distribution of F_{ST} values across

6461 SNPs showed a large dominance of very low F_{ST} values throughout the genome

(Fig. 2). The highest per locus F_{ST} was only 0.126, while 5006 of the 6461 loci

revealed F_{ST} near zero. The modest geographic differentiation was also well

illustrated by a hierarchical AMOVA, which showed that the genetic variance was

explained essentially by within-individual variations (99.7%). Finally, we found very

low overall Wright's inbreeding coefficient (F_{IS}= 0.0014, p < 0.005 in Nyabessan and

F_{IS} = 0.0025, p < 0.005 in Olama) (Table 2) suggesting that allelic frequencies within

both populations were in accordance with proportions expected under the Hardy-

Weinberg equilibrium.

3.3. Genetic diversity and demographic history

The estimates of the overall nucleotide diversity (π = 0.0020 and π = 0.0016,

respectively, in Olama and Nyabessan) (Table 2) were within the range of average

values found in other African *Anopheles* species using RADseq approaches (Fouet et

al., Unpublished data; Kamdem et al., Unpublished data (a, b); O'Loughlin et al.,

2014). Notorious demographic expansions have been described in natural

populations of this insect clade (Donnelly et al., 2001), and the values of π observed

in *An. moucheti* likely reflect the level of genetic diversity of a population with large

effective size. The great genetic diversity of *An. moucheti* was also illustrated by the

percentage of polymorphic sites. Of the 6461 variant sites, 89.60% were

polymorphic in Olama and 34.82% in Nyabessan (Table 2). The difference observed

between the two locations can be related to the sample size (n = 19 in Nyabessan

and n = 59 in Olama) or to demographic particularities that persists between the

two geographic sites despite a massive gene flow. To infer the demographic history

of *An. moucheti*, we examined the Allele Frequency Spectrum (AFS), summarized as

the distribution of the major allele in one population. This approach was a surrogate

to model-based methods that provide powerful examinations of the history of

genetic diversity by modeling the AFS at genome-wide SNP variants, but that

couldn't be implemented here due to the lack of a reference genome assembly. The

frequency distribution of the major allele p (Fig. 3) indicates that the majority of

polymorphic loci are highly frequent in Olama and Nyabessan as shown by the

predominance of SNPs at frequencies equal to 1. Ranges of allele frequencies are

also similar in both locations (between 0.47 and 1 in Olama and between 0.34 and 1

in Nyabessan). These frequency ranges are expected for old populations at

equilibrium capable of accumulating high amount of genetic diversity.

3.4. Polymorphic chromosomal inversions and local adaptation

When paracentric inversions are involved in local adaptation, high values of genetic

divergence are often observed within inversion loci in natural populations.

Cytogenetic analyses of the polytene chromosome of *An. moucheti* have identified

three polymorphic chromosomal inversions within samples collected from the sites

we have studied (Sharakhova et al., 2014). However, the weak overall population

structure and the very low F_{ST} values we have detected throughout the genome are

clear indicators of the absence of local adaptation. Interestingly, this finding also

suggests that none of the polymorphic inversions described previously is actually

segregating among our samples, as high values of F_{ST} are absent even within

inversion loci. We provided further support to this hypothesis by performing LD

analyses. First, we found a globally low LD in the *An. moucheti* genome (average

genome-wide r^2 = 0.0149) as expected in highly polymorphic populations with large

effective size. We next used LDna to cluster the LD values and to identify Single

Outlier Clusters (SOC) that can be associated with distinct or multiple evolutionary

phenomena in the *An. moucheti* history. We set the parameters to collect and screen

a high number of SOCs using 2569 highly filtered SNPs, which allowed us to identify

20 independent LD blocks in our samples (Fig. 4). In principle, when these blocks

are associated with important events in the evolutionary history of a species,

downstream analyses can reveal clear pattern reflecting the underlying process

(Kemppainen et al., 2015). This has been illustrated for example by studies

demonstrating that SNPs within SOCs generated by polymorphic inversions in

Anopheles baimaii clearly separate the three expected karyotypes (inverted

homozygotes, heterozygotes and standard homozygotes) (Kemppainen et al., 2015).

We conducted downstream analyses with a PCA using SNPs identified within the

SOCs. As shown in Fig S2, although individuals were occasionally spread along three

PCA axes, no distinct cluster could be identified from any of the 20 SOCs. These

results were consistent with the absence of segregating inversions and local

adaptation in our samples and corroborated low F_{ST} values observed throughout the

genome. Precisely, in our data, we couldn't identify polymorphic inversions whose

karyotype frequencies change between Olama and Nyabessan due to a differential

adaption between the two sites. Some of the different SOCs identified can be

associated with other processes that were not captured by our analytical approach;

others are probably methodological artifacts associated with the LDna pipeline

(Kemppainen et al., 2015).

4. Discussion

We have analyzed the genome-wide polymorphism and characterized some of the baseline population genomic parameters in *An. moucheti*, an important malaria vector in rural areas across the African rainforest. We found very little differentiation among our samples, with most of the genetic variation distributed within individuals. Although a more substantial sampling will be necessary to fully dissect the population genetic structure of this species, our finding likely reflects the current dynamic of *An. moucheti* populations in Cameroon. It is worth mentioning that we have surveyed a total of 28 locations across the country (Fig 1A), some of which were known from several past surveys to harbor *An. moucheti* populations (Antonio-Nkondjio et al., 2013, 2009, 2008, 2006, 2002; Kengne et al., 2007), but we confirmed the presence of the species in only 2 villages. Extant populations of *An. moucheti* are distributed in patches of favorable habitats along river networks where larval populations breed. Our results indicate that despite this apparent fragmentation, connectivity and gene flow are high among population aggregates. The weak population genetic structure of *An. moucheti* observed with genome-wide markers corroborated results obtained with microsatellites and allozymes (Antonio-Nkondjio et al., 2008, 2002). A survey of eight microsatellite loci revealed that the highest F_{ST} among Cameroonian populations was as low as 0.003. Nevertheless, a substantial differentiation was found between samples from different countries consistent with an isolation-by-distance model (Antonio-Nkondjio et al., 2008). It is clear that a deep sequencing of continental populations is necessary to further clarify the status of these putative subpopulations. However, samples collected at

lower spatial scales like ours are also very relevant as they can allow robust

inferences about ongoing selective processes that cannot be captured at continental

scale. Although RADseq samples only a small fraction of the genome and certain

signatures of selection are likely missing when reduce representation sequencing

approaches are used, it has been shown that such approaches can effectively capture

strong footprints of selection across genomes of *Anopheles* mosquitoes (Fouet et al.,

Unpublished data; Kamdem et al., Unpublished data a). We have found that

signatures of selection are rare in the genome of *An. moucheti* populations from the

Cameroonian rainforest. Populations remain largely undifferentiated throughout the

genome, with F_{ST} values near zero across the vast majority of variations suggesting

that no local adaptation is ongoing. This perception is further supported by the

absence of segregating linkage disequilibrium blocks between geographic locations.

The characterization of chromosomal inversions with cytogenetic methods can be

laborious and challenging (Kirkpatrick, 2010; Sharakhova et al., 2014). So far, three

paracentric polymorphic inversions have been discovered in *An. moucheti* in

Cameroon (Sharakhova et al., 2014). The ecological, behavioral or functional roles of

these inversion polymorphisms remain unknown. We have implemented a recently

designed method that uses Next Generation Sequencing and LD estimates to

indirectly identify paracentric inversions whose karyotype frequencies varies

among populations due to local adaptation (Kemppainen et al., 2015). Our LD

analyses revealed the presence of a few LD clusters that are however not associated

with inversions. On the other hand, the low overall LD observed across the genome

reflected the significant genetic polymorphism that seems to prevail within *An.*

moucheti populations. This polymorphism translates into exceptional levels of overall genetic diversity and very high percentage of polymorphic sites that are in the range of values observed in other mosquito species undergoing significant demographic expansions (Donnelly et al., 2001; Fouet et al., Unpublished data; Kamdem et al., Unpublished data a). The amount of neutral genetic diversity is often viewed as a correlate of the adaptive potential of a species (Orr and Unckless, 2008). Although the relationship is more complex in reality, estimates of neutral genetic diversity are commonly used in conservation biology as an intuitive conceptual and management framework to assess the genetic resilience of endangered species (Bonin et al., 2007; Latta IV et al., 2010). Our population genomic analyses have depicted *An. moucheti* as a species with a great genetic diversity and hence a sustainable long-term adaptive resilience. Implications of our findings in malaria epidemiology and control can be very significant. First, *An. moucheti* is essentially endophilic and is particularly sensitive to the principal measures currently employed to control malaria in Sub-Saharan Africa such as the massive use of Insecticide Treated Nets (ITNs) and Indoor Residual insecticide Spraying (IRS). For example, estimates of population effective size in one village in Equatorial Guinea indicated that both mass distribution of ITNs and IRS campaigns resulted in a decline of approximately 55% of *An. moucheti* (Athrey et al., 2012). However, the great genetic diversity and the massive gene flow we observed within populations could easily enable this vector to challenge population declines and recover from shallow bottlenecks. Moreover, most insecticide resistance mechanisms found in insects exploit standing genetic variation to rapidly respond to the evolutionary

challenge by increasing the frequency of existing variations rather than relying on

infrequent *de novo* mutations (Messer and Petrov, 2013). As a result, despite the

current sensitivity of *An. moucheti* to common insecticides, the significant amount of

standing genetic variation provides the species with a great potential to challenge

insecticide-based interventions and other types of human-induced stress.

5. Conclusions

Recent advances in sequencing allow sensitive genomic data to be generated for

virtually any species (Ellegren, 2014). However, the most important information we

can obtain from population resequencing approaches often depends on the

availability and the quality of genomic resources such as a well-annotated reference

genome. The reduced genome sequencing strategy (RADseq) offers a cost-effective

strategy that can be used to effectively study the genetic variation in a broad range

of species from yeast to plants, insects, etc., in the absence of a reference genome.

We have extended this approach to the study of the genetic structure of an

understudied mosquito species with a great epidemiological significance. We have

provided both significant baseline population genomic data and the methodological

validation of one approach that should motivate further studies on this species and

other understudied anopheline mosquitoes lacking genomic resources.

Acknowledgements

Funding for this project was provided by the University of California Riverside and

NIH grants 1R01AI113248 and 1R21AI115271 to BJW. We thank populations and

authorities of the locations surveyed for their kind collaboration. We thank the

anonymous reviewers for their careful reading of our manuscript and their many

insightful comments and suggestions.

References

Alexander, D.H., Novembre, J., Lange, K., 2009. Fast model-based estimation of

ancestry in unrelated individuals. Genome Res. 19, 1655–1664.

Antonio-Nkondjio, C., Demanou, M., Etang, J., Bouchite, B., 2013. Impact of cyfluthrin

(Solfac EW050) impregnated bed nets on malaria transmission in the city of

Mbandjock : lessons for the nationwide distribution of long-lasting insecticidal

nets (LLINs) in Cameroon. Parasit. Vectors 6, 10. doi:10.1186/1756-3305-6-10

Antonio-Nkondjio, C., Kerah, C.H., Simard, F., Awono-Ambene, P., Chouaibou, M.,

Tchuinkam, T., Fontenille, D., 2006. Complexity of the malaria vectorial system

in Cameroon: contribution of secondary vectors to malaria transmission. J. Med.

Entomol. 43, 1215–1221.

doi:10.1603/0022-2585(2006)43[1215:COTMVS]2.0.CO;2

Antonio-Nkondjio, C., Ndo, C., Awono-Ambene, P., Ngassam, P., Fontenille, D., Simard,

F., 2007. Population genetic structure of the malaria vector Anopheles moucheti

in south Cameroon forest region. Acta Trop. 101, 61–68.

doi:10.1016/j.actatropica.2006.12.004

Antonio-Nkondjio, C., Ndo, C., Costantini, C., Awono-Ambene, P., Fontenille, D.,

Simard, F., 2009. Distribution and larval habitat characterization of Anopheles

moucheti, Anopheles nili, and other malaria vectors in river networks of

southern Cameroon. Acta Trop. 112, 270–276.

doi:10.1016/j.actatropica.2009.08.009

Antonio-Nkondjio, C., Ndo, C., Kengne, P., Mukwaya, L., Awono-Ambene, P.,

Fontenille, D., Simard, F., 2008. Population structure of the malaria vector

Anopheles moucheti in the equatorial forest region of Africa. Malar. J. 7, 120.

doi:10.1186/1475-2875-7-120

Antonio-Nkondjio, C., Simard, F., Cohuet, A., Fontenille, D., 2002. Morphological

variability in the malaria vector, Anopheles moucheti, is not indicative of

speciation: evidences from sympatric south Cameroon populations. Infect.

Genet. Evol. 2, 69–72. doi:10.1016/S1567-1348(02)00084-9

Athrey, G., Hodges, T.K., Reddy, M.R., Overgaard, H.J., Matias, A., Ridl, F.C.,

Kleinschmidt, I., Caccone, A., Slotman, M. a, 2012. The effective population size

of malaria mosquitoes: large impact of vector control. PLoS Genet. 8, e1003097.

doi:10.1371/journal.pgen.1003097

Ayala, D., Fontaine, M.C., Cohuet, A., Fontenille, D., Vitalis, R., Simard, F., 2011.

Chromosomal inversions, natural selection and adaptation in the malaria vector

Anopheles funestus. Mol. Biol. Evol. 28, 745–758. doi:10.1093/molbev/msq248

Ayala, F.J., Coluzzi, M., 2005. Chromosome speciation: humans, Drosophila, and

mosquitoes. Proc. Natl. Acad. Sci. U. S. A. 102 Suppl, 6535–42.

doi:10.1073/pnas.0501847102

Bonin, A., Nicole, F., Pompanon, F., Miaud, C., Taberlet, P., 2007. Population adaptive

index: A new method to help measure intraspecific genetic diversity and

prioritize populations for conservation. Conserv. Biol. 21, 697–708.

doi:10.1111/j.1523-1739.2007.00685.x

Catchen, J., Hohenlohe, P. a, Bassham, S., Amores, A., Cresko, W.., 2013. Stacks: An

analysis tool set for population genomics. Mol. Ecol. 22, 3124–3140.

doi:10.1111/mec.12354

Catchen, J.M., Amores, A., Hohenlohe, P., Cresko, W., Postlethwait, J.H., 2011. Stacks:

building and genotyping Loci de novo from short-read sequences. G3

(Bethesda). 1, 171–82. doi:10.1534/g3.111.000240

Clarkson, C.S., Weetman, D., Essandoh, J., Yawson, A.E., Maslen, G., Manske, M., Field,

S.G., Webster, M., Antão, T., MacInnis, B., Kwiatkowski, D., Donnelly, M.J., 2014.

Adaptive introgression between Anopheles sibling species eliminates a major

genomic island but not reproductive isolation. Nat. Commun. 5, 4248.

doi:10.1038/ncomms5248

Donnelly, M.J., Licht, M.C., Lehmann, T., 2001. Evidence for recent population

expansion in the evolutionary history of the malaria vectors Anopheles

arabiensis and Anopheles gambiae. Mol. Biol. Evol. 18, 1353–1364.

doi:10.1093/oxfordjournals.molbev.a003919

Ellegren, H., 2014. Genome sequencing and population genomics in non-model

organisms. Trends Ecol. Evol. 29, 51–63. doi:10.1016/j.tree.2013.09.008

Excoffier, L., Smouse, P.E., Quattro, J.M., 1992. Analysis of molecular variance

inferred from metric distances among DNA haplotypes: Application to human

mitochondrial DNA restriction data. Genetics 131, 479–491.

doi:10.1007/s00424-009-0730-7

Fontaine, M.C., Pease, J.B., Steele, a., Waterhouse, R.M., Neafsey, D.E., Sharakhov, I. V.,

Jiang, X., Hall, a. B., Catteruccia, F., Kakani, E., Mitchell, S.N., Wu, Y.-C., Smith, H.

a., Love, R.R., Lawniczak, M.K., Slotman, M. a., Emrich, S.J., Hahn, M.W., Besansky,

N.J., 2015. Extensive introgression in a malaria vector species complex revealed

by phylogenomics. Science (80). 347, 1258524–1258524.

doi:10.1126/science.1258524

Fouet, C., Kamdem, C., White, B.J., 2016. Chromosomal inversions facilitate

chromosome-scale evolution in Anopheles funestus. bioRxiv.

Gillies, M.T., Coetzee, M., 1987. A supplement to the Anophelinae of Africa south of

the Sahara. The South African Institute for Medical Research, Johannesburg.

Gillies, M.T., De Meillon, B., 1968. The Anophelinae of Africa South of the Sahara,

Second Edi. ed. Publications of the South African Institute for Medical Research,

Johannesburg.

Harrisson, K. a., Pavlova, A., Telonis-Scott, M., Sunnucks, P., 2014. Using genomics to

characterize evolutionary potential for conservation of wild populations. Evol.

Appl. n/a-n/a. doi:10.1111/eva.12149

Jombart, T., 2008. adegenet: a R package for the multivariate analysis of genetic

markers. Bioinformatics 24, 1403–1405.

Kamdem, C., Fouet, C., Gamez, S., White, B.J., 2016a. Pollutants and insecticides drive

local adaptation in African malaria mosquitoes. bioRxiv.

Kamdem, C., Fouet, C., Gamez, S., White, B.J., 2016b. Genomic signatures of

introgression at late stages of speciation in a malaria mosquito. bioRxiv.

Kemppainen, P., Knight, C.G., Sarma, D.K., Hlaing, T., Prakash, A., Maung Maung, Y.N.,

Somboon, P., Mahanta, J., Walton, C., 2015. Linkage disequilibrium network

analysis (LDna) gives a global view of chromosomal inversions, local adaptation

and geographic structure. Mol. Ecol. Resour. n/a-n/a. doi:10.1111/1755-

0998.12369

Kengne, P., Antonio-Nkondjio, C., Awono-Ambene, H.P., Simard, F., Awolola, T.S.,

Fontenille, D., 2007. Molecular differentiation of three closely related members

of the mosquito species complex, Anopheles moucheti, by mitochondrial and

ribosomal DNA polymorphism. Med. Vet. Entomol. 21, 177–182.

doi:10.1111/j.1365-2915.2007.00681.x

Kirkpatrick, M., 2010. How and why chromosome inversions evolve. PLoS Biol. 8.

doi:10.1371/journal.pbio.1000501

Kirkpatrick, M., Barton, N., 2006. Chromosome inversions, local adaptation and

speciation. Genetics 173, 419–434. doi:10.1534/genetics.105.047985

Latta IV, L.C., Fisk, D.L., Knapp, R.A., Pfrender, M.E., 2010. Genetic resilience of

Daphnia populations following experimental removal of introduced fish.

Conserv. Genet. 11, 1737–1745. doi:10.1007/s10592-010-0067-y

Lewontin, R.C., Kojima, K., 1960. The evolutionary dynamics of complex

polymorphisms. Evolution (N. Y). doi:10.2307/2405995

Meirmans, P., Van Tienderen, P., 2004. GENOTYPE and GENODIVE: two programs for

the analysis of genetic diversity of asexual organisms. Mol. Ecol. Notes 4, 792–

794.

Messer, P.W., Petrov, D., 2013. Population genomics of rapid adaptation by soft

selective sweeps. Trends Ecol. Evol. 28, 659–669.

doi:10.1016/j.tree.2013.08.003

Norris, L.C., Main, B.J., Lee, Y., Collier, T.C., Fofana, A., Cornel, A.J., Lanzaro, G.C., 2015.

Adaptive introgression in an African malaria mosquito coincident with the

increased usage of insecticide-treated bed nets. Proc. Natl. Acad. Sci.

201418892. doi:10.1073/pnas.1418892112

O'Loughlin, S.M., Magesa, S., Mbogo, C., Mosha, F., Midega, J., Lomas, S., Burt, A., 2014.

Genomic Analyses of Three Malaria Vectors Reveals Extensive Shared

Polymorphism but Contrasting Population Histories. Mol. Biol. Evol. 1–14.

doi:10.1093/molbev/msu040

Orr, H.A., Unckless, R.L., 2008. Population extinction and the genetics of adaptation.

Am. Nat. 172, 160–9. doi:10.1086/589460

Paradis, E., Claude, J., Strimmer, K., 2004. Analyses of Phylogenetics and Evolution in

R language. Bioinformatics 20, 289–290.

Peterson, B.K., Weber, J.N., Kay, E.H., Fisher, H.S., Hoekstra, H.E., 2012. Double Digest

RADseq: An Inexpensive Method for De Novo SNP Discovery and Genotyping in

Model and Non-Model Species. PLoS One 7, e37135.

doi:10.1371/journal.pone.0037135

Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M., Bender, D., Maller, J.,

Sklar, P., de Bakker, P., Daly, M., Sham, P., 2007. PLINK: a toolset for whole-

genome association and population-based linkage analysis. Am. J. Hum. Genet.

81.

Sharakhova, M. V., Antonio-Nkondjio, C., Xia, a., Ndo, C., Awono-Ambene, P., Simard,

F., Sharakhov, I. V., 2014. Polymorphic chromosomal inversions in Anopheles

moucheti, a major malaria vector in Central Africa. Med. Vet. Entomol. 28, 337-

340. doi:10.1111/mve.12037

Team, R.D.C., 2008. R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria.

Weir, B.S., Cockerham, C.C., 1984. Estimating F-statistics for the analysis of

population structure. Evolution (N. Y). 38, 1358–1370.

World Health Organization, 2013. World malaria report 2013. World Health

WHO/HTM/GM, 238. doi:ISBN 978 92 4 1564403

Author contributions

Conceived and designed the experiments: CF CK BJW. Performed the experiments:

CF CK SG BJW. Analyzed the data: CF CK BJW. Wrote the paper: CF CK BJW.

Competing interests

The authors declare that they have no competing interests.

Tables

Table 1: Information on *An. moucheti* samples included in this study.

Sampling locations	Geographic coordinates	Sampling methods			Total
		HLC-OUT	HLC-IN	LC	
Nyabessan	2°24'00"N, 10°24'00"E	21	15	1	37
Olama	3°26'00"N, 11°17'00"E	30	31	0	61
Total					98

HLC-OUT, human landing catches performed outdoor; HLC-IN, human landing catches performed indoor; LC, larval collection

548 **Table 2**: Population genomic parameters based on 6461 variant sites reflecting the

549 genetic diversity and conformity to Hardy-Weingberg equilibrium.

	Number of individuals*	Sites	Observed heterozygosity	F_{IS}	Nucleotide diversity (π)	% polymorphic sites
Variant positions						
Olama	54.12	6 461	0.0445	0.0631	0.0505	89.60
Nyabessan	16.78	6 461	0.0334	0.0372	0.0402	34.82
All positions						
Olama	54.47	165 975	0.0017	0.0025	0.0020	3.49
Nyabessan	16.86	165 975	0.0013	0.0014	0.0016	1.36

* Mean number of individuals per locus in this population (as estimated by Stacks v 1.35)

 Figure legends

Figure 1: Relationship between *An. moucheti* individuals from Olama and

Nyabessan. (A) Map of the study site showing both the locations surveyed (small

black dots) and the two villages (large red and blue squares) where *An. moucheti*

samples were collected. (B) and (C) Plots of the ADMIXTURE cross-validation error

and the Bayesian Information Criterion (BIC) (DAPC) as a function of the number of

genetic clusters indicating that k = 1. The lowest BIC and CV error indicate the

suggested number of clusters. (D) and (E) Absence of genetic structure within

populations illustrated by neighbor-joining and PCA. The percentage of variance

explained by each PCA axis is indicated.

Figure 2: Frequency distribution of F_{ST} between Olama and Nyabessan across 6461

SNP loci and plot of these F_{ST} values along arbitrary positions in the genome.

Figure 3: Allele Frequency Spectrum for 6461 SNP loci in Nyabessan and Olama

populations. The x-axis presents the frequency of the major allele and the y-axis the

frequency distribution of loci in each class of the major allele frequency.

Figure 4: LDna analyses on 2569 SNPs showing the presence of 20 Single Outlier

Clusters (SOCs) of linkage disequilibrium in *An. moucheti*. The graph presents the

results obtained with values of the two parameters: φ (which controls when clusters

are defined as outliers) and $|E|_{min}$, the minimum number of edges required for a LD

cluster to be considered as an outlier, indicated on top. LD thresholds are shown on

the x-axis.

Figures

Figure 1

Figure 2

Figure 3

Figure 4

Supplemental Material

Figure S1: Selection of individuals included in final analyses based on the average

per individual sequencing coverage. Neighbor-joining tree (A) and PCA (B)

indicating spurious population structure due to individuals with low sequencing

coverage in Olama (red) and Nyabessan (blue).

Figure S2: PCA indicating the population genetic structure inferred from SNPs

within the 20 Single Outlier Clusters (SOCs) of linkage disequilibrium identified in

An. moucheti (red: Olama; blue: Nyabessan).

Figure S1:

Figure S2:

601
602

603 **Table S1:** Distribution of the number of reads among sequenced individuals.
604 <u>Individuals below the dashed line were excluded from analysis.</u>

Mosquito ID	Total number of reads	Ambiguous/low quality reads	Retained reads	Sampling location
43	5190686	140706	5044739	Olama
850	4211705	136872	4066857	Nyabessan
27	2888406	77609	2800516	Olama
97	2773623	56956	2714480	Olama
94	2729590	61848	2660187	Olama
26	1915405	45856	1866121	Olama
82	1910651	50411	1854108	Olama
37	1764370	51133	1706506	Olama
95	1720323	37181	1678873	Olama
93	1450516	31318	1406481	Olama
25	1334643	45439	1285047	Olama
79	1263586	35615	1222166	Olama
80	1259353	42228	1202777	Olama
98	957698	20322	934945	Olama
49	874718	20033	852633	Olama
36	867344	21542	842175	Olama
99	853444	17206	834357	Olama
84	836989	18434	816949	Olama
76	850380	26481	814198	Olama
31	782599	18231	759965	Olama
86	761099	17235	741314	Olama
81	735291	13915	716347	Olama
1007	732960	19269	710564	Nyabessan
48	725668	17270	705049	Olama
743	703141	15098	686817	Nyabessan
46	639840	16191	621897	Olama
42	634781	16180	617726	Olama
92	593819	12260	579932	Olama
34	575175	11846	561789	Olama
90	579585	17480	556841	Olama
868	572514	16379	553831	Nyabessan
47	569810	12524	551627	Olama
28	557485	12363	534835	Olama
30	552365	13813	534263	Olama
38	541076	13190	525801	Olama
L_627	531803	12874	516440	Nyabessan
32	520421	11244	507425	Olama
78	510457	11112	493998	Olama
19	511063	15054	493676	Olama

87	507445	10896	492038	Olama
85	498180	10747	485508	Olama
792	478211	10753	465355	Nyabessan
35	467060	9616	455391	Olama
33	463964	10001	453130	Olama
60	465276	11304	452748	Olama
77	428787	10568	415155	Olama
45	391371	8565	380806	Olama
61	372074	10772	360550	Olama
851	363537	10079	352761	Nyabessan
40	346713	6421	339007	Olama
89	350001	12049	331665	Olama
15	341218	8157	331365	Olama
91	337294	12007	320481	Olama
44	323246	7406	313719	Olama
731	315382	10965	302496	Nyabessan
100	311056	5531	301968	Olama
729	295394	6781	287687	Nyabessan
18	290682	8206	281904	Olama
39	225981	6085	217620	Olama
73	220793	6020	212919	Olama
70	189567	4906	183785	Olama
756	188521	4639	182974	Nyabessan
75	181946	6799	173364	Olama
730	178968	5081	171828	Nyabessan
758	177709	5115	170782	Nyabessan
732	161697	4019	149839	Nyabessan
742	149159	3223	144972	Nyabessan
793	132552	4456	126135	Nyabessan
14	109314	3398	104900	Olama
96	99548	2293	95813	Olama
741	94635	2614	90106	Nyabessan
724	88884	3176	83605	Nyabessan
16	85486	2665	81748	Olama
88	83193	3414	78289	Olama
777	80457	2346	77695	Nyabessan
41	80127	2363	75238	Olama
72	76135	2366	73512	Olama
723	71212	2266	67258	Nyabessan
17	67037	2188	64443	Olama
795	65448	1778	62623	Nyabessan
869	57496	2654	54157	Nyabessan
760	45620	1610	43127	Nyabessan

727	43892	1402	40694	Nyabessan
71	37279	1233	35811	Olama
794	36669	1359	34439	Nyabessan
754	35476	1137	33712	Nyabessan
726	35274	1149	33230	Nyabessan
757	37431	1289	32648	Nyabessan
761	29783	947	28368	Nyabessan
833	27881	800	26582	Nyabessan
776	24245	783	22874	Nyabessan
755	24690	1012	22274	Nyabessan
753	19802	1301	17379	Nyabessan
728	17829	713	16750	Nyabessan
725	16069	706	14676	Nyabessan
810	14088	610	12963	Nyabessan
762	12307	568	11575	Nyabessan
763	11562	901	10374	Nyabessan

605