**Extensive genetic diversity among populations of the malaria**

**mosquito *Anopheles moucheti* revealed by population genomics**

Caroline Fouet[1]*, Colince Kamdem[1], Stephanie Gamez[1], Bradley J. White[1,2]*

[1]Department of Entomology, University of California, Riverside, CA 92521

[2]Center for Disease Vector Research, Institute for Integrative Genome Biology,

University of California, Riverside, CA 92521

*Corresponding authors: caroline.fouet@ucr.edu; bwhite@ucr.edu

**Key words:** *Anopheles moucheti*, population genomics, RADseq, *de novo* assembly

**Abstract**

**Background**: Recent intensive efforts to control malaria in African countries expose

vector populations to additional adaptive challenges. The main malaria mosquitoes

of the continent display an array of adaptive strategies to cope with such challenges.

The development of genomic resources will empower genetic studies that are

crucial to understand the evolutionary history and adaptive potential of these

vectors.

**Methodology/Principal findings**: Here we constructed double-digest Restriction

Associated DNA (ddRAD) libraries and generated 6461 Single Nucleotide

Polymorphisms (SNPs) that we used to explore the population structure and

demographic history of wild-caught *Anopheles moucheti* from Cameroon. The

genome-wide distribution of allelic frequencies among sampled populations best

fitted that of an old population at equilibrium, characterized by a weak genetic

structure and extensive genetic diversity, presumably due to a large long term

effective population size. In contrast to other important African malaria vectors,

polymorphic chromosomal inversions play little role in the genome architecture and

evolutionary adaptation of *An. moucheti*.

**Conclusions/Significance**: Our study provides the first investigation of the genetic

structure and diversity in *An. moucheti* at the genomic scale. Despite a weak genetic

structure and absence of adaptive divergence, the adaptive potential of this

mosquito remains significant owing to a great diversity and standing genetic

variation that can be used to face current vector control measures and other rapid

anthropogenic and environmental changes.

## Background

*An. moucheti sensu lato* is a group of three related mosquito species (*An. moucheti moucheti*, *An. moucheti nigeriensis*, and *An. moucheti bervoetsi*) distributed across forested areas in Sub-Saharan Africa and distinguishable from each other by slight morphological differences [1]. The nominal species *An. moucheti moucheti* (hereafter *An. moucheti*) is the most widespread and one of the leading malaria vectors in the equatorial region [2–4]. The species breeds in slow moving streams and rivers close to villages where it contributes to high malaria transmission and often outcompete other main malaria vectors species. Despite this epidemiological significance, the evolutionary history and the evolvability of this mosquito remain understudied. Early investigations of the genetic structure based on allozymes and microsatellites detected no significant differentiation among *An. moucheti* populations over substantial geographic areas. This finding was particularly intriguing given the widespread distribution of *An. moucheti* in the evergreen forest areas of Central Africa. Moreover, African anopheline mosquito populations are increasingly exposed to intense selective pressure associated with the use of insecticides and insecticide-impregnated materials associated with recent expansive malaria control campaigns [5]. Such pressures are strong driving forces that often contribute to rapid diversification of vector populations at the scales of a few decades [6–8]. As a result, a more detailed characterization of the genomic architecture of *An. moucheti* will be crucial to: (1) understand the genetic bases allowing populations of this species to adapt to such a wide geographic area and (2)

predict the evolutionary potential of this vector in face of selective constraints

imposed by rapidly changing environments.

Thanks to recent progresses in sequencing technology, high-resolution sequence

information can be generated for virtually any living organism. However, while

significant progresses have been made in genomic studies of other African

anopheline mosquitoes [9], *An. moucheti* has lagged behind mainly because of the

absence of genomic resources. So far, this species crucially lacks a reference genome

assembly, a physical or linkage map and laboratory strains that are essential to

generate high-quality sequencing information and to enable robust interpretations

of natural polymorphisms.

To start filling this gap, we have conducted the first screening of genetic variation

across the genome of wild populations of *An. moucheti*. We have performed a high-

throughput sequencing of reduced representation libraries in 87 wild-caught

individuals from Cameroon and used a *de novo* assembly to identify thousands of

RAD loci scattered throughout the genome. Using high-quality Single Nucleotide

Polymorphisms (SNPs) identified within these loci, we have investigated the genetic

relatedness and population history of our samples. We found that populations of *An.*

*moucheti* are characterized by extensive gene flow and a great genetic diversity. This

vector appears particularly adapted to challenge the selective pressures imposed by

vector controls and rapid human-induced environmental modifications.

## Methods

### Mosquito sampling and sequencing

This study included two *An. moucheti* populations from the Cameroonian equatorial rainforest. A total of 98 mosquitoes (97 adults and 1 larva) were collected in August and November 2013 Olama and Nyabessan, respectively (Table 1). The two locations are separated by ~200 km (Fig. 1A) and are crossed respectively by the Nyong and the Ntem rivers that provide the breeding sites for *An. moucheti* larvae. Specimens were identified as *An. moucheti moucheti* using morphological identification keys and a diagnostic PCR [1,10,11]. We extracted genomic DNA of specimens using either the DNeasy Blood and Tissue kit (Qiagen) or the Zymo Research MinPrep kit. We used 10ul (~50ng) of genomic DNA to prepare double-digest Restriction-site Associated DNA libraries following a modified protocol of Peterson et al.,2012 [12]. *MluC1* and *NlaIII* restriction enzymes were used to digest DNA of individual mosquitoes yielding RAD-tags of different sizes to which short unique DNA sequences  (barcodes) were ligated to enable the identification of reads belonging to each specimen. The digestion products were purified and pooled. DNA fragments with size ~400bp were selected and amplified via PCR. The distribution of fragment sizes was checked on a BioAnalyzer (Agilent Technologies, Inc., USA) before sequencing. The sequencing was performed on an Illumina HiSeq2000 platform (Illumina Inc., USA) (Genomic Core Facility, University of California, Riverside) to yield single-end reads of 101 bp.

### *De novo* assembly and SNP discovery

We used the bioinformatics pipeline Stacks v1.35 [13] to process Illumina short

reads. The program *process_radtags* was first used to sort the reads according to the

barcodes and to trim all reads to 96-bp in length by removing index and barcode

sequences from the ends of the reads. Reads with ambiguous barcodes, those that

did not contain the *NlaIII* recognition site and those with low-quality scores

(average Phred score < 33) were excluded. The program *ustacks* was then utilized

for a *de novo* assembly enabling the identification of consensus RAD loci in each

individual in our populations. We allowed a maximum of 2 nucleotide mismatches

between stacks (M parameter in *ustacks*) and we required a minimum of three reads

to create a stack (m parameter in *ustacks*). Using the *cstacks* program, a catalogue of

loci was built to synchronize variations across all individuals in our populations.

Finally, we utilized the *populations* program to calculate population genetic

parameters and output SNPs in different formats. To avoid bias associated with less

informative SNPs or possible false positive SNPs (due to sequencing or pipeline

errors), only RAD loci scored in at least 70-80% of individuals were retained for

further analyses.

**Population genomic analyses**

SNP files outputted by the *populations* program were used to assess the population

genetic structure with a Principal Component Analysis (PCA) and a Neighbor-joining

tree analysis using respectively the R packages *adegenet* and *ape* [14,15]. We also

explored patterns of ancestry and admixture in *An. moucheti* individuals in

ADMIXTURE v1.23 [16] with 10-fold cross-validation for k assumed ancestral

populations (k= 1 through 6). The optimal number of clusters was confirmed using

the Discriminant Analysis of Principal Component (DAPC) method which explores

the number of genetically distinct groups by running a k-means clustering

sequentially with increasing numbers and by comparing different clustering

solutions using Bayesian Information Criterion (BIC) [14]. We examined the

population genetic diversity, conformity to Hardy-Weinberg equilibrium and

demographic background using several statistics calculated with the *populations*

program. Precisely, to assess the global genetic diversity per population, we

calculated the overall nucleotide diversity ($\pi$) and the frequency of polymorphic

sites within population. To make inferences on the demographic history and to test

for departures from Hardy–Weinberg equilibrium, we utilized the Allele frequency

spectrum and the Wright's inbreeding coefficient ($F_{IS}$). To quantify the geographic

and genetic differentiation between allopatric populations, we estimated the

genome-wide average $F_{ST}$ [17] on 2000 randomly selected SNPs in Genodive v1.06

[18]. We also conducted an hierarchical Analysis of Molecular Variance (AMOVA)

[19] on the same SNP set to quantify the effects of the geographic origin on the

genetic variance among individuals. The statistical significance of $F_{ST}$ and AMOVA

was assessed with 10000 permutations. Finally, to examine the genomic

architecture divergence, we inspected the genome-wide distribution of locus-

specific estimates of $F_{ST}$.

**Identification of polymorphic chromosomal inversions**

The neutral recombination rate is notoriously reduced in genomic regions bearing

chromosomal inversions, which results in elevated linkage disequilibrium (LD) in

those regions relative to the rest of the genome. Thus, assessing genome-wide

patterns of LD can reveal clusters of highly correlated SNPs (LD blocks)

corresponding potentially to chromosomal inversions. The R package LDna [20]

allow the examination of the distinct LD network clusters within the genome of

nonmodel species without the need of a linkage map or reference genome. We

calculated LD, estimated as the $r^2$ correlation coefficient between all pairs of SNPs, in

PLINK v1.09 [21]. To avoid spurious LD due to the strong correlation between SNPs

located on the same RAD locus, we randomly selected only one SNP within each RAD

locus resulting in a dataset of 1056 variants containing less than 10% missing data.

LDna was then used to identify clusters of highly correlated SNPs that were

interpreted in light of the population demographic history, the population genetic

structure and previous cytological studies of *An. moucheti*.

**Results**

**De novo assembly**

In total, 518,218 unique 96-bp RAD loci were identified from *de novo* assembly of

reads in 98 individuals. We retained 946 loci that were present in all sampled

populations and in at least 75% of individuals in every population, and we identified

3027 high-quality biallelic SNPs from these loci.

**Population genetic structure**

First, we tested for the presence of cryptic genetic subdivision within *An. moucheti*

with PCA, NJ trees and the ADMIXTURE ancestry model. A NJ tree constructed from

a matrix of Euclidian distance using allele frequencies at 3027 genome-wide SNPs

showed a putative subdivision of *An. moucheti* populations in two genetic clusters

(Fig. S1). The first three axes of PCA also revealed a number of outlier individuals

separated from a main cluster (Fig. S1). However, when we ranked our sequenced

individuals based on the number of sequencing reads, we noticed that one of the

putative genetic clusters corresponded to a group of individuals having the lowest

sequencing coverage. We excluded all these individuals and reduced our dataset to

78 individuals. We conducted a *de novo* assembly and analyzed the relationship

between the 78 remaining individuals at 6461 genome-wide SNPs using PCA, NJ

trees and ADMIXTURE. Both the k-means clustering (DAPC) and the variation of the

cross-validation error as a function of the number of ancestral populations in

ADMIXTURE revealed that the polymorphism of *An. moucheti* resulted from only one

ancestral population (k = 1) (Fig. 1E,F). PCA and NJ depicted a homogeneous cluster

comprising all 78 individuals providing additional evidence of the lack of genetic or

geographic structuring among populations. Unsurprisingly, the overall $F_{ST}$ was

remarkably low between populations from the two sampling locations Olama and

Nyabessan ($F_{ST}$ = 0.008, p < 0.005). Similarly, the distribution of $F_{ST}$ values across the

6461 SNPs showed a large dominance of very low $F_{ST}$ values (Fig. 2). The highest per

locus $F_{ST}$ was only 0.126, while 5006 of the 6461 loci revealed $F_{ST}$ near zero. The

modest geographic differentiation was also well illustrated by a hierarchical

AMOVA, which showed that the genetic variance was explained essentially by

within-individual variations (99.7%). Finally, we found very low overall Wright's

inbreeding coefficient ($F_{IS}$= 0.0014 in Nyabessan and $F_{IS}$ = 0.0025 in Olama) (Table

2) suggesting that allelic frequencies within both populations were in accordance

with proportions expected under the Hardy-Weinberg equilibrium.

**Genetic diversity and demographic history**

The estimates of the overall nucleotide diversity ($\pi$ = 0.0020 and $\pi$ = 0.0016,

respectively, in Olama and Nyabessan) (Table 2) were within the range of average

values found in other African *Anopheles* species using RAD-seq approaches [8,22–

24]. Notorious demographic expansions have been described in natural populations

of this insect clade [25], and the values of $\pi$ observed in *An. moucheti* likely reflect

the level genetic diversity of a population with large effective size. The great genetic

diversity of *An. moucheti* was also illustrated by the percentage of polymorphic sites.

Of the 6461 variant sites, 89.60% were polymorphic in Olama and 34.82% in

Nyabessan (Table 2). The difference observed between the two locations can be

related to the sample size (n = 19 in Nyabessan and n = 59 in Olama) or to

demographic particularities that persists between the two geographic sites despite a

massive gene flow. To infer the demographic history of *An. moucheti*, we examined

the Allele Frequency Spectrum (AFS), summarized as the distribution of the major

allele in one population. This approach represented a substitute to model-based

methods that provide powerful examinations of the history of genetic diversity by

modeling the AFS at genome-wide SNP variants, but that couldn't be implemented

here due to the lack of reference genome assembly. The frequency distribution of

the major allele *p* (Fig. 3) indicates that the majority of loci that are polymorphic

between Olama and Nyabessan are fixed within each population as shown by the

predominance of SNPs at frequencies equal to 1. Ranges of allele frequencies are

similar in Olama and Nyabessan (between 0.47 and 1 in Olama and between 0.34

and 1 in Nyabessan). These frequency ranges are expected for old populations at

equilibrium capable of accumulating high amount of genetic diversity.

**Polymorphic chromosomal inversions**

LD analyses at 1056 highly filtered SNPs revealed a globally low LD in the *An.*

*moucheti* genome (average genome-wide $r^2$ = 0.0149) as expected in highly

polymorphic populations with large effective size. We used LDna to cluster the LD

values and to identify Single Outlier Clusters (SOC) that can be associated with

distinct or multiple evolutionary phenomena in the *An. moucheti* history. Results

showed the presence of two independent LD blocks in our samples (Fig. 4).

Cytogenetic analyses have identified three polymorphic chromosomal inversions in

Cameroon samples [26] and the two LD clusters could well correspond to linked

SNPs within inversions. Studies on *Anopheles baimaii* [20] have shown that when

SOCs are generated by polymorphic inversions, SNPs within the SOCs can clearly

separate the three expected karyotypes (inverted homozygotes, heterozygotes and

uninverted homozygotes). We conducted downstream analyses with a PCA using

respectively 30 and 34 SNPs identified within the two SOCs. As shown on Fig,

although individuals were spread along three PCA axes, no distinct cluster could be

identified for the first SOC. likewise, except three outlier individuals, the second SOC

formed a single genetic cluster (Fig. 4B,C). The two LD clusters were apparently not

associated with polymorphic inversions, but caution must be used when

interpreting this result as evidence of absence of inversions in our samples because

other LDna analyses conducted on *An. funestus* and *An. gambiae* which contain tens

of polymorphic inversions did not systematically identified the segregating

karyotypes of key adaptive inversions (results not shown). However, based on our

findings and previous cytogenetic studies, we can reasonably think that inversion

polymorphisms are less important in the genome architecture *An. moucheti*

compared with that of other important African malaria vectors such as *An. gambiae*

and *An. funestus* whose populations often segregate along clines of multiple

polymorphic inversions in nature.

**Discussion**

We have analyzed genome-wide polymorphism and characterized some of the baseline population genomic parameters in *An. moucheti*, an important malaria vector in the African rainforest. We found very little differentiation among our samples, with most of the genetic variation distributed within individuals. Although a more substantial sampling will be necessary to fully appreciate the population genetic structure of this species, our sampling likely reflects the current dynamic of *An. moucheti* populations in Cameroon. It is worth mentioning that we have surveyed a total of 28 locations across the country, some of which were known from several past surveys to harbor *An. moucheti* populations [1–4,27,28], but we confirmed the presence of the species in only 2 villages. Extant populations of *An. moucheti* are distributed in patches of favorable habitats along river networks where larval populations breed. Our results indicate that despite this apparent fragmentation, connectivity and gene flow are high among population aggregates. The weak population genetic structure of *An. moucheti* observed with genome-wide markers corroborated results obtained with microsatellites and allozymes [2,3]. A survey of eight microsatellite loci revealed that the highest $F_{ST}$ among Cameroonian populations was as low as 0.003. Nevertheless, a substantial differentiation was found between samples from different countries consistent with an isolation-by-distance model [3]. It is clear that a deep sequencing of continental populations is necessary to further clarify the status of these putative subpopulations that seem to be endemic to different countries. But, samples collected at lower spatial scales like ours are also relevant as they can allow robust inferences about ongoing selective

processes. We have found that signatures of selection are very rare in the genome of

*An. moucheti* populations from the Cameroonian rainforest. Populations remain

largely undifferentiated throughout the genome, with $F_{ST}$ values near zero across the

vast majority of segregating sites. Interestingly, polymorphic chromosomal

inversions that are the prime source of adaptive polymorphisms in many other

important malaria vectors seem to be less vital in the evolution of *An. moucheti*. The

characterization of chromosomal inversions with cytogenetic methods can be

laborious, particularly with regard to the identification of new inversions [26,29].

We have implemented a recently designed method that used Next Generation

Sequencing and LD estimates to identify paracentric inversions. Although the

method holds great promise in identifying inversions in genomes of nonmodel

species like *An. moucheti*, care should be taken when interpreting the results. So far,

only three paracentric polymorphic inversions have been discovered in *An. moucheti*

thanks to cytogenetics [26]. The ecological, behavioral or functional roles of these

inversion polymorphisms remain unknown. In keeping with cytogenetic

observations, LD analyses revealed the presence of only a few LD clusters that could

not be unambiguously associated with inversions. On the other hand, the low overall

LD observed across the genome reflected the significant genetic polymorphism that

seems to prevail in *An. moucheti* populations. This polymorphism translates into

exceptional levels of overall genetic diversity and very high percentage of

polymorphic sites that are in the range of values found in other mosquito species

undergoing a significant demographic expansion [8,24,25]. One of our main goals

was to know if the genetic diversity of *An. moucheti* can provide cues about its

extensive spatial distribution and can help to predict its environmental resilience. In

principle, evolutionary responses of species to human-induced or natural changes

rely largely on available heritable variation, which reflects the evolutionary

potential and adaptability to novel environments [30]. Therefore, the screening of

genome-wide variation is supposed to be a sensible approach that may provide a

generalized measure of evolutionary potential in species like *An. moucheti* for which

direct ecological, evolutionary or functional tests are impossible [31]. Our

population genomic analyses have depicted *An. moucheti* as a species with a great

genetic diversity and consequently a sustainable long-term adaptive resilience.

Implications of our findings in malaria epidemiology and control can be very

significant. First, *An. moucheti* is essentially endophilic and is particularly sensitive

to the principal measures currently employed to control malaria in Sub-Saharan

Africa such as the massive use of Insecticide Treated Nets (ITNs) and Indoor

Residual insecticide Spraying (IRS). For example, estimates of population effective

size in one village in Equatorial Guinea indicated that both mass distribution of ITNs

and IRS campaigns resulted in a decline of approximately 55% of *An. moucheti* [32].

However, the great genetic diversity and the massive gene flow we observed within

populations could easily enable *An. moucheti* to challenge population declines and

recover from shallow bottlenecks. Moreover, most insecticide resistance

mechanisms found in insects exploit standing genetic variation that allow to rapidly

respond to the evolutionary challenge by increasing the frequency of existing

variations rather than that of *de novo* mutations [33]. As a result, despite the current

sensitivity of *An. moucheti* to common insecticides, the significant amount of

standing genetic variation provides the species with a great potential to challenge

insecticides and other types of human-induced stress.

Recent advances in sequencing allow sensitive genomic data to be generated for

virtually any species [30]. However, the most important information we can obtain

from population resequencing approaches often depends on the availability and the

quality of genomic resources such as a well-annotated reference genome.

Nevertheless, the reduced genome sequencing strategy (RAD-seq) offers a cost-

effective strategy that can been used to effectively study the genetic variation in a

broad range of species from yeast to plants, insects, etc., in the absence of reference

genome. We have extended this approach to the study of the genetic structure of an

understudied epidemiologically important mosquito species. We have provided

both significant baseline population genomic data and the methodological validation

of one approach that should motivate further studies on this species and other

understudied anopheline mosquitoes lacking genomic resources.

**Acknowledgements**

Funding for this project was provided by the University of California Riverside and

NIH grants 1R01AI113248 and 1R21AI115271 to BJW. We thank populations and

authorities of the locations surveyed for their kind collaboration.

**References**

1. Kengne P, Antonio-Nkondjio C, Awono-Ambene HP, Simard F, Awolola TS,

Fontenille D. Molecular differentiation of three closely related members of the

mosquito species complex, Anopheles moucheti, by mitochondrial and ribosomal

DNA polymorphism. Med. Vet. Entomol. 2007;21:177–82.

2. Antonio-Nkondjio C, Simard F, Cohuet A, Fontenille D. Morphological variability in

the malaria vector, Anopheles moucheti, is not indicative of speciation: evidences

from sympatric south Cameroon populations. Infect. Genet. Evol. 2002;2:69–72.

3. Antonio-Nkondjio C, Ndo C, Kengne P, Mukwaya L, Awono-Ambene P, Fontenille

D, et al. Population structure of the malaria vector Anopheles moucheti in the

equatorial forest region of Africa. Malar. J. 2008;7:120.

4. Antonio-Nkondjio C, Ndo C, Costantini C, Awono-Ambene P, Fontenille D, Simard

F. Distribution and larval habitat characterization of Anopheles moucheti,

Anopheles nili, and other malaria vectors in river networks of southern Cameroon.

Acta Trop. 2009;112:270–6.

5. Bowen HL. Impact of a mass media campaign on bed net use in Cameroon. Malar.

J. Malaria Journal; 2013;12:36.

6. Norris LC, Main BJ, Lee Y, Collier TC, Fofana A, Cornel AJ, et al. Adaptive

introgression in an African malaria mosquito coincident with the increased usage of

insecticide-treated bed nets. Proc. Natl. Acad. Sci. 2015;201418892.

7. Clarkson CS, Weetman D, Essandoh J, Yawson AE, Maslen G, Manske M, et al.

Adaptive introgression between Anopheles sibling species eliminates a major

genomic island but not reproductive isolation. Nat. Commun. 2014;5:4248.

8. Kamdem C, Fouet C, Gamez S, White BJ. Pollutants and insecticides drive local

adaptation in African malaria mosquitoes. bioRxiv. 2016;

9. Neafsey DE, Waterhouse RM, Abai MR, Aganezov SS, Alekseyev MA, Allen JE, et al.

Highly evolvable malaria vectors: The genomes of 16 Anopheles mosquitoes. Science

(80-. ). 2015;347:1258522–1258522.

10. Gillies MT, De Meillon B. The Anophelinae of Africa South of the Sahara. Second

Edi. Johannesburg: Publications of the South African Institute for Medical Research;

1968.

11. Gillies MT, Coetzee M. A supplement to the Anophelinae of Africa south of the

Sahara. Johannesburg: The South African Institute for Medical Research; 1987.

12. Peterson BK, Weber JN, Kay EH, Fisher HS, Hoekstra HE. Double Digest RADseq:

An Inexpensive Method for De Novo SNP Discovery and Genotyping in Model and

Non-Model Species. PLoS One. 2012;7:e37135.

13. Catchen J, Hohenlohe P a., Bassham S, Amores A, Cresko W. Stacks: An analysis

tool set for population genomics. Mol. Ecol. 2013;22:3124–40.

14. Jombart T. adegenet: a R package for the multivariate analysis of genetic

markers. Bioinformatics. 2008;24:1403–5.

15. Paradis E, Claude J, Strimmer K. Analyses of Phylogenetics and Evolution in R

language. Bioinformatics. 2004;20:289–90.

16. Alexander DH, Novembre J, Lange K. Fast model-based estimation of ancestry in

unrelated individuals. Genome Res. 2009;19:1655–64.

17. Weir BS, Cockerham CC. Estimating F-statistics for the analysis of population

structure. Evolution (N. Y). 1984;38:1358–70.

18. Meirmans P, Van Tienderen P. GENOTYPE and GENODIVE: two programs for the

analysis of genetic diversity of asexual organisms. Mol. Ecol. Notes. 2004;4:792–4.

19. Excoffier L, Smouse PE, Quattro JM. Analysis of molecular variance inferred from

metric distances among DNA haplotypes: Application to human mitochondrial DNA

restriction data. Genetics. 1992;131:479–91.

20. Kemppainen P, Knight CG, Sarma DK, Hlaing T, Prakash A, Maung Maung YN, et

al. Linkage disequilibrium network analysis (LDna) gives a global view of

chromosomal inversions, local adaptation and geographic structure. Mol. Ecol.

Resour. 2015;n/a – n/a.

21. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira M, Bender D, et al. PLINK:

a toolset for whole-genome association and population-based linkage analysis. Am. J.

Hum. Genet. 2007;81.

22. O'Loughlin SM, Magesa S, Mbogo C, Mosha F, Midega J, Lomas S, et al. Genomic

Analyses of Three Malaria Vectors Reveals Extensive Shared Polymorphism but

Contrasting Population Histories. Mol. Biol. Evol. 2014;1–14.

23. Kamdem C, Fouet C, Gamez S, White BJ. Genomic signatures of introgression at

late stages of speciation in a malaria mosquito. bioRxiv. 2016;

24. Fouet C, Kamdem C, White BJ. Chromosomal inversions facilitate chromosome-

scale evolution in Anopheles funestus. bioRxiv. 2016;

25. Donnelly MJ, Licht MC, Lehmann T. Evidence for recent population expansion in

the evolutionary history of the malaria vectors Anopheles arabiensis and Anopheles

gambiae. Mol. Biol. Evol. 2001;18:1353–64.

26. Sharakhova M V., Antonio-Nkondjio C, Xia a., Ndo C, Awono-Ambene P, Simard F,

et al. Polymorphic chromosomal inversions in Anopheles moucheti, a major malaria

vector in Central Africa. Med. Vet. Entomol. 2014;28:337–40.

27. Antonio-Nkondjio C, Kerah CH, Simard F, Awono-Ambene P, Chouaibou M,

Tchuinkam T, et al. Complexity of the malaria vectorial system in Cameroon:

contribution of secondary vectors to malaria transmission. J. Med. Entomol.

2006;43:1215–21.

28. Antonio-Nkondjio C, Demanou M, Etang J, Bouchite B. Impact of cyfluthrin

(Solfac EW050) impregnated bed nets on malaria transmission in the city of

Mbandjock : lessons for the nationwide distribution of long-lasting insecticidal nets

(LLINs) in Cameroon. Parasit. Vectors. Parasites & Vectors; 2013;6:10.

29. Kirkpatrick M. How and why chromosome inversions evolve. PLoS Biol. 2010;8.

30. Ellegren H. Genome sequencing and population genomics in non-model

organisms. Trends Ecol. Evol. Elsevier Ltd; 2014;29:51–63.

31. Harrisson K a., Pavlova A, Telonis-Scott M, Sunnucks P. Using genomics to

characterize evolutionary potential for conservation of wild populations. Evol. Appl.

2014;n/a – n/a.

32. Athrey G, Hodges TK, Reddy MR, Overgaard HJ, Matias A, Ridl FC, et al. The

effective population size of malaria mosquitoes: large impact of vector control. PLoS

Genet. 2012;8:e1003097.

33. Messer PW, Petrov D. Population genomics of rapid adaptation by soft selective

sweeps. Trends Ecol. Evol. Elsevier Ltd; 2013;28:659–69.

## Author contributions

426 **Author contributions**

427 Conceived and designed the experiments: CF CK BJW. Performed the experiments:

428 CF CK SG BJW. Analyzed the data: CF CK BJW. Wrote the paper: CF CK BJW.

430 **Competing interests**

431 The authors declare that they have no competing interests.

## Tables

**Table 1**: Information on *An. moucheti* samples included in this study.

| Sampling locations | Geographic coordinates | Sampling methods | | | Total |
|---|---|---|---|---|---|
| | | **HLC-OUT** | **HLC-IN** | **LC** | |
| Nyabessan | 2°24'00"N, 10°24'00"E | 21 | 15 | 1 | 37 |
| Olama | 3°26'00"N, 11°17'00"E | 30 | 31 | 0 | 61 |
| Total | | | | | 98 |

HLC-OUT, human landing catches performed outdoor; HLC-IN, human landing catches performed indoor; LC, larval collection

**Table 2**: Population genomic parameters based on 6461 variant sites reflecting the genetic diversity and conformity to Hardy-Weingberg equilibrium.

| | n | Sites | Observed heterozygosity | $F_{IS}$ | $\pi$ | % polymorphic sites |
|---|---|---|---|---|---|---|
| **Variant positions** | | | | | | |
| **Olama** | 54.12 | 6 461 | 0.0445 | 0.0631 | 0.0505 | 89.60 |
| **Nyabessan** | 16.78 | 6 461 | 0.0334 | 0.0372 | 0.0402 | 34.82 |
| **All positions** | | | | | | |
| **Olama** | 54.47 | 165 975 | 0.0017 | 0.0025 | 0.0020 | 3.49 |
| **Nyabessan** | 16.86 | 165 975 | 0.0013 | 0.0014 | 0.0016 | 1.36 |

 **Figures**

**Figure 1**: Relationship between *An. moucheti* individuals from Olama and Nyabessan. (A) Map of the study site showing both the locations surveyed (small black dots) and the two villages (large red and blue squares) where *An. moucheti* samples were collected. (B) and (C) Absence of genetic structure within populations illustrated by neighbor-joining and PCA. The percentage of variance explained by each PCA axis is indicated. (D) and (E) Plots of the ADMIXTURE cross-validation error and the Bayesian Information Criterion (BIC) (DAPC) as a function of the number of genetic clusters indicating that k = 1. The lowest BIC and CV error indicate the suggested number of clusters.

**Figure 2**: Frequency distribution of $F_{ST}$ between Olama and Nyabessan across 6461

SNP loci and plot of these $F_{ST}$ values along arbitrary positions in the genome.

**Figure 3**: Allele Frequency Spectrum for 6461 SNP loci in Nyabessan and Olama

populations. The x-axis presents the frequency of the major allele and the y-axis the

frequency distribution of loci in each class of the major allele frequency.

**Figure 4**: LDna analyses on 1056 SNPs showing the presence of two Single Outlier

Clusters (SOCs) of linkage disequilibrium in *An. moucheti*. (A) Graph presenting the

results obtained with values of the two parameters: $\varphi$ (which controls when clusters

are defined as outliers) and $|E|_{min}$, the minimum number of edges required for a LD

cluster to be considered as an outlier, indicated on top. LD thresholds are shown on

the x-axis. (B) and (C) PCA indicating the population genetic structure inferred from

SNPs within the two SOCs (red: Olama; blue: Nyabessan).

**Supplemental Material**

**Figure S1**: Selection of individuals included in final analyses based on the average

per individual sequencing coverage. (A) and (B) Neighbor-joining tree and PCA

indicating spurious population structure due to individuals with low sequencing

coverage in Olama (red) and Nyabessan (blue).