1

3 Fitness costs of noise in biochemical reaction networks and the evolutionary limits of

4 cellular robustness

6 J. David Van Dyken
7
8
9 Department of Biology, Institute of Theoretical and Mathematical Ecology, University of
10 Miami, Coral Gables, FL 33146
11
12 vandyken@bio.miami.edu
13

20 Keywords: gene expression noise; drift barrier hypothesis; robustness; noise propagation;

21 allometric scaling; coarse-grained cell model; linear noise approximation; Michaelis-

22 Menten reaction; non-linear stochastic model

ABSTRACT

Gene expression is inherently noisy, but little is known about whether noise affects cell

function or, if so, how and by how much. Here I present a theoretical framework to

quantify the fitness costs of gene expression noise and identify the evolutionary and

synthetic targets of noise control. I find that gene expression noise reduces fitness by

slowing the average rate of nutrient uptake and protein synthesis. This is a direct

consequence of the hyperbolic (Michaelis-Menten) kinetics of most biological reactions,

which I show cause "hyperbolic filtering", a process that diminishes both the average rate

and noise propagation of stochastic reactions. Interestingly, I find that transcriptional

noise directly slows growth by slowing the average translation rate. Perhaps surprisingly,

this is the largest fitness cost of transcriptional noise since translation strongly filters

mRNA noise, making protein noise largely independent of transcriptional noise,

consistent with empirical data. Translation, not transcription, then, is the primary target of

protein noise control. Paradoxically, selection for protein-noise control favors increased

ribosome-mRNA binding affinity, even though this increases translational bursting.

However, I find that the efficacy of selection to suppress noise decays faster than linearly

with increasing cell size. This predicts a stark, cell-size-mediated taxonomic divide in

selection pressures for noise control: small unicellular species, including most

prokaryotes, face fairly strong selection to suppress gene expression noise, whereas larger

unicells, including most eukaryotes, experience extremely weak selection. I suggest that

this taxonomic discrepancy in selection efficacy contributed to the evolution of greater

gene-regulatory complexity in eukaryotes.

ARTICLE SUMMARY

Gene expression is a probabilistic process, resulting in random variation in mRNA and

protein abundance among cells called "noise". Understanding how noise affects cell

function is a major problem in biology. Here I present theory demonstrating that gene

expression noise slows the average rate of cell division. Furthermore, by modeling

stochastic gene expression with non-linearity, I identify novel mechanisms of cellular

robustness. However, I find that the cost of noise, and therefore the strength of selection

favoring robustness, decays faster than linearly with increasing cell size. This may help

explain the vast differences in gene-regulatory complexity between prokaryotes and

eukaryotes.

INTRODUCTION

The probabilistic nature of chemical reactions, along with the small volumes of living

cells, create substantial molecular randomness called "noise" (Berg 1978; McAdams and

Arkin 1997). Gene expression noise, for example, is measured as variation in mRNA or

protein abundance within a single cell over time (Taniguchi et al. 2010), or, equivalently,

among cells in an isogenic cell population (Elowitz et al. 2002). Because the abundance

of most mRNAs and proteins are on the order of 1 to 1000 copies per cell (Taniguchi et

al. 2010), a random birth or death of a single copy can instantaneously change the cellular

concentration by 0.1-100%. These random fluctuations are amplified by bursting, the

synthesis of multiple mRNAs or proteins in brief pulses (Raser and O'Shea 2004; Raser

and O'Shea 2005; Raj et al. 2006; Pedraza and Paulsson 2008; Thattai and van

Oudenaarden 2001). Using fluorescent probes and fluorescently tagged proteins,

researchers have quantified the noise statistics of mRNA and/or proteins in *E. coli*,

budding yeast, and mammalian cell lines, demonstrating the ubiquity of gene expression

noise and its considerable variability across the genome and among taxa (Elowitz et al.

2002; Blake et al. 2003; Golding et al. 2005; Raj et al. 2006; Zenklusen, Larson, and

Singer 2008; Taniguchi et al. 2010; Newman et al. 2006; Balázsi, van Oudenaarden, and

Collins 2011).

A central question in cell biology and evolution is whether or not cellular noise

has a functional consequence. Answering this question is necessary for establishing

rational design criteria for synthetic genetic circuits and for providing evolutionary

explanations for empirical patterns of noise among genes and taxa. For example, why

does *E. coli* have an average mRNA Fano factor (variance/mean) of $F = 1.6$ (Taniguchi et

al. 2010), which is just above the theoretical minimum of $F = 1$, whereas mammalian

cells have 40-100 times stronger mRNA noise (Raj et al. 2006; Balázsi,

van Oudenaarden, and Collins 2011; Dar et al. 2012)? While the mechanistic causes of

this discrepancy are becoming clearer (Suter et al. 2011), the evolutionary forces

permitting such a discrepancy remain murky.

Mounting empirical evidence suggests that natural selection has acted to limit

gene expression noise in many cases (Metzger et al. 2015; Newman et al. 2006; Alemu et

al. 2014; Batada and Hurst 2007; Fraser et al. 2004; Lehner 2008), implying that noise is

often costly to cells. Yet identifying and quantifying this cost remains a major challenge.

Perhaps the most widely invoked cost of cellular noise is imprecision, the lack of fine

control over cellular processes. Evolutionarily, this cost of noise is typically modeled as

random phenotypic deviations from the adaptive optimum, which is opposed by

stabilizing selection (Wagner et al. 1997; Draghi and Whitlock 2015). Several molecular

mechanisms linking imprecision to suboptimal fitness deviations have been proposed.

The dosage-control hypothesis proposes that noise disrupts control over the

stoichiometric balance of interacting proteins (Fraser et al. 2004; Lehner 2008).

Consistent with this hypothesis, proteins that form complexes have lower average noise

than other proteins (Fraser et al. 2004; Lehner 2008; Bar-Even et al. 2006), but models

currently do not predict the magnitude of this effect so that quantitative correspondence

between predictions and observations remain unknown. More commonly, researchers

focus on how noise disrupts the fine control over cell states regulated by bistable

(ON/OFF) switches (Balázsi, van Oudenaarden, and Collins 2011; Csete and Doyle

2002). Noise in the input parameter can cause the cell to flip stochastically from the

desired to undesired state, or to respond slowly to environmental cues, creating a

maladaptive mismatch between the cell state and the selecting environment. However,

the least noisy genes, those presumably experiencing the strongest selection to suppress

noise, are constitutively expressed and involved in core cell functions, not control over

bistability (Fraser et al. 2004; Lehner 2008; Bar-Even et al. 2006). This cost is also

difficult to quantify in any general way because its effects are highly context dependent,

to the extent that in some contexts the cost of imprecision can change signs, becoming

evolutionarily beneficial (Acar, Mettetal, and van Oudenaarden 2008; Rao, Wolf, and

Arkin 2002; Kussell and Leibler 2005; Eldar and Elowitz 2010; Kaern et al. 2005; Thattai

and Van Oudenaarden 2004).

As an alternative, here I demonstrate a general, readily quantifiable and consistent

cost of cellular noise. I show that noise in substrate abundance slows the average rate of

product formation in Michaelis-Menten-type (i.e., non-cooperative bimolecular) chemical

reactions, which constitute most reactions in the cell including those most closely tied to

fitness: nutrient uptake and protein synthesis. This result is a straightforward consequence

of the non-linearity of bimolecular reactions. While it is well known that noise alters the

mean of nonlinear stochastic systems, this effect is typically ignored, giving rise to the

widespread use of linear approximations (e.g., the "linear noise approximation" (LNA)

(van Kampen 2007)), particularly in studies of gene expression noise (Thattai and van

Oudenaarden 2001; Shahrezaei and Swain 2008; Paulsson 2004). The LNA is well-

justified when applied to investigating steady-state concentrations in noisy chemical

networks, since the difference between linear and non-linear predictions at steady-state

are on the order of a single molecule. However, fitness in living cells is determined by

rates—not concentrations--integrated over an enormous number of reactions over the

lifetime of a cell and its genetic lineage. For example, an *E. coli* cell must carry out a

minimum of 2.5×10^6 translation reactions to produce a single daughter cell. A minor

slowing of reaction rates within cells, if heritable, may pose a non-negligible selective

cost. From this perspective, the effects of noise on fitness cannot be ignored, and indeed,

I find that they can be substantial in small cells.

RESULTS

The goal of the paper is to quantify the strength of selection favoring the attenuation of

gene expression noise, and to identify the targets of noise attenuation. The paper proceeds

as follows. First, I show that noise slows the average rate of biochemical reactions that

have non-linear, hyperbolic kinetics. Then, I extend this result to networks of coupled

non-linear reactions in order to determine how noise effects total flux through

biochemical networks, as well as how the network architecture itself affects the

propagation of noise. This enables us to quantify the cost of noise, measured as the loss in

average rate of end-product formation, and to quantify the noise statistics of the system. I

then apply the model to gene expression, focusing on the consequences of non-linear,

hyperbolic translation kinetics on the propagation of mRNA noise to the protein level,

and to the average rate of biomass synthesis. Finally, I investigate how the volume of

reaction compartments and whole-cells influences the cost of noise, concluding that the

cost of noise, and therefore the selection pressures favoring low gene expression noise,

decays rapidly with increasing cell size, leading to a testable prediction about levels of

gene expression noise across unicellular species.

Noise-induced slowdown of Michaelis-Menten reactions. The purpose of this section is

to begin building the theory from the simplest component of a network: a single isolated

reaction. We will see that input noise slows the reaction rate of a single, isolated reaction

obeying Michaelis-Menten type kinetics. In the following section, the analytical results

from this section are put on a more rigorous theoretical footing and then extended to

networks of coupled reactions in open systems.

The cornerstone of biochemistry is the Michaelis-Menten (MM) reaction, which

describes the enzyme catalyzed conversion of substrate, *S*, into product, *P*. The rate of

product formation is described by the hyperbolic MM equation (Michaelis and Menten

1913; Briggs and Haldane 1925; Ingalls 2013):

$$\frac{dp}{dt} = V = V_{max} \frac{s}{K_m + s}$$ 1

s, *e* and *p* are reactant, enzyme and product concentration, respectively (following

convention lowercase denotes concentration, which unless otherwise noted will take units

of molecules per cell), $V_{max} = k_{cat}e$ is the asymptotic reaction rate (k_{cat} is the number of

substrate molecules converted to product per enzyme per unit time), and K_m, the

Michaelis or half-saturation constant, is the substrate concentration at $V = V_{max}/2$. This

equation is derived from a system of elementary reactions by applying the law of mass

action and the quasi-steady state approximation (QSSA) (Briggs and Haldane 1925;

Ingalls 2013) (Sup. Mat.). The QSSA is a separation of timescales approximation that

eliminates fast variables by setting them to their steady-state. The law of mass action is

valid in cases where the timescale of molecular diffusion is fast relative to the reaction

rate, so that the system behaves as if it were well-mixed. The rate of enzymatic reactions

in cells is far below the diffusion limit (Bar-Even et al 2006), so that mass action is a

reasonable approximation in most cases. For stochastic reactions, wherein *s*, *e*, and *p*

experience random number fluctuations, Eqn. 1 takes on a probabilistic interpretation as

the "reaction propensity" (van Kampen 2007) giving the transition probability for $n_P \rightarrow$

$n_P + 1$ and $n_S \rightarrow n_S - 1$, where *n* denotes particle number. In general, the QSSA is valid

for stochastic systems under the same conditions for which it is valid in deterministic

systems (Rao and Arkin 2003; Sanft, Gillespie, and Petzold 2011), notwithstanding

finite-volume corrections (Grima 2009b; Grima 2009a; Grima 2010) (see below).

Because Eqn. 1 is hyperbolic in *s*, Jensen's inequality proves that, for random *s*,

$\langle V(s) \rangle \leq V(\langle s \rangle)$ (angle brackets denote ensemble average). That is, the average reaction

rate is less than or equal to a noise-free (i.e., deterministic) reaction with $s = \langle s \rangle$ (Figs. 1,

S1). This is seen by expanding Eqn. 1 to second order in Taylor series about $\langle s \rangle$ and

taking the expectation (see SI),

$$\langle V(s) \rangle = \underbrace{V_{max} \frac{\langle s \rangle}{K_m + \langle s \rangle}}_{V(\langle s \rangle)} - \underbrace{\frac{\langle \epsilon_S^2 \rangle}{\langle s \rangle^2} V_{max} a(1-a)^2}_{noise-induced\ slowdown} + O(\epsilon_S^3) \qquad\qquad 2$$

$a = K_m/(K_m + \langle s \rangle)$ is the "kinetic order" of the MM reaction evaluated at the average

substrate concentration, and $(\eta^2 = \langle \epsilon_S^2 \rangle / \langle s \rangle^2)$ the squared coefficient of variation, which

is often simply called "the noise". The leading order correction in Eqn. 2 takes the sign of

the second derivative of Eqn. 1, which is everywhere negative for hyperbolic transition

rates, thus demonstrating the noise-induced slowdown of MM type reactions.

The kinetic order parameter, *a,* is a key parameter that appears repeatedly below.

a is identical to the steady-state fraction of unbound enzyme. In metabolic control

analysis (MCA), *a* is equivalent to the "elasticity coefficient" (Kacser and Burns 1973).

In general, *a* measures the nonlinearity of the reaction rate at a given mean substrate

concentration. First and zeroth order reactions ($a = 1$ and 0, respectively) are both linear,

and as a consequence there is no noise-induced slowdown, consistent with Jensen's

Inequality. The rate of a first order reaction is linearly dependent on the substrate

concentration, whereas the rate of a zeroth order reaction is independent of substrate

concentration. In a MM type reaction, the reaction is in its first order regime when

substrate is scarce, and in the zeroth order regime when substrate is abundant (the flat

part of the rate law curve in Fig. 1).

Figure 1: **Noise-induced slowdown of hyperbolic reactions.** Stochastic
simulations employing Gillespie's (Gillespie 1977) exact stochastic simulation algorithm
(SSA) of the microscopic system of elementary reactions for the Michaelis-Menten
reaction (see SI). Points represent the rate of product formation over time steps 100-800
(providing a burn-in more than 10 times the half life of P or S) ensemble averaged over
1000 realizations of the stochastic process. Results fit well with the QSSA model (Eqn. 3,
solid lines in **a**, **b** and thin black line in **c**). The substrate burst input, b, is varied to alter
the substrate noise level. Substrate input rate, k_{in} is divided by b so that mean substrate is
not affected by bursting, and is varied to generate different substrate concentrations. All
other parameters are constant ($e = 1$, $k_{ON} = 1$, $k_{OFF} = 19.5$, $k_{cat} = 0.5$, $\delta_S = 0.1$, $\delta_P = 0.1$). **A)**
The rate law curve demonstrating that, for a given average substrate concentration,
substrate noise slows the reaction, but only if the reaction is not in its first or zeroth order
regime (low or high substrate, respectively). Note that the accuracy of the stochastic
QSSA requires that s, $K_m << F$, which fails for the red points in **a** and **b** (where $K_m = 20$
and $F = 20.5$). Inset is a zoomed in view. **B)** A geometric illustration of noise-induced
slowdown, showing that variance (noise) in a state variable reduces the average of a
hyperbolic function (i.e., Jensen's inequality). **C)** 100 realizations of the stochastic
process (gray) compared to the deterministic (thick black line) and stochastic (thin black
line) predictions. Red traces in inset are example outputs from a single realization of the
substrate dynamics for a short time window.

A more rigorous and general expansion method is used in the following section

("Noisy Non-linear Reaction Networks in Finite Volumes"). For now, there are two main

points: 1) reaction kinetics depend not only on average molecule concentrations, but on

the mean and variance in molecule numbers, and 2) for a given average substrate

concentration, substrate noise reduces the rate of biochemical reactions. The difference

between the average stochastic rate and deterministic rate grows monotonically with

increasing noise, and equality occurs only with zero noise or zero curvature of $V(s)$.

Mechanistically, this can be understood as follows: positive fluctuations in substrate

oversaturate available catalysts leaving many unbound S molecules that are not actively

creating product, causing inefficiency. As a consequence, the same amount of substrate

would produce a faster net reaction rate if spread out evenly over the same time interval

(i.e., the same mean with zero noise).

If the substrate statistics are known, then the probability density of V with random

s can be solved explicitly for a number of common probability distributions. mRNA and

protein counts are typically gamma distributed within a cell over time or among cells in a

clonal population (Taniguchi et al. 2010). The gamma distribution has shape parameter α

$= CV^{-2}$ and rate parameter $\beta = F$, where $CV^2 = \sigma_x^2/\langle x \rangle^2$ and $F = \sigma_x^2/\langle x \rangle$ are the squared

coefficient of variation (the noise) and the Fano factor (often called "the noise strength"),

respectively (Taniguchi et al. 2010). F and CV^2 are common measures of noise intensity,

with convenient mechanistic interpretations in gene expression: F is closely related to the

burst size (e.g., number of proteins synthesized per mRNA lifetime) and CV^{-2} to the burst

frequency (e.g., the number of mRNA's synthesized per protein lifetime). F is

independent of cell volume, whereas CV^2 decreases with cell volume (they are intensive

and extensive variables, respectively). For gamma distributed s, the probability density

function of the Michaelis-Menten reaction rate (for $V < V_{max}$) is exactly,

$$Pr(V = v) = \frac{V_{max}}{k_m} \frac{v^{\alpha-1}e^{-v/z\beta}}{z(z\beta)^\alpha \Gamma(\alpha)}, \qquad z = \frac{V_{max} - v}{k_m} \qquad\qquad 3$$

Eqn. 3 is verified by Monte Carlo simulations implemented with Gillespie's exact

stochastic simulation algorithm (SSA) (Gillespie 1977) (Fig. 1). Eqn. 3 fits data from

stochastic simulations very well provided that $F, K_m \ll s$. Equation 2 gives a very good

approximation to 3 under this same restriction (Fig. S2), which provides an important

benchmark for the accuracy of the more rigorous expansion approximations used below.

Figure 1 shows that the effect of noise on MM-type reactions is to increase the

observed Michaelis constant. Therefore, a stochastic formulation of Eqn. 1 that accounts

for noisy substrate while retaining the classical form is obtained simply by substituting

into Eqn. 1 an "effective", stochastic Michaelis constant, $\widetilde{K}_m$, where $\widetilde{K}_m \geq K_m$ with

equality occurring only with zero noise, zero curvature of the rate law, and macroscopic

reaction volume (S6).

Noisy Non-linear Reaction Networks in Finite Volumes. In the previous section, we

showed that substrate noise slows the rate of product formation in a single hyperbolic

reaction in a closed system. To apply this result, the statistics of the substrate species are

required. We now take a more rigorous approach and generalize the noise-induced

slowdown phenomenon to reaction networks in open systems. Further, the model allows

one to predict the noise statistics of the system solely with knowledge of the network

structure and reaction kinetics. Of particular interest is how network architectures may

create feedbacks that buffer the system flux against noise-induced slowdown. The

following theory builds on the work of van Kampen (van Kampen 2007) and Elf and

Erhenberg (Elf and Ehrenberg 2003) (see also Grima (Grima 2010)).

The time evolution of the joint probability density of all N nodes in a chemical

reaction network with R edges is given by the following "system-size expansion" (van

Kampen 2007) of the multidimensional chemical master equation (derived in the SI):

$$\overbrace{\frac{\partial \Pi(\vec{\epsilon},t)}{\partial t} = -\sum_{i,k}^{N} J_{ik}\frac{\partial(\epsilon_k\Pi)}{\partial\epsilon_i} + \frac{1}{2}\sum_{i,k=1}^{N} D_{ik}\frac{\partial^2\Pi}{\partial\epsilon_i\partial\epsilon_k}}^{linear\ Fokker-Planck\ equation}$$

$$+\underbrace{\frac{\Omega^{-1/2}}{2}\left(-\sum_{i,k,l=1}^{N}\frac{\partial J_{ik}}{\partial\overline{x}_l}\frac{\partial(\epsilon_l\epsilon_k\Pi)}{\partial\epsilon_i} + \sum_{i,k,l=1}^{N}\frac{\partial D_{ik}}{\partial\overline{x}_l}\frac{\partial^2(\epsilon_l\Pi)}{\partial\epsilon_i\partial\epsilon_k}\right)}_{correction\ for\ non-linearity\ and\ finite\ volume} + O(\Omega^{-1}) \qquad 4$$

282 $\Pi = \Pi(\vec{\epsilon},t)$ is the joint probability density of all reacting species, Ω is the reaction

283 volume, ϵ_i is the random perturbation of the i^{th} species from $\overline{x}_i$, its macroscopic steady-

284 state (overbars will denote the macroscopic steady-state value of a variable throughout),

285 S_{ij} is the stoichiometric coefficient of species i in the j^{th} reaction, and

286
$$J_{ik} = \sum_{j=1}^{R} S_{ij}\frac{\partial f_j(\vec{\overline{x}})}{\partial\overline{x}_k}, \quad D_{ik} = \sum_{j=1}^{R} S_{ij}S_{kj}\,f_j(\vec{\overline{x}})$$

are entries of the Jacobian and diffusion matrices, respectively, of the continuous

deterministic system of equations. Equation 4 is a linearization of the chemical master

equation obtained by applying the ansatz that concentrations fluctuate about their

macroscopic steady-state value, with fluctuation size scaling inversely with the square

root of the reaction volume, Ω (van Kampen 2007). This inverse square-root dependence

of noise on volume is a formal statement of the Law of Large Numbers, and is formally

equivalent to the relationship between sample size and the standard deviation of a

sampling distribution. One thus substitutes, $x_i = \overline{x}_i + \Omega^{-1/2}\epsilon_i$, for each species i in the

deterministic system of equations, and then expands each transition rate, $V_j(x)$, in powers

of $\Omega^{-1/2}$ about $\overline{x}_i$, truncating the series at the desired order (van Kampen 2007). The terms

of order $O(\Omega^0)$ in (4) give the "linear noise approximation" (LNA) (van Kampen 2007),

which is equivalent to the standard linear Fokker-Planck equation, while the terms in Ω^{-}

$^{1/2}$ account for noise-induced deviations caused by non-linear rate laws in finite volumes,

which we show cannot be ignored in living cells.

To apply 4, one begins by writing a macroscopic description of a reaction network

in QSSA reduced form. Here we analyze a generalized MM scheme (S17) in an open

system obeying the following macroscopic system of ODEs (Sup. Mat.):

304
$$\frac{ds}{dt} = b_S k_{in,S} - hV - \delta_S s, \qquad \frac{de}{dt} = b_E k_{in,E} - \delta_E e, \qquad \frac{dp}{dt} = V - \delta_P p \qquad\qquad 5$$

δ_i and $k_{in,i}$ are the decay and input rates, respectively, of i, and $V = k_{cat} es/(K_m + s)$.

Substrate (S) and enzyme (E) are fed into the system via a burst process with burst size b_S

and b_E, respectively ($n_S \rightarrow n_S + b_S$; $n_E \rightarrow n_E + b_E$). $h = \{0,1\}$ is an indicator variable

that toggles between reactions where S is consumed ($h = 1$), as in most metabolic

reactions, and reactions where S is not consumed ($h = 0$), as with template-mediated

reactions such as translation where E denotes ribosomes and S mRNA, which is not

consumed in translation.

From System 5 and Eqn 4, we obtain the following steady-state average

concentrations of each species in a generalized, open MM reaction network (Sup. Mat.):

314
$$\langle s \rangle = \bar{s} + h \underbrace{\left[a \frac{\langle \epsilon_S^2 \rangle}{\bar{s}} - K_m \frac{\langle \epsilon_S \epsilon_E \rangle}{\bar{e}\bar{s}} \right] A\Omega^{-1}}_{stochastic\ amplification}, \qquad \langle e \rangle = \bar{e} ,$$

315
$$\langle p \rangle = \bar{p} - \frac{\delta_S}{\delta_P}\left[a \frac{\langle \epsilon_S^2 \rangle}{\bar{s}} - K_m \frac{\langle \epsilon_S \epsilon_E \rangle}{\bar{e}\bar{s}} \right] A\Omega^{-1}, \qquad\qquad 6$$

316
$$\langle V \rangle = \bar{V} - \delta_S \underbrace{\left[a \frac{\langle \epsilon_S^2 \rangle}{\bar{s}} - K_m \frac{\langle \epsilon_S \epsilon_E \rangle}{\bar{e}\bar{s}} \right] A\Omega^{-1}}_{noise-induced\ slowdown}$$

Where $A = (k_{cat}\bar{e}\bar{s}/(hk_{cat}\bar{e}K_m + \delta_S(K_m + \bar{s})^2))$ and $a = K_m/(K_m + \bar{s})$ is the MM

reaction order. Note that in the general case, $\langle s \rangle$ is increased and $\langle p \rangle$ is decreased by

noise compared to the macroscopic system. The non-linear mesoscopic correction terms

vanish when $s\rightarrow0$ (first order kinetics), $s\rightarrow$Infinity (zeroth order kinetics) or when

$\Omega\rightarrow$Infinity (macroscopic volume). Two of the second moments (Sup. Mat.) for the

general MM case can be written compactly:

323
$$\langle\epsilon_E^2\rangle = \frac{1}{2}\bar{e}(b_E + 1), \quad \langle\epsilon_S\epsilon_E\rangle = -h\langle\epsilon_E^2\rangle\frac{k_{cat}\bar{s}}{hak_{cat}\bar{e}+(\delta_E+\delta_P)(K_m+\bar{s})} \qquad 7$$

While the present approach builds on that derived by Grima (Grima 2010) the

results and conclusions differ: Grima (Grima 2010) found that noise had no effect on

product concentration in MM reactions. There are two reasons for this discrepancy. First,

Grima assumed no dilution of the substrate ($\delta_S = 0$). As can be seen from equation 6, this

causes the finite volume correction term for $\langle p \rangle$ to vanish. Thus, Grima's results are in

agreement with the present results for this special case, and all of the results in his papers

are correct. However, following (Grima 2010) exactly for $\delta_S > 0$ also gives zero

stochastic deviation of the product concentration (see SI), which is incorrect. Here is

why: when the system-size expansion of the chemical master equation is applied to a

system of elementary reactions, as in (Grima 2010) and most other methods, the rate of

product formation, V, is proportional to the intermediate complex concentration, c.

Because c is unimolecular, the method reads this rate as linear (its second derivative is

zero), and so the mean cannot be affected by noise (as proven by Jensen's inequality).

However, in MM reactions V is decidedly non-linear. Indeed, it is hyperbolic. Thus, the

elementary reaction conceals an essential non-linearity of the system, and will give

incorrect results in the general case. To rectify this, one must apply Eqn. 4 to a QSSA

reduced system of biochemical equations as shown in the SI. Note that the use of

elementary or QSSA reactions is not up to choice: the elementary reaction scheme will

give the wrong answer. Note also that applying the method of (Grima 2010) to a QSSA

reduced system also gives incorrect results, for reasons discussed in the SI.

Noise propagation in gene expression with hyperbolic filtering. Because gene

expression noise is costly, there will ostensibly be selection pressure for noise control.

But how can a cell suppress gene expression noise? That is, what are the evolutionary

targets of noise-control? To answer these questions, we must model the process of gene

expression, form transcription through translation, to identify how noise is generated and

filtered.

We deviate from previous models of gene expression noise by assuming that

translation obeys non-linear reaction kinetics. Previous models of gene expression noise

have considered translation as a first-order process (Thattai and van Oudenaarden 2001;

Shahrezaei and Swain 2008; Paulsson 2004; Bar-Even et al. 2006; Pedraza and Paulsson

2008; McAdams and Arkin 1997). However, translation in living cells may actually be

closer to zeroth-order than first-order, with a reaction order of $a = 0.1$-0.2 in log-phase

growth in *E. coli* and budding yeast, based on the fraction of unoccupied ribosomes

(Arava et al. 2003; Zenklusen, Larson, and Singer 2008; Ingolia 2014). We now

investigate the effects of hyperbolic translation kinetics on gene expression noise.

Following previous models of gene expression noise (Thattai and van

Oudenaarden 2001; Shahrezaei and Swain 2008; Paulsson 2004; Bar-Even et al. 2006;

Pedraza and Paulsson 2008; McAdams and Arkin 1997), we assume for now that

translation is initiation limited (see "Cell fitness" section below for generalization), such

that it relies only on the concentrations of ribosomes, r, and mRNA, m. If mRNA is

transcribed in bursts of size b_t, then the steady-state variance and Fano factor of mRNA

are (Sup. Mat.),

$$\sigma_{mRNA}{}^2 = \tfrac{1}{2}\overline{m}(b_t + 1), \qquad F_{mRNA} = \tfrac{1}{2}(b_t + 1) \qquad\qquad 8$$

which is a special case of (Elf and Ehrenberg 2003) and which is in excellent agreement

with simulations (Figs 2, 3).

Figure 2: **Protein noise with hyperbolic filtering.** Translation propagates mRNA noise
to the protein level, but hyperbolic reaction kinetics (0<*a*<1) can substantially ameliorate
this noise and cause highly non-linear relationships between noise and underlying

parameters. Points represent ensemble variance of protein number from the last of 110
time steps over 10,000 realizations of the SSA (see SI), lines represent theoretical
prediction (from Eqns 9 and S21). **A, C)** Increasing ribosomal binding affinity (i.e.,
reducing K_m) has two opposing effects on protein noise: it reduces reaction order but
increases translational bursting. Which effect dominates depends on the population's
starting point relative to the protein noise peak at a* = 2/3. Most species have
translational $a < 2/3$, suggesting that increasing ribosomal binding affinity will actually
reduce protein noise, even though it increases bursting. **B, D)** Increasing mRNA
abundance, holding mRNA noise constant, monotonically decreases protein noise for all
parameter values, as both the reaction order and translational burst parameters decrease
with increasing mRNA (Sup. Mat. for theoretical derivation of this result). Parameters: e
= 5, k_{ON} =1, k_{OFF} = 19.5 (varied in **A)**, k_{cat} = 0.5, δ_S = 0.1, δ_P = 0.1, k_{in} = 4/b (varied in **B**).
K_m was tuned by varying k_{OFF}, s was tuned by varying k_{in}, while mRNA noise was held
constant with a burst input of either b =10 or 15.

388
389

Figure 3: **Effect of network architecture on the cost of noise.** Theoretical predictions
(Eqns. 6-8, S20, S21) for metabolic reactions (dashed lines; $h = 1$, $\delta_S = 0$) and translation
(solid lines; $h =0$, $\delta_S = 0.1$) compared to ensemble average (blue) and variance (red) of

1,000-5,000 SSA realizations (points). **A)** Noise-induced slowdown causes an

accumulation of substrate ("stochastic amplification") which completely buffers reaction

flux for metabolic reactions when substrate dilution/degradation is zero ($\delta_S = 0$) **(B)**.

Because mRNA is not consumed in translation ($h = 0$), its concentration is independent of

reaction rate so does not experience stochastic amplification. Thus, translation is not

buffered by network feedback. **C,D)** The effect of network architecture on substrate (C)

and product (D) noise. e = 1, k_{ON} =1, k_{OFF} = 19.5, k_{cat} = 0.5, , δ_P = 0.01, and k_{in} = 5/b

(solid lines) or 0.357/b (dashed lines).

The variance in protein numbers at steady-state is, $\langle \epsilon^2_{protein} \rangle \cong \bar{p}(1 +$

$a^2 b_T F_{mRNA} + CV^2_R \bar{p}/2)$, where the parameter $a^2 = [K_m/(K_m + m)]^2$ is the squared reaction

order and , $CV^2_{ribo} = \sigma^2_{ribo}/\bar{r}^2$ is the ribosomal noise (Sup. Mat.). The translational burst

parameter, b_T, is the number of proteins synthesized by a single mRNA molecule over its

lifetime, which with hyperbolic translation kinetics is, $b_T = k_T r/((K_m + m)\delta_S)$.

Previous work assumed first-order translation, such that $b_T = k_T r/(K_m \delta_S)$ (Thattai and

van Oudenaarden 2001; Shahrezaei and Swain 2008). Introducing non-linearity causes

the translational burst size to become a non-linear, decreasing function of mRNA

abundance: all else equal, an mRNA molecule competing with fewer other mRNA's will

be translated more often in its lifetime, increasing its burst size. Alternatively, in the

zeroth order regime ($m \gg K_m$), translational bursting goes to zero because ribosomes are

so oversaturated with mRNA's that any given mRNA is more likely to be degraded than

translated. The protein noise strength and noise, then, are, respectively,

415
$$F_{protein} \cong 1 + a^2 b_T F_{mRNA} + CV^2_{ribo} \bar{p}/2$$

416
$$CV^2_{protein} \cong \underbrace{\frac{1}{\bar{p}} + \frac{a^2 b_T F_{mRNA}}{\bar{p}}}_{\substack{intrinsic \\ noise}} + \underbrace{\frac{CV^2_{ribo}}{2}}_{\substack{extrinsic \\ noise}}$$
9

Eqn. 9 agrees well with simulation results (Figs. 2, 3) and retrieves the result of

Thattai and van Oudenaarden (Thattai and van Oudenaarden 2001) ($F_{protein} \cong 1 + b_T$)

as a special case when there is no transcriptional bursting ($b_t = 1 \Rightarrow F_{mRNA} = 1$), no

ribosome fluctuations ($CV^2_{ribo} = 0$), and translation is first order (i.e., linear) in mRNA

abundance ($a = 1$). The first terms in $F_{protein}$ and $CV^2_{protein}$ (1 and $1/\bar{p}$, respectively) are

simply Poisson terms reflecting low copy number fluctuations (Paulsson 2004; Pedraza

and Paulsson 2008), while the second and third terms introduce factors that increase noise

above Poisson levels. The second terms ($a^2 b_T F_{mRNA}$ and $a^2 b_T F_{mRNA}/\bar{p}$, respectively)

incorporate the consequences of translational and transcriptional bursting. These terms

propagate mRNA fluctuations, F_{mRNA}, to the protein level. Importantly, hyperbolic

translation kinetics ($0 < a < 1$) dampen the effect of mRNA fluctuations on protein noise,

thus attenuating the propagation of mRNA noise via a process that might be called

"hyperbolic filtering". Together the first and second terms are considered "intrinsic

noise" (Swain, Elowitz, and Siggia 2002; Raser and O'Shea 2005; Paulsson 2004; Bar-

Even et al. 2006; Sanchez and Golding 2013). The last term ($CV^2_{ribo}\bar{p}/2$ and $CV^2_{ribo}/2$)

accounts for fluctuations in ribosome numbers, which create positively correlated

fluctuations among protein species called "extrinsic noise" (Swain, Elowitz, and Siggia

2002; Raser and O'Shea 2005; Paulsson 2004; Bar-Even et al. 2006; Sanchez and

Golding 2013). This extrinsic noise term dominates for abundant proteins (large *p*),

consistent with the observation that protein noise scales inversely with protein abundance

for low copy number proteins, but is independent of abundance for high copy number

proteins (Blake et al. 2003; Raser and O'Shea 2004; Bar-Even et al. 2006).

Translation initiation in *E. coli* and yeast has a reaction order of $a \sim 0.1$-0.2 (3,

49), which is comfortably to the left of the protein noise peak at $a^* = 2/3$ (see Sup. Mat.;

Fig 2a). This implies that, contrary to predictions based on first-order translation

initiation models (Thattai and van Oudenaarden 2001; Raser and O'Shea 2005), evolving

or engineering increased binding affinity of mRNA to ribosomes (i.e., reduced

translational K_m) will actually lower protein noise, not increase it. This occurs because

reducing K_m has two complementary effects on protein noise: it increases translational

bursting, b_T, but reduces reaction order, $a,$ which attenuates noise via a phenomenon we

term "zeroth-order insensitivity". The latter effect is not accounted for in previous gene

expression models, but has implications for the evolutionary and synthetic targets of

noise-amelioration (see Discussion). Hyperbolic filtering also has practical implications

when making inferences from data. For example, translational burst size, b_T, can be

estimated from Eqn 9 with knowledge of the mRNA and protein statistics; however,

assuming first order reaction kinetics ($a = 1$) can lead to large underestimates of this

parameter from data. For example, for $a \sim 1/5$, assuming first order translation kinetics

can lead to a 25-fold underestimate of translational burst size.

Correlation between mRNA and protein noise with hyperbolic translational

filtering: We now quantify the effect of hyperbolic translation kinetics on the correlation

between mRNA and protein noise. Emprical work in yeast, E. coli and human cell lines

have found that there is no correlation between mRNA and protein levels in the cell, or

between mRNA and protein noise (Chen et al. 2002; Ghaemmaghami et al. 2003;

Taniguchi et al. 2010; Gygi et al. 1999). Here I show that this lack of correlation can be

explained, in part, by modeling translation as a hyperbolic reaction.

Assuming as before that mRNA lifetimes are much shorter than protein lifetimes

($\delta_{mRNA} \gg \delta_{protein}$), the covariance between mRNA and protein noise, , is found to be,

$\sigma_{mp} \cong ab_T\sigma^2_{mRNA}$. The linear regression coefficient, $\beta_{mp} \equiv \sigma_{mp}/\sigma^2_{mRNA}$ is then,

$\beta_{mp} \cong ab_T$. Both of these show that hyperbolic translation kinetics ($a < 1$) diminish the

statistical association between mRNA and proteins. This is most usefully illustrated by

the Pearson correlation coefficient, $\rho \equiv \sigma_{mp}/\sigma_{mRNA}\sigma_{protein}$, which is approximately

equal to,

470
$$\rho \cong a\sqrt{\left(\frac{\delta_{protein}}{\delta_{mRNA}}\right)\left(\frac{b_T F_{mRNA}}{1 + a^2 b_T F_{mRNA} + CV^2_{ribo}\bar{p}/2}\right)} \qquad 10$$

Consistent with the theoretical predictions of Tanaguchi et al (3), Eqn. 10 shows that the

mRNA-protein correlation is increased with increasing transcriptional noise strength,

F_{mRNA}, is diminished with larger differences in lifetimes of mRNA and protein

($\delta_{protein}/\delta_{mRNA} < 1$), and is diminished by extrinsic noise ($CV^2_{ribo}\bar{p}/2$). In addition, the

covariance, linear regression coefficient, and correlation between mRNA and protein

fluctuations all increase with increasing translational bursting, b_T. The most important

novel result here is that the correlation between mRNA noise and protein noise is

diminished by the hyperbolic filtering of translation (that is, when $a < 1$).

Network architecture, feedback, and buffering: While noise slows a reaction for a

given substrate concentration, network feedbacks cause adjustment to substrate levels

that may in turn alter reaction rates. Specifically, if a reaction proceeds more slowly, then

substrate levels accumulate, which then cause the reaction to speed up again. Accounting

for this feedback is crucial in quantifying the costs of noise. We now ask: How exactly

does this feedback influence the total flux through noisey reaction networks? To address

this problem, we now analyze how network architecture creates feedbacks that alter

chemical flux, potentially compensating for noise-induced slowdown. We treat three

limiting cases: 1) $h = 1$, $\delta_S = 0$; 2) $h = 1$, $\delta_S > 0$; and 3) $h = 0$.

Cases 1 ($h = 1$, $\delta_S = 0$): Stochastic substrate amplification and buffering: One

important feedback in noisy networks is the amplification of substrate concentration by

noise (Eqn. 6), which has been studied in a process called "stochastic focusing"

(Paulsson, Berg, and Ehrenberg 2000). Eqn. 6 shows that stochastic amplification will

occur whenever substrate is consumed by the reaction ($h = 1$), as in metabolic reactions.

Consequently, this steady-state adjustment of substrate creates intrinsic buffering of

reaction flux that ameliorates noise-induced slowdown: the reaction is slowed by noise,

allowing substrate to accumulate, which then speeds up the reaction. The extent of

buffering depends on the substrate dilution parameter, δ_S. In the MM network studied by

Grima (Grima 2010; Grima 2009b), for example, the substrate entered the reaction

volume at rate k_{in} and exited only once consumed by the focal reaction at rate V. Flux

balance requires that $V = k_{in}$ in such a system, so that the reaction rate in steady-state is

unaffected by noise, as found by Grima (Grima 2010; Grima 2009b). Thus, under the flux

constraint $V = k_{in}$, stochastic amplification completely buffers the system against noise-

induced slowdown (Fig 3).

Case 2 (h = 1, δ_S > 0): Incomplete amplification and buffering. However, the

precise balancing of V with k_{in} does not apply in living cells, where substrate exits the

system via multiple alternative channels (δ_S > 0), primarily dilution by cell division, but

also through excretion, titration by other reactions and enzymatic degradation.

Extracellular substrate also experiences non-zero δ_S, as it is diluted by bulk flow (such as

in a chemostat) or by uptake from neighboring cells. Accounting for these effects

modifies the flux constraint to: $k_{in} = V + \delta_S s$. With this network architecture, stochastic

amplification compensates for some of the diminution of V by noise, but not all.

Consequently, stochastic amplification fails to completely buffer metabolic reactions in

living cells against noise-induced slowdown.

Case 3 (h = 0, δ_S > 0): Gene expression is especially sensitive to noise-induced

slow-down; no stochastic amplification, no buffering. The $V = k_{in}$ flux constraint is

entirely abolished in template-mediated reactions ($h = 0$), such as protein synthesis via

translation of mRNA. Because mRNA is not consumed by translation, steady-state

mRNA concentration is set by the flux constraint $k_{in} = \delta_S s$, which is independent of

translation rate, V. Thus, stochastic amplification does not occur in translation and so

cannot buffer translation from mRNA noise (Fig. 3). As a consequence, gene expression

is unusually sensitive to noise.

Cell fitness. Noise slows biochemical reactions, but how does it affect fitness? We now

consider a reaction network representing a coarse-grained model of cell growth. We then

analyze this reaction network in the stochastic case using the method developed above in

order to quantify the effects of gene expression noise on the average rate of cell division.

During balanced exponential growth, it has been shown empirically that fitness,

here defined as the division rate of a cell, is equivalent to the total rate of protein

synthesis by translation in a cell (Shahrezaei and Marguerat 2015; Scott et al. 2010; Scott

et al. 2014). We denote the rate of cell division by the parameter λ. In general, biomass

synthesis involves the import of extracellular nutrients followed, ultimately, by their

conversion into new biomass via translation. Translation initiation involves the binding of

mRNA, in abundance m, to a ribosome, in abundance r, followed by elongation via the

polymerization addition of amino acids (aa) to a growing protein chain. Elongation rate is

limited by the intracellular nutrient, in abundance s_I, in scarcest supply (i.e., von Leibig's

Law of the Minimum (Droop 1974; De Baar 1994; Tilman 1982)), which is typically

either ATP or aa-charged tRNA's. In eutrophic conditions, where nutrients are saturating,

translation is initiation-limited (Shah et al. 2013), whereas in oligotrophic conditions,

where nutrients are scarce, translation is elongation-limited (Tuller et al. 2010).

This coarse-grained cell model follows a QSSA system of ODE's (S22) with

growth rate,

$$542 \qquad \frac{dp}{dt} = k_T r \frac{s_I m}{K_1 + K_2 s_I + K_3 m + s_I m} - \lambda p \qquad\qquad 11$$

which is equivalent to a compulsory-order two-substrate enzymatic reaction with

independent (i.e., non-cooperative) binding events (Ingalls 2013). k_T is the translation

elongation rate per mRNA per ribosome (which absorbs the number of S_I per protein),

K_1/ K_2/ K_3 are affinity constants, and λ is the protein dilution rate, which during steady-

state growth is equivalent to the rate of cell division. Eutrophic and oligotrophic regimes,

corresponding to initiation- and elongation-limited translation, respectively, are found

from this equation in the limit as s_I or m, respectively, goes to infinity. Eqn. 11 can be

rearranged to form a Monod equation, which maps growth rate to extracellular nutrient

supply with a number of practical and conceptual benefits (S24).

The expected steady-state rate of cell division (cells/time) is found by setting Eqn.

11 to zero and solving for λ,

$$\langle \lambda \rangle = \frac{\langle V \rangle}{p_{ss}}$$ 12

p_{ss} is defined as the species-specific total protein content of a cell (protein/cell), which is

assumed constant for a given species, V is the synthesis rate of proteins (protein/time).

Fitness with noise. Following the method above, we write down a QSSA reduced system

of equations for growth rate according to our coarse-grained cell model (S22), giving a

system in five equations describing the time evolution of mRNA, uptake proteins,

extracellular substrate, intracellular substrate and protein biomass. We assume that

mRNA and extracellular nutrients undergo a linear birth-death process, supplied with

burst inputs of size b_t and b_{SE} respectively, and ignore ribosome fluctuations.

Intracellular nutrients are consumed by protein synthesis, but mRNA is not. Expected

fitness, then, is (Sup. Mat.):

$$\langle \lambda \rangle \cong \frac{\bar{V}}{p_{ss}}\left[1 - \underbrace{\overbrace{\frac{\langle \epsilon_m^2 \rangle}{\bar{m}}\frac{A_1}{\Omega}}^{mRNA\ noise} - \overbrace{\frac{\langle \epsilon_{PU}^2 \rangle}{\bar{p}_U}\frac{A_2}{\Omega}}^{protein\ noise}}_{cost\ of\ gene\ expression\ noise} - \overbrace{\frac{\langle \epsilon_{SE}^2 \rangle}{\overline{s_E}}\frac{A_3}{\Omega}}^{extracellular\ noise}\right]$$ 13

Where the A_i's are positive constants given in the SI. In reality, the protein noise term

will contain contributions from all proteins involved in the pathway importing and

processing nutrients to their final form utilized in translation. The MCA model of Wang

and Zhang (Wang and Zhang 2011) then fits within this term. The extracellular noise

term quantifies the strength of selection favoring phenotypic "bet hedging" (Starrfelt and

Kokko 2012) strategies in response to environmental unpredictability.

Surprisingly, transcriptional noise directly reduces fitness. This occurs because

fluctuations in mRNA levels reduce the average translation rate. This is counter to the

typical view, which holds that transcriptional noise impinges on cell function only to the

extent that it propagates to generate protein noise. Unlike the cost of protein or

extracellular nutrient noise, which require a number of parameter estimates that are

highly context-dependent, the cost of mRNA noise is easily estimated from available data

if we assume that cells are growing in their eutrophic regime ($s_I \rightarrow$ infinity), where

growth rate is limited by translation initiation. Taking the eutrophic limit of Eqn 11

gives, $dp/dt \cong k_T rm/(K_m + m) - \lambda p$, which is simply a MM reaction with mRNA

serving as the substrate and ribosomes as the catalyst. Noting that fluctuations in total

ribosome number and mRNA number are uncorrelated, we find from equations 4, 7 and

9,

$$\langle \lambda \rangle \cong \frac{\bar{V}}{p_{ss}} \left[1 - \underbrace{\frac{(b_t+1)}{2\bar{m}} a(1-a)\Omega^{-1}}_{cost\ of\ trancriptional\ bursting} \right] \qquad 14$$

Where $\bar{V} = k_T rm/(K_m + m)$. The cost of transcriptional bursting vanishes with zeroth-

or first-order translation ($a = 0$ and 1, respectively) and/or as cell volume and/or mRNA

abundance become large.

During balanced exponential growth of *E. coli* in rich medium: $m \sim 1400$

molecules, $r \sim 21,000$ molecules, and $k_T \sim 21aa/s \sim 4.6$ proteins/min (given a mean

protein length of 275*aa*). The total protein count of a cell is $p_{ss} = 2.4 \times 10^6$ molecules.

During growth in nutrient rich medium, $(1 - a) \sim 0.85$ (Arava et al. 2003; Ingolia 2014;

Zenklusen, Larson, and Singer 2008). Applying these parameter estimates to $\bar{V}/p_{ss}$ gives

a doubling time for *E. coli* of ~30 minutes. Similar calculations with parameters for the

budding yeast *Saccharomyces cerevisiae (m ~* 15,000-60,000, *r ~* 243,000, k_T *~* 15*aa/s ~*

2 proteins/min (average protein length 450*aa*), $p_{ss} = 5x10^7$, *a* = .85) gives a doubling

time of ~120 minutes. Both values are in close agreement with experiments in nutrient

rich media, providing an important benchmark.

In an *E. coli* cell, according to Eqn. 14, a mutation causing one unit increase in the

total mRNA Fano factor results in an average loss of 7.5 proteins/min, 223 proteins/cell

division, and $1.5x10^{-4}$ cells/generation, such that selection against mRNA noise is more

than 10,000 times stronger than random drift in this species (the effective population size

determines the strength of random allele frequency fluctuations and is $N_e = 2x10^8$ in *E.*

coli). In contrast, the same mutation in yeast, which has 40-fold higher *m*, has a cost of

$2.1x10^{-6}$ cells/generation, which is only about 20 times stronger than random drift ($N_e =$

$1x10^7$ in *S. cerevisiae*). This illustrates how fitness magnifies small differences in

reaction rates by integrating over the whole cell and cell lifetime, and highlights the

importance of cell volume (or more directly, molecule number) in influencing the

efficacy of selection against noise.

Rapid decay of selection efficacy with cell volume. As we just showed, the cost of

noise depends on reaction volume. Because unicellular species differ over several orders

of magnitude in their cell volumes, it may be possible to use these two facts in order to

make predictions about taxonomic patterns of gene expression noise.

The fate of a mutation that modifies cellular noise depends on the relative

magnitude of selection and random genetic drift. Selection efficiently overcomes random

drift provided that $2N_e|s| > 1$ (Kimura 1957; Kimura 1962). The boundary at $2N_e|s| = 1$

defines the" drift barrier" separating efficient and inefficient selection regimes (Sung et

al. 2012; Lynch 2007b). The drift barrier acts as a kind of evolutionary attractor

preventing organismal perfection: recurrent deleterious mutations with effects smaller

than $2N_e|s|$ push species away from their adaptive optima, and selection is too weak to

push them back. Thus, the balance between recurrent mutation, drift and selection

establishes a suboptimal phenotypic steady-state. We now derive a scaling relationship

between cell volume and selection efficacy demonstrating that the drift boundary for

noise attenuation scales superlinearly with cell volume.

Selection strength scales inversely with cell size. Selection strength, s, is defined as the

difference in absolute fitness of mutant (*m*) and resident (*r*) alleles scaled by the mean

fitness of the population. For a rare mutant we have: $s = (\lambda_m - \lambda_r)/\lambda_r$. This is simply

the difference in the absolute number of daughter cells produced by the mutant per

resident doubling. For simplicity, we assume a haploid population and that the mutant

and resident genotypes differ only in transcriptional noise. When the only source of noise

is intrinsic mRNA fluctuations, the selection coefficient is approximately (see SI for

derivation),

$$s \cong \Delta F_{mRNA} \frac{K_m}{(K_m + m)^2} \Omega_c^{-1} = \Delta CV_{mRNA}^2 a(1 - a)\Omega_c^{-1}$$
    15

Ω_c denotes the volume of the reaction compartment. The sign of selection is determined

by $\Delta F = (F_r - F_m)$ or $\Delta CV^2 = (CV_r^2 - CV_m^2)$, the difference in Fano factors (noise

strength) or squared coefficients of variation (noise), respectively, between resident and

mutant alleles. A mutation that increases transcriptional noise will be deleterious (s < 0),

while a decrease in noise is advantageous (s>0). More generally, $s \cong \sum_i^n (\Delta F_i A_i) \Omega_{ci}^{-1}$,

where the sum is over all *n* sources of noise affected by the mutation, the A_i are constants,

and Ω_{ci} is the reaction volume of the i^{th} noise source. Note that selection cannot act on

noise levels in first or zeroth order reactions (*a* = 1 or 0, respectively), since noise does

not alter the mean in these cases. Noise will, however, alter the heritability (e.g.,

penetrance) of a trait (Wang and Zhang 2011), which will reduce the response to

selection, but we do not consider this effect here. Importantly, the strength of selection

against noise explicitly depends on reaction volume.

Drift strength scales positively with cell size. A species population size tends to scale

inversely with its body size: a species must divide a limited resource pool amongst its

members, resulting either in a large number of small individuals or a small number of

large ones (Damuth 1981; Damuth 1987). For multicellular species the empirical scaling

relationship between body mass and population size, known as "Damuth's Law", follows

a power law with a scaling exponent of -¾, likely reflecting the ¾ power scaling of body

size with energetic demands (Damuth 1981; Damuth 1987). However, in unicellular

species' metabolic rate scales either linearly (eukaryotes) or superlinearly (prokaryotes)

with cell mass (DeLong et al. 2010). Assuming that N_e scales linearly with *N*, that mass

scales linearly with cell volume (i.e., cells have constant density), then $N_e = B\Omega_w^{-g_{drift}}$,

where the subscript on Ω_w denotes whole cell volume and g_{drift} = [3/4, 5/4] is the scaling

exponent. The constant, *B,* is proportional to the population size of a species with unit

volume, and absorbs cell density as the proportionality constant between mass and

volume.

Combining forces: Selection efficacy scales superlinearly with cell size. Combining

these results implies that the efficacy of selection against noise is proportional to

$(\Omega_w^{-g_{drift}}\Omega_c^{-1})$. For translation, it is possible to establish a relationship between Ω_c and

Ω_w, allowing us to reduce the number of independent variables.

668   Protein synthesis occurs in the cytoplasm/cytosol, suggesting that reaction-

compartment volume may simply equal whole-cell volume ($\Omega_c \sim \Omega_w$) for translation. This

relationship appears to hold between *E. coli* and yeast (m = 1500 and 60,000,

respectively, with a 40 fold difference in cell volume) and within mammalian cells as

they change volume throughout the cell cycle (Kempe et al. 2015; Padovan-Merhar et al.

2015). However, Gillooly et al (Gillooly et al. 2005) found that RNA concentration scales

as body mass to the -1/4 power over a broad range of eukaryotes, corresponding to RNA

number scaling with mass to the ¾ power. Although unicells were underrepresented in

their sample, this agrees well with the scaling of mRNA between *E. coli* and the amoeba

Dictyostelium discoideum where m = 155,000 and cell size is approximately 500-1000

fold larger than *E. coli*, such that $\Omega_c \sim \Omega_w^{2/3} \ to \ \Omega_w^{3/4}$.

679   Thus, for translation, we assume that reaction volume scales with whole cell

volume according to, $\Omega_c = \Omega_w^{-g_{sel.}}$, where g_{sel} = [3/4, 1]. Therefore, the efficacy of

selection against mRNA noise can be written as an allometric scaling law:

$$N_e|s| \cong \Delta CV^2_{mRNA} a(1-a)B\Omega_w^{-g} \qquad\qquad 16$$

The superlinear scaling exponent, $g = g_{sel} + g_{drift} = [3/2, 2¼]$, makes selection against

noise extremely weak in large cells. For example, all else equal, selection against a

mutation that increases expression noise is thirty thousand to one million times stronger

in *E. coli* than in a species, such as the amoeba *Dictyostelium discoideum*, which is about

1000 times larger (Fig. 4).

Figure 4: **Power-law scaling of selection efficacy with cell volume. A,B)** The predicted
efficacy of selection, $N_e|s|$, for *E. coli*, *S. cerevisae* and *Dictyostelium disoicdeum*, as a
function of the jump size in noise caused by a mutation, $|\Delta F|$, for mutations acting at a
single locus (left-hand column) and mutations acting on the noise of the global
transcriptome (right-hand column). The mRNA concentrations and effective population
size (N_e) estimates used for the figure are found in Table S1 ($m = 1500$, 60,000 and
155,000, respectively, and ploidy-adjusted $N_e = 2x10^8$, $1x10^7$ and $1.1x10^5$, respectively).
Reaction order was assumed to be a = 0.8 for all three species. The "drift boundary" is
$2N_e|s| = 1$. **C,D)** The predicted noise jump size at the drift boundary as a function of cell
volume. The green region represents a range of values for the scaling exponent (-2 to -
3/2), reaction order (0.75-0.9), and in **C)** the gene number (3250-12,500). Above the
green region (or individual points) selection is strong, below mutations are effectively
neutral.

Solving for $|\Delta F|$ at the drift boundary ($2N_e|s| = 1$), gives the jump size in noise

strength "visible" to natural selection. If the only cost of transcriptional noise is reduced

translation, then the selectively impermissible jump sizes in transcriptional noise follows

the scaling relationship,

$$|\Delta F_{mRNA}{}^*| > b\Omega_w{}^g \qquad\qquad\qquad 17$$

where $b = (K_m + m)^2 / BK_m$. Mutations causing jumps smaller than this critical size are

"effectively neutral", and thus invisible to selection acting on translation rate. Adaptive

mutations that decrease noise by an amount less than this jump size cannot efficiently

spread, while deleterious mutations with effect size below this threshold cannot be

efficiently purged by selection.

Parameterizing our model with literature estimates of cell volume, effective

population size, genome size, ribosomal occupancy, and translation kinetics, we predict

that *E. coli* should be strongly selected to minimize intrinsic transcriptional noise simply

because this noise slows translation. Whereas the cost of transcriptional noise is too weak

per gene (on average) in *S. cerevisiae* or *Dictyostelium* for selection to maintain intrinsic

noise at the Poisson minimum if noise-induced slowdown of translation is the only cost

of mRNA noise. Fig 4 shows that in cells below about 5fL (most prokaryotes), selection

favors suppression of intrinsic transcriptional noise to its theoretical minimum defined by

Poisson statistics, even if the only cost of transcription noise is reduced translation rate.

But because of the superlinear scaling of selection efficacy, cells only 100 times larger

(most eukaryotes) cannot efficiently oppose intrinsic noise jumps even as large as $\Delta F =$

1000 (again, if this is the only selection pressure for noise-minimization acting).

Alternatively, jumps in global transcriptional noise (i.e., extrinsic noise) have a much

greater selective effect, meaning that traits affecting extrinsic noise, such as ribosome

fluctuations (Eqn 9), should be under strong selection for noise-amelioration even in

moderately sized species.

DISCUSSION

Biochemical reactions are non-linear, but are typically modeled as linear in stochastic

models of gene expression due to technical limitations (van Kampen 2007). Here we have

introduced a novel procedure for easily and more accurately incorporating non-linearity

into models of coupled stochastic chemical reactions. We have found that hyperbolic

reaction kinetics filter input noise in such a way that the average reaction rate is slowed

by noise, while attenuating noise propagation. These results have a number of

implications.

A drift barrier hypothesis for gene expression noise and robustness. Cell size places

severe constraints on the evolution of noise suppression. The magnitude of noise-induced

slowdown of Michaelis-Menten reactions is inversely proportional to the reaction

volume, which follows intuitively from the fact that, for a given molecular density,

greater volumes contain more molecules and thus generate less noise. Lynch (Lynch

2007b) has previously noted the "genomic perils of evolving large body size". Large

bodied species tend to be less efficient at purging deleterious mutations due to the

positive scaling of genetic drift strength with body size, leading to a number of broad

taxonomic patterns including greater genomic mutation rate, transposon abundance and

intron number in unicellular eukaryotes and metazoans relative to prokaryotes and

viruses (Lynch 2007b; Lynch 2007a; Sung et al. 2012). Unique to gene expression noise,

cell size affects not only drift but also fitness, resulting in a negative superlinear scaling

of selection efficacy with cell size (Eqns. 15-17). Unicellular species span more than four

orders of magnitude in cell volume (~0.1 μm^3 Mycoplasma to more than 1000 μm^3 in

amoebazoans and other protists), with a sharp taxonomic divide separating prokaryotes

and eukaryotes. "Typical" cell diameters are 0.5-5um for prokaryotes and 10-100um for

unicellular eukaryotes. Taking a conservative value of a 100-fold average volume

difference between prokaryotes and eukaryotes, this translates to a 1000-10,000-fold

difference in the average efficacy of selection against noise for non-compartmentalized

processes (Eqn. 16).

Of course, numerous other fitness effects of gene expression noise occur (Eqn. 13

and (Fraser et al. 2004; Wang and Zhang 2011; Lehner 2008; Raj and van Oudenaarden

2008; Balázsi, van Oudenaarden, and Collins 2011; Raser and O'Shea 2005)), but these

must all scale with cell (or compartment) volume, such that selection for noise-

amelioration will often be weaker in eukaryotes than prokaryotes. It is possible that the

greater transcriptional complexity in eukaryotic cells, which employ multisubunit

regulatory complexes, enhancer regions, heterochromatic gene silencing, nuclear

localization into "transcription factories" and other features that greatly increase the

burstiness of transcription, evolved only once greater cell size freed populations from the

selective constraint of noise minimization. The cause of the size disparity between

prokaryotes and eukaryotes is thought to be due to differences in cellular respiration:

prokaryotes synthesize ATP via the electron transport chain embedded in the cell's

plasma membrane, which requires a high surface area to volume ratio for the whole cell

in order to balance energy production and demand. Respiration via mitochondria free

eukaryotes from this constraint, allowing them to achieve large cell sizes without

bankrupting the cell's energy budget. Thus, the evolution of mitochondria may have

paved the way for the greater transcriptional sophistication of eukaryotes.

Translation attenuates the propagation of transcriptional noise: Gene expression

noise is generated at two levels: transcription and translation. While the molecular

mechanisms controlling noise at each of these levels are becoming increasingly well

understood, much remains unclear about the propagation of noise between levels. I have

shown here that hyperbolic translation kinetics greatly attenuate the propagation of

mRNA noise to the protein level. Given parameter estimates of translation kinetics in E.

coli and yeast, the results here imply that hyberbolic translation will reduce mRNA noise

by between 25 and 100 fold, substantially attenuating the propagation of mRNA noise to

the protein level. The hyperbolic filtering of mRNA noise by translation also manifests as

a substantial reduction in the correlation between mRNA and protein abundance (Eqn.

10). This result is consistent with data from *E. coli* showing that there is zero correlation

between a gene's mRNA noise and its protein noise (Tanaguchi et al 2010). We have

shown that hyperbolic translation kinetics can help explain this lack of correlation, since

the correlation coefficient between mRNA and protein noise is proportional to the

reaction order parameter, a, which is between 0.1 and 0.2 in E. coli, implying a large

reduction in the correlation. Geometrically, one can visualize this diminution of

correlation by observing the Michaelis-Menten rate curve: on the flat part of the curve, all

mRNA concentrations lead to the same rate of protein synthesis, thus creating a zero

correlation between mRNA and protein.

A major implication of these results is that the vast majority of all intrinsic protein

noise comes from translation, not transcription. Consequently, if protein noise, rather

than mRNA noise, determines the functional consequences of gene expression noise, then

natural selection on transcriptional noise will be extremely weak, and synthetic systems

desiring control of protein noise should target translation, not transcription. We have thus

come to two conclusions about transcriptional noise: 1) mRNA noise directly inhibits

growth by slowing translation and 2) very little mRNA noise propagates to the protein

level. These two predictions cast new light on the functional consequences of

transcriptional noise in living cells.

Survival of the flattest. We have shown that hyperbolic reactions, by filtering out high-

amplitude input fluctuations, cause a reduction in both signal (average output) and noise

(relative output variance) compared to first-order reactions. The strength of selection

favoring organismal robustness is determined by the former, while the target of selection

for increased robustness is determined by the latter. Both the strength of noise and its cost

are mediated by the reaction order, $a = K_m/(K_m + s)$ (Eqns. 6, 9), making this the obvious

target for noise-attenuation. Noise-induced slowdown is minimal for first and zeroth

order reactions ($a = 1, 0$, respectively), while noise itself is minimal only in zeroth order

($a = 0$). Indeed, zeroth order kinetics completely filter out super-Poisson intrinsic input

noise (Eqn. 9; Fig. 2), making the output entirely insensitive to input noise. We might call

this phenomenon "zeroth order insensitivity". Selection for robustness, then, should push

reactions towards their zeroth order regime. Geometrically, the zeroth order regime

corresponds to the flat part of the rate curve, such that adaptive robustness by this

mechanism promotes "survival of flattest", a term previously invoked to describe

selection for robustness via very different mechanisms (Wilke et al. 2001; Codoñer et al.

2006).

There are two targets of modification for reaction order minimization: increasing

substrate abundance, s, which moves the substrate distribution to the flatter part of the

rate curve, or reducing K_m, which shifts the rate curve itself. Importantly, these strategies

are not equivalent (Fig. 2). Decreasing K_m (while holding s constant) causes a unimodal

change in product noise because K_m has two complementary effects: it reduces reaction

order, a, which filters noise, but it also increases product burst size, b_T (Fig. 2).

Alternatively, increasing mRNA concentration (with K_m held constant) monotonically

decreases protein noise because both reaction order, a, and translational bursting, b_T,

decline. This strategy increases fitness monotonically, thus avoiding the mRNA/protein

noise trade-off. Importantly, increasing robustness by increasing substrate concentration

is different from increasing robustness by decreasing substrate noise. To highlight this,

one could force noise strength to stay constant while increasing mean substrate. The

result would be increased robustness without a decrease in noise strength. This strategy

undoubtedly generates other costs not accounted for in the current model, such as the

energetic cost of increased mRNA synthesis. Future work should explicitly examine the

evolutionary feasibility of cellular robustness via this mechanism in light of its potential

trade-offs.

Robustness via compartmentalization: The present theory provides new insight into the

recent argument that compartmentalization of reactions in eukaryotic cells is a strategy

for noise control (Stoeger, Battich, and Pelkmans 2016). If cell compartments are filled

with reactants via active transport by membrane transporters, then the resulting

hyperbolic filtering (described above) will attenuate copy number fluctuations of

reactants potentially down to the Poisson minimum, though not lower (e.g., Eqns 9 and

S21). But hyperbolic filtering also slows the average rate of transport, creating

inefficiency. In addition, reducing the reaction volume increases the relative size of

fluctuations (the noise), which slows the absolute average rate of reactions within the

compartment. Future work is necessary to determine exactly when the conflicting costs

and benefits of hyperbolic filtering favor reaction compartmentalization as a strategy for

cellular robustness. In eukaryotes, this takes on another dimension, because mRNA is

actively transported out of the nucleus and into the cytoplasm before translation. The

hyperbolic filtering of transcriptional noise at this level will greatly ameliorate

transcriptional noise even before mRNA is filtered by translation. Future work should

model the consequences of this serial transcriptional noise filtering in eukaryotes.

Metabolic flux. Wang and Zhang (Wang and Zhang 2011) used metabolic control

analysis (MCA) to study the effect of enzyme fluctuations on metabolic flux, concluding

that enzyme noise diminishes fitness in pathways with more than one enzyme. This result

arises from the hyperbolic relationship between enzyme concentration and steady state

flux in multi-enzyme metabolic pathways (Kacser and Burns 1973), and so, like the

present paper, is a straightforward consequence of nonlinear averaging. However, MCA

models assume first-order enzyme kinetics (Kacser and Burns 1973). In light of the

present results, it is not immediately clear if this assumption makes the results of (Wang

and Zhang 2011) an underestimate of the true cost of gene expression noise, because

hyperbolic rate laws cause noise-induced slowdown at each pathway step, or an

overestimate of cost, because hyperbolic rate laws attenuate noise at each step, thus

damping noise propagation through the pathway. Future work should distinguish between

these two possibilities.

The crucial assumption: All of the major results of this paper rely on the validity of

hyperbolic reaction kinetics as an accurate description of *in vivo* biochemical reactions.

This assumption is not guaranteed, especially for translation. Translation is a unique

biochemical reaction because a single substrate molecule (mRNA) is simultaneously

bound to multiple enzymes (ribosomes), and it is therefore easy to imagine that

translation kinetics may not be well-described by Michaelis-Menten reaction kinetics.

Therefore, the present theoretical results must remain in question until future empirical

work carefully measures the *in vivo* rate law of protein translation.

METHODS

All Monte Carlo simulations were run on the elementary (microscopic) system of

reactions (S17) using Gillespie's Exact Stochastic Simulation Algorithm (SSA) (Gillespie

1977) implemented in *Mathematica* v10.3.0.0 with the open source xSSAlite package,

freely available from the xCellerator project (http://www.xlr8r.info/SSA/). Each

realization of the simulation is a Markov Chain with irregular (exponentially distributed)

time intervals. For statistical analysis, each realization was regularized using the

TimeSeriesAggregate[] command. Mean[] and Variance[] commands of the regularized

time series were used to compute the moments of the Markov Chain. Data points in

figures 1,2 and 3 are ensemble averages or variances over 1000-10,000 simulated

realizations of the stochastic process; each realization was run for at least 10 times the

half life of S or P (half-life = $\ln(2)/\delta$), whichever was longer, and the value for that

realization was s or p during the final second of the simulation run.

References
895

Acar, Murat, Jerome T Mettetal, and Alexander van Oudenaarden. 2008. "Stochastic
Switching as a Survival Strategy in Fluctuating Environments." *Nature Genetics* 40
(4): 471–75.
Alemu, Elfalem Y, Joseph W Carl, Héctor Corrada Bravo, and Sridhar Hannenhalli.
2014. "Determinants of Expression Variability." *Nucleic Acids Research*.
doi:10.1093/nar/gkt1364.
Arava, Yoav, Yulei Wang, John D Storey, Chih Long Liu, Patrick O Brown, and Daniel
Herschlag. 2003. "Genome-Wide Analysis of mRNA Translation Profiles in
Saccharomyces Cerevisiae." *Proceedings of the National Academy of Sciences* 100
(7): 3889–94. doi:10.1073/pnas.0635171100.
Balázsi, Gábor, Alexander van Oudenaarden, and James J Collins. 2011. "Cellular
Decision Making and Biological Noise: From Microbes to Mammals." *Cell* 144 (6):
910–25. doi:http://dx.doi.org/10.1016/j.cell.2011.01.030.
Bar-Even, Arren, Johan Paulsson, Narendra Maheshri, Miri Carmi, Erin O'Shea, Yitzhak
Pilpel, and Naama Barkai. 2006. "Noise in Protein Expression Scales with Natural
Protein Abundance." *Nat Genet* 38 (6): 636–43.
doi:http://www.nature.com/ng/journal/v38/n6/suppinfo/ng1807_S1.html.
Batada, Nizar N, and Laurence D Hurst. 2007. "Evolution of Chromosome Organization
Driven by Selection for Reduced Gene Expression Noise." *Nat Genet* 39 (8). Nature
Publishing Group: 945–49.
doi:http://www.nature.com/ng/journal/v39/n8/suppinfo/ng2071_S1.html.
Berg, Otto G. 1978. "A Model for the Statistical Fluctuations of Protein Numbers in a
Microbial Population." *Journal of Theoretical Biology* 71 (4): 587–603.
doi:http://dx.doi.org/10.1016/0022-5193(78)90326-0.
Blake, William J, Mads Kaern, Charles R Cantor, and J J Collins. 2003. "Noise in
Eukaryotic Gene Expression." *Nature* 422 (6932): 633–37.
doi:http://www.nature.com/nature/journal/v422/n6932/suppinfo/nature01546_S1.ht
ml.
Briggs, George Edward, and John Burdon Sanderson Haldane. 1925. "A Note on the
Kinetics of Enzyme Action." *Biochemical Journal* 19 (2): 338.

Chen, Guoan, Tarek G Gharib, Chiang-Ching Huang, Jeremy M G Taylor, David E
Misek, Sharon L R Kardia, Thomas J Giordano, et al. 2002. "Discordant Protein and
mRNA Expression in Lung Adenocarcinomas." *Molecular & Cellular Proteomics* 1
(4): 304–13. doi:10.1074/mcp.M200008-MCP200 .
Codoñer, Francisco M, José-Antonio Darós, Ricard V Solé, and Santiago F Elena. 2006.
"The Fittest versus the Flattest: Experimental Confirmation of the Quasispecies
Effect with Subviral Pathogens." *PLoS Pathog* 2 (12): e136.
Csete, Marie E, and John C Doyle. 2002. "Reverse Engineering of Biological
Complexity." *Science* 295 (5560): 1664–69. doi:10.1126/science.1069981.
Damuth, John. 1981. "Population Density and Body Size in Mammals." *Nature* 290
(5808): 699–700. http://dx.doi.org/10.1038/290699a0.
———. 1987. "Interspecific Allometry of Population Density in Mammals and Other
Animals: The Independence of Body Mass and Population Energy-use." *Biological*
Journal of the Linnean Society 31 (3): 193–246.
Dar, Roy D, Brandon S Razooky, Abhyudai Singh, Thomas V Trimeloni, James M
McCollum, Chris D Cox, Michael L Simpson, and Leor S Weinberger. 2012.
"Transcriptional Burst Frequency and Burst Size Are Equally Modulated across the
Human Genome." *Proceedings of the National Academy of Sciences* 109 (43).
National Acad Sciences: 17454–59.
De Baar, H J W. 1994. "Von Liebig's Law of the Minimum and Plankton Ecology
(1899–1991)." *Progress in Oceanography* 33 (4): 347–86.
DeLong, John P, Jordan G Okie, Melanie E Moses, Richard M Sibly, and James H
Brown. 2010. "Shifts in Metabolic Scaling, Production, and Efficiency across Major
Evolutionary Transitions of Life." *Proceedings of the National Academy of Sciences*
107 (29): 12941–45. doi:10.1073/pnas.1007783107.
Draghi, Jeremy, and Michael Whitlock. 2015. "Robustness to Noise in Gene Expression
Evolves despite Epistatic Constraints in a Model of Gene Networks." *Evolution* 69
(9). Wiley Online Library: 2345–58.
Droop, M R. 1974. "The Nutrient Status of Algal Cells in Continuous Culture." *Journal*
of the Marine Biological Association of the United Kingdom 54 (04): 825–55.
Eldar, Avigdor, and Michael B Elowitz. 2010. "Functional Roles for Noise in Genetic
Circuits." *Nature* 467 (7312). Nature Publishing Group, a division of Macmillan
Publishers Limited. All Rights Reserved.: 167–73.
http://dx.doi.org/10.1038/nature09326.
Elf, J, and M Ehrenberg. 2003. "Fast Evaluation of Fluctuations in Biochemical
Networks with the Linear Noise Approximation." *Genome Research* 13 (11): 2475–
84. doi:10.1101/gr.1196503.
Elowitz, Michael B, Arnold J Levine, Eric D Siggia, and Peter S Swain. 2002.
"Stochastic Gene Expression in a Single Cell." *Science* 297 (5584): 1183–86.
doi:10.1126/science.1070919.
Fraser, Hunter B, Aaron E Hirsh, Guri Giaever, Jochen Kumm, and Michael B Eisen.
2004. "Noise Minimization in Eukaryotic Gene Expression." *PLoS Biol* 2 (6). Public
Library of Science: e137. doi:10.1371/journal.pbio.0020137.
Ghaemmaghami, Sina, Won-Ki Huh, Kiowa Bower, Russell W Howson, Archana Belle,
Noah Dephoure, Erin K O'Shea, and Jonathan S Weissman. 2003. "Global Analysis
of Protein Expression in Yeast." *Nature* 425 (6959): 737–41.

http://dx.doi.org/10.1038/nature02046.

Gillespie, Daniel T. 1977. "Exact Stochastic Simulation of Coupled Chemical
Reactions." *The Journal of Physical Chemistry* 81 (25): 2340–61.

Gillooly, James F, Andrew P Allen, James H Brown, James J Elser, Carlos Martinez del
Rio, Van M Savage, Geoffrey B West, William H Woodruff, and H Arthur Woods.
2005. "The Metabolic Basis of Whole-Organism RNA and Phosphorus Content."
Proceedings of the National Academy of Sciences of the United States of America
102 (33). National Acad Sciences: 11923–27.

Golding, Ido, Johan Paulsson, Scott M Zawilski, and Edward C Cox. 2005. "Real-Time
Kinetics of Gene Activity in Individual Bacteria." *Cell* 123 (6): 1025–36.
doi:http://dx.doi.org/10.1016/j.cell.2005.09.031.

Grima, R. 2009a. "Investigating the Robustness of the Classical Enzyme Kinetic
Equations in Small Intracellular Compartments." *Bmc Systems Biology* 3. doi:Artn
10110.1186/1752-0509-3-101.

———. 2009b. "Noise-Induced Breakdown of the Michaelis-Menten Equation in
Steady-State Conditions." *Physical Review Letters* 102 (21). doi:ARTN
21810310.1103/PhysRevLett.102.218103.

———. 2010. "An Effective Rate Equation Approach to Reaction Kinetics in Small
Volumes: Theory and Application to Biochemical Reactions in Nonequilibrium
Steady-State Conditions." *Journal of Chemical Physics* 133 (3). doi:Artn
03510110.1063/1.3454685.

Gygi, Steven P, Yvan Rochon, B Robert Franza, and Ruedi Aebersold. 1999.
"Correlation between Protein and mRNA Abundance in Yeast." *Molecular and*
Cellular Biology 19 (3). Am Soc Microbiol: 1720–30.

Ingalls, Brian P. 2013. *Mathematical Modeling in Systems Biology: An Introduction*.
Cambridge, Massachusetts: The MIT Press.

Ingolia, Nicholas T. 2014. "Ribosome Profiling: New Views of Translation, from Single
Codons to Genome Scale." *Nat Rev Genet* 15 (3). Nature Publishing Group, a
division of Macmillan Publishers Limited. All Rights Reserved.: 205–13.
doi:10.1038/nrg3645.

Kacser, H and, and JA34 Burns. 1973. "The Control of Flux." In *Symp. Soc. Exp. Biol.*,
27:65–104.

Kaern, Mads, Timothy C Elston, William J Blake, and James J Collins. 2005.
"Stochasticity in Gene Expression: From Theories to Phenotypes." *Nat Rev Genet* 6
(6): 451–64. http://dx.doi.org/10.1038/nrg1615.

Kempe, H, A Schwabe, F Crémazy, P J Verschure, and F J Bruggeman. 2015. "The
Volumes and Transcript Counts of Single Cells Reveal Concentration Homeostasis
and Capture Biological Noise." *Molecular Biology of the Cell* 26 (4): 797–804.
doi:10.1091/mbc.E14-08-1296.

Kimura, Motoo. 1957. "Some Problems of Stochastic Processes in Genetics." *The Annals*
of Mathematical Statistics 28 (4). Institute of Mathematical Statistics: 882–901.
http://www.jstor.org/stable/2237051.

———. 1962. "On the Probability of Fixation of Mutant Genes in a Population."
Genetics 47 (6): 713–19. http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1210364/.

Kussell, Edo, and Stanislas Leibler. 2005. "Phenotypic Diversity, Population Growth, and
Information in Fluctuating Environments." *Science* 309 (5743): 2075–78.

Lehner, Ben. 2008. "Selection to Minimise Noise in Living Systems and Its Implications
for the Evolution of Gene Expression." *Molecular Systems Biology* 4 (1).
doi:10.1038/msb.2008.11.
Lynch, Michael. 2007a. "The Frailty of Adaptive Hypotheses for the Origins of
Organismal Complexity." *Proceedings of the National Academy of Sciences* 104
(suppl 1): 8597–8604.
———. 2007b. *The Origins of Genome Complexity*. Sunderland, MA: Sinauer
Associates, Inc.
McAdams, Harley H., and Adam Arkin. 1997. "Stochastic Mechanisms in
Gene Expression." *Proceedings of the National Academy of Sciences* 94 (3): 814–
19. http://www.pnas.org/content/94/3/814.abstract.
Metzger, Brian P H, David C Yuan, Jonathan D Gruber, Fabien Duveau, and Patricia J
Wittkopp. 2015. "Selection on Noise Constrains Variation in a Eukaryotic
Promoter." *Nature* 521 (7552). Nature Publishing Group, a division of Macmillan
Publishers Limited. All Rights Reserved.: 344–47.
doi:10.1038/nature14244http://www.nature.com/nature/journal/v521/n7552/abs/natu
re14244.html#supplementary-information.
Michaelis, Leonor, and Maud L Menten. 1913. "Die Kinetik Der Invertinwirkung."
Biochem. Z 49 (333-369): 352.
Newman, John R S, Sina Ghaemmaghami, Jan Ihmels, David K Breslow, Matthew
Noble, Joseph L DeRisi, and Jonathan S Weissman. 2006. "Single-Cell Proteomic
Analysis of S. Cerevisiae Reveals the Architecture of Biological Noise." *Nature* 441
(7095): 840–46.
doi:http://www.nature.com/nature/journal/v441/n7095/suppinfo/nature04785_S1.ht
ml.
Padovan-Merhar, O, G P Nair, A G Biaesch, A Mayer, S Scarfone, S W Foley, A R Wu,
L S Churchman, A Singh, and A Raj. 2015. "Single Mammalian Cells Compensate
for Differences in Cellular Volume and DNA Copy Number through Independent
Global Transcriptional Mechanisms." *Molecular Cell* 58 (2): 339–52.
doi:10.1016/j.molcel.2015.03.005.
Paulsson, Johan. 2004. "Summing up the Noise in Gene Networks." *Nature* 427 (6973):
415–18.
doi:http://www.nature.com/nature/journal/v427/n6973/suppinfo/nature02257_S1.ht
ml.
Paulsson, Johan, Otto G Berg, and Måns Ehrenberg. 2000. "Stochastic Focusing:
Fluctuation-Enhanced Sensitivity of Intracellular Regulation." *Proceedings of the
National Academy of Sciences* 97 (13): 7148–53. doi:10.1073/pnas.110057697.
Pedraza, Juan M, and Johan Paulsson. 2008. "Effects of Molecular Memory and Bursting
on Fluctuations in Gene Expression." *Science* 319 (5861): 339–43.
doi:10.1126/science.1144331.
Raj, Arjun, Charles S Peskin, Daniel Tranchina, Diana Y Vargas, and Sanjay Tyagi.
2006. "Stochastic mRNA Synthesis in Mammalian Cells." *PLoS Biol* 4 (10).
doi:doi:10.1371/journal.pbio.0040309.
Raj, Arjun, and Alexander van Oudenaarden. 2008. "Nature, Nurture, or Chance:
Stochastic Gene Expression and Its Consequences." *Cell* 135 (2): 216–26.
doi:http://dx.doi.org/10.1016/j.cell.2008.09.050.

Rao, Christopher V, and Adam P Arkin. 2003. "Stochastic Chemical Kinetics and the Quasi-Steady-State Assumption: Application to the Gillespie Algorithm." *The Journal of Chemical Physics* 118 (11): 4999–5010.

Rao, Christopher V, Denise M Wolf, and Adam P Arkin. 2002. "Control, Exploitation and Tolerance of Intracellular Noise." *Nature* 420 (6912): 231–37. http://dx.doi.org/10.1038/nature01258.

Raser, Jonathan M, and Erin K O'Shea. 2004. "Control of Stochasticity in Eukaryotic Gene Expression." *Science* 304 (5678): 1811–14. doi:10.1126/science.1098641.

———. 2005. "Noise in Gene Expression: Origins, Consequences, and Control." *Science* 309 (5743): 2010–13. doi:10.1126/science.1105891.

Sanchez, Alvaro, and Ido Golding. 2013. "Genetic Determinants and Cellular Constraints in Noisy Gene Expression." *Science* 342 (6163): 1188–93. doi:10.1126/science.1242975.

Sanft, Kevin R, Daniel T Gillespie, and Linda R Petzold. 2011. "Legitimacy of the Stochastic Michaelis?? Menten Approximation." *Systems Biology, IET* 5 (1): 58–69.

Schwanhausser, Bjorn, Dorothea Busse, Na Li, Gunnar Dittmar, Johannes Schuchhardt, Jana Wolf, Wei Chen, and Matthias Selbach. 2011. "Global Quantification of Mammalian Gene Expression Control." *Nature* 473 (7347). Nature Publishing Group, a division of Macmillan Publishers Limited. All Rights Reserved.: 337–42. doi:http://www.nature.com/nature/journal/v473/n7347/abs/10.1038-nature10098-unlocked.html#supplementary-information.

Scott, Matthew, Carl W Gunderson, Eduard M Mateescu, Zhongge Zhang, and Terence Hwa. 2010. "Interdependence of Cell Growth and Gene Expression: Origins and Consequences." *Science* 330 (6007): 1099–1102.

Scott, Matthew, Stefan Klumpp, Eduard M Mateescu, and Terence Hwa. 2014. "Emergence of Robust Growth Laws from Optimal Regulation of Ribosome Synthesis." *Molecular Systems Biology* 10 (8): 747.

Shah, Premal, Yang Ding, Malwina Niemczyk, Grzegorz Kudla, and Joshua B Plotkin. 2013. "Rate-Limiting Steps in Yeast Protein Translation." *Cell* 153 (7): 1589–1601. doi:http://dx.doi.org/10.1016/j.cell.2013.05.049.

Shahrezaei, Vahid, and Samuel Marguerat. 2015. "Connecting Growth with Gene Expression: Of Noise and Numbers." *Current Opinion in Microbiology* 25: 127–35. doi:http://dx.doi.org/10.1016/j.mib.2015.05.012.

Shahrezaei, Vahid, and Peter S Swain. 2008. "Analytical Distributions for Stochastic Gene Expression." *Proceedings of the National Academy of Sciences* 105 (45): 17256–61. doi:10.1073/pnas.0803850105.

Starrfelt, Jostein, and Hanna Kokko. 2012. "Bet-Hedging—a Triple Trade-off between Means, Variances and Correlations." *Biological Reviews* 87 (3). Blackwell Publishing Ltd: 742–55. doi:10.1111/j.1469-185X.2012.00225.x.

Stoeger, Thomas, Nico Battich, and Lucas Pelkmans. 2016. "Passive Noise Filtering by Cellular Compartmentalization." *Cell* 164 (6): 1151–61. doi:http://dx.doi.org/10.1016/j.cell.2016.02.005.

Sung, Way, Matthew S Ackerman, Samuel F Miller, Thomas G Doak, and Michael Lynch. 2012. "Drift-Barrier Hypothesis and Mutation-Rate Evolution." *Proceedings of the National Academy of Sciences* 109 (45): 18488–92. doi:10.1073/pnas.1216223109.

Suter, David M, Nacho Molina, David Gatfield, Kim Schneider, Ueli Schibler, and Felix
Naef. 2011. "Mammalian Genes Are Transcribed with Widely Different Bursting
Kinetics." *Science* 332 (6028): 472–74.
http://science.sciencemag.org/content/332/6028/472.abstract.
Swain, Peter S, Michael B Elowitz, and Eric D Siggia. 2002. "Intrinsic and Extrinsic
Contributions to Stochasticity in Gene Expression." *Proceedings of the National
Academy of Sciences* 99 (20): 12795–800. doi:10.1073/pnas.162041399.
Taniguchi, Yuichi, Paul J Choi, Gene-Wei Li, Huiyi Chen, Mohan Babu, Jeremy Hearn,
Andrew Emili, and X Sunney Xie. 2010. "Quantifying E. Coli Proteome and
Transcriptome with Single-Molecule Sensitivity in Single Cells." *Science* 329
(5991): 533–38. doi:10.1126/science.1188308.
Thattai, Mukund, and Alexander van Oudenaarden. 2001. "Intrinsic Noise in Gene
Regulatory Networks." *Proceedings of the National Academy of Sciences* 98 (15):
8614–19. doi:10.1073/pnas.151588598.
Thattai, Mukund, and Alexander Van Oudenaarden. 2004. "Stochastic Gene Expression
in Fluctuating Environments." *Genetics* 167 (1): 523–30.
Tilman, D. 1982. *Resource Competition and Community Structure*. Princeton, NJ:
Princeton University Press.
Tuller, Tamir, Asaf Carmi, Kalin Vestsigian, Sivan Navon, Yuval Dorfan, John
Zaborske, Tao Pan, Orna Dahan, Itay Furman, and Yitzhak Pilpel. 2010. "An
Evolutionarily Conserved Mechanism for Controlling the Efficiency of Protein
Translation." *Cell* 141 (2): 344–54. doi:http://dx.doi.org/10.1016/j.cell.2010.03.031.
van Kampen, N G. 2007. *Stochastic Processes in Physics and Chemistry*. Third Edit.
New York: Elsevier.
Wagner, G, xfc, P nter, Ginger Booth, and Homayoun Bagheri-Chaichian. 1997. "A
Population Genetic Theory of Canalization." *Evolution* 51 (2). [Society for the
Study of Evolution, Wiley]: 329–47. doi:10.2307/2411105.
Wang, Zhi, and Jianzhi Zhang. 2011. "Impact of Gene Expression Noise on Organismal
Fitness and the Efficacy of Natural Selection." *Proceedings of the National
Academy of Sciences* 108 (16): E67–76. doi:10.1073/pnas.1100059108.
Wilke, Claus O, Jia Lan Wang, Charles Ofria, Richard E Lenski, and Christoph Adami.
2001. "Evolution of Digital Organisms at High Mutation Rates Leads to Survival of
the Flattest." *Nature* 412 (6844): 331–33.
Zenklusen, Daniel, Daniel R Larson, and Robert H Singer. 2008. "Single-RNA Counting
Reveals Alternative Modes of Gene Expression in Yeast." *Nat Struct Mol Biol* 15
(12). Nature Publishing Group: 1263–71.
doi:http://www.nature.com/nsmb/journal/v15/n12/suppinfo/nsmb.1514_S1.html.