DeepWAS: Directly integrating regulatory information into GWAS using

deep learning supports master regulator MEF2C as risk factor for major

depressive disorder

Gökcen Eraslan[*,1], Janine Arloth[*,1,2], Jade Martins[2], Stella Iurato[2], Darina Czamara[2], Elisabeth B. Binder[2,4], Fabian J.

Theis[1,3], Nikola S. Mueller[1,#]

[1] Institute of Computational Biology, Helmholtz Zentrum München Neuherberg 85764, Germany

[2] Department of Translational Research in Psychiatry, Max Planck Institute of Psychiatry, Munich 80804, Germany

[3] Department of Mathematics, Technische Universität München Garching 85748, Germany

[4] Department of Psychiatry and Behavioral Sciences, Emory University School of Medicine, Atlanta GA 30322, USA

[*] These authors contributed equally to this work.

[#]To whom the correspondence should be addressed: nikola.mueller@helmholtz-muenchen.de

Abstract

Background: Genome-wide association studies (GWAS) identify genetic variants

predictive of common diseases but this does not directly inform on molecular

mechanisms. The recently developed deep learning-based method DeepSEA uses

DNA sequences to predict regulatory effects for up to 1000 functional units, namely

regulatory elements and chromatin features in specific cell-types from the ENCODE

project.

Results: We here describe "DeepWAS", a conceptually new GWAS approach that

integrates these predictions to identify SNP sets per functional units prior to association

analysis based on multiple regression. To test the power of this approach, we use

genotype data from a major depressive disorder (MDD) case/control sample (total

N=1,537). DeepWAS identified 177 regulatory SNPs moderating 122 functional units.

MDD regulatory SNPs were located mostly in promoters, intronic and distal intergenic

regions and validated with public data. Blood regulatory SNPs were experimentally

annotated with methylation quantitative trait loci (QTLs), expression quantitative trait

methylation loci and expression QTLs and replicated in an independent cohort. Joint

integrative analysis of regulatory SNPs and the independently identified annotations

were connected through transcription factors MEF2A, MEF2C and ATF2, regulating a

network of transcripts previously linked to other psychiatric disorders. In the latest

GWAS for MDD, the *MEF2C* gene itself is within the top genome-wide significant locus.

Conclusions: DeepWAS is a novel concept with the power to directly identify individual

regulatory SNPs from genotypes. In a proof of concept study, MEF2C was identified as

a master-regulator in major depression, a finding complementary to recent depression

GWAS data, underlining the power of DeepWAS.

Keywords: GWAS concept, functional data, chromatin features, cell-type specificity,

ENCODE, deep learning, multiple regression, major depressive disorder, MEF2,

MEF2C

Background

Genome-wide association studies (GWAS) have been highly successful in identifying

genetic variants associated with risk for common diseases[1]. However, going from pure

association to mechanistic insight has been a much more challenging task. The

identification of the true functional/causal variants within association signals is

hampered by the linkage disequilibrium (LD) block structure of the genome. Due to this

structure, the true functional variants can most often not be discerned within associated

blocks, which can span several gene loci. One approach to better infer functional

variants has been the annotation of association signals by their correlation with more

proximal molecular read-outs such as gene expression or DNA methylation in the form

of expression and methylation quantitative trait locus (eQTL and meQTL)

approaches[2,3]. Very recent approaches such as binding QTL (bQTL) studies for

transcription factor (TF) binding now add additional regulatory information[4]. While

these approaches can indicate regulatory effects of associated SNPs, they can also not

pinpoint single functional variants within an LD block.

To further close the gap of missing functional annotation, additional post-processing

approaches, so called functional GWAS[5], have been used. The majority of the

previously published functional GWAS studies are based on the overlap of SNPs with

cis-regulatory elements such as promoters and enhancers (see Tak and Farnham[6] for

a comprehensive review). These functional GWAS indicate that for common disease,

the majority of associated SNPs reside in non-coding, regulatory regions[5,7,8]. One

drawback of these methods is that the actual impact of the variant on regulatory

elements is not assessed and thus not taken into account for the annotation, which

does not go beyond positional overlap. For example, two SNPs that localize to the same

chromatin immunoprecipitation with massively parallel sequencing[9] (ChIP-seq) peak

of a TF might have opposing effects or no effects at all. To try to resolve this, *in silico*

approaches predicting disruption of transcription factor binding motifs[10,11] have been

used, however, our understanding of actual binding based on known motifs is still

limited. Given the fact that classical GWAS thus test many variants that are highly

unlikely to be functionally relevant, methods integrating functional knowledge of SNP

into GWAS could not only allow prioritize relevant variants but also to increase power

for such association studies that now often need tens of thousands of cases for robust

signals[1].

Recent advances in systems genetics, harnessing the predictive power of deep

learning, might have the capacity to enhance the performance of functional SNP

prioritization methods. Zhou J and Troyanskaya OG have developed a deep learning

method called "DeepSEA" uses only DNA sequence information to predict effects on

regulatory chromatin features such as histone marks, TF binding or the presence of

open chromatin[12]. For this experimental, publicly available data from the ENCODE

project[13] and the Roadmap Epigenomics Project[14] for cell type-specific TF binding,

histone modifications and chromatin states was used. This type of functional sequence

annotation is superior to pure overlap methods as it computes allele-specific differences

in the effects on these regulatory elements and thus discerns SNPs with functional

impact, at least in the given cell lines, from those just residing within the annotated

element. Furthermore, it allows incorporating cell-type specific regulatory effects of such

variants, adding another critical layer to understand disease mechanisms that are often

tissue-specific.

In this manuscript we present a conceptually new approach fusing classical and

functional GWAS. We obtain regulatory information on SNPs by generating sets of

SNPs in "functional units" using *deep* learning and then performing functional unit-*W*ide

*A*ssociation *S*tudies (deepWAS). First, we extracted DNA sequences centred on a SNP

to predict close to 1000 allele-specific regulatory effects of chromatin features in various

cell types (a pair of one chromatin feature in one cell type is further called "functional

unit") using DeepSEA[12]. Second, the resulting significant regulatory SNPs were then

used to identify sets of SNPs characterized by their joint moderation of a functional unit.

Finally, we identify regulatory SNPs, short "deepSNPs", each coupled to a functional

unit by associating each set of predicted regulatory SNPs with a trait or disease. To that

end we use a multiple regression model with SNP selection using L1 ("LASSO")

penalization[15]. By testing regulatory SNPs within each of the confined functional units,

we controlled not only for correlations induced by LD but also for possible joint, thereby

again correlated, SNP effects on two or more chromatin features or cell types for which

LASSO would select one representative SNP. This optimized variant selection improves

our power to identify SNPs that may play a role in the etiology of the disease.

For a proof of concept, we used data from a published GWAS for major depressive

disorder (MDD)[16]. Heritability for this disorder has been reported to be up to 40%[17]

and thus comparable with other common diseases, GWAS have, however, only been

successful very recently when including over 110,000 cases and 300,000 controls

identifying 17 independent SNPs associated with MDD at genome-wide significant

level[18,19]. Nonetheless, the required sample size for each discovered disease locus is

much higher than for other psychiatric disorders such as schizophrenia, in which 128

independent SNPs have been identified with only 35,000 cases[20]. This difference may

be attributed to the possibly higher biological heterogeneity not covered with current

phenotyping methods and the large relevance of the environment as a risk factor for

MDD. A recent study from our group has reported the importance of genetic variants in

specific enhancer regions conferring risk for major depression, likely by gene x stress

interactions[21]. DeepWAS, with its focus on variants altering functional regulatory

elements, could thus help to promote our understanding of the genetic mechanisms

underlying depression risk.

127   Results

128   DeepWAS approach

To identify SNP sets that potentially affect common regulatory functions in a cell type

specific manner, we need to investigate their genomic location and sequence alterations

within so called "functional units". We denote a "functional unit" as a gene-regulatory

state of a specific cell type characterized by the mapping of one chromatin feature onto

the genome in the absence or presence of additional stimulating/treatment conditions,

see (**Figure 1A**). For example, the functional unit "p53-HeLa" was defined by all

genomic regions covered by ChIP-seq peaks of the TF p53 within baseline conditions in

the HeLa cell line. For this manuscript, the 919 DeepSEA features [12] are considered

as functional units representing data from the ENCODE project [13] covering

combinations of 201 different experimental annotations of epigenetically relevant

information. This includes the mapping of different TFs, various DNase hypersensitive

(DHS) regions and histone marks across 31 cell lines and 17 treatment conditions.

Next, one kilobase DNA sequences were generated for each SNP for both reference

and alternative alleles. We used the pre-trained DeepSEA model[12] to predict the

probabilities of a SNP allele to moderate a functional unit. DeepSEA is a method that

uses deep neural networks and was trained to predict membership of 919 functional

units given a one kilobase genomic sequence. To select SNPs moderating a functional

unit, we used the metric provided by DeepSEA that estimates the impact of a variant on

the functional read out by comparing the probabilities of allele-specific regulation[12].

Therefore, the authors used one million random SNPs from the 1,000 Genomes project

[22] as a background distribution to calculate e-values for each functional unit by

assessing the proportion of random variants with a bigger effect than that of the

observed variants. SNPs with significant e-values, thus potentially having allele-specific

effects on a functional unit, were defined as a functional-unit specific SNP set. In other

words, only SNPs with a predicted and significant difference in allele-specific regulatory

effect for a given genomic feature in a cell type (functional unit) were retained for the

subsequent association analysis. When overlapping the available genotypes (measured

and imputed) in the case/control samples with the functional-unit specific SNP sets, 919

sets of genotypes filtered by regulatory effect in a functional unit remained for the

analysis (**Figure 1B**).

For the final step in the DeepWAS, we used the functional genotypes of the case/control

sample and possible confounders to associate SNPs set to a disease or trait (**Figure**

1C). The number of tested SNPs is massively reduced using the above-described

functional annotation based on ENCODE data and this variant filtering now enables us

to employ regularized regression methods with multiple SNPs. Therefore, each set of

regulatory SNPs in one functional unit was subjected to a logistic regression model with

L1 regularization (LASSO). In other words, we fitted 919 functional unit-based models to

again select regulatory SNPs associated to a trait or disease. For models with at least

one non-zero coefficient for a SNP, we further implemented a permutation-based

significance test with controlled false positive rate. From all models that withstood this

permutation-based multiple testing correction, we finally identified the deepSNPs that

are defined as significant non-zero regulatory SNPs associated to the trait or disease

and moderating a functional unit. The advantage of DeepWAS over traditional GWAS

(**Figure 1D**) is thus twofold, 1) it includes putative regulatory mechanisms in the GWAS

analysis from the start and 2) it controls false discovery error by reducing multiple

testing.

DeepWAS of major depressive disorder

We used post-quality control genotype data from 1,537 individuals (739 controls and

798 cases) of a cohort recruited for recurrent major depressive disorder (MDD). The

data were imputed to the 1000 Genomes Phase 1 reference panel [22] using SHAPEIT2

[23] and IMPUTE2 [24]. The resulting data set contained more than 1.7 million variants

with MAFs of at least 5% and non-missing genotypes. As expected, a classical GWAS

approach failed to identify genome-wide significant associations [16] (**Additional file 1:**

Figure S1).

For the DeepWAS approach, a total, 6,685 SNPs and 50,704 SNP-functional unit pairs

were retained after filtered for allele-specific effects in a given functional unit at a

significance level of $5 \cdot 10^{-5}$. These SNPs were then tested for association with

depression within 919 LASSO models for each functional unit (see above) including age

and sex as additional covariates (number of SNP predictors in 919 LASSO models:

min=17, median=51, max=213. See **Additional file 1: Figure S2**). Of these models,

193 had at least one non-zero coefficient for a SNP (number of SNPs with non-zero

coefficients in 193 models: min=1, median=2 and max=22). Permutation-based

significance test identified 122 out of these 193 models to be below the FDR-adjusted p-

value threshold of 0.1 (**Additional file 1: Figure S2** and **Additional file 2 and 3**). In

other words, deepWAS identified 122 functional units associated with MDD containing

in total 177 deepSNPs moderating 74 chromatin features in 31 cell lines.

We first analysed, whether the MDD deepSNPs were enriched for specific chromatin

features or tissue (**Figure 2A**). Not surprisingly, cell lines with a large number of

investigated chromatin features such as GM12878, K562 and HepG2 exhibited more

significant models since more chromatin features are covered in the data generated

from these cell lines. Interestingly, TFs such as c-Myc and EBF1 that are more relevant

to cell proliferation did not show significant associations. CTCF on the other hand, an

transcriptional regulator relevant for chromatin conformations important for the function

of long-range enhancers showed consistent effects across many cell lines. In fact,

genetic variants in long range enhancer have been shown to be highly relevant for

common disorders [14] and also major depression [21]. Furthermore, NF-κB and

glucocorticoid response element related deepSNPs only become apparent in the

stimulation condition, underlining the importance of going beyond pure baseline

characterization. Both systems and transcription factors, NF-κB – a major immune

regulator and the glucocorticoid receptor – central for the stress response, have been

previously implicated in the pathogenesis of MDD [21,25,26] (**Figures 2A and**

Additional file 1: Figure S2).

We next investigated the genomic regions in which these deepSNPs reside. Using the

UCSC knownGene annotations [27], we observed that vast majority of deepSNPs are in

promoters, intronic regions and distal intergenic regions that might indicate enhancers

(**Figure 2B**). We then examined the enrichment of deepSNPs in tissue-specific *cis-*

regulatory elements using genome annotations of the 15-state ChromHMM model[14].

The promoters are significantly enriched for deepSNPs in 126 out of 127 epigenomes

(99%) spanning all tissue groups, whereas the enhancers are only enriched in

"Astrocytes Primary Cells - E125" (FDR-adjusted p-value: 0.093) and in "A549 EtOH

0.02pct Lung Carcinoma Cell Line - E114" (FDR-adjusted p-value: 0.006) epigenomes

(**Additional file 1: Figure S4**).

To test whether the deepSNPs could be validated for their predicted allele-specific

effects in experimental data, we evaluated the effect of cell line-matched deepSNPs on

TF binding using previously published binding QTLs (bQTL) [4]. The bQTLs were

identified in GM12878 (Lymphoblastic cell line (LCL) from B-Lymphocytes) using a

pooled ChIP-seq approach of 60 Lymphoblastic Yoruban cell lines to identify SNPs

significantly altering binding of TFs by comparing the allele frequencies in sequencing

data before and after the ChIP experiment. Out of 26 deepSNPs in GM12878, we

identified eight SNPs (31 %) that were also experimentally validated bQTLs for JunD,

Pu.1, NF-κB and POU2f1 (**Additional file 4**).

237 Blood deepSNPs overlap with methylation and expression QTLs

Allele-specific effects on chromatin features likely associate with altered transcriptional

activation and this is reflected in differences in DNA methylation levels and gene

transcription. To test, whether the deepSNPs are indeed preferentially transcriptionally

relevant variants, we used DNA methylation and mRNA expression data from peripheral

blood in the recMDD samples (see Methods) as well as a second independent sample

(MPIP, see Methods). We focused on the detailed analysis of 26 GM12878 deepSNPs

(**Figure 3**), as GM12878 is among the best-characterized cell lines, derived from blood

cells, B-lymphocytes and amongst the top regulatory cell line identified by the MDD

deepWAS.

First, we conducted a methylation quantitative trait locus (meQTL) analysis using whole

blood methylation levels of a subset of the recMDD cohort (n=257 individuals). SNP-

CpG associations were carried out in a 3Mb *cis*-window around the deepSNPs to

assess the effect of nearby CpG sites. From the 26 deepSNPs, we identified 17

significant meQTLs containing seven unique deepSNPs (27 %) and 17 CpGs after

Bonferroni-correction for all tested CpGs per SNP (**Table 1**). The majority (>71%) of the

meQTL SNPs were associated with more than one DNA methylation site (one to four

CpG sites per SNP, **Table 1**). Notably, 25 deepSNPs (96 %) were nominally associated

with differential methylation levels ($p<0.05$). The most significant meQTL is located in an

enhancer region and all four identified CpGs in regions epigenetically associate with

gene silencing (**Figure 4A**). The genotypes at the intergenic SNP rs9293528 were

significantly associated with the methylation levels of four moderately correlated CpGs

located approximately 30-55 kb away from the SNP: cg1193710, cg14617041,

cg14727987 and cg26653990 (**Figures 4B** and **4C**). Sixteen out of the 17 meQTLs

were validated in the independent cohort (MPIP cohort, n=229 samples, see Methods),

i.e., showed a significant meta-analysis p-value (see Methods) showing equally or more

significant associations than in the discovery cohort alone (**Figure 4D** and **Table 1**).

Interestingly, deepSNPs are also associated with MDD-specific methylation changes. In

total we found five deepSNP-CpG pairs with significant differences in the allele-specific

methylation profiles between MDD cases and controls in recMDD cohort (see Methods).

Two of these MDD specific meQTLs are replicated in the independent MPIP cohort (see

Additional file 1: Table S1 and **Additional file 1: Figure S5**).

Second, we investigated the effects of the identified 17 CpG methylation sites

associated to MDD deepSNPs on matched gene expression levels, if CpG and

transcripts were within 1.5Mb of genomic distance. This was investigated in the MPIP

cohort, as gene expression data were only available in this sample. Among these, we

identified three expression quantitative methylation loci (eQTM) comprising two CpGs

and three gene expression probes after genome-wide Bonferroni-correction for the

number of tested transcripts per CpG (**Additional file 1: Table S2**) and 11 eQTMs (8

CpGs and 9 transcripts) at nominal level. These are: cg19235974-*CACNA2D4 (Calcium*

voltage-gated channel auxiliary subunit alpha2delta 4), cg19235974-*CACNA1C (Calcium*

voltage-gated channel subunit alpha1 C) and cg21290162-*MSRA (Methionine sulfoxide*

reductase A). *CACNA1C* has been associated with schizophrenia, bipolar disorder and

major depression[28], *CACNA2D4* with late onset bipolar disorder[29] and *MRSA* with

both bipolar disorder and schizophrenia [30,31].

Finally, we associated the 26 deepSNPs to gene expression directly, by conducting an

expression quantitative trait locus (eQTL) analysis using the MPIP cohort data (n=289

samples). We examined transcripts within a *cis*-window of 1.5Mb upstream and

downstream of the deepSNPs for an association with whole blood gene expression

profiles. We identified three deepSNPs with significant eQTL: rs12541159-*MSRA*,

rs1868881-*TIMM10* and rs4646797-*ALDH3A2* (**Additional file 1: Table S3**).

Interestingly, at nominal significance level we found 27 eQTLs (12 deepSNPs and 27

transcripts).

Overall, the investigation of the functional consequences of a hematopoietic tissue-

relevant set of deepSNPs in experimental data indicates that these indeed preferentially

tag meQTLs and eQTLs and are thus transcriptionally relevant.

Predicted regulatory effects of deepSNPs in MDD

Finally, we integrated the analysis results of meQTLs, eQTLs, eQTMs in peripheral

blood with our GM12878 deeSNPs to explore power of the deepWAS approach for

identifying novel, putative disease mechanisms. We combined all pairwise links of

meQTL (SNP-CpG), eQTL (SNP-gene), eQTM (CpG-gene) and deepSNP-chromatin

features and illustrated them in a network (**Figure 5A**).

One of six deepSNPs, rs12541159, had a meQTL and eQTL and at the same time

harboured an eQTM, in which the CpG site was affecting the transcriptional levels of the

same gene (**Figure 5B**). By regulating the methylation and expression profiles of

multiple genes via common TFs, deepSNPs may represent putative master regulators

of in a given disease. The intergenic deepSNP rs12541159 was predicted to affect

binding of the MEF2C TF, correlated with blood methylation levels of two intergenic

CpGs (cg21290162 cg11269159) and one exonic CpG cg27411982 with blood gene

expression of *MSRA* resigning 517 kb (**Figure 5B,** box 1 and **5C**). The CpGs

themselves showed no correlation to each other (**Figure 5D).** The transcriptional levels

of *MSRA* were associated with increased cg21290162 methylation (**Figure 5E**)*.* This,

motivated us to infer causality and we investigated whether the CpG site could fully

explain the observed association between the SNP and *MSRA* expression using causal

inference testing[32]. We found partial mediation of the effect of rs12541159 on *MSRA*

expression by DNA methylation status of CpG site cg21290162 (**Figure 5E**, p = 0.043).

Strikingly, the deepWAS results connected the independently identified six SNPs of the

nine molecular QTLs (see Figure 5A) through a family of TFs, namely MEF2A (*myocyte*

enhancer factor 2A), MEF2C (*myocyte enhancer factor 2C*) and ATF2 (*activating*

transcription factor 2). Predicted regulatory effect to identical SNPs suggest TF co-

localization, which has been not been reported so far. We thus utilized ReMap

annotation tool [33] to study co-localization and identified a significantly overlap of TF

ChIP-seq peaks for these three TFs (**Additional file 1: Figure S6**). Interestingly, a

SNP in the locus encompassing the MEF2C gene is the top locus of the largest GWAS

for MDD so far, with a $p < 10^{-16}$[19]. Similarly, we identify a regulatory effect of a second

TF, P300, of which the respective EB300 gene was a GWAS locus in the discovery

cohort[19].

Discussion

In classical GWAS all SNPs are tested independently and in a fully genome-wide

manner, thereby implicitly assuming that any SNP can moderate the function of any cell

state at any time. It is now clear that disease associations especially to common

disorders are driven by SNPs altering the function of regulatory elements. Hence it is

likely not necessary to test all SNPs in GWAS but instead useful to focus on functional

annotation to help prioritize putative risk variants[14].

In this manuscript, we address this idea by directly integrating the functional data into

the GWAS approach itself. To that end, we employed the powerful DeepSEA method

[12] that uses raw DNA sequence data to predict regulatory effects of 919 chromatin

features (from the ENCODE project) in various cell types, termed "functional unit" here.

These predictions enable determining allele specific effects of single SNPs by

comparing predictions of reference and alternative allele sequences (**Figure 1**). As

deep learning refers to the idea of learning multiple levels of representations and

relevant features from the raw input, it eliminates the need for manual feature

extraction, such as extracting k-mer frequencies for DNA sequence analysis [34]. To our

best knowledge, our study is the first to use deep learning-based predictors to identify

regulatory SNPs in GWAS. All identified deepSNPs were predicted to have a specific

regulatory effect of a chromatin feature in a defined cell type (functional unit), thereby

allowing having directly testable hypotheses of regulatory mechanisms in a trait or

disease.

Our deepWAS approach is superior to simply overlapping peaks or tracks of chromatin

features to GWAS signals, even though it is computationally more expensive due to the

deep learning-based predictions, cross-validated LASSO models and permutation-

based model selection. In contrast to annotation-based methods, data-driven methods,

such as DeepSEA, have a high predictive power on a single base resolution (by

comparing predictions of reference and alternative allele sequences) and not only for

larger stretches of sequence underlying one or several chromatin features. It thus

enables us to systematically assess the effects of sub-threshold SNPs that are missed

in classical approaches. DeepWAS does not identify genomic loci containing many

possible regulatory SNPs, but identifies single deepSNPs with predicted allele-specific

regulatory effect in a function unit, a cell-type and chromatin feature.

The second major methodological advantage of deepWAS is the implementation of

multiple (multi-SNP) regression models with L1 regularization (called LASSO

regression) selecting only a few SNPs, so-called deepSNPs. The deepSNPs of one

functional unit are only selected when they jointly associate with the disease or trait.

Multiple regression models are relatively new in the field of GWAS, nonetheless already

show promising result [35,36]. A related multi-SNP approach used pre-clustering of LD

blocks prior to LASSO modelling [37]. Models with millions of SNPs are less powerful to

associate SNPs, most probably due to the high correlation structure induced by LD. The

power of a pre-selection of regulatory SNP sets is underlined by the fact that when all

predicted regulatory SNPs (neglecting the information on functional units) were used as

input to a single LASSO regression model only one SNP (rs8180478 in osteoblast-

H2A.Z functional unit) showed significant association with MDD. This is in contrast to

the 177 deepSNPs identified using the pre-selection models.

Currently deepWAS has two main limitations. First, the proposed study was designed

for best-guess genotype data (i.e. 0-1-2 encoding). Since LASSO models require full

genotype information for all samples, the numbers of SNPs entering the analysis had to

be reduced due to quality control procedures. To address this issue, we are planning to

extend the method for dosage data (i.e. probabilistic representation of genotypes) to

increase the range of predicted regulatory effects. This will also necessitate as re-

evaluation of the LASSO model as many more SNPs in high LD will enter the analysis.

The second limitation is the current comprehensiveness of regulatory element

catalogues like ENCODE and Roadmap. ENCODE lacks for example a number of

relevant disease-specific stimulation conditions as well as disease-related tissues, in

our case brain tissues, which are important for psychiatric disease. Our previous and

related studies indicated the importance to test SNPs in stimulated conditions, like

TNFalpha and glucocorticoids receptor agonist dexamethasone conditions activating

NF-κB and GR, respectively [21]. Data from cell lines or bulk tissues will miss variants

with effects only in rare cell types or cell type specific effects in native tissue. It is

therefore important to be able to retrain the DeepSEA neural network with additional

publicly available chromatin features and as well as newly generated experimental data.

This will be possible using the deepWAS code publicly available at DOI:

10.5281/zenodo.59282. Additional data, from efforts such as the PsychENCODE will

make deepWAS even more powerful in the future.

The power of DeepWAS could be demonstrated in our small GWAS sample for

MDD[16]. Only using a subset of SNPs in functionally confined, regulatory units, we

identify risk variants for MDD even though the classical GWAS in this small sample was

negative (**Additional file 1: Figure S1**). The results in this small MDD sample illustrate

the power of deepWAS. 1) DeepSNPs were identified in cell types and enhancers

previously shown to be relevant for depression. Furthermore, using the stimulation

conditions, the importance of both immune system regulator NF-κB and the

glucocorticoid receptor system for depression was also observed in this study. Both

systems have been implicated by a large number of previous publications [21,25,26]. 2)

deepSNPs for MDD were highly likely to be functional in experimental data and impact

both DNA methylation and gene expression (**Figure 4** and **Table 1, Additional file 1:**

Table S2 and Table S3) as shown by their effects meQTL, eQTL and eQTM analyses.

Interestingly, for some of these, differential genotypic effects by major depression status

were observed (**Additional file 1: Figure S5** and **Table S1**). 3) The integrated analysis

of eQTL, meQTL and eQTM results with the GM12878 deepSNPs, identified the MEF2

TF family, including MEF2C, as important risk factors in MDD (**Figure 5**). The MEF2 TF

family was already identified to play a major role in neuronal plasticity, which is an

important component in disease development of stress-related disorders, like MDD and

other psychiatric disorders. Chen and colleagues[38] identified the TF MEF2 as a

master regulator of developmental metaplasticity, which is important to guide

developmental structural and functional neuronal plasticity. Additional evidence was

found by Barbosa et al [39]. Relating MEF2 to activity-dependent dendritic spine growth

and suggesting that this TF may suppress memory formation [39,40]. Most relevant for

this study, SNPs in the locus encoding MEF2C are the top signal in the latest meta-

analysis for major depression with over 130,000 MDD cases and 310,000 controls [19]

and minimum p-value of $9.99*10^{-16}$. Our study now identified SNPs altering the binding

of this TF to target transcript as relevant for MDD. This implies that not only SNPs in the

MEF2C locus itself, as seen in the meta-analysis for MDD [19] but also SNPs altering

it's binding in target transcripts, as observed in our deepWAS, are associated with major

depression. This shows that our approach focused on regulatory SNPs was able to

generate complementary functional data to the largest classical GWAS for MDD to date

suggesting that not only transcriptional regulation of MEF2C itself but also of its

downstream targets is relevant for MDD.

Our results supports the transcription factor MEF2C as a master regulators in MDD. In

fact we observed that transcripts affected by differential MEF regulation included a

network of genes previously implicated in psychiatric disorders. This includes

CACNA1C, a gene encoding a voltage gated calcium channel that has reported

genome-wide significant associations for schizophrenia as well as associations for

bipolar disorder and major depression in previous studies[28] as well as another

voltage-gated calcium channel, *CACNA2D4* that has been linked to late onset bipolar

disorder [29]. *MRSA* has been associated with both bipolar disorder and schizophrenia

[30,31] .

Conclusions

Our data indicate that deepWAS, a method combining classical GWAs with deep

learning based functional SNP annotation is a powerful tool to uncover disease

mechanisms for common disorders, including relevant cell types. It also allows to

directly identify functional SNPs by having a single base resolution and not being limited

by the LD structure of the locus. With ever increasing amounts of available functional

data the deepWAS approach will become even more valuable and allow to integrate

both publicly as well as unpublished data generated by individuals labs. DeepWAS is a

versatile tool, publicly available and together with available code from DeepSEA

applicable to any GWAS data.

Figures

Figure 1

460
Figure 1: **DeepWAS approach. (A)** The first step of the workflow corresponds to a

regulatory variant filtering procedure where only the SNPs with significant regulatory

effect are retained in the analysis. Prediction of significant regulatory variants is

performed using a pretrained deep neural network, namely DeepSEA, and e-values are

calculated by comparing the regulatory effects of observed SNPs to randomly selected

SNPs. **(B)** We define the functional genotype as the intersection of all SNPs assessed

(genotyped and imputed) in a sample and the preselected SNPs from DeepSEA. **(C)**

The association between disease and retained regulatory variants are interrogated via

regularized logistic regression. A regression model is fitted for each functional unit that

is define by tested TF/histone mark/DNase effects, the cell line and the specific

treatment e.g. NF-κB -GM12878-TNFa) where only the SNPs with a regulatory effect in

the specific cell type context are included as covariates. **(D)** The classical GWAS design

focuses on allele frequency differences and does not include information on the

regulatory impact of the tested variants.

Figure 2: **Significance of DeepWAS models and overlap of deepSNPs with**

functional annotations (A) Results of the significance test is given in heatmap. Every

functional unit is represented by the p-value, which shows the FDR-adjusted

significance of the corresponding LASSO model. Missing values, represented in white,

show functional units for which no data were available. The models of the functional

units with LASSO coefficients with p > 0.1 and those without any non-zero coefficients

are shown in grey and those with p < 0.1 in shades from yellow to red, with red

representing the most significant models. **(B)** Overlap of deepSNPs with genomic

elements is performed using UCSC knownGene annotations. Vast majority of

deepSNPs overlap with non-coding genomic regions.

Figure 3: Transcription factors harbouring deepSNPs in GM12878 cell line. In data

from GM12878 26 deepSNPs are predicted to affect the binding to 12 TFs. Group of

SNPs that are overlapping exclusively with a group of TFs are highlighted

Figure 4: **meQTL results for the GM12878 deepSNP rs9293528. (A)** Overview of the

rs9293528 meQTL locus on chromosome 5. Top panel, ideogram for chromosome 5. A

red box indicates the region shown enlarged in the bottom panels. Bottom panels:

Location of CpGs (n=4) that are significantly associated with rs9293528 genotypes,

SNP position, genes nearby (based on USCS knownGene annotation) and annotation

of chromatin states (based on GM12878 15-state core ChromHMM model). **(B)** Matrix

of Pearson's correlation coefficients between the DNAm levels (beta values) of

cg1193710, cg14617041, cg14727987 and cg26653990. **(C)** meQTLs between

rs9293528 and cg1193710, cg14617041, cg14727987 and cg26653990 within the

recMDD cohort. **(D)** Replication of those meQTLs in the MPIP cohort. Y-axis in C and D

displays the DNAm level of a particular CpG site and the x-axis indicates rs9293528

genotype.

Figure 5: Joint analysis of all significant meQTL, eQTL, eQTM and deepWAS

results: (A) Schematic illustration of our joint analysis. Grey box highlighting the MDD

disease effect. **(B)** Network visualization of the joint analysis. Edges represent the

association relation of deepSNPs, CpGs, TFs and transcripts. The grey circle highlights

the deepSNP rs12541159, which coincides all types of QTLs. **(C)** Regional plot for the

deepSNP rs12541159 locus: with top panel illustrating the ideogram for chromosome 8.

Bottom panels the location of CpGs (n=3) that are significantly associated with

rs12541159 genotypes, SNP position, gene expression probe position sig. association

with rs12541159, genes nearby (based on USCS knownGene annotation) and

annotation of chromatin states (based on GM12878 ChromHMM). **(D)** Matrix of

Pearson's correlation coefficients between the DNAm levels (beta values) of

cg27411982, cg11269159 and cg21290162. **(E)** Relationship between cg21290162

methylation, MSRA expression, and deepSNP rs12541159 genotype in MPIP samples,

including mediation effect rs12541159 → cg21290162 → MSRA expression.

Tables

CpG	SNP	Chr	SNP Position	SNP Location	Genes near SNP	CpG Position	CpG Location	Genes near CpG	P-value	Validation in MPIP
cg02138778	rs112478139	20	25110965	intergenic	VSX1,LOC284798	25172805	intergenic	LOC284798,ENTPD6	6.72e-06	yes
cg20318748	rs112478139	20	25110965	intergenic	VSX1,LOC284798	25605228	upstream	NANP	0.00306	yes
cg03677069	rs1240390	10	88725984	intergenic	SNCG,ADIRF	88718366	UTR5	SNCG	0.0148	yes
cg09552892	rs1240390	10	88725984	intergenic	SNCG,ADIRF	88718324	UTR5	SNCG	0.0229	yes

cg10376827	rs1240390	10	88725984	intergenic	SNCG,ADIRF	88730374	exonic	ADIRF	0.0142	yes
cg21012874	rs1240390	10	88725984	intergenic	SNCG,ADIRF	88718443	UTR5	SNCG	0.000177	yes
cg11269159	rs12541159	8	9769098	intergenic	MIR124-1,MSRA	9788795	intergenic	MIR124-1,MSRA	0.00265	yes
cg21290162	rs12541159	8	9769098	intergenic	MIR124-1,MSRA	9741960	intergenic	TNKS,LINC00599	0.000191	yes
cg27411982	rs12541159	8	9769098	intergenic	MIR124-1,MSRA	10470102	exonic	RP1L1	0.0105	no
cg14602222	rs12816718	12	969800	intronic	WNK1	1025664	exonic	RAD52	0.00226	yes
cg19235974	rs12816718	12	969800	intronic	WNK1	1063197	intergenic	RAD52,ERC1	0.0386	yes
cg10979063	rs2150175	6	97269213	intronic	GPR63	97285872	upstream	GPR63	0.000352	yes
cg09316306	rs5743565	4	38805983	UTR5	TLR1	38807387	upstream	TLR1	4.9e-05	yes
cg11937107	rs9293528	5	72456267	intergenic	TMEM171,TMEM174	72511061	intergenic	TMEM174,FOXD1	9.81e-10	yes
cg14617041	rs9293528	5	72456267	intergenic	TMEM171,TMEM174	72415688	upstream	TMEM171	0.00791	yes
cg14727987	rs9293528	5	72456267	intergenic	TMEM171,TMEM174	72485684	intergenic	TMEM174,FOXD1	6.44e-07	yes
cg26653990	rs9293528	5	72456267	intergenic	TMEM171,TMEM174	72510846	intergenic	TMEM174,FOXD1	0.0104	yes

524 **Table 1:** Significant GM12878 deepSNPs representing a meQTL (GSK cohort). P-values were

525 corrected according to Bonferroni.

526 Additional files

Additional file 1: Supplementary Figures S1-S6 and Tables S1-S3 (PDF)

Additional file 2: List of MDD deepWAS results. Table lists deepWAS results for MDD.

Triplet of the first three columns together form a functional unit. (XLS)

• Cell Line: Name of cell line of the functional unit

• Chromatin Feature: Name of chromatin feature of the functional unit

• Treatment: Name of Treatment of the cell line of the functional unit

• SNP: The deepSNPs jointly associated to MDD using deepWAS

Additional file 3: Significance of LASSO models with number of predictors in each

model (XLS)

• Cell line: Name of cell line of the functional unit

• Chromatin feature: Name of chromatin feature of the functional unit

• Treatment: Name of Treatment of the cell line of the functional unit

• pval.adj: FDR-adjusted p-values which represent the significance of a regression

model

• pval: p-values which represent the significance of a regression model

• devratio: Deviance ratio used as a test statistic for quantify the significance of

regression model

• n.snps: Number of SNPs that are used as predictors in regression models

• n.nonzero.snps: Number of predictors selected by LASSO approach (i.e.

covariates with non-zero coefficients)

Additional file 4: Overlap of bQTLs and MDD deepSNPs. Overlap of MDD deepSNPs

in cell line GM12878 with public bQTL study in same cells. (XLS)

• Chr: Chromosome number

• Pos.h19: Genomic position of SNP in hg19 assembly

• Rsid: SNP identifier

• bQTL.ref: Reference allele base extracted from bQTL study

• bQTL.alt: Alternative allele base extracted from bQTL study

• bQTL.pval: p-value extracted from bQTL study

• Higher Binding Allele: Base of higher binding allele extracted from bQTL study

• bQTL.TF: Transcription factor extracted from bQTL study

• deepwas_TF: Transcription factor identified using MDD deepWAS and rsid

Methods

Clinical Samples

recMDD

The sample consists of 1,537 Caucasian individuals (with 84%, n=1294 of German

origin) recruited at the Max-Planck Institute of Psychiatry (MPIP) in Munich, Germany

and two satellite hospitals in the Munich metropolitan area (BKH Augsburg and Klinikum

Ingolstadt): 739 controls (500 females, 239 males) and 798 cases diagnosed with

recurrent major depression (526 females, 272 males). Please see Muglia et al.[16] for

more detail in sample recruitment and characterization. All subjects are independent

from the MPIP collection below.

MPIP

The group of subjects consists of 289 Caucasian individuals of the Max-Planck Institute

of Psychiatry (MPIP), 93 women and 196 men. Recruitment strategies and further

characterization of the MPIP cohort have been described previously[21]. One hundred

sixty of them were healthy (115 men, 45 women), 129 (81 men, 48 women) had a

depressive disorder treated at the MPIP's hospital in Munich.

Genotype data and Imputation (GSK & MPIP)

Human DNA of the GSK and MPIP cohort samples was isolated from EDTA blood

samples using the Gentra Puregene Blood Kit (Qiagen) with standardized protocols.

Genome-wide SNP genotyping was performed using Illumina HumanHap550 Quad

(GSK), Illumina Human610-Quad (MPIP) and OmniExpress (MPIP) genotyping

BeadChips according to the manufacturer's standard protocols. Quality control (QC) of

genotyped data was conducted in PLINK 1.90b3s [20] separately for each cohort and

genotyping BeadChip. QC steps on samples included removal of individuals with

genotyping rate < 2 %, cryptic relatives (PI-HAT > 0.05), and genetic outliers (distance

in first two MDS components from mean > 2 SD). QC steps on variants included

removal of variants with call rate < 2 %, MAF < 5% and HWE test p-value ≤ 10-6.

Furthermore, variants on non-autosomal chromosomes were excluded, resulting in

GSK: 481,178, MPIP: 481,762 (Human610-Quad) and 544,908 (OmniExpress) SNPs.

These sets of SNPs comprised the input for imputation, which was performed

separately for each cohort and genotyping BeadChip using IMPUTE v2[24] with the

following parameters: 1000G phase I reference panel (released in June 2014, ALL

samples), SHAPEIT [23] phasing. QC of imputed probabilities was conducted in

QCTOOL 1.4 (http://www.well.ox.ac.uk/~gav/qctool/). Imputed SNPs were excluded if

the HWE test p-value ≤ 10^{-6} and the info metric < 0.8. Called genotypes ("best guess

genotypes'" with a probability of 70%) were used for further analysis. SNP sets of the

MPIP cohort were merged together. GSK and merged MPIP SNPs were further

processed in PLINK and variants were excluded if their MAF < 5% and best guess

genotypes are not called (call rate = 100%). This yielded a total of 1.78 Mio GSK SNPs

and 3 Mio MPIP SNPs. To annotate SNPs for the closest genes, we used Annovar

version Februar 2016 [41] with the UCSC knownGene annotations. SNP coordinates are

given according to hg19.

Prediction of regulatory effects

In this study, we employed DeepSEA[12] to determine the SNPs that might play an

important role in MDD by acting through the alteration of regulatory elements.

Pretrained DeepSEA network (v0.94) was downloaded from

http://deepsea.princeton.edu/help/ and the predictions of 1.7M SNPs were generated

using NVIDIA GeForce GTX TITAN X GPU (Maxwell) in 10 hours. Next step, filtering of

regulatory SNPs, was performed using generated so-called "e-value" files that represent

the significance of the regulatory effect of SNPs. We applied an e-value cutoff of

$5 \cdot 10^{-5}$ where we only take the SNPs that are associated to at least one functional unit

into consideration e.g. (rs1035271, GM12878, MEF2C). Set of SNPs that have an

impact on functional unit k is depicted here as set S_k.

We then employed a simple probabilistic genotype encoding where allele-specific

regulatory probability of the reference or alternative alleles is used if genotype is

reference or alternative homozygous, respectively. If the genotype of the individual is

heterozygous then the mean of two allele probabilities are used. Here, we refer to the

genotype matrix (1,537 x 1.7M) as G where the rows are individuals and columns are

SNPs. Genotypes in the G matrix is encoded such that $G_{ij} \in \{0, 1, 2\}$ where we simply

count an arbitrarily chosen allele. We then define D as a tensor representing the allele-

specific probabilities for each SNP, functional unit and allele. For example $D_{jk}^{(ref)}$ refers

to the probability of reference allele of SNP_j in the context of functional unit k. Therefore,

the SNP encoding scheme can be described in terms of G and D matrices:

$$X_{ijk} = \begin{cases} D_{jk}^{(ref)} & if\ G_{ij} = 0 \\ D_{jk}^{(alt)} & if\ G_{ij} = 2 \\ \dfrac{D_{jk}^{(ref)} + D_{jk}^{(alt)}}{2} & if\ G_{ij} = 1 \end{cases} \qquad G_{ij} \in \{0, 1, 2\}$$

or equivalently

$$X_{ijk} = G_{ij}\left(\frac{D_{jk}^{(alt)} - D_{jk}^{(ref)}}{2}\right) + D_{jk}^{(ref)}$$

Here the subscripts i, j and k represent individual, SNP, and functional unit indices,

respectively. Resulting encoded matrix is 1,537 x 6,143,515 where we store all SNP-

functional unit pairs (919 x 6,685) for all individuals using "bigmemory" R package[32] in

order to limit the memory consumption. During model fitting, however, only the list of

SNPs in sets S_k are used.

DeepWAS

Penalized regression models

Compared to the classical GWAS approaches where the trait of interest is regressed

separately on each SNP, regularized regression approaches provide an alternative way

to handle high dimensional data and to identify SNPs associated with the trait of interest

using variable selection. Here we utilize L1-regularized logistic regression (LASSO) for

variable selection. 919 different LASSO models are fitted for each functional unit in

order to estimate the statistical association between the disease status (y=0 for controls,

and 1 for cases) and SNPs in the context of a specific cell line:

$$logit\ P(y_i = 1) = \beta_{0k} + \sum_{j \in S_k} \beta_{jk} X_{ijk} + \beta_{sex,k} sex_i + \beta_{age,k} age_i$$

where k is used as a model index. Note also that the regression coefficients are model

specific and therefore indexed with k. In each LASSO model, only the SNPs that are

significantly affecting the binding of a specific TF in a specific cell line are included as

covariates. This is represented in the equation by the summation over the elements of

S_k.

Note that the simple probabilistic encoding does not affect the variable selection,

because the effect goes away during the standardization of the design matrix except for

possible sign flips. Yet, it leads to the effect sizes that are scaled based on the allele-

specific regulatory probabilities reported by DeepSEA since the effect sizes in logistic

regression are reported in the scale of original covariates.

The parameters (β) of the logistic regression are optimized with L1 regularization:

$$min_\beta \sum_{i=1}^{M} \left[y_i - \beta_{0k} - \sum_{j \in S_k} \beta_{jk} X_{ijk} + \beta_{sex,k} sex_i + \beta_{age,k} age_i \right]^2 + \lambda \|\beta\|_1$$

where M is the number of individuals, λ parameter represents the strength of the

regularization and β vector represents model parameters. We fitted L1-regularized

logistic regression model (LASSO) using glmnet R package[42]. Regularization

parameter λ is determined by 100-fold cross validation after which we used

"lambda.1se" value to determine non-zero parameters.

Permutation-based significance test

We used a permutation-based approach to select significant LASSO models with non-

zero coefficients and to control false discovery rate. We generated 1000 random

permutations of the response variable and fitted LASSO models on each. λ parameter

in each model is determined by 10-fold cross validation in order to reduce the

computational cost. After fitting LASSO models on the permuted data, deviance ratio is

used as a test statistic.

$$deviance\ ratio\ =\ 1 - \frac{D}{D_{null}}\ \text{where } D = 2\ (L_S\ -\ L_P) \text{ and } D_{null} = 2\ (L_S\ -\ L_N)$$

Here L_S, L_P and L_N are the log likelihoods of saturated model (model with a free

parameter per individual), proposed model and null model (intercept model),

respectively. Empirical p-values are calculated as the number of permuted models with

a deviance ratio greater than or equal to the proposed model divided by 1000. After

FDR[34] multiple testing correction with threshold of 10%, models with lower adjusted p-

values are retained.

Functional annotation of deepSNPs

Enrichment of Roadmap cis-regulatory elements

In order to quantify the enrichment of deepSNPs in *cis*-regulatory elements identified by

Roadmap Epigenomics Project[14], we first downloaded the segmentation files of core

15-state model for 127 epigenomes were downloaded from Roadmap epigenomics web

portal

(http://egg2.wustl.edu/roadmap/data/byFileType/chromhmmSegmentations/ChmmMode

ls/coreMarks/jointModel/final/all.mnemonics.bedFiles.tgz) in BED format. Then we

sampled random genomic ranges with the same size distribution as promoter (State 1)

and enhancer (State 7) regions from the background distribution using GAT[44]. The

background distribution was defined as the intersection of mappable regions of the

genome

(https://www.cgat.org/downloads/public/gat/datasets/hg19/mapability_36.filtered.bed.gz)

and genomic segments that belong to the first 13 chromatin states of ChromHMM

model (State 14: Weak repressed Polycomb and State 15: Quiescent are excluded). In

order to limit the computation time, 1000 samples were drawn and p-values smaller

than 10^{-3} were extrapolated from normal distribution using "--pvalue-method=norm"

command line option.

bQTL overlap

bQTL results were downloaded from "Supplementary Table 2" of Tehranchi et al.[4] and

intersection with GM12878 deepSNPs is computed in R version 3.3.0.

Transcription factor co-localization

ReMap project[33] provides a consolidated dataset for ChIP-seq peaks of several

transcription factors including the ENCODE peaks as well as various public datasets. In

order to quantify TF-TF colocalization patterns, we downloaded non-redundant peaks

for 11 individual TFs in BED format from the web interface (http://tagc.univ-

mrs.fr/remap/index.php?page=download) and then ran the annotation interface

(http://tagc.univ-mrs.fr/remap/index.php?page=annotation) for each BED file separately.

Downloaded TSV files with –log10 transformed e-values are combined and plotted with

corrplot R package[43]. Because very high –log10-transformed e-values obscure

visualization, e-values are binarized with a cutoff of 10^{-3}. Out of 12 TFs affected by 26

GM12878 deepSNPs 2 TFs are not shown in **Additional file 1: Figure S6** since p300

peaks are not available in ReMap dataset and CTCF did not show any significant

association with other TFs.

DNA methylation (GSK & MPIP)

For a subset of GSK (n=257 with 53% MDD cases) and MPIP cohort (n=229 with 52%

MDD cases) genomic DNA was extracted from whole blood using the Gentra Puregene

Blood Kit (QIAGEN). DNA quality and quantity of both was assessed by NanoDrop 2000

Spectrophotometer (Thermo Scientific) and Quant-iT Picogreen (Invitrogen). Genomic

DNA was bisulfite converted using the Zymo EZ-96 DNA Methylation Kit (Zymo

Research) and DNA methylation levels were assessed for >480,000 CpG sites using

the Illumina HumanMethylation450 BeadChip array. Hybridization and processing was

performed according to manufacturer's instructions. Quality control of methylation data,

including intensity read outs, filtering (detection p-value > 0.01 in at least 75% of the

samples), cellular composition estimation according to[44], as well as beta calculation

was done using the minfi Bioconductor R package version 1.18.2. Filtered beta values

were reduced by eliminating any CpG sites on sex chromosomes, as well as probes

found to have SNPs at the CpG site itself or in the single-base extension site with a

MAF ≥ 1% in the 1,000 Genomes Project European population and/or non-specific

binding probes according to[45]. Additionally, we performed a re-alignment of the array

probe sequences using Bismark[45,46]. This yielded a total of around 425,000 GSK and

MPIP CpG sites for further analysis. The data were then normalized with functional

normalization[47]. Technical batch effects were identified by inspecting the association

of the first principal components of the methylation levels with plate and plate position

for the GSK sample as well as processing (experiment) date for the MPIP sample as

technical batch. The data were then adjusted using ComBat[48]. To annotate CpGs for

the closest genes, we used Annovar version February 2016[41] with the UCSCknown

gene annotation. CpG coordinates are given according to hg19.

Gene expression (MPIP)

Whole blood RNA of the MPIP sample (n=289 individuals) was collected using

PAXgene Blood RNA Tubes (PreAnalytiX) and processed as described previously[21].

The RNA was then hybridized to Illumina HT-12 v3 and v4 expression Bead Chips

(Illumina, San Diego, CA). Raw probe intensities were exported using Illumina's

GenomeStudio and further statistical processing was carried out using R version 3.3.0.

All 29,075 probes present on both BeadChips (v3 vs. v4), excluding X and Y

chromosomes as well as cross-hybridizing probes identified by using the Re-Annotator

pipeline [21,49] were first filtered with a detection p-value of 0.05 in at least 50% of the

samples, leaving 11,994 autosomal expression array probes. Subsequently, each probe

was transformed and normalized through variance stabilization and normalization

(VSN)[50]. Technical batch effects were identified by inspecting the association of the

first principal components of the expression levels for all known batch effects and then

adjusted using ComBat with slide, amplification round, array version, and amplification

plate column as fixed effects. The position of the gene expression probe and gene

symbols were annotated using the Re-Annotator pipeline based on hg19. Blood cell

counts were estimated according to CellCODE [51].

Statistical analysis of gene expression and methylation data

Methylation QTL (meQTL) analysis

For each of the 26 deepSNP CpG sites within a 3Mb window around the SNP position

were selected for the meQTL analysis. Due to low frequency of minor allele

homozygotes for many SNPs, all SNP genotypes were coded and evaluated as a

dominant model. Linear regression was used to measure the relationship between the

DNA methylation (beta values) and the number of minor alleles (coded 0 and 1),

including covariates for age, disease-state, gender and blood cell counts. So we

defined the model for a single meQTL mapping as follows: $CpG = \beta_0 + \beta_{sex}sex +$

$\beta_{age}age + \beta_{CD8T}CD8T + \beta_{CD4T}CD4T + \beta_{NKcell}NKcell + \beta_{Bcell}Bcell + \beta_{Mono}Mono +$

$\beta_{Gran}Gran + \beta_{disesase\ status}disease\ status + \beta_{SNP}SNP + e,$ where e is the general

error term for any residual variation not explained by the rest of the model.

In addition a differential meQTL analysis by the disease status was performed.

Therefore an additive model adjusting for age, gender and blood cell counts was

applied in controls and cases separately:

$CpG = \beta_0 + \beta_{sex}sex + \beta_{age}age + \beta_{CD8T}CD8T + \beta_{CD4T}CD4T + \beta_{NKcell}NKcell +$

$\beta_{Bcell}Bcell + \beta_{Mono}Mono + \beta_{SNP}SNP + e.$

P-values were adjusted for the total number of CpG sites that were within the tested

region surrounding the SNP. Case specific meQTLs were defined as meQTLs, which

show a significant P-value in cases but not in controls (nominal p-value> 0.05). GSK

and MPIP data were analysed independently and a validation of meQTL results was

carried out with a sample size-weighted Z-score meta-analysis [52]. This method

accounts for different sizes and suggests the robustness the meQTLs.

Top hits were plotted with easyGgplot2 (https://github.com/kassambara/easyGgplot2),

corrplot R package [43] and Gviz 1.16.1 bioconductor package [53].

Expression QTL (eQTL) analysis

For each of deepSNP all transcripts beginning or ending within 1Mb up- or downstream

of the SNP were determined. Associations between genotype (coded 0, 1 and 2) and

expression levels (normalized and batch corrected log2 gene expression values) were

determined in the MPIP data set by linear regression, using sex, age, disease status

and blood cell estimates as covariates:

$GEX =$

$\beta_0 + \beta_{sex}sex + \beta_{age}age + \beta_{Neutrophil}Neutrophil + \beta_{Mono}Mono + \beta_{Bcell}Bcell +$

$\beta_{Tcell}Tcell + \beta_{NKcell}NKcell + \beta_{Plasmacell}Plasmacell + \beta_{DendriticCell}DendriticCell +$

$\beta_{diseas\ status}disease\ status + \beta_{SNP}SNP + e.$

To correct for multiple testing, p-values were corrected for the number of transcripts per

SNP.

Expression quantitative trait methylation (eQTM) analysis

For each of CpG site of the significant DeepQTL SNP- CpG pairs all transcripts

beginning or ending within 1Mb up- or downstream of the CpG were determined.

Associations between CpG and expression levels were determined in the MPIP data set

by linear regression, using sex, age, disease status and blood cell estimates as

covariates:

$CpG = \beta_0 + \beta_{sex}sex + \beta_{age}age + \beta_{CD8T}CD8T + \beta_{CD4T}CD4T + \beta_{NKcell}NKcell +$

$\beta_{Bcell}Bcell + \beta_{Mono}Mono + \beta_{Gran}Gran + \beta_{disesase\ status}disease\ status + \beta_{GEX}GEX +$

$e.$

To correct for multiple testing, p-values were first corrected for the number of transcripts

per SNP window.

Joint analysis

For the joint analysis we first measured the overlap of deepSNPs with eQTL, meQTL

data and secondly for all overlapping pairs we calculated the overlap of meQTL CpGs

and eQTM CpGs. For the deepSNPs, CpG, transcripts triplets we then assessed

possible causal relationships by using a causal inference test [32].

The network was plotted using the R package igraph [54]. The triplet was plotted using

ggplot2[55], corrplot R package [43] and Gviz 1.16.1 Bioconductor package [53].

Declarations

Availability of data and material

Data from MPIP gene expression experiment are deposited at the GEO repository under GEO: GSE64930 and methylation under: GSE74414. DeepWAS source code is available at DOI: 10.5281/zenodo.59282 through Zenodo.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was funded by the German Federal Ministry of Education and Research through the Research Consortium Integrated Network IntegraMent (grant 01ZX1314H, to GE), the LiSyM Verbundprojekt Pillar II/III (grant 031L0047, to NSM) under the auspices of the e:Med Programme as well as the European Research Council (grant 281338 to EBB)

Ethics approval and consent to participate

Both studies (recMDD and MPIP) were approved by the local ethics committee and all individuals gave written informed consent. All experimental methods comply with the Helsinki Declaration.

Authors' contributions

841 NSM conceived and designed the study. GE and JA performed the research, analysed

842 the data, developed the computational methods and created figures. GE, JA, EBB, FJT

843 and NSM wrote the paper. JM and DC performed imputation of recMDD and MPIP

844 genotypes. SI performed the pre-processing of the recMDD methylation data. EBB and

845 FJT contributed to the study design.

Acknowledgements

848 We thank Michael Laimighofer for feedback on the statistical approach. We thank

849 Matthias Heinig for valuable comments on the manuscript. We would like to thank

850 Martin Preusse for useful discussions on the approach.

References

1. Visscher PM, Brown MA, McCarthy MI, Yang J. Five Years of GWAS Discovery. Am.
J. Hum. Genet. 2012;90:7–24.

2. Gamazon ER, Wheeler HE, Shah KP, Mozaffari SV, Aquino-Michaels K, Carroll RJ,
et al. A gene-based association method for mapping traits using reference
transcriptome data. Nat. Genet. Nature Research; 2015;47:1091–.

3. Gamazon ER, Badner JA, Cheng L, Zhang C, Zhang D, Cox NJ, et al. Enrichment of
cis-regulatory gene expression SNPs and methylation quantitative trait loci among
bipolar disorder susceptibility variants. Mol. Psychiatry. Nature Publishing Group;
2013;18:340–6.

4. Tehranchi AK, Myrthil M, Martin T, Hie BL, Golan D, Fraser HB. Pooled ChIP-Seq
Links Variation in Transcription Factor Binding to Complex Disease Risk. Cell.
2016;165:730–41.

5. Edwards SL, Beesley J, French JD, Dunning AM. Beyond GWASs: Illuminating the
Dark Road from Association to Function. The American Journal of Human Genetics.
2013;93:779–97.

6. Tak YG, Farnham PJ. Making sense of GWAS: using epigenomics and genome
engineering to understand the functional relevance of SNPs in non-coding regions of
the human genome. Epigenetics & Chromatin 2015 8:1. BioMed Central; 2015;8:1.

7. Maurano MT, Humbert R, Rynes E, Thurman RE, Haugen E, Wang H, et al.
Systematic Localization of Common Disease-Associated Variation in Regulatory DNA.
Science. American Association for the Advancement of Science; 2012;337:1190–5.

8. Hindorff LA, Sethupathy P, Junkins HA, Ramos EM, Mehta JP, Collins FS, et al.
Potential etiologic and functional implications of genome-wide association loci for
human diseases and traits. Proc. Natl. Acad. Sci. U.S.A. National Acad Sciences;
2009;106:9362–7.

9. Robertson G, Hirst M, Bainbridge M, Bilenky M, Zhao Y, Zeng T, et al. Genome-wide
profiles of STAT1 DNA association using chromatin immunoprecipitation and massively
parallel sequencing. Nature Methods. Nature Publishing Group; 2007;4:651–7.

10. Coetzee SG, Coetzee GA, Hazelett DJ. motifbreakR: an R/Bioconductor package
for predicting variant effects at transcription factor binding sites. Bioinformatics. Oxford
University Press; 2015;31:btv470–3849.

11. Thomas-Chollier M, Hufton A, Heinig M, O'Keeffe S, Masri NE, Roider HG, et al.
Transcription factor binding predictions using TRAP for the analysis of ChIP-seq data
and regulatory SNPs. Nature Protocols. Nature Research; 2011;6:1860–9.

12. Zhou J, Troyanskaya OG. Predicting effects of noncoding variants with deep
learning-based sequence model. Nature Methods. Nature Research; 2015;12:931–4.

13. Consortium TEP. An integrated encyclopedia of DNA elements in the human
genome. Nature. Nature Research; 2012;489:57–74.

14. Consortium RE, Kundaje A, Meuleman W, Ernst J, Bilenky M, Yen A, et al.
Integrative analysis of 111 reference human epigenomes. Nature. Nature Research;
2015;518:317–30.

15. Tibshirani R. Regression shrinkage and selection via the Lasso. Journal of the
Royal Statistical Society Series B-Methodological. 1996;58:267–88.

16. Muglia P, Tozzi F, Galwey NW, Francks C, Upmanyu R, Kong XQ, et al. Genome-
wide association study of recurrent major depressive disorder in two European case-
control cohorts. Mol. Psychiatry. Nature Publishing Group; 2010;15:589–601.

17. Kendler KS, Gatz M, Gardner CO, Pedersen NL. A Swedish national twin study of
lifetime major depression. Am J Psychiatry. American Psychiatric Publishing;

2006;163:109–14.

18. Levinson DF, Mostafavi S, Milaneschi Y, Rivera M, Ripke S, Wray NR, et al. Genetic
Studies of Major Depressive Disorder: Why Are There No Genome-wide Association
Study Findings and What Can We Do About It? Biol. Psychiatry. 2014;76:510–2.

19. Hyde CL, Nagle MW, Tian C, Chen X, Paciga SA, Wendland JR, et al. Identification
of 15 genetic loci associated with risk of major depression in individuals of European
descent. Nat. Genet. Nature Research; 2016.

20. Consortium SWGOTPG, Ripke S, Neale BM, Corvin A, Walters JT, Farh K-H, et al.
Biological Insights From 108 Schizophrenia-Associated Genetic Loci. Nature. Europe
PMC Funders; 2014;511:421–7.

21. Arloth J, Bogdan R, Weber P, Frishman G, Menke A, Wagner KV, et al. Genetic
Differences in the Immediate Transcriptome Response to Stress Predict Risk-Related
Brain Function and Psychiatric Disorders. Neuron. 2015;86:1189–202.

22. Consortium T1GP. A map of human genome variation from population-scale
sequencing. Nature. Nature Research; 2011;473:544–4.

23. Delaneau O, Zagury J-F, Marchini J. Improved whole-chromosome phasing for
disease and population genetic studies. Nature Methods. Nature Research; 2013;10:5–
6.

24. Howie BN, Donnelly P, Marchini J. A Flexible and Accurate Genotype Imputation
Method for the Next Generation of Genome-Wide Association Studies. Schork NJ,
editor. PLOS Genet. Public Library of Science; 2009;5:e1000529.

25. Wohleb ES, Franklin T, Iwata M, Duman RS. Integrating neuroimmune systems in
the neurobiology of depression. Nat. Rev. Neurosci. Nature Research; 2016;17:497–
511.

26. Mostafavi S, Battle A, Zhu X, Potash JB, Weissman MM, Shi J, et al. Type I
interferon signaling genes in recurrent major depression: increased expression detected
by whole-blood RNA sequencing. Mol. Psychiatry. Nature Publishing Group;
2014;19:1267–74.

27. Hsu F, Kent WJ, Clawson H, Kuhn RM, Diekhans M, Haussler D. The UCSC Known
Genes. Bioinformatics. Oxford University Press; 2006;22:1036–46.

28. Bhat S, Dao DT, Terrillion CE, Arad M, Smith RJ, Soldatov NM, et al. CACNA1C
(Cav1.2) in the pathophysiology of psychiatric disease. Prog. Neurobiol. 2012;99:1–14.

29. Van Den Bossche MJ, Strazisar M, De Bruyne S, Bervoets C, Lenaerts A-S, De
Zutter S, et al. Identification of a CACNA2D4 deletion in late onset bipolar disorder
patients and implications for the involvement of voltage-dependent calcium channels in
psychiatric disorders. Am. J. Med. Genet. B Neuropsychiatr. Genet. Wiley Subscription

Services, Inc., A Wiley Company; 2012;159B:465–75.

30. Ni P, Ma X, Lin Y, Lao G, Hao X, Guan L, et al. Methionine sulfoxide reductase A
(MsrA) associated with bipolar I disorder and executive functions in A Han Chinese
population. J Affect Disord. 2015;184:235–8.

31. Walss-Bass C, Soto-Bernardini MC, Johnson-Pais T, Leach RJ, Ontiveros A,
Nicolini H, et al. Methionine sulfoxide reductase: a novel schizophrenia candidate gene.
Am. J. Med. Genet. B Neuropsychiatr. Genet. Wiley Subscription Services, Inc., A Wiley
Company; 2009;150B:219–25.

32. Millstein J, Zhang B, Zhu J, Schadt EE. Disentangling molecular relationships with a
causal inference test. BMC Genet. BioMed Central; 2009;10.

33. Griffon A, Barbier Q, Dalino J, van Helden J, Spicuglia S, Ballester B. Integrative
analysis of public ChIP-seq experiments reveals a complex multi-cell regulatory
landscape. Nucl. Acids Res. Oxford University Press; 2015;43:e27–7.

34. Lee D, Karchin R, Beer MA. Discriminative prediction of mammalian enhancers from
DNA sequence. Genome Res. Cold Spring Harbor Lab; 2011;21:2167–80.

35. Waldmann P, Mészáros G, Gredler B, Fuerst C, Sölkner J. Evaluation of the lasso
and the elastic net in genome-wide association studies. Frontiers in Genetics. Frontiers;
2013;4.

36. Ayers KL, Cordell HJ. SNP Selection in genome-wide and candidate gene studies
via penalized logistic regression. Genetic Epidemiology. Wiley Subscription Services,
Inc., A Wiley Company; 2010;34:879–91.

37. Dehman A, Ambroise C, Neuvial P. Performance of a blockwise approach in
variable selection using linkage disequilibrium information. BMC Bioinformatics. BioMed
Central; 2015;16.

38. Chen SX, Cherry A, Tari PK, Podgorski K, Kwong YKK, Haas K. The Transcription
Factor MEF2 Directs Developmental Visually Driven Functional and Structural
Metaplasticity. Cell. 2012;151:41–55.

39. Barbosa AC, Kim M-S, Ertunc M, Adachi M, Nelson ED, McAnally J, et al. MEF2C, a
transcription factor that facilitates learning and memory by negative regulation of
synapse numbers and function. Proc. Natl. Acad. Sci. U.S.A. National Acad Sciences;
2008;105:9391–6.

40. Pfeiffer BE, Zang T, Wilkerson JR, Taniguchi M, Maksimova MA, Smith LN, et al.
Fragile X Mental Retardation Protein Is Required for Synapse Elimination by the
Activity-Dependent Transcription Factor MEF2. Neuron. 2010;66:191–7.

41. Wang K, Li M, Hakonarson H. ANNOVAR: functional annotation of genetic variants
from high-throughput sequencing data. Nucl. Acids Res. Oxford University Press;

2010;38:–e164.

42. Friedman J, Hastie T, Tibshirani R. glmnet: Lasso and elastic-net regularized
generalized linear models. R package version; 2009.

43. Wie T. Corrplot: Visualization of a correlation matrix-R package version 0.71.
Retrieved August; 2013.

44. Houseman E, Accomando WP, Koestler DC, Christensen BC, Marsit CJ, Nelson
HH, et al. DNA methylation arrays as surrogate measures of cell mixture distribution.
BMC Bioinformatics. BioMed Central; 2012;13:86.

45. Chen Y-A, Lemire M, Choufani S, Butcher DT, Grafodatskaya D, Zanke BW, et al.
Discovery of cross-reactive probes and polymorphic CpGs in the Illumina Infinium
HumanMethylation450 microarray. Epigenetics. Taylor & Francis; 2014;8:203–9.

46. Krueger F, Andrews SR. Bismark: a flexible aligner and methylation caller for
Bisulfite-Seq applications. Bioinformatics. Oxford University Press; 2011;27:1571–2.

47. Fortin J-P, Labbe A, Lemire M, Zanke BW, Hudson TJ, Fertig EJ, et al. Functional
normalization of 450k methylation array data improves replication in large cancer
studies. Genome Biol. BioMed Central; 2014;15:1.

48. Johnson WE, Li C, Rabinovic A. Adjusting batch effects in microarray expression
data using empirical Bayes methods. Biostatistics. Oxford University Press;
2006;8:118–27.

49. Arloth J, Bader DM, Röh S, Altmann A. Re-Annotator: Annotation Pipeline for
Microarray Probe Sequences. Xue Y, editor. PLOS ONE. Public Library of Science;
2015;10:e0139516.

50. Lin SM, Du P, Huber W, Kibbe WA. Model-based variance-stabilizing transformation
for Illumina microarray data. Nucl. Acids Res. Oxford University Press; 2007;36:e11–1.

51. Chikina M, Zaslavsky E, Sealfon SC. CellCODE: A robust latent variable approach
to differential expression analysis for heterogeneous cell populations. Bioinformatics.
Oxford University Press; 2015;31:1584–91.

52. Evangelou E, Ioannidis JPA. Meta-analysis methods for genome-wide association
studies and beyond. Nature Reviews Genetics. Nature Research; 2013;14:379–89.

53. Hahne F, Ivanek R. Visualizing Genomic Data Using Gviz and Bioconductor.
Methods Mol. Biol. New York, NY: Springer New York; 2016;1418:335–51.

54. Csardi G, Nepusz T. The igraph software package for complex network research.
InterJournal. 2006.

55. Wickham H. ggplot2. Cham: Springer; 2016.

1007