Thermal niche evolution across replicated *Anolis* lizard adaptive radiations

Alex R. Gunderson[†*], D. Luke Mahler[§], and Manuel Leal[ʝ]

†Department of Environmental Science, Policy, and Management, University of California, Berkeley,

California 94720-3140

*Corresponding author: e-mail: alexrgunderson@gmail.com

§Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, Ontario, M5S 3B2

CA

ʝ Division of Biological Sciences, University of Missouri, 105 Tucker Hall, Columbia, MO 65211.

Running title: Physiology of adaptive radiation

Keywords: *Anolis*, thermal biology, physiology, thermal tolerance, thermal performance, evolutionary

rates, sympatry, species richness

Data accessibility: If accepted, data herein will be archived at Dryad (http://datadryad.org)

Abstract

Elucidating how ecological and evolutionary mechanisms interact to produce and maintain biodiversity is

a fundamental problem in evolutionary ecology. We investigate this issue by focusing on how

physiological evolution affects performance and species coexistence along the thermal niche axis in

replicated radiations of *Anolis* lizards, groups best known for resource partitioning based on

morphological divergence. We find repeated divergence in thermal physiology within these radiations,

and that this divergence significantly affects performance within natural thermal environments.

Morphologically similar species that co-occur invariably differ in their thermal physiology, providing

evidence that physiological divergence facilitates species co-existence within anole communities. Despite

repeated divergence in traits of demonstrable ecological importance, phylogenetic comparative analyses

indicate that physiological traits have evolved more slowly than key morphological traits related to the

structural niche. Phylogenetic analyses also reveal that physiological divergence is correlated with

divergence in broad-scale habitat climatic features commonly used to estimate thermal niche evolution,

but that the latter incompletely predicts variation in the former. We provide comprehensive evidence for

repeated adaptive evolution of physiological divergence within *Anolis* adaptive radiations, including the

complementary roles of physiological and morphological divergence in promoting community-level

diversity. We recommend greater integration of performance-based traits into analyses of climatic niche

evolution, as they facilitate a more complete understanding of the phenotypic and ecological

consequences of climatic divergence.

Introduction

A mechanistic understanding of how divergent phenotypic evolution facilitates species coexistence is a fundamental problem in ecology and evolution [1]. This problem comes into particular focus in the study of adaptive radiation, a process for which a hallmark is the repeated evolution of niche differences that allow closely related species to partition resources and stably co-occur [2-4]. Traditionally, most research on trait evolution during adaptive radiation has focused on morphological traits associated with dietary resource acquisition or structural habitat use, due to the clear role such adaptations play in mediating competition and facilitating coexistence [3, 5-7]. Physiological adaptation to abiotic conditions has received comparatively less attention in adaptive radiation studies, particularly in animals [3, 8]. Such adaptations are more typically studied for their role in generating divergence among geographically disjunct lineages experiencing different climates [9, 10]. However, physiological divergence can also play an important role in adaptive radiation by facilitating fine-scale partitioning of the local climatic niche [11], providing an additional resource axis along which a radiating clade may diversify [12, 13]. Indeed, such abiotic niche divergence has been hypothesized to underlie many impressive radiations, including bolitoglossine salamanders [14, 15], *Liolaemus* lizards [16-18], *Drosophila* flies [19], *Petrolisthes* crabs [20] and lake whitefish [21].

If physiological adaptation to abiotic conditions contributes to adaptive radiation, physiological divergence should affect performance in the different abiotic regimes that species occupy. Therefore, knowledge of both physiological traits related to the fundamental abiotic niche (broadly, the range of conditions a species could occupy) and abiotic conditions (at scales appropriate for the organisms under study [22]) are required to make the important phenotype-to-performance linkages established for many classic adaptive radiations driven by morphological divergence [3, 5, 23]. Despite the power of performance-based analyses, one of most common tools applied in recent investigations of abiotic niche divergence is correlative ecological niche modeling (ENM), which estimates species' realized niches (*i.e.*, the range of conditions that species occupy) using information about species occurrences and broad-scale climate conditions [24-28]. Though correlative ENMs are powerful for addressing many questions, they

also have characteristics that limit the inferences that can be made from their application [29-34]. In

particular, correlative ENMs provide no information about phenotypic traits (physiological or otherwise)

linked to performance [30]. Additionally, ENMs are limited in their ability to elucidate the role of abiotic

niche divergence in coexistence, as they do not provide fine-scale information about microclimate use.

Such factors can be of paramount importance for coexistence in radiating clades, as species with identical

or largely overlapping ranges may be able to partition resources via adaptation to different

microenvironments [11, 35].

Here, we investigate the contribution of thermal niche divergence to the replicated *Anolis* lizard

adaptive radiations of Puerto Rico and Jamaica by analyzing patterns and performance consequences of

physiological divergence across 16 species (Fig. 1). Greater Antillean *Anolis* are best known for

morphological diversification along the fundamental structural niche axis [36-38], but morphological

evolution cannot fully explain the patterns of divergence and community composition observed across the

Greater Antilles [23]. Divergence in the thermal niche is also proposed to be an important component of

Anolis diversification based primarily on realized niche estimates from natural history and ENM studies

[12, 13, 24, 39-44]. However, we still know little about how physiology evolves during realized niche

divergence [45] and the extent to which physiological evolution facilitates species co-existence.

We first use data from the Puerto Rican *cristatellus* species group to investigate repeated

divergence in thermal physiology and performance between species that occupy distinct thermal

environments. Next, we integrate these data with measurements of operative thermal conditions in Puerto

Rico to explore the performance consequences of physiological divergence. We then consider spatial

patterns of physiological variation to evaluate the contribution of physiological evolution to fine-scale

niche partitioning and community-level species co-existence across Puerto Rico and Jamaica. Finally, we

conduct phylogenetic analyses of rates of physiological and morphological evolution to compare the pace

of disparification between traits associated with the fundamental thermal niche and the fundamental

structural niche, and we employ phylogenetic methods to estimate the strength of the association between

fundamental and broad-scale realized thermal niche divergence.

Methods

We measured physiological traits for 304 individuals across 16 species from three islands (Fig. 1; see

Supplementary Table 1 for species information). Throughout, we focus primarily on the eight Puerto

Rican members of the *cristatellus* group, for two reasons. First, phylogenetic and natural history data

indicate that these species diversified *in situ* and form four sister species pairs [46-48], with each pair

representing an independent divergence in realized thermal niche based on the degree to which they

occupy open versus shaded perches (which throughout we refer to as "warm niche" and "cool niche"

species, respectively)[12, 41, 49, 50]. Second, we have detailed measurements of operative thermal

environments in Puerto Rico to investigate performance consequences of physiological divergence.

Experiments in Puerto Rico were conducted May 20 – June 20 2011, October 8-21 2011, and May 13-28

2012 at the Mata de Plátano field station. On Jamaica, we sampled five of the six extant endemic species,

plus the introduced *A. sagrei*. As with the Puerto Rican species, the endemic Jamaican anoles evolved *in*

situ from a single colonization event [51]. Experiments in Jamaica occurred from February 25 – March 13

2013 at Green Castle Estates, a privately owned farm and nature preserve in St. Mary Parish. Prior to

experiments, lizards were kept individually in plastic cages (18 x 11 x 15 cm) with a wooden dowel perch

on an approximately 12L:12D light schedule. Lizards were watered daily and fed crickets or Phoenix

worms every other day. The exception was *A. acutus* from St. Croix, which were brought to Durham, NC

and housed following Gunderson and Leal [52].

Heat tolerance (CT_{max}) was measured following Leal and Gunderson [53], with N = 9-11

individuals/species (Supplementary Table S1). Briefly, lizards (all captured three days prior to

measurements) were warmed under a heat lamp while monitoring their body temperatures with a wire

thermocouple probe placed inside the cloaca. Lizards were flipped onto their backs at one-degree body

temperature intervals starting at 34°C, and CT_{max} was recorded as the body temperature at which they lost

righting ability. Warming rates were very similar among species, with mean rates ranging from 1.8 to

2.6°C/min (Fig. S1). We modeled variation in log-transformed CT_{max} among species using analysis of

variance (ANOVA). Planned orthogonal contrasts were applied to test *a priori* predictions about

divergence in CT_{max} between sister species (in Puerto Rico) and divergence among sympatric,

morphologically similar species (Puerto Rico and Jamaica).

Temperature-dependent sprint speeds were measured for thirteen species (N = 9-13

individuals/species, Supplementary Table S1) following Gunderson and Leal [52]. Briefly, we analyzed

high-speed video (120 frames/s) of lizards running up a 2 m wooden racetrack set at a 37° angle and

marked with a line every 12.5 cm. Lizards were run 2-4 times at each temperature (two runs minimum,

additional runs were added if the lizard stopped or jumped off of the track during a trial). Speed for a

lizard at a given temperature was taken as the fastest 25 cm interval at that temperature [54, 55]. Puerto

Rican lizards were run at five body temperatures in the following randomized order: 32, 22, 27, 17, and

35°C. Conditions were the same for Jamaican lizards, except the 17°C temperature was excluded. All

individuals were captured one day prior to the start of trials, and each temperature treatment was

administered on a different day over five straight days (four days for Jamaican lizards). No animals were

used for both heat tolerance and sprint performance. Target body temperatures were achieved by placing

lizards into a calibrated chilling-heating incubator (Cole-Parmer©, Vernon Hills, IL, USA) for 30 min

prior to a run [52].

The body temperature of maximum performance (the "optimal temperature," T_{opt}) was estimated

for each individual [sensu 56]. Thermal performance curves are non-linear [57], and we initially fit two

different non-linear models to the sprint data for each individual: a second order polynomial and a

modified Gaussian function ($P(T) = e^{-(\frac{T-T_{opt}}{2 \bullet \sigma})^2}$) commonly used to fit performance curves for ectotherms

[58], including *Anolis* sprinting [52, 59]. The Gaussian model provided relatively poor fits to the data

(much higher residual standard errors than the second-order polynomial model (Fig. S2)). Therefore, T_{opt}

estimates from the second-order polynomial models were used in analyses. T_{opt} estimates were

constrained such that T_{opt} had to occur within the range of experimental temperatures. This is a

conservative assumption that likely leads us to underestimate divergence in thermal physiology between

species (see Results). T_{opt} data were heavily skewed (see Results), so we compared T_{opt} among species by

assessing overlap in bootstrapped 95% confidence intervals (999 bootstrap replicates using the adjusted

bootstrap percentile method due to the skew in our data [60]). Analyses were conducted with the "boot"

and "boot.ci" functions, respectively, from the "boot" package in R [61].

We calculated the expected physiological performance of Puerto Rican species under natural

warm (open xeric forest) and cool (shaded mesic forest) operative thermal regimes found on Puerto Rico

[62]. Operative temperature distributions provide a quantitative description of a thermal environment

integrating the physical features of the organism [63], representing the distribution of available body

temperatures in a habitat. Operative temperature distributions came from a previous study [52] and were

measured with copper lizard models [50] during the peak breeding season (summer) for Puerto Rican

anoles [64]. Mean operative temperature in the mesic forest (28.9°C) is significantly cooler than that in

the xeric forest (33.4°C) by 4.5°C [52].

We estimated two performance metrics under the different operative environments: 1) relative

sprint performance capacity [52, 59, 63], and 2) probability of overheating (the percentage of operative

temperature observations over CT_{max}). For relative performance capacity, we generated species-level

thermal performance curves using our sprint speed and CT_{max} data. The curves were a combination of two

functions fit to the data: one describing performance below T_{opt}, and the other describing performance

from T_{opt} to CT_{max} [52, 58, 59]. The former function was a second-order polynomial fit to the sprint data,

and the latter function took the form: $P(T) = 1.0 - \left(\dfrac{T - T_{opt}}{CT_{max} - T_{opt}} \right)^2$ [52, 59]. Curves were scaled to relative

performance by setting sprint speed at T_{opt} to 1.0 (see Fig. S3 for fitted curves). Performance in each

habitat was calculated by applying the curves to operative temperatures [52, 59, 63].

To compare rates of evolution of thermal physiology to ecomorphological traits involved in

fundamental structural niche partitioning, we used a phylogenetic comparative approach. Rate of

evolution of heat tolerance (CT_{max}, with expanded species sampling, see Supplementary Table S1) was

compared to rates for body size (snout-vent-length, or SVL), head length, femur length, and adhesive

toepad width for the 4[th] hindtoe. We used the maximum clade credibility phylogeny and morphological

data of Mahler *et al*. [47] for these analyses. Shape data (head length, femur length, and toepad width)

consisted of residuals from phylogenetic regressions of each shape variable on SVL, a standard measure

of lizard body size; these residuals were obtained using the phyl.resid function in phytools [65]. We used

the approach of Adams [66] to conduct pairwise rate comparisons for all combinations of these five traits.

We used likelihood ratio tests to compare a model in which the two traits could evolve at different rates to

a model in which the traits were constrained to evolve at a single rate. Rates were estimated using species

means of natural log-transformed values for all traits (such that rate variation is represented in terms of

relative change in proportion to the trait mean; this is essential for comparison of traits measured using

different units [66]). For these analyses, we co-estimated among-trait covariances (similar results were

obtained with covariances fixed at zero).

To test for an association between fundamental and broad-scale realized thermal niche estimates,

we conducted a phylogenetically-controlled correlation test between CT_{max} and geo-referenced Worldclim

temperature data for each species based on collection localities. We used the first principal component

axis from the *Anolis* climate temperature dataset of Algar and Mahler (2015), in which a phylogenetic

principal component analysis (PCA) was used to condense the 11 Worldclim variables related to

temperature (at 1 km resolution) into PCA axes (Supplementary Table S2). The first PC axis loaded

positively for all variables related to temperature (e.g., mean annual temperature, mean maximum

temperature; Supplementary Table S2), and can therefore be considered a composite estimate of habitat

temperature. Phylogenetically-controlled correlation analyses between CT_{max} and individual Worldclim

variables that loaded strongly with PC1 yielded similar results (Supplementary Figure S4).

Results

All Puerto Rican sister species pairs have diverged in at least one physiological trait (T_{opt}, CT_{max}, or both),

and in the predicted direction in all cases (Fig. 2). T_{opt} differed between the *A. evermanni*/*A. stratulus* and

A. cristatellus/*A. cooki* species pairs (non-overlapping bootstrapped 95% CIs), but not between the *A.*

poncensis/*A. gundlachi* or *A. pulchellus*/*A. krugi* species pairs (Fig. 2A). We note that the T_{opt} of *A. cooki*

and *A. stratulus* were likely underestimated because most individuals had a T_{opt} of 35°C solely because

that was our maximum test temperature. CT_{max} differed among Puerto Rican species ($F_{7,72} = 42.74$; $P <$

0.001), with significant divergence in three of the four species pairs: *A. evermanni*/*A. stratulus*, *A.*

poncensis/*A. gundlachi*, and *A. pulchellus*/*A. krugi* (all $P < 0.001$), but not in the *A. cristatellus*/*A. cooki*

species pair ($P = 0.490$; Fig. 2B).

Physiological divergence influenced predicted performance in natural environments. In three of

four species pairs, the cool niche species had higher predicted performance than the warm niche species

in the shaded operative thermal environment, with mean performance advantages of 8-14% depending on

the species pair and time of day (Fig. 3). The lone exception was the *A. pulchellus*/*A. krugi* species pair,

for which both have similar performance estimates (Fig. 3G). Warm niche species tend to have a

performance advantage in the open operative environment, particularly during midday hours (from 10:00-

13:00; Fig. 3). The *A. cristatellus*/*A. cooki* species pair is the exception (Fig. 3D). However, *A. cooki*

midday performance is likely underestimated due to our probable underestimation of its T_{opt} (see above).

No species are under threat of overheating in the shaded operative environment (Fig. 4A, C, E,

G). However, all species have overheating risk in the open operative environment (Fig. 4B, D, F, H). For

three of the four sister-species pairs (*A. stratulus*/*A. evermanni*, *A. poncensis*/*A. gundlachi*, and *A.*

pulchellus/*A. krugi*), the cool niche species has a higher probability of overheating (Fig. 4B, F, H).

Difference in overheating probability are particularly acute during midday, when cool niche species have

overheating probabilities at least twice that of their warm niche counterparts (G-tests, all $P < 0.05$). The

A. cristatellus/*A. cooki* species pair, which had no divergence in CT_{max}, was the only exception (Fig. 4D).

Significant physiological divergence is also apparent among Jamaican species. Among species

mean T_{opt} ranges from 29.5 to 34.6°C (Fig. 5A) while CT_{max} ranges from 35.8-41.5°C (Fig. 5B; ANOVA,

$F_{3,36} = 13.6$, $P < 0.001$). Note that T_{opt} values of *A. grahami* and *A. opalinus* are likely underestimated due

to our upper experimental limit of 35°C (see above).

Across Puerto Rico and Jamaica, sympatric species that share the same ecomorphology invariably

differ in thermal physiology (Table 1). Thermal tolerance has evolved significantly more slowly than

some, but not all, ecomorphological traits (Table 2). CT_{max} evolved significantly more slowly than body

size (SVL; $P < 0.001$) and relative limb length (femur; $P = 0.029$), but did not differ from head length (P

$= 0.105$) or toepad width ($P = 0.112$).

CT_{max} was significantly positively correlated with geo-referenced climatic temperatures as

represented by temperature principal component axis 1 ($P = 0.006$, $r^2 = 0.45$; Fig. 6; see Supplementary

Figure S4).

Discussion

Abiotic niche evolution has been invoked as an important component of many evolutionary radiations,

but the functional consequences of abiotic niche divergence are generally poorly known. We demonstrate

that the Puerto Rican *Anolis* adaptive radiation is accompanied by physiological divergence in both heat

tolerance and sensitivity to sub-lethal temperature variability. Warm-niche species (determined based on

preference for open versus shaded perches) have either higher heat tolerance, higher optimal

physiological temperatures, or both, compared to cool-niche species. The anoles of Jamaica also

diversified physiologically, with species evolving essentially the full range of heat tolerances and optimal

temperatures observed in their Puerto Rican counterparts (Fig. 5). These results highlight the possibility

that physiological adaptation can be pervasive within evolutionary radiations, including those for which

morphological divergence is widespread.

Repeated divergence in physiological traits under similar conditions is itself evidence of

adaptation [67]. However, we provide more direct evidence of adaptation by estimating physiological

performance of species in warm and cool operative environments on Puerto Rico. Cool niche species have

higher predicted performance in shaded operative environments than their warm niche sister species in

three of the four species pairs (Fig. 3), with the lone exception a species pair for which there was no

divergence in T_{opt}. The pattern was less pronounced in the open operative environment, with warm niche

species having a sizable performance advantage in only two species pairs (Fig. 3). However, the

performance advantage for warm niche species becomes apparent when considering overheating risk.

During midday hours in the warmer open environment, warm niche species in three of the four species

pairs had less than half the overheating risk of cool niche species (Fig. 4). The sole exception was the *A.*

cristatellus/*A. cooki* species pair, which exhibited no divergence in thermal tolerance (Fig. 2B). One

implication of these results is that what appears to be modest physiological evolution (CT_{max} and T_{opt}

divergence of 2-4°C) can still have significant performance consequences.

The high community-level species diversity associated with *Anolis* and other adaptive radiations

is facilitated by fine-scale resource partitioning [3]. Sympatric anoles are known to behaviorally partition

thermal microhabitats [12, 13], and our findings indicate that this is facilitated by physiological

divergence. When co-occurring anole species share the same fundamental structural niche (*i.e.*,

ecomorphological microhabitat specialization), they invariably differ in thermal physiology (Table 1).

This pattern is also maintained for introduced species: in Jamaica, the endemic trunk-ground species *A.*

lineatopus is found in sympatry with the introduced trunk-ground species *A. sagrei*, and these species

differ significantly in thermal tolerance (Table 1). These data highlight that, even in radiations for which

morphological divergence is important in facilitating coexistence, physiological divergence can also play

an important role, providing novel axes of diversification [68].

We found that thermal tolerance evolved more slowly than some ecomorphological traits (*i.e.*,

SVL and femur length), and evolved at similar rates to others (*i.e.*, head length and toe-pad width; Table

2). In general, divergence along the fundamental thermal niche axis appears to have occurred more slowly

than divergence along the fundamental structural niche axis, at least within the Puerto Rican and Jamaican

anole radiations. This finding is inconsistent with a recent analysis that found that thermal niche,

estimated as field body temperature, evolves more quickly than morphology in Caribbean anoles [43].

Discrepancy between these results may be due to the different traits analyzed and/or the different species

included, or the fact that that study estimated rates of absolute, rather than proportional, trait change (Fig.

4). Regardless, the observation that physiology evolves slowly, with repeated divergence of relatively

small magnitude, is consistent with our above suggestion that large changes in physiology are not

necessary for divergence to be ecologically important. These results also indicate that caution should be

exhibited when inferring the importance of traits based on comparisons of evolutionary rate estimates in

the absence of data on how phenotypic changes map to performance change under natural conditions.

We found that fundamental and broad-scale realized thermal niche estimates were correlated, as

CT_{max} was significantly positively correlated with geo-referenced temperature data across species (Fig. 6).

Therefore, the realized thermal niches estimated in correlative ENM studies can capture some of the

underlying variability in the fundamental thermal niche. Nonetheless, the majority of variability in the

data (~55%) is left unexplained, and there are clear cases where realized and fundamental niche estimates

diverge. For example, *A. lineatopus* and *A. gundlachi* have very similar CT_{max} (mean CT_{max} = 36.3°C and

36.2°C, respectively), but occur in very different realized thermal niches (temperature PC1 = 0.14 and -

2.66, respectively). Conversely, *A. opalinus* and *A. pulchellus* occur in very similar realized niches

(temperature PC1 = -0.53 and -0.51, respectively) but have very different CT_{max} (mean CT_{max} = 37.5°C

and 41.3°C, respectively). These results underscore the fact that broad-scale climatic conditions are not

necessarily reflective of the underlying thermal biology of the organism being considered. Much of this

discrepancy is likely due to behavioral thermoregulation, which allows taxa in the same geographic

location to experience very different body temperatures and those in different geographic locations to

experience very similar body temperatures [35, 69].

We have shown that fundamental thermal niche divergence is important in the Puerto Rican and

Jamaican *Anolis* adaptive radiations. However, the importance of such divergence is likely not restricted

to these islands. For example, species of *cybotes* group trunk-ground anoles in Hispaniola have diverged

in thermal tolerance along an elevational gradient [45, 70], and variation in thermal physiology also

occurs among non-Antillean anoles [71, 72]. Water loss rates also differ among anole species and

populations [73-79]. In this context, our results suggest that physiological divergence may be as important

as morphological divergence for diversification and coexistence in this classic radiation.

Achieving a mechanistic understanding of how evolutionary and ecological processes interact to

promote the production and maintenance of biodiversity is a long-standing goal in evolutionary ecology.

Given the reality of ongoing climate change, the need to understand these processes has become

immediate, particularly with respect to temperature-dependent traits [80]. By focusing on physiological

traits that link temperature to performance, we demonstrate that the evolution of thermal physiology can

facilitate adaptive radiation by contributing to *in situ* performance tradeoffs and species co-existence.

While our trait- and performance-based analyses would not be possible using the type of broad-scale

climatic data used in ENMs studies alone, we find that climate-scale realized niche features correlate with

fundamental niche features. Nonetheless, climatic data provide imperfect signal with respect to the

evolution of functional traits, likely due to the ability of species to behaviorally augment their effective

thermal environments. We suggest that deeper insights about the contribution of climatic niche

divergence to evolutionary radiations will emerge when correlative niche data are used in tandem with

experimental physiological studies within an integrative research program.

Acknowledgements

We thank J. Losos and two anonymous reviewers for helpful comments on this manuscript, A. Algar for

help with data, D. Steinberg for field help and the El Verde and Mata de Plátano field stations, Ridge to

Reef farm, and Green Castle Estates for logistical support. Permits were provided by the Puerto Rican

Departmento de Recursos Naturales y Ambientales, the St. Croix Division of Fish and Wildlife, and the

Jamaican National Environment and Planning Agency.

Funding. This research was funded by NSF DDIG #1110570 to A.R.G.

Ethics. This research was conducted with the approval of the Duke University Animal Care and Use

Committee permit #A106-10-04.

Author Contributions. A.R.G. and M.L. designed the study. A.R.G. collected the physiological data

and conducted all statistical analyses that were not phylogenetically controlled. L.D.M. conducted all

phylogenetic comparative analyses. A.R.G. led the writing with contributions from M.L. and L.D.M.

Competing interests. We have no competing interests.

Data accessibility. Physiological data are deposited in Dryad (DOI:

https://doi.org/10.5061/dryad.688jj72)

References

1.   Hutchinson G.E. 1959 Homage to Santa Rosalia or why are there so many kinds of animals? *The*
American Naturalist **93**(870), 145-159.
2.   Simpson G.G. 1953 *Major features of evolution*. New York, Columbia University Press.
3.   Schluter D. 2000 *The ecology of adaptive radiation*, Oxford University Press.
4.   McPeek M.A. 2008 The ecological dynamics of clade diversification and community assembly.
The American Naturalist **172**(6), E270-E284.
5.   Grant P.R. 1999 *Ecology and evolution of Darwin's finches*, Princeton University Press.
6.   Streelman J.T., Danley P.D. 2003 The stages of vertebrate evolutionary radiation. *Trends in*
Ecology & Evolution **18**(3), 126-131.
7.   Meyer A. 1993 Phylogenetic relationships and evolutionary processes in East African cichlid
fishes. *Trends in Ecology & Evolution* **8**(8), 279-284.
8.   Nosil P. 2012 *Ecological speciation*, Oxford University Press.
9.   Chown S., Gaston K., Robinson D. 2004 Macrophysiology: large-scale patterns in physiological
traits and their ecological implications. *Functional Ecology* **18**(2), 159-167.
10.   Gaston K.J., Chown S.L., Calosi P., Bernardo J., Bilton D.T., Clarke A., Clusella-Trullas S.,
Ghalambor C.K., Konarzewski M., Peck L.S. 2009 Macrophysiology: a conceptual reunification. *The*
American Naturalist **174**(5), 595-612.
11.   Kaspari M., Clay N.A., Lucas J., Yanoviak S.P., Kay A. 2015 Thermal adaptation generates a
diversity of thermal limits in a rainforest ant community. *Global Change Biology* **21**(3), 1092-1102.
12.   Rand A.S. 1964 Ecological distribution in anoline lizards of Puerto Rico. *Ecology* **45**, 745-752.
13.   Ruibal R. 1961 Thermal relations of five species of tropical lizards. *Evolution* **15**(1), 98-111.
14.   Wake D.B. 1987 Adaptive radiation of salamanders in Middle American cloud forests. *Annals of*
the Missouri Botanical Garden **74**, 242-264.
15.   Kozak K.H., Wiens J.J. 2007 Climatic zonation drives latitudinal variation in speciation
mechanisms. *Proceedings of the Royal Society of London B: Biological Sciences* **274**(1628), 2995-3003.
16.   Labra A., Pienaar J., Hansen T.F. 2009 Evolution of thermal physiology in *Liolaemus* lizards:
adaptation, phylogenetic inertia, and niche tracking. *The American Naturalist* **174**(2), 204-220.
17.   Bonino M.F., Azócar D.L.M., Tulli M.J., Abdala C.S., Perotti M.G., Cruz F.B. 2011 Running in
cold weather: morphology, thermal biology, and performance in the southernmost lizard clade in the
world (*Liolaemus lineomaculatus* section: Liolaemini: Iguania). *Journal of Experimental Zoology Part A:*
Ecological Genetics and Physiology **315**(8), 495-503.
18.   Pincheira-Donoso D. 2011 Predictable variation of range-sizes across an extreme environmental
gradient in a lizard adaptive radiation: evolutionary and ecological inferences. *PloS One* **6**(12), e28942.
19.   Kellermann V., van Heerwaarden B., Sgrò C.M., Hoffmann A.A. 2009 Fundamental evolutionary
limits in ecological traits drive *Drosophila* species distributions. *Science* **325**(5945), 1244-1246.
20.   Stillman J.H., Somero G.N. 2000 A comparative analysis of the upper thermal tolerance limits of
eastern Pacific porcelain crabs, genus *Petrolisthes*: influences of latitude, vertical zonation, acclimation,
and phylogeny. *Physiological and Biochemical Zoology* **73**(2), 200-208.
21.   Dalziel A.C., Martin N., Laporte M., Guderley H., Bernatchez L. 2015 Adaptation and
acclimation of aerobic exercise physiology in Lake Whitefish ecotypes (*Coregonus clupeaformis*).
Evolution **69**(8), 2167-2186.
22.   Potter K.A., Woods H.A., Pincebourde S. 2013 Microclimatic challenges in global change
biology. *Global change biology* **19**, 2932–2939.
23.   Losos J.B. 2009 *Lizards in an evolutionary tree: ecology and adaptive radiation of anoles*,
University of California Press.
24.   Knouft J.H., Losos J.B., Glor R.E., Kolbe J.J. 2006 Phylogenetic analysis of the evolution of the
niche in lizards of the *Anolis sagrei* group. *Ecology* **87**(sp7), S29-S38.
25.   Quintero I., Wiens J.J. 2013 Rates of projected climate change dramatically exceed past rates of
climatic niche evolution among vertebrate species. *Ecology Letters* **16**(8), 1095-1103.

26.   Warren D.L., Glor R.E., Turelli M. 2008 Environmental niche equivalency versus conservatism: quantitative approaches to niche evolution. *Evolution* **62**(11), 2868-2883.

27.   Lawing A.M., Polly P.D., Hews D.K., Martins E.P., Collar D.C., Bronstein J.L. 2016 Including Fossils in Phylogenetic Climate Reconstructions: A Deep Time Perspective on the Climatic Niche Evolution and Diversification of Spiny Lizards (*Sceloporus*). *The American Naturalist* **188**(2), 000-000.

28.   Peterson A.T., Soberón J., Pearson R.G., Anderson R.P., Nakamura M., Martinez-Meyer E., Araújo M.B. 2011 *Ecological niches and geographic distributions*. Princeton, New Jersey, USA, Princeton University Press.

29.   Guisan A., Thuiller W. 2005 Predicting species distribution: offering more than simple habitat models. *Ecology Letters* **8**(9), 993-1009.

30.   Buckley L.B., Urban M.C., Angilletta M.J., Crozier L.G., Rissler L.J., Sears M.W. 2010 Can mechanism inform species' distribution models? *Ecology Letters* **13**(8), 1041-1054.

31.   Alvarado-Serrano D.F., Knowles L.L. 2014 Ecological niche models in phylogeographic studies: applications, advances and precautions. *Molecular Ecology Resources* **14**(2), 233-248.

32.   Kearney M., Porter W. 2009 Mechanistic niche modelling: combining physiological and spatial data to predict species' ranges. *Ecology letters* **12**(4), 334-350.

33.   Warren D.L., Cardillo M., Rosauer D.F., Bolnick D.I. 2014 Mistaking geography for biology: inferring processes from species distributions. *Trends in Ecology & Evolution* **29**(10), 572-580.

34.   Araújo M.B., Peterson A.T. 2012 Uses and misuses of bioclimatic envelope modeling. *Ecology* **93**(7), 1527-1539.

35.   Bogert C.M. 1949 Thermoregulation in reptiles, a factor in evolution. *Evolution* **3**(3), 195-211.

36.   Losos J.B., Jackman T.R., Larson A., de Queiroz K., Rodríguez-Schettino L. 1998 Contingency and determinism in replicated adaptive radiations of island lizards. *Science* **279**(5359), 2115-2118.

37.   Williams E.E. 1983 Ecomorphs, faunas, island size, and diverse end points in island radiations of Anolis. In *Lizard ecology: studies of a model organism* (pp. 326-370. Cambridge, Harvard University Press.

38.   Mahler D.L., Ingram T., Revell L.J., Losos J.B. 2013 Exceptional convergence on the macroevolutionary landscape in island lizard radiations. *Science* **341**(6143), 292-295.

39.   Williams E.E. 1972 The origin of faunas. Evolution of lizard congeners in a complex island fauna: a trial analysis. *Evolutionary Biology* **6**, 47-89.

40.   Heatwole H., Lin T.-H., Villalón E., Muñiz A., Matta A. 1969 Some aspects of the thermal ecology of Puerto Rican anoline lizards. *Journal of Herpetology* **3**, 65-77.

41.   Huey R.B., Webster T.P. 1976 Thermal biology of *Anolis* lizards in a complex fauna: the *cristatellus* group on Puerto Rico. *Ecology* **57**, 985-994.

42.   Losos J.B., Leal M., Glor R.E., de Queiroz K., Hertz P.E., Schettino L.R., Lara A.C., Jackman T.R., Larson A. 2003 Niche lability in the evolution of a Caribbean lizard community. *Nature* **424**(6948), 542-545.

43.   Hertz P.E., Arima Y., Harrison A., Huey R.B., Losos J.B., Glor R.E. 2013 Asynchronous evolution of physiology and morphology in *Anolis* lizards. *Evolution* **67**(7), 2101-2113.

44.   Algar A.C., Mahler D.L. 2015 Area, climate heterogeneity, and the response of climate niches to ecological opportunity in island radiations of *Anolis* lizards. *Global Ecology and Biogeography* **doi:10.1111 /geb.12327**.

45.   Muñoz M.M., Stimola M.A., Algar A.C., Conover A., Rodriguez A.J., Landestoy M.A., Bakken G.S., Losos J.B. 2014 Evolutionary stasis and lability in thermal physiology in a group of tropical lizards. *Proceedings of the Royal Society B: Biological Sciences* **281**(1778), 20132433.

46.   Nicholson K.E., Glor R.E., Kolbe J.J., Larson A., Blair Hedges S., Losos J.B. 2005 Mainland colonization by island lizards. *Journal of Biogeography* **32**(6), 929-938.

47.   Mahler D.L., Revell L.J., Glor R.E., Losos J.B. 2010 Ecological opportunity and the rate of morphological evolution in the diversification of Greater Antillean anoles. *Evolution* **64**(9), 2731-2745.

48.   Gamble T., Geneva A.J., Glor R.E., Zarkower D. 2014 *Anolis* sex chromosomes are derived from a single ancestral pair. *Evolution* **68**(4), 1027-1041.

49.   Hertz P.E. 1992 Temperature regulation in Puerto Rican *Anolis* lizards: a field test using null hypotheses. *Ecology* **73**, 1405-1417.

50.   Hertz P.E. 1992 Evaluating thermal resource partitioning by sympatric lizards *Anolis cooki* and *A. cristatellus*: a field test using null hypotheses. *Oecologia* **90**(1), 127-136.

51.   Jackman T.R., Larson A., De Queiroz K., Losos J.B. 1999 Phylogenetic relationships and tempo of early diversification in *Anolis* lizards. *Systematic Biology* **48**(2), 254-285.

52.   Gunderson A.R., Leal M. 2012 Geographic variation in vulnerability to climate warming in a tropical Caribbean lizard. *Functional Ecology* **26**(4), 783-793.

53.   Leal M., Gunderson A.R. 2012 Rapid change in the thermal tolerance of a tropical lizard. *The American Naturalist* **180**(6), 815-822.

54.   Huey R.B. 1983 Natural variation in body temperature and physiological performance in a lizard (*Anolis cristatellus*). In *Advances in herpetology and evolutionary biology: essays in honor of Ernest E Williams* (pp. 484-490.

55.   Losos J.B. 1990 Ecomorphology, performance capability, and scaling of West Indian *Anolis* lizards: an evolutionary analysis. *Ecological Monographs* **60**, 369-388.

56.   Logan M.L., Cox R.M., Calsbeek R. 2014 Natural selection on thermal performance in a novel thermal environment. *Proceedings of the National Academy of Sciences* **111**(39), 14165-14169.

57.   Angilletta M.J. 2009 *Thermal adaptation: a theoretical and empirical synthesis*, Oxford University Press.

58.   Deutsch C.A., Tewksbury J.J., Huey R.B., Sheldon K.S., Ghalambor C.K., Haak D.C., Martin P.R. 2008 Impacts of climate warming on terrestrial ectotherms across latitude. *Proceedings of the National Academy of Sciences* **105**(18), 6668-6672.

59.   Huey R.B., Deutsch C.A., Tewksbury J.J., Vitt L.J., Hertz P.E., Pérez H.J.Á., Garland T. 2009 Why tropical forest lizards are vulnerable to climate warming. *Proceedings of the Royal Society B: Biological Sciences* **276**(1664), 1939-1948.

60.   Efron B. 1987 Better bootstrap confidence intervals. *Journal of the American Statistical Association* **82**(397), 171-185.

61.   Canty A., Ripley B. 2017 boot: bootstrap R (S-Plus) functions. *R package v 13-20*.

62.   Ewel J.J., Whitmore J.L. 1973 Ecological life zones of Puerto Rico and US Virgin Islands. In *Forest Service Research Paper ITF-18* (U.S. Department of Agriculture.

63.   Hertz P.E., Huey R.B., Stevenson R. 1993 Evaluating temperature regulation by field-active ectotherms: the fallacy of the inappropriate question. *The American Naturalist* **142**, 796-818.

64.   Gorman G.C., Licht P. 1974 Seasonality in ovarian cycles among tropical *Anolis* lizards. *Ecology* **55**, 360-369.

65.   Revell L.J. 2012 phytools: an R package for phylogenetic comparative biology (and other things). *Methods in Ecology and Evolution* **3**(2), 217-223.

66.   Adams D.C. 2013 Quantifying and comparing phylogenetic evolutionary rates for shape and other high-dimensional phenotypic data. *Systematic Biology* **62**, 181-192.

67.   Harvey P.H., Pagel M.D. 1991 *The comparative method in evolutionary biology*, Oxford University Press.

68.   Leal M., Fleishman L.J. 2002 Evidence for habitat partitioning based on adaptation to environmental light in a pair of sympatric lizard species. *Proceedings of the Royal Society of London Series B: Biological Sciences* **269**(1489), 351-359.

69.   Huey R.B., Hertz P.E., Sinervo B. 2003 Behavioral drive versus behavioral inertia in evolution: a null model approach. *The American Naturalist* **161**(3), 357-366.

70.   Hertz P.E., Huey R.B. 1981 Compensation for altitudinal changes in the thermal environment by some *Anolis* lizards on Hispaniola. *Ecology* **62**(3), 515-521.

71.   Logan M.L., Huynh R.K., Precious R.A., Calsbeek R.G. 2013 The impact of climate change measured at relevant spatial scales: new hope for tropical lizards. *Global Change Biology* **19**, 3093-3102.

72.   van Berkum F.H. 1986 Evolutionary patterns of the thermal sensitivity of sprint speed in *Anolis*
lizards. *Evolution* **40**, 594-604.
73.   Gunderson A.R., Siegel J., Leal M. 2011 Tests of the contribution of acclimation to geographic
variation in water loss rates of the West Indian lizard *Anolis cristatellus*. *Journal of Comparative*
Physiology B **181**(7), 965-972.
74.   Hillman S., Gorman G. 1977 Water loss, desiccation tolerance, and survival under desiccating
conditions in 11 species of Caribbean *Anolis*. *Oecologia* **29**(2), 105-116.
75.   Sexton O.J., Heatwole H. 1968 An experimental investigation of habitat selection and water loss
in some anoline lizards. *Ecology* **49**, 762-767.
76.   Hertz P., Arce-Hernandez A., Ramirez-Vazquez J., Tirado-Rivera W., Vazquez-Vives L. 1979
Geographical variation of heat sensitivity and water loss rates in the tropical lizard, *Anolis gundlachi*.
Comparative Biochemistry and Physiology Part A: Physiology **62**(4), 947-953.
77.   Hertz P.E. 1980 Responses to dehydration in *Anolis* lizards sampled along altitudinal transects.
Copeia **1980**, 440-446.
78.   Hillman S., Gorman G.C., Thomas R. 1979 Water loss in *Anolis* lizards: evidence for acclimation
and intraspecific differences along a habitat gradient. *Comparative Biochemistry and Physiology Part A:*
Physiology **62**(2), 491-493.
79.   Dmi'el R., Perry G., Lazell J. 1997 Evaporative water loss in nine insular populations of the lizard
Anolis cristatellus group in the British Virgin Islands. *Biotropica* **29**(1), 111-116.
80.   Huey R.B., Kearney M.R., Krockenberger A., Holtum J.A., Jess M., Williams S.E. 2012
Predicting organismal vulnerability to climate warming: roles of behaviour, physiology and adaptation.
Philosophical Transactions of the Royal Society B: Biological Sciences **367**(1596), 1665-1679.

524
525

Table 1. Physiology differences between co-occurring species that occupy the same or similar structural niches (*i.e*., share the same or similar ecomorphology). For CT_{max}, P-values are given based on one-way ANOVA with planned orthogonal contrasts. For T_{opt}, "*" indicates that the bootstrapped 95% confidence intervals did not overlap, "N.S." indicates no significant difference. "†" indicates that the species is an introduced member of the community.

Table 2. Estimated rates of evolution for heat tolerance (CT_{max}) and morphological traits related to ecomorphological divergence. Rate estimates (bold) are in the diagonal of the matrix. Upper off-diagonals contain estimates of pairwise evolutionary covariances among traits, lower off-diagonals contain P-values for likelihood ratio test comparisons of 2-rate versus 1-rate models for each trait pair. P-values less than 0.05 indicate support for a 2-rate model over a 1-rate model.

555
556

560   Table 1

	co-occurring species	ecomorph	physiological trait	
			CT_{max}	T_{opt}
A. stratulus	*A. evermani*	trunk-crown	<0.001	*
A. cristatellus	*A. cooki*	trunk-ground	0.490	*
A. cristatellus	*A. gundlachi*	trunk-ground	<0.001	n.s.
A. krugi	*A. pulchellus*	grass-bush	<0.001	n.s.
A. lineatopis	*A. grahami*	trunk-ground/trunk-crown	<0.001	n.s.
A. lineatopis	*A. sagrei*[†]	trunk-ground	<0.001	n.s.

569   Table 2.

	CT_{max}	SVL	head length	femur length	toepad width
CT_{max}	**5.80E-05**	-4.90E-05	-1.60E-05	-5.50E-05	-9.90E-06
SVL	<0.001	**2.30E-03**	8.90E-06	-1.80E-05	2.70E-05
head length	0.105	<0.001	**2.70E-05**	1.30E-05	-6.30E-06
femur length	0.029	<0.001	0.001	**1.40E-04**	2.20E-05
toepad width	0.112	<0.001	0.003	0.832	**1.30E-04**

Figure 1: Phylogenetic relationships of the species included in this study, along with the islands from

which they were sampled.

Figure 2: Thermal physiology of four sister-species pairs of Puerto Rican anole. Each species pair is

represented in alternating grey and white regions of the plot with warm niche species in red and cool

niche species in blue. (A) Optimal sprint performance temperatures (T_{opt}). (B) Heat tolerance limits

(CT_{max}). "*", significant difference between members of a species pair; N.S., no significant difference

between members of a species pair. See Methods for analysis details.

Figure 3: Predicted relative physiological performance of sister-species pairs of Puerto Rican *Anolis*

under shaded and open habitat operative thermal environments found on Puerto Rico during morning,

midday, and afternoon/evening hours. Warm niche species in red, cool niche species in blue. (A, B) *A.*

stratulus/A. evermanni, (C, D) *A. cooki/A. cristatellus* (E, F) *A. poncensis/A. gundlachi* (G, H) *A.*

pulchellus/A. krugi.

Figure 4: Predicted overheating probability of sister-species pairs of Puerto Rican *Anolis* under shaded

and open habitat operative thermal environments found on Puerto Rico during morning, midday, and

afternoon/evening hours. Warm niche species in red, cool niche species in blue. (A, B) *A. stratulus/A.*

evermanni, (C, D) *A. cooki/A. cristatellus* (E, F), *A. poncensis/A. gundlachi*, (G, H) *A. pulchellus/A. krugi*.

Figure 5: Thermal physiology of endemic Jamaican anoles. (A) Optimal sprinting temperature (T_{opt}).

Anolis valencienni does not have a T_{opt} estimate because this species could not be induced to run. (B) Heat

tolerance (CT_{max}).

Figure 6: Relationship between CT_{max} and climate temperature PC1 from Algar and Mahler (2015).

631 Figure 1

(a)

(b)

638   Figure 2.

647
648
649

time of day

658 Figure 3

667   Figure 4

675   Figure 5

701 Figure 6.