A functional analysis of the contribution of climatic niche divergence to adaptive

radiation

Alex R. Gunderson[†‡*], D. Luke Mahler[§], and Manuel Leal[¶]

†Romberg Tiburon Center and Department of Biology, San Francisco State University,

3150 Paradise Dr., Tiburon, California 94920 and ‡Department of Integrative Biology,

University of California, Berkeley, California 94720-3140

*Corresponding author: e-mail: alexrgunderson@gmail.com

§Department of Ecology and Evolutionary Biology, University of Toronto, Toronto,

Ontario, M5S 3B2 CA

¶ Division of Biological Sciences, University of Missouri, 105 Tucker Hall, Columbia,

MO 65211. Email: lealm@missouri.edu

Running title: Physiology of adaptive radiation

Keywords: *Anolis*, thermal biology, physiology, thermal tolerance, thermal performance,

evolutionary rates, sympatry, species richness

Data accessibility: If accepted, data herein will be archived at Dryad

(http://datadryad.org)

Abstract

Analyses of evolutionary radiations have traditionally focused on traits related to

resource acquisition, habitat use, and mate choice. Nonetheless, incorporation of climatic

niche divergence has become more common in recent years, driven by the development

of correlative ecological niche models (ENMs) that allow niche characterization based

solely on geo-referenced locality data. However, ENMs require no data on organismal

traits that link climatic conditions to performance, and thus have little to say about how

organisms phenotypically adapt to different climatic conditions, what the performance

consequences of phenotypic divergence are, and how phenotypic divergence influences

resource partitioning within communities of co-occurring species. Here, we conduct a

functional (*i.e.*, physiological) analysis of climatic niche divergence within Greater

Antillean *Anolis* radiations. We find that physiological divergence occurred repeatedly

within these radiations, that there are performance consequences of physiological

divergence within natural habitats, and that physiological divergence facilitates species

co-existence within species-rich communities. We also use phylogenetic methods to

compare rates of evolution of ecologically relevant physiological and morphological

traits, and find that physiology has evolved either more slowly or as fast (but not faster)

than morphology. Our analysis demonstrates the benefit of a functional approach to

understanding macroevolutionary patterns of climatic niche divergence.

Introduction

During adaptive radiations, the proliferation of species is accompanied by adaptive phenotypic diversification, leading to a pattern in which multiple closely related but ecologically distinct species coexist (Simpson 1953; Schluter 2000). The vast majority of research on adaptive radiation, and evolutionary radiation in general, has focused on morphological traits directly linked to resource acquisition, structural habitat use, and mating display (Schluter 2000; Seehausen et al. 2008; Fleishman et al. 2009; Arnegard et al. 2010). Classic examples include divergence in jaw morphology in African cichlids (Meyer 1993) and bill shape in Darwin's finches (Grant 1999) to specialize on different food sources, and body shape divergence in Caribbean *Anolis* lizards to specialize on different structural microhabitats (*e.g.*, twigs, trunks and grasses) (Williams 1983; Losos 2009).

In contrast, adaptation to aspects of the climatic environment, such as temperature and water availability, have traditionally received much less attention in studies of evolutionary radiations, particularly in animals (Schluter 2000). This has been changing in recent years due to the development of global scale climate databases and correlative ecological niche models (ENMs) that allow one to estimate parameters of a species' abiotic ecological niche based on geo-referenced species locality data (Peterson et al. 2011). Due largely to their relative ease of use and the fact that they facilitate the incorporation of ecologically intractable or understudied species into ecological and evolutionary analyses, ENMs are now commonly used to calculate rates and frequencies of climatic niche divergence and to infer adaptive diversification (Knouft et al. 2006; Warren et al. 2008; Quintero and Wiens 2013; Lawing et al. 2016).

However, ENMs also have several disadvantages that limit the inferences that can

be drawn from their application (Guisan and Thuiller 2005; Kearney and Porter 2009;

Buckley et al. 2010; Araújo and Peterson 2012; Alvarado‑Serrano and Knowles 2014;

Warren et al. 2014). For example, they provide no information about what phenotypic

traits, if any, have evolved to facilitate the occupation of different climatic niches. In

other words, functional analyses using the ENM approach alone are not possible

(Buckley et al. 2010). In addition, the climatic data used in ENMs are typically collected

at scales much larger than those at which organisms experience and utilize their habitats

(Potter et al. 2013). Thus, allopatric populations from broadly different climates could

actually experience very similar abiotic conditions because of differences in microhabitat

use (Bogert 1949; Huey et al. 2003); conversely, different species that are broadly

sympatric could actually experience very different abiotic conditions because of

differential microhabitat use (Kaspari et al. 2015).

When species adapt to different climates, the phenotypes most likely to diverge

are physiological traits related to tolerance for, and sensitivity to, abiotic conditions

(Hochachka and Somero 2002). For example, physiological variation is extensive across

environmental gradients (Levin et al. 2001; Hochachka and Somero 2002; Angilletta

2009; Sunday et al. 2011). Physiological divergence can also lead to reproductive

isolation between populations (Matute et al. 2009; Keller and Seehausen 2012) and

facilitate fine-scale resource partitioning among co-occurring taxa (Rand 1964),

characteristics that can hypothetically facilitate adaptive radiation (Schluter 2000).

Indeed, the evolution of thermal physiology is hypothesized to play an important role in

the radiation of a number of groups including bolitoglossine salamanders (Wake 1987;

Kozak and Wiens 2007), *Liolaemus* lizards (Labra et al. 2009; Bonino et al. 2011;

Pincheira-Donoso 2011), *Drosophila* flies (Kellermann et al. 2009) and *Petrolisthes* crabs

(Stillman and Somero 2000). Still, a comprehensive understanding of the contribution of

physiological adaptation to the process of adaptive radiation is currently lacking.

Greater Antillean *Anolis* lizards are a textbook example of adaptive radiation, best

known for the repeated morphological evolution of distinct perch type (*i.e.*, structural

microhabitat) specialists, known as ecomorphs (Williams 1983; Losos et al. 1998; Mahler

et al. 2013). Nonetheless, morphological specialization cannot fully explain the Greater

Antillean radiation because most cladogenic events are not associated with

ecomorphological divergence (Losos 2009). Furthermore, many *Anolis* communities

support more than one species of a given ecomorph (Williams 1983; Losos et al. 2003),

and thus morphological specialization cannot fully explain the diversity of *Anolis*

communities.

Climatic niche divergence is proposed to be a major contributor to the species

richness of Greater Antillean anoles based primarily on body temperature and ENM

studies (Ruibal 1961; Rand 1964; Heatwole et al. 1969; Williams 1972; Huey and

Webster 1976; Losos et al. 2003; Knouft et al. 2006; Hertz et al. 2013; Algar and Mahler

2015). However, we know relatively little about how the occupation of different thermal

niches has been facilitated by physiological evolution during this radiation (Muñoz et al.

2014), and the extent to which physiological divergence has facilitated the co-existence

of species within diverse anole communities. For example, what components of

physiology evolve during adaptive radiations? Additionally, what are the geographical

patterns of physiological divergence? In other words, are physiologically divergent

species geographically separated, or do they occur sympatrically? The latter scenario

would indicate that physiological divergence can promote the maintenance of high

community-level diversity by facilitating species co-existence. Finally, how does the

tempo of evolution of physiological traits compare to that of morphological traits

important to divergence along other ecological axes (*e.g.*, the structural niche axis)?

Here, we take a functional approach to investigate the physiological and

ecological consequences of climatic niche divergence in West Indian *Anolis* lizards. Our

analyses focus primarily on the *cristatellus* species group from Puerto Rico, but we also

conduct analyses that include data from other Puerto Rican anoles as well as species from

St. Croix and Jamaica (Fig. 1). First, we use data from the *cristatellus* species group to

investigate divergence in thermal physiology between species that occupy distinct

thermal niches. Next, we integrate these data with measurements of operative thermal

environments in Puerto Rico to explore the performance consequences of this

physiological variation. We then consider geographical patterns of physiological trait

divergence to evaluate the contribution of thermal adaptation to niche partitioning and

community-level species co-existence across Puerto Rico and Jamaica. Finally, we

conduct phylogenetically-based analyses of phenotypic rates of evolution with

physiological and morphological traits to compare macro-evolutionary patterns of

diversification between traits associated with divergence along the thermal and structural

niche axes.

Methods

STUDY SPECIES

We measured physiological traits for a total of 304 individuals representing 16 species of

anole from the islands of Puerto Rico, St. Croix (U.S. Virgin Islands), and Jamaica (Fig.

1; see Supplementary Table 1 for sample sizes and natural history information for each

species). Nine species are from Puerto Rico, eight of which are members of the

cristatellus species group that diverged *in situ* on the greater Puerto Rico bank following

a single colonization event (Jackman et al. 1999; Losos 2009). According to recent

phylogenetic analyses these species most likely fall into four sister-species pairs

(Nicholson et al. 2005; Mahler et al. 2010; Gamble et al. 2014) (Fig. 1). Each species

within a given pair occupies a different thermal niche (Rand 1964; Huey and Webster

1976; Hertz 1992b, a). We sampled six species from Jamaica, including five of the six

extant endemic species on the island, plus the introduced *A. sagrei*. As with the Puerto

Rican *cristatellus* group, all endemic Jamaican species evolved *in situ* as the result of a

single colonization event (Jackman et al. 1999). Throughout, we focus primarily on the

Puerto Rican *cristatellus* radiation because of the fortuitous pattern of species

relationships (*i.e.*, four sister species pairs with different thermal niches) and because we

have detailed measurements of operative thermal environments in Puerto Rico with

which to investigate performance consequences of physiological divergence.

Experiments in Puerto Rico were conducted May 20 – June 20 2011, October 8-

21 2011, and May 13-28 2012 at the Mata de Plátano field station in the Arecibo

municipality (18.414037 N; 66.728582 W; elevation 150 m). Most lizard collection

occurred at and around Mata de Platano (*A. cristatellus*, *A. krugi*, *A. pulchellus*, and *A.*

culvieri) and El Verde Research Station (18.322416 N; 65.818052 W, elevation 350 m;

A. gundlachi, *A. evermanni*, and *A. stratulus*) along the north coast. Two species occur

only on the southwest coast (*A. poncensis* and *A. cooki*), and were collected at Bosque

Estatal de Boquerón (17.936091 N; 67.192854 W, elevation 5 m) and Bosque Estatal de

Guánica (17.963301 N; 66.889752 W, elevation 5 m), respectively. In Jamaica, animal

collection and experiments occurred from February 25 – March 13 2013 at Green Castle

Estates, a privately owned 1600 square acre farm and nature preserve in St. Mary Parish

(18.287927 N; 76.807951 W, elevation 50 m). Prior to experiments, lizards were kept

individually in plastic cages (18 x 11 x 15 cm) with a wooden dowel for a perch on an

approximately 12L:12D light schedule. Lizards were watered daily and fed crickets or

Phoenix worms every other day. The only exception was *A. acutus* from St. Croix

(captured at 17.753274 N; 64.866563 W, elevation 172 m), which were brought back to

Durham, NC and housed following Gunderson and Leal (2012).

HEAT TOLERANCE

Heat tolerance (CT_{max}) was measured following Leal and Gunderson (2012). Briefly, the

tip of a thermocouple probe (~1.5 m, 40 gauge) was placed inside the cloaca and secured

to the tail with surgical tape so that body temperature could be measured continuously

during the experiment with a model HH603A Type T (sensitivity $\pm$ 0.1°C) digital

thermometer (Omega, Stamford, CT, USA). The lizard was loosely tethered to a piece of

cardboard (29L, 21W cm) with dental floss looped around the waist and body

temperatures were raised by placing lizards under a 150w double-mirror incandescent

light bulb. Lizards were flipped onto their backs at one-degree intervals starting at 34°C

to test righting ability. If the lizard righted itself, it was placed back under the light bulb

and warmed to the next temperature. If not, the trial was terminated. CT_{max} was recorded

as the last temperature at which the lizard righted itself + 0.5°C (Leal and Gunderson

2012). Mean warming rates during experiments were very similar among species, ranging

from 1.8 to 2.6°C/min (Fig. S1). All lizards were captured three days prior to CT_{max}

measurement. We modeled variation in CT_{max} among species using Analysis of Variance

(ANOVA), with separate analyses for the species on Puerto Rico and Jamaica. Data were

log-transformed prior to analysis. Planned orthogonal contrasts were applied to test *a*

priori predictions about divergence in CT_{max} between sister species with different thermal

niches (in Puerto Rico) and divergence among sympatric, morphologically similar

species (Puerto Rico and Jamaica).

SPRINT PERFORMANCE

Temperature-dependent sprint speeds were measured for thirteen species (*A. cuvieri*, *A.*

acutus, and *A. valencienni* were excluded) following Gunderson and Leal (2012). Briefly,

lizards were filmed at high speed (120 frames/s; Sony© Handycam, Tokyo, Japan) while

sprinting up a 2 m wooden racetrack marked with a line every 12.5 cm. The ramp was set

at a 37° angle as anoles will jump instead of run on a horizontal surface (A. Gunderson,

personal observation). Videos were watched frame-by-frame to determine speed. Lizards

were run 2-4 times at each temperature (two runs were the minimum, additional runs

were added if the lizard stopped or jumped off of the track during a run). Runs for a

lizard at a given temperature were a minimum of 40 minutes apart. Speed for a lizard at a

given temperature was taken as the fastest 25 cm run at that temperature (Huey 1983;

Losos 1990). In Puerto Rico, lizards were run at five body temperatures in the following

randomized order: 32, 22, 27, 17, and 35°C. In Jamaica, lizards were run at the same

temperatures in the same order, except the 17°C temperature was excluded. All

individuals were captured one day prior to the start of trials, and each temperature

treatment was administered on a different day over five straight days (four days for

Jamaican lizards). Body temperatures were achieved by placing lizards into a calibrated

chilling-heating incubator (Cole-Parmer©, Vernon Hills, IL, USA) for 30 min prior to a

run (Gunderson and Leal 2012).

The body temperature of maximum physiological performance (the "optimal

temperature," T_{opt}) was estimated for each individual (sensu Logan et al. 2014). Thermal

performance curves are non-linear (Angilletta 2009), and we initially fit two different

non-linear models to the sprint data for each individual: a second order polynomial and a

modified Gaussian function ($P(T) = e^{-(\frac{T-T_{opt}}{2 \cdot \sigma})^2}$) that has been commonly used to fit

performance curves for ectotherms (Deutsch et al. 2008), including *Anolis* sprint

performance (Huey et al. 2009; Gunderson and Leal 2012). The Gaussian model provided

relatively poor fits to the data, with residual standard errors much higher than the second-

order polynomial model for all individuals (Fig. S2). Therefore, we use T_{opt} estimates

from the second-order polynomial models for all analyses. T_{opt} estimates were

constrained such that T_{opt} had to occur within the range of experimental temperatures.

Thus, if the model predicted a T_{opt} greater than 35°C, we set that individual's T_{opt} to

35°C. This is a conservative assumption that likely leads us to underestimate divergence

in thermal physiology between species (see Results).

We were unable to use ANOVA for analyses of T_{opt} data because for four of our

species, most individuals ran fastest at the highest test temperature, yielding heavily

skewed distributions (see Results). Thus, we compared T_{opt} among species by assessing

overlap in 95% confidence intervals. We estimated confidence intervals based on 999

bootstrap replicates using the adjusted bootstrap percentile method due to the skew in our

data (Efron 1987). Bootstrap resampling and 95% CI estimation were calculated in R

using the "boot" and "boot.ci" functions, respectively, from the "boot" package.

PERFORMANCE CONSEQUENCES OF PHYSIOLOGICAL DIVERGENCE

To evaluate the performance consequences of physiological divergence between Puerto

Rican sister species pairs, we calculated the expected physiological performance of each

species under natural warm and cool temperature regimes found on Puerto Rico. Two

primary habitat types on Puerto Rico are shaded, cool mesic forest and more open, warm

xeric forest (Ewel and Whitmore 1973). All four "warm niche" species in the species

pairs occur in xeric forest, and all four "cool niche" species in the species pairs occur in

mesic forest. In addition, two of the four "warm niche" species (*A. pulchellus* and *A.*

stratulus) can occur in mesic forest where they favor open areas (Hertz 1979 and personal

observation), and one "cool niche" species (*A. cristatellus*) can occur in xeric forest

where it favors shaded areas (Gunderson and Leal 2012). From a previous study

(Gunderson and Leal 2012), we have a data set of operative temperature distributions

from mesic and xeric forest measured with copper lizard models (Hertz 1992a) during the

summer, the peak of the breeding season for Puerto Rican anoles (Gorman and Licht

1974). The operative temperature distribution is a quantitative description of a thermal

environment from the organism's perspective (Hertz et al. 1993) and can be thought of as

a distribution of potential body temperatures in each habitat. Mean operative temperature

in the mesic habitat is significantly cooler than that in the xeric habitat, by 4.5°C

(Gunderson and Leal 2012).

We estimated the physiological performance in the mesic and xeric operative

thermal environments using two metrics: 1) mean relative physiological performance

capacity (Hertz et al. 1993; Huey et al. 2009; Gunderson and Leal 2012), and 2) threat of

overheating, the percentage of operative temperature observations over CT_{max}. To

estimate mean relative performance capacity, we generated a thermal performance curve

for each species using our sprint performance and CT_{max} data sets. Thermal performance

curves were a combination of two functions: one describing performance below T_{opt}, and

the other describing performance from T_{opt} to CT_{max} (Deutsch et al. 2008; Huey et al.

2009; Gunderson and Leal 2012). The former function was a second-order polynomial fit

to the sprint data of each species. The latter function took the form:

$P(T) = 1.0 - \left(\dfrac{T - T_{opt}}{CT_{max} - T_{opt}} \right)^2$ (Huey et al. 2009; Gunderson and Leal 2012). Fitted curves for

each species can be seen in Fig. S3. The curves were scaled to represent relative

performance by setting sprint speed at T_{opt} to 1.0. With these curves, we calculated the

mean relative physiological performance capacity of each species in each habitat (Hertz

et al. 1993; Huey et al. 2009; Gunderson and Leal 2012). Theory predicts that relative

performance is expected to remain high as species adapt to different thermal

environments (Angilletta et al. 2006; Martin and Huey 2008).

RATES OF EVOLUTION

To compare rates of evolution of thermal niche traits to morphological traits involved in

structural niche partitioning (*sensu* Hertz et al. 2013), we used a phylogenetic

comparative approach. Rate of evolution of heat tolerance (CT_{max}) was compared to rates

for body size (snout-vent-length, or SVL), head length, femur length, and adhesive

toepad width for the 4[th] hindtoe. We used the maximum clade credibility phylogeny of

Mahler et al. (2010) for these analyses. Morphological data are also from Mahler et al.

(2010) and are described in greater detail therein. Shape data (head length, femur length,

and toepad width) were size-corrected using phylogenetic regression in phytools (Revell

2012) prior to analysis. We used the phylogenetic approach described by Adams (2013)

to conduct pairwise rate comparisons for all combinations of these five traits. For each

comparison, we used likelihood ratio tests to compare a model in which the two traits

were free to evolve at different rates to a model in which the traits were constrained to

evolve at a single rate. Rates were estimated using species means of natural log-

transformed values for all traits (such that variation is represented in terms of

proportional change) and permitting co-estimation of among-trait covariances (similar

results were obtained with covariances fixed at zero).

Results

PHYSIOLOGICAL DIVERGENCE BETWEEN PUERTO RICAN SPECIES

PAIRS

Physiological divergence is apparent in at least one trait (T_{opt}, CT_{max}, or both) between all

sister species pairs on Puerto Rico, and in all cases the pattern of divergence is in the

direction predicted based on the species' thermal niches.

Significant differences in T_{opt} were observed for the *A. evermanni*/*A. stratulus*

and *A. cristatellus*/*A. cooki* species pairs (*i.e.*, no overlap in 95% confidence intervals),

but not in the *A. poncensis*/*A. gundlachi* or *A. pulchellus*/*A. krugi* species pairs (Fig. 2A).

However, it should be noted that our analysis almost certainly underestimates the T_{opt} of

A. cooki and *A. stratulus*, which can be seen by looking at Fig. 2A; in both of these

species, nearly all of the individuals had a T_{opt} of 35°C solely because that was our

maximum test temperature.

CT_{max} differed significantly among Puerto Rican species ($F_{7,72} = 42.74$; $P <$

0.001). Planned orthogonal contrasts demonstrated significant divergence in three of the

four species pairs: *A. evermanni*/*A. stratulus*, *A. poncensis*/*A. gundlachi*, and *A.*

pulchellus/*A. krugi* (all $P < 0.001$), but not in the *A. cristatellus*/*A. cooki* species pair ($P =$

0.490; Fig. 2B).

PHYSIOLOGICAL PERFORMANCE IN DIFFERENT THERMAL REGIMES

For three of the four sister-species pairs, the cool niche species is predicted to have higher

physiological performance capacity throughout the day within the shaded operative

thermal environment, with the mean performance advantage being 8-14% depending on

the species pair and time of day (Fig. 3). The lone exception occurs with the *A.*

pulchellus/*A. krugi* sister species pair, for which performance estimates are similar for

both species throughout the day (Fig. 3G). In the open habitat operative environment, the

warm niche species tend to have a performance advantage, particularly during midday

hours (from 10:00-13:00; Fig. 3). The exception in this case is the *A. cristatellus*/*A. cooki*

species pair, for which *A. cristatellus* has higher predicted performance capacity early

and late in the day, and both species have similar predicted performance during the

midday hours (Fig. 3D). However, midday performance of *A. cooki* is likely

underestimated due to our probable underestimation of the T_{opt} of this species (see

above).

None of the species are under threat of overheating in the shaded operative

thermal environment (Fig. 4A, C, E, G). However, all species have a threat of

overheating in the open habitat operative thermal environment (Fig. 4B, D, F, H). For

three of the four sister-species pairs (*A. stratulus*/*A. evermanni*, *A. poncensis*/*A.*

gundlachi, and *A. pulchellus*/*A. krugi*), the cool niche species has a higher probability of

overheating throughout the day (Fig. 4B, F, H). The difference in probability of

overheating is particularly acute during the midday hours, when the cool niche species in

these sister-species pairs have probabilities of overheating that are at least twice that of

warm niche species (all of which are significant differences; G-tests, all $P < 0.05$). The

only sister-species pair for which this relationship does not hold is *A. cristatellus*/*A.*

cooki, for which there was no divergence in CT_{max} (Fig. 4D).

PHYSIOLOGICAL EVOLUTION WITHIN THE JAMAICAN RADIATION

Significant differences in thermal physiology were also present among Jamaican species.

Mean T_{opt} ranges from approximately 29.5 to 34.6°C (Fig. 5A; true T_{opt}s of *A. grahami*

and *A. opalinus* likely underestimated due to our experimental upper measurement limit

of 35°C, see above), while CT_{max} ranges from 35.8-41.5°C among species (Fig. 5B;

ANOVA, $F_{3,36} = 13.6$, $P < 0.001$).

PHYSIOLOGICAL DIFFERENCES BETWEEN SYMPATRIC SPECIES

Across Puerto Rico and Jamaica, sympatric species that occupy the same structural niche

(*i.e.*, are morphologically similar) invariably differ in thermal physiology (Table 1).

Three of these sympatric pairs are sister species from Puerto Rico described above (*i.e.*,

the trunk-crown ecomorph species *A. stratulus*/*A. evermanni*; the trunk-ground species *A.*

cristatellus/*A. cooki;* the grass-bush species *A. krugi*/*A. pulchellus*). The trunk-ground

species *A. cristatellus* and *A. gundlachi* also co-occur in some areas of Puerto Rico and

differ in CT_{max} (P<0.001). On Jamaica, the trunk-ground anole *A. lineatopus* and trunk-

crown anole *A. grahami* differ in CT_{max}, as do *A. lineatopus* and the introduced trunk-

ground anole *A. sagrei* (both P<0.001; Table 1).

RATES OF PHYSIOLOGICAL AND MORPHOLOGICAL EVOLUTION

Thermal tolerance has evolved more slowly than some, but not all, morphological traits

associated with ecomorphological divergence (Table 2). CT_{max} evolved significantly

more slowly than body size (SVL; $P < 0.001$) and relative limb length (femur; $P =$

0.029). No differences in evolutionary rates were found between CT_{max} and either head

length ($P = 0.105$) or toepad width ($P = 0.112$).

Discussion

Evolution of the climatic niche has emerged as an important component of many

evolutionary radiations, but the functional consequences of climate niche divergence are

generally unknown. We demonstrate here that climatic niche divergence among species

in the *Anolis* adaptive radiation is accompanied by physiological divergence in both

temperature tolerance and sensitivity to sub-lethal temperature variability. Species that

occupy warmer niches have either higher heat tolerance, higher optimal temperatures for

physiological performance, or both, compared to species that occupy cooler thermal

niches. Such physiological diversification has occurred repeatedly, as is most easily seen

among four sister-species pairs present on Puerto Rico that differ in thermal niche (Fig.

2). Divergence in thermal physiology is also present on Jamaica, with species spanning

essentially the full range of heat tolerances and optimal temperatures found on Puerto

Rico (Fig. 5).

The repeated divergence of thermal physiology among species occupying distinct

thermal niches is itself evidence of adaptation (Harvey and Pagel 1991). However, we

also provide more direct evidence of the adaptive significance of physiological

divergence by estimating physiological performance parameters for Puerto Rican species

based on detailed measurements of warm and cool thermal environments found on the

island. In shaded habitat, cool niche species have higher predicted mean performance

capacities than their warm niche sister species in three of the four species pairs (Fig. 3).

The only exception is the *A. krugi*/*A. pulchellus* species pair, for which there was no

divergence in optimal temperature. Within the warmer open habitat the pattern was less

pronounced, with warm niche species appearing to have a sizable performance advantage

in only two of the species pairs (Fig. 3). The advantage of physiological divergence for

warm-niche species becomes much more apparent when considering overheating risk.

During midday hours in the warmer open habitat, warm niche species in three of the four

species pairs had less than half the overheating risk of cool-niche species (Fig. 4). The

lone exception was the *A. cristatellus*/*A. cooki* species pair, which had no divergence in

thermal tolerance (Fig. 2B). No species was at risk of overheating in the cooler, shaded

habitat (Fig. 4).

Our analysis of physiological performance in different thermal environments

assumes that animals are randomly utilizing perch sites with respect to available

operative temperatures. Anoles are known to behaviorally thermoregulate (Hertz et al.

1993), in effect modulating the thermal environment that they experience by avoiding

perch sites with operative temperatures that are physiologically detrimental (Huey et al.

2003). However, behavioral thermoregulation is costly, and animals living within

environments in which they do not need to make behavioral adjustments should be at a

substantial advantage (Huey 1974; Huey and Slatkin 1976; Sears et al. 2011; Sears and

Angilletta 2015). Thus, cool niche lizards in more open areas are likely to have reduced

space and time available for normal activity compared to the more heat-tolerant warm

niche species, and the same will hold true for warm-niche lizards in cooler areas

(Gunderson and Leal 2015, 2016).

The high community-level species diversity associated with many adaptive

radiations, including *Anolis*, is at least partially enabled by fine-scale resource

partitioning among constituent taxa (Schluter 2000). Sympatric anoles are known to

partition thermal microhabitats (Ruibal 1961; Rand 1964), and our analysis indicates that

this is facilitated by physiological divergence. When anole species that share the same

structural niche (*i.e.*, share the same or similar ecomorphology) co-occur, they invariably

differ in temperature-dependent physiological traits (Table 1). This pattern is not only

maintained for ecologically similar species that are endemic members of the community,

but also for introduced species: in Jamaica, the endemic trunk-ground species *A.*

lineatopus is found in sympatry with the introduced trunk-ground species *A. sagrei*, and

these species differ significantly in thermal tolerance (Table 1).

We found that physiological thermal tolerance has evolved more slowly than

some morphological traits (SVL and femur length), and has evolved at similar rates to

other morphological traits (head length and toe-pad width; Table 2). This result is at odds

with a recent analysis which found that divergence in thermal physiology, estimated as

field body temperature, occurs more quickly than morphological divergence in anoles

(Hertz et al. 2013). However, we note that in that study, estimated rates for thermal and

morphological traits are not directly comparable because they were not standardized to

the same scale before analysis (Adams 2013). Our results demonstrate that high rates of

evolutionary change are not necessary for traits to be ecologically important, as the

relatively slow divergence that we have observed in anole thermal tolerance has

important performance consequences with respect to overheating risk in warm habitats

(Fig. 4).

A critical question when assessing patterns of evolution is determining the

potential for observed divergence to be driven by phenotypic plasticity as opposed to

heritable genetic change. With respect to thermal physiology, direct evidence from anoles

and broader patterns from reptiles indicate that the magnitude of divergence that we have

observed is unlikely to be the result of acclimation to different thermal regimes. For

example, metabolic rates of the Puerto Rican *A. cristatellus* and *A. gundlachi*, as well as

A. sagrei introduced to Florida, do not change in response to acclimation temperature

(Rogowitz 1996b, a). Tests for acclimation in thermal tolerance in Greater Antillean

anoles have also suggested no (Leal and Gunderson 2012; Muñoz et al. 2014) or limited

(Kolbe et al. 2012) plasticity in this physiological trait. These results are consistent with

two recent meta-analyses of plasticity in thermal physiology, both of which demonstrated

that reptiles have among the lowest acclimation capacities of all ectotherms (Gunderson

and Stillman 2015; Seebacher et al. 2015). For example, among reptiles CT_{max} changes an

average of only 0.1°C for every 10°C change in mean body temperature (Gunderson and

Stillman 2015).

We have provided comprehensive evidence for the importance of physiological

divergence in the Puerto Rican and Jamaican *Anolis* adaptive radiations. However, the

importance of physiological divergence is likely not restricted to these islands. For

example, divergence in thermal tolerances have been demonstrated for trunk-ground

anoles of the *cybotes* group that occur along an elevational gradient in Hispaniola (Hertz

and Huey 1981; Muñoz et al. 2014), and variation in thermal physiology has also been

demonstrated among some mainland anoles (van Berkum 1986; Logan et al. 2013).

Physiological divergence is also not restricted to thermal traits, as there is significant

divergence in water loss rates among species and populations of anoles, which is often

correlated with thermal niche (Sexton and Heatwole 1968; Hillman and Gorman 1977;

Hertz et al. 1979; Hillman et al. 1979; Hertz 1980; Dmi'el et al. 1997; Gunderson et al.

2011). Though more work is required, the general pattern emerging is that physiological

divergence along the climatic niche axis may be as important as morphological

divergence along the structure niche axis in this classic radiation.

Our results highlight the benefits of a functional approach to investigating

climatic niche evolution. By focusing on physiological traits that link organismal

performance to abiotic conditions, we were able to demonstrate how phenotypes evolve

in response to climatic environments and estimate the performance consequences of

observed divergence in natural habitats. In addition, our functional perspective allowed

us to determine the contribution of climatic niche divergence to microhabitat partitioning

and species co-existence. We suggest that the current trend to use ENMs, which contain

no information about organismal phenotypes or fine-scale habitat use, should only be a

first step in a more comprehensive research program to determine how and why climatic

niche divergence contributes to evolutionary radiations.

Acknowledgements

David Steinberg provided help in the field. Logistical support in Puerto Rico was

provided by the El Verde Biological Station and Mata de Plátano Field Station. We also

thank the owners of Ridge to Reef farm in St. Croix and Green Castle Estates in Jamaica

for allowing us to stay and work on their properties. Permits were provided by The

Departmento de Recursos Naturales y Ambientales of Puerto Rico, the St. Croix Division

of Fish and Wildlife, and the Jamaican National Environment and Planning Agency.

Funding was provided by a NSF Doctoral Dissertation Improvement Grant (1110570) to

A.R.G.

References

Adams, D. C. 2013. Quantifying and comparing phylogenetic evolutionary rates for
shape and other high-dimensional phenotypic data. Systematic Biology
62:181-192.
Algar, A. C. and D. L. Mahler. 2015. Area, climate heterogeneity, and the response of
climate niches to ecological opportunity in island radiations of *Anolis* lizards.
Global Ecology and Biogeography doi:10.1111 /geb.12327.
Alvarado‑Serrano, D. F. and L. L. Knowles. 2014. Ecological niche models in
phylogeographic studies: applications, advances and precautions. Molecular
Ecology Resources 14:233-248.
Angilletta, M. J. 2009. Thermal adaptation: a theoretical and empirical synthesis.
Oxford University Press.
Angilletta, M. J., A. F. Bennett, H. Guderley, C. A. Navas, F. Seebacher, and R. S. Wilson.
2006. Coadaptation: A Unifying Principle in Evolutionary Thermal Biology.
Physiological and Biochemical Zoology 79:282-294.
Araújo, M. B. and A. T. Peterson. 2012. Uses and misuses of bioclimatic envelope
modeling. Ecology 93:1527-1539.
Arnegard, M. E., P. B. McIntyre, L. J. Harmon, M. L. Zelditch, W. G. Crampton, J. K.
Davis, J. P. Sullivan, S. Lavoué, and C. D. Hopkins. 2010. Sexual signal
evolution outpaces ecological divergence during electric fish species
radiation. The American Naturalist 176:335-356.
Bogert, C. M. 1949. Thermoregulation in reptiles, a factor in evolution. Evolution
3:195-211.
Bonino, M. F., D. L. M. Azócar, M. J. Tulli, C. S. Abdala, M. G. Perotti, and F. B. Cruz.
2011. Running in cold weather: morphology, thermal biology, and
performance in the southernmost lizard clade in the world (*Liolaemus*
lineomaculatus section: Liolaemini: Iguania). Journal of Experimental Zoology
Part A: Ecological Genetics and Physiology 315:495-503.
Buckley, L. B., M. C. Urban, M. J. Angilletta, L. G. Crozier, L. J. Rissler, and M. W. Sears.
2010. Can mechanism inform species' distribution models? Ecology Letters
13:1041-1054.
Deutsch, C. A., J. J. Tewksbury, R. B. Huey, K. S. Sheldon, C. K. Ghalambor, D. C. Haak,
and P. R. Martin. 2008. Impacts of climate warming on terrestrial ectotherms
across latitude. Proceedings of the National Academy of Sciences 105:6668-
6672.
Dmi'el, R., G. Perry, and J. Lazell. 1997. Evaporative water loss in nine insular
populations of the lizard *Anolis cristatellus* group in the British Virgin Islands.
Biotropica 29:111-116.
Efron, B. 1987. Better bootstrap confidence intervals. Journal of the American
Statistical Association 82:171-185.
Ewel, J. J. and J. L. Whitmore. 1973. Ecological life zones of Puerto Rico and US Virgin
Islands. Forest Service Research Paper ITF-18. U.S. Department of
Agriculture.

Fleishman, L. J., M. Leal, and M. H. Persons. 2009. Habitat light and dewlap color
diversity in four species of Puerto Rican anoline lizards. Journal of
Comparative Physiology A 195:1043-1060.
Gamble, T., A. J. Geneva, R. E. Glor, and D. Zarkower. 2014. *Anolis* sex chromosomes
are derived from a single ancestral pair. Evolution 68:1027-1041.
Gorman, G. C. and P. Licht. 1974. Seasonality in ovarian cycles among tropical *Anolis*
lizards. Ecology 55:360-369.
Grant, P. R. 1999. Ecology and evolution of Darwin's finches. Princeton University
Press.
Guisan, A. and W. Thuiller. 2005. Predicting species distribution: offering more than
simple habitat models. Ecology Letters 8:993-1009.
Gunderson, A. R. and M. Leal. 2012. Geographic variation in vulnerability to climate
warming in a tropical Caribbean lizard. Functional Ecology 26:783-793.
Gunderson, A. R. and M. Leal. 2015. Patterns of thermal constraint on ectotherm
activity. The American Naturalist 185:653-664.
Gunderson, A. R. and M. Leal. 2016. A conceptual framework for understanding
thermal constraints on ectotherm activity with implications for predicting
responses to global change. Ecology Letters 19:111-120.
Gunderson, A. R., J. Siegel, and M. Leal. 2011. Tests of the contribution of acclimation
to geographic variation in water loss rates of the West Indian lizard *Anolis*
cristatellus. Journal of Comparative Physiology B 181:965-972.
Gunderson, A. R. and J. H. Stillman. 2015. Plasticity in thermal tolerance has limited
potential to buffer ectotherms from global warming. Proceedings of the Royal
Society of London B: Biological Sciences 282:20150401.
Harvey, P. H. and M. D. Pagel. 1991. The comparative method in evolutionary
biology. Oxford University Press.
Heatwole, H., T.-H. Lin, E. Villalón, A. Muñiz, and A. Matta. 1969. Some aspects of the
thermal ecology of Puerto Rican anoline lizards. Journal of Herpetology 3:65-
77.
Hertz, P., A. Arce-Hernandez, J. Ramirez-Vazquez, W. Tirado-Rivera, and L. Vazquez-
Vives. 1979. Geographical variation of heat sensitivity and water loss rates in
the tropical lizard, *Anolis gundlachi*. Comparative Biochemistry and
Physiology Part A: Physiology 62:947-953.
Hertz, P. E. 1979. Sensitivity to high temperatures in three west Indian grass anoles
(sauria, iguanidae), with a review of heat sensitivity in the genus< i>
Anolis</i>. Comparative Biochemistry and Physiology Part A: Physiology
63:217-222.
Hertz, P. E. 1980. Responses to dehydration in *Anolis* lizards sampled along
altitudinal transects. Copeia 1980:440-446.
Hertz, P. E. 1992a. Evaluating thermal resource partitioning by sympatric lizards
Anolis cooki and *A. cristatellus*: a field test using null hypotheses. Oecologia
90:127-136.
Hertz, P. E. 1992b. Temperature regulation in Puerto Rican *Anolis* lizards: a field test
using null hypotheses. Ecology 73:1405-1417.

Hertz, P. E., Y. Arima, A. Harrison, R. B. Huey, J. B. Losos, and R. E. Glor. 2013.
Asynchronous evolution of physiology and morphology in *Anolis* lizards.
Evolution 67:2101-2113.
Hertz, P. E. and R. B. Huey. 1981. Compensation for altitudinal changes in the
thermal environment by some *Anolis* lizards on Hispaniola. Ecology 62:515-
521.
Hertz, P. E., R. B. Huey, and R. Stevenson. 1993. Evaluating temperature regulation
by field-active ectotherms: the fallacy of the inappropriate question. The
American Naturalist 142:796-818.
Hillman, S. and G. Gorman. 1977. Water loss, desiccation tolerance, and survival
under desiccating conditions in 11 species of Caribbean *Anolis*. Oecologia
29:105-116.
Hillman, S., G. C. Gorman, and R. Thomas. 1979. Water loss in *Anolis* lizards: evidence
for acclimation and intraspecific differences along a habitat gradient.
Comparative Biochemistry and Physiology Part A: Physiology 62:491-493.
Hochachka, P. W. and G. N. Somero. 2002. Biochemical Adaptation: Mechanism and
process in physiological evolution. Oxford University Press.
Huey, R. B. 1974. Behavioral thermoregulation in lizards: importance of associated
costs. Science 184:101-103.
Huey, R. B. 1983. Natural variation in body temperature and physiological
performance in a lizard (*Anolis cristatellus*). Pp. 484-490. Advances in
herpetology and evolutionary biology: essays in honor of Ernest E. Williams.
Huey, R. B., C. A. Deutsch, J. J. Tewksbury, L. J. Vitt, P. E. Hertz, H. J. Á. Pérez, and T.
Garland. 2009. Why tropical forest lizards are vulnerable to climate warming.
Proceedings of the Royal Society B: Biological Sciences 276:1939-1948.
Huey, R. B., P. E. Hertz, and B. Sinervo. 2003. Behavioral drive versus behavioral
inertia in evolution: a null model approach. The American Naturalist
161:357-366.
Huey, R. B. and M. Slatkin. 1976. Cost and benefits of lizard thermoregulation.
Quarterly Review of Biology 51:363-384.
Huey, R. B. and T. P. Webster. 1976. Thermal biology of *Anolis* lizards in a complex
fauna: the *cristatellus* group on Puerto Rico. Ecology 57:985-994.
Jackman, T. R., A. Larson, K. De Queiroz, and J. B. Losos. 1999. Phylogenetic
relationships and tempo of early diversification in *Anolis* lizards. Systematic
Biology 48:254-285.
Kaspari, M., N. A. Clay, J. Lucas, S. P. Yanoviak, and A. Kay. 2015. Thermal adaptation
generates a diversity of thermal limits in a rainforest ant community. Global
Change Biology 21:1092-1102.
Kearney, M. and W. Porter. 2009. Mechanistic niche modelling: combining
physiological and spatial data to predict species' ranges. Ecology letters
12:334-350.
Keller, I. and O. Seehausen. 2012. Thermal adaptation and ecological speciation.
Molecular Ecology 21:782-799.
Kellermann, V., B. van Heerwaarden, C. M. Sgrò, and A. A. Hoffmann. 2009.
Fundamental evolutionary limits in ecological traits drive *Drosophila* species
distributions. Science 325:1244-1246.

Knouft, J. H., J. B. Losos, R. E. Glor, and J. J. Kolbe. 2006. Phylogenetic analysis of the
615   evolution of the niche in lizards of the *Anolis sagrei* group. Ecology 87:S29-
616   S38.
617   Kolbe, J. J., P. S. VanMiddlesworth, N. Losin, N. Dappen, and J. B. Losos. 2012. Climatic
618   niche shift predicts thermal trait response in one but not both introductions
619   of the Puerto Rican lizard *Anolis cristatellus* to Miami, Florida, USA. Ecology
620   and Evolution 2:1503-1516.
621   Kozak, K. H. and J. J. Wiens. 2007. Climatic zonation drives latitudinal variation in
622   speciation mechanisms. Proceedings of the Royal Society of London B:
623   Biological Sciences 274:2995-3003.
624   Labra, A., J. Pienaar, and T. F. Hansen. 2009. Evolution of thermal physiology in
625   *Liolaemus* lizards: adaptation, phylogenetic inertia, and niche tracking. The
626   American Naturalist 174:204-220.
627   Lawing, A. M., P. D. Polly, D. K. Hews, E. P. Martins, D. C. Collar, and J. L. Bronstein.
628   2016. Including Fossils in Phylogenetic Climate Reconstructions: A Deep
629   Time Perspective on the Climatic Niche Evolution and Diversification of
630   Spiny Lizards (*Sceloporus*). The American Naturalist 188:000-000.
631   Leal, M. and A. R. Gunderson. 2012. Rapid change in the thermal tolerance of a
632   tropical lizard. The American Naturalist 180:815-822.
633   Levin, L. A., R. J. Etter, M. A. Rex, A. J. Gooday, C. R. Smith, J. Pineda, C. T. Stuart, R. R.
634   Hessler, and D. Pawson. 2001. Environmental influences on regional deep-
635   sea species diversity. Annual Review of Ecology and Systematics 32:51-93.
636   Logan, M. L., R. M. Cox, and R. Calsbeek. 2014. Natural selection on thermal
637   performance in a novel thermal environment. Proceedings of the National
638   Academy of Sciences 111:14165-14169.
639   Logan, M. L., R. K. Huynh, R. A. Precious, and R. G. Calsbeek. 2013. The impact of
640   climate change measured at relevant spatial scales: new hope for tropical
641   lizards. Global Change Biology 19:3093-3102.
642   Losos, J. B. 1990. Ecomorphology, performance capability, and scaling of West
643   Indian *Anolis* lizards: an evolutionary analysis. Ecological Monographs
644   60:369-388.
645   Losos, J. B. 2009. Lizards in an evolutionary tree: ecology and adaptive radiation of
646   anoles. University of California Press.
647   Losos, J. B., T. R. Jackman, A. Larson, K. de Queiroz, and L. Rodríguez-Schettino. 1998.
648   Contingency and determinism in replicated adaptive radiations of island
649   lizards. Science 279:2115-2118.
650   Losos, J. B., M. Leal, R. E. Glor, K. de Queiroz, P. E. Hertz, L. R. Schettino, A. C. Lara, T.
651   R. Jackman, and A. Larson. 2003. Niche lability in the evolution of a Caribbean
652   lizard community. Nature 424:542-545.
653   Mahler, D. L., T. Ingram, L. J. Revell, and J. B. Losos. 2013. Exceptional convergence
654   on the macroevolutionary landscape in island lizard radiations. Science
655   341:292-295.
656   Mahler, D. L., L. J. Revell, R. E. Glor, and J. B. Losos. 2010. Ecological opportunity and
657   the rate of morphological evolution in the diversification of Greater Antillean
658   anoles. Evolution 64:2731-2745.

Martin, T. L. and R. B. Huey. 2008. Why "suboptimal" is optimal: Jensen's inequality
and ectotherm thermal preferences. The American Naturalist 171:E102-
E118.
Matute, D. R., C. J. Novak, and J. A. Coyne. 2009. Temperature‐based extrinsic
reproductive isolation in two species of *Drosophila*. Evolution 63:595-612.
Meyer, A. 1993. Phylogenetic relationships and evolutionary processes in East
African cichlid fishes. Trends in Ecology & Evolution 8:279-284.
Muñoz, M. M., M. A. Stimola, A. C. Algar, A. Conover, A. J. Rodriguez, M. A. Landestoy,
G. S. Bakken, and J. B. Losos. 2014. Evolutionary stasis and lability in thermal
physiology in a group of tropical lizards. Proceedings of the Royal Society B:
Biological Sciences 281:20132433.
Nicholson, K. E., R. E. Glor, J. J. Kolbe, A. Larson, S. Blair Hedges, and J. B. Losos. 2005.
Mainland colonization by island lizards. Journal of Biogeography 32:929-938.
Peterson, A. T., J. Soberón, R. G. Pearson, R. P. Anderson, M. Nakamura, E. Martinez-
Meyer, and M. B. Araújo. 2011. Ecological niches and geographic
distributions. Princeton University Press, Princeton, New Jersey, USA.
Pincheira-Donoso, D. 2011. Predictable variation of range-sizes across an extreme
environmental gradient in a lizard adaptive radiation: evolutionary and
ecological inferences. PloS One 6:e28942.
Potter, K. A., H. A. Woods, and S. Pincebourde. 2013. Microclimatic challenges in
global change biology. Global change biology 19:2932–2939.
Quintero, I. and J. J. Wiens. 2013. Rates of projected climate change dramatically
exceed past rates of climatic niche evolution among vertebrate species.
Ecology Letters 16:1095-1103.
Rand, A. S. 1964. Ecological distribution in anoline lizards of Puerto Rico. Ecology
45:745-752.
Revell, L. J. 2012. phytools: an R package for phylogenetic comparative biology (and
other things). Methods in Ecology and Evolution 3:217-223.
Rogowitz, G. L. 1996a. Evaluation of thermal acclimation and altitudinal variation of
metabolism in a Neotropical lizard, *Anolis gundlachi*. Copeia 1996:535-542.
Rogowitz, G. L. 1996b. Evaluation of thermal acclimation of metabolism in two
eurythermal lizards, *Anolis cristatellus* and *A. sagrei*. Journal of Thermal
Biology 21:11-14.
Ruibal, R. 1961. Thermal relations of five species of tropical lizards. Evolution 15:98-
111.
Schluter, D. 2000. The ecology of adaptive radiation. Oxford University Press.
Sears, M. W. and M. J. Angilletta. 2015. Costs and benefits of thermoregulation
revisited: both the heterogeneity and spatial structure of temperature drive
energetic costs. The American Naturalist 185:E94-E102.
Sears, M. W., E. Raskin, and M. J. Angilletta. 2011. The world is not flat: defining
relevant thermal landscapes in the context of climate change. Integrative and
Comparative Biology 51:666-675.
Seebacher, F., C. R. White, and C. E. Franklin. 2015. Physiological plasticity increases
resilience of ectothermic animals to climate change. Nature Climate Change
5:61-66.

Seehausen, O., Y. Terai, I. S. Magalhaes, K. L. Carleton, H. D. Mrosso, R. Miyagi, I. van
der Sluijs, M. V. Schneider, M. E. Maan, and H. Tachida. 2008. Speciation
through sensory drive in cichlid fish. Nature 455:620-626.
Sexton, O. J. and H. Heatwole. 1968. An experimental investigation of habitat
selection and water loss in some anoline lizards. Ecology 49:762-767.
Simpson, G. G. 1953. Major features of evolution. Columbia University Press, New
York.
Stillman, J. H. and G. N. Somero. 2000. A comparative analysis of the upper thermal
tolerance limits of eastern Pacific porcelain crabs, genus *Petrolisthes*:
influences of latitude, vertical zonation, acclimation, and phylogeny.
Physiological and Biochemical Zoology 73:200-208.
Sunday, J. M., A. E. Bates, and N. K. Dulvy. 2011. Global analysis of thermal tolerance
and latitude in ectotherms. Proceedings of the Royal Society B: Biological
Sciences 278:1823-1830.
van Berkum, F. H. 1986. Evolutionary patterns of the thermal sensitivity of sprint
speed in *Anolis* lizards. Evolution 40:594-604.
Wake, D. B. 1987. Adaptive radiation of salamanders in Middle American cloud
forests. Annals of the Missouri Botanical Garden 74:242-264.
Warren, D. L., M. Cardillo, D. F. Rosauer, and D. I. Bolnick. 2014. Mistaking geography
for biology: inferring processes from species distributions. Trends in Ecology
& Evolution 29:572-580.
Warren, D. L., R. E. Glor, and M. Turelli. 2008. Environmental niche equivalency
versus conservatism: quantitative approaches to niche evolution. Evolution
62:2868-2883.
Williams, E. E. 1972. The origin of faunas. Evolution of lizard congeners in a complex
island fauna: a trial analysis. Evolutionary Biology 6:47-89.
Williams, E. E. 1983. Ecomorphs, faunas, island size, and diverse end points in island
radiations of Anolis. Pp. 326-370. Lizard ecology: studies of a model
organism. Harvard University Press, Cambridge.
733
734

Table 1. Physiology differences between co-occurring species that occupy the same or

similar structural niches (*i.e.*, share the same or similar ecomorphology). For CT_{max}, P-

values are given based on one-way ANOVA with planned orthogonal contrasts. For T_{opt},

"*" indicates that the bootstrapped 95% confidence intervals did not overlap, "N.S."

indicates no significant difference. "†" indicates that the species is an introduced member

of the community.

Table 2. Estimated rates of evolution for heat tolerance (CT_{max}) and morphological traits

related to ecomorphological divergence. Rate estimates (bold) are in the diagonal of the

matrix. Upper off-diagonals contain estimates of pairwise evolutionary covariances

among traits, lower off-diagonals contain P-values for likelihood ratio test comparisons

of 2-rate versus 1-rate models for each trait pair. P-values less than 0.05 indicate support

for a 2-rate model over a 1-rate model.

765
766

770   Table 1

	Co-occurring species	Ecomorph	Physiological divergence	
			CT_{max}	T_{opt}
A. stratulus	*A. evermani*	Trunk-crown	<0.001	$*$
A. cristatellus	*A. cooki*	Trunk-ground	0.490	$*$
A. cristatellus	*A. gundlachi*	Trunk-ground	<0.001	N.S.
A. krugi	*A. pulchellus*	Grass-bush	<0.001	N.S.
A. lineatopis	*A. grahami*	Trunk-ground/Trunk-crown	<0.001	N.S.
A. lineatopis	*A. sagrei*[†]	Trunk-ground	<0.001	N.S.

779 Table 2.

	CT_{max}	SVL	Head length	Femur length	Toepad width
CTmax	**5.80E-05**	-4.90E-05	-1.60E-05	-5.50E-05	-9.90E-06
SVL	<0.001	**2.30E-03**	8.90E-06	-1.80E-05	2.70E-05
Head length	0.105	<0.001	**2.70E-05**	1.30E-05	-6.30E-06
Femur length	0.029	<0.001	0.001	**1.40E-04**	2.20E-05
Toepad width	0.112	<0.001	0.003	0.832	**1.30E-04**

Figure 1: Phylogenetic relationships of the species included in this study, along with the

islands from which they were sampled.

Figure 2: Thermal physiology of four sister-species pairs of Puerto Rican anole. Each

species pair is represented in alternating grey and white regions of the plot with warm

niche species in red and cool niche species in blue. (A) Optimal sprint performance

temperatures (T_{opt}). (B) Heat tolerance limits (CT_{max}). "*", significant difference between

members of a species pair; N.S., no significant difference between members of a species

pair. See Methods for analysis details.

Figure 3: Predicted relative physiological performance of sister-species pairs of Puerto

Rican *Anolis* under shaded and open habitat operative thermal environments found on

Puerto Rico during morning, midday, and afternoon/evening hours. Warm niche species

in red, cool niche species in blue. (A, B) *A. stratulus*/*A. evermanni*, (C, D) *A. cooki*/*A.*

cristatellus, (E, F) *A. poncensis*/*A. gundlachi* (G, H) *A. pulchellus*/*A. krugi*.

Figure 4: Predicted overheating probability of sister-species pairs of Puerto Rican *Anolis*

under shaded and open habitat operative thermal environments found on Puerto Rico

during morning, midday, and afternoon/evening hours. Warm niche species in red, cool

niche species in blue. (A, B) *A. stratulus*/*A. evermanni*, (C, D) *A. cooki*/*A. cristatellus* (E,

F)*, A. poncensis*/*A. gundlachi*, (G, H) *A. pulchellus*/*A. krugi*.

Figure 5: Thermal physiology of endemic Jamaican anoles. (A) Optimal sprinting

temperature (T_{opt}). *Anolis valencienni* does not have a T_{opt} estimate because this species

could not be induced to run. (B) Heat tolerance (CT_{max}).

863   Figure 1.

871   Figure 2.

877
878

Time of day

881   Figure 3

Time of day

887   Figure 4

Figure 5