Thermal niche evolution across replicated *Anolis* lizard adaptive radiations

Alex R. Gunderson[†‡*], D. Luke Mahler[§], and Manuel Leal[ʃ]

†Romberg Tiburon Center and Department of Biology, San Francisco State University, 3150 Paradise

Dr., Tiburon, California 94920 and ‡Department of Integrative Biology, University of California,

Berkeley, California 94720-3140

*Corresponding author: e-mail: alexrgunderson@gmail.com

§Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, Ontario, M5S 3B2

CA

ʃ Division of Biological Sciences, University of Missouri, 105 Tucker Hall, Columbia, MO 65211.

Running title: Physiology of adaptive radiation

Keywords: *Anolis*, thermal biology, physiology, thermal tolerance, thermal performance, evolutionary

rates, sympatry, species richness

Data accessibility: If accepted, data herein will be archived at Dryad (http://datadryad.org)

Abstract

How evolutionary and ecological forces interact to promote the maintenance of biodiversity is a fundamental biological question. Here, we investigate this question by focusing on physiological evolution along the thermal niche axis in replicated radiations of *Anolis* lizards, groups that are best known for resource partitioning based on morphological evolution. We find repeated divergence in thermal physiology within these radiations, and that this divergence has performance consequences within natural habitats. In addition, morphologically similar species that co-occur invariably differ in their thermal physiology, providing evidence that physiological divergence facilitates species co-existence within these communities. Despite their ecological importance, phylogenetic comparative analyses suggest that thermal niche traits have evolved more slowly than key morphological traits related to the structural niche. Phylogenetic analyses also reveal that physiological (*i.e.*, fundamental) thermal niche evolution is correlated with realized thermal niche evolution based on geo-referenced locality data, but the latter incompletely predicts variation in the former. We provide comprehensive evidence for the importance of physiological divergence within *Anolis* adaptive radiations, including the complementary roles of physiological and morphological divergence in maintaining community-level diversity. We recommend greater integration of fundamental niche traits into analyses of climatic niche evolution, as they facilitate a more complete understanding of the phenotypic and ecological consequences of climatic divergence.

Introduction

A mechanistic understanding of how divergent phenotypic evolution facilitates species coexistence is a fundamental problem in ecology and evolution [1]. This problem comes into particular focus in the study of adaptive radiation, a process for which a hallmark is the repeated evolution of niche differences that allow closely related species to partition resources and stably co-occur [2-4]. Traditionally, most research on trait evolution during adaptive radiation has focused on morphological traits associated with dietary resource acquisition or structural habitat use, due to the clear role such adaptations play in mediating competition and facilitating coexistence [3, 5-7]. Physiological adaptation to abiotic conditions has received comparatively less attention in adaptive radiation studies, particularly in animals [3, 8]. Such adaptations are more typically studied for their role in generating divergence among geographically disjunct lineages experiencing different climates [9, 10]. However, physiological divergence can also play an important role in adaptive radiation by facilitating fine-scale partitioning of the local climatic niche [11], providing an additional resource axis along which a radiating clade may diversify [12, 13]. Indeed, such abiotic niche divergence has been hypothesized to underlie many impressive radiations, including bolitoglossine salamanders [14, 15], *Liolaemus* lizards [16-18], *Drosophila* flies [19], *Petrolisthes* crabs [20] and Lake Whitefish [21].

One of the most common tools applied in recent investigations of abiotic niche divergence is correlative ecological niche modeling (ENM), which estimates species' realized niches (*i.e.*, the range of conditions that species occupy) using information about species occurrences and broad-scale climate conditions [22-26]. Though correlative ENMs are powerful for addressing many questions, they also have characteristics that limit the inferences that can be made from their application [27-32]. With respect to adaptive radiations, one limitation is that correlative ENMs are based on geo-referenced species locality data and therefore provide no information about phenotypic traits (physiological or otherwise) linked to performance [28]. Performance-based traits relate to the fundamental niche (broadly, the range of conditions a species could occupy). Knowledge of performance-based traits is critical for understanding how species are able to occupy the habitats that they do with respect to both abiotic factors and other

species. ENMs may also be limited in what they can tell us about the role of abiotic niche divergence in

coexistence, as they cannot provide fine-scale information about micro-environmental niche use or abiotic

tolerances. Such factors can be of paramount importance for coexistence in radiating clades, as species

with identical or largely overlapping ranges may be able to partition resources via adaptation to different

thermal microenvironments [11, 33].

Here, we investigate the contribution of thermal niche divergence to the replicated *Anolis* lizard

adaptive radiations of Puerto Rico and Jamaica by analyzing patterns and performance consequences of

physiological divergence across 16 species (Fig. 1). Greater Antillean *Anolis* are best known for

morphological diversification along the fundamental structural niche axis [34-36], but morphological

evolution cannot fully explain the patterns of divergence and community composition observed across the

Greater Antilles [37]. Divergence in the thermal niche is also proposed to be an important component of

Anolis diversification based primarily on realized niche estimates from natural history and ENM studies

[12, 13, 22, 38-43]. However, we still know little about how physiology evolves during realized niche

divergence [44] and the extent to which physiological evolution facilitates species co-existence.

We first use data from the Puerto Rican *cristatellus* species group to investigate repeated

divergence in thermal physiology and performance between species that occupy distinct thermal

environments. Next, we integrate these data with measurements of operative thermal conditions in Puerto

Rico to explore the performance consequences of physiological divergence. We then consider spatial

patterns of physiological variation to evaluate the contribution of physiological evolution to fine-scale

niche partitioning and community-level species co-existence across Puerto Rico and Jamaica. Finally, we

conduct phylogenetic analyses of rates of physiological and morphological evolution to compare

diversification patterns between traits associated with the fundamental thermal niche and the fundamental

structural niche, and we employ phylogenetic methods to estimate the strength of the evolutionary

association between fundamental and realized thermal niche divergence. For the latter, we estimate

realized thermal niche characteristics of each species from geo-referenced bioclimatic data commonly

used in correlative ENM analyses.

Methods

We measured physiological traits for 304 individuals across 16 species from three islands (Fig. 1; see

Supplementary Table 1 for species information). Throughout, we focus primarily on the eight Puerto

Rican members of the *cristatellus* group, for two reasons. First, phylogenetic and natural history data

indicate that these species diversified *in situ* and form four sister species pairs [45-47], with each pair

representing an independent divergence in realized thermal niche based on the degree to which they

occupy open versus shaded perches (which throughout we refer to as "warm niche" and "cool niche"

species, respectively)[12, 40, 48, 49]. Second, we have detailed measurements of operative thermal

environments in Puerto Rico to investigate performance consequences of physiological divergence.

Experiments in Puerto Rico were conducted May 20 – June 20 2011, October 8-21 2011, and May 13-28

2012 at the Mata de Plátano field station. On Jamaica, we sampled five of the six extant endemic species,

plus the introduced *A. sagrei*. As with the Puerto Rican species, the endemic Jamaican anoles evolved *in*

situ from a single colonization event [50]. Experiments in Jamaica occurred from February 25 – March 13

2013 at Green Castle Estates, a privately owned farm and nature preserve in St. Mary Parish. Prior to

experiments, lizards were kept individually in plastic cages (18 x 11 x 15 cm) with a wooden dowel perch

on an approximately 12L:12D light schedule. Lizards were watered daily and fed crickets or Phoenix

worms every other day. The exception was *A. acutus* from St. Croix, which were brought to Durham, NC

and housed following Gunderson and Leal [51].

124   Heat tolerance (CT_{max}) was measured following Leal and Gunderson [52]. Briefly, lizards were

warmed under a heat lamp while monitoring their body temperatures with a wire thermocouple probe

placed inside the cloaca. Lizards were flipped onto their backs at one-degree intervals starting at 34°C,

and CT_{max} was recorded as the temperature at which they lost righting ability. Warming rates were very

similar among species, ranging from 1.8 to 2.6°C/min (Fig. S1). All lizards were captured three days prior

to CT_{max} measurement. We modeled variation in log-transformed CT_{max} among species using analysis of

variance (ANOVA). Planned orthogonal contrasts were applied to test *a priori* predictions about

divergence in CT_{max} between sister species (in Puerto Rico) and divergence among sympatric,

morphologically similar species (Puerto Rico and Jamaica).

Temperature-dependent sprint speeds were measured for thirteen species following Gunderson

and Leal [51]. Briefly, we analyzing high-speed video (120 frames/s) of lizards running up a 2 m wooden

racetrack set at a 37° angle and marked with a line every 12.5 cm. Lizards were run 2-4 times at each

temperature (two runs minimum, additional runs were added if the lizard stopped or jumped off of the

track during a trial). Speed for a lizard at a given temperature was taken as the fastest 25 cm interval at

that temperature [53, 54]. Puerto Rican lizards were run at five body temperatures in the following

randomized order: 32, 22, 27, 17, and 35°C. Conditions were the same for Jamaican lizards, except the

17°C temperature was excluded. All individuals were captured one day prior to the start of trials, and each

temperature treatment was administered on a different day over five straight days (four days for Jamaican

lizards). No animals were used for both heat tolerance and sprint performance. Target body temperatures

were achieved by placing lizards into a calibrated chilling-heating incubator (Cole-Parmer©, Vernon

Hills, IL, USA) for 30 min prior to a run [51].

The body temperature of maximum performance (the "optimal temperature," T_{opt}) was estimated

for each individual [sensu 55]. Thermal performance curves are non-linear [56], and we initially fit two

different non-linear models to the sprint data for each individual: a second order polynomial and a

modified Gaussian function ($P(T) = e^{-(\frac{T - T_{opt}}{2 \cdot \sigma})^2}$) commonly used to fit performance curves for ectotherms

[57], including *Anolis* sprinting [51, 58]. The Gaussian model provided relatively poor fits to the data

(much higher residual standard errors than the second-order polynomial model (Fig. S2)). Therefore, T_{opt}

estimates from the second-order polynomial models were used in analyses. T_{opt} estimates were

constrained such that T_{opt} had to occur within the range of experimental temperatures. This is a

conservative assumption that likely leads us to underestimate divergence in thermal physiology between

species (see Results). T_{opt} data were heavily skewed (see Results), so we compared T_{opt} among species by

assessing overlap in bootstrapped 95% confidence intervals (999 bootstrap replicates using the adjusted

bootstrap percentile method due to the skew in our data [59]). Analyses were conducted with the "boot"

and "boot.ci" functions, respectively, from the "boot" package in R.

We calculated the expected physiological performance of Puerto Rican species under natural

warm (open xeric forest) and cool (shaded mesic forest) operative thermal regimes found on Puerto Rico

[60]. Operative temperature distributions provide a quantitative description of a thermal environment

integrating the physical features of the organism [61], representing the distribution of available body

temperatures in a habitat. Operative temperature distributions came from a previous study [51] and were

measured with copper lizard models [49] during the peak breeding season (summer) for Puerto Rican

anoles [62]. Mean operative temperature in the mesic forest is significantly cooler than that in the xeric

forest, by 4.5°C [51].

We estimated two performance metrics under the different operative environments: 1) relative

sprint performance capacity [51, 58, 61], and 2) probability of overheating (the percentage of operative

temperature observations over CT_{max}). For relative performance capacity, we generated species-level

thermal performance curves using our sprint speed and CT_{max} data. The curves were a combination of two

functions fit to the data: one describing performance below T_{opt}, and the other describing performance

from T_{opt} to CT_{max} [51, 57, 58]. The former function was a second-order polynomial fit to the sprint data,

and the latter function took the form: $P(T) = 1.0 - \left(\dfrac{T - T_{opt}}{CT_{max} - T_{opt}} \right)^2$ [51, 58]. Curves were scaled to relative

performance by setting sprint speed at T_{opt} to 1.0 (see Fig. S3 for fitted curves). Performance in each

habitat was calculated by applying the curves to operative temperatures [51, 58, 61].

To compare rates of evolution of thermal physiology to ecomorphological traits involved in

fundamental structural niche partitioning, we used a phylogenetic comparative approach. Rate of

evolution of heat tolerance (CT_{max}) was compared to rates for body size (snout-vent-length, or SVL), head

length, femur length, and adhesive toepad width for the 4[th] hindtoe. We used the maximum clade

credibility phylogeny and morphological data of Mahler *et al*. [46] for these analyses. Shape data (head

length, femur length, and toepad width) were size-corrected using phylogenetic regression in phytools

[63] prior to analysis. We used the approach of Adams [64] to conduct pairwise rate comparisons for all

combinations of these five traits. We used likelihood ratio tests to compare a model in which the two

traits could evolve at different rates to a model in which the traits were constrained to evolve at a single

rate. Rates were estimated using species means of natural log-transformed values for all traits (such that

rate variation is represented in terms of relative change in proportion to the trait mean; this is essential for

comparison of traits measured using different units [64]). For these analyses, we co-estimated among-trait

covariances (similar results were obtained with covariances fixed at zero).

To test for an association between fundamental and realized thermal niche estimates, we

conducted a phylogenetically-controlled correlation test between CT_{max} and geo-referenced Worldclim

temperature data for each species based on collection localities. We used the first principal component

axis from the *Anolis* climate temperature dataset of Algar and Mahler (2015), in which a phylogenetic

principal component analysis (PCA) was used to condense the 11 Worldclim variables related to

temperature into PCA axes (Supplementary Table S2). The first PC axis loaded positively for all variables

related to temperature (e.g., mean annual temperature, mean maximum temperature; Supplementary Table

S2), and can therefore be considered a composite estimate of habitat temperature.

Results

All Puerto Rican sister species pairs have diverged in at least one physiological trait (T_{opt}, CT_{max}, or both),

and in the predicted direction in all cases (Fig. 2). T_{opt} differed between the *A. evermanni*/*A. stratulus* and

A. cristatellus/*A. cooki* species pairs (non-overlapping bootstrapped 95% CIs), but not between the *A.*

poncensis/*A. gundlachi* or *A. pulchellus*/*A. krugi* species pairs (Fig. 2A). We note that the T_{opt} of *A. cooki*

and *A. stratulus* were likely underestimated because most individuals had a T_{opt} of 35°C solely because

that was our maximum test temperature. CT_{max} differed among Puerto Rican species ($F_{7,72} = 42.74$; $P <$

0.001), with significant divergence in three of the four species pairs: *A. evermanni*/*A. stratulus*, *A.*

poncensis/A. gundlachi, and *A. pulchellus/A. krugi* (all $P < 0.001$), but not in the *A. cristatellus/A. cooki*

species pair ($P = 0.490$; Fig. 2B).

Physiological divergence influenced predicted performance in natural environments. In three of

four species pairs, the cool niche species had higher predicted performance than the warm niche species

in the shaded operative thermal environment, with mean performance advantages of 8-14% depending on

the species pair and time of day (Fig. 3). The lone exception was the *A. pulchellus/A. krugi* species pair,

for which both have similar performance estimates (Fig. 3G). Warm niche species tend to have a

performance advantage in the open operative environment, particularly during midday hours (from 10:00-

13:00; Fig. 3). The *A. cristatellus/A. cooki* species pair is the exception (Fig. 3D). However, *A. cooki*

midday performance is likely underestimated due to our probable underestimation of its T_{opt} (see above).

No species are under threat of overheating in the shaded operative environment (Fig. 4A, C, E,

G). However, all species have overheating risk in the open operative environment (Fig. 4B, D, F, H). For

three of the four sister-species pairs (*A. stratulus/A. evermanni*, *A. poncensis/A. gundlachi*, and *A.*

pulchellus/A. krugi), the cool niche species has a higher probability of overheating (Fig. 4B, F, H).

Difference in overheating probability are particularly acute during midday, when cool niche species have

overheating probabilities at least twice that of their warm niche counterparts (G-tests, all $P < 0.05$). The

A. cristatellus/A. cooki species pair, which had no divergence in CT_{max}, was the only exception (Fig. 4D).

Significant physiological divergence is also apparent among Jamaican species. Among species

mean T_{opt} ranges from 29.5 to 34.6°C (Fig. 5A) while CT_{max} ranges from 35.8-41.5°C (Fig. 5B; ANOVA,

$F_{3,36} = 13.6$, $P < 0.001$). Note that T_{opt} values of *A. grahami* and *A. opalinus* are likely underestimated due

to our upper experimental limit of 35°C (see above).

Across Puerto Rico and Jamaica, sympatric species that share the same ecomorphology invariably

differ in thermal physiology (Table 1). Thermal tolerance has evolved significantly more slowly than

some, but not all, ecomorphological traits (Table 2). CT_{max} evolved significantly more slowly than body

size (SVL; $P < 0.001$) and relative limb length (femur; $P = 0.029$), but did not differ from head length (P

$= 0.105$) or toepad width ($P = 0.112$).

CT_{max} was significantly positively correlated with geo-referenced climatic temperatures as

represented by temperature principal components axis 1 ($P = 0.006$, $r^2 = 0.45$; Fig. 6).

Discussion

Abiotic niche evolution has been invoked as an important component of many evolutionary radiations,

but the functional consequences of abiotic niche divergence are generally unknown. We demonstrate that

the Puerto Rican *Anolis* adaptive radiation is accompanied by physiological divergence in both heat

tolerance and sensitivity to sub-lethal temperature variability. Warm-niche species (determined based on

preference for open versus shaded perches) have either higher heat tolerance, higher optimal

physiological temperatures, or both, compared to cool-niche species. The anoles of Jamaica also

diversified physiologically, with species evolving essentially the full range of heat tolerances and optimal

temperatures observed in their Puerto Rican counterparts (Fig. 5).

Repeated divergence in physiological traits under similar conditions is itself evidence of

adaptation [65]. However, we provide more direct evidence of adaptation by estimating physiological

performance of species in warm and cool operative environments on Puerto Rico. Cool niche species have

higher predicted performance in shaded operative environments than their warm niche sister species in

three of the four species pairs (Fig. 3), with the lone exception a species pair for which there was no

divergence in T_{opt}. The pattern was less pronounced in the open operative environment, with warm niche

species having a sizable performance advantage in only two species pairs (Fig. 3). However, the

performance advantage for warm niche species becomes apparent when considering overheating risk.

During midday hours in the warmer open environment, warm niche species in three of the four species

pairs had less than half the overheating risk of cool niche species (Fig. 4). The lone exception was the *A.*

cristatellus/*A. cooki* species pair, which exhibited no divergence in thermal tolerance (Fig. 2B).

The high community-level species diversity associated with *Anolis* and other adaptive radiations

is facilitated by fine-scale resource partitioning [3]. Sympatric anoles are known to behaviorally partition

thermal microhabitats [12, 13], and our analysis indicates that this is facilitated by physiological

divergence. When co-occurring anole species share the same fundamental structural niche (*i.e.*,

ecomorphological microhabitat specialization), they invariably differ in thermal physiology (Table 1).

This pattern is not only maintained for endemic members of the community, but also for introduced

species: in Jamaica, the endemic trunk-ground species *A. lineatopus* is found in sympatry with the

introduced trunk-ground species *A. sagrei*, and these species differ significantly in thermal tolerance

(Table 1). These data highlight that, even in radiations for which morphological divergence is important

in facilitating coexistence, physiological divergence can also play an important role, providing an

orthogonal axis of diversification.

We found that thermal tolerance evolved more slowly than some ecomorphological traits (SVL

and femur length), and evolved at similar rates to others (head length and toe-pad width; Table 2). In

general, divergence along the fundamental thermal niche axis appears to have occurred more slowly than

divergence along the fundamental structural niche axis, at least within the Puerto Rican and Jamaican

anole radiations. This result is inconsistent with a recent analysis that found that thermal niche, estimated

as field body temperature, evolves more quickly than morphology in Caribbean anoles [42]. Discrepancy

between these results may be due to the different traits analyzed and/or the different species included, or

the fact that that study estimated rates of absolute, rather than proportional, trait change. Regardless, our

results suggest that high rates of evolutionary change are not necessary for traits to be ecologically

important, as the relatively slow divergence that we find in anole thermal tolerance has important

performance consequences with respect to overheating risk in warm habitats (Fig. 4).

We found that evolution of the fundamental and realized thermal niche are correlated, as CT_{max}

was significantly positively correlated with geo-referenced temperature data across species (Fig. 6).

Therefore, the realized thermal niches estimated in correlative ENM studies can capture some of the

underlying variability in the fundamental thermal niche. Nonetheless, the majority of variability in the

data (~55%) is left unexplained, and there are clear cases where realized and fundamental niche estimates

diverge. For example, *A. lineatopus* and *A. gundlachi* have very similar CT_{max} (mean CT_{max} = 36.3°C and

36.2°C, respectively), but occur in very different realized thermal niches (temperature PC1 = 0.14 and -

2.66, respectively). Conversely, *A. opalinus* and *A. pulchellus* occur in very similar realized niches

(temperature PC1 = -0.53 and -0.51, respectively) but have very different CT_{max} (mean CT_{max} = 37.5°C

and 41.3°C, respectively). These results underscore the fact that broad-scale climatic conditions are not

necessarily reflective of the underlying thermal biology of the organism being considered. Much of this

discrepancy is likely due to behavioral thermoregulation, which allows taxa in the same geographic

location to experience very different body temperatures and those in different geographic locations to

experience very similar body temperatures [33, 66].

We have shown that fundamental thermal niche divergence is important in the Puerto Rican and

Jamaican *Anolis* adaptive radiations. However, the importance of such divergence is likely not restricted

to these islands. For example, species of *cybotes* group trunk-ground anoles in Hispaniola have diverged

in thermal tolerance along an elevational gradient [44, 67], and variation in thermal physiology also

occurs among non-Antillean anoles [68, 69]. Water loss rates also differ among anole species and

populations [70-76]. In this context, our results suggest that physiological divergence may be as important

as morphological divergence for diversification and coexistence in this classic radiation.

Achieving a mechanistic understanding of how evolutionary and ecological processes interact to

promote the production and maintenance of biodiversity is a long-standing goal in biology. Given the

reality of ongoing climate change, the need to understand these processes has become immediate,

particularly with respect to temperature-dependent traits [77]. By focusing on physiological traits that link

temperature to performance, we demonstrate that the evolution of thermal physiology can facilitate

adaptive radiation by contributing to *in situ* performance tradeoffs and species co-existence. While our

trait- and performance-based analyses would not be possible using the type of broad-scale climatic data

used in ENMs studies alone, we find that climate-scale realized niche features correlate with fundamental

niche features. Nonetheless, climatic data provide imperfect signal with respect to the evolution of

functional traits, likely due to the ability of species to behaviorally augment their effective thermal

environments. We suggest that deeper insights about the contribution of climatic niche divergence to

evolutionary radiations will emerge when correlative ENMs are used in tandem with experimental

physiological studies within an integrative research program.

Acknowledgements

We thank Jonathan Losos for helpful comments on this manuscript, D. Steinberg for field help and the El

Verde and Mata de Plátano field stations, Ridge to Reef farm, and Green Castle Estates for logistical

support. Permits were provided by the Puerto Rican Departmento de Recursos Naturales y Ambientales,

the St. Croix Division of Fish and Wildlife, and the Jamaican National Environment and Planning

Agency. Funding was provided by NSF DDIG #1110570 to A.R.G.

References

1.   Hutchinson G.E. 1959 Homage to Santa Rosalia or why are there so many kinds of
animals? *The American Naturalist* **93**(870), 145-159.
2.   Simpson G.G. 1953 *Major features of evolution*. New York, Columbia University Press.
3.   Schluter D. 2000 *The ecology of adaptive radiation*, Oxford University Press.
4.   McPeek M.A. 2008 The ecological dynamics of clade diversification and community
assembly. *The American Naturalist* **172**(6), E270-E284.
5.   Grant P.R. 1999 *Ecology and evolution of Darwin's finches*, Princeton University Press.
6.   Streelman J.T., Danley P.D. 2003 The stages of vertebrate evolutionary radiation. *Trends*
in Ecology & Evolution **18**(3), 126-131.
7.   Meyer A. 1993 Phylogenetic relationships and evolutionary processes in East African
cichlid fishes. *Trends in Ecology & Evolution* **8**(8), 279-284.
8.   Nosil P. 2012 *Ecological speciation*, Oxford University Press.
9.   Chown S., Gaston K., Robinson D. 2004 Macrophysiology: large-scale patterns in
physiological traits and their ecological implications. *Functional Ecology* **18**(2), 159-167.
10.   Gaston K.J., Chown S.L., Calosi P., Bernardo J., Bilton D.T., Clarke A., Clusella-Trullas
S., Ghalambor C.K., Konarzewski M., Peck L.S. 2009 Macrophysiology: a conceptual
reunification. *The American Naturalist* **174**(5), 595-612.
11.   Kaspari M., Clay N.A., Lucas J., Yanoviak S.P., Kay A. 2015 Thermal adaptation
generates a diversity of thermal limits in a rainforest ant community. *Global Change Biology*
21(3), 1092-1102.

12.   Rand A.S. 1964 Ecological distribution in anoline lizards of Puerto Rico. *Ecology* **45**,
745-752.
13.   Ruibal R. 1961 Thermal relations of five species of tropical lizards. *Evolution* **15**(1), 98-
111.
14.   Wake D.B. 1987 Adaptive radiation of salamanders in Middle American cloud forests.
Annals of the Missouri Botanical Garden **74**, 242-264.
15.   Kozak K.H., Wiens J.J. 2007 Climatic zonation drives latitudinal variation in speciation
mechanisms. *Proceedings of the Royal Society of London B: Biological Sciences* **274**(1628),
2995-3003.
16.   Labra A., Pienaar J., Hansen T.F. 2009 Evolution of thermal physiology in *Liolaemus*
lizards: adaptation, phylogenetic inertia, and niche tracking. *The American Naturalist* **174**(2),
204-220.
17.   Bonino M.F., Azócar D.L.M., Tulli M.J., Abdala C.S., Perotti M.G., Cruz F.B. 2011
Running in cold weather: morphology, thermal biology, and performance in the southernmost
lizard clade in the world (*Liolaemus lineomaculatus* section: Liolaemini: Iguania). *Journal of*
Experimental Zoology Part A: Ecological Genetics and Physiology **315**(8), 495-503.
18.   Pincheira-Donoso D. 2011 Predictable variation of range-sizes across an extreme
environmental gradient in a lizard adaptive radiation: evolutionary and ecological inferences.
PloS One **6**(12), e28942.
19.   Kellermann V., van Heerwaarden B., Sgrò C.M., Hoffmann A.A. 2009 Fundamental
evolutionary limits in ecological traits drive *Drosophila* species distributions. *Science* **325**(5945),
1244-1246.
20.   Stillman J.H., Somero G.N. 2000 A comparative analysis of the upper thermal tolerance
limits of eastern Pacific porcelain crabs, genus *Petrolisthes*: influences of latitude, vertical
zonation, acclimation, and phylogeny. *Physiological and Biochemical Zoology* **73**(2), 200-208.
21.   Dalziel A.C., Martin N., Laporte M., Guderley H., Bernatchez L. 2015 Adaptation and
acclimation of aerobic exercise physiology in Lake Whitefish ecotypes (*Coregonus*
clupeaformis). *Evolution* **69**(8), 2167-2186.
22.   Knouft J.H., Losos J.B., Glor R.E., Kolbe J.J. 2006 Phylogenetic analysis of the evolution
of the niche in lizards of the *Anolis sagrei* group. *Ecology* **87**(sp7), S29-S38.
23.   Quintero I., Wiens J.J. 2013 Rates of projected climate change dramatically exceed past
rates of climatic niche evolution among vertebrate species. *Ecology Letters* **16**(8), 1095-1103.
24.   Warren D.L., Glor R.E., Turelli M. 2008 Environmental niche equivalency versus
conservatism: quantitative approaches to niche evolution. *Evolution* **62**(11), 2868-2883.
25.   Lawing A.M., Polly P.D., Hews D.K., Martins E.P., Collar D.C., Bronstein J.L. 2016
Including Fossils in Phylogenetic Climate Reconstructions: A Deep Time Perspective on the
Climatic Niche Evolution and Diversification of Spiny Lizards (*Sceloporus*). *The American*
Naturalist **188**(2), 000-000.
26.   Peterson A.T., Soberón J., Pearson R.G., Anderson R.P., Nakamura M., Martinez-Meyer
E., Araújo M.B. 2011 *Ecological niches and geographic distributions*. Princeton, New Jersey,
USA, Princeton University Press.
27.   Guisan A., Thuiller W. 2005 Predicting species distribution: offering more than simple
habitat models. *Ecology Letters* **8**(9), 993-1009.
28.   Buckley L.B., Urban M.C., Angilletta M.J., Crozier L.G., Rissler L.J., Sears M.W. 2010
Can mechanism inform species' distribution models? *Ecology Letters* **13**(8), 1041-1054.

29.   Alvarado-Serrano D.F., Knowles L.L. 2014 Ecological niche models in phylogeographic
studies: applications, advances and precautions. *Molecular Ecology Resources* **14**(2), 233-248.
30.   Kearney M., Porter W. 2009 Mechanistic niche modelling: combining physiological and
spatial data to predict species' ranges. *Ecology letters* **12**(4), 334-350.
31.   Warren D.L., Cardillo M., Rosauer D.F., Bolnick D.I. 2014 Mistaking geography for
biology: inferring processes from species distributions. *Trends in Ecology & Evolution* **29**(10),
572-580.
32.   Araújo M.B., Peterson A.T. 2012 Uses and misuses of bioclimatic envelope modeling.
Ecology **93**(7), 1527-1539.
33.   Bogert C.M. 1949 Thermoregulation in reptiles, a factor in evolution. *Evolution* **3**(3),
195-211.
34.   Losos J.B., Jackman T.R., Larson A., de Queiroz K., Rodríguez-Schettino L. 1998
Contingency and determinism in replicated adaptive radiations of island lizards. *Science*
279(5359), 2115-2118.
35.   Williams E.E. 1983 Ecomorphs, faunas, island size, and diverse end points in island
radiations of Anolis. In *Lizard ecology: studies of a model organism* (pp. 326-370. Cambridge,
Harvard University Press.
36.   Mahler D.L., Ingram T., Revell L.J., Losos J.B. 2013 Exceptional convergence on the
macroevolutionary landscape in island lizard radiations. *Science* **341**(6143), 292-295.
37.   Losos J.B. 2009 *Lizards in an evolutionary tree: ecology and adaptive radiation of*
anoles, University of California Press.
38.   Williams E.E. 1972 The origin of faunas. Evolution of lizard congeners in a complex
island fauna: a trial analysis. *Evolutionary Biology* **6**, 47-89.
39.   Heatwole H., Lin T.-H., Villalón E., Muñiz A., Matta A. 1969 Some aspects of the
thermal ecology of Puerto Rican anoline lizards. *Journal of Herpetology* **3**, 65-77.
40.   Huey R.B., Webster T.P. 1976 Thermal biology of *Anolis* lizards in a complex fauna: the
cristatellus group on Puerto Rico. *Ecology* **57**, 985-994.
41.   Losos J.B., Leal M., Glor R.E., de Queiroz K., Hertz P.E., Schettino L.R., Lara A.C.,
Jackman T.R., Larson A. 2003 Niche lability in the evolution of a Caribbean lizard community.
Nature **424**(6948), 542-545.
42.   Hertz P.E., Arima Y., Harrison A., Huey R.B., Losos J.B., Glor R.E. 2013 Asynchronous
evolution of physiology and morphology in *Anolis* lizards. *Evolution* **67**(7), 2101-2113.
43.   Algar A.C., Mahler D.L. 2015 Area, climate heterogeneity, and the response of climate
niches to ecological opportunity in island radiations of *Anolis* lizards. *Global Ecology and*
Biogeography **doi:10.1111 /geb.12327**.
44.   Muñoz M.M., Stimola M.A., Algar A.C., Conover A., Rodriguez A.J., Landestoy M.A.,
Bakken G.S., Losos J.B. 2014 Evolutionary stasis and lability in thermal physiology in a group
of tropical lizards. *Proceedings of the Royal Society B: Biological Sciences* **281**(1778),
20132433.
45.   Nicholson K.E., Glor R.E., Kolbe J.J., Larson A., Blair Hedges S., Losos J.B. 2005
Mainland colonization by island lizards. *Journal of Biogeography* **32**(6), 929-938.
46.   Mahler D.L., Revell L.J., Glor R.E., Losos J.B. 2010 Ecological opportunity and the rate
of morphological evolution in the diversification of Greater Antillean anoles. *Evolution* **64**(9),
2731-2745.
47.   Gamble T., Geneva A.J., Glor R.E., Zarkower D. 2014 *Anolis* sex chromosomes are
derived from a single ancestral pair. *Evolution* **68**(4), 1027-1041.

48. Hertz P.E. 1992 Temperature regulation in Puerto Rican *Anolis* lizards: a field test using
null hypotheses. *Ecology* **73**, 1405-1417.
49. Hertz P.E. 1992 Evaluating thermal resource partitioning by sympatric lizards *Anolis*
cooki and *A*. *cristatellus*: a field test using null hypotheses. *Oecologia* **90**(1), 127-136.
50. Jackman T.R., Larson A., De Queiroz K., Losos J.B. 1999 Phylogenetic relationships and
tempo of early diversification in *Anolis* lizards. *Systematic Biology* **48**(2), 254-285.
51. Gunderson A.R., Leal M. 2012 Geographic variation in vulnerability to climate warming
in a tropical Caribbean lizard. *Functional Ecology* **26**(4), 783-793.
52. Leal M., Gunderson A.R. 2012 Rapid change in the thermal tolerance of a tropical lizard.
The American Naturalist **180**(6), 815-822.
53. Huey R.B. 1983 Natural variation in body temperature and physiological performance in
a lizard (*Anolis cristatellus*). In *Advances in herpetology and evolutionary biology: essays in*
honor of Ernest E Williams (pp. 484-490.
54. Losos J.B. 1990 Ecomorphology, performance capability, and scaling of West Indian
Anolis lizards: an evolutionary analysis. *Ecological Monographs* **60**, 369-388.
55. Logan M.L., Cox R.M., Calsbeek R. 2014 Natural selection on thermal performance in a
novel thermal environment. *Proceedings of the National Academy of Sciences* **111**(39), 14165-
14169.
56. Angilletta M.J. 2009 *Thermal adaptation: a theoretical and empirical synthesis*, Oxford
University Press.
57. Deutsch C.A., Tewksbury J.J., Huey R.B., Sheldon K.S., Ghalambor C.K., Haak D.C.,
Martin P.R. 2008 Impacts of climate warming on terrestrial ectotherms across latitude.
Proceedings of the National Academy of Sciences **105**(18), 6668-6672.
58. Huey R.B., Deutsch C.A., Tewksbury J.J., Vitt L.J., Hertz P.E., Pérez H.J.Á., Garland T.
2009 Why tropical forest lizards are vulnerable to climate warming. *Proceedings of the Royal*
Society B: Biological Sciences **276**(1664), 1939-1948.
59. Efron B. 1987 Better bootstrap confidence intervals. *Journal of the American Statistical*
Association **82**(397), 171-185.
60. Ewel J.J., Whitmore J.L. 1973 Ecological life zones of Puerto Rico and US Virgin
Islands. In *Forest Service Research Paper ITF-18* (U.S. Department of Agriculture.
61. Hertz P.E., Huey R.B., Stevenson R. 1993 Evaluating temperature regulation by field-
active ectotherms: the fallacy of the inappropriate question. *The American Naturalist* **142**, 796-
818.
62. Gorman G.C., Licht P. 1974 Seasonality in ovarian cycles among tropical *Anolis* lizards.
Ecology **55**, 360-369.
63. Revell L.J. 2012 phytools: an R package for phylogenetic comparative biology (and other
things). *Methods in Ecology and Evolution* **3**(2), 217-223.
64. Adams D.C. 2013 Quantifying and comparing phylogenetic evolutionary rates for shape
and other high-dimensional phenotypic data. *Systematic Biology* **62**, 181-192.
65. Harvey P.H., Pagel M.D. 1991 *The comparative method in evolutionary biology*, Oxford
University Press.
66. Huey R.B., Hertz P.E., Sinervo B. 2003 Behavioral drive versus behavioral inertia in
evolution: a null model approach. *The American Naturalist* **161**(3), 357-366.
67. Hertz P.E., Huey R.B. 1981 Compensation for altitudinal changes in the thermal
environment by some *Anolis* lizards on Hispaniola. *Ecology* **62**(3), 515-521.

68.   Logan M.L., Huynh R.K., Precious R.A., Calsbeek R.G. 2013 The impact of climate
change measured at relevant spatial scales: new hope for tropical lizards. *Global Change Biology*
19, 3093-3102.
69.   van Berkum F.H. 1986 Evolutionary patterns of the thermal sensitivity of sprint speed in
Anolis lizards. *Evolution* **40**, 594-604.
70.   Gunderson A.R., Siegel J., Leal M. 2011 Tests of the contribution of acclimation to
geographic variation in water loss rates of the West Indian lizard *Anolis cristatellus*. *Journal of*
Comparative Physiology B **181**(7), 965-972.
71.   Hillman S., Gorman G. 1977 Water loss, desiccation tolerance, and survival under
desiccating conditions in 11 species of Caribbean *Anolis*. *Oecologia* **29**(2), 105-116.
72.   Sexton O.J., Heatwole H. 1968 An experimental investigation of habitat selection and
water loss in some anoline lizards. *Ecology* **49**, 762-767.
73.   Hertz P., Arce-Hernandez A., Ramirez-Vazquez J., Tirado-Rivera W., Vazquez-Vives L.
1979 Geographical variation of heat sensitivity and water loss rates in the tropical lizard, *Anolis*
gundlachi. *Comparative Biochemistry and Physiology Part A: Physiology* **62**(4), 947-953.
74.   Hertz P.E. 1980 Responses to dehydration in *Anolis* lizards sampled along altitudinal
transects. *Copeia* **1980**, 440-446.
75.   Hillman S., Gorman G.C., Thomas R. 1979 Water loss in *Anolis* lizards: evidence for
acclimation and intraspecific differences along a habitat gradient. *Comparative Biochemistry and*
Physiology Part A: Physiology **62**(2), 491-493.
76.   Dmi'el R., Perry G., Lazell J. 1997 Evaporative water loss in nine insular populations of
the lizard *Anolis cristatellus* group in the British Virgin Islands. *Biotropica* **29**(1), 111-116.
77.   Huey R.B., Kearney M.R., Krockenberger A., Holtum J.A., Jess M., Williams S.E. 2012
Predicting organismal vulnerability to climate warming: roles of behaviour, physiology and
adaptation. *Philosophical Transactions of the Royal Society B: Biological Sciences* **367**(1596),
1665-1679.

Table 1. Physiology differences between co-occurring species that occupy the same or similar structural

niches (*i.e.*, share the same or similar ecomorphology). For CT_{max}, P-values are given based on one-way

ANOVA with planned orthogonal contrasts. For T_{opt}, "*" indicates that the bootstrapped 95% confidence

intervals did not overlap, "N.S." indicates no significant difference. "†" indicates that the species is an

introduced member of the community.

Table 2. Estimated rates of evolution for heat tolerance (CT_{max}) and morphological traits related to

ecomorphological divergence. Rate estimates (bold) are in the diagonal of the matrix. Upper off-diagonals

contain estimates of pairwise evolutionary covariances among traits, lower off-diagonals contain P-values

for likelihood ratio test comparisons of 2-rate versus 1-rate models for each trait pair. P-values less than

0.05 indicate support for a 2-rate model over a 1-rate model.

527
528

532   Table 1

| co-occurring species | | ecomorph | physiological trait | |
			CT_{max}	T_{opt}
A. stratulus	*A. evermani*	trunk-crown	<0.001	*
A. cristatellus	*A. cooki*	trunk-ground	0.490	*
A. cristatellus	*A. gundlachi*	trunk-ground	<0.001	n.s.
A. krugi	*A. pulchellus*	grass-bush	<0.001	n.s.
A. lineatopis	*A. grahami*	trunk-ground/Trunk-crown	<0.001	n.s.
A. lineatopis	*A. sagrei*[†]	trunk-ground	<0.001	n.s.

541 Table 2.

	CT_{max}	SVL	head length	femur length	toepad width
CT_{max}	**5.80E-05**	-4.90E-05	-1.60E-05	-5.50E-05	-9.90E-06
SVL	<0.001	**2.30E-03**	8.90E-06	-1.80E-05	2.70E-05
head length	0.105	<0.001	**2.70E-05**	1.30E-05	-6.30E-06
femur length	0.029	<0.001	0.001	**1.40E-04**	2.20E-05
toepad width	0.112	<0.001	0.003	0.832	**1.30E-04**

Figure 1: Phylogenetic relationships of the species included in this study, along with the islands from

which they were sampled.

Figure 2: Thermal physiology of four sister-species pairs of Puerto Rican anole. Each species pair is

represented in alternating grey and white regions of the plot with warm niche species in red and cool

niche species in blue. (A) Optimal sprint performance temperatures (T_{opt}). (B) Heat tolerance limits

(CT_{max}). "*", significant difference between members of a species pair; N.S., no significant difference

between members of a species pair. See Methods for analysis details.

Figure 3: Predicted relative physiological performance of sister-species pairs of Puerto Rican *Anolis*

under shaded and open habitat operative thermal environments found on Puerto Rico during morning,

midday, and afternoon/evening hours. Warm niche species in red, cool niche species in blue. (A, B) *A.*

stratulus/A. evermanni, (C, D) *A. cooki/A. cristatellus*, (E, F) *A. poncensis/A. gundlachi* (G, H) *A.*

pulchellus/A. krugi.

Figure 4: Predicted overheating probability of sister-species pairs of Puerto Rican *Anolis* under shaded

and open habitat operative thermal environments found on Puerto Rico during morning, midday, and

afternoon/evening hours. Warm niche species in red, cool niche species in blue. (A, B) *A. stratulus/A.*

evermanni, (C, D) *A. cooki/A. cristatellus* (E, F), *A. poncensis/A. gundlachi*, (G, H) *A. pulchellus/A. krugi*.

Figure 5: Thermal physiology of endemic Jamaican anoles. (A) Optimal sprinting temperature (T_{opt}).

Anolis valencienni does not have a T_{opt} estimate because this species could not be induced to run. (B) Heat

tolerance (CT_{max}).

Figure 6: Relationship between CT_{max} and climate temperature PC1 from Algar and Mahler (2015).

603   Figure 1

610   Figure 2.

relative performance capacity

time of day

Figure 3

Figure 4

673   Figure 5

699 Figure 6.

702