1 **290 Metagenome-assembled Genomes from the Mediterranean Sea: Ongoing Effort to**
2 **Generate Genomes from the *Tara Oceans* Dataset**
3

4 **Authors**
5 Benjamin J. Tully[1‡], Rohan Sachdeva[2], Elaina D. Graham[2], and John F. Heidelberg[1,2]
6
‡ - corresponding author, tully.bj@gmail.com
1 – Center for Dark Energy Biosphere Investigations, University of Southern California, Los
Angeles, CA 90089
2 – Department of Biological Sciences, University of Southern California, Los Angeles, CA
90089

Abstract
The *Tara Oceans* Expedition has provided large, publicly-accessible microbial metagenomic
datasets from a circumnavigation of the globe. Utilizing several size fractions from the samples
originating in the Mediterranean Sea, we have used current assembly and binning techniques to
reconstruct 290 putative high-quality metagenome-assembled bacterial and archaeal genomes,
with an estimated completion of ≥50%, and an additional 2,786 bins, with estimated completion
of 0-50%. We have submitted our results, including initial taxonomic and phylogenetic
assignments for the putative high-quality genomes, to open-access repositories (iMicrobe and
FigShare) for the scientific community to use in ongoing research.

Introduction
Microorganisms are a major constituent of the biology within the world's oceans and act as the
important linchpins in all major global biogeochemical cycles[1]. Marine microbiology is among
the disciplines at the forefront of pushing advancements in understanding how microorganisms
respond to and impact the local and large-scale environments. An estimated 10^{29} Bacteria and
Archaea[2] reside in the oceans and an immense amount of poorly constrained, and ever evolving
genetic diversity.
The *Tara Oceans* Expedition (2003-2010) encompassed a major endeavor to add to the
body of knowledge collected during previous global ocean surveys to sample the genetic
potential of microorganisms [3]. To accomplish this goal, members of *Tara Oceans* sampled
planktonic organisms (viruses to fish larvae) at two major depths, the surface ocean and the
mesopelagic. The amount of data collected was expansive and included 35,000 samples from
210 ecosystems[3]. The *Tara Oceans* Expedition generated and publically released 7.2 Tbp of
metagenomic data from 243 ocean samples from throughout the global ocean, specifically
targeting the smallest members of the ocean biosphere, the viruses, Bacteria and Archaea, and
picoeukaryotes[4]. Initial work on these fractions produced a large protein database, totaling > 40
million nonredundant protein sequences and identified >35,000 microbial operational taxonomic
units (OTUs)[4].
Leveraging the publically available metagenomic sequences from the "girus" (giant virus;
0.22-1.6 μm), "bacteria" (0.22-1.6 μm), and "protist" (0.8-5 μm) size fractions, we have
performed a new joint assembly of these samples using current sequence assemblers (Megahit[5])
and methods (combining assemblies from multiple sites using Minimus2[6]). These metagenomic
assemblies were binned using a strictly coverage based binning algorithm[7] in to 290 high-quality
(low contamination) microbial genomes, ranging from 50-100% estimated completion.

Environmentally derived genomes representing the most abundant microorganisms are
imperative for a number of downstream applications, including comparative genomes,
metatranscriptomics, and metaproteomics. This series of genomic data can allow for the
recruitment of environmental "-omic" data and provide linkages between functions and
phylogenies. This method was initially performed on the seven sites from the Mediterranean Sea
containing microbial metagenomic samples (TARA007, -009, -018, -023, -025 and -030), but
will continue through the various Longhurst provinces[8,9] sampled during the *Tara Oceans*
project (Figure 1). All of the assembly data is publically available, including the initial Megahit
assemblies for each site from the various size fractions and depths and putative (minimal quality
control) genomes within iMicrobe (http://imicrobe.us).

57
Materials and Methods

59
60 A generalized version of the following workflow is presented in Figure 2.

61
Sequence Retrieval and Assembly
All sequences for the reverse and forward reads from each sampled site and depth within the
Mediterranean Sea were accessed from European Molecular Biology Laboratory (EMBL)
utilizing their FTP service (Table 1). Paired-end reads from different filter sizes from each site
and depth (e.g., TARA0007, girus filter fraction, sampled at the deep chlorophyll maximum)
were assembled using Megahit[5] (v1.0.3; parameters: --preset, meta-sensitive). To keep consistent
with TARA sample nomenclature, "bacteria" or "BACT" will be used to encompass the size
fraction 0.22-1.6 μm. All of the Megahit assemblies were pooled in to two tranches based on
assembly size, ≤1,999bp, and ≥2,000bp. Longer assemblies (≥2kb) with ≥99% semi-global
identity were combined using CD-HIT-EST (v4.6; -T 90 -M 500000 -c 0.99 -n 10). The reduced
set of contiguous DNA fragments (contigs) was then cross-assembled using Minimus2[6] (AMOS
v3.1.0; parameters: -D OVERLAP=100 MINID=95).

74
Metagenome-assembled Genomes
Sequence reads were recruited against a subset of contigs (≥7.5kb) constructed during the
secondary assembly (Megahit + Minimus2) for each of the *Tara* samples using Bowtie2[10]
(v4.1.2; default parameters). Utilizing the SAM file output, read counts for each contig were
determined using featureCounts[11] (v1.5.0; default parameters). Coverage was determined for all
contigs by dividing the number of recruited reads by the length of the contig (reads/bp). Due to
the low coverage nature of the samples, in order to effectively delineate between contig coverage
patterns, the coverage values were transformed by multiplying by five (determined through
manual tuning). Transformed coverage values were then utilized to cluster contigs in to bins
utilizing BinSanity (parameters: -p -3, -m 4000, -v 400, -d 0.9) [7]. Bins were assessed for the
presence of putative microbial genomes using CheckM[12] (v1.0.3; parameters: lineage_wf). Bins
were split in to three categories: (1) putative high quality genomes (≥50% complete and ≤10%
cumulative redundancy [% contamination – (% redundancy × % strain heterogeneity ÷ 100)]);
(2) bins with "high" contamination (≥50% complete and ≥10% cumulative redundancy); and (3)
low completion bins (<50% complete). The high contamination group were additionally binned
using the BinSanity refinement method (refine-contaminated-log.py; parameters: -p 'variable', -
m 2000, -v 200, -d 0.9), which utilizes affinity propagation[13] to cluster contigs within a bin based
on tetranucleotide frequencies and %G+C.

To determine the preference values needed to successfully bin the high contamination
bins, 15 bins were assessed manually using the number of marker occurrences determined by
CheckM. Bins containing approximately two genomes, three genomes, and bins with more
genomes used a preference of -1000 (-p -1000), -500 (-p -500), and -100 (-p -100), respectively.
The 15 manually assessed bins were used to train a decision tree within scikit-learn[14] (default
parameters, DecisionTreeClassifier) to assign parameters to the other bins. The resulting bins
were added to one of the three categories: putative high quality genomes, high contamination
bins, and low completion bins. The high contamination bins were processed for a third time with
the BinSanity refinement step utilizing a preference of -100 (-p -100). These bins were given
final assignments to either the putative high quality genomes (some putative genomes had >10%
cumulative contamination, but have been designated) or low completion bins. Bins determined to
be low completion bins were reserved for an additional round of binning (see below).
After this initial round of binning, all contigs not assigned to putative high-quality
genomes were assessed using BinSanity using raw coverage values. Two additional rounds of
refinement were performed (as above) with the first round of refinement using the decision tree
to determine preference and the second round using a set preference of -10 (-p -10). Following
this binning phase, contigs were assigned to high quality bins (*e.g.*, *Tara* **Med**iterranean genome
1, referred to as TMED1, etc.), low completion bins with at least five contigs (0-50% complete;
TMEDlc1, etc. lc, **l**ow **c**ompletion), or were not placed in a bin (Supplemental Table 1 & 2).
112
Taxonomic and Phylogenetic Assignment of High Quality Genomes
The bins representing the high quality genomes were assessed for taxonomy and phylogeny
using multiple methods to provide a quick reference for selecting genomes of interest. Taxonomy
as assigned using the putative placement provided via CheckM during the pplacer[15] step of the
analysis to the lowest taxonomic placement (parameters: tree_qa -o 2). This step was also
performed for all low completion bins. A second taxonomic assignment was determined using a
method modified from Albersten, *et al*. (2013)[16], wherein putative coding DNA sequences
(CDSs) were determined using Prodigal[17] (v2.6.3; parameters: -m -o -p meta -q). The putative
CDS were searched against the NCBI non-redundant (NR) database (accessed March 2016)
using DIAMOND[18] (v0.8.11.73; parameters: -f xml -k 5 --sensitive -e 1e-10) and the output was
processed using MEGAN[19] (v4; parameters: recompute toppercent = 5, recompute minsupport =
1, collapse rank = species, select nodes = all) to determine the last common ancestor for the top
five matches. Using a script from the Multi-Metagenome package (hmm.majority.vote.pl;
https://github.com/MadsAlbertsen/multimetagenome; parameters: -n -l 4 [-l 5, -l 6, or -l 7]), each
contig was assigned a consensus taxonomic identification at approximately the Phylum, Class,
Order, and Family levels. A consensus for all contigs at each taxonomic level was determined. If
at any level a tie was achieved between possible assignments, it has been denoted with a "T" in
the genome table.
Two separate attempts were made to assign the high quality genomes a phylogenetic
assignment. High quality genomes were searched for the presence of the full-length 16S rRNA
gene sequence using RNAmmer[20] (v1.2; parameters: -S bac -m ssu). All full-length sequences
were aligned to the SILVA SSU reference database (Ref123) using the SINA web portal
aligner[21] (https://www.arb-silva.de/aligner/). These alignments were loaded in to ARB[22] (v6.0.3),
manually assessed, and added to the non-redundant 16S rRNA gene database (SSURef123
NR99) using ARB Parsimony (Quick) tool (parameters: default). A selection of the nearest
neighbors to the *Tara* genome sequences were selected and used to construct a 16S rRNA

phylogenetic tree. Genome-identified 16S rRNA sequences and SILVA reference sequences
were aligned using MUSCLE[23] (v3.8.31; parameters: -maxiters 8) and processed by the
automated trimming program trimAL[24] (v1.2rev59; parameters: -automated1). Automated
trimming results were assessed manually in Geneious[25] (v6.1.8) and trimmed where necessary
(positions with >50% gaps) and re-aligned with MUSCLE (parameters: -maxiters 8). An
approximate maximum likelihood (ML) tree with pseudo-bootstrapping was constructed using
FastTree[26] (v2.1.3; parameters: -nt -gtr -gamma; Figure 3).
High-quality genomes were assessed for the presence of the 16 ribosomal markers genes
used in Hug, *et al*. (2016)[27]. Putative CDSs were determined using Prodigal (v2.6.3; parameters:
-m -p meta) and were searched using HMMs for each marker using HMMER[28] (v3.1b2;
parameters: hmmsearch --cut_tc --notextw). If a genome had multiple copies of any single
marker gene, neither was considered, and only genomes with ≥8 markers were used to construct
a phylogenetic tree. Markers identified from the high quality genomes were combined with
markers from 1,729 reference genomes that represent the major bacterial phylogenetic groups (as
presented by IMG[29]). Archaeal reference sequences were not included; however, none of the
putative archaeal environmental genomes had a sufficient number of markers for inclusion on the
tree. Each marker gene was aligned using MUSCLE (parameters: -maxiters 8) and automatically
trimmed using trimAL (parameters: -automated1). Automated trimming results were assessed (as
above) and re-aligned with MUSCLE, as necessary. Final alignments were concatenated and
used to construct an approximate ML tree with pseudo-bootstrapping with FastTree (parameters:
-gtr -gamma; Figure 4).
160
Relative Abundance of High Quality Genomes
To set-up a baseline that could approximate the "microbial" community (Bacteria, Archaea and
viruses) present in the various *Tara* metagenomes, which included filter sizes specifically
targeting both protists and giruses, reads were recruited against all contigs generated from the
Minimus2 and Megahit assemblies ≥2kb using Bowtie2 (default parameters). Some assumptions
were made that contigs <2kb would include, low abundance bacteria and archaea, bacteria and
archaea with high degrees of repeats/assembly poor regions, fragmented picoeukaryotic
genomes, and problematic read sequences (low quality, sequencing artefacts, etc.). All relative
abundance measures are relative to the number of reads recruited to the assemblies ≥2kb. Read
counts were determined using featureCounts (as above). Length-normalized relative abundance
values were determined for each high quality genome for each sample:

172
$$\frac{\frac{Reads}{bp}\,per\ genome}{\sum \frac{Reads}{bp}\,all\ genomes} \times \frac{\sum Recruited\ reads\ to\ genomes}{\sum Recruited\ reads\ to\ all\ contigs\ (\geq 2kb)} \times 100$$

173
Available Through iMicrobe
In keeping with the open-access nature of the *Tara Oceans* project, all of the data generated for
this analysis is publically available through iMicrobe (http://data.imicrobe.us/project/view/261),
including: all contigs generated using Megahit from each sample; all contigs from Minimus2 +
Megahit output used for binning and community assessment, ≥2kb and ≥7.5kb; a table that
details statistics, taxonomy, and phylogeny for the high quality genomes; the putative genome
contigs and Prodigal-predicted nucleotide and protein putative CDS FASTA files. Additional
files, such as, the ribosomal marker HMM profiles, reference genome markers, high quality
genome markers, final concatenated MUSCLE alignment, FastTree Newick file, contig read

count data, relative abundance matrix for genomes from all samples, low completion bins, and
contigs without a bin, as well as, additional data files, have been provided and are available
through FigShare (https://dx.doi.org/10.6084/m9.figshare.3545330). Digital locations of data
files and contents can be found on Supplemental Table 3.
187
Results
Assembly
The initial Megahit assembly was performed on the publicly available reads for *Tara* stations
007, 009, 018, 023, 025, 030. Starting with 147-744 million reads per sample, the Megahit
assembly process generated 1.2-4.6 million assemblies with a mean N_{50} and longest contig of
785bp and 537kb, respectively (Table 1). In general, the assmeblies generated from the *Tara*
samples targeting the protist size fraction (0.8-5 μm) had a shorter N_{50} value than the bacteria
size fractions (mean: 554bp vs 892bp, respectively). Assemblies from the Megahit assembly
process were pooled and separated by length. Of the 42.6 million assemblies generated during
the first assembly, 1.5 million were ≥2kb in length (Table 2). Several attempts were made to
assemble the shorter contigs, but publicly available overlap-consensus assemblers (Newbler [454
Life Sciences], cap3[30], and MIRA[31]) failed on multiple attempts. Processing the ≥2kb assemblies
from all of the samples through CD-HIT-EST reduced the total to 1.1 million contigs ≥2kb. This
group of contigs was subjected to the secondary assembly through Minimus2, generating
158,414 new contigs (all ≥2kb). The secondary contigs were combined with the Megahit contigs
that were not assembled by Minimus2. This provided a contig dataset consisting of 660,937
contigs, all ≥2kb in length (Table 2; further referred to as data-rich-contigs).
205
Binning
The set of data-rich-contigs was used to recruit the metagenomic reads from each sample using
Bowtie2. The data-rich-contigs recruited 15-81% of the reads depending on the sample. In
general, the protist size fraction recruited substantially fewer reads than the girus and bacteria
size fractions (mean: 19.8% vs 75.0%, respectively) (Table 1). For the protist size fraction, the
"missing" data for these recruitments likely results from the poor assembly of more complex and
larger eukaryotic genomes. The fraction of the reads that do not recruit in the girus and bacterial
size fraction samples could be accounted for by the large number of low quality assemblies (200-
500bp) and reads that could not be assembled due to low abundance or high complexity (Table
2). Coverage was determined as total reads per base pair, based on the number of reads recruited
to each contig.
Unsupervised binning was performed using both transformed and raw coverage values
for a subset of 95,506 contigs from the data-rich-contigs that were ≥7.5kb (referred to further as
binned-contigs) utilizing the tool BinSanity. An iterative process was performed that first used
coverage to generate putative bins, and then after removing putative high-quality genomes
(≥50% complete and <10% redundancy), based on estimates of redundancy through CheckM,
used two passes through the BinSanity refinement process, utilizing sequence composition.
Binning using the transformed coverage data generated 237 putative high-quality genomes (12
putative genomes are of slightly lower quality with >10% redundancy and have been noted)
containing 15,032 contigs. Contigs not in putative genomes were re-binned through the iterative
use of BinSanity based on raw coverage values, generating 53 additional putative high-quality
genomes encompassing 3,348 contigs. In total, 290 putative high-quality genomes were
generated with 50-100% completion (mean: 69%) with a mean length and number of putative

CDS of 1.7Mbp and 1,699, respectively (iMicrobe; Supplemental Table 1). All other contigs
were grouped in to bins with at least five contigs, but with estimated completion of 0-50% (2,786
low completion bins; 74,358 contigs; Supplemental Table 2) or did not bin (2,732 contigs).
Nearly a quarter of the low completion bins (24.7%) have an estimated completion of 0%.
233
Taxonomy, Phylogeny, & Potential Organisms of Interest
The 290 putative high-quality genomes had a taxonomy assigned to it via CheckM during the
pplacer step. All of the genomes, except for 20, had an assignment to at least the Phylum level,
and 83% of the genomes had an assignment to at least the Class level. Additionally, all of the
genomes were assigned putative taxonomies using a consensus method of the taxonomies
assigned to the putative CDS on a contig. The genomes were assigned four levels of taxonomic
information, roughly equivalent to Phylum, Order, Class, and Family. Due to the nature of this
method, especially at lower taxonomic levels, it is possible for a small number of assignments to
greatly influence the results. Because all of these methods have inherent biases, consistency
across several results should be viewed as reinforcing support for the accuracy of the genome,
while inconsistent results should not be used as evidence of an incorrectly binned genome.
Attempts were made to provide phylogenetic information for as many genomes as
possible. Genomes were assessed for the presence of full-length 16S rRNA genes. In total, 37
16S rRNA genes were detected in 35 genomes (mean 16S rRNA gene copy number, 1.05). 16S
rRNA genes can prove to be problematic during the assembly steps due the high level of
conservation that can break contigs[32] (Figure 3). Additionally, the conserved regions of the 16S
rRNA, depending on the situation, can over- or under-recruit reads, resulting in coverage
variations that can misplace contigs in to the incorrect genome. As such, several of the 16S
rRNA phylogenetic placements support the taxonomic assignments, while some are
contradictory. Further analysis should allow for the determination of the most parsimonious
result.
Beyond the 16S rRNA gene, genomes were searched for 16 conserved, syntenic
ribosomal markers. Sufficient markers (≥8) were identified in 193 of the genomes (67%) and
placed on a tree with 1,729 reference sequences (Figure 4). Phylogenies were then assigned to
the lowest taxonomic level that could be confidently determined.
The taxonomic and phylogenetic assignments are provided to give downstream users a
guide for determining which genomes prove to be most interesting for further analysis. Highest
confidence should be given to genomes with multiple lines of evidence supporting an assignment
and additional confirmation should be gathered for those with multiple conflicting results. These
putative results reveal a number of genomes were generated that represent multiple clades for
which environmental genomic information remains limited, including: *Planctomycetes*,
Verrucomicrobia, *Marinimicrobia*, *Cyanobacteria*, and uncultured groups within the *Alpha*- and
Gammaproteobacteria.
267
Relative Abundance
Based on the assembly and recruitment results, the assumption was made that the data-rich-
contigs and their corresponding reads represent the dominant portion of the microbial (bacterial,
archaeal, and viral) community and that reads that did not recruit represent eukaryotes, low
quality assemblies, and/or less dominant portions of the microbial community. A length-
normalized relative abundance value was determined for each genome in each sample based on

the number of reads recruited to the data-rich-contigs. The relative abundance for the individual
genomes was determined based on this portion of the read dataset.
In general, the genomes and their underlying contigs had low coverage (<1X coverage)
and low relative abundance (maximum relative abundance = 1.9% for TMED155 a putative
Cyanobacteria in TARA023-PROT-SRF; Supplemental Table 1). The high-quality genomes
accounted for 1.57-25.16% of the approximate microbial community as determined by the data-
rich-contigs (mean = 13.69%), with the ten most abundant genomes representing 0.61-10.31%
(Table 1).
Almost all of the contigs in the binned-contigs were low coverage, only a small subset of
6,350 contigs (6.6%) had >1X coverage in at least one sample. Of these contigs, 1,962 were
assigned to putative high-quality genomes, while the other contigs were placed in the low
completion bins. Further, an additional 22,470 contigs (Total bp = 79,422,500bp, mean =
3,535bp, and longest contig = 7,498bp) within data-rich-contigs (1.3%) had greater than >1X
coverage, but were not included in the binning protocol.
288
Concluding Statement
The goal of this project was to provided preliminary putative genomes from the *Tara Oceans*
microbial metagenomic datasets. The 290 putative high-quality genomes and 2,786 low
completion bins were created using the 20 samples and six stations from the Mediterranean Sea.
We will continue to generate putative high-quality genomes from additional *Tara Oceans*
dataset, starting with the Red Sea and Arabian Sea in the near future.
We would like to take some time to highlight to interesting results created within this
dataset. For new genomes from environmental organisms, this project created approximately 14
new *Cyanobacteria* genomes within the genera *Prochlorococcus* and *Synechococcus* and 33 new
SAR11 genomes. Three unconfirmed members related to the Candidate Phyla Radiation (CPR)
as determined by placement of an internal node between the *Parcubacteria* and *Microgenomates*
(with long-branch characteristics; TMED88) and a node basal to the CPR genomes, potentially
related to the *Wirthbacteria* (TMED70 and TMED22), on the concatenated ribosomal marker
tree. Additionally, there are putative genomes from the marine *Euryarchaeota* (n = 11),
Verrucomicrobia (n = 17), *Planctomycetes* (n = 14), and *Marinimicrobia* (n ≈ 5).
Some additional perplexing results include, TMED58 a putative *Deltaproteobacteria*
with taxonomic assignments from the NCBI NR database to the *Myoviridae*. This result occurs
due the presence of a few large contigs assigned to the bacteria and many small contigs assigned
to the virus. However, if these two entities should be binned together remains unresolved. Lastly,
the low completion bins may house distinct viral genomes. Of particular interest may be the 40
bins with 0% completion (based on single-copy marker genes), but that contain >500kb of
genetic material (including 3 bins with >1Mb). These large bins lacking markers may be good
candidates for research in to the marine "giant viruses" and episomal DNA sources (plasmids,
etc.).
It should be noted, researchers using this dataset should be aware that all of the genomes
generated from these samples (and additional stations, which are on-going) should be used as a
resource with some skepticism towards the results being an absolute. Like all results for
metagenome-assembled genomes, these genomes represent a best-guess approximation of a
taxon from the environment[33]. Researchers are encouraged to confirm all claims through various
genomic analyses and accuracy may require the removal of conflicting sequences.
319

Acknowledgements
We would like to thank iMicrobe and FigShare for hosting data for this research. We are
indebted to the *Tara Oceans* project and team for their commitment to open-access data that
allows data aficionados to indulge in the data and attempt to add to the body of science contained
within. And we thank the Center for Dark Energy Biosphere Investigations (C-DEBI) for
providing funding to BJT and JFH (OCE-0939654).

Author Contributions
BJT conceived of the project, performed all of the methods and analyses, and wrote the
manuscript. RS provided the origins of the workflow and invaluable feedback during the
execution of the methods and analyses. EDG provided feedback and troubleshooting using the
pre-release version of BinSanity. JFH provided funding. RS and JFH contributed to manuscript
editing and polishing. All authors have read the submitted draft of the manuscript.

Legends

Table 1. Megahit sequencing, recruitment to Megahit + Minimus2 contigs ≥2kb, and relative
abundance of high-quality genome results for each sample

Table 2. Statistics and the number contigs/assemblies at various steps during processing

Figure 1. Map illustrating the locations and size fractions sampled for the *Tara Oceans*
Mediterranean Sea datasets. Girus, 'giant virus' size fraction (0.22-1.6 μm). Bact, 'bacteria' size
fraction (0.22-1.6 μm). Prot, 'protist' size fraction (0.8-5.0 μm).

Figure 2. Workflow used to process *Tara Oceans* Mediterranean Sea metagenomic datasets.

Figure 3. FastTree approximate maximum-likelihood phylogenetic tree constructed with 37 and
785 16S rRNA genes from putative high-quality genomes and references, respectively.

Figure 4. Cladogram of a FastTree approximate maximum-likelihood phylogenetic tree
constructed using 16 syntenic, single-copy marker genes for 193 high-quality genomes and 1,729
reference genomes. Leaves denoting the position of the TMED genomes have been indicated by
extending beyond the edge of the tree.

Supplemental Table 1. Statistics and taxonomic and phylogenetic assignments for the putative
high-quality genomes

Supplemental Table 2. Statistics and CheckM taxonomy for low completion bins

Supplemental Table 3. Data file names, descriptions, and digital locations

References

1.   Falkowski, P. G., Fenchel, T. & DeLong, E. F. The Microbial Engines That Drive Earth's
Biogeochemical Cycles. *Science* **320,** 1034–1039 (2008).

2. Whitman, W. B., Coleman, D. C. & Wiebe, W. J. Prokaryotes: The unseen majority. *Proc.*
Natl. Acad. Sci. U.S.A. **95,** 6578–6583 (1998).
3. Karsenti, E. *et al.* A Holistic Approach to Marine Eco-Systems Biology. *Plos Biol* **9,**
e1001177–5 (2011).
4. Sunagawa, S. *et al.* Ocean plankton. Structure and function of the global ocean
microbiome. *Science* **348,** 1261359–1261359 (2015).
5. Li, D. *et al.* MEGAHIT v1.0: A fast and scalable metagenome assembler driven by
advanced methodologies and community practices. *Methods* **102,** 3–11 (2016).
6. Treangen, T. J., Sommer, D. D., Angly, F. E., Koren, S. & Pop, M. Next generation
sequence assembly with AMOS. *Curr Protoc Bioinformatics* **Chapter 11,** Unit 11.8
(2011).
7. Graham, E. D., Heidelberg, J. F. & Tully, B. J. BinSanity: Unsupervised Clustering of
Environmental Microbial Assemblies Using Coverage and Affinity Propagation. *bioRxiv*
(2016). doi:10.1101/069567
8. Longhurst, A., Sathyendranath, S., Platt, T. & Caverhill, C. An Estimate of Global
Primary Production in the Ocean From Satellite Radiometer Data. *Journal of Plankton*
Research **17,** 1245–1271 (1995).
9. Longhurst, A. *PROVINCES: THE SECONDARY COMPARTMENTS. Ecological*
Geography of the Sea 103–114 (Elsevier Inc., 2006). doi:10.1016/B978-0-12-455521-
1.50008-5
10. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Meth* **9,**
357–359 (2012).
11. Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics* **30,** 923–930 (2014).
12. Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W. CheckM:
assessing the quality of microbial genomes recovered from isolates, single cells, and
metagenomes. *Genome Res.* **25,** 1043–1055 (2015).
13. Frey, B. J. & Dueck, D. Clustering by passing messages between data points. *Science* **315,**
972–976 (2007).
14. Pedregosa, F. *et al.* Scikit-learn: Machine Learning in Python. *Journal of Machine*
Learning Research **12,** 2825–2830 (2011).
15. Matsen, F. A., Kodner, R. B. & Armbrust, E. V. pplacer: linear time maximum-likelihood
and Bayesian phylogenetic placement of sequences onto a fixed reference tree. *BMC*
Bioinformatics **11,** 538 (2010).
16. Albertsen, M. *et al.* Genome sequences of rare, uncultured bacteria obtained by
differential coverage binning of multiple metagenomes. *Nat Biotechnol* **31,** 533–538
(2013).
17. Hyatt, D., LoCascio, P. F., Hauser, L. J. & Uberbacher, E. C. Gene and translation
initiation site prediction in metagenomic sequences. *Bioinformatics* **28,** 2223–2230
(2012).
18. Buchfink, B., Xie, C. & Huson, D. H. Fast and sensitive protein alignment using
DIAMOND. *Nat Meth* **12,** 59–60 (2014).
19. Huson, D. H., Mitra, S., Ruscheweyh, H.-J., Weber, N. & Schuster, S. C. Integrative
analysis of environmental sequences using MEGAN4. *Genome Res.* **21,** 1552–1560
(2011).
20. Lagesen, K. *et al.* RNAmmer: consistent and rapid annotation of ribosomal RNA genes.

412   *Nucleic Acids Res.* **35,** 3100–3108 (2007).

413   21.   Pruesse, E., Peplies, J. & Glöckner, F. O. SINA: accurate high-throughput multiple
414   sequence alignment of ribosomal RNA genes. *Bioinformatics* **28,** 1823–1829 (2012).

415   22.   Ludwig, W. *et al.* ARB: a software environment for sequence data. *Nucleic Acids Res.* **32,**
416   1363–1371 (2004).

417   23.   Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and high
418   throughput. *Nucleic Acids Res.* **32,** 1792–1797 (2004).

419   24.   Capella-Gutiérrez, S., Silla-Martínez, J. M. & Gabaldón, T. trimAl: a tool for automated
420   alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25,** 1972–1973
421   (2009).

422   25.   Kearse, M. *et al.* Geneious Basic: an integrated and extendable desktop software platform
423   for the organization and analysis of sequence data. *Bioinformatics* **28,** 1647–1649 (2012).

424   26.   Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2--approximately maximum-likelihood
425   trees for large alignments. *PLoS ONE* **5,** e9490 (2010).

426   27.   Hug, L. A. *et al.* A new view of the tree of life. *Nature Microbiology* 1–6 (2016).
427   doi:10.1038/nmicrobiol.2016.48

428   28.   Finn, R. D., Clements, J. & Eddy, S. R. HMMER web server: interactive sequence
429   similarity searching. *Nucleic Acids Res.* **39,** W29–W37 (2011).

430   29.   Markowitz, V. M. *et al.* The integrated microbial genomes (IMG) system. *Nucleic Acids*
431   *Res.* **34,** D344–8 (2006).

432   30.   Huang, X. & Madan, A. CAP3: A DNA sequence assembly program. *Genome Res.* **9,**
433   868–877 (1999).

434   31.   Chevreux, B. *et al.* Using the miraEST assembler for reliable and automated mRNA
435   transcript assembly and SNP detection in sequenced ESTs. *Genome Res.* **14,** 1147–1159
436   (2004).

437   32.   Miller, J. R., Koren, S. & Sutton, G. Assembly algorithms for next-generation sequencing
438   data. *Genomics* **95,** 315–327 (2010).

439   33.   Sharon, I. & Banfield, J. F. Microbiology. Genomes from metagenomics. *Science* **342,**
440   1057–1058 (2013).

Table 1. Megahit sequencing, recruitment to Megahit + Minimus2 contigs ≥2kb, and relative abundance of high-quality genome results for each sample

TARA Sample Site	Size Fraction (Girus, Bacteria, or Protist)	Depth (Surface or DCM*)	No. of reads assembly	No. of initial Megahit assembly	N50[a] (bp; initial Megahit assembly)	Longest initial Megahit contigs (bp)	Recruitment (% to ≥2kb Megahit+Minimus2 contigs)	Relative abundance° of high-quality genomes (%)	Relative abundance° of ten most abundant genomes (%)
TARA007	Girus	DCM	178,519,830	1,318,470	828	220,754	72.84	14.64	6.35
TARA007	Girus	Surface	221,166,612	1,308,847	861	211,946	81.74	14.83	6.12
TARA007	Protist	DCM	744,458,992	4,667,618	654	188,635	19.45	8.60	3.18
TARA007	Protist	Surface	265,432,098	2,590,120	564	18,444	25.58	1.57	0.61
TARA009	Girus	DCM	416,553,274	2,796,841	831	1,643,839	69.48	14.16	6.32
TARA009	Girus	Surface	489,617,426	1,787,467	929	1,142,851	68.85	12.29	4.76
TARA009	Protist	DCM	329,036,110	1,938,636	613	95,724	22.07	13.35	4.20
TARA009	Protist	Surface	370,813,078	1,700,350	588	292,050	22.53	15.97	6.17
TARA018	Bacteria	DCM	408,021,182	2,520,645	840	1,573,060	76.22	11.49	3.18
TARA018	Bacteria	Surface	414,976,308	2,604,031	816	2,086,508	75.80	11.03	3.02
TARA023	Bacteria	DCM	147,400,552	1,273,576	830	213,456	76.08	13.29	4.09
TARA023	Bacteria	Surface	149,566,010	1,237,617	825	134,179	75.98	13.82	4.01
TARA023	Protist	DCM	508,610,652	2,707,801	734	336,689	28.23	25.07	7.83
TARA023	Protist	Surface	397,044,232	2,246,571	593	397,140	23.00	25.16	10.31
TARA025	Bacteria	DCM	386,627,816	2,516,865	806	388,546	69.77	14.55	5.35
TARA025	Bacteria	Surface	457,560,422	2,326,838	857	330,773	75.57	10.99	3.18
TARA030	Bacteria	DCM	346,837,034	1,968,945	1097	508,775	80.16	10.31	2.57
TARA030	Bacteria	Surface	478,785,582	1,639,697	1194	204,976	77.70	7.26	2.64
TARA030	Protist	DCM	426,896,616	1,620,343	616	478,892	15.12	17.83	5.13
TARA030	Protist	Surface	430,029,974	1,838,588	628	287,782	22.36	17.60	6.73

*DCM - deep chlorophyll maximum

[a]N50 - length of DNA sequence above which 50% of the total is contained

°relative abundance - determined using the reads recruited to the contigs ≥2kb in length (data-rich-contigs)

Table 2. Statistics and the number contigs/assemblies at various steps during processing

Contig Grouping	No. of contigs	N50*	Total sequence (bp)
Megahit assemblies 200-499bp	24,999,285	n.d.	9,293,098,676
Megahit assemblies 500-1,999bp	16,103,221	n.d.	13,382,057,993
Megahit assemblies ≥2kb	1,517,360	4,658	6,691,877,664
Megahit assemblies ≥2kb (post-CD-HIT-EST)	1,126,975	4,520	4,894,479,496
Minimus2 contigs	158,414	15,394	1,727,079,865
Minimus2 + unassembled Megahit contigs ≥2kb (data-rich-contigs)	660,937	5,466	3,612,405,904
Minimus2 + unassembled Megahit contigs ≥7.5kb (binned-contigs)	95,506	20,556	1,725,063,313

*N50 - length of DNA sequence above which 50% of the total is contained

Figure 1. Map illustrating the locations and size fractions sampled for the Tara Oceans Mediterranean Sea datasets. Girus, 'giant virus' size fraction (0.22-1.6 μm). Bact, 'bacteria' size fraction (0.22-1.6 μm). Prot, 'protist' size fraction (0.8-5.0 μm)

Figure 2. Workflow used to process Tara Oceans Mediterranean Sea metagenomic datasets. Black hash boxes, program or tool used with parameters.

Figure 3. FastTree approximate maximum-likelihood phylogenetic tree constructed with 37 and 785 16S rRNA genes from putative high-quality genomes and references, respectively.

Figure 4. Cladogram of a FastTree approximate maximum-likelihood phylogenetic tree constructed using 16 syntenic, single-copy marker genes for 193 high-quality genomes and 1,729 reference genomes. Leaves denoting the position of the TMED genomes have been indicated by extending beyond the edge of the tree.