290 Metagenome-assembled Genomes from the Mediterranean Sea: a resource for marine microbiology

Authors
Benjamin J. Tully[1‡], Rohan Sachdeva[2], Elaina D. Graham[2], and John F. Heidelberg[1,2]

‡ - corresponding author, tully.bj@gmail.com
1 – Center for Dark Energy Biosphere Investigations, University of Southern California, Los Angeles, CA 90089
2 – Department of Biological Sciences, University of Southern California, Los Angeles, CA 90089

Abstract
The *Tara Oceans* Expedition has provided large, publicly-accessible microbial metagenomic datasets from a circumnavigation of the globe. Utilizing several size fractions from the samples originating in the Mediterranean Sea, we have used current assembly and binning techniques to reconstruct 290 putative high-quality metagenome-assembled bacterial and archaeal genomes, with an estimated completion of ≥50%, and an additional 2,786 bins, with estimated completion of 0-50%. We have submitted our results, including initial taxonomic and phylogenetic assignments, for the putative high-quality genomes to open-access repositories for the scientific community to use in ongoing research.

Introduction
Microorganisms are a major constituent of the biology within the world's oceans and act as the important linchpins in all major global biogeochemical cycles[1]. Marine microbiology is among the disciplines at the forefront of advancements in understanding how microorganisms respond to and impact the local and large-scale environments. An estimated 10^{29} Bacteria and Archaea[2] reside in the oceans and represent an immense amount of poorly constrained, and ever evolving genetic diversity.

The *Tara Oceans* Expedition (2003-2010) encompassed a major endeavor to add to the body of knowledge collected during previous global ocean surveys to sample the genetic potential of microorganisms[3]. To accomplish this goal, *Tara Oceans* sampled planktonic organisms (viruses to fish larvae) at two major depths, the surface ocean and the mesopelagic. The amount of data collected was expansive and included 35,000 samples from 210 ecosystems[3]. The *Tara Oceans* Expedition generated and publically released 7.2 Tbp of metagenomic data from 243 ocean samples from throughout the global ocean, specifically targeting the smallest members of the ocean biosphere, the viruses, Bacteria and Archaea, and picoeukaryotes[4]. Initial work on these fractions produced a large protein database, totaling >40 million nonredundant protein sequences and identified >35,000 microbial operational taxonomic units (OTUs)[4].

Leveraging the publically available metagenomic sequences from the "girus" (giant virus; 0.22-1.6 µm), "bacteria" (0.22-1.6 µm), and "protist" (0.8-5 µm) size fractions, we have performed a new joint assembly of these samples using current sequence assemblers (Megahit[5]) and methods (combining assemblies from multiple sites using Minimus2[6]). These metagenomic assemblies were binned using BinSanity[7] in to 290 high-quality (low contamination) microbial genomes, ranging from 50-100% estimated completion. Environmentally derived genomes are imperative for a number of downstream applications, including comparative genomes,

metatranscriptomics, and metaproteomics. This series of genomic data can allow for the
recruitment of environmental "-omic" data and provide linkages between functions and
phylogenies. This method was initially performed on the seven sites from the Mediterranean Sea
containing microbial metagenomic samples (TARA007, -009, -018, -023, -025 and -030), but
will continue through the various Longhurst provinces[8] sampled during the *Tara Oceans* project
(Figure 1). All of the assembly data is publically available, including the initial Megahit
assemblies for each site from the various size fractions and depths and putative (minimal quality
control) genomes.
55
56   **Materials and Methods**
57
58   A generalized version of the following workflow is presented in Figure 2.
59
Sequence Retrieval and Assembly
All sequences for the reverse and forward reads from each sampled site and depth within
the Mediterranean Sea were accessed from European Molecular Biology Laboratory (EMBL)
utilizing their FTP service (Table 1). Paired-end reads from different filter sizes from each site
and depth (e.g., TARA0007, girus filter fraction, sampled at the deep chlorophyll maximum)
were assembled using Megahit[5] (v1.0.3; parameters: --preset, meta-sensitive). To keep consistent
with TARA sample nomenclature, "bacteria" or "BACT" will be used to encompass the size
fraction 0.22-1.6 μm. All of the Megahit assemblies were pooled in to two tranches based on
assembly size, ≤1,999bp, and ≥2,000bp. Longer assemblies (≥2kb) with ≥99% semi-global
identity were combined using CD-HIT-EST (v4.6; -T 90 -M 500000 -c 0.99 -n 10). The reduced
set of contiguous DNA fragments (contigs) was then cross-assembled using Minimus2[6] (AMOS
v3.1.0; parameters: -D OVERLAP=100 MINID=95).
72
Metagenome-assembled Genomes
Sequence reads were recruited against a subset of contigs (≥7.5kb) constructed during the
secondary assembly (Megahit + Minimus2) for each of the *Tara* samples using Bowtie2[9] (v4.1.2;
default parameters). Utilizing the SAM file output, read counts for each contig were determined
using featureCounts[10] (v1.5.0; default parameters). Coverage was determined for all contigs by
dividing the number of recruited reads by the length of the contig (reads/bp). Due to the low
coverage nature of the samples, in order to effectively delineate between contig coverage
patterns, the coverage values were transformed by multiplying by five (determined through
manual tuning). Transformed coverage values were then utilized to cluster contigs in to bins
utilizing BinSanity[7] (parameters: -p -3, -m 4000, -v 400, -d 0.9). Bins were assessed for the
presence of putative microbial genomes using CheckM[11] (v1.0.3; parameters: lineage_wf). Bins
were split in to three categories: (1) putative high quality genomes (≥50% complete and ≤10%
cumulative redundancy [% contamination – (% contamination × % strain heterogeneity ÷ 100)]);
(2) bins with "high" contamination (≥50% complete and ≥10% cumulative redundancy); and (3)
low completion bins (<50% complete).
The high contamination bins containing approximately two genomes, three genomes, or
≥4 genomes used the BinSanity refinement method (Binsanity-refine; -m 2000, -v 200, -d 0.9)
with variable preference values (-p) of -1000, -500, and -100, respectively. The resulting bins
were added to one of the three categories: putative high quality genomes, high contamination
bins, and low completion bins. The high contamination bins were processed for a third time with

the Binsanity-refine utilizing a preference of -100 (-p -100). These bins were given final
assignments to either the putative high quality genomes (some putative genomes had >10%
cumulative contamination, but have been designated) or low completion bins.
96   Any contigs not assigned to putative high-quality genomes were assessed using
BinSanity using raw coverage values. Two additional rounds of refinement were performed with
the first round of refinement using preference values based on the estimated number of
contaminating genomes (as above) and the second round using a set preference of -10 (-p -10).
Following this binning phase, contigs were assigned to high quality bins (*e.g.*, **Tara**
Mediterranean genome 1, referred to as TMED1, etc.), low completion bins with at least five
contigs (0-50% complete; TMEDlc1, etc. lc, **l**ow **c**ompletion), or were not placed in a bin
(Supplemental Table 1 & 2).
104
105   *Taxonomic and Phylogenetic Assignment of High Quality Genomes*
106   The bins representing the high quality genomes were assessed for taxonomy and
107   phylogeny using multiple methods to provide a quick reference for selecting genomes of interest.
108   Taxonomy as assigned using the putative placement provided via CheckM during the pplacer[12]
109   step of the analysis to the lowest taxonomic placement (parameters: tree_qa -o 2). This step was
110   also performed for all low completion bins.
111   Two separate attempts were made to assign the high quality genomes a phylogenetic
assignment. High quality genomes were searched for the presence of the full-length 16S rRNA
gene sequence using RNAmmer[13] (v1.2; parameters: -S bac -m ssu). All full-length sequences
were aligned to the SILVA SSU reference database (Ref123) using the SINA web portal
aligner[14] (https://www.arb-silva.de/aligner/). These alignments were loaded in to ARB[15] (v6.0.3),
manually assessed, and added to the non-redundant 16S rRNA gene database (SSURef123
NR99) using ARB Parsimony (Quick) tool (parameters: default). A selection of the nearest
neighbors to the *Tara* genome sequences were selected and used to construct a 16S rRNA
phylogenetic tree. Genome-identified 16S rRNA sequences and SILVA reference sequences
were aligned using MUSCLE[16] (v3.8.31; parameters: -maxiters 8) and processed by the
automated trimming program trimAL[17] (v1.2rev59; parameters: -automated1). Automated
trimming results were assessed manually in Geneious[18] (v6.1.8) and trimmed where necessary
(positions with >50% gaps) and re-aligned with MUSCLE (parameters: -maxiters 8). An
approximate maximum likelihood (ML) tree with pseudo-bootstrapping was constructed using
FastTree[19] (v2.1.3; parameters: -nt -gtr -gamma; Figure 3).
126   High-quality genomes were assessed for the presence of the 16 ribosomal markers genes
used in Hug, *et al*. (2016)[20]. Putative CDSs were determined using Prodigal (v2.6.3; parameters:
-m -p meta) and were searched using HMMs for each marker using HMMER[21] (v3.1b2;
parameters: hmmsearch --cut_tc --notextw). If a genome had multiple copies of any single
marker gene, neither was considered, and only genomes with ≥8 markers were used to construct
a phylogenetic tree. Markers identified from the high quality genomes were combined with
markers from 1,729 reference genomes that represent the major bacterial phylogenetic groups (as
presented by IMG[22]). Archaeal reference sequences were not included; however, none of the
putative archaeal environmental genomes had a sufficient number of markers for inclusion on the
tree. Each marker gene was aligned using MUSCLE (parameters: -maxiters 8) and automatically
trimmed using trimAL (parameters: -automated1). Automated trimming results were assessed (as
above) and re-aligned with MUSCLE, as necessary. Final alignments were concatenated and

used to construct an approximate ML tree with pseudo-bootstrapping with FastTree (parameters:
-gtr -gamma; Figure 4).
140
Relative Abundance of High Quality Genomes
To set-up a baseline that could approximate the "microbial" community (Bacteria, Archaea and
viruses) present in the various *Tara* metagenomes, which included filter sizes specifically
targeting both protists and giruses, reads were recruited against all contigs generated from the
Minimus2 and Megahit assemblies ≥2kb using Bowtie2 (default parameters). Some assumptions
were made that contigs <2kb would include, low abundance bacteria and archaea, bacteria and
archaea with high degrees of repeats/assembly poor regions, fragmented picoeukaryotic
genomes, and problematic read sequences (low quality, sequencing artefacts, etc.). All relative
abundance measures are relative to the number of reads recruited to the assemblies ≥2kb. Read
counts were determined using featureCounts (as above). Length-normalized relative abundance
values were determined for each high quality genome for each sample:

$$\frac{\frac{Reads}{bp}\,per\,genome}{\sum \frac{Reads}{bp}\,all\,genomes} \times \frac{\sum Recruited\,reads\,to\,genomes}{\sum Recruited\,reads\,to\,all\,contigs\,(\geq 2kb)} \times 100$$

152
Data Availability
This project has been deposited at DDBJ/ENA/GenBank under the BioProject accession
no. #### and drafts of genomes are available with accession no. #####-#####. Additional files
have been provided and are available through FigShare
(https://dx.doi.org/10.6084/m9.figshare.3545330), such as: all contigs from Minimus2 + Megahit
output used for binning and community assessment; contig read counts per sample; the putative
genome contigs and Prodigal-predicted nucleotide and protein putative CDS FASTA files; the
ribosomal marker HMM profiles; reference genome markers; high quality genome markers; low
completion bins, and contigs without a bin. All contigs generated using Megahit from each
sample are available through iMicrobe (http://data.imicrobe.us/project/view/261).

163
Results
Assembly
The initial Megahit assembly was performed on the publicly available reads for *Tara*
stations 007, 009, 018, 023, 025, 030. Starting with 147-744 million reads per sample, the
Megahit assembly process generated 1.2-4.6 million assemblies with a mean N_{50} and longest
contig of 785bp and 537kb, respectively (Table 1). In general, the assemblies generated from the
Tara samples targeting the protist size fraction (0.8-5 μm) had a shorter N_{50} value than the
bacteria size fractions (mean: 554bp vs 892bp, respectively). Assemblies from the Megahit
assembly process were pooled and separated by length. Of the 42.6 million assemblies generated
during the first assembly, 1.5 million were ≥2kb in length (Table 2). Several attempts were made
to assemble the shorter contigs, but publicly available overlap-consensus assemblers (Newbler
[454 Life Sciences], cap3[23], and MIRA[24]) failed on multiple attempts. Processing the ≥2kb
assemblies from all of the samples through CD-HIT-EST reduced the total to 1.1 million contigs
≥2kb. This group of contigs was subjected to the secondary assembly through Minimus2,
generating 158,414 new contigs (all ≥2kb). The secondary contigs were combined with the

Megahit contigs that were not assembled by Minimus2. This provided a contig dataset consisting
of 660,937 contigs, all ≥2kb in length (Table 2; further referred to as data-rich-contigs).
181
Binning
The set of data-rich-contigs was used to recruit the metagenomic reads from each sample
using Bowtie2. The data-rich-contigs recruited 15-81% of the reads depending on the sample. In
general, the protist size fraction recruited substantially fewer reads than the girus and bacteria
size fractions (mean: 19.8% vs 75.0%, respectively) (Table 1). For the protist size fraction, the
"missing" data for these recruitments likely results from the poor assembly of more complex and
larger eukaryotic genomes. The fraction of the reads that do not recruit in the girus and bacterial
size fraction samples could be accounted for by the large number of low quality assemblies (200-
500bp) and reads that could not be assembled due to low abundance or high complexity (Table
2).
Unsupervised binning was performed using both transformed and raw coverage values
for a subset of 95,506 contigs from the data-rich-contigs that were ≥7.5kb (referred to further as
binned-contigs). Binning using the transformed coverage data generated 237 putative high-
quality genomes (12 putative genomes are of slightly lower quality with >10% redundancy and
have been noted) containing 15,032 contigs (Supplemental Information S1). Contigs not in
putative genomes were re-binned based on raw coverage values, generating 53 additional
putative high-quality genomes encompassing 3,348 contigs. In total, 290 putative high-quality
genomes were generated with 50-100% completion (mean: 69%) with a mean length and number
of putative CDS of 1.7Mbp and 1,699, respectively (Supplemental Information S1). All other
contigs were grouped in to bins with at least five contigs, but with estimated completion of 0-
50% (2,786 low completion bins; 74,358 contigs; Supplemental Information S2) or did not bin
(2,732 contigs). Nearly a quarter of the low completion bins (24.7%) have an estimated
completion of 0%.
205
Taxonomy, Phylogeny, & Potential Organisms of Interest
The 290 putative high-quality genomes had a taxonomy assigned to it via CheckM during
the pplacer step. All of the genomes, except for 20, had an assignment to at least the Phylum
level, and 83% of the genomes had an assignment to at least the Class level (Supplemental
Information S1).
Phylogenetic information was determined for as many genomes as possible. Genomes
were assessed for the presence of full-length 16S rRNA genes. In total, 37 16S rRNA genes were
detected in 35 genomes. 16S rRNA genes can prove to be problematic during the assembly steps
due the high level of conservation that can break contigs[25] (Figure 3). Additionally, the
conserved regions of the 16S rRNA, depending on the situation, can over- or under-recruit reads,
resulting in coverage variations that can misplace contigs in to the incorrect genome. As such,
several of the 16S rRNA phylogenetic placements support the taxonomic assignments, while
some are contradictory. Further analysis should allow for the determination of the most
parsimonious result.
Beyond the 16S rRNA gene, genomes were searched for 16 conserved, syntenic
ribosomal markers. Sufficient markers (≥8) were identified in 193 of the genomes (67%) and
placed on a tree with 1,729 reference sequences (Figure 4). Phylogenies were then assigned to
the lowest taxonomic level that could be confidently determined. These putative results reveal a
number of genomes were generated that represent multiple clades for which environmental

genomic information remains limited, including: *Planctomycetes*, *Verrucomicrobia*,
Marinimicrobia, *Cyanobacteria*, and uncultured groups within the *Alpha-* and
Gammaproteobacteria.
228
Relative Abundance
A length-normalized relative abundance value was determined for each genome in each
sample based on the number of reads recruited to the data-rich-contigs. The relative abundance
for the individual genomes was determined based on this portion of the dataset (Supplemental
Information S3). In general, the genomes had low relative abundance (maximum relative
abundance = 1.9% for TMED155 a putative *Cyanobacteria* at site TARA023 from the protistan
size fraction sampled at the surface; Supplemental Table 1). The high-quality genomes
accounted for 1.57-25.16% of the approximate microbial community as determined by the data-
rich-contigs (mean = 13.69%), with the ten most abundant genomes in a sample representing
0.61-10.31% (Table 1).
239
Concluding Statement
The goal of this project was to provided preliminary putative genomes from the *Tara*
Oceans microbial metagenomic datasets. The 290 putative high-quality genomes and 2,786 low
completion bins were created using the 20 samples and six stations from the Mediterranean Sea.
Initial assessment of the phylogeny of these metagenomic-assembled genomes indicates
several new genomes from environmentally relevant organisms, including, approximately 14
new *Cyanobacteria* genomes within the genera *Prochlorococcus* and *Synechococcus* and 33 new
SAR11 genomes. Additionally, there are putative genomes from the marine *Euryarchaeota* (n =
11), *Verrucomicrobia* (n = 17), *Planctomycetes* (n = 14), and *Marinimicrobia* (n ≈ 5).
Additionally, the low completion bins may house distinct viral genomes. Of particular interest
may be the 40 bins with 0% completion (based on single-copy marker genes), but that contain
>500kb of genetic material (including 3 bins with >1Mb). These large bins lacking markers may
be good candidates for research in to the marine "giant viruses" and episomal DNA sources
(plasmids, etc.).
It should be noted, researchers using this dataset should be aware that all of the genomes
generated from these samples should be used as a resource with some skepticism towards the
results being an absolute. Like all results for metagenome-assembled genomes, these genomes
represent a best-guess approximation of a taxon from the environment[26]. Researchers are
encouraged to confirm all claims through various genomic analyses and accuracy may require
the removal of conflicting sequences.
260

261 **Acknowledgements**
262   We are indebted to the *Tara Oceans* project and team for their commitment to open-
263 access data that allows data aficionados to indulge in the data and attempt to add to the body of
264 science contained within. And we thank the Center for Dark Energy Biosphere Investigations (C-
265 DEBI) for providing funding to BJT and JFH (OCE-0939654).
266

Author Contributions
BJT conceived of the project, performed all of the methods and analyses, and wrote the
manuscript. RS provided the origins of the workflow and invaluable feedback during the
execution of the methods and analyses. EDG provided feedback and troubleshooting using the

pre-release version of BinSanity. JFH provided funding. RS and JFH contributed to manuscript
editing and polishing. All authors have read the submitted draft of the manuscript.
273
Legends
275
276 Table 1. Statistics for Megahit assemblies, recruitment to data-rich-contigs, and relative
277 abundance of high-quality genome results for each sample
278
279 Table 2. Assembly statistics at various steps during processing
280
281 Figure 1. Map illustrating the locations and size fractions sampled for the *Tara Oceans*
282 Mediterranean Sea datasets. Girus, 'giant virus' size fraction (0.22-1.6 μm). Bact, 'bacteria' size
283 fraction (0.22-1.6 μm). Prot, 'protist' size fraction (0.8-5.0 μm).
284
285 Figure 2. Workflow used to process *Tara Oceans* Mediterranean Sea metagenomic datasets.
286
287 Figure 3. FastTree approximate maximum-likelihood phylogenetic tree constructed with 37 and
288 785 16S rRNA genes from putative high-quality genomes and references, respectively.
289
290 Figure 4. Cladogram of a FastTree approximate maximum-likelihood phylogenetic tree
291 constructed using 16 syntenic, single-copy marker genes for 193 high-quality genomes and 1,729
292 reference genomes. Leaves denoting the position of the TMED genomes have been indicated by
293 extending beyond the edge of the tree. Sequence alignment is available in Supplemental
294 Information S4. Phylogenetic tree newick file is available in Supplemental Information S5.
295
Supplemental Information
297
298 Supplemental Information S1. Statistics, taxonomic and phylogenetic assignments for the
299 putative high-quality genomes.
300
301 Supplemental Information S2. Statistics and CheckM taxonomy for low completion bins.
302
303 Supplemental Information S3. Relative abundance values determined for each genome based the
304 length-normalized fraction of reads recruited to the genome relative to reads recruited for the
305 data-rich-contigs.
306
307 Supplemental Information S4. Concatenated MUSCLE alignment file of 16 ribosomal marker
308 proteins used to construct Figure 4.
309
310 Supplemental Information S5. Newick file of concatenated 16 ribosomal marker proteins,
311 including FastTree determined local support values using the Shimodaira-Hasegawa test.
312
References
314
1.   Falkowski, P. G., Fenchel, T. & DeLong, E. F. The Microbial Engines That Drive Earth's
Biogeochemical Cycles. *Science* **320,** 1034–1039 (2008).

2. Whitman, W. B., Coleman, D. C. & Wiebe, W. J. Prokaryotes: The unseen majority. *Proc.*
Natl. Acad. Sci. U.S.A. **95,** 6578–6583 (1998).
3. Karsenti, E. *et al.* A Holistic Approach to Marine Eco-Systems Biology. *Plos Biol* **9,**
e1001177–5 (2011).
4. Sunagawa, S. *et al.* Ocean plankton. Structure and function of the global ocean
microbiome. *Science* **348,** 1261359–1261359 (2015).
5. Li, D. *et al.* MEGAHIT v1.0: A fast and scalable metagenome assembler driven by
advanced methodologies and community practices. *Methods* **102,** 3–11 (2016).
6. Treangen, T. J., Sommer, D. D., Angly, F. E., Koren, S. & Pop, M. Next generation
sequence assembly with AMOS. *Curr Protoc Bioinformatics* **Chapter 11,** Unit 11.8
(2011).
7. Graham, E. D., Heidelberg, J. F. & Tully, B. J. BinSanity: unsupervised clustering of
environmental microbial assemblies using coverage and affinity propagation. *PeerJ* **5,**
e3035–19 (2017).
8. Longhurst, A., Sathyendranath, S., Platt, T. & Caverhill, C. An Estimate of Global
Primary Production in the Ocean From Satellite Radiometer Data. *Journal of Plankton*
Research **17,** 1245–1271 (1995).
9. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Meth* **9,**
357–359 (2012).
10. Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics* **30,** 923–930 (2014).
11. Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W. CheckM:
assessing the quality of microbial genomes recovered from isolates, single cells, and
metagenomes. *Genome Res.* **25,** 1043–1055 (2015).
12. Matsen, F. A., Kodner, R. B. & Armbrust, E. V. pplacer: linear time maximum-likelihood
and Bayesian phylogenetic placement of sequences onto a fixed reference tree. *BMC*
Bioinformatics **11,** 538 (2010).
13. Lagesen, K. *et al.* RNAmmer: consistent and rapid annotation of ribosomal RNA genes.
Nucleic Acids Res. **35,** 3100–3108 (2007).
14. Pruesse, E., Peplies, J. & Glöckner, F. O. SINA: accurate high-throughput multiple
sequence alignment of ribosomal RNA genes. *Bioinformatics* **28,** 1823–1829 (2012).
15. Ludwig, W. *et al.* ARB: a software environment for sequence data. *Nucleic Acids Res.* **32,**
1363–1371 (2004).
16. Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and high
throughput. *Nucleic Acids Res.* **32,** 1792–1797 (2004).
17. Capella-Gutiérrez, S., Silla-Martínez, J. M. & Gabaldón, T. trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25,** 1972–1973
(2009).
18. Kearse, M. *et al.* Geneious Basic: an integrated and extendable desktop software platform
for the organization and analysis of sequence data. *Bioinformatics* **28,** 1647–1649 (2012).
19. Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2--approximately maximum-likelihood
trees for large alignments. *PLoS ONE* **5,** e9490 (2010).
20. Hug, L. A. *et al.* A new view of the tree of life. *Nature Microbiology* 1–6 (2016).
doi:10.1038/nmicrobiol.2016.48
21. Finn, R. D., Clements, J. & Eddy, S. R. HMMER web server: interactive sequence
similarity searching. *Nucleic Acids Res.* **39,** W29–W37 (2011).

22.   Markowitz, V. M. *et al.* The integrated microbial genomes (IMG) system. *Nucleic Acids*
364   *Res.* **34,** D344–8 (2006).
365   23.   Huang, X. & Madan, A. CAP3: A DNA sequence assembly program. *Genome Res.* **9,**
366   868–877 (1999).
367   24.   Chevreux, B. *et al.* Using the miraEST assembler for reliable and automated mRNA
368   transcript assembly and SNP detection in sequenced ESTs. *Genome Res.* **14,** 1147–1159
369   (2004).
370   25.   Miller, J. R., Koren, S. & Sutton, G. Assembly algorithms for next-generation sequencing
371   data. *Genomics* **95,** 315–327 (2010).
372   26.   Sharon, I. & Banfield, J. F. Microbiology. Genomes from metagenomics. *Science* **342,**
373   1057–1058 (2013).
374

Figure 1. Map illustrating the locations and size fractions sampled for the Tara Oceans Mediterranean Sea datasets. Girus, 'giant virus' size fraction (0.22-1.6 μm). Bact, 'bacteria' size fraction (0.22-1.6 μm). Prot, 'protist' size fraction (0.8-5.0 μm)

Figure 4. Cladogram of a FastTree approximate maximum-likelihood phylogenetic tree constructed using 16 syntenic, single-copy marker genes for 193 high-quality genomes and 1,729 reference genomes. Leaves denoting the position of the TMED genomes have been indicated by extending beyond the edge of the tree. Sequence alignment is available in Supplemental Information S4. Phylogenetic tree newick file is available in Supplemental Information S5.

Figure 2. Workflow used to process Tara Oceans Mediterranean Sea metagenomic datasets.
Black hash boxes, program or tool used with parameters.

Figure 3. FastTree approximate maximum-likelihood phylogenetic tree constructed with 37 and 785 16S rRNA genes from putative high-quality genomes and references, respectively.

Table 1. Statistics for Megahit assemblies, recruitment to data-rich-contigs, and relative abundance of high-quality genome results for each sample

TARA Sample Site	Size Fraction (Girus, Bacteria, or Protist)	Depth (Surface or DCM*)	No. of reads	No. of initial Megahit assembly	N50[a] (bp; initial Megahit assembly)	Longest initial Megahit assembly (bp)	Recruitment (% data-rich-contigs)	Relative abundance[o] of high-quality genomes (%)	Relative abundance[o] of ten most abundant genomes (%)
TARA007	Girus	DCM	178,519,830	1,318,470	828	220,754	72.84	14.64	6.35
TARA007	Girus	Surface	221,166,612	1,308,847	861	211,946	81.74	14.83	6.12
TARA007	Protist	DCM	744,458,992	4,667,618	654	188,635	19.45	8.60	3.18
TARA007	Protist	Surface	265,432,098	2,590,120	564	18,444	25.58	1.57	0.61
TARA009	Girus	DCM	416,553,274	2,796,841	831	1,643,839	69.48	14.16	6.32
TARA009	Girus	Surface	489,617,426	1,787,467	929	1,142,851	68.85	12.29	4.76
TARA009	Protist	DCM	329,036,110	1,938,636	613	95,724	22.07	13.35	4.20
TARA009	Protist	Surface	370,813,078	1,700,350	588	292,050	22.53	15.97	6.17
TARA018	Bacteria	DCM	408,021,182	2,520,645	840	1,573,060	76.22	11.49	3.18
TARA018	Bacteria	Surface	414,976,308	2,604,031	816	2,086,508	75.80	11.03	3.02
TARA023	Bacteria	DCM	147,400,552	1,273,576	830	213,456	76.08	13.29	4.09
TARA023	Bacteria	Surface	149,566,010	1,237,617	825	134,179	75.98	13.82	4.01
TARA023	Protist	DCM	508,610,652	2,707,801	734	336,689	28.23	25.07	7.83
TARA023	Protist	Surface	397,044,232	2,246,571	593	397,140	23.00	25.16	10.31
TARA025	Bacteria	DCM	386,627,816	2,516,865	806	388,546	69.77	14.55	5.35
TARA025	Bacteria	Surface	457,560,422	2,326,838	857	330,773	75.57	10.99	3.18
TARA030	Bacteria	DCM	346,837,034	1,968,945	1097	508,775	80.16	10.31	2.57
TARA030	Bacteria	Surface	478,785,582	1,639,697	1194	204,976	77.70	7.26	2.64
TARA030	Protist	DCM	426,896,616	1,620,343	616	478,892	15.12	17.83	5.13
TARA030	Protist	Surface	430,029,974	1,838,588	628	287,782	22.36	17.60	6.73

*DCM - deep chlorophyll maximum

[a]N50 - length of DNA sequence above which 50% of the total is contained

[o]relative abundance - determined using the reads recruited data-rich-contigs

Table 2. Assembly statistics at various steps during processing

Contig Grouping	No. of contigs	N50*	Total sequence (bp)
Megahit assemblies 200-499bp	24,999,285	n.d.	9,293,098,676
Megahit assemblies 500-1,999bp	16,103,221	n.d.	13,382,057,993
Megahit assemblies ≥2kb	1,517,360	4,658	6,691,877,664
Megahit assemblies ≥2kb (post-CD-HIT-EST)	1,126,975	4,520	4,894,479,496
Minimus2 contigs	158,414	15,394	1,727,079,865
Minimus2 + unassembled Megahit contigs ≥2kb (data-rich-contigs)	660,937	5,466	3,612,405,904
Minimus2 + unassembled Megahit contigs ≥7.5kb (binned-contigs)	95,506	20,556	1,725,063,313

*N50 - length of DNA sequence above which 50% of the total is contained