Non-Mendelian inheritance of SNP markers reveals extensive chromosomal translocations in dioecious hops (*Humulus lupulus* L.)

Dong Zhang, Nicholi J. Pitra, Mark C. Coles, Edward S. Buckler, Paul D. Matthews

Hopsteiner, S.S. Steiner, Inc., New York, New York, 10065 (M.C.C, P.D.M, N.J.P, D.Z.), Institute for Genomic Diversity, Cornell University, Ithaca, New York, 14853 (E.S.B, D.Z), Agricultural Research Service, United States Department of Agriculture, Ithaca, New York, 14853, (E.S.B.)

Footnotes:
E.S.B. provided guidance on statistical analyses and project design, N.J.P. prepared sequencing libraries, N.P. and D.Z. analyzed data and interpreted results, M.C.C. collected samples and prepared DNA extracts, P.D.M created germplasm resources, devised and directed the studies, D.Z., N.J.P. wrote and P.D.M edited the manuscript.

The study was funded by Hopsteiner, S.S. Steiner, Inc.

Corresponding author emails: zhangdong20046@gmail.com, pmatthews@hopsteiner.com

Abstract

Genome-wide meiotic recombination structures, sex chromosomes, and candidate genes for sex determination were discovered among *Humulus* spp. by application of a novel, high-density molecular marker system: ~1.2M single nucleotide polymorphisms (SNPs) were profiled with genotyping-by-sequencing (GBS) among 4512 worldwide accessions, including 4396 cultivars and landraces and 116 wild accessions of hops. Pre-qualified GBS markers were validated by inferences on families, population structures and phylogeny. Candidate genes discovered for several traits, including sex and drought stress-resistance, demonstrate the quality and utility of GBS SNPs for genome-wide association studies (GWAS) and Fst analysis in hops. Most importantly, pseudo-testcross mappings in F1 families delineated non-random linkage of Mendelian and non-Mendelian markers: structures that are indicative of unusual meiotic events which may have driven the evolution and cultivation of hops.

Introduction

The Cannabaceae family of flowering plants has a rich history of contributions to humanity, with the promise of still greater contributions as result new commercial values and invigorated research in two members, *Humulus lupulus* (hop) (2n = 20) and *Cannabis sativa* (hemp, marijuana) (2n = 20) (van Bakel et al., 2011), which diverged around 27.8 Myr (Laursen, 2015). Hop (*H. lupulus*) is a high-climbing, herbaceous perennial, dioecious vine, and has a long history of use as flavoring and stability agent in beer as well as nutraceutical medicine, bio-fuel fermentations and animal fodder (Siragusa et al., 2008). For example, studies of specific hop-derived prenylflavonoids in prevention of cancer, dyslipidemia, and postmenopausal systems spawn interest in metabolic engineering and marker-directed breeding in hop (Ososki and Kennelly, 2003; Stevens and Page, 2004; Nagel et al., 2008; Miranda et al., 2016). The properties of its reproductive system, such as dioecy and obligate outcrossing, high heterozygosity and a large genome size (~2.6Gb), and complex sex-determination system (Neve, 1958), render challenges of genetic dissection of complex traits in hops.

Wild *H. lupulus* is represented by at least five extant species: (1) var. *lupulus* for

European wild hops; (2) var. *cordifolius* mainly distributed in Japan, and vars. (3)

neomexicanus (in the Southwestern U.S.), (4) *pubescens* (in the Eastern/Midwestern U.S.)

and (5) *lupuloides* (throughout the northern Great Plains); spreading throughout North

America. Asian and North American wild hops resemble each other morphologically,

suggesting a genetically closer relationship, while they differ more so from European

hops (Murakami et al., 2006). Many contemporary cultivars are hybrids of North

American and European genetic materials, in which North American hops have been

characterized by their higher bitterness and aroma (Reeves and Richards, 2011) than

European cultivars. In other crops, breeding programs have successfully exploited novel

genetic variations from wild exotic germplasms into modern cultivars (Tanksley and

McCouch, 1997; Bradshaw, 2016) to gain desirable traits such as favored flavors, drought

tolerance, and disease resistance. Successes with wild resources and predictions of

climate change have spurred resurgence in conservation biology of plant genetic

resources (Castañeda-Álvarez et al., 2016; Gruber, 2016).

Molecular marker systems have been developed for hops and applied in genetic mapping

of families (reviewed in Henning et al., 2015), including non-referenced GBS markers

(Matthews et al., 2013) and GWAS applied to disease resistance (Henning et al., 2015)

and sex determination (Hill et al., 2016), using whole genome-referenced GBS markers.

However, genetic phenomenon in hops still is under-explained. For example: (1)

significant segregation distortion from Mendelian segregation expectations have been

repeatedly reported in mapping populations, indicating that the segregation bias was due

to genetic properties rather than genotyping errors (Seefelder et al., 2000; McAdam et al.,

2013); (2) Unusual female-biased sex ratios have been observed in controlled crosses,

where high pollen loads were applied (Jakse et al., 2008). In other genetic systems,

segregation distortion is a result of various patterns of meiotic drive and chromosomal

(re)arrangements (Taylor and Ingvarsson, 2003). Examples of meiotic drive include the B

chromosomes observed in insects (Fontana and Vickery, 1973), the X-linked meiotic

drive in Drosophila species (Lyttle, 1993) and the t-haplotype in mice (Silver, 1993).

Some well-known examples of chromosomal (re)arrangements in plants include the

neocentromeres (knobs) of maize (Buckler et al., 1999), the translocation heterozygosity

in some species of the genus *Clarkia* (Snow, 1960), *Oenothera* (Rauwolf et al., 2008;

Golczyk et al., 2014) and *Viscum* (mistletoe) (Wiens and Barlow, 1975; Rauwolf et al.,

2008).

One classical cytogenetic analysis (Sinotô, 1929) suggested that at least 4 chromosomes

in the male of var. *cordifolius* are involved in reciprocal translocation, detected by a

chromosome chain in a shape of 'zigzag' during the first meiotic metaphase. Our

characterization with an unprecedented system of high density genome-wide SNP

markers allows detailed examination of translocation heterozygosity, which predisposes

rediscovery of the mode of inheritance in this species. Cytogenetic investigations in

Clarkia (Snow, 1960) and *Oenothera* (Rauwolf et al., 2008; Golczyk et al., 2014) may

provide comparative insight on putative translocation heterozygosity in *Humulus*.

However, except for Japanese wild hops (var. *cordifolius*), heterozygotes in males were

reported neither in European wild types (var. *lupulus*) nor in North American wild types

(var. *neomexicanus*) (Winge, 1932; Jacobsen, 1957; Shephard and Parker, 2000). To date,

size ratio of sex chromosomes in wild hops is not fully explained: a description of

sex-chromosome variability across subspecies in hops is widely accepted (Shephard and

Parker, 2000). Moreover, lack of cytogenetic studies in modern hybrids confuses the

mode of genetic inheritance in hops. New insight into genetic parameters in *Humulus* is

herein offered by a next generation sequencing (NGS) platform applied to an

unprecedented 4512 accessions, including 22 sibling and halfsibling families, genotyped

with GBS SNP marker system, comprising 1,235,148 SNPs. Previously reported NGS

GBS studies in hop (Matthews, 2013, Henning et al., 2015; Hill et al., 2016) have focused

on smaller (511) association panels and reduced marker sets, have not addressed

population structure. Furthermore, filtering against SNPs in segregation distortion (SD),

thus, ignoring rather than characterizing chromosome regions in SD, resulted in a

low-power genetic system that dubiously supported candidate gene discoveries. We

present a qualified, high-density marker system applied across a structured association

panel and 22 families, analysis of population structure among exotic vs. cultivated

accessions within the panel, and a set of strongly supported candidates associated with

sex determination. Inclusion, rather than ignorance, of markers in strong SD, has led to

testable hypotheses of chromosome structure and recombination constraints in hop, which

requires re-assessment of breeding strategies. The need for new, classical cytogenetic

studies is implicated by NGS exploration.

Results

Phylogenetic relationships of modern cultivars and North American indigenous

exotics

European var. *lupulus* is the ancestor of most commercial hops used today, thereby

commercial cultivars retain a large proportion of var. *lupulus* genome. In addition, the

genetic diversity of hop crop has been contributed by mostly male donors from North

America and Asia. To understand the phylogenetic relatedness in hop races, we focused

on a subset of 251 accessions, consisting of 183 modern cultivars (CV) including all

progenitors of F1 families in this study and 68 wild hops. The neighbor-joining tree

(Figure 1a) shows three distinct clusters. The modern cultivars were clustered together,

indicating a common derivation in domestication of hops. The other two clusters reflect

geographical origins of North American wild hops (Figure 1b), in which one group

(SW_wild) includes 22 Southwestern U.S. wild hops (represented by var. *neomexicanus*),

and the other group contains 20 wild hops (represented by var. *lupuloides*) from Northern

U.S./Canada (N_wild) and 3 (represented by var. *pubescens*) from Midwestern U.S.

(MW_wild). Seven wild individuals from Kazakhstan are intermediate among the modern

cultivars, consistent with a previous inference (Murakami et al., 2006) of a close genetic

relationship between wild hops from Europe and the Altai region (close to western China,

located on boundaries of Russia, Mongolia, Kazakhstan and China).

The level of population differentiation, fixation index (Fst), was measured across the

three clusters. SW_wild exhibits relatively close genetic relationship (Fst = 0.1663) with

N_wild, apparently supporting relatively close ancestry and geographical origins of the

two wild populations. Genetic distinction between the modern cultivars and the North

American wild hops is evident: [Fst (CV vs. SW_wild) = 0.31; Fst (CV vs. N_wild) =

0.295].

To demonstrate the population structure of F1 families (N ≥ 60) (Figure 2a) in our dataset, we used a nonlinear algorithm (implemented in Python scikit-learn), t-Distributed Stochastic Neighbor Embedding (t-SNE) (Maaten and Hinton, 2008), for dimension reduction of the IBS-based distance matrix. The F1 families derived from genetically divergent progenitors can be easily distinguished from one another, while the half-sibling families exhibit ambiguous clustering patterns. A network of pedigree (Figure 2b) reflects that the F1 families in the current dataset were mostly derived from genetically related varieties.

Segregation distortion in hybrids

Ubiquitous presence of non-Mendelian factors results from inherent mechanisms in *H. lupulus* or from genotyping errors. While genotyping errors are random, the genuinely distorted markers can exhibit pronounced correlation with Mendelian segregation markers. On the basis of clustering of pairwise Spearman's correlation in pseudo-testcross (Pt) markers in three F1 families, we hypothesize that (1) severe SD tends to occur near breakpoints to favor translocation complexes; (2) patterns of linkage can differ across the three populations; (3) a large scale, perhaps genome-wide, meiotic chromosomal complex might occur in the progenitors of the three populations; and (4) translocation heterozygosity may be a ubiquitous phenomenon in hybrids of hops, implicating its genetics and biological significance throughout the cultivation history of hops.

Pseudo-testcross in families "144" and "247" shows multiple 'super' linkage groups in terms of their size and inter-marker correlation (Figure 3a,S2a). In family "265", linkage groups tend to have equal size (Figure S2b), but exhibit relatively high correlation to one another. Alignments across the three sets of clusters (before phasing coupling groups) (Figure 3b,3c) show most of anchor (common) markers were distinctly clustered. Using a nonlinear algorithm (implemented in Python scikit-learn), locally linear embedding (LLE) (Roweis et al., 2000), the consistent clustering patterns (Figure 4) were implicated by projection of genetic maps into spatial coordinates. Moreover, we observed that the

Louvain method performs extremely well to phase groups in coupling. The effectiveness

of the clustering methods indicates the correlation across linkage groups was caused by

real meiotic events.

The loci with $5\% \leq MAF < 15\%$, deviated significantly from the 1:3 expectation for Pt

markers, account for 28.3%, 49% and 48.3% in families "144", "247" and "265"

respectively, in which proportions of the distorted loci correlated (rho $\geq$ 0.3) to the

Mendelian segregation markers ($15\% \leq MAF \leq 35\%$) are 78.3%, 48.9% and 71.8%.

Spatial coordinates of Pt markers, in accordance with correlation heatmaps, offer a

complete picture (Figure 4) of genetic linkage with and without inclusion of SD markers,

which appear to play a role in bridging Mendelian markers. Such linkage patterns were

constantly observed in the three families (Figure 4, S3).

We show the correlation across the 5 largest linkage groups (Figure 5a) in family "265"

based on the spatial representation of the positions of the markers with $15\% \leq MAF \leq$

35%. The model (Figure 5b) clearly depicts distortion decreases, when the markers are

more distal to the convergence areas, and indicates that the progenitors of family "265"

experienced translocation to form a large chromosomal complex.

One linkage group (LG) in one family corresponding to multiple groups in the other

family, would suggest corresponding loci were involved in chromosomal rearrangement

in progenitor of the former family. One striking case (Figure 6) can be found in LG2.1 in

"144" corresponding to two coupling LGs (2.1 and 2.2) in "265". Two control

correspondences (LG1.1-LG1.2 and LG3.1-LG3.1) were used to illustrate the power of

the clustering approaches. However, such one-to-multiple correspondence was seldom

observed across the three families. That may reflect the conservation of chromosomes

positioning in the heterozygotes complex and invariable occurrence of the translocation

heterozygotes in the progenitors of the three families.

GWAS for sex determination

Sexual phenotype regulation is a particularly important problem in dioecious plants,

herein exemplified by hop (*H. lupulus*). Male and female hop flowers can be easily

distinguished; the male flower closely resembles a typical perfect flower, while female

inflorescence meristems produce flowers arranged in 'cones' (Shephard and Parker,

2000).

We used a mixed linear model to assess evidence of phenotype-genotype association. In

families "247" (N = 364, N_{male} = 30) and "265" (N = 95, N_{male} = 13), linkage group (LG)

4 (Figure 7a,S4) consistently shows the most striking association with sex, even though

"265" has a small effective population size. This signal was additionally supported by Fst

mapping in "247" (Figure 7b). However, pseudo-testcross accounts for part of association

signals. To perform genome-wide scan, we assessed association between 356,527

markers and 850 individuals (N_{male} = 129, N_{female} = 721), as described in Methods.

A total of 588 SNPs with $P \leq 10^{-7}$ were identified (Figure 7c,7d). We noted that LG4 and

other LGs respectively account for 38.6% and 0.0% of the association markers,

reinforcing the importance of LG4 in sex expression in hops. On the basis of pairwise

Spearman's rho, we observed high correlation among the 588 SNPs, implicating that the

association markers mostly come from one linkage disequilibrium (LD) block. Adding up

scaffolds showing association approximates ~9.75Mb of the mapping resolution

accounting for ~0.38% of the hop genome.

To identify gene candidates for hop's sex, we aligned by BLAST the top 100 scaffolds

against UniProt and NCBI protein databases to search for genes with known function

involving in the mechanism of sex determination (Table S4). A total of 8 gene candidates

(Ruegger et al., 1998; Ishiguro et al., 2002; Koizuka et al., 2003; Hála et al., 2008;

Shimizu et al., 2008; Zhang et al., 2008; Chen et al., 2010; Zhang et al., 2011) were

obtained (Figure 7d), in which 7 blast hits have 81%-99% length matching to the target

proteins and 7 hits encompass ≥ 1 the significant association site ($p \leq 10^{-25}$).

In hop, a significant deformation of the apical meristem, the producer of flower primordia

cells, has been observed during the transition to the reproductive phase, resulting in the apparent morphological differences between male and female flowers (Shephard and Parker, 2000). Two notable examples, relating to the floral structures, are (1) a glucose-regulated protein 94 (GRP94)-like protein on scaffold LD152823 that is known in Arabidopsis affecting shoot apical meristems, floral meristems and pollen tube elongation (Ishiguro et al., 2002); and (2) a Squamosa-like protein, identified on scaffold LD147778, has essential roles in vegetative phase change and flower development in multiple plants (Chen et al., 2010).

Genetic differences and phenotypic variation across populations

To assess genetic contributions to between-population phenotypic differences, we used Fst analysis to characterize genetic variations across var. *neomexicanus*, var. *lupuloides* and CV. Previously cloned genes in *Humulus* were highlighted to suggest that with respect to essential chemical composition and drought tolerance, hotspots with unusually high or low Fst values deserve a great deal of attention.

A linkage map-based view of Fst highlights two notable patterns (Figure 8). First, the degree of genetic variation, as expected, is much greater in CV vs. *neomexicanus*/*lupuloides* than in *neomexicanus* vs. *lupuloides*. Regions that exhibit above-average population differentiation in CV vs. *neomexicanus* typically also exhibit above-average population differentiation in CV vs. *lupuloides*. Second, the 5 largest linkage groups account for a large proportion of genetic variation between populations. The 90[th] percentile of Fst values (Figure 8d), referred to as 0.63 in CV vs. *neomexicanus* and 0.62 in CV vs. *lupuloides*, were used to define the significant genetic difference.

In beer brewing, the chemical composition, resins imparting bitterness and essential oils contributing to flavor and aroma, determine the quality and flavor of hops. The key brewing resins include alpha-acids and beta-acids, also referred to as humulones and lupulones respectively. It is generally accepted that North American hops have high content of alpha acids, rendering bitter beers, while European hops have lower content of resins and essential oil. Enhancement of alpha acids content in the contemporary cultivars was accomplished by crossing European hops (var. *lupulus*) to North American donors,

one of which resembles var. *lupuloides* from Canada (Neve, 1991; Reeves and Richards,

2011). Genes affecting resins and essential oil can be found in the regions presenting

genetic divergence between North American hops and adapted varieties.

In hop, two homologues of chalcone synthase (*CHS*) in bitter acid biosynthesis have been

cloned (Figure S5), referred to as *CHS_VPS* (GenBank: AB015430.1) (Okada and Ito,

2001) and *CHS_H1* (GenBank: CAC19808.1) (Matoušek et al., 2002) respectively. Both

genes are highly active in lupulin glands, to serve as catalysts of the synthesis reactions of

alpha-acid and beta-acid. Significant genetic differences (Fst $\geq$ 0.6) between cultivars

and North American wild types emerge in a region adjacent to (~40Kb away from)

CHS_H1. In contrast, a surrounding region (~15Kb away from) of *CHS_VPS* harbors

moderate Fst (= ~0.4) in CV vs. *neomexicanus*, and low Fst (= ~0.1) in CV vs. *lupuloides*.

The differences of population differentiation in the surrounding regions of the two *CHS*

homologues may reflect genetic introgression and differential allele selection resulting

from domestication towards higher alpha acid yields.

Var. *neomexicanus* is traditionally grown in rain-fed or supplementary irrigated areas,

and has capability to withstand periods of water deficit and to yield an economic return to

farmers. In contrast, hops frequently occurring in the high latitude of the northern North

America favor plenty of water and sunlight. Likewise, heavy irrigation is required in

modern hop crops. Hence, genic regions exhibiting the significant population

differentiation in CV vs. *neomexicanus*, but not in CV vs. *lupuloides* can be prioritized to

identify gene candidates affecting drought tolerance.

Two intriguing proteins were identified on scaffold "LD161390" (Figure 9a), which are a

H. lupulus basic leucine zipper transcription factor Long Hypocotyl 5 (*bZIP HY5*)

(GenBank: CBY88800.1) (Matousek et al., 2010) and a light-harvesting chlorophyll

(*LHCII*) a/b binding protein (*LHCB*) (NCBI RefSeq: XP_002307004.1).

HY5 has a notable relationship with the phytohormone abscisic acid (ABA), a plant

hormone in response to environmental stress, such as high salinity, drought and low

temperature (Nakagawa H, Ohmiya K, 1996). Via binding and promoting *ABI5*, encoding

a bZIP transcription factor, *HY5* can mediate *ABA* sensitivity (Chen et al., 2008) to induce

stomatal closure, thus reducing transpiration and preventing water loss from leaves.

The putative *LHCB*, physically close to the *HY5*-like gene, embeds a mutation with

striking Fst in CV vs. *neomexicanus*, but not in CV vs. *lupuloides*. *LHCB* involves in the

steps of photosynthesis by capturing sunlight and balancing excitation energy between

photosystems I (PSI) and II (PSII). It is clear that over-expression of a *LHCB* is a key to

enhance stomatal sensitivity to ABA in green alga (*Dunaliella salina*) (Liu and Shen,

2004).

A previously *C. sativa* (marijuana) *LHCII* protein, *CP47* (NCBI RefSeq:

YP_009143579.1), is present on scaffold "LD140230" (Figure 9b), near markers showing

the pronounced population divergence between *neomexicanus* and CV/*lupuloides*. In

sorghum, *CP47* serves as a connection between the main light harvesting complex LHCII

and reaction center of PSII. The characteristic adaptation of sorghum to drought may be

partly related to downregulation of *CP47* during drought stress and high irradiance

(Masojídek et al., 1991).

Discussion

Hop crop acreage and usage is rapidly expanding and diversifying because of a

bourgeoning craft brewing industry. Hop breeding programs have been long attempting to

exploit genetic resources for bitter flavor, aroma and disease resistance. However, a

worsening drought and unseasonably hot weather may decentralize the targets of

breeding programs soon. For example, in Europe and the US, most hop farms

experienced severe water shortage in 2015. Like many other crops, exploitation of novel

genetic variation in response to drought stress is of paramount importance for a

sustainable hop production system.

Understanding genetic recombination is essential for speed and accuracy of plant

breeding. Indeed, it is generally difficult to breed new commercial hop varieties through

mass selection and crossing. Our findings show that a large scale, perhaps genome-wide,

chromosomal rearrangement may occur in the progenitors of F1 families. Translocation

heterozygosity can extend linkage to nonhomologous chromosomes, and favor severe

segregation distortion accumulated near the translocation breakpoints (Taylor and

Ingvarsson, 2003; Farré et al., 2011). Such high degree recombination suppression may

hinder effectively selection of desired, novel allele combinations. Inasmuch, breeding

strategies favoring introgression of diversity deserve re-emphasis in addition to extended

attention to recombination and segregation phenomena in traits. Manipulation of genetic

recombination in hops also deserves further focus.

At least 57 species of flowering plants were characterized by permanent translocation

heterozygotes (Holsinger and Ellstrand, 1984), in which *Clarkia* (2n = 18) may provide a

comparative system to hypothesize meiotic configurations in *Humulus*. Translocation

polymorphism has been observed in at least 14 of the 34 known species in *Clarkia*

(Snow, 1960). Judging from cytogenetic analyses in 9 natural populations of *Clarkia*

dudleyana, none or few translocation heterozygotes occurred within individual colonies,

while extensive translocation heterozygotes invariably arose from hybrids derived from

cytologically distinct races (Snow, 1960). The largest heterozygotes complex consists of

18 chromosomes. For *H. lupulus* wild types, until now, researchers have still been at odds

about the size ratios of sex chromosomes. Except for a tetravalent in var. *cordifolius*, no

meiotic chromosome associations were reported in other wild types, suggesting that the

five wild *H. lupulus* species are cytologically differentiated. Segregation distortion

presenting in cultivated mapping populations may stem from historical hybridization

across isolated populations of hops.

Based on the projection of SNPs into a 3D coordinate system using LLE, the

nondisjunction of the 5 largest LGs in "265" provides an example of funnel-shaped

representation of a heterozygotes complex. Specifically, chromosome segments distal to

and close to translocation breakpoints can be represented by wider and narrower end of

the funnel-shaped structure respectively. Apparently, two ends of the funnel also

characterize clusters of SNPs with similar allele frequencies. There is a need for

cytogenetic studies for drawing conclusions regarding the exact meiotic configuration in

Humulus hybrids for both males and females. Whether the translocation heterozygosity is

sex-linked deserves further investigation.

Translocation heterozygosity may have an important connection to the significantly

distorted sex-ratio in favor of females in hops. Likewise, female-biased sex ratios have

been found in Mistletoe, another notable dioecious case of translocation heterozygosity.

Known that, to maintain heterozygosity, *Oenothera*, a notable monoecious case of

translocation heterozygosity, utilizes a system of balanced lethal to purge the lethal

homozygotes (Steiner, 1956; Harte, 1994), which is referred to as "recessive lethals". In

the context of XY system, heteromorphism of sex chromosomes dictates that males are

more severely affected than females by "X-linked recessive lethals", because males only

have one copy of the X chromosome. Hence, *H. lupulus* may use a system of balanced

lethals at the expense of male offspring to preserve genetic heterozygosity in hybrids.

Our results are compelling for translocation heterozygosity studies in light of

high-density molecular markers in many other biota. For example, such large scale

recombination suppression is also presented in at least 10 species of termite, some types

of centipede, and perhaps all of the monotremes (Holsinger and Ellstrand, 1984; Rowell,

1987; Rens et al., 2004). Beyond homologous crossover, translocation heterozygosity has

shown considerable evolutionary interest and selective advantage in its own right.

Materials and Methods

Plant materials

Hops used in this study were grown under standard agronomic conditions at the Golden

Gate Ranches, S.S. Steiner, Inc, Yakima, WA. The un-domesticated, exotic hops are from

the National Clonal Germplasm Repository in Corvallis, Oregon (accession details in

Table S1-S3). Fifty milligrams of young leaf tissues were extracted in a 96 well block

using Qiagen Plant DNeasy Kits and was tested for quality, quantity, and purity, prior to

library preparations, using a Agilent 2100 Bioanalyzer (Applied Biosystems, Foster City,

CA) and Life Technologies (Carlsbad, CA) Qubit 3.0 Fluorometer. The GBS libraries

were prepared using the *Ape*K1 enzyme according to Elshire, et al. (Elshire et al., 2011).

Pools of 96 accessions were sequenced on one lane of an Illumina HighSeq 2000

(Illumina, San Diego, CA)

SNP calling and quality control

The reference sequence refers to a draft haploid genome sequence of Shinshu Wase (SW)

(Natsume et al., 2015), which is a modern cultivar bred from a seeding selection cross

between Saazer and White Vine-OP**.** The draft genome, with a total size of 2.05 Gb,

consists of ~130,000 scaffolds covering approximately 80% of the estimated genome size

of hop (2.57 Gb).

Tassel 5 GBS v2 Pipeline (Glaubitz et al., 2014) was applied to identify tags with at least

10x total coverage, and to call SNPs. Tag sequences were mapped to the reference

genome using BWA aligner.

One main source of erroneous SNP calling is misalignment caused by incomplete

reference genome, gene duplication and low-complexity regions. To filter out erroneous

SNPs due to misalignment, we used two criteria: (1) SNPs with an excessive coverage

(e.g. read depth > 127) can be false positives. For GBS data, the maximum read depth for

one genotype is unlikely to exceed 127 (Glaubitz et al., 2014). Indeed, we observed that

heterozygosity rates and MAF are significantly increased when read coverage exceeds

127 (Figure S1). (2) The orientation of paired reads of the cultivar Apollo (unpublished

data), a highly used maternal line in our F1 families, was used to detect false positive

SNPs caused by gene duplications. Paired-end alignment was generated by BWA Sampe.

Identification of correctly aligned regions was based on SAM flags indicating reads

mapped in proper pairs. Using criteria (2) was able to detect ~73% SNPs with the

excessive coverage.

Pseudo-testcross

Three F1 families were used to conduct pseudo-testcross (Pt) recombination mappings,

including (1) "144" (N = 179) derived from a cross between Nugget (maternal line) and

Male50 (paternal line); (2) "247" (N = 364) derived from two parental lines, Super

Galena and Male15; (3) "265" (N = 95) derived from a cross between Chinook and

Male57. Using markers heterozygous in the maternal line and null in the paternal line,

three genetic map sets were constructed, consisting of 3551 SNPs for "144", 2369 SNPs

for "247" and 4506 SNPs for "265".

Our analyses followed the main steps in HetMappS pipelines (Hyma et al., 2015).

Specifically, (1) to remove contaminants, identity by state (IBS) based distance matrices

calculated by TASSEL (Bradbury et al., 2007) were used to identify outliers for each

family; (2) SNPs having both parental genotypes (e.g. AA×Aa) with read depth ≥ 4 were

retained for the next step; (3) in progeny, SNPs with average read depth ≥ 4 and with site

coverage $\geq 50\%$ were retained for the next step; (4) to eliminate the effect of

under-calling heterozygotes and sequencing errors, we masked progeny genotypes with

depth=1, and converted genotypes aa to Aa because genotype aa cannot exist for parental

genotypes AA×Aa in Pt; (5) after correction, SNPs with $15\% \leq MAF \leq 35\%$ were

selected to create linkage groups, and SNPs with $5\% \leq MAF < 15\%$ were deemed the

pronounced SD markers; (6) to cluster and order markers, an adjacency matrix with

Spearman's correlation (rho) were derived from the remaining SNPs; (7) on the basis of

absolute values of rho, the Louvain method (Blondel et al., 2008) implemented in

NetworkX (http://networkx.github.io/) was applied to detect communities (clusters). The

Louvain method is an efficient algorithm for community detection in large networks. A

similar method, modulated modularity clustering (MMC) (Stone and Ayroles, 2009), has

been successfully applied to construct linkage groups; (8) to identify coupling phase from

each "absolute rho" cluster, negative values of rho were set to zero, and the Louvain

method was applied to positive values of rho (Hyma et al., 2015); (9) MSTmap (Wu et

al., 2008) was used to provide a sub-optimal solution of genetic ordering within each

linkage group.

Putative 10×2 linkage groups in coupling were obtained in each F1 family. Linkage

groups may not represent one chromosome due to pseudo-linkage resulting from

chromosomal rearrangement, as discussed in Results.

Genome-wide association studies (GWAS)

An association population includes 850 individuals, in which 837 (116 males and 721 females) are progeny in 6 F1 families and 13 are paternal lines. Male and female were encoded as '1' and '0' individually. A total of 356,527 SNPs with coverage $\geq$ 50% and MAF $\geq$ 5% were retained. The Mixed Linear Model (MLM) (Bradbury et al., 2007; Lipka et al., 2012) was used to assess genotype-phenotype association. The Bonferroni method was used to adjust the significance cutoff for an overall probability of 0.05 for type I error.

Additional files

Supplementary Figures. The file contains supplementary figure S1-S5.

Supplementary Tables. The file contains supplementary tables. (**Table S1** Pedigrees of genotyped F1 populations. **Table S2** Cultivar and landrace accessions. **Table S3** Wild exotic accessions. **Table S4** BLASTX hits for scaffolds encompassing sex association (P $\leq$ 10^{-10}) SNPs. Eight strongly supported gene candidates are highlighted.).

Acknowledgments

We thank Buckler lab and Qi Sun's group at Cornell for helpful discussions. We thank the growers at Golden Gate ranches for cultivation of experimental plants.

Literature Cited

Bakel H V, Stout JM, Cote AG, Tallon CM, Sharpe AG, Hughes TR, Page JE (2011) The draft genome and transcriptome of Cannabis sativa The draft genome and transcriptome of Cannabis sativa. Genome Biol 12: R102

Blondel VD, Guillaume J-L, Lambiotte R, Lefebvre E (2008) Fast unfolding of communities in large networks. J Stat Mech Theory Exp 10008: 6

Bradbury PJ, Zhang Z, Kroon DE, Casstevens TM, Ramdoss Y, Buckler ES (2007) TASSEL: software for association mapping of complex traits in diverse samples. Bioinformatics 23: 2633–5

Bradshaw JE (2016) Use of Sexual Reproduction in Base Broadening and Introgression. Plant Breed. Past, Present Futur. Springer. pp 483–527

Buckler ES, Phelps-Durr TL, Buckler CS, Dawe RK, Doebley JF, Holtsford TP (1999)

Meiotic drive of chromosomal knobs reshaped the maize genome. Genetics 153: 415–26

Castañeda-Álvarez NP, Khoury CK, Achicanoy HA, Bernau V, Dempewolf H, Eastwood RJ, Guarino L, Harker RH, Jarvis A, Maxted N, et al (2016) Global conservation priorities for crop wild relatives. Nat Plants 2: 1–6

Chen H, Zhang J, Neff MM, Hong S-W, Zhang H, Deng X-W, Xiong L (2008) Integration of light and abscisic acid signaling during seed germination and early seedling development. Proc Natl Acad Sci U S A 105: 4495–4500

Chen X, Zhang Z, Liu D, Zhang K, Li A, Mao L (2010) SQUAMOSA promoter-binding protein-like transcription factors: Star players for plant growth and development. J Integr Plant Biol 52: 946–951

Elshire RJ, Glaubitz JC, Sun Q, Poland JA, Kawamoto K, Buckler ES, Mitchell SE (2011) A robust, simple genotyping-by-sequencing (GBS) approach for high diversity species. PLoS One 6: 1–10

Farré A, Benito IL, Cistué L, de Jong JH, Romagosa I, Jansen J (2011) Linkage map construction involving a reciprocal translocation. Theor Appl Genet 122: 1029–1037

Fontana PG, Vickery VR (1973) Segregation-distortion in the B-chromosome system of Tettigidea lateralis (say) (Orthoptera: Tetrigidae). Chromosoma 43: 75–98

Glaubitz JC, Casstevens TM, Lu F, Harriman J, Elshire RJ, Sun Q, Buckler ES (2014) TASSEL-GBS: A high capacity genotyping by sequencing analysis pipeline. PLoS One. doi: 10.1371/journal.pone.0090346

Golczyk H, Massouh A, Greiner S (2014) Translocations of chromosome end-segments and facultative heterochromatin promote meiotic ring formation in evening primroses. Plant Cell 26: 1280–93

Gruber K (2016) Re-igniting the green revolution with wild crops. Nat Plants 2: 1–4

Hála M, Cole R, Synek L, Drdová E, Pecenková T, Nordheim A, Lamkemeyer T, Madlung J, Hochholdinger F, Fowler JE, et al (2008) An exocyst complex functions in plant cell growth in Arabidopsis and tobacco. Plant Cell 20: 1330–1345

Harte C (1994) Oenothera Contributions of a Plant to Biology. Monogr Theor Appl Genet. doi: 10.1017/CBO9781107415324.004

Henning JA, Gent DH, Twomey MC, Townsend MS, Pitra NJ, Matthews PD (2015) Precision QTL mapping of downy mildew resistance in hop (Humulus lupulus L .). doi: 10.1007/s10681-015-1356-9

Hill ST, Coggins J, Liston A, Hendrix D, Henning JA (2016) Genomics of the hop
pseudo-autosomal regions. Euphytica. doi: 10.1007/s10681-016-1655-9

Holsinger KE, Ellstrand NC (1984) The Evolution and Ecology of Permanent
Translocation Heterozygotes. Am Nat 124: 48–71

Hyma KE, Barba P, Wang M, Londo JP, Acharya CB, Mitchell SE, Sun Q, Reisch B,
Cadle-Davidson L (2015) Heterozygous Mapping Strategy (HetMappS) for High
Resolution Genotyping-By-Sequencing Markers: A Case Study in Grapevine. PLoS
One. doi: 10.1371/journal.pone.0134880

Ishiguro S, Watanabe Y, Ito N, Nonaka H, Takeda N, Sakai T, Kanaya H, Okada K (2002)
SHEPHERD is the Arabidopsis GRP94 responsible for the formation of functional
CLAVATA proteins. EMBO J 21: 898–908

Jacobsen P (1957) The sex chromosomes in Humulus. Hereditas 43: 357–370

Jakse J, Stajner N, Kozjak P, Cerenak A, Javornik B (2008) Trinucleotide microsatellite
repeat is tightly linked to male sex in hop (Humulus lupulus L.). Mol Breed 21:
139–148

Koizuka N, Imai R, Fujimoto H, Hayakawa T, Kimura Y, Kohno-Murase J, Sakai T,
Kawasaki S, Imamura J (2003) Genetic characterization of a pentatricopeptide
repeat protein gene, orf687, that restores fertility in the cytoplasmic male-sterile
Kosena radish. Plant J 34: 407–415

Laursen L (2015) Botany: The cultivation of weed. Nature 525: S4–S5

Lipka AE, Tian F, Wang Q, Peiffer J, Li M, Bradbury PJ, Gore MA, Buckler ES, Zhang
Z (2012) GAPIT: genome association and prediction integrated tool. Bioinformatics
28: 2397–9

Liu X-D, Shen Y-G (2004) NaCl-induced phosphorylation of light harvesting chlorophyll
a/b proteins in thylakoid membranes from the halotolerant green alga, Dunaliella
salina. FEBS Lett 569: 337–40

Lyttle TW (1993) Cheaters sometimes prosper: distortion of mendelian segregation by
meiotic drive. Trends Genet 9: 205–210

Maaten L Van Der, Hinton G (2008) Visualizing Data using t-SNE. J Mach Learn Res 9:
2579–2605

Masojídek J, Trivedi S, Halshaw L, Alexiou A, Hall DO (1991) The synergistic effect of
drought and light stresses in sorghum and pearl millet. Plant Physiol 96: 198–207

Matousek J, Kocábek T, Patzak J, Stehlík J, Füssy Z, Krofta K, Heyerick A, Roldán-Ruiz

I, Maloukh L, De Keukeleire D (2010) Cloning and molecular analysis of HlbZip1
and HlbZip2 transcription factors putatively involved in the regulation of the lupulin
metabolome in hop (Humulus lupulus L.). J Agric Food Chem 58: 902–12

Matoušek J, Novák P, Bříza J, Patzak J, Niedermeierová H (2002) Cloning and
characterisation of chs-specific DNA and cDNA sequences from hop (*Humulus
lupulus* L.). Plant Sci 162: 1007–1018

Matthews PD, Coles MC, Pitra NJ (2013) Next Generation Sequencing for a Plant of
Great Tradition : Application of NGS to SNP Detection and Validation in Hops
(*Humulus lupulus* L .). BrewingScience 66: 185–191

McAdam EL, Freeman JS, Whittock SP, Buck EJ, Jakse J, Cerenak A, Javornik B, Kilian
A, Wang C-H, Andersen D, et al (2013) Quantitative trait loci in hop (Humulus
lupulus L.) reveal complex genetic architecture underlying variation in sex, yield
and cone chemistry. BMC Genomics 14: 360

Miranda CL, Elias VD, Hay JJ, Choi J, Reed RL, Stevens JF. Xanthohumol improves
dysfunctional glucose and lipid metabolism in diet-induced obese C57BL/6J mice.
Archives of biochemistry and biophysics. 2016 Jun 1;599:22-30.

Murakami A, Darby P, Javornik B, Seigner E, Lutz A, Svoboda P (2006) Molecular
phylogeny of wild Hops , *Humulus lupulus* L . Heredity (Edinb) 97: 66–74

Nagel J, Culley LK, Lu Y, Liu E, Matthews PD, Stevens JF, Page JE (2008) EST
Analysis of Hop Glandular Trichomes Identifies an O-Methyltransferase That
Catalyzes the Biosynthesis of Xanthohumol[W][OA]. Plant Cell 20: 186–200

Nakagawa H, Ohmiya K HT (1996) A rice bZIP protein, designated OSBZ8, is rapidly
induced by abscisic acid. Plant J 217–227

Natsume S, Takagi H, Shiraishi a., Murata J, Toyonaga H, Patzak J, Takagi M,
Yaegashi H, Uemura a., Mitsuoka C, et al (2014) The Draft Genome of Hop
(Humulus lupulus), an Essence for Brewing. Plant Cell Physiol 0: 1–14

Neve RA (1958) Sex Chromosomes in the Hop Humulus lupulus. Nature 181: 1084 –
1085

Neve RA (1991) Hops. London Chapman Hall

Okada Y, Ito K (2001) Cloning and analysis of valerophenone synthase gene expressed
specifically in lupulin gland of hop (Humulus lupulus L.). Biosci Biotechnol
Biochem 65: 150–155

Ososki AL, Kennelly EJ (2003) Phytoestrogens: a review of the present state of research.
Phyther Res 17: 845–869

Rauwolf U, Golczyk H, Meurer J, Herrmann RG, Greiner S (2008) Molecular Marker
Systems for Oenothera Genetics. Genetics 180: 1289–1306

Reeves PA, Richards CM (2011) Species Delimitation under the General Lineage
Concept: An Empirical Example Using Wild North American Hops (Cannabaceae:
Humulus lupulus). Syst Biol 60: 45–59

Rens W, Grützner F, O'brien PCM, Fairclough H, Graves JAM, Ferguson-Smith MA
(2004) Resolution and evolution of the duck-billed platypus karyotype with an
X1Y1X2Y2X3Y3X4Y4X5Y5 male sex chromosome constitution. Proc Natl Acad
Sci U S A 101: 16257–16261

Roweis ST, Saul LK, Roweis ST (2000) Nonlinear Dimensionality Reduction by Locally
Linear Embedding. Science 290: 2323–2326

Rowell DM (1987) Complex sex-linked translocation heterozygosity: Its genetics and
biological significance. Trends Ecol Evol 2: 242–246

Ruegger M, Dewey E, Gray WM, Hobbie L, Turner J, Estelle M (1998) The TIR1 protein
of Arabidopsis functions in auxin response and is related to human SKP2 and yeast
Grr1p. Genes Dev 12: 198–207

Seefelder S, Ehrmaier H, Schweizer G, Seigner E (2000) Male and female genetic linkage
map of hops Humulus lupulus. Plant Breed 119: 249–255

Shephard H, Parker J (2000) Sexual development and sex chromosomes in hop. New
Phytol 148: 397–411

Shimizu KK, Ito T, Ishiguro S, Okada K (2008) MAA3 (MAGATAMA3) helicase gene
is required for female gametophyte development and pollen tube guidance in
Arabidopsis thaliana. Plant Cell Physiol 49: 1478–1483

Silver LM (1993) The peculiar journey of a selfish chromosome: mouse t haplotypes and
meiotic drive. Trends Genet 9: 250–254

Sinotô Y (1929) Chromosome Studies in Some Dioecious Plants, with Special Reference
to the Allosomes. Cytologia (Tokyo) 1: 109–191

Siragusa GR, Haas GJ, Matthews PD, Smith RJ, Buhr RJ, Dale NM, Wise MG. (2008)
Antimicrobial activity of lupulone against Clostridium perfringens in the chicken
intestinal tract jejunum and caecum. Journal of antimicrobial chemotherapy. Apr
1;61(4):853-8.

Snow R (1960) Chromosomal Differentiation in Clarkia dudleyana. Am J Bot 47:
302–309

Steiner E (1956) New aspects of the balanced lethal mechanism in oenothera. Genetics

Stevens JF, Page JE (2004) Xanthohumol and related prenylflavonoids from hops and
beer: To your good health! Phytochemistry 65: 1317–1330

Stone E a., Ayroles JF (2009) Modulated modularity clustering as an exploratory tool for
functional genomic inference. PLoS Genet. doi: 10.1371/journal.pgen.1000479

Tanksley SD, McCouch SR (1997) Seed banks and molecular maps: unlocking genetic
potential from the wild. Science 277: 1063–1066

Taylor DR, Ingvarsson PK (2003) Common features of segregation distortion in plants
and animals. Genetica 117: 27–35

Wiens D, Barlow BA (1975) Permanent Translocation Heterozygosity and Sex
Determination in East African Mistletoes. Science (80-) 187: 1208–1209

Winge O (1932) The Nature of Sex Chromosomes. Proc Sixth Int Congr Genet 343–355

Wu Y, Bhat PR, Close TJ, Lonardi S (2008) Efficient and accurate construction of
genetic linkage maps from the minimum spanning tree of a graph. PLoS Genet. doi:
10.1371/journal.pgen.1000212

Zhang Y, Feng S, Chen F, Chen H, Wang J, McCall C, Xiong Y, Deng XW (2008)
Arabidopsis DDB1-CUL4 ASSOCIATED FACTOR1 forms a nuclear E3 ubiquitin
ligase with DDB1 and CUL4 that is involved in multiple plant developmental
processes. Plant Cell 20: 1437–1455

Zhang Z, Zhang Y, Tan H, Wang Y, Li G, Liang W, Yuan Z, Hu J, Ren H, Zhang D
(2011) RICE MORPHOLOGY DETERMINANT encodes the type II formin FH5 and
regulates rice morphogenesis. Plant Cell 23: 681–700

Figure Legends

Figure 1 Population structure of 251 hop accessions and geographic origins of the U.S. wild types. 183 modern cultivars are indicated by red color. 68 wild hops are color-coded by geographic origins. (a) Neighbor-joining tree of the 251 hop accessions. (b) The state names are followed by sample counts. Three state groups ("MT, ND, SD, NE, IA, KS, MO", "CO, AZ, NM" and "MA") are color-coded to distinguish from one another.

Figure 2 Population structure and pedigree network of F1 families. (a) t-SNE plot for 20 F1 families (N $\geq$ 60). (b) The overview of pedigree for genotyped F1 families.

Figure 3 Linkage groups for the maternal line of family "144" and correspondence across 3 genetic map sets. The degrees of Spearman's correlation (rho) are color-coded. (a) Unphased and phased (linkage for grandparents) groups are bounded by white and black frames individually. (b) Alignment of unphased groups between "144" and "247". (c) Alignment of unphased groups between "144" and "265".

Figure 4 Linkage of Mendelian (15% $\leq$ MAF $\leq$ 30%) and non-Mendelian Pt markers (5% $\leq$ MAF < 15%), based on Spearman's correlation (rho). In each sub-figure, clustering patterns without (left) and with (right) inclusion of segregation distortion are presented by LLE (top) and the Louvain Modularity (bottom). Mendelian markers in two linkage groups are indicated by blue and red colors individually. Segregation distortion (SD) markers are indicated by yellow color.

Figure 5 Linkage patterns of the 5 largest linkage groups in family "265", based on spatial coordinates defined by LLE. (a) Linkage groups are color-coded. (b) Markers with 0.15 $\leq$ MAF < 0.2 and 0.2 $\leq$ MAF $\leq$ 0.3 are distinguished by cyan and gray colors.

Figure 6 One-to-two genetic correspondence between "144" and "265". LG2.1 in "144" corresponds to LG2.1 and LG2.2 in "265" (top sub-figure). Two instances of one-to-one correspondence (LG1.1-LG1.2 and LG3.1-LG3.1) are added for control. Spatial representations of linkage groups (bottom sub-figure) in the two families were derived from LLE.

Figure 7 Association studies and Fst mapping of sex determination in hop. (a) Linkage group-based Manhattan-plot of MLM for sex determination in family "247" (N = 364,

N_{male} = 30). Light and deep colors are used to distinguish two phases (linkage for

grandparents) in coupling. (b) Manhattan-plot of Fst in females vs. males in "247". (c)

Log Quantile-Quantile (QQ) plot of 356,526 association tests (SNPs) for sex

determination in 850 individuals (N_{male} = 129, N_{female} = 721). (d) Correlation among 588

association (P ≤ 10^{-7}) markers, the proportions of 588 markers in LG4, other LGs and

unmapped data set, and 8 gene candidates for sex determination in hop.

Figure 8 Linkage group (in family "247")-based Fst heatmaps and the overall Fst

distribution. Population differentiation (a) between modern cultivars (CV) and var.

neomexicanus; (b) between CV and var. *lupuloides*; (c) between var. *neomexicanus* and

var. *lupuloides*. (d) Spectrum of the overall Fst distribution.

Figure 9 Three gene candidates for drought tolerance in hops. Between-population Fst

values are indicated on the scaffolds where the three candidates are located. (a) The

approximate positions of *bZIP HY5* (GenBank: CBY88800.1) (Matousek et al., 2010) and

LHCB (NCBI RefSeq: XP_002307004.1) on scaffold "LD161390". (b) The approximate

position of *CP47* (NCBI RefSeq: YP_009143579.1) on scaffold "LD140230"

a

b

● Canada	● United States (MA)	● F1 parents	● Kazakhstan
● United States (MT,ND,SD,NE,IA,KS,MO)	● Unknown	● United States (CO,AZ,NM)	

a

Population structure of crosses

Coord 2 (t-SNE) vs Coord 1 (t-SNE)

Legend:
- 265
- 264
- TeaMaker x USDA_21422M
- 241
- 149
- Zeus_1
- 106
- Cascade
- Chinook
- DG
- Nugget
- 144
- DF
- Zeus_2
- 248
- 255
- 247
- 249
- 147
- 145

b

TeaMaker x USDA_21422M

255
241
247 248
149 145
106 144 147 249
264
Chinook
265
DG   DF

Legend:
- Mother
- Father
- Progeny

a LGs 1.1 and 4.1 (F1: 144 maternal LGs)

b LGs 2.1 and 8.1 (F1: 247 maternal LGs)

c LGs 10.1 and 10.2 (F1: 265 maternal LGs)

d LGs 2.1 and 2.2 (F1: 265 maternal LGs)

🟡	3.1	⚫ 9.1
🔴	10.1	🟠 4.1
🔵	2.1	

⚪	$0.2 \leq MAF \leq 0.3$
🔵	$0.15 \leq MAF < 0.2$

a

bZIP HY5   LHCB

LD161390

b

LHCII CP47

LD140230

CV vs. neomexicanus
CV vs. lupuloides