Premature termination codons signal targeted gene repair by nonsense-mediated gene editing in *E. coli*

Xiaolong Wang[1][*], Xuxiang Wang[1], Haibo Peng[1], Chunyan Li[1], Yalei Wang[1], Gang Chen[1], Jianye Zhang[1]

College of Life Sciences, Ocean University of China, Qingdao, 266003, P. R. China

[*] To whom correspondence should be addressed: Xiaolong Wang, Department of Biotechnology, Ocean University of China, No. 5 Yushan Road, Qingdao, 266003, Shandong, P. R. China, Tel: 0086-139-6969-3150, E-mail: Xiaolong@ouc.edu.cn. .

Abstract

Frameshift mutations yield truncated proteins, leading to loss-of-function, genetic disorders or even death. Reverse mutations, which restore the wild-type phenotype of a mutant, have been assumed to be far rarer than a forward mutation. However, in this study, screening tests in *E. coli* showed that the revertants of a frameshift mutation are detected more frequently than expected. Sanger and NGS sequencing of the plasmids revealed that the reversion of the frameshift mutation is not likely caused by random mutagenesis but by an active and targeted gene repair. Molecular studies suggest that premature termination codons (PTCs) in the nonsense mRNAs signal the repair of the frameshift mutation. A genome resequencing indicates that the genome sequence of the revertant is stable, suggesting that the proofreading/mismatch repair system is not defective but active in the revertant. Transcriptome profiling identified upregulated genes/pathways probably involved in the repair of the frameshift mutation, including DNA replication, RNA processing, RNA editing, mismatch repair, and homologous recombination. Based on these data and an in-depth review of previous studies, we hypothized a model for frameshift repair referred to as *nonsense-mediated gene editing* (*NMGE*): nonsense mRNAs are recognized by PTC signaling, edited by RNA editing and then used to direct the repair of their defective coding gene through mismatch repair and homologous recombination. Moreover, NMGE may also serve as a driving force for molecular evolution and a new source of genetic diversity leads to the widespread existence of frameshift homologs.

1. Introduction

DNA replication and DNA repair happen daily in cell division and multiplication. DNA damage, point mutations, insertions and deletions (indels) can be induced by chemical and physical mutagens, such as radiation, pollutants, or toxins. In addition, because of the imperfect nature of DNA replication, mutations and indels also occur spontaneously as replication errors or slipped-strand mispairing. In a protein-coding gene, if the size of an indel is not a multiple-of-three, it causes a frameshift mutation, which results in a substantial change of its encoded amino acid sequence; in addition, a number of nonsense codons are often produced downstream the frameshift mutation, and thus, it has been generally believed that a frameshifted gene yields truncated and dysfunctional proteins [1], and therefore lead to genetic disorders or even death [2-5].

The molecular mechanisms/pathways involved in the repair of DNA damages and point mutations have been intensively studied [6], including mismatch repair, base- or nucleotide-excision, homologous recombination, and non-homologous end-joining. But no molecular mechanism has been reported for the repair of frameshift mutations (hereafter *frameshift repair*). The maintenance of the reading frames of coding genes is fundamentally important for cells. A reverse mutation phenomenon was discovered as early as in the 1960s [7], however, both forward and backward mutations have been explained by random mutagenesis. In principle, the rate of a reverse mutation should be much lower than that of a corresponding forward mutation. However, here we found that the reversion of a frameshift mutation occurs far more frequently than that was expected in *E. coli* and it seems that it is not random mutagenesis but involves an active and targeted gene repair.

2. Materials and Methods

2.1 Frameshift mutagenesis and back mutation screening

A plasmid containing a frameshifted β-lactamase gene (*bla*), pBR322-(bla-), was constructed by site-directed mutagenesis (Fig 1): in plasmid pBR322, a G:C base pair, located in the upstream (+136) of the wild-type *bla* gene (*bla+*), was deleted by using overlapping extension polymerase chain reaction (OE-PCR). Competent cells of

Escherichia coli strain DH5α were transformed with pBR322-(bla-), propagated in tetracycline broth (TCB), and then diluted cultures were plated on a tetracycline plate (TCP) and ampicillin plates (ACPs) to screen for revertants. The recovery rates were calculated by $r = N_1/N_0$, where N_0 is the total number of bacteria and N_1 is the number of revertants. The revertants were propagated in ampicillin broth (ACB) at 37°C with 200 rpm shaking and with 1 mL of overnight seed culture in 10 mL ACB. The growth rates of the revertants were evaluated by the doubling time during their exponential growth phase. Their plasmids DNA were extracted and their *bla* genes were Sanger sequenced.

2.2 Construction and expression of a PTC-free frameshifted bla

A *PTC-free frameshifted bla*, denoted as *bla#*, was derived from *bla-* by replacing each nonsense codon with a sense codon according to the *readthrough rules* (Table 1), and by adding a stop codon (TAA) in the 3'-end. The *bla#* gene was chemically synthesized by *Sangon Biotech, Co. Ltd* (*Shanghai*), inserted into an expression vector, pET28a, and transformed into competent cells of *E. coli* strain BL21. Plasmid pET28a contains a kanamycin resistance gene, the transformed bacteria were therefore plated on a kanamycin plate (KCP), the transformants were propagated in kanamycin broth (KCB), plated on ACPs to screen for revertants, and then their *bla* genes were Sanger sequenced. The expression of the frameshifted BLA was induced by 0.1 mM IPTG. Their total protein samples were extracted, frameshifted BLA was purified by nickel column chromatography and analyzed by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE). The activity of BLA was measured by iodometry.

2.3 Genome resequencing and structure/variation analysis of the E. coli strains

Genomic DNA was extracted from wild-type (*bla+*) and revertant (*bla**) samples. Library preparation and genome sequencing were conducted by a commercial service provided by *Novogene Co. Ltd*. Paired-end reads were obtained on an Illumina HiSeq 250PE platform. Clean reads were obtained by removing adapters and low-quality sequences. For each sample, clean reads were mapped onto the reference genome of *E. coli* K12 MG1655 (NC_000913.3) and the plasmid pBR322, to identify single nucleotide polymorphisms (SNPs), indels and structure variations (SVs). Samtools

(v0.1.18) was used to sort and reorder the bam alignments [8]. Genome analysis tool kit (GATK) was used for SNP calling [9]. Circos was used to display the coverage of reads and the distribution of SNPs and indels on the ring diagram [10]. BreakDancer was used to detect structure variations [11, 12].

2.4 Transcriptome analysis of the E. coli strains

Total RNAs were extracted from a wild type (*bla+*), a frameshift (*bla-*) and two revertants (*bla**) samples. Library preparation and RNA sequencing were conducted by a commercial service (*Novogene Co. Ltd*). For each sample, the high-quality clean reads were mapped onto the annotated reference genome of *E. coli* K12 MG1655 (NC_000913.3) and plasmid pBR322, and the expression levels of each gene were calculated and compared to each other to identify differentially expressed genes (DEGs), enriched gene ontology (GO) terms and Kyoto encyclopedia of genes and genomes (KEGG) pathways.

(1) *Quantification of gene expression level*

Bowtie (v2-2.2.3) was used for index building and aligning the clean reads to the reference genome [13]. HTSeq (v0.6.1) was used to count the numbers of clean reads mapped to each gene [14]. And then the expected number of Fragments PerKilobase of transcript sequence per million base pairs sequenced (FPKM) of each gene was calculated based on their lengths and read counts.

(2) *Differential expression analysis*

Prior to differential expression gene analysis, the read counts were adjusted by the edgeR program for each strain [15]. Differential expression of the read counts of two conditions was performed using the DEGSeq R package (1.20.0) [16]. The P values were adjusted by the Benjamini & Hochberg method. The threshold for significantly differential expression was set as: corrected P-value (q-value) < 0.005 and log2 (fold change) $\geqslant$ 1.

(3) *GO and KEGG enrichment analysis of differentially expressed genes*

Gene Ontology (GO) enrichment analysis was implemented by the GOseq R package [17], in which the potential gene length bias was corrected. GO terms with corrected P-value less than 0.05 were considered significantly enriched by DEGs.

KOBAS 2.0 [18] was used to test the statistical enrichment of DEGs in KEGG pathways (http://www.genome.jp/kegg/).

3. Results and Analysis

3.1 The growth of the frameshift mutant and the revertants

The plasmid pBR322 contains two wild-type resistance genes, a β-lactamase gene (*bla+*) and a tetracycline resistance gene (*tet+*). Using OE-PCR, one single G:C base pair in the upstream of *bla+* was deleted (Fig 1). In the resulting frameshift *bla* (*bla-*), seventeen nonsense codons appeared. Because the active sites of BLA, including the substrate-binding site, all located downstream the deletion, this frameshift mutation is a loss-of-function. The plasmid containing a wild-type or a frameshifted *bla* gene was transformed into *E. coli* (strain DH5*α* and BL21), the transformants were propagated in tetracycline broth (TCB), dilution series were plated on tetracycline plate (TCPs) to estimate the total number of viable bacteria and plated on ampicillin plates (ACPs) to test their ampicillin resistance. The wild-type (*bla+*) bacterial grew well on both ACP and TCP (Fig 2A, left); and it was expected that the mutant (*bla-*) would not grow on ACP. However, a few ampicillin-resistant colonies (*bla**) were observed (Fig 2A, middle). The possibility of carryover- or cross-contamination was ruled out by blank controls (Fig 2A, right). So, these ampicillin-resistant colonies are considered as revertants whose *bla-* genes were repaired by reverse mutations.

The test was repeated by different operators for over one hunred times and the revertants are found highly reproducible in most tests. In addition, although most of the revertants can grow in ACB, the growth rates of the revertants are often very much slower when compared to the wild-type strain: by culturing at 37°C with 200 rpm shaking, it takes 12 to 24 hrs for the wild type but up to 36 to 48 hrs for the revertants to reach the late log phase, and many of the revertants failed to grow when they were subcultured in ACB, even without adding ampicillin.

Hitherto, it seems that there is nothing unusual because the reverse mutations are a fairly common phenomenon, arising from simple random mutations (Fig 3): in DNA replication, natural or induced mutations occur randomly, a forward mutation causes a

defective mutant or a loss-of-function, and a reverse mutation restores the wild-type phenotype. Under this view, these revertants survived only because they were 'lucky': their defective *bla* gene restored coincidently by a reverse mutation, while most of the other bacteria died without a reverse mutation.

3.2 frameshift repair seems to be not a random reversion but a targeted gene repair

The classical random mutagenesis model for reverse mutation sounds faultless in explaining the reversions of point mutations. However, in a 'thought experiment', we noticed that the model is insufficient to explain the frameshift repair observed herein: (1) in the model, a bacterium survives if its frameshifted *bla* gene is restored through random mutagenesis. However, a reverse mutation must occur in the process of DNA replication and/or repair in a living cell. If the mutant itself cannot live in ampicillin, the reverse mutation must occur not after but before they were cultured in ampicillin; (2) a frameshift mutation can only be repaired by an appropriate insertion or deletion restoring the reading frame of the defective gene. However, even if relatively small, a bacterial genome consists of millions of base pairs and thousands of genes. If the cells do not 'know' which gene is defective, there is no or little chance to repair it through random mutagenesis; (3) As determined by whole-genome sequencing [19], the baseline mutation rate of the wild-type *E. coli* is $\sim 10^{-3}$ per genome (or $\sim 10^{-9}$ per nucleotide) per generation. In *E. coli*, the rate of point mutation is very low, and the frameshift mutation rate is even ten-times lower than that of point mutations [19]. It is assumed that a reverse mutation occurs at a lower rate than a forward mutation. For a certain frameshift mutation, the reverse mutation rate should be extremely low ($<10^{-10}$) if it depends on random mutagenesis, as insertions/deletions are far rarer than base substitutions. Therefore, a frameshift revertant should be very difficult to be detected. However, through repeated screening tests, we found that the revertants of *bla-* can always be detected easily, and it seems that the reversion of a frameshift occurs much more frequently than expected. The average recovery rate of *bla-* measured in DH5α was 2.61×10^{-8} per cell division and reached up to 1.19×10^{-7} per cell division in BL21, both significantly higher than the expectation.

If a reverse mutation is caused by random mutagenesis, it is bound to be inhibited by the proofreading and mismatch repair system (PR-MRS). However, as mentioned above, actually the opposite is true: the reverse rate is not lower but 4-5 times higher in BL21 (which has a functional PR-MRS) than that in DH5α, which has a defective PR-MRS, strongly suggesting that the PR-MRS system does not inhibit but promote the reversion. Therefore, it is highly possible that an efficient mechanism specific to the repair of frameshift mutation has been developed in the history of life. However, how a frameshifted mutation is being identified and repaired at the molecular level? It is yet unknown but the PR-MRS system may play a vital role in the process.

3.3 Sanger sequencing of the bla genes of the revertants

To investigate how the frameshift gene is repaired, the revertants were picked and subcultured in ACB. The initial revertants usually grew very slowly, and many of their subcultures failed to grow, but the surviving subcultures grow faster and faster. Their plasmid DNA was extracted, and their *bla* were Sanger sequenced. Unexpectedly, in these initial revertants and subcultures, the *bla* genes are often neither a wild type nor a revertant but seem to be still frameshifted (Fig 2B, top). It is puzzling how these *bla* frameshifts could be functional.

To clarify this problem, the revertants were subcultured, propagated and their *bla* were re-sequenced for over 300 replications. In the subcultures, the sequencing results of the *bla* genes also often seems to be frameshifted, but their Sanger diagrams often contain two sets of superimposed peaks (Fig 2B, middle), indicating that they consist of two types of *bla* sequences: one is the original frameshifted *bla*, and the other is the repaired one. By visual inspection of the Sanger diagrams, the repaired sequences of the revertants (*bla**) were read from the secondary peaks. Sometimes a sequencing diagram may contain three or more sets of peaks, but the sequences are nonreadable. In the final fast-growing subcultures, most of their *bla* copies were repaired, and their Sanger diagrams contain few superimposed peaks (Fig 2B, bottom).

As shown in Fig 2C, by aligning *bla** sequences against *bla+* and *bla-* sequences, it was found that they contain a variety of indels and substitutions, especially G:C←

→A:T transitions (Fig 2C, circles). Previously, we developed a novel tool for multiple sequence alignment, *codon and aa unified alignment* (CAUSA) [20], which can align and visualize coding sequences and their encoded protein sequences in the same user interface. As shown in Fig 2D, CAUSA demonstrates that the protein sequences translated from the *bla** genes are all different from each other but all highly similar to that of the wild-type BLA, even if they are interrupted by some stop signals. In other words, the frameshifted genes are restored not completely and immediately, but partially and progressively, resulting in a variety of *frameshift homologs*.

3.4 The genome of the revertant is not highly variable but very stable

As is well known, an extraordinarily high rate of spontaneous mutagenesis occurs over its whole genome in a mutator, such as *E. coli* strain mutD5, due to its defective proofreading and mismatch repair during DNA replication [21]. The *E. coli* strain we used in this study, either DH5α or BL21, is not a mutator, but the frameshift/revertants obtained may change into mutators if their proofreading and mismatch repair system is damaged during culturing and subculturing so that high-rate of mutagenesis can occur in their genomes, not only in *bla* but also in other genes.

To clarify whether a revertant may change into a mutator, the genome sequences of a frameshift (*bla*-) and a revertant (*bla**) were sequenced by high-throughput NGS sequencing. The clean reads were mapped onto the reference genome sequence of the *E. coli* strain K12 and compared with each other (Fig 4A). As shown in Fig 4B and Table 2, the SNP/indel levels of the revertant and frameshift tested are both very low, suggesting that their proofreading/mismatch repair system is not defective. Therefore, they are not mutators but adopt a stringent proofreading and mismatch repair during the replication/repair of their DNA. In addition, if the variations observed in *bla** are caused by random mutagenesis, the number of variations observed in *bla* should be proportional to the number of variations observed in the whole genome, and therefore, the number of variations observed in the revertant genome should be higher than that in the frameshift genome. However, as shown in Fig 4B and Table 2, the numbers of SNPs/indels of the revertant are not higher but even lower than those of the frameshift. In other words, the number of *bla* variations are independent of that of the variations

observed in the whole genome, suggesting that the high-level of variations of *bla** are

not caused by nonspecific, whole-genome random mutagenesis, but more likely to be

an active gene repair specifically targeting the frameshifted *bla*.

3.5 The other gene in the same plasmid does not change

To clarify whether tetracycline-resistant (*tet*), the other gene in the same plasmid,

changes or not, it was also Sanger sequenced but no obvious overlapping peaks were

observed in their sequencing diagrams, demonstrating that overlapping peaks oftenly

observed in *bla** are not the result of a nonspecific mutagenesis but more likely to be

an active gene repair targeting specifically to the frameshifted *bla* gene.

To validate that the *tet* gene does not change as fast as the *bla* gene, we further

mapped the NGS reads of the genome or transcriptome onto the plasmid. As shown in

Fig 4D, by visual inspection, very few variations can be observed in the *bla* genes of

the wild type, but a lot of insertion, deletion, and substitutions can be observed in the

bla genes, as well as in their transcripts, in both the frameshift and the revertant; as

shown in Fig 4E, however, almost no or only a few variations were observed in the *tet*

genes or their transcripts, neither in the frameshift nor in the revertant.

The SNPs were called by using bcftools/vcfutils.pl and summarized in Table 3

and Supplementary data 7. Such kind of random variations is hard to distinguish from

sequencing errors, filtering of the raw SNPs may discard the real changes. We

therefore used unfiltered raw SNPs, but discarded those obviously ambiguous ones, to

count and evaluate the numbers and densities of SNPs of the two genes in the plasmid

pbr322. In DNA-level, the density of raw SNPs is very high in both genes. the density

of *bla* variation is slightly higher than that of *tet* in the frameshift; however, is

significantly higher than that of *tet* in the revertant; in RNA level, the number of raw

SNPs is smaller than those in DNA but the difference between *tet* and *bla* is more

obvious: the density of variation is lower in *bla* than in *tet* in the wild type but

significantly higher in *bla* than in *tet* in the revertant,. In a word, DNA- and

RNA-level variations are both higher in *bla* than in *tet*, either in the frameshift or the

revertant but not in the wild type cells, demonstrating that these variations are likely

to be caused by an active gene repair targeting specifically to the frameshifted *bla*.

3.6 PTCs signal the repair of the frameshift mutation

We speculated that it was the PTCs presented in the nonsense mRNAs that signal the repair of frameshift mutations. To validate this, a *PTC-free bla-* (denoted as *bla#*) was derived from *bla-* by replacing each nonsense codon with an appropriate sense codon in accordance with the *in vivo readthrough rules* (Table 1). The *bla#* gene was synthesized chemically, cloned in the plasmid vector pET-28a and expressed in *E. coli* strain BL21. The *bla#* transformants were plated on ACPs, but no revertant was found. Their *bla#* genes were Sanger sequenced but few variation were observed, suggesting that the *bla#* gene is not repaired as efficiently as the *bla-* gene. Analysis of total and purified protein by SDS-PAGE revealed a protein band corresponding to the predicted molecular mass (34-kDa), representing a defective BLA expressed from the PTC-free frameshifted *bla* (Fig 5A). No lactamase activity was detected by iodimetry test of the defective product (Fig 5B), confirming that the frameshift is indeed a loss-of-function when the PTCs are removed. In other words, the *bla#* gene is kept unchanged and expressed a frameshifted, dysfunctional product in its wrong reading frame. The only difference between *bla#* and *bla-* is that: *bla-* consists of a certain number of PTCs but *bla#* does not. Therefore, it must be the PTCs that signaled the repair of the frameshift mutation.

3.7 Identifying genes/proteins involved in frameshift repair

We speculated that frameshift repair requires the upregulation of relevant genes and pathways. By transcriptome sequencing and profiling, as shown in Fig 6A, the gene expression patterns of these strains are significantly different from each other. When compared to the wild-type (*W*), respectively 292 or 247 genes displayed upregulated transcription (Fig 6A, Supplementary dataset 1-2) in the frameshift (*FS*) or revertant (*RE*). In the pathway enrichment analysis, respectively 65 or 58 pathways were identified as upregulated (Supplementary dataset 3-4). Most of the upregulated genes or pathways do not likely to be involved in the frameshift repair but are related to nutrition or ampicillin resistance, probably due to the different growth conditions of these different cells. However, we manually selected the upregulated genes/pathways that most probably involve in frameshift repair. The criterion for selection is whether

the gene/pathway is involved in the processing of DNA or RNA, the storages of the genetic information. As shown in Fig 6B-6C and Table 4-5, the selected upregulated genes/pathways include *DNA replication* (Fig 6D)*, RNA degradation/processing* (Fig 6E)*, mismatch repair* and *homologous recombination* (Fig 6F).

In addition, the selenocysteinyl-tRNA-specific translation factor (*selB*) gene is found upregulated in one of the two revertants tested (Supplementary dataset 5). SelB is a specialized translation factor that takes the place of elongation factor EF-Tu, in the case of selenocysteines being inserted into a peptide chain at the site of a UGA codon [22, 23]. The upregulation of *selB* in the revertant suggested that the nonsense codons remained in the mRNAs might be readthrough in the revertants.

By carefully inspecting the list of differentially expressed genes (Supplementary dataset 6), we found that the expression of the cytosine or isoguanine deaminase gene (*codA*) and the adenosine deaminase gene (*add*) were upregulated in the frameshift, and those of the cytidine/deoxycytidine deaminase gene (*cdd*) and the tRNA-specific adenosine deaminase gene (*tadA*) were also upregulated in the revertant:

(1) In *E. coli*, cytosine or isoguanine deaminase (CodA) catalyzes the hydrolytic deamination of cytosine to uracil, and also the deamination of isoguanine, a mutagenic oxidation product of adenine, and of isocytosine, in DNA [24, 25]. In eukaryotes, cytosine deaminase is involved in RNA editing [26].

(2) In *E. coli*, cytidine/deoxycytidine deaminase (Cdd) scavenges exogenous and endogenous cytidine and 2'-deoxycytidine for uridine monophosphate (UMP) synthesis [27-32]. In eukaryotes, cytidine/deoxycytidine deaminase is also involved in RNA editing [33, 34].

(3) In *E. coli*, adenosine deaminase (Add) is an inducible enzyme that converts adenine, adenosine, and deoxyadenosine to guanine nucleotides [35-37]. In eukaryotes, adenosine deaminase (ADAR2) is an important enzyme that is responsible for RNA editing [38-50].

(4) The tRNA-specific adenosine deaminase (TadA) catalyzes the deamination of adenosine to inosine at the wobble position 34 of tRNA-Arg2. TadA was first identified in yeast [51], which is found widely existing in many bacteria

[52-55]. TadA the first prokaryotic RNA editing enzyme, which is identified in *E. coli* [55]. TadA is related to the eukaryotic RNA editing enzymes, including ADAR1 and ADAR2 [51], which widely exists in many eukaryotic species, such as human [56], mouse [57] and fruit fly [58].

As above mentioned, Sanger sequencing showed that the base substitutions in the repaired *bla* are mostly G:C←→A:T transitions, suggesting that they are derived from the deamination of DNA bases. It is well known that DNA deamination generates deoxyinosine from deoxyadenosine, which links DNA mismatch repair and RNA editing [59]. Here we show that DNA deamination and mismatch repair pathways are upregulated in the frameshift, suggesting that both RNA-editing and mismatch-repair might be involved in frameshift repair. These genes are also upregulated in the revertant, suggesting that gene repair is not ceased but still ongoing in the revertants.

4. Discussion

4.1 A new model for the repair of frameshift mutations

The mechanisms for the repair of DNA damages and point mutations have been well-understood. However, the molecular mechanism for the recognition and repair of frameshift mutations is yet unknown. In the above, the expression of the PTC-free *bla* indicated that PTCs signaled the repair of the frameshift mutation. We postulated that a transcript of the frameshift gene (a nonsense mRNA) must be involved in the repair, because presumably these PTCs can only be recognized in a nonsense mRNA in the process of translation but not likely possible in a DNA double helix.

To gain further insight into this very complex problem, we conducted an in-depth survey of previous studies that covered a wide range of prokaryotic and eukaryotic organisms. By presuming that frameshift repair is widely existing and the mechanism is basically conserved among species, here we integrate plenty of fragmentary pieces of evidence together and propose a hypothetical model for frameshift repair referred to as *nonsense-mediated gene editing* (NMGE), consisting five major steps (Fig 7):

Step 1. *Nonsense mRNA recognition*: in principle, a frameshift coding gene cannot be recognized at the DNA level, and therefore must be transcribed firstly. The

transcripts of the frameshift gene (nonsense mRNAs) are recognized by the RNA surveillance system [60-66]. When a nonsense mRNA is translated, the ribosomes are blocked when they encounter a stop codon [67]. A stop codon emerges inside a frameshift gene or an mRNA is called a premature termination codon (PTC) [68-70], also known as a hidden stop codon (HSC) [1].

Step 2. *Nonsense mRNA processing*: In eukaryotes, there are several nonsense mRNA processing pathways [71-73], in which nonsense-mediated mRNA decay (NMD) has been intensively studied [74-77]. NMD targeting mRNAs harboring PTCs for degradation enhances their decay rate and reduces the abundance of the mRNA transcribed from that gene. In bacteria, it is not yet clear how nonsense mRNAs are processed, but it is known that the quality of their mRNAs is also tightly controlled [78]. Therefore, we speculate that NMD may also exist in bacteria, although it has not yet been characterized. Nevertheless, in both prokaryotes and eukaryotes, nonsense mRNAs must be processed by one of the pathways, either degradation (step 2), translation (step 3), or editing (step 4).

Step 3. *Nonsense mRNA translation*: Nonsense mRNAs are not always fully degraded but sometimes translated. Frameshifted or nonsense-mutated coding genes can be expressed through some special mechanisms, including programmed translational frameshifting [79-86] and translational readthrough [87-90]. Especially, if the target gene is essential, a cell survives only if it produces functional proteins by translating a nonsense transcript/mRNA and dies if no protein is produced or the products are truncated or nonfunctional.

Step 4. *RNA editing*: nonsense mRNAs must be involved in the recognition and repair of the frameshift mutation, but a nonsense mRNA is defective by itself, and thus, cannot template the repair of its coding gene directly, therefore, it is likely that a nonsense mRNA must be modified by RNA editing prior to templating the repair. Nonsense mRNAs are recognized by PTC signaling, but it ought to be impossible for cells to locate the frameshift-causing indel exactly, either in a genomic DNA or a transcript, and therefore, often it is not possible to restore the original coding sequence. Therefore, the cells have to insert, delete and substitute nucleotide(s) in

the target mRNA in or near to a PTC randomly, resulting in a variety of variants.

After editing, the translation is restored and a functional BLA is produced if the

mRNA is repaired.

Step 5. *RNA-directed DNA repair*: to repair the frameshifted coding gene, mRNAs are

transported back to chromosomal DNA to localize its defective coding sequence.

the defective coding gene is repaired by using an edited mRNA as the template. A

bacterium recovers if functional mRNAs and proteins are produced by a normal

transcription and translation process.

As abovementioned, substantial evidence suggests a direct link between RNA

editing and DNA repair, which can also support this NMGE model. However, further

systematic experimental investigations are needed to elucidate this process in detail

and validate it in other species, especially the eukaryotes.

4.2 ***The NMGE model better explains the variations of the bla genes***

As shown by a question mark in Fig 7, there exist two possible strategies: (1)

direct NMGE: using an original nonsense mRNA to identify the defective coding

sequence, and then editing it by DNA-level mutagenesis; however, whether or not the

edited CDS is repaired, it must be transcribed and translated again to be functional; (2)

indirect NMGE: the nonsense mRNAs are first edited, and then transported back to

DNA to identify the CDS and to template the gene repair, *i.e.*, a nonsense mRNA is

functionalized through RNA editing prior to directing the repair of its own CDS.

Compared to the direct strategy, indirect NMGE seems more reasonable, because the

repair of the coding DNA sequences happens only in a living cell which has gained a

functional mRNA by RNA editing.

As described above, distinct types of bases were substituted, inserted or deleted in

the *bla* genes of the revertants. These editing events are mostly located near to or in

the downstream of the frameshift-causing indel, but not in the upstream (Fig 2B-2C).

This phenomenon can be well-explained by the PTC-signaled editing of the nonsense

mRNAs: PTCs presented downstream the indel may not only signal the recognition of

nonsense mRNA but also serve as flags for RNA editing. Because the PTCs appear in

the nonsense mRNAs only in downstream of the indel, PTC-signaled RNA editing is

restricted near to or in downstream of the indel, may also restrict DNA repair to the

corresponding region in the coding gene. Therefore, a frameshift mutation is repaired

indirectly by editing nonsense mRNAs rather than the CDS directly. It is possible that

RNA editing may sometimes cause even more problems in a nonsense mRNA, while

these non-repaired mRNAs can be recycled or degraded. This process is repeated in a

cell until the gene is repaird or the PTCs are eliminated. When the cells are cultured in

ampicillin-containing media, the survival of a cell depends on whether it can produce

enough functional proteins before it was killed. By transcribing, each copy of a target

gene can produce many copies of nonsense mRNAs, each can be edited in a different

way without changing the coding sequences, the probability of successfully producing

a functional mRNA by editing nonsense mRNAs can therefore be much higher than

that of random mutagenesis directly in their coding sequences. This mechanism brings

cells potentially a greater chance of survival when a fatal frameshift mutation occurs.

4.3 *The NMGE model better explains the growth of the revertants*

As above mentioned, frameshift repair is not generally a sudden + complete event

but a progressive process. Interestingly, the sizes of wild-type colonies are uniform

(Fig 2A), showing that they are uniform in growth rates; while the sizes of revertant

colonies varied greatly, indicating varied growth rates associated with the diversified

activities of their BLA. Comparing with random mutagenesis, NMGE better explains

not only the high-level variations of *bla* but also the diversified growth rates of the

revertants:

(1). In the initial revertants, whose *bla* genes were still frameshifted, survival can

be explained in three possible ways: (a) the frameshifted *bla* is translated by

ribosomal frameshifting, producing a wild-type BLA; (b) the frameshifted *bla*

is translated by *translational readthrough*, producing a functional, partially

frameshifted BLA; (c) the initial slowly-growing revertants may also contain a

few copies of repaired *bla*, with the proportion of the repaired *bla* being too

low to be detected by Sanger sequencing. As pBR322 is a multi-copy plasmid,

it might be sufficient to support the survival of the host in ampicillin even

when only a small proportion of its copies are repaired. The host bacteria grow

slowly, and many of them failed to grow, probably due to these abnormal processes are very time- and resource-consuming, as cell energy and resources would be depleted for gene repair if the defective *bla* is not repaired soon.

(2). By NMGE, the *bla* genes were repaired progressively, not only before but also after culturing in the ampicillin media. As the substrate of BLA, ampicillin is neither the cause nor the terminator of the gene repair, but a selection pressure to screen and display the revertants, which is the result of gene repair.

(3). NMGE is triggered by the PTCs presented in the *bla* mRNAs, and the aim of NMGE is to eliminate these PTCs. In the initial revertants, there are so many PTCs in the *bla* mRNAs that the probability of producing a functional product by mRNA editing is very low, so their growth rates and survival rates are very low, as most revertants died before their *bla* genes were repaired.

(4). Later, in revertants whose *bla* are being repaired, with less and less remaining PTCs, energy and resources are used less and less for gene repair, more and more for growth, their growth/survival rates recover gradually.

(5). Finally, gene repair completes when PTCs are eliminated. The reading frames of the *bla* genes are recovered but, usually, their sequences are not the same as the original wild-type sequence but may change into a variety of variant types, and their growth/survival rates also vary greatly.

4.4 *Previous studies are supportive to the NMGE model*

In the above, we proposed that NMGE depends on RNA-directed DNA repair but is triggered by PTCs, and the nonsense mRNAs are edited prior to directing the repair of the coding DNA. When a frameshift mutation occurs, the defective gene is repaired not by direct gene editing but by NMGE. The above proposed NMGE model well explains our observations and is supported by numerous previous studies. In fact, the model is an integration of several key links, which have already been quite clear:

(1). mRNA decay is linked to other nonsense mRNA processing pathways

Nonsense-mediated mRNA decay (NMD) is the best-studied nonsense mRNA processing pathway [71-73]. In eukaryotes, NMD uses the presence of an exon junction complex (EJC) downstream of PTCs as a second signal to distinguish a PTC

from a true stop codon [91]. In addition to NMD, the nonsense mRNAs may also be subject to other nonsense-mediated pathways, such as *translational repression* [92], *transcriptional silencing* [93], and *alternative splicing* [94], all of which are connected to NMD [92-94]. In eukaryotes, the NMD regulating factors, including three interacting proteins, UPF1, UPF2 (also known as NMD2) and UPF3, are encoded by highly conserved genes originally identified in yeast [95].

Frameshifted or nonsense-mutated coding genes can be expressed through some special mechanisms, including ribosomal frameshifting [79-86] and translational readthrough [87-90]. Ribosomal frameshifting occurs in all known organisms, from *E. coli* to mammals [96-99], whereby the ribosome is guided toward a triplet codon that is shifted one basepair position to the upstream (+1) or downstream (−1). In yeast, telomere maintenance is globally controlled by the ribosomal frameshifting process and NMD [86]. Translational readthrough is a process wherein a nonsense mRNA is translated by translating PTCs each into a specific amino acid [100-104]. Mutations in *upf* genes caused nonsense suppression and subsequent readthrough in yeast and 5-20 fold decreases in the rates of NMD [105], suggesting a direct link between NMD, nonsense suppression and readthrough. The three Upf proteins all interact with Sup35, and Upf1 interact further with a second release factor, Sup45 (eRF1), and binding of an SMG-1-Upf1-eRF1-eRF3 (SURF) complex to EJC triggers Upf1 phosphorylation and NMD [106]. In addition, the nonsense suppression caused by *upf* mutations is additive to those by *sup35* and *sup45* mutations [107], suggesting that nonsense suppression is a nonsense-mRNA processing pathway that is alternative to NMD. In summary, although nonsense mRNAs usually lead to translation errors and NMD, when NMD is inhibited, they can also be translated by translational frameshifting, nonsense suppression or translational readthrough.

(2). RNA can direct DNA repair

All types of RNAs are subject to translation, degradation or processing, and a variety of molecular/cellular mechanisms are involved in RNA processing [108]. In the last few decades, studies have established a link between DNA repair and RNA processing. For example, several proteins respond to DNA damage and base-excision

repair enzymes, such as SMUG1, APE1, and PARP1, have been shown to participate in RNA surveillance and processing [109].

As is known, RNA is transcribed in the nucleus and then transported to the cytoplasm in eukaryotes [110]. Recently, there is growing evidence that RNAs can also be transported back to the nucleus to repair DNA double-strand breaks (DSBs), a process known as *RNA-directed DNA Repair* [111-116]. In yeast [115, 116] and human embryonic kidney (HEK-293) cells [117], not only a synthetic DNA/RNA hybrid but also an RNA-only oligonucleotide can serve as a template for DNA synthesis for the repair of a DSB in a homologous DNA. In *E. coli*, short RNA patches can also direct DNA modification [117, 118]. However, here we report that mRNA can direct precise and targeted repair of a frameshift mutation by itself. Although this process is not yet clear in detail at the molecular level, mismatch repair and homologous recombination are likely to be involved, as they are both upregulated in the frameshift and the revertant tested (Fig 6F).

We here demonstrated that many G:C→A:T substitutions were observed in the repaired bla in the revertants; in addition, four deaminase genes (codA, cdd, add and tadA) and the mismatch repair genes/pathways were upregulated in the frameshift and/or the revertant. We, therefore, speculate that mismatch repair and RNA editing are both involved in frameshift repair. The details of the mechanism of nonsense mRNA editing and how it connects to DNA repair has not been made clear.

(3). RNA editing is linked to DNA repair

RNA editing has been studied extensively and intensively [119-123], indicating that RNA editing is widely existing in *E. coli* [33], kinetoplastids [124], trypanosomes [125], plants [126, 127], mammals [128] and human [129-136]. Although there are a few different mechanisms of RNA editing, such as A-to-I editing [137-140], C-to-U editing [141-145], and insertion/deletion of uridine [146-148], the basic schematism are highly conserved among species. In particular, the nonsense *apoB* mRNA editing complex suppresses its own decay by NMD [149], suggesting that nonsense mRNAs can be subjected to RNA editing, which is linked to the NMD pathway.

Both DNA mismatch repair and RNA editing involve deamination, the removal of an amine group from a DNA/RNA molecule. Enzymes that catalyze this reaction are called DNA/RNA deaminases. Deamination of DNA deoxyadenosine bases generates deoxyinosine, create missense mutations predisposing humans to cancer, and interfere with other molecular genetic processes. Cytidine deamination was first recognized as a mechanism for RNA editing in wheat [150, 151]. The characterization of the tRNA-specific adenosine deaminases (ADATs) has led to the suggestion that these enzymes, as well as the cytidine/adenosine deaminases (CDARs/ADARs) acting on pre-mRNAs, belong to a superfamily of RNA dependent deaminases [152]. The first example of C→U editing is the apolipoprotein B (apo B) mRNA, in which single, site-specific cytidine deamination introduces a stop codon (UAA) into the reading frame of the edited transcript [153]. It is well known that deamination based RNA editing plays many important roles, and aberrant or dysregulated RNA editing might contribute to genomic instability in cancer [154]. In eukaryotes, several studies have provided evidence that deamination is related to both DNA repair and RNA editing. For example, endonuclease V, which is highly conserved from *E. coli* to human [59], is involved in excision repair that removes deoxyinosine from DNA.

TadA, the first prokaryotic deaminase for RNA editing identified in *E. coli,* displays sequence similarity to the yeast tRNA deaminase subunit Tad2p [55]. In addition, cytidine deaminase (CodA) has been used as a molecular model for RNA editing and a mechanism for RNA substrate recognition [33]. These studies suggest that RNA editing is conserved between prokaryotes and eukaryotes, and a direct link between the regulation of mismatch repair and RNA editing, and therefore support our speculation that mismatch repair and RNA editing may both involve in frameshift repair. However, the details of the mechanism of nonsense mRNA editing and how it connects to DNA repair has not been made clear.

Previously, we have reported that the genetic code, the protein-coding genes and genomes of some species are optimal for frameshift tolerance [155]. Widespread premature termination codons were recently observed in populations of humans [68] and fruit flies [156], suggesting ongoing evolutionary plasticity at the genetic level in

modern human and other species. In this study, we demonstrate that PTCs signal the

frameshift repair. Frameshift mutations in protein-coding genes are repaired through

NMGE, resulting in a variety of frameshift homologs, some of which are functional

and preserved in nature. Therefore, the widespread existence of frameshift homologs

among species may be considered as natural evidence for this model. In short, NMGE

is not only the molecular mechanism for frameshift repair, but also be a driving force

for molecular evolution and a source of genetic diversity.

Author Contributions

10 Xiaolong Wang conceived the study, designed the experiments, analyzed the data and

11 wrote the paper. Xuxiang Wang, Chunyan Li, Haibo Peng, and Yalei Wang performed the

12 experiments. Gang Chen and Jianye Zhang provided some equipment, discussed the paper

13 and gave some suggestions.

Acknowledgments:

15 This study was supported by The *National Natural Science Foundation of China* (Grant

16 No. 31571369).

Figure legends

Fig 1. The introduction of a frameshift mutation in the upstream of the *bla* gene in the

plasmid pBR322. Sanger sequencing diagrams: (A) the wild type (*bla+*); (B) the frameshift

mutant (*bla-*). (C) Alignment of the nucleotide sequence of *bla+* and *bla-*.

Fig 2. Growth of the wild-type/frameshift *E. coli* and Sanger sequencing of the *bla* genes:

(A) Growth of the wild-type and frameshift *E. coli* on ampicillin- or tetracycline plates: *bla+*: wild

type; *bla-*: frameshift; *bla**: revertants; *blank*: blank control; ACP: ampicillin-containing plates;

ACP: tetracycline-containing plates.

(B) The Sanger diagram of the *bla* genes of the revertants: *Top*: in the initial revertants, most of

the *bla* genes are still frameshifted; *Middle*: in the later subcultures, sequencing diagrams contains

two sets of overlapping peaks, the main peaks are the frameshifted *bla*, the second peaks are the

repaired *bla*; *Bottom*: in the final fast-growing subcultures, sequencing diagrams contains very few

overlapping peaks.

(C) The ClustalW alignment of the wild-type, frameshifted and repaired *bla* gene sequences: ***bla+***:

wild-type; *bla-*: frameshift; *bla*A-E*: different (independent) revertants; *box*: the base G deleted in

the OE-PCR; *underlined*: the bases (in blue) inserted or deleted in the revertants; *circles*: base

substitutions (G:C←→A:T transitions).

(D) The codon and amino acid unified alignment (CAUSA) of *bla* and their translated protein

sequences: *underlined*: the base G deleted in codon 46; *red box*: stop codons; *red star*: stop

signals.

Fig 3. The traditional *random mutagenesis* model for frameshift repair: *Step 1*: a frameshift

mutation occurs in a coding gene, many hidden stop codons (*red bars*) emerge in the CDS, and in

the nonsense mRNAs, cause *translational termination* and resulted in truncated products; *Step 2*:

the coding gene is repaired by a *reverse mutation* through random mutagenesis, the reading frame

is restored immediately, stop codons are hidden (*green bars*), and translation proceeds, producing

functional mRNAs and proteins, which are identical and same to the original genes and proteins.

Fig 4. NGS resequencing and variation analysis the genomes of the frameshift/revertant:

(A) The genome-wide variation map of the frameshift /revertant: to visualize the levels of read

coverages and the distribution of mutations, Circos software was used to show the clean reads of

each sample, the depth of coverage (number of clean reads covering a given base position of the

reference sequence), and the SNP/indel, using the reference genome sequence as a ruler. The

outermost circle: the coordinate position of the reference genome, and then, outside-in, the

distribution of indel/SNPs (red dots), the depth of coverage (grey), the GC content (green) and the

GC skew value (pink) in the reference genome, respectively. There are two samples, the frameshift

(*bla-*) and the revertant (*bla**), so there are two sets of indel/SNP and reads coverage depth

mapped onto the reference genome sequences.

(B) The distribution of the numbers of SNPs/indels of the frameshift/revertant on the reference

genome: The abscissa axis shows the chromosome coordination of the reference genome sequence,

and the ordinate axis represents the SNP/indel density on the genome sequence in per 10 kb. The

left scale: SNP (red peaks); the right scale: indel (blue peaks). See Table 2 for data.

(C) Mapping of reads for *bla* coding sequences and RNA transcripts. See Table 3 for data.

(D) Mapping of reads for *tet* coding sequences and RNA transcripts. See Table 3 for data.

Fig 5. Expression of the PTC-free frameshifted *bla* (*bla#*) in *E. coli* BL21:

(A) SDS-PAGE of the lysates of pET28a-*bla#* recombinant *E. coli* cells; Lane 1: protein marker;

Lane 2 and 3: whole-cell lysates of uninduced and induced cells; lane 4 and 5: the supernatant of

uninduced and induced cells; lane 6 and 7: the precipitate of uninduced and induced cells;

(B) Detection of β-lactamase activity by iodimetry: (1) the revertants of *E. coli* BL21/pET28a-bla-

caused the fading of the iodine/starch solution, suggesting that their *bla-* was repaired, produced

active β-lactamases, and transformed ampicillin into penicillium thiazole acid, which binds with

starch and competitively inhibits the binding of iodine to starch; (2) *E. coli* BL21/pET28a-bla# did

not cause fading, suggesting that it was not repaired but expressed a frameshifted BLA (BLA#),

which is an inactive product; (3) the negative control (*E. coli* BL21), no β-lactamase, no fading.

Fig 6. The transcriptome analysis of different *E. coli* strains: *W-1*: wildtype; *FS-2*: frameshift;

R-3: a revertant; *RE-4*: a subculture of the revertant; (A) The number of differentially expressed

genes (DEGs) and the cluster analysis of DEGs; (B) the most enriched GO terms by comparing

the frameshift (FS_2) with the wild type (W_1); (C) the most enriched GO terms by comparing

the revertant (R_3) with the frameshift (FS_2); (D) DNA replication pathway: (E) genes

upregulated in the RNA degradation/processing pathway; (F) genes upregulated in the mismatch

repair and homologous recombination pathways. *Underlined* in B, C: GO terms involved in the

processing of DNA or RNA, which probably also involve in frameshift repair; in D, E, F: *green*

box: genes involved in the pathways; *rex box*: genes upregulated in the frameshift or revertant.

Fig 7. The new model for frameshift repair: nonsense-mediated gene editing (NMGE): *Step 1*:

when the frameshifted gene is transcribed and translated, the nonsense mRNAs containing many

PTCs (*yellow bars*) are identified by mRNA surveillance through PTCs signaling; *Step 2*: some of

the nonsense mRNAs causing *translational termination* are degraded by NMD; *Step 3*: some of the nonsense mRNAs are translated by translational frameshifting or readthrough; *Step 4*: some of the nonsense mRNAs are repaired by *RNA editing*, PTCs are replaced by sense codons (*green bars*) through base insertion, deletion, and substitution; *Step 5*: some of the original or repaired nonsense mRNAs are transported to identify and localize the defective coding gene; *Step 6*: the coding gene is repaired by *RNA-directed DNA repair*, the reading frame is restored partially, and the translation proceeds, the remaining stop codons were readthrough, producing functional mRNAs and proteins with many variations, and so that the host recovered gradually.

2   Table 1. The natural suppressor tRNA for nonsense mutations (the *readthrough rules*).

Site	tRNA (AA)	Wild type		Correction	
		Code	**Anti-code**	**Code**	**Anti-code**
supD	Ser (S)	→ UCG	CGA←	→ UAG	CUA←
supE	Gln (Q)	→ CAG	CUG←	→ UAG	CUA←
supF	Tyr (Y)	→ UAC	GUA←	→ UAG	CUA←
supG	Lys (K)	→ AAA	UUU←	→ UAA	UUA←
supU	Trp (W)	→ UGG	CCA←	→ UGA	UCA←

5   Table 2. The summary of SNP/indel variations in the *bla**/- genome

Sample	Indels			SNPs				
	INS	DEL	Total	ts	tv	ts/tv	Total	Density
bla-	2	4	6	43	27	1.59	70	0.02/Kb
bla*	2	3	5	42	27	1.56	69	0.01/Kb

11   Note: INS: insertion; DEL: deletion; ts: transitions; tv: transversions;

14   Table 3. The count and density of raw SNPs of the two genes in the plasmid pbr322

Count of SNPs	tet (1191 bp)		bla (861 bp)	
	Count	Density	Count	Density
Frameshift-DNA	76	63.81/Kb	60	69.69/Kb
Revertant-DNA	70	58.77/Kb	68	78.98/Kb
Wildtype-RNA	5	4.20/Kb	2	2.32/Kb
Revertant-RNA	6	5.04/Kb	6	6.97/Kb

17   Table 4. Possible NMGE-relevant genes and pathways that were upregulated in the frameshift

Pathway Term	Database	ID	Input number	Background number	P-Value	Corrected P-Value
ABC transporters	KEGG	eco02010	32	171	0.0001	0.0090
RNA degradation	KEGG	eco03018	1	15	0.7419	0.9762
DNA replication	KEGG	eco03030	1	17	0.7822	0.9762
Mismatch repair	KEGG	eco03430	1	22	0.8575	0.9796

Homologous recombination	KEGG	eco03440	1	27	0.9069	0.9796

3   Table 5. Possible NMGE-relevant genes and pathways that were upregulated in the revertant

Pathway Term	Database	ID	Input number	Background number	P-Value	Corrected P-Value
ABC transporters	KEGG	eco02010	32	171	3.53E-05	0.0018
Homologous recombination	KEGG	eco03440	2	27	0.6539	0.9519
RNA degradation	KEGG	eco03018	3	15	0.1017	0.5470
DNA replication	KEGG	eco03030	1	17	0.7564	0.9519
Mismatch repair	KEGG	eco03430	1	22	0.8357	0.9519

References

1. Seligmann, H. and D.D. Pollock, *The ambush hypothesis: hidden stop codons prevent off-frame gene reading.* DNA Cell Biol, 2004. **23**(10): p. 701-5.

2. Owens, M., et al., *SOS1 frameshift mutations cause pure mucosal neuroma syndrome, a clinical phenotype distinct from multiple endocrine neoplasia type 2B.* Clin Endocrinol (Oxf), 2016. **84**(5): p. 715-9.

3. Yan, S.E., et al., *In-depth analysis of hyaline fibromatosis syndrome frameshift mutations at the same site reveal the necessity of personalized therapy.* Hum Mutat, 2013. **34**(7): p. 1005-17.

4. Amiel, J., et al., *Polyalanine expansion and frameshift mutations of the paired-like homeobox gene PHOX2B in congenital central hypoventilation syndrome.* Nat Genet, 2003. **33**(4): p. 459-61.

5. Kang, S., et al., *GLI3 frameshift mutations cause autosomal dominant Pallister-Hall syndrome.* Nat Genet, 1997. **15**(3): p. 266-8.

6. Jensen, N.M., et al., *An update on targeted gene repair in mammalian cells: methods and mechanisms.* J Biomed Sci, 2011. **18**: p. 10.

7. Streisinger, G., et al., *Frameshift mutations and the genetic code. This paper is dedicated to Professor Theodosius Dobzhansky on the occasion of his 66th birthday.* Cold Spring Harb Symp Quant Biol, 1966. **31**: p. 77-84.

8. Li, H., et al., *The Sequence Alignment/Map format and SAMtools.* Bioinformatics, 2009. **25**(16): p. 2078-9.

9. DePristo, M.A., et al., *A framework for variation discovery and genotyping using next-generation DNA sequencing data.* Nat Genet, 2011. **43**(5): p. 491-8.

10. Krzywinski, M., et al., *Circos: an information aesthetic for comparative genomics.* Genome Res, 2009. **19**(9): p. 1639-45.

11. Fan, X., et al., *BreakDancer: Identification of Genomic Structural Variation from Paired-End Read Mapping.* Curr Protoc Bioinformatics, 2014. **45**: p. 15 6 1-11.

12. Chen, K., et al., *BreakDancer: an algorithm for high-resolution mapping of genomic structural variation.* Nat Methods, 2009. **6**(9): p. 677-81.

13. Langmead, B., *Aligning short sequencing reads with Bowtie.* Curr Protoc Bioinformatics, 2010. **Chapter 11**: p. Unit 11 7.

14. Anders, S., P.T. Pyl, and W. Huber, *HTSeq--a Python framework to work with high-throughput sequencing data.* Bioinformatics, 2015. **31**(2): p. 166-9.

15. Robinson, M.D., D.J. McCarthy, and G.K. Smyth, *edgeR: a Bioconductor package for differential expression analysis of digital gene expression data.* Bioinformatics, 2010. **26**(1): p. 139-40.

16. Wang, L., et al., *DEGseq: an R package for identifying differentially expressed genes from RNA-seq data.* Bioinformatics, 2010. **26**(1): p. 136-8.

17. Young, M.D., et al., *Gene ontology analysis for RNA-seq: accounting for selection bias.* Genome Biol, 2010. **11**(2): p. R14.

18. Xie, C., et al., *KOBAS 2.0: a web server for annotation and identification of enriched pathways and diseases.* Nucleic Acids Res, 2011. **39**(Web Server issue): p. W316-22.

19.   Lee, H., et al., *Rate and molecular spectrum of spontaneous mutations in the bacterium Escherichia coli as determined by whole-genome sequencing.* Proc Natl Acad Sci U S A, 2012. **109**(41): p. E2774-83.

20.   Wang, X. and C. Yang, *CAUSA 2.0: accurate and consistent evolutionary analysis of proteins using codon and amino acid unified sequence alignments.* PeerJ PrePrints **3**.

21.   Schaaper, R.M., *Mechanisms of mutagenesis in the Escherichia coli mutator mutD5: role of DNA mismatch repair.* Proc Natl Acad Sci U S A, 1988. **85**(21): p. 8126-30.

22.   Itoh, Y., S. Sekine, and S. Yokoyama, *Crystal structure of the full-length bacterial selenocysteine-specific elongation factor SelB.* Nucleic Acids Res, 2015. **43**(18): p. 9028-38.

23.   Baron, C. and A. Bock, *The length of the aminoacyl-acceptor stem of the selenocysteine-specific tRNA(Sec) of Escherichia coli is the determinant for binding to elongation factors SELB or Tu.* J Biol Chem, 1991. **266**(30): p. 20375-9.

24.   Ireton, G.C., et al., *The structure of Escherichia coli cytosine deaminase.* J Mol Biol, 2002. **315**(4): p. 687-97.

25.   Danielsen, S., et al., *Characterization of the Escherichia coli codBA operon encoding cytosine permease and cytosine deaminase.* Mol Microbiol, 1992. **6**(10): p. 1335-44.

26.   MacGinnitie, A.J., S. Anant, and N.O. Davidson, *Mutagenesis of apobec-1, the catalytic subunit of the mammalian apolipoprotein B mRNA editing enzyme, reveals distinct domains that mediate cytosine nucleoside deaminase, RNA binding, and RNA editing activity.* J Biol Chem, 1995. **270**(24): p. 14768-75.

27.   Vita, A., et al., *Cytidine deaminase from Escherichia coli B. Purification and enzymatic and molecular properties.* Biochemistry, 1985. **24**(21): p. 6020-4.

28.   Ashley, G.W. and P.A. Bartlett, *Purification and properties of cytidine deaminase from escherichia coli.* J Biol Chem, 1984. **259**(21): p. 13615-20.

29.   von Dippe, P.J., et al., *Deoxycytidine transport in the presence of a cytidine deaminase inhibitor and the transport of uracil in Escherichia coli B.* J Biol Chem, 1975. **250**(10): p. 3666-71.

30.   Hosono, H. and S. Kuno, *The purification and properties of cytidine deaminase from Escherichia coli.* J Biochem, 1973. **74**(4): p. 797-803.

31.   Cohen, R.M. and R. Wolfenden, *Cytidine deaminase from Escherichia coli. Purification, properties and inhibition by the potential transition state analog 3,4,5,6-tetrahydrouridine.* J Biol Chem, 1971. **246**(24): p. 7561-5.

32.   Ashley, G.W. and P.A. Bartlett, *Inhibition of Escherichia coli cytidine deaminase by a phosphapyrimidine nucleoside.* J Biol Chem, 1984. **259**(21): p. 13621-7.

33.   Navaratnam, N., et al., *Escherichia coli cytidine deaminase provides a molecular model for ApoB RNA editing and a mechanism for RNA substrate recognition.* J Mol Biol, 1998. **275**(4): p. 695-714.

34.   Navaratnam, N., et al., *Evolutionary origins of apoB mRNA editing: catalysis by a cytidine deaminase that has acquired a novel RNA-binding motif at its active site.* Cell, 1995. **81**(2): p. 187-95.

35.   Nygaard, P., *Adenosine deaminase from Escherichia coli.* Methods Enzymol, 1978. **51**: p. 508-12.

36.   Koch, A.L. and G. Vallee, *The properties of adenosine deaminase and adenosine nucleoside phosphorylase in extracts of Escherichia coli.* J Biol Chem, 1959. **234**(5): p. 1213-8.

37. Remy, C.N. and S.H. Love, *Induction of adenosine deaminase in Escherichia coli.* J Bacteriol, 1968. **96**(1): p. 76-85.

38. Kim, U., et al., *Molecular cloning of cDNA for double-stranded RNA adenosine deaminase, a candidate enzyme for nuclear RNA editing.* Proc Natl Acad Sci U S A, 1994. **91**(24): p. 11457-61.

39. Kim, U. and K. Nishikura, *Double-stranded RNA adenosine deaminase as a potential mammalian RNA editing factor.* Semin Cell Biol, 1993. **4**(4): p. 285-93.

40. Oakes, E., et al., *Adenosine Deaminase That Acts on RNA 3 (ADAR3) Binding to Glutamate Receptor Subunit B Pre-mRNA Inhibits RNA Editing in Glioblastoma.* J Biol Chem, 2017. **292**(10): p. 4326-4335.

41. Ward, S.V., et al., *RNA editing enzyme adenosine deaminase is a restriction factor for controlling measles virus replication that also is required for embryogenesis.* Proc Natl Acad Sci U S A, 2011. **108**(1): p. 331-6.

42. Herbert, A., et al., *A Z-DNA binding domain present in the human editing enzyme, double-stranded RNA adenosine deaminase.* Proc Natl Acad Sci U S A, 1997. **94**(16): p. 8421-6.

43. Yang, J.H., et al., *Purification and characterization of a human RNA adenosine deaminase for glutamate receptor B pre-mRNA editing.* Proc Natl Acad Sci U S A, 1997. **94**(9): p. 4354-9.

44. Maas, S., et al., *Structural requirements for RNA editing in glutamate receptor pre-mRNAs by recombinant double-stranded RNA adenosine deaminase.* J Biol Chem, 1996. **271**(21): p. 12221-6.

45. Polson, A.G., B.L. Bass, and J.L. Casey, *RNA editing of hepatitis delta virus antigenome by dsRNA-adenosine deaminase.* Nature, 1996. **380**(6573): p. 454-6.

46. Dabiri, G.A., et al., *Editing of the GLuR-B ion channel RNA in vitro by recombinant double-stranded RNA adenosine deaminase.* EMBO J, 1996. **15**(1): p. 34-45.

47. Hurst, S.R., et al., *Deamination of mammalian glutamate receptor RNA by Xenopus dsRNA adenosine deaminase: similarities to in vivo RNA editing.* RNA, 1995. **1**(10): p. 1051-60.

48. Wang, Y., et al., *Genomic organization and chromosomal location of the human dsRNA adenosine deaminase gene: the enzyme for glutamate-activated ion channel RNA editing.* J Mol Biol, 1995. **254**(2): p. 184-95.

49. Chen, Z., A. Hirano, and T.C. Wong, *Isolation and characterization of intranuclear ribonucleoprotein complexes associated with double-stranded RNA adenosine deaminase from brain cells: implications for RNA-editing and hypermutation of viral RNA in the CNS.* J Neurovirol, 1995. **1**(3-4): p. 295-306.

50. Lai, F., R. Drakas, and K. Nishikura, *Mutagenic analysis of double-stranded RNA adenosine deaminase, a candidate enzyme for RNA editing of glutamate-gated ion channel transcripts.* J Biol Chem, 1995. **270**(29): p. 17098-105.

51. Gerber, A., et al., *Tad1p, a yeast tRNA-specific adenosine deaminase, is related to the mammalian pre-mRNA editing enzymes ADAR1 and ADAR2.* EMBO J, 1998. **17**(16): p. 4780-9.

52.   Losey, H.C., A.J. Ruthenburg, and G.L. Verdine, *Crystal structure of Staphylococcus aureus tRNA adenosine deaminase TadA in complex with RNA.* Nat Struct Mol Biol, 2006. **13**(2): p. 153-9.

53.   Kuratani, M., et al., *Crystal structure of tRNA adenosine deaminase (TadA) from Aquifex aeolicus.* J Biol Chem, 2005. **280**(16): p. 16002-8.

54.   Lee, W.H., et al., *Crystal structure of the tRNA-specific adenosine deaminase from Streptococcus pyogenes.* Proteins, 2007. **68**(4): p. 1016-9.

55.   Wolf, J., A.P. Gerber, and W. Keller, *tadA, an essential tRNA-specific adenosine deaminase from Escherichia coli.* EMBO J, 2002. **21**(14): p. 3841-51.

56.   Maas, S., A.P. Gerber, and A. Rich, *Identification and characterization of a human tRNA-specific adenosine deaminase related to the ADAR family of pre-mRNA editing enzymes.* Proc Natl Acad Sci U S A, 1999. **96**(16): p. 8895-900.

57.   Maas, S., Y.G. Kim, and A. Rich, *Sequence, genomic organization and functional expression of the murine tRNA-specific adenosine deaminase ADAT1.* Gene, 2000. **243**(1-2): p. 59-66.

58.   Keegan, L.P., et al., *The properties of a tRNA-specific adenosine deaminase from Drosophila melanogaster support an evolutionary link between pre-mRNA editing and tRNA modification.* Mol Cell Biol, 2000. **20**(3): p. 825-33.

59.   Kuraoka, I., *Diversity of Endonuclease V: From DNA Repair to RNA Editing.* Biomolecules, 2015. **5**(4): p. 2194-206.

60.   Ge, Z., et al., *Polypyrimidine tract binding protein 1 protects mRNAs from recognition by the nonsense-mediated mRNA decay pathway.* Elife, 2016. **5**.

61.   Schweingruber, C., et al., *Nonsense-mediated mRNA decay - mechanisms of substrate mRNA recognition and degradation in mammalian cells.* Biochim Biophys Acta, 2013. **1829**(6-7): p. 612-23.

62.   Niu, D.K. and J.L. Cao, *Nucleosome deposition and DNA methylation may participate in the recognition of premature termination codon in nonsense-mediated mRNA decay.* FEBS Lett, 2010. **584**(16): p. 3509-12.

63.   Muhlemann, O., et al., *Recognition and elimination of nonsense mRNA.* Biochim Biophys Acta, 2008. **1779**(9): p. 538-49.

64.   Muhlemann, O., *Recognition of nonsense mRNA: towards a unified model.* Biochem Soc Trans, 2008. **36**(Pt 3): p. 497-501.

65.   Muhlrad, D. and R. Parker, *Recognition of yeast mRNAs as "nonsense containing" leads to both inhibition of mRNA translation and mRNA degradation: implications for the control of mRNA decapping.* Mol Biol Cell, 1999. **10**(11): p. 3971-8.

66.   Zhang, J. and L.E. Maquat, *Evidence that the decay of nucleus-associated nonsense mRNA for human triosephosphate isomerase involves nonsense codon recognition after splicing.* RNA, 1996. **2**(3): p. 235-43.

67.   Hwang, J. and Y.K. Kim, *When a ribosome encounters a premature termination codon.* BMB Rep, 2013. **46**(1): p. 9-16.

68.   Fujikura, K., *Premature termination codons in modern human genomes.* Sci Rep, 2016. **6**: p. 22468.

69.   Wong, T.Y., et al., *Role of premature stop codons in bacterial evolution.* J Bacteriol, 2008. **190**(20): p. 6718-25.

70. Cirulli, E.T., et al., *A whole-genome analysis of premature termination codons.* Genomics, 2011. **98**(5): p. 337-42.

71. Le Hir, H., et al., *The exon-exon junction complex provides a binding platform for factors involved in mRNA export and nonsense-mediated mRNA decay.* EMBO J, 2001. **20**(17): p. 4987-97.

72. Gonzalez, C.I., W. Wang, and S.W. Peltz, *Nonsense-mediated mRNA decay in Saccharomyces cerevisiae: a quality control mechanism that degrades transcripts harboring premature termination codons.* Cold Spring Harb Symp Quant Biol, 2001. **66**: p. 321-8.

73. Zhang, J. and L.E. Maquat, *Evidence that translation reinitiation abrogates nonsense-mediated mRNA decay in mammalian cells.* EMBO J, 1997. **16**(4): p. 826-33.

74. Son, H.G., et al., *RNA surveillance via nonsense-mediated mRNA decay is crucial for longevity in daf-2/insulin/IGF-1 mutant C. elegans.* Nat Commun, 2017. **8**: p. 14749.

75. Fang, Y., et al., *Nonsense-mediated mRNA decay of collagen -emerging complexity in RNA surveillance mechanisms.* J Cell Sci, 2013. **126**(Pt 12): p. 2551-60.

76. Behm-Ansmant, I. and E. Izaurralde, *Quality control of gene expression: a stepwise assembly pathway for the surveillance complex that triggers nonsense-mediated mRNA decay.* Genes Dev, 2006. **20**(4): p. 391-8.

77. Wilusz, C.J., W. Wang, and S.W. Peltz, *Curbing the nonsense: the activation and regulation of mRNA surveillance.* Genes Dev, 2001. **15**(21): p. 2781-5.

78. Richards, J., et al., *Quality control of bacterial mRNA decoding and decay.* Biochim Biophys Acta, 2008. **1779**(9): p. 574-82.

79. Xu, J., R.W. Hendrix, and R.L. Duda, *Conserved translational frameshift in dsDNA bacteriophage tail assembly genes.* Molecular Cell, 2004. **16**(1): p. 11-21.

80. Baykal, U., A.L. Moyne, and S. Tuzun, *A frameshift in the coding region of a novel tomato class I basic chitinase gene makes it a pseudogene with a functional wound-responsive promoter.* Gene, 2006. **376**(1): p. 37-46.

81. Chen, J., et al., *Dynamic pathways of -1 translational frameshifting.* Nature, 2014. **512**(7514): p. 328-32.

82. Gaudin, C., et al., *Structure of the RNA signal essential for translational frameshifting in HIV-1.* J Mol Biol, 2005. **349**(5): p. 1024-35.

83. Bekaert, M., et al., *Identification of programmed translational -1 frameshifting sites in the genome of Saccharomyces cerevisiae.* Genome Res, 2005. **15**(10): p. 1411-20.

84. Licznar, P., et al., *Programmed translational -1 frameshifting on hexanucleotide motifs and the wobble properties of tRNAs.* EMBO J, 2003. **22**(18): p. 4770-8.

85. Klobutcher, L.A. and P.J. Farabaugh, *Shifty ciliates: frequent programmed translational frameshifting in euplotids.* Cell, 2002. **111**(6): p. 763-6.

86. Advani, V.M., A.T. Belew, and J.D. Dinman, *Yeast telomere maintenance is globally controlled by programmed ribosomal frameshifting and the nonsense-mediated mRNA decay pathway.* Translation (Austin), 2013. **1**(1): p. e24418.

87. Claisse, M.L., M. Boguta, and W. Zagorski, *Translational readthrough of a termination codon in the yeast mitochondrial mRNA VAR1 as a result of mutation in the release factor mRF1.* Acta Biochim Pol, 2005. **52**(1): p. 129-37.

88. Halvey, P.J., D.C. Liebler, and R.J. Slebos, *A Reporter System for Translational Readthrough of Stop Codons in Human Cells.* FEBS Open Bio, 2012. **2**: p. 56-59.

89.   Eswarappa, S.M., et al., *Programmed translational readthrough generates antiangiogenic VEGF-Ax.* Cell, 2014. **157**(7): p. 1605-18.

90.   Staff, P.G., *Correction: Functional Translational Readthrough: A Systems Biology Perspective.* PLoS Genet, 2016. **12**(11): p. e1006434.

91.   Scofield, D.G., X. Hong, and M. Lynch, *Position of the final intron in full-length transcripts: determined by NMD?* Mol Biol Evol, 2007. **24**(4): p. 896-9.

92.   Isken, O., et al., *Upf1 phosphorylation triggers translational repression during nonsense-mediated mRNA decay.* Cell, 2008. **133**(2): p. 314-27.

93.   Ward, A.J., et al., *Nonsense-mediated decay as a terminating mechanism for antisense oligonucleotides.* Nucleic Acids Res, 2014. **42**(9): p. 5871-9.

94.   Lewis, B.P., R.E. Green, and S.E. Brenner, *Evidence for the widespread coupling of alternative splicing and nonsense-mediated mRNA decay in humans.* Proc Natl Acad Sci U S A, 2003. **100**(1): p. 189-92.

95.   He, F., A.H. Brown, and A. Jacobson, *Upf1p, Nmd2p, and Upf3p are interacting components of the yeast nonsense-mediated mRNA decay pathway.* Mol Cell Biol, 1997. **17**(3): p. 1580-94.

96.   Namy, O., et al., *A mechanical explanation of RNA pseudoknot function in programmed ribosomal frameshifting.* Nature, 2006. **441**(7090): p. 244-7.

97.   Dinman, J.D., *Control of gene expression by translational recoding.* Adv Protein Chem Struct Biol, 2012. **86**: p. 129-49.

98.   Dinman, J.D., *Mechanisms and implications of programmed translational frameshifting.* Wiley Interdiscip Rev RNA, 2012. **3**(5): p. 661-73.

99.   Ketteler, R., *On programmed ribosomal frameshifting: the alternative proteomes.* Front Genet, 2012. **3**: p. 242.

100.   Kopelowitz, J., et al., *Influence of codon context on UGA suppression and readthrough.* J Mol Biol, 1992. **225**(2): p. 261-9.

101.   Weiss, R.B., *Ribosomal frameshifting, jumping and readthrough.* Curr Opin Cell Biol, 1991. **3**(6): p. 1051-5.

102.   Engelberg-Kulka, H., L. Dekel, and M. Israeli-Reches, *Regulation of the escherichia coli tryptophan operon by readthrough of UGA termination codons.* Biochem Biophys Res Commun, 1981. **98**(4): p. 1008-15.

103.   Seidman, J.S., B.D. Janssen, and C.S. Hayes, *Alternative fates of paused ribosomes during translation termination.* J Biol Chem, 2011. **286**(36): p. 31105-12.

104.   Su, D., Y. Li, and V.N. Gladyshev, *Selenocysteine insertion directed by the 3'-UTR SECIS element in Escherichia coli.* Nucleic Acids Res, 2005. **33**(8): p. 2486-92.

105.   Leeds, P., et al., *Gene products that promote mRNA turnover in Saccharomyces cerevisiae.* Mol Cell Biol, 1992. **12**(5): p. 2165-77.

106.   Kashima, I., et al., *Binding of a novel SMG-1-Upf1-eRF1-eRF3 complex (SURF) to the exon junction complex triggers Upf1 phosphorylation and nonsense-mediated mRNA decay.* Genes Dev, 2006. **20**(3): p. 355-67.

107.   Keeling, K.M., et al., *Leaky termination at premature stop codons antagonizes nonsense-mediated mRNA decay in S. cerevisiae.* RNA, 2004. **10**(4): p. 691-703.

108.   Isken, O. and L.E. Maquat, *Quality control of eukaryotic mRNA: safeguarding cells from abnormal mRNA function.* Genes Dev, 2007. **21**(15): p. 1833-56.

109.   Vohhodina, J., D.P. Harkin, and K.I. Savage, *Dual roles of DNA repair enzymes in RNA biology/post-transcriptional control.* Wiley Interdiscip Rev RNA, 2016.

110.   Nakielny, S. and G. Dreyfuss, *Transport of proteins and RNAs in and out of the nucleus.* Cell, 1999. **99**(7): p. 677-90.

111.   Trott, D.A. and A.C. Porter, *Hypothesis: transcript-templated repair of DNA double-strand breaks.* Bioessays, 2006. **28**(1): p. 78-83.

112.   Keskin, H., C. Meers, and F. Storici, *Transcript RNA supports precise repair of its own DNA gene.* RNA Biol, 2016. **13**(2): p. 157-65.

113.   Keskin, H., et al., *Transcript-RNA-templated DNA recombination and repair.* Nature, 2014. **515**(7527): p. 436-9.

114.   Wei, L., et al., *DNA damage during the G0/G1 phase triggers RNA-templated, Cockayne syndrome B-dependent homologous recombination.* Proc Natl Acad Sci U S A, 2015. **112**(27): p. E3495-504.

115.   Storici, F., *RNA-mediated DNA modifications and RNA-templated DNA repair.* Curr Opin Mol Ther, 2008. **10**(3): p. 224-30.

116.   Storici, F., et al., *RNA-templated DNA repair.* Nature, 2007. **447**(7142): p. 338-41.

117.   Shen, Y., et al., *RNA-driven genetic changes in bacteria and in human cells.* Mutat Res, 2011. **717**(1-2): p. 91-8.

118.   Thaler, D.S., G. Tombline, and K. Zahn, *Short-patch reverse transcription in Escherichia coli.* Genetics, 1995. **140**(3): p. 909-15.

119.   Kugita, M., et al., *RNA editing in hornwort chloroplasts makes more than half the genes functional.* Nucleic Acids Res, 2003. **31**(9): p. 2417-23.

120.   Bock, R., *Sense from nonsense: how the genetic information of chloroplasts is altered by RNA editing.* Biochimie, 2000. **82**(6-7): p. 549-57.

121.   Horvath, A., E.A. Berry, and D.A. Maslov, *Translation of the edited mRNA for cytochrome b in trypanosome mitochondria.* Science, 2000. **287**(5458): p. 1639-40.

122.   Yoshinaga, K., et al., *Extensive RNA editing and possible double-stranded structures determining editing sites in the atpB transcripts of hornwort chloroplasts.* Nucleic Acids Res, 1997. **25**(23): p. 4830-4.

123.   Sloof, P., et al., *RNA editing in mitochondria of cultured trypanosomatids: translatable mRNAs for NADH-dehydrogenase subunits are missing.* J Bioenerg Biomembr, 1994. **26**(2): p. 193-203.

124.   Simpson, L. and J. Shaw, *RNA editing and the mitochondrial cryptogenes of kinetoplastid protozoa.* Cell, 1989. **57**(3): p. 355-66.

125.   Alfonzo, J.D., O. Thiemann, and L. Simpson, *The mechanism of U insertion/deletion RNA editing in kinetoplastid mitochondria.* Nucleic Acids Res, 1997. **25**(19): p. 3751-9.

126.   Takenaka, M., et al., *RNA editing in plants and its evolution.* Annu Rev Genet, 2013. **47**: p. 335-52.

127.   Graves, P.V., et al., *Direct protein sequencing of wheat mitochondrial ATP synthase subunit 9 confirms RNA editing in plants.* J Mol Biol, 1990. **214**(1): p. 1-6.

128.   Slavov, D., M. Clark, and K. Gardiner, *Comparative analysis of the RED1 and RED2 A-to-I RNA editing genes from mammals, pufferfish and zebrafish.* Gene, 2000. **250**(1-2): p. 41-51.

129.   Zougman, A., et al., *Evidence for insertional RNA editing in humans.* Curr Biol, 2008. **18**(22): p. 1760-5.

130.   Athanasiadis, A., A. Rich, and S. Maas, *Widespread A-to-I RNA editing of Alu-containing mRNAs in the human transcriptome.* PLoS Biol, 2004. **2**(12): p. e391.

131.   Wang, Y., et al., *Systematic characterization of A-to-I RNA editing hotspots in microRNAs across human cancers.* Genome Res, 2017.

132.   Picardi, E., D.S. Horner, and G. Pesole, *Single cell transcriptomics reveals specific RNA editing signatures in the human brain.* RNA, 2017.

133.   Xu, G. and J. Zhang, *Human coding RNA editing is generally nonadaptive.* Proc Natl Acad Sci U S A, 2014. **111**(10): p. 3769-74.

134.   Bazak, L., et al., *A-to-I RNA editing occurs at over a hundred million genomic sites, located in a majority of human genes.* Genome Res, 2014. **24**(3): p. 365-76.

135.   Shtrichman, R., et al., *Altered A-to-I RNA editing in human embryogenesis.* PLoS One, 2012. **7**(7): p. e41576.

136.   Bahn, J.H., et al., *Accurate identification of A-to-I RNA editing in human by transcriptome sequencing.* Genome Res, 2012. **22**(1): p. 142-50.

137.   Amariglio, N. and G. Rechavi, *A-to-I RNA editing: a new regulatory mechanism of global gene expression.* Blood Cells Mol Dis, 2007. **39**(2): p. 151-5.

138.   Silberberg, G., et al., *Deregulation of the A-to-I RNA editing mechanism in psychiatric disorders.* Hum Mol Genet, 2012. **21**(2): p. 311-21.

139.   Terajima, H., et al., *ADARB1 catalyzes circadian A-to-I editing and regulates RNA rhythm.* Nat Genet, 2017. **49**(1): p. 146-151.

140.   Valente, L. and K. Nishikura, *ADAR gene family and A-to-I RNA editing: diverse roles in posttranscriptional gene regulation.* Prog Nucleic Acid Res Mol Biol, 2005. **79**: p. 299-338.

141.   Blanc, V. and N.O. Davidson, *C-to-U RNA editing: mechanisms leading to genetic diversity.* J Biol Chem, 2003. **278**(3): p. 1395-8.

142.   Bourara, K., S. Litvak, and A. Araya, *Generation of G-to-A and C-to-U changes in HIV-1 transcripts by RNA editing.* Science, 2000. **289**(5484): p. 1564-6.

143.   Giege, P. and A. Brennicke, *RNA editing in Arabidopsis mitochondria effects 441 C to U changes in ORFs.* Proc Natl Acad Sci U S A, 1999. **96**(26): p. 15324-9.

144.   Sharma, S., et al., *Transient overexpression of exogenous APOBEC3A causes C-to-U RNA editing of thousands of genes.* RNA Biol, 2016: p. 1-8.

145.   Fossat, N., et al., *C to U RNA editing mediated by APOBEC1 requires RNA-binding protein RBM47.* EMBO Rep, 2014. **15**(8): p. 903-10.

146.   Simpson, L., S. Sbicego, and R. Aphasizhev, *Uridine insertion/deletion RNA editing in trypanosome mitochondria: a complex business.* Rna, 2003. **9**(3): p. 265-76.

147.   Kable, M.L., et al., *RNA editing: a mechanism for gRNA-specified uridylate insertion into precursor mRNA.* Science, 1996. **273**(5279): p. 1189-95.

148.   Ringpis, G.E., J. Stagno, and R. Aphasizhev, *Mechanism of U-insertion RNA editing in trypanosome mitochondria: characterization of RET2 functional domains by mutational analysis.* J Mol Biol, 2010. **399**(5): p. 696-706.

149.   Chester, A., et al., *The apolipoprotein B mRNA editing complex performs a multifunctional cycle and suppresses nonsense-mediated decay.* EMBO J, 2003. **22**(15): p. 3971-82.

150.   Yu, W. and W. Schuster, *Evidence for a site-specific cytidine deamination reaction involved in C to U RNA editing of plant mitochondria.* J Biol Chem, 1995. **270**(31): p. 18227-33.

151.   Blanc, V., S. Litvak, and A. Araya, *RNA editing in wheat mitochondria proceeds by a deamination mechanism.* FEBS Lett, 1995. **373**(1): p. 56-60.

152.   Gerber, A.P. and W. Keller, *RNA editing by base deamination: more enzymes, more targets, new mysteries.* Trends Biochem Sci, 2001. **26**(6): p. 376-84.

153.   Young, S.G., *Recent progress in understanding apolipoprotein B.* Circulation, 1990. **82**(5): p. 1574-94.

154.   Anant, S. and N.O. Davidson, *Hydrolytic nucleoside and nucleotide deamination, and genetic instability: a possible link between RNA-editing enzymes and cancer?* Trends Mol Med, 2003. **9**(4): p. 147-52.

155.   Wang, X., et al., *Why are frameshift homologs widespread within and across species?* bioRxiv, 2016.

156.   Yang, H., et al., *Expression profile and gene age jointly shaped the genome-wide distribution of premature termination codons in a Drosophila melanogaster population.* Mol Biol Evol, 2015. **32**(1): p. 216-28.

Fig 1. The frameshift mutagenesis of the *bla* gene in the plasmid pBR322

Fig 2A

Fig 2B

Initial revertants (grow Slowly)

Frameshift: A G T G G G T T A C A T C *A A C T G G A T C T C A A C A G C G G T A A G*

Wild-type: A G T G G G T T A C A T C G A A C T G G A T C T C A A C A G C G G T A A G

Later subculture (grow faster)

Frameshift: A G T G G G T T A C A T C *A A C T G G A T C T C A A C A G C G G T A A G .*
(main peak)

Revertants: A G T G G G T T A C A T C C A A C T G G A T C T C A A C A G C G G T A A G
(2nd peak)

Wild-type: A G T G G G T T A C A T C G A A C T G G A T C T C A A C A G C G G T A A G

Final subculture (grow fast)

Revertants: A G T G G G T T A C A T C G A A C T G G A T C T C A A A A G C G G T A A G

Wild-type: A G T G G G T T A C A T C G A A C T G G A T C T C A A C A G C G G T A A G

Fig 2C

```
                        20                    40                    60
bla+  : ATGAGTATTCAACATTTCCGTGTCGCCCTTATTCCCTTTTTTGCGGCATTTTGCCTTCCTGTTTTTGCTCACCCAGA : 77
bla-  : ATGAGTATTCAACATTTCCGTGTCGCCCTTATTCCCTTTTTTGCGGCATTTTGCCTTCCTGTTTTTGCTCACCCAGA : 77
bla*A : ATGAGTATTCAACATTTCCGTGTCGCCCTTATTCCCTTTTTTGCGGCATTTTGCCTTCCTGTTTTTGCTCACCCAGA : 77
bla*B : ATGAGTATTCAACATTTCCGTGTCGCCCTTATTCCCTTTTTTGCGGCATTTTGCCTTCCTGTTTTTGCTCACCCAGA : 77
bla*C : ATGAGTATTCAACATTTCCGTGTCGCCCTTATTCCCTTTTTTGCGGCATTTTGCCTTCCTGTTTTTGCTCACCCAGA : 77
bla*D : ATGAGTATTCAACATTTCCGTGTCGCCCTTATTCCCTTTTTTGCGGCATTTTGCCTTCCTGTTTTTGCTCACCCAGA : 77
bla*E : ATGAGTATTCAACATTTCCGTGTCGCCCTTATTCCCTTTTTTGCGGCATTTTGCCTTCCTGTTTTTGCTCACCCAGA : 77

                        80                   100                   120                   140
bla+  : AACGCTGGTGAAAGTAAAAGATGCTGAAGATCAGTTGGGTGCACGAGTGGG-TTACATCGAACTGGAT-CTCAACAG : 152
bla-  : AACGCTGGTGAAAGTAAAAGATGCTGAAGATCAGTTGGGTGCACGAGTGGG-TTACATC-AACTGGAT-CTCAACAG : 151
bla*A : AACGCTGGTGAAAGTAAAAGATGCTGAAGATCAGTTGGGTGCACGAGTGGG-TTACATC-AACTGGAT-CTCAACAA : 151
bla*B : AACGCTGGTGAAAGTAAAAGATGCTGAAGATCAGTTGGGTGCACGAGTGGGGTACCTTC--ACTGGATTCCAACAC : 152
bla*C : AACGCTGGTGAAAGTAAAAGATGCTGAAGATCAGTTGGGTGCACGAGTGGGGTTACTTC--ACTGGATTCTCACCAC : 152
bla*D : AACGCTGGTGAAAGTAAAAGATGCTGAAGATCAGTTGGGTGCACGAGTGGG-TTACATC-AACTGGAT-CTCAACAG : 151
bla*E : AACGCTGGTGAAAGTAAAAGATGCTGAAGATCAGTTGGGTGCACGAGTGGG-TTACATC-AACTGGAT-CTCAACAG : 151

                       160                   180                   200                   220
bla+  : CGGTAAGATCCTTGAGAGTTTTCGCCCCGAAGAACGTTTTCCAATGATGAGCACTTTTAAAGTTCTGCTATGTGGCG : 229
bla-  : CGGTAAGATCCTTGAGAGTTTTCGCCCCGAAGAACGTTTTCCAATGATGAGCACTTTTAAAGTTCTGCTATGTGGCG : 228
bla*A : CGGGAAAATCCTTGAGAGTTTTCCCCCCCAAAAACGTTTCCAATGGTGAACACTTTTAAAGTTCTGGTATGGGGCG : 228
bla*B : CGTAAAAATCCTGGAAATTTTCCCCCCAAAAAAC-TTTTCCAATGGAGAACACTTTTAAAGTTCGGCTAGGGGCC : 228
bla*C : CGGTAAAATCCTTGAAAGTTTTCCCCCCAAAAACGTTTTCCAAGGATAACAACTTTTAAAGTTCGGCTATG-GGCC : 228
bla*D : CGGTAAGATCCTTGAGAGTTTTCGCCCCGAAGAACGTTTTCCAATGATGAACACTTTTAAAGTTCTGCTATGTGGCG : 228
bla*E : CGGTAAGATCCTTGAGAGTTTTCGCCCCGAAGAACGTTTTCCAATGATGAGCACTTTTAAAGTTCTGCTATGTGGCG : 228
```

Fig 2D

Fig 3   *Random mutagenesis*

Fig 4A

bla-

bla*

NC_000913

Fig 4B

bla-

bla*

Fig 4C

bla-

bla*

bla+
transcripts

bla*
transcripts

Fig 4D

tet CDS
bla-

tet CDS
bla*

tet **transcripts**
bla-

tet **transcripts**
bla*

Fig 5. *Expression of the PTC-free frameshifted bla (bla#)*

Fig 6. Transcriptome Analysis

6A Numbers of differentially expressed genes (DGE)

A: FS_2vsW_1
B: R_3vsW_1
C: RE_4vsW_1

6B

6C

The Most Enriched GO Terms (R_3vsFS_2)

6D genes upregulated in DNA replication

6E genes upregulated in RNA degradation/processing

6F genes upregulated in mismatch repair/homologous recombination

Fig 7 *Nonsense-mediated gene editing (NMGE)*