Identification of *Klebsiella* capsule synthesis loci from whole genome

data

Kelly L. Wyres[1,2,Ψ], Ryan R. Wick[1,2], Claire Gorrie[1,2,], Adam Jenney[3], Rainer Follador[4],

Nicholas R. Thomson[5,6] and Kathryn E. Holt[1,2,Ψ]

[1]Centre for Systems Genomics, University of Melbourne, Parkville, Victoria 3010,

Australia

[2]Department of Biochemistry and Molecular Biology, Bio21 Molecular Science and

Biotechnology Institute, University of Melbourne, Parkville, Victoria 3010, Australia

[3]Department Infectious Diseases and Microbiology Unit, The Alfred Hospital,

Melbourne, Victoria 3004, Australia

[4]LimmaTech Biologics AG, Schlieren, Switzerland

[5]The Wellcome Trust Sanger Institute, Hinxton, Cambridge, UK

[6]London School of Hygiene and Tropical Medicine, Keppel Street London, UK

[Ψ] **Corresponding authors**

Abstract

Background: *Klebsiella pneumoniae* and close relatives are a growing cause of healthcare-associated infections for which increasing rates of multi-drug resistance are a major concern. The *Klebsiella* polysaccharide capsule is a major virulence determinant and epidemiological marker. However, little is known about capsule epidemiology since serological typing is not widely accessible, and many isolates are serologically non-typeable. Molecular methods for capsular typing are needed, but existing methods lack sensitivity and specificity and fail to take advantage of the information available in whole-genome sequence data, which is increasingly being generated for surveillance and investigation of *Klebsiella*.

Methods: We investigated the diversity of capsule synthesis loci (K loci) among a large, diverse collection of 2503 genome sequences of *K. pneumoniae* and closely related species. We incorporated analyses of both full-length K locus DNA sequences and clustered protein coding sequences to identify, annotate and compare K locus structures, and we propose a novel method for identifying K loci based on full locus information extracted from whole genome sequences.

Results: A total of 134 distinct K loci were identified, including 31 novel types. Comparative analysis of K locus gene content detected 508 unique protein coding gene clusters that appear to reassort via homologous recombination, generating novel K locus types. Extensive nucleotide diversity was detected among the *wzi* and *wzc* genes, both within and between K loci, indicating that current typing schemes

based on these genes are inadequate. As a solution, we introduce *Kaptive*, a novel

software tool that automates the process of identifying K loci from large sets of

Klebsiella genomes based on full locus information.

Conclusions: This work highlights the extensive diversity of *Klebsiella* K loci and the

proteins that they encode. We propose a standardised K locus nomenclature for

Klebsiella, present a curated reference database of all known K loci, and introduce a

tool for identifying K loci from genome data (https://github.com/katholt/Kaptive).

These developments constitute important new resources for the *Klebsiella*

community for use in genomic surveillance and epidemiology.

Key Words:

Klebsiella, capsule, K locus, genomic epidemiology, polysaccharide variation

Background

Klebsiella pneumoniae, *Klebsiella variicola* and *Klebsiella quasipneumoniae* are ubiquitous, encapsulated Gram-negative bacteria. They can be carried asymptomatically in the human gut or nasopharynx [1] but are also opportunistic pathogens, frequently associated with human disease and recognised as a significant threat to global health. Antimicrobial resistance, particularly multi-drug resistance and resistance to the carbapenems, is a major concern. Notably, there are a number of multi-drug resistant clones, which are distributed world-wide and are reported to cause outbreaks of healthcare-associated infections [2,3]. There are also increasing reports of invasive, community-acquired *K. pneumoniae* disease in many Asian countries [4]. While this phenomenon is not yet well understood, it is associated with 'hypervirulent' *K. pneumoniae* strains expressing specific capsular serotypes known as K1, K2 and K5 [5,6].

In order to control the emerging threat of *K. pneumoniae sensu stricto, K. variicola* and *K. quasipneumoniae* (hereafter collectively referred to as *K. pneumoniae* unless otherwise stated), there is an urgent requirement for genome-based surveillance. Recent advances in understanding *K. pneumoniae* population structure [7,8] highlight immense genomic diversity and provide a framework for tracking this pathogen. Useful strategies involve analyses of lineages or multi-locus sequence types in combination with resistance and virulence gene characterisation, e.g. using tools such as SRST2 [9] and BIGSdb [8]. Additionally, there have been successful *K. pneumoniae* outbreak investigations using genomic analysis [3,10]. However, methods for tracking *K. pneumoniae* capsular variation are currently lacking.

The polysaccharide capsule is the outer most layer of the *K. pneumoniae* cell, which

protects the bacterium from desiccation, phage and protist predation [11]. The

capsule is also a key virulence determinant due to its antiphagocytic properties [12–

14]. There are 77 immunologically distinct *K. pneumoniae* capsule types (K-types)

defined by serology, mostly based on work done in the 1950s-70s [15–17]. However,

serological typing requires specialist techniques and reagents not available to most

microbiology laboratories, so it is very rarely applied. Furthermore, between 10%

and 70% of *K. pneumoniae* isolates are serologically non-typeable, either because

they express a novel capsule (most commonly for clinical isolates) or are non-

capsulated [6,18,19].

K. pneumoniae employ a Wzy-dependent capsule synthesis process [11,20] and the

genes required for capsule synthesis and assembly are located at the capsule

polysaccharide synthesis locus (K locus). The K locus varies in length from 10 to 30

kbp [21–26] and includes a set of 'common' genes in the terminal regions, which

encode the core capsule biosynthesis machinery (e.g. *galF, wzi, wza, wzb, wzc*, *gnd*

and *ugd*). The central region of the K locus is highly variable, encoding the capsule-

specific sugar synthesis, processing and export proteins plus the core assembly

components Wzx (flippase) and Wzy (capsule repeat unit polymerase). The *wzx* and

wzy genes are more diverse than those of the other core assembly components and

do not have a fixed position within the K locus [22].

K locus nucleotide sequences and annotations are now available for a large number

of *K. pneumoniae* isolates, including the 77 K-type reference strains [3,21–23,25,27–

29]. Serological K-types are generally defined by distinct sets of genes in the variable

central region of the K locus. This is usually due to the presence of entirely different

sets of protein coding sequences; however two types (K22 and K37) are

distinguished by a single point mutation resulting in a premature stop codon that

affects acetyltransferase function [22].

A number of molecular K-typing schemes have been developed that take advantage

of the conserved K locus structure: restriction fragment length polymorphism ('C-

typing') [30], *wzi* and *wzc* typing [31,32] and capsule-specific *wzy* PCR-based typing

[25,33]. C-typing comprises PCR amplification of a large region of the K locus (from

upstream of *wzi* to within *gnd*), followed by *Hinc*II restriction. In contrast, *wzi* and

wzc typing each comprise PCR amplification and nucleotide sequencing of regions of

a single gene, *wzi* and *wzc* respectively. Within the *wzi* scheme, unique alleles are

associated to specific K-types [32]. Within the *wzc* scheme, K-types are assigned

based on the level of *wzc* nucleotide similarity to a reference sequence, with a

threshold of 94% [31]. These molecular typing methods are less technically

challenging than serological techniques and are more discriminatory [30–32].

However, none of the methods have been widely adopted and regardless of the

method, a substantial proportion of isolates remain non-typeable. As a

consequence, the true extent of *K. pneumoniae* capsule diversity remains unknown.

Here we report the K loci from a collection of 2503 *K. pneumoniae*. We identify 31

novel K loci, and provide evidence that limited additional diversity remains to be

discovered in *K. pneumoniae*. We define a standardised nomenclature for *Klebsiella*

K loci, provide a curated reference database and introduce *Kaptive*, a tool for rapid

identification of reference K loci from genome data, which will greatly facilitate

genomic surveillance efforts and evolutionary investigations of this important

pathogen.

Methods

We obtained a total of 2600 *K. pneumoniae* genomes (2021 publicly available

genomes and 579 novel genomes from a diverse set of isolates collected in

Australia). Sequence reads were generated locally or obtained from the European

Nucleotide Archive (accessions listed in **Table S1, Additional File 1**); 916 genomes

that were publicly available as assembled contigs only were downloaded from

PATRIC [34] and the NCTC3000 project [35]. For isolates sequenced in this study (n =

579) DNA was extracted and libraries prepared using the Nextera® XT 96 barcode

DNA kit and 125 bp paired-end sequence reads were generated on the Illumina

HiSeq 2500 platform.

All paired-end read sets were filtered to remove reads with mean Phred quality

score <30, then assembled *de novo* using SPAdes v3 [36]. Genomes were excluded

from the study if they were duplicate samples, if there was evidence of

contamination or mixed culture measured by: (i) <50% reads mapped to the NTUH-

K2044 reference chromosome (accession: AP006725.1); (ii) the ratio of

heterozygous/homozygous SNP calls compared to the reference chromosome

exceeded 20%; (iii) the total assembly length was >6.5Mb, or >6.0Mb with evidence

of >1% non-*Klebsiella* read contamination as determined by MetaPhlAn [37]; or (v)

the assembly was low quality i.e total length <5Mb.

Existing high quality K locus reference sequences

K locus nucleotide sequences for each of the 77 K-type references and two

serologically non-typeable strains published previously [21–26] were obtained from

GenBank or directly from the authors (accessions in **Table S2, Additional File 2**). A

total of 12 additional K locus sequences had been published prior to the K-type

references [3,27,29,38]; we have compared these loci to those of the 77 K-type

references [21–26] and identified seven that were novel. These seven novel loci plus

17 distinct loci described in our recent survey [28] were added to the non redundant

list of K locus reference sequences, resulting in a total of 103 loci (see **Table S2,**

Additional File 2).

Identification of novel K loci

In order to identify novel K loci we first classified each genome by similarity to

previously known loci. BLASTn [39] was used to search each genome assembly for

sequences with similarity to those of annotated K locus coding sequences (CDS)

usually located between *galF* and *ugd* inclusive (minimum coverage 80%, minimum

identity 50%). Transposase CDS present in the published K locus reference

sequences were excluded from this analysis since they are not K locus specific. Up to

three missing CDS were tolerated for K locus assignment, to allow for assembly

problems and insertion sequence (IS) insertions (see **Figure S1, Additional File 3**).

This approach can successfully distinguish the 77 K-type reference loci (with the

exception of K22 and K37).

Genomes that could not be assigned a K locus were investigated further: BLASTn was

used to identify the *galF* and *ugd* genes within the assembly, and single contig loci

were extracted. The assembly graph viewer Bandage [40] was used to identify K loci

that did not assemble on a single contig or where *galF* and/or *ugd* where missing.

Loci were clustered, with identity and coverage thresholds of 90%, using CD-HIT-EST

[41,42]. A representative sequence from each cluster was annotated with Prokka

[43], using all proteins in the 77 reference K-type loci as the primary annotation

database. Novel K locus sequences were deposited in Genbank (accessions:

LT603702 - LT603735; also included in the Kaptive database at

https://github.com/katholt/Kaptive/).

Recombination in K loci was investigated by aligning nucleotide sequences for the

eight common genes (extracted from the reference annotations) using MUSCLE [44].

This generated a 9944 bp concatenated gene alignment which was used as input for

maximum likelihood (ML) phylogenetic inference with RAxML [45] (best scoring tree

from five runs each of 1000 bootstrap replicates with gamma model of rate

heterogeneity), and recombination analysis using ClonalFrameML [46] (run for 1000

simulations and using the ML phylogeny as the starting tree).

Amino acid clustering

Predicted amino acid sequences of all annotated K locus coding regions were

translated from the DNA sequences using BioPython and clustered with CD-HIT

[41,42] (90%, 80%, 70%, 60%, 50%, 40% identity). We explored the co-occurrence of

predicted protein clusters present in three or more K loci each (excluding the

common proteins and the initiating glycosyltransferases, WbaP and WcaJ, n = 115

clusters for analysis): Pairwise Jaccard similarity scores were calculated as $J(A, B) =$

$A \cap B / A \cup B$ and were used to draw a weighted edge graph with the igraph R

package v 1.0.1 [47]. A weight threshold was determined empirically as 0.61 and all

edges for which $J < 0.61$ were removed.

Wzc and wzi nucleotide sequence determination

We characterised *wzi* and *wzc* sequence diversity and explored the association with

K loci. We used SRST2 [9] to determine *wzi* alleles defined in the *K. pneumoniae*

BIGSdb [48]. BLASTn was used to determine alleles in genomes available only as

assemblies. Novel alleles were submitted to the *K. pneumoniae* BIGSdb for official

designation. *Wzc* sequences were extracted from genome assemblies by BLASTn

search against a database of previously published alleles [31]. Sequences were

aligned with MUSCLE [44] and pairwise nucleotide divergences calculated.

Kaptive, a tool for identification of K loci in genome data

We developed an extended procedure for identification and assessment of full-

length K loci among bacterial genomes based on BLAST analysis of assemblies. The

procedure has been automated and is implemented in a freely available open source

software tool, *Kaptive* (https://github.com/katholt/Kaptive). For full details see

Additional File 3, Figures S2 and S3.

Comparison of Kaptive, wzi and wzc typing results

Kaptive, wzi and *wzc* typing were applied to the 86 genomes that had matched

serological typing information available (**Table S3, Additional File 4**). *Wzi* alleles

were determined as described above, and used to predict serotypes by comparison

to the *K. pneumoniae* BIGSdb. *Wzc* sequences were extracted as above and genomes

were assigned to serotypes if the sequence was <6% divergent from the

corresponding reference [31].

Results

We analysed K loci in a final dataset comprising 2503 high quality genomes, including

2298 *K. pneumoniae sensu stricto*, 144 *K. variicola*, 57 *K. quasipneumoniae* and four

unclassified *Klebsiella* spp. (**Table 1** and **Table S1, Additional File 1**). Also included

were 10 publicly available genomes representing the more distantly related

Klebsiella oxytoca (**Table S4, Additional File 3**), as we hypothesised that these

organisms may share capsule synthesis loci with *K. pneumoniae*. Isolates were

collected between 1932 and 2014 (see **Figure S4, Additional File 3**) and from eight

different geographic regions spanning six continents (see **Figure S5, Additional File**

3).

Table 1: *K. pneumoniae* genomes investigated in this study

Dataset	Count	Reference	Notes
Bialek-Davenet *et al.*	33	[8]	Investigation of multi-drug resistant and hypervirulent clones
Bowers *et al.*	160	[49]	Isolates mostly of clonal group 258
Davis *et al.*	77	[50]	Isolates from human UTI and animal meats in Arizona, USA
Deleo *et al.*	69	[29]	Isolates of clonal group 258
Ellington	185	[51]	Multidrug resistant isolates from a hospital in Cambridge, UK
Holt *et al.*	274	[7]	Global diversity study
Lee *et al.*	27	[52]	Isolates from pyogenic liver abscess disease, Singapore
NCTC3000	81	[35]	Isolates from the Public Health England NCTC reference collection
PATRIC	811	[34]	Genome assemblies submitted to GenBank
Stoesser *et al.* 2013	69	[53]	Isolates from health-care associated infections, Oxford, UK
Stoesser *et al.* 2014	55	[54]	Isolates collected during an outbreak in Nepal
Struve *et al.*	67	[55]	Predominantly isolates of clonal group 23
The *et al.*	76	[3]	Isolates from two outbreaks in Nepal
Wand *et al.*	35	[56]	Historical isolate collection
Novel isolates	484	This study	Diverse collection of Australian hospital surveillance isolates

A total of 1371 *K. pneumoniae* genomes could be putatively assigned to 63 of the 77

K-type reference loci, and a further 918 to one of the other 25 previously published K

loci. Among the remaining 213 genomes, 106 were assigned to 29 novel K loci,

bringing the total number of known K loci to 132. A further six genomes harboured

deletion mutants of known K loci, two had IS variants of known loci and one had an

IS variant of a novel locus (see below). For 93 genomes (3.7%), no K locus could be

determined; however we found evidence of the presence of three or more K locus-

associated genes in all such genomes and consider the lack of assignment was likely

attributable to low quality sequence data (low read depth and/or fragmented

assembly) rather than complete K locus deletion. Complete details of K locus

assignments to all genomes are given in **Table S1** in **Additional File 1**.

A locus previously associated with K66 was identified in one *K. oxytoca* genome and

the K74 locus in four *K. oxytoca* genomes; these matches were very close to the *K.*

pneumoniae reference sequences (100% coverage and >93% nucleotide identity in

all cases). Two novel K loci were identified in three *K. oxytoca* genomes, increasing

the total known *Klebsiella* K loci to 134 (**Table S4, Additional File 3**).

We estimated the extent to which we had captured the repertoire of K locus

diversity in the *K. pneumoniae* population (**Figure 1**). The rarefaction curves were

estimated from (i) the full genome set for which K loci were assigned (n = 2410; grey

lines in **Figure 1**); (ii) a 'non-redundant' genome set from which highly biased sub-

samples such as outbreaks were removed (n = 1081; blue lines in **Figure 1**); and (iii)

genomes from the non-redundant set representing each of the distinct species, *K.*

pneumoniae sensu stricto, K. variicola and *K. quasipneumoniae* (black lines in **Figure**

1). In comparison to that for the full genome set (grey), the non-redundant (blue)

curve better represents the true diversity of the *K. pneumoniae sensu lato*

population. Note that neither reached the total number of known K loci, since 13 of

the serologically defined K loci [22] were not represented at all in our 2503 genomes.

The rarefaction curves for each of the three *Klebsiella* species within the non-

redundant dataset were highly similar to one another, indicating similar levels of

capsule diversity within each species (**Figure 1**). There was no strong evidence of

species specificity: across our entire genome collection 46 distinct K loci were

identified only among *K. pneumoniae sensu stricto*, while three and two were

identified only among *K. variicola* and *K. quasipneumoniae*, respectively, however

these differences are likely an artefact of the much larger sample size currently

available for *K. pneumoniae sensu stricto*.

K locus nomenclature

We used a standardised K locus nomenclature based on that proposed for

Acinetobacter baumannii [57]. Each distinct *Klebsiella* K locus was designated as KL

(K locus) and a unique numeric identifier. The K-type reference K loci were assigned

the same numeric identifier as the corresponding K-type, for example K1 is encoded

by the KL1 locus. K loci for which capsule types have not yet been phenotypically

defined were assigned identifiers starting from 101 (note KL101 and KL102

correspond to those previously named as KN1 and KN2).

K loci with IS insertions disrupting the region were distinguished from orthologous

IS-free variants by using -1, -2. This nomenclature was consistently applied to the 10

K-type reference K loci published previously that include one or more ISs

[21,22,25,26]. Deletion variants derived from a known K locus were given the suffix -

D1, -D2, etc.

K locus reference database

We curated a K locus reference database of complete annotated K locus sequences

for all of the 134 loci (**Table S2, Additional File 2**). Where possible sequences were

included at their full length, from the start of *galF* to the end of *ugd*. Where a

previously published sequence did not span the full length of the locus or contained

an IS, we substituted the complete, IS-free K locus sequence from a genome in our

collection if available (39 of 51). Where no naturally occurring IS-free variants were

available, we manually generated an IS-free synthetic sequence (**Table S2,**

Additional File 2). IS-free sequences are included in a primary K locus reference

database, while all available IS or deletion variant K locus references are included in

an accompanying variant database, both available at

https://github.com/katholt/Kaptive.

K locus structures

All of the novel K loci identified in this study conformed to the common structure

described previously (**Figure S6**) [20–22]. We also identified six deletion variants

within our genome collection (KL5-D1, KL20-D1, KL30-D1, KL62-D1, KL106-D1 and

KL107-D1). Each putative deletion variant was missing several common genes but

the remaining regions showed a high degree of similarity to other apparently

complete K loci, which we suggest represent the ancestral forms. Isolate NCTC10004

(recorded as serotype K11 in the UK National Culture Type Collection) and four other

genomes carried a K locus that was nearly identical to the previously published K11 K

locus reference sequence [22]. However, the latter lacked the essential *wzx* gene

plus two other neighbouring genes, and was not identified among any other

genomes. We assume the NCTC10004 locus represents the full length KL11 locus and

designate the original K11 reference as KL11-D1 (note it is unclear whether the

original sequenced reference isolate had retained the ability to produce a capsule,

since the serological typing was performed decades earlier [22]). In four of the

deletion variants the deleted region was replaced by an IS, which may have

mediated the deletion.

Of the other IS related variants, KL157-1 contained an IS*903* family IS without an

obvious deletion. In addition, we identified two novel IS variants of K-type reference

loci (KL15-1 and KL22-1), and five IS-free variants of K-type reference loci (KL3, KL6,

KL38, KL57, KL81), plus one other previously published K locus (KL103). In total,

seven IS insertions were associated with neither a deletion nor a rearrangement

event. In contrast, the KL22-1 locus included a translocation of part of the

lipopolysaccharide (LPS) locus to the centre of the K locus, plus an inversion of the 3'

portion of the K locus (**Figure 2**). The translocated and inverted regions were bound

at each end by a copy of IS*Kpn26*.

ClonalFrameML analysis of the nucleotide sequences of common K locus genes

identified a high number of putative recombination events (n=382) between the

reference K loci. These events were not distributed equally across the nucleotide

alignment (**Figure 3A**); rather the genes closest to the central variable region of the K

locus were affected by a greater number of recombination events compared to

those at the ends of the locus.

Variation in K locus gene content

A total of 2675 predicted proteins from 134 complete K loci were clustered at

various identity levels (see **Methods**), resulting in 1496 to 508 clusters. As the

identity threshold was reduced the number of clusters continued to fall and showed

no signs of stabilising, even between 50% and 40% identity (**Figure S7**), and we

believe the latter is a lower bound for sensible comparison. At 40% identity, the core

capsule assembly proteins GalF, Wzi, Wza, Wzb, Gnd and Ugd each formed a single

cluster, and were present in nearly all loci (**Figure 3**). The Wzc sequences clustered

into two groups, and each locus encoded one Wzc protein (except KL50). In contrast,

Wzx (flippase) clustered into 42 groups and Wzy (capsule repeat unit polymerase)

clustered into 83 groups, highlighting the extreme diversity of these proteins

compared to the other core capsule assembly machinery proteins (**Figure 3**).

There were 374 clusters among the remaining proteins, almost all of which were

associated with sugar synthesis and processing (**Figure 3**). The initiating sugar

transferase proteins WbaP (undecaprenyl-phosphate galactosephosphotransferase)

and WcaJ (undecaprenyl-phosphate glucose phosphotransferase) were grouped into

two clusters. These proteins are considered essential for capsule synthesis.

Concordantly each locus encoded a single protein from one of these two clusters.

RmlB, RmlA, RmlD and RmlC, which are associated with synthesis and processing of

rhamnose and typically encoded together in a single operon, were each represented

by a single cluster. Similarly, the mannose synthesis and processing proteins, ManC

and ManB, were grouped into a single cluster each. The associated operons *rmlBADC*

and *manCB* were present in 55 and 73 K loci, respectively (14 loci contained both

operons, **Figure 3**). In contrast, 360 of the remaining 366 protein clusters were

present in fewer than ten K loci each (**Figure 3E**).

Co-occurrence analysis identified 18 correlated groups of K locus proteins, ranging in

size from two to five protein clusters (pairwise Jaccard similarity ≥0.61 for all pairs in

the group, **Figure 4** and **Table S5, Additional File 5**). One group included the four Rml

protein clusters; interestingly this group also included a WcaA glycosyltransferase,

which was present in 67.3% of *rmlBADC*-containing K loci and no *rmlBADC*-negative

loci (X-squared = 70.09, p-value < 2.2e-16 by two-sided proportion test). Similarly

another group included the ManCB proteins and the putative mannosyl transferase,

WbaZ, which was present in 65.8% of *manCB*-containing K loci and one *manCB*-

negative locus (X-squared = 56.159, p-value = 6.683e-14 by two-sided proportion

test). In addition, several groups included proteins for which the associated genes

were located sequentially in their K loci (e.g. *wckG*, *wckH* and *wzx* in KL12, KL29 and

KL42) consistent with linked gene transfer.

Diversity of wzc and wzi gene sequences

We confidently assigned *wzi* alleles to 2461 *K. pneumoniae* genomes, including 390

distinct alleles, 218 of which were novel. Median pairwise nucleotide divergence was

7%. Among the non-redundant genome set there were 54 *wzi* alleles represented by

at least five genomes and of these, 15 (28%) were associated with more than one K

locus type (**Table S1, Additional File 1**). Much of the *wzi* allelic variation appeared to

result from accumulation of mutations within K loci. Among the 67 K loci for which

we had ≥5 representative sequences, 64 (95.5%) were associated with two or more

wzi alleles, and there was a general trend towards increasing *wzi* allelic diversity with

increasing K locus representation (**Figure 5**).

We extracted *wzc* sequences from 1041 of 1082 genomes in the non-redundant

genomes set (**Figure 6**). In general, genomes sharing the same K locus (n=6262

pairwise observations) showed lower *wzc* nucleotide divergence than those with

different K loci (n=491,775 pairwise observations), but the distributions overlapped

substantially (**Figure 6**). Notably, there were five distinct combinations of K loci for

which one or more pairs harboured *wzc* sequences that were <6% divergent (the

cut-off for K-type assignment as described in [31]; KL1 and KL112, KL9 and KL45,

KL15 and KL52, KL30 and KL104, KL40 and KL135). Conversely, some K loci (KL45,

KL112) had more than 25% *wzc* nucleotide diversity between representatives of the

same K locus.

Kaptive – <u>ca</u>psule locus (K locus) <u>t</u>yping and <u>v</u>ariant <u>e</u>valuation from genome data

To facilitate easy identification of K loci from genome assemblies, we developed the

command-line software tool *Kaptive,* which is an extension of the analysis procedure

described above, as shown in **Figure 7** (also see **Additional File 3**). We used *Kaptive*

with our primary *Klebsiella* K locus reference database to rapidly type the K loci in

our collection of 2503 *Klebsiella* genomes, and obtained confident K locus calls for

2412 genomes (96.4%, see **Additional File 3** for further details).

We compared the K locus calls from *Kaptive*, *wzc* and *wzi* typing to serological typing

results for 86 isolates for which both genome and serology data were available

([7,18,35], see **Table S3, Additional File 4**). Five of six isolates that were non-

typeable by serological techniques were identified by *Kaptive* as carrying KL16, KL54,

KL81, KL111 and KL149. The KL16, KL54 and KL81 calls were in agreement with *wzc*

and *wzi* typing results; the other two K loci were not present in the *wzi* or *wzc*

schemes and so were not typeable by those methods. Among the 80 serologically

typeable isolates, the three molecular methods were generally in agreement with

one another, although concordance with recorded phenotypes was quite low (65-

74%, **Table S3**). Call rates were highest for *Kaptive* (95%), followed by *wzc* (89%) and

wzi typing (75%).

Discussion

The number of distinct *Klebsiella* K loci (now 134) is striking and exceeds that

described for capsule synthesis loci in other bacterial species such as *A. baumannii,*

Streptococcus pneumoniae and *Neisseria meningitidis.* Furthermore, the diversity is

an order of magnitude greater than that recently described for *Klebsiella* LPS, the

other major *Klebsiella* surface antigen [28]. This suggests that the K locus is subject

to strong diversifying selection. Given that these bacteria are not obligate pathogens

and are ubiquitous in non-host-associated environments [58,59], it seems likely that

the factors driving selection are not immune pressures from humans or other hosts,

but may include phage and/or protist predation.

Two novel K loci were identified from *K. oxytoca*, a close relative of *K. pneumoniae*.

The KL66 and KL74 K-type reference loci were also identified among *K. oxytoca*

genomes. Little is known about *K. oxytoca* capsules, though a report from Japan in

2012 identified several other *K. pneumoniae*-associated capsules among *K. oxytoca*

isolates from blood and bile infections [60]. Together these findings indicate that *K.*

oxytoca is able to exchange genetic material with *K. pneumoniae.* Therefore, *K.*

oxytoca represents a potential reservoir of virulence, drug resistance and other

genes for *K. pneumoniae,* and warrants greater research attention.

Our analysis confirms there are strong constraints on the structure of K loci, which

generally include *galF, cpsACP, wzi, wza, wzb* and *wzc* at the 5' end, *gnd* and *ugd* at

the 3' end, and a highly variable set of genes in the centre (**Figure 3**). Our data also

reveal the extensive diversity of proteins encoded in the variable central region, with

499 unique proteins identified across the 134 K loci. These genes ranged in

frequency from 0.7% to 54.7% of the K loci. Among those represented in at least

three K loci, approximately half co-occurred in groups ranging from two to five

genes.

The molecular evolutionary events driving K locus diversification are not yet well

understood, but likely include a combination of point mutation, IS-mediated

rearrangements, and homologous recombination within the locus, resulting in the

mosaic structure summarised in **Figure 3**. It has been shown, in both *A. baumannii*

[61,62] and *S. pneumoniae* [63], that recombination within the capsule synthesis

locus can drive capsule exchange between distinct clones. We recently speculated

that this may also be true for *K. pneumoniae* [27] and the recombination analysis

presented here supports this theory. The genes closest to the central variable region

of the K locus (i.e. *wzb, wzc* and *gnd*), showed evidence of the greatest number of

recombination events, consistent with the hypothesis that they act as regions of

homology for recombination events that shuffle the central region of the locus.

The prediction of capsule phenotypes from genome data is complex, as capsule

expression is a highly regulated process that involves loci outside the K locus region

[64], and so presence of an identical K locus sequence does not guarantee an

identical phenotype. However it is likely that K loci encoding distinct sets of proteins

are associated with distinct capsule phenotypes, as is the case for the vast majority

of K-type reference strains [22]. Therefore, our data suggest that there are at least

134 distinct *Klebsiella* capsule types. Note that this is a lower bound estimate since

there are likely additional K loci in the wider population that were not in the current

genome collection, and our analysis did not capture differences that may arise from

point mutations and small-scale insertions or deletions (e.g. in the case of K22 and

K37 described previously [22]). Furthermore, while we did not attempt to thoroughly

characterise IS variants, several such variants were apparent. The potential

functional impacts of IS insertions likely vary depending on their location in the

locus, but may include up-regulation, loss of capsule production and/or more subtle

changes in sugar structures [65–67]. However, functional studies are required to

understand these effects and to improve the prediction of phenotypes.

Serological typing of *Klebsiella* isolates is notoriously difficult and rarely performed.

We were able to compare genotypes (whole-locus typing using *Kaptive*, as well as

wzi and *wzc* typing schemes) with phenotypes on just 86 isolates for which both

sequences and serotypes were available. Of the 19 discordant genotype vs

phenotype results, two were due to deletion variants and were resolved by running

Kaptive with the K locus variants database. Interestingly, one of these isolates was

non-typeable by serology, *wzi* or *wzc* typing, but recognized as a specific K locus

deletion variant by *Kaptive*. This highlights a benefit of our whole-locus typing

approach; it provides epidemiologically relevant information even when the K locus

is interrupted. Another isolate was serologically typed as K54 but genotyped by

Kaptive as KL113, which has sequence homology with KL54 (>84% nucleotide

identity over 76% of the locus) and may encode a serologically similar or cross-

reacting capsule. The other cases of discordance had no obvious explanation,

however it is likely that some result from serological typing errors or from mutations

arising during subculture (as identified for the K11 reference isolate above), neither

of which we were able to check. Some discordance may also be due to unpredictable

serological cross-reactions.

Given the problems with serotyping and the comparative robustness and

widespread access to genome sequencing, we anticipate that genotyping will remain

the preferred method for tracking capsular diversity in *Klebsiella*. Due to the

extensive diversity and potential for ongoing evolution, we strongly advocate for

classification based on complete, or near complete K locus sequences, rather than

single genes such as *wzi* or *wzc,* which can be misled by substitutions and horizontal

gene transfer. *Kaptive* analyses the full-length K locus nucleotide sequences and

assesses the presence of all K locus associated genes by protein BLAST search, thus

the approach is resilient to spurious results that may arise due to sequence

divergence. Furthermore, the information provided allows users to determine

confidence in the results and to identify putative novel K loci or variants of known

loci if desired (see **Additional File 2**).

Conclusions

We report an investigation of K loci among a large collection of 2513 *Klebsiella*

genomes. We identified 31 novel K loci, increasing the total number of known loci to

134, almost twice the number of serologically defined K-types. We defined a

standardised *Klebsiella* K locus nomenclature and developed a curated reference

database, which captures the majority of the extensive diversity in the *K.*

pneumoniae population. Lastly, we developed a simple program, *Kaptive,* for the

detection of reference K loci from genome assemblies. These new resources will

greatly facilitate evolutionary investigations and genomic surveillance efforts for this

and other important bacterial pathogens.

List of Abbreviations

K-type: capsule type; K locus: capsule synthesis locus; IS: insertion sequence; CDS:

coding sequence; LPS: lipopolysaccharide

Additional Files

Additional file 1 (Excel spreadsheet, xlsx)

Table S1. *K. pneumoniae* genome data analysed in this study. Accession numbers, K

locus designations and summarised *Kaptive* typing results are provided.

Additional file 2 (Excel spreadsheet, xlsx)

Table S2. *Klebsiella* K locus references, accession numbers and isolate names.

Additional file 3 (Word document, docx)

Supplementary Methods and Results

Table S4. *K. oxytoca* genomes analysed in this study and K locus designations.

Figures S1 – S7.

Additional file 4 (Excel spreadsheet, xlsx)

Table S3. *Kaptive*, *wzi, wzc* and serological typing results for 86 *K. pneumoniae*

genomes for which serological typing information were available.

Additional file 5 (Excel spreadsheet, xlsx)

Table S5. Jaccard similarity scores for 115 K locus protein clusters for which the

associated genes were present in at least three K loci.

Declarations

Availability of data and material

The datasets generated and/or analysed during the current study are available in the

European Nucleotide Archive and/or PATRIC genome database, accession numbers

are listed in **Table S1**. Novel K locus nucleotide sequences have been deposited in

GenBank (accessions listed in **Table S2**) and are also distributed together with our re-

annotated and curated set of all known K loci at https://github.com/katholt/Kaptive.

Competing interests

None declared.

Funding

This work was funded by the NHMRC of Australia (Fellowship #1061409 and Project

#1043822 to K.E.H.) and Wellcome Trust (Grant 098051 to the W.T.S.I.).

Authors' Contributions

KLW and KEH designed the study, collected and analysed data, designed the *Kaptive*

tool and wrote the manuscript. RRW designed and implemented the *Kaptive* tool

and wrote the manuscript. CG and AJ generated serological and sequence data for

Australian isolates. RF and NT contributed to data analysis and interpretation. All

authors read and approved the final manuscript.

References

1. Lin YT, Wang YP, Wang F Der, Fung CP. Community-onset *Klebsiella pneumoniae*

pneumonia in Taiwan: Clinical features of the disease and associated microbiological

characteristics of isolates from pneumonia and nasopharynx. Front Microbiol.

2015;6:1–8.

2. Munoz-Price LS, Poirel L, Bonomo RA, Schwaber MJ, Daikos GL, Cormican M, et al.

Clinical epidemiology of the global expansion of *Klebsiella pneumoniae*

carbapenemases. Lancet Infect Dis. 2013;13:785–96.

3. The HC, Karkey A, Thanh DP, Boinett CJ, Cain AK, Ellington M, et al. A high-

resolution genomic analysis of multidrug- resistant hospital outbreaks of *Klebsiella*

pneumoniae. EMBO Molec Med. 2015;7:227–39.

4. Siu LK, Yeh KM, Lin JC, Fung CP, Chang FY. *Klebsiella pneumoniae* liver abscess: A

new invasive syndrome. Lancet Infect Dis. 2012;12:881–5.

5. Fung CP, Hu BS, Chang FY, Lee SC, Kuo BI, Ho M, et al. A 5-year study of the

seroepidemiology of *Klebsiella pneumoniae*: high prevalence of capsular serotype K1

in Taiwan and implication for vaccine efficacy. J Infect Dis. 2000;181:2075–9.

6. Tsay R-W, Siu LK, Fung C-P, Chang F-Y. Characteristics of bacteremia between

community-acquired and nosocomial *Klebsiella pneumoniae* infection. Arch Intern

Med. 2002;162:1021–7.

7. Holt KE, Wertheim H, Zadoks RN, Baker S, Whitehouse CA, Dance D, et al. Genomic

analysis of diversity, population structure, virulence, and antimicrobial resistance in

Klebsiella pneumoniae, an urgent threat to public health. Proc Natl Acad Sci U S A.

2015;112:E3574–81.

8. Bialek-Davenet S, Criscuolo A, Ailloud F, Passet V, Jones L, Delannoy-Vieillard AS,

et al. Genomic definition of hypervirulent and multidrug-resistant *Klebsiella*

pneumoniae clonal groups. Emerg Infect Dis. 2014;20:1812–20.

9. Inouye M, Dashnow H, Raven L, Schultz MB, Pope BJ, Tomita T, et al. SRST2: Rapid

genomic surveillance for public health and hospital microbiology labs. Genome Med.

2014;6:90.

10. Snitkin ES, Zelazny AM, Thomas PJ, Stock F, Program NCS, Henderson DK, et al.

Tracking a hospital outbreak of carbapenem-resistant *Klebsiella pneumoniae* with

whole-genome sequencing. Sci Transl Med. 2012;4:148ra116.

11. Whitfield C. Biosynthesis and assembly of capsular polysaccharides in *Escherichia*

coli. Annu Rev Biochem. 2006;75:39–68.

12. Allen PM, Fisher D, Saunders JR, Hart CA. The role of capsular polysaccharide

K21b of *Klebsiella* and of the structurally related colanic-acid polysaccharide of

Escherichia coli in resistance to phagocytosis and serum killing. J Med Microbiol.

1987;24:363–70.

13. Kabha K, Nissimov L, Athamna A, Keisari Y, Parolis H, Parolis LA, et al.

Relationships among capsular structure, phagocytosis, and mouse virulence in

Klebsiella pneumoniae. Infect Immun. 1995;63:847–52.

14. Lee CH, Chang CC, Liu JW, Chen RF, Yang KD. Sialic acid involved in

hypermucoviscosity phenotype of *Klebsiella pneumoniae* and associated with

resistance to neutrophil phagocytosis. Virulence 2014;5:673–9.

15. Edwards PR, Fife MA. Capsule types of *Klebsiella*. J Infect Dis. 1952;91:92–104.

16. Edmunds PN. Further *Klebsiella* capsule types. J Infect Dis. 1954;94:65–71.

17. Ørskov IDA, Fife-Asbury MA. New *Klebsiella* capsular antigen, K82, and the

deletion of five of those previously assigned. Int J Syst Bacteriol. 1977;27:386–7.

18. Jenney AW, Clements A, Farn JL, Wijburg OL, McGlinchey A, Spelman DW, et al.

Seroepidemiology of *Klebsiella pneumoniae* in an Australian tertiary hospital and its

implications for vaccine development. J Clin Microbiol. 2006;44:102–7.

19. Cryz SJ, Mortimer PM, Mansfield V, Germanier R. Seroepidemiology of *Klebsiella*

bacteremic isolates and implications for vaccine development. J Clin Microbiol.

1986;23:687–90.

20. Rahn A, Drummelsmith J, Whitfield C. Conserved organization in the *cps* gene

clusters for expression of *Escherichia coli* group 1 K antigens: relationship to the

colanic acid biosynthesis locus and the *cps* genes from *Klebsiella pneumoniae*. J

Bacteriol. 1999;181:2307–713.

21. Shu HY, Fung CP, Liu YM, Wu KM, Chen YT, Li LH, et al. Genetic diversity of

capsular polysaccharide biosynthesis in *Klebsiella pneumoniae* clinical isolates.

Microbiology. 2009;155:4170–83.

22. Pan Y-J, Lin T-L, Chen C-T, Chen Y-Y, Hsieh P-F, Hsu C-R, et al. Genetic analysis of

capsular polysaccharide synthesis gene clusters in 79 capsular types of *Klebsiella* spp.

Nat Sci Rep. Nature Publishing Group; 2015;5:15573.

23. Chuang YP, Fang CT, Lai SY, Chang SC, Wang JT. Genetic determinants of capsular

serotype K1 of *Klebsiella pneumoniae* causing primary pyogenic liver abscess. J Infect

Dis. 2006;193:645–54.

24. Arakawa Y, Wacharotayankun R, Nagatsuka T, Ito H, Kato N, Ohta M. Genomic

organization of the *Klebsiella pneumoniae cps* region responsible for serotype K2

capsular polysaccharide synthesis in the virulent strain Chedid. J Bact.

1995;177:1788–96.

25. Pan YJ, Fang HC, Yang HC, Lin TL, Hsieh PF, Tsai FC, et al. Capsular polysaccharide

synthesis regions in *Klebsiella pneumoniae* serotype K57 and a new capsular

serotype. J Clin Microbiol. 2008;46:2231–40.

26. Fevre C, Passet V, Deletoile A, Barbe V, Frangeul L, Almeida AS, et al. PCR-based

identification of *Klebsiella pneumoniae* subsp. rhinoscleromatis, the agent of

rhinoscleroma. PLoS Negl Trop Dis. 2011;5:e1052.

27. Wyres KL, Gorrie C, Edwards DJ, Wertheim HFL, Hsu LY, Van Kinh N, et al.

Extensive capsule locus variation and large-scale genomic recombination within the

Klebsiella pneumoniae clonal group 258. Genome Biol Evol. 2015;7:1267–79.

28. Follador R, Heinz E, Wyres KL, Ellington MJ, Kowarik M, Holt KE, et al. The

diversity of *Klebsiella pneumoniae* surface polysaccharides. MGen. 2016;Online

ahead of print.

29. Deleo FR, Chen L, Porcella SF, Martens CA, Kobayashi SD, Porter AR, et al.

Molecular dissection of the evolution of carbapenem-resistant multilocus sequence

type 258 *Klebsiella pneumoniae*. Proc Natl Acad Sci U S A. 2014;111:4988–93.

30. Brisse S, Issenhuth-Jeanjean S, Grimont PA. Molecular serotyping of *Klebsiella*

species isolates by restriction of the amplified capsular antigen gene cluster. J Clin

Microbiol. 2004;42:3388–98.

31. Pan YJ, Lin TL, Chen YH, Hsu CR, Hsieh PF, Wu MC, et al. Capsular types of

Klebsiella pneumoniae revisited by *wzc* sequencing. PLoS One 2013;8:e80670.

32. Brisse S, Passet V, Haugaard AB, Babosan A, Kassis-Chikhani N, Struve C, et al. *wzi*

gene sequencing, a rapid method for determination of capsular type for *Klebsiella*

strains. J Clin Microbiol. 2013;51:4073–8.

33. Yu W-L, Fung C-P, Ko W-C, Cheng K-C, Lee C-C, Chuang Y-C. Polymerase chain

reaction analysis for detecting capsule serotypes K1 and K2 of *Klebsiella pneumoniae*

causing abscesses of the liver and other sites. J Infect Dis. 2007;195:1235–6.

34. Wattam AR, Abraham D, Dalay O, Disz TL, Driscoll T, Gabbard JL, et al. PATRIC,

the bacterial bioinformatics database and analysis resource. Nucleic Acids Res.

2014;42:581–91.

35. NCTC3000 Project. Wellcome Trust Sanger Inst. website. 2016

36. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al.

SPAdes: a new genome assembly algorithm and its applications to single-cell

sequencing. J Comput Biol. 2012;19:455–77.

37. Segata N, Waldron L, Ballarini A, Narasimhan V, Jousson O, Huttenhower C.

Metagenomic microbial community profiling using unique clade-specific marker

genes. Nat Methods. 2012;9:811–4.

38. Chen L, Mathema B, Pitout JD, DeLeo FR, Kreiswirth BN. Epidemic *Klebsiella*

pneumoniae ST258 Is a hybrid strain. MBio 2014;5:e01355–14.

39. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al.

BLAST+: architecture and applications. BMC Bioinformatics 2009/12/17 ed.

2009;10:421.

40. Wick RR, Schultz MB, Zobel J, Holt KE. Bandage: interactive visualization of *de*

novo genome assemblies. Bioinformatics. 2015;31:3350–2.

41. Fu L, Niu B, Zhu Z, Wu S, Li W. CD-HIT: Accelerated for clustering the next-

generation sequencing data. Bioinformatics. 2012;28:3150–2.

42. Li W, Godzik A. CD-Hit: A fast program for clustering and comparing large sets of

protein or nucleotide sequences. Bioinformatics. 2006;22:1658–9.

43. Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics 2014;

5;30:2068-9.

44. Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high

throughput. Nucleic Acids Res. 2004;32:1792–7.

45. Stamatakis A. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses

with thousands of taxa and mixed models. Bioinformatics. 2006;22:2688–90.

46. Didelot X, Wilson DJ. ClonalFrameML: Efficient inference of recombination in

whole bacterial genomes. PLoS Comp Biol. 2015;11:e1004041.

47. Csardi G, Nepusz T. The igraph software package for complex network research.

InterJournal 2006;Complex Sy:1695.

48. Institut Pasteur. *Klebsiella pneumoniae* BIGSdb 2016 [cited 2016 Apr 1]. Available

from: http://bigsdb.web.pasteur.fr/klebsiella/klebsiella.html

49. Bowers JR, Kitchel B, Driebe EM, MacCannell DR, Roe C, Lemmer D, et al.

Genomic analysis of the emergence and rapid global dissemination of the clonal

group 258 *Klebsiella pneumoniae* pandemic. PLoS One 2015;10:e0133727.

50. Davis GS, Waits K, Nordstrom L, Weaver B, Aziz M, Gauld L, et al. Intermingled

Klebsiella pneumoniae populations between retail meats and human urinary tract

infections. Clin Infect Dis 2015;61:892–9.

51. Ellington MJ. *Klebsiella pneumoniae* collection from Cambridge University

Hospitals NHS Foundation Trust. 2016;Manuscript in preparation.

52. Lee R, Molton JS, Wyres KL, Gorrie C, Wong J, Hoh CH, et al. Differential host

susceptibility and bacterial virulence factors driving *Klebsiella* liver abscess in an

ethnically diverse population. Sci Rep. 2016;In press.

53. Stoesser N, Batty EM, Eyre DW, Morgan M, Wyllie DH, Del Ojo Elias C, et al.

Predicting antimicrobial susceptibilities for *Escherichia coli* and *Klebsiella*

pneumoniae isolates using whole genomic sequence data. J Antimicrob Chemother.

2013 [cited 2013 Jun 1];68:2234–44.

54. Stoesser N, Giess A, Batty EM, Sheppard AE, Walker AS, Wilson DJ, et al. Genome

sequencing of an extended series of NDM-producing *Klebsiella pneumoniae* isolates

from neonatal infections in a Nepali hospital characterizes the extent of community-

versus hospital- associated transmission in an endemic setting. Antimicrob Agents

Chemother. 2014;58:7347–57.

55. Struve C, Roe CC, Stegger M, Stahlhut SG, Hansen DS, Engelthaler DM, et al.

Mapping the evolution of hypervirulent *Klebsiella pneumoniae*. MBio. 2015;6:1–12.

56. Wand ME, Baker KS, Benthall G, McGregor H, McCowen JWI, Deheer-Graham A,

et al. Characterization of pre-antibiotic era *Klebsiella pneumoniae* Isolates with

respect to antibiotic/disinfectant susceptibility and virulence in *Galleria mellonella*.

Antimicrob Agents Chemother. 2015;59:3966–72.

57. Kenyon JJ, Hall RM. Variation in the complex carbohydrate biosynthesis loci of

Acinetobacter baumannii genomes. PLoS One. 2013;8:e62160.

58. Podschun R, Ullmann U. *Klebsiella* spp. as nosocomial pathogens: epidemiology,

taxonomy, typing methods, and pathogenicity factors. Clin Microbiol Rev.

1998;11:589–603.

59. Bagley ST. Habitat association of *Klebsiella* species. Infect Contr. 1985;6:52–8.

60. Ishihara Y, Yagi T, Mochizuki M, Ohta M. Capsular types, virulence factors and

DNA types of Klebsiella oxytoca strains isolated from blood and bile. Kansenshogaku

Zasshi. 2012;86:1221–126.

61. Holt K, Kenyon JJ, Hamidian M, Schultz MB, Pickard DJ, Dougan G, et al. Five

decades of genome evolution in the globally distributed, extensively antibiotic

resistant *Acinetobacter baumannii* global clone 1. MGen. 2016;

62. Schultz MB, Thanh DP, Do Hoan NT, Wick RR, Ingle DJ, Hawkey J, et al. Repeated

local emergence of carbapenem resistant *Acinetobacter baumannii* in a single

hospital ward. MGen. 2016;1–15.

63. Wyres KL, Lambertsen LM, Croucher NJ, McGee L, von Gottberg A, Liñares J, et al.

Pneumococcal capsular switching: An historical perspective. J Infect Dis.

2013;207:439-49.

64. Hsu CR, Lin TL, Chen YC, Chou HC, Wang JT. The role of *Klebsiella pneumoniae*

rmpA in capsular polysaccharide synthesis and virulence revisited. Microbiology.

2011;157:3446–57.

65. Salter SJ, Gould KA, Lambertsen LM, Hanage WP, Antonio M, Turner P, et al.

Variation at the capsule locus, *cps*, of mistyped and non-typeable *Streptococcus*

pneumoniae isolates. Microbiology. 2012;158:1560–9.

66. Bentley SD, Aanensen DM, Mavroidi A, Saunders D, Rabbinowitsch E, Collins M,

et al. Genetic analysis of the capsular biosynthetic locus from all 90 pneumococcal

serotypes. PLoS Genet. 2006;2:262–9.

67. Uria MJ, Zhang Q, Li Y, Chan A, Exley RM, Gollan B, et al. A generic mechanism in

Neisseria meningitidis for enhanced resistance against bactericidal antibodies. J Exp

Med. 2008;205:1423–34.

Figure Legends

Figure 1: Rate of discovery of distinct K loci with increasing genome sample size.

Curves indicate the accumulation of distinct K loci (mean ± SE) in different genome

sets. Grey; all genomes (n=2410, excluding *K. oxytoca*). Blue; non-redundant

genomes (n=1081, excludes genomes from investigations of disease outbreaks and

specific clonal groups). Black; species specific genome sets (*K. pneumoniae* refers to

K. pneumoniae sensu stricto, SE not shown). Inset box shows a zoomed view of the

bottom-left section of the plot, as indicated by the dashed box.

Figure 2: Example K locus structures and comparisons

Coding sequences are represented as arrows coloured by predicted function of the

protein product and labelled with gene names where known. Grey bars indicate

regions of similarity identified by BLAST comparison, darker shading indicates higher

sequence identity. **(A)** Comparison of deletion variant KL15-D1 and insertion

sequence variant KL15-1 with the synthetic KL15 locus. **(B)** Comparison of the

insertion sequence variant KL22-1 with the K-type reference KL22 locus. The

downstream LPS (lipopolysaccharide synthesis) operon (pink arrows) has been

translocated into the K locus.   35

Figure 3: Composition and diversity of *Klebsiella* K loci

(A) Putative recombination events among the common K locus genes. Values plotted

are relative number of events per 100 bp window, inferred using ClonalFrameML. **(B)**

Representation of a generalised K locus structure. Arrows represent K locus coding

regions coloured by predicted protein product as in **Figure 2**. Percentage values

indicate the number of reference K loci containing each gene (total 134 references).

Note that 13 of the K locus references partially or completely exclude *ugd*, although

it is known to be present in 11 of these loci [22]. Thus we counted these 11 as

containing *ugd*. The locations within this structure at which *wzx* (C), *wzy* (D) and

sugar processing genes (E) have been found to occur are indicated. **(C, D, E)** Diversity

of proteins encoded by *wzx* (C), *wzy* (D) and sugar processing genes (E) annotated

amongst the 134 K locus reference sequences. Bar charts indicate the frequency of

each predicted protein cluster.

Figure 4: Co-occurrence of *wzx, wzy* and sugar processing genes across reference K

loci

Nodes represent genes, labelled by name (HYP: hypothetical protein) and coloured

by protein product as in Figures 2 and 3: Wzx (red), Wzy (Orange), mannose

synthesis/processing proteins (dark purple), rhamnose synthesis/processing proteins

(light purple), other proteins (green). Edge widths are proportional to Jaccard index

(*J*) and are shown for all pairs where *J* ≥0.61. Numbers represent co-occurrence

group assignments as defined in **Table S5, Additional File 5**.

Figure 5: Allelic diversity of *wzi*

Within K locus *wzi* allelic diversity increases with total K locus representation. The

blue line represents the least-squares regression and grey shading indicates the 95%

confidence interval.

Figure 6: *wzc* nucleotide diversity

Barplots showing distribution of pairwise *wzc* nucleotide divergence for pairs of

genomes with the same (light blue) or different (dark blue) K loci. The inset box

shows a zoomed view of the lower end of the distribution.

Figure 7. Summary of the *Kaptive* analysis procedure

Kaptive takes as input a set of annotated reference K loci in GenBank format and one

or more genome assemblies, each as a single FASTA file of contigs. *Kaptive* performs

a series of BLAST searches to identify the best-match K locus in the query genome

and assess the presence of genes annotated in the best-match locus (expected

genes) and those annotated in other loci (unexpected genes) both within and

outside the putative K locus region of the query assembly. The output is a FASTA file

containing the nucleotide sequence(s) of the K locus region(s) for each query

assembly and a table summarising the best-match locus, gene content and potential

problems with the match (e.g. the assembly K locus region is fragmented, expected

genes are missing from the K locus region or at low identity, or unexpected genes

are present) for each query assembly. CDS = coding sequence.