Classification: (Major) Biological sciences, (Minor) Neuroscience

Title: Astrocytes locally translate transcripts in their peripheral processes

Authors: Kristina Sakers[1-3], Allison M. Lake[2,3], Rohan Khazanchi[2,3], Rebecca Ouwenga[1-3], Michael J. Vasek[2,3], Adish Dani[4], Joseph D. Dougherty[2,3]*

Affiliations:

[1]Division of Biology and Biomedical Sciences, Washington University School of Medicine, 660 S. Euclid Ave, Saint Louis MO, 63108, USA.

[2]Department of Genetics, Washington University School of Medicine, 660 S. Euclid Ave, Saint Louis MO, 63108, USA.

[3]Department of Psychiatry, Washington University School of Medicine, 660 S. Euclid Ave, Saint Louis MO, 63108, USA.

[4]Department of Pathology and Immunology, Washington University School of Medicine, 660 S. Euclid Ave, Saint Louis MO, 63108, USA.

*Correspondence to:

Joseph Dougherty, PhD

Department of Genetics

Washington University School of Medicine

Campus Box 8232

660 S. Euclid Ave, Saint Louis MO, 63108, USA.

jdougherty@genetics.wustl.edu

(314)362-3774

Keywords: Astrocyte, local translation, synapse, TRAP, RNA-sequencing

Abstract:

Local translation in neuronal processes is key to the alteration of synaptic strength that contributes to long term potentiation and learning and memory. Here, we present evidence that astrocytes have ribosomes in their peripheral and perisynaptic processes, and that *de novo* protein synthesis occurs in the astrocyte periphery. We also developed a new biochemical approach to profile and define a set of candidate transcripts that are locally translated in astrocyte processes, several of which were validated *in vivo* using *in situ* hybridization of sparsely labeled cells. Computational analyses indicate that localized translation is both sequence dependent and enriched for particular biological functions. This includes novel pathways such as fatty acid synthesis as well as pathways consistent with known roles for astrocyte processes, such as GABA and glutamate metabolism. Finally, enriched transcripts also include key glial regulators of synaptic refinement, suggesting that local production of astrocyte proteins may support microscale alterations of adjacent synapses.

Significance Statement:

Cellular compartments are specialized for particular functions. In astrocytes, the peripheral processes, particularly near synapses, contain proteins specialized for reuptake of neurotransmitters and ions, and have been shown to alter their morphology in response to activity. Regulated transport of a specific subset of nuclear-derived mRNAs to particular compartments is thought to support the specialization of these compartments and allow for local regulation of translation. In neurons, local translation near activated synapses is thought to generate the proteins needed for the synaptic alterations that constitute memory. We demonstrate that astrocytes also have sequence-dependent local translation in their peripheral processes, including transcripts with roles in regulating synapses. This suggests local translation in astrocyte processes may also play a role in modulating synapses.

Introduction:

Astrocytes are known to be indispensable in the development and maintenance of the synapse- a structure described as tripartite due to the critical contribution from peripheral processes of these cells(1). There is evidence that astrocytes determine synapse number(2, 3), and strength(4, 5), and complementary indications that neurons mediate maturation of astrocytes processes(6). Yet, our understanding of these interactions is incomplete.

The localized synthesis of proteins in neurons is instrumental for synaptic modulation(7) and neurite outgrowth(8, 9). Because of the distances neurites must extend to reach a target cell, they have the ability to harbor mRNAs and ribosomes proximal to spines and axons(10, 11) where local translation occurs in a precise spatial, temporal and activity-dependent manner. Additionally, oligodendrocytes also selectively localize some myelin associated mRNAs to specialized domains in their peripheral processes for local translation(12). Outside of the nervous system, localized protein synthesis has been detected in a variety of biological systems, including drosophila oocytes and migrating fibroblasts, suggesting that this phenomenon is a widely used regulatory system(13, 14).

Local translation in both neurons and oligodendrocyte processes suggests that other CNS cells may require this adaptation to compartmentalize functions within the cell. Peripheral astrocyte processes (PAPs) reach lengths comparable to that of some neurites and one astrocyte territory can contact up to 100,000 synapses(15). Further, the dynamic nature of PAPs at synapses requires

rapid responsiveness to synaptic changes(5, 16). Therefore, we hypothesized that astrocytes also
utilize local translation. We reasoned that several conditions must be met to establish that local
translation occurs in astrocytes: First, both ribosomes and mRNA must be present in PAPs.
Second, new protein translation must occur in PAPs. Finally, even if protein synthesis is detected,
if local translation in astrocytes does indeed support either local adaptation or specialization of
function as it does in other cells, then it should be regulated by specific sequence features, and
therefore be enriched for specific transcripts. Here, we provide biochemical and imaging evidence
for these criteria, and define for the first time the peripheral translatome of the astrocyte.

Astrocyte ribosomes and mRNA exist in their peripheral processes, *in vivo*.

Previous work has indicated that ribosome-like structures exist in PAPs, using electron
microscopy(17). We used transgenic mice where ribosomal protein Rpl10a is fused with EGFP
(Aldh1L1-EGFP/Rpl10a) specifically within astrocytes to localize the large subunit of ribosomes
in vivo(18). In cortical astrocytes, we found that EGFP/Rpl10a extends beyond Gfap+ processes
and into peripheral processes surrounded by Aqp4, which marks astrocyte membranes and vascular
endfeet(19) (Fig. 1A,B). As translation requires both large and small subunit, we confirmed
peripheral localization of endogenous small ribosomal subunits via immunofluorescence (IF)
against Rps16 (Fig. S1), indicating the tagged Rpl10a is not simply mislocalized in the Aldh1L1-
EGFP/Rpl10a mice. We also asked if we could detect tagged ribosomes in perisynaptic PAPs
using a recently-developed approach leveraging stochastic optical reconstruction microscopy
(STORM), to allow for ultrastructural level resolution compatible with multicolor fluorescent
labeling.(20) Astrocyte EGFP/Rpl10a was found within 100 nm of synapses, which are defined by
apposition of pre- and post-synaptic markers (Fig. 1C). This confirms the earlier EM work
indicating ribosome-like structures are found in the peripheral processes of astrocytes, and
suggests that the EGFP tag might be used for purification of ribosome bound transcripts, if mRNA
is indeed present and translating on these ribosomes.

Therefore, we next sought to confirm that at least some mRNAs are found in PAPs. We chose to
focus on the localization of *Slc1a2* (Glt-1) for two reasons: first, Glt-1 protein is highly expressed
in PAPs(21) and second, previous colormetric *in situ* hybridization (ISH) suggested its peripheral
localization(22). As both Aqp4 antibody and the pan astrocyte Aldh1L1-EGFP/Rpl10a were
suboptimal for easily tracking the processes of single astrocytes, we developed a viral method to
sparsely GFP-label astrocytes in non-transgenic mice (Fig. 2A). Using this approach, we
confirmed that *Slc1a2* mRNA is found throughout the astrocyte, including in distal GFAP negative
PAPs (Fig. 2B).

Astrocyte processes synthesize proteins in their distal tips.

After demonstrating the presence of both ribosomes and mRNA in PAPs, we next tested whether
distal translation occurs by using puromycin to label actively-translated peptides in acute brain
slices (Fig. 3A). Puromycin, a tRNA structural analog, incorporates into translating ribosomes and
puromycylates the growing peptide(23). Using an antibody against puromycin for IF in sparsely
labeled astrocytes (Fig. 2A, S2A), we were able to detect nascent translation within PAPs, which
was blocked by pretreatment with another translation inhibitor, anisomycin, indicating that
labeling requires active translation (Fig. 3B). We quantified the total translation occurring
throughout the astrocyte by measuring the fluorescence intensity of puromycin in the astrocyte
(determined by colocalization with GFP) at increasing radii from the nucleus of the astrocyte. We
empirically determined that a 27um radius captures the majority of the astrocyte, without

substantially extending to another astrocyte territory (Fig S2B), which is consistent with previous
data regarding a mean astrocyte diameter of 56 microns(24). Given the short duration of
incubation, we concluded that puromyclated peptides in PAPs were made locally rather than
transported. Quantification of puromyclation in astrocytes, indicates that on average 73% of
translation in an astrocyte occurs > 9 microns from the nucleus (Fig. 3C), and does not taper
significantly as it extends to the periphery.

PAP-TRAP reveals hundreds of enriched ribosome-bound transcripts in PAPs.

In neurons, local translation appears to be transcript-specific, often generating proteins with
synaptic roles. If translation in PAPs does indeed have a physiological role, then a specific subset
of astrocyte transcripts should be translated there. In other words, the localization of RNAs to the
periphery should be sequence dependent, and consistent with the known roles of the PAP in
glutamate homeostasis and other processes. To capture ribosome-bound mRNA from PAPs, we
applied translating ribosome affinity purification(TRAP) to a synaptoneurosome(SN) preparation
from Aldh1L1:EGFP/Rpl10a mice, a method we have abbreviated 'PAP-TRAP' (Fig. 4A). While
SN fractions have been previously studied largely for containing synaptic structures, previous
work has indicated at least one peripherally localized astrocyte mRNA is present in them(25) and
our preliminary RNA sequencing of this fraction suggested substantial contribution of mRNA
from non-neuronal cells(*data not shown).* Here, immunoblot confirmed depletion of nuclear
protein LaminB2 and enrichment, as expected, of Psd95 in the SN fraction (Fig. 4B). Additionally,
we show the fraction contains Ezrin, a constituent of PAPs(26), and astrocytic EGFP/Rpl10a (Fig.
4B). Thus, we used this affinity tag to harvest astrocyte ribosomes from the fraction, and performed
RNA-seq on the PAP-TRAP sample and three comparison samples (Table S1) to identify PAP-
enriched transcripts (Fig. 4A). Our analysis identified 224 transcripts that are significantly
enriched on PAP ribosomes (Fig. 4C, Table S2), including *Slc1a2*, and 116 that are
depleted('Soma' transcripts, Fig. 4C, Table S3). These data establish that PAP-TRAP samples
contain a distinct profile of ribosome-bound mRNAs, which includes the significant enrichment
of mRNAs for proteins, such as *Slc1a2*(Glt-1), known to be found in PAPs (Fig. 4D).
Unfortunately, PAPs have no sufficiently distinctive stable structure equivalent to a post-synaptic
density that can be used to readily verify their presence in SN fractions by electron microscopy.
Therefore, we instead focused on using an independent anatomical method *in vivo* to validate the
novel predications of the PAP-TRAP purification experiment. Using FISH, we successfully
confirmed the peripheral localization of several of these new candidates (*Cpe, Clu, Glul,*and
Sparc) (Fig. 5).

PAP-enriched transcripts suggest physiological roles for local translation including
modulation of neurotransmitter metabolism.

To understand what purpose local translation might serve in astrocytes, we did a pathway analysis
to identify consistent biological functions of the PAP enriched transcripts (Fig. 6A). We found that
PAP-localized transcripts had overrepresentation of genes mediating Glutamate and GABA
metabolism, consistent with PAP functions of Glutamate transport(*Slc1a2, Slc1a3*)(21, 27) and
metabolism(*Glul*). We also identified a set of enzymes representing multiple steps in a pathway
for biosynthesis of unsaturated fatty acids (*Scd1, Scd2, Fads1, Fads2, Elovl5, Hadha)* suggesting
there may be some local regulation of fatty acid production either for signaling or membrane use.
We also identified several motor and cytoskeletal proteins(e.g *Kif1c, Myo1D*) local translation of
which could play a role in morphological remodeling of astrocyte processes. Finally, we also
noticed an enrichment for genes and members of gene families known to regulate synapse

number(*Mertk, Sparc, Thbs4*)(28–30) suggesting that synapse formation and elimination may be
mediated in part by local translation of signaling molecules in astrocytes. Importantly, astrocyte
soma transcripts have largely non-overlapping GO categories with PAP transcripts and are
enriched for transcription regulators and amino acid catabolism (Fig. S4).

PAP-enriched transcripts have longer 3'UTRs and are higher expressed than Soma-localized
transcripts.

For dendritically localized transcripts, as well as the localization of β-actin in fibroblasts,
sequence-specific features are commonly found within the 3'UTR that mediate their
localization(31–33). Therefore, we asked whether PAP-enriched transcripts contain sequence-
specific features or characteristics that may mediate localized translation. We found no differences
in individual nucleotide or GC content (Fig. S5). However, PAP-enriched transcripts were
significantly higher expressed in cortical astrocytes compared to 'Soma' transcripts (Fig. 6B). In
neurons, locally translated mRNAs, often sequestered by RNA-binding proteins(RBPs) into
granules for transport and protection, have significantly longer 3'UTRs(34). This may enable more
sequence motifs or secondary structure for binding RBPs. We found that PAP-enriched transcripts
are also significantly longer, specifically in their 3'UTRs (Fig. 6C). Additionally, RBPs may
recognize secondary structure rather than specific sequence. We used the RNA-folding algorithm,
ViennaRNA(35), to investigate the stability of transcript 3'UTRs in the PAP and Soma transcripts.
Interestingly, we found that PAP-enriched 3'UTRs are predicted to have more stable secondary
structures (Fig 6D). The greater length and structure suggests that sequences in 3' UTRs may also
serve to regulate the localized enrichment of specific mRNAs in PAPs.

Discussion

Here we provide, to our knowledge, the first evidence of localized translation of specific transcripts
in astrocytes. We utilize a novel biochemical method to enrich for ribosome-bound transcripts in
PAPs and provide independent *in vivo* validation of several of the transcripts, confirming the PAP-
TRAP method is capable of selecting for peripherally enriched astrocyte mRNA. Additionally,
we have provided evidence that translation indeed occurs outside of the astrocyte soma, as
indicated by labeling of new protein synthesis with a brief pulse with puromycin. With this
approach we have confirmed that peripherally localized puromycylated peptides extend to the very
distal processes of the astrocyte.

Our data also indicate that PAP-enriched candidates have a median higher expression and a
significantly longer and more stable 3'UTR than those transcripts not enriched in the PAP. These
findings suggest that localization information exists in the 3'UTR, as is the case with many
dendritically-localized transcripts(32). Understanding of the mechanisms governing this
localization will provide insight to the phenomenon and opportunities to test the physiological
relevance of localized translation in astrocytes. Interestingly, we also detected a few neuron-
derived transcripts in the PAP-localized transcripts(Table S2), but we conclude that the removal
of these transcripts from the data does not change the overall conclusions of the PAP-localized
transcripts (Fig. S7, Table S4). The possible mechanisms for this signal may reveal interesting
biology and are of interest for future study.

We also note that our current list of locally translated candidates includes several disease genes,
most notably the Alzheimer's associated gene *Apoe* appears to be robustly enriched in the PAP-
TRAP sample. Further, at least two transcripts reported to be disrupted using TRAP in astrocytes
in Amyotrophic Lateral Sclerosis (ALS) models (*Heyl*, and *Vim*)(36), as well as *Slc1a2,*which is

downregulated in the motor cortex of ALS patients(37), are enriched in the PAP-TRAP sample. It
is possible that dysregulation of local translation may contribute to disease progression and
vulnerability of adjacent neurons. The PAP-TRAP method should be readily adaptable study this
possibility in mouse models of neurological disorders.

While local translation in neurons has been described for decades, many years later very few
studies have been able to address whether or not local translation specifically, rather than local
translation more generally had a physiological role. Local translation was first shown to be
important in the induction of long term potentiation (LTP), whereby physical separation of the
dendritic layer in the hippocampus was sufficient to induce LTP, but translation was required as
LTP was blocked by pretreatment with a protein synthesis inhibitor (7). Moreover, mutation of the
dendrite-targeting element of *CamKIIα* resulted in decreased performed in spatial memory tasks
such as the Morris Water Maze(38). Though we do not yet have the tools to directly probe the
physiological role of astrocyte translation, our pathway analyses provide interesting clues. We
found that the transcripts for secreted protein precursors are found in PAPs (*Clu*, *Cpe* and *Sparc*),
all of which have been identified as enriched in astrocyte-conditioned media by mass
spectrometry(39). This suggests that these glial derived soluble signaling molecules may be locally
synthesized before their extracellular release. Further, we noted transcripts for multiple proteins
with known roles in maintaining synapse numbers were enriched on PAP ribosomes (*Sparc, Mertk,*
Thbs4). It has been shown that Sparc negatively regulates excitatory synaptogenesis by
antagonizing Hevin(29). Further, Mertk is a required engulfment receptor for astrocyte-mediated
synapse elimination, *in vitro* and *in vivo*(28). Together, these findings provide evidence consistent
with the exciting possibility that localized translation might have a role in a localized astrocyte-
mediated synaptic modulation. Overall our data overwhelmingly indicate that astrocytes are
capable of a sequence-regulated localized translation. We posit here that, whereas neurons are
thought to require localized translation to compensate for the great length of their dendrites and
axons, astrocytes contacting multiple synapses may use local translation to allow for them to
respond to or modulate the activity of specific synapses.

Materials and Methods
Animals
All procedures were performed in accordance with the Washington University Institutional
Animal Care and Use Committee (IACUC) and the Animal Studies Committee (ASC). Mice were
maintained in standard housing conditions with food and water provided ad libitum. B6.FVB-
Tg(Aldh1L1-EGfp/Rpl10a)^{JD130Htz}/J animals(40) were bred to pure C57BL/6J mice (Jackson strain
000664) to obtain litters of GFP+ and GFP- animals.
For sparse labeling of astrocytes for microscopy, AAV9-CBA-IRES-GFP injections were
performed in P2 pups. 2μl of 10^{12} vector genome (vg) / mL virus was injected bilaterally in the
cortex, 1.5mm from the midline in two regions: 1mm caudal to bregma and 2mm rostral to lambda.
Injections were performed with a 33g needle (Hamilton #7803-05) with a 50μl Hamilton syringe
(#7655-01). Mice were aged until 3-4 postnatal weeks prior to CO_2 euthanasia and transcardial
perfusion with PBS, followed by 4% paraformaldehyde in PBS.
250
Acute Slice Preparation and Drug Treatment
Acute cortical slices (300μm) were prepared in artificial cerebrospinal fluid (aCSF, in
mM: 125 NaCl, 25 glucose, 25 NaHCO$_3$, 2.5 KCl, 1.25 NaH$_2$PO$_4$, equilibrated with 95% oxygen-
5% CO_2 plus 0.5 CaCl$_2$, 3 MgCl$_2$; 320 mosmol). 3μM Puromycin (Tocris #40-895-0) was added

to slices and allowed to incubate for 10 minutes at 37C. 1mM Anisomycin (Sigma #A9789) in
aCSF was added to slices 30 minutes before Puromycin treatment at 37°C.

<u>Immunofluorescence (IF)</u>
40μm sections were cryosectioned after 4% paraformaldehyde transcardial perfusions.
Sections were incubated with either: Rabbit GFAP (Dako, 1:1000), Rabbit Rps16 (Sigma, 1:1000),
Rabbit Aqp4 (Santa Cruz, 1:100), Mouse Puromycin (Kerafast, 1:1000) followed by detection with
appropriate Alexa conjugated secondary antibodies (Invitrogen) and nuclei were counterstained
with DAPI. For dual FISH/IF, GFP was detected with Chicken GFP (Aves) at 1:1000, and detected
as above. For STORM IF: 14μm sections were cryosectioned after flash freezing on dry ice and
images as previously described(41). Cryosections were immunolabelled with primary antibodies,
Chicken anti-GFP (Invitrogen), Mouse anti-Bassoon (SAP7F407, Novus Biologicals) and Rabbit
anti-Homer1 (Synaptic Systems, Germany) followed by anti-chicken, anti-rabbit and anti-mouse
secondary antibodies raised in donkey (Jackson Immunoresearch). Secondary reagents were
custom conjugated with acceptor and reporter fluorophore dye pairs Cy3-Alexa647, Cy2-
Alexa647 and Alexa405-Alexa647. Immunolabelled sections were overlaid with a buffer
containing 100mM Tris-HCL pH8.0, 150mM NaCl and containing an oxygen scavenging system
comprised of glucose, glucose oxidase, catalase and the reducing agent 2-mercaptoethylamine
(Bates *et al.*, 2007). After removing excess buffer, edges of the coverslip were sealed with nail
polish before imaging. Representative images are from at least triplicate experiments.

<u>Fluorescent In Situ Hybridization (FISH)</u>
FISH was performed on 14μm slide-mounted cryosections after 4% paraformaldehyde
fixation of either Aldh1l1:eGFP/Rpl10a or AAV9-CBA-IRES-GFP animals. Digoxigenin-labeled
antisense probes were made from purified PCR products containing a T7 promoter site in the
appropriate orientation. In vitro transcription was performed using T7 polymerase (Promega
#P2075) and Dig RNA labeling Mix (Roche #11277073910) according to manufacturer's
instructions. Primers for probe template were as follows:
Slc1a2: 5' GTTCCCTCCCAATCTGGTAGA 3', 5'
*TAATACGACTCACTATAGGG*CCGTGGCTGTGATGCTTATT 3'
Cpe: 5' TTTTCCAAAGCTTGGCTCGC 3', 5'
TAATACGACTCACTATAGGGTGTGATTGCCAGGTAGCCAG 3'
Clu: 5' GCTCTGGAGGACACTAGGGA 3', 5'
TAATACGACTCACTATAGGGCCCATAGTGGGAGGGAGACA 3'
Sparc: 5' GCCTGGATTCCGAGCTG 3', 5'
TAATACGACTCACTATAGGGCCTCCAGGGCAATGTACTTGT 3'
Glul: 5' CGGGCCTGCTTGTATGCTG 3', 5'
TAATACGACTCACTATAGGGTCTTCTCCTGGCCGACAGTCC 3'
Hybridization was performed at 65C overnight in a humid chamber, followed by washes
and H_2O_2 block as described in(42).
Detection was performed using Sheep Anti-Dig-POD (Roche #11207733910) followed by
Tyramide Signal Amplification Cyanine 3 Tyramide (PerkinElmer #NEL704A001KT) as per
manufacturer's instructions. Slides were then incubated with antibodies against additional specific
proteins (see IF above) for dual FISH/IF.

 Microscopy

Confocal microscopy was performed on an UltraVIEW VoX spinning disk microscope
(PerkinElmer) or an AxioImager Z2 (Zeiss). SIM microscopy was performed on a Nikon n-SIM
microscope.

STORM images were acquired on a custom built single molecule-imaging rig. Briefly, the
imaging rig was constructed on an active dampening vibration isolation table (Newport) and
around a Nikon TiE inverted microscope. Lasers, 642nm (Vortran laser, CA), 561nm, 488nm
(Sapphire, Coherent Inc., CA) and 405nm (Cube, Coherent Inc., CA) were shuttered individually
via mechanical shutters as well as an acousto optical tunable filter (AOTF, Crystal technologies,
PA). Lasers were combined, expanded, collimated and focused at the back focal plane of the
microscope. The diameter of the laser beam was cut using a field diaphragm to just cover a
256x256 pixel area of the CCD camera. A mechanical translation stage enabled objective type
total internal reflection (TIR) illumination via a 100X, 1.4NA objective (UPlanSApo, Olympus).
The Nikon perfect focus system was used to lock focus at the coverslip surface during imaging. A
quad band dichroic mirror zt/zet-405/488/561/640rpc (Chroma Technology, VT) was used to
excite with laser light and fluorescence collected by the same objective was filtered using
et705/72m emission filter for Alexa647. Images were acquired on a back illuminated EM-CCD
camera (DU-897, iXon+, Andor Technology, UK), using custom software. 3D astigmatism
imaging was performed as described before by placing a cylindrical lens in front of the camera.
Sparse single molecule images were acquired at 60Hz frequency using an imaging and activator
laser sequence: one frame of weak activator laser (561/488/405) followed by three frames of
642nm laser at 560W/cm^2, repeated as a train for each activator dye-antibody combination. The
amount of activation lasers was adjusted to ensure sparse single molecule events in each camera
frame. Raw image stacks were analyzed to determine the centroid positions of fluorescent intensity
peaks and these STORM localizations were rendered as images using custom software.

Quantification of imaging data.
To determine the distribution of translation occurring in the astrocyte, we drew concentric rings
around the nucleus with increasing radii of 3μm. We empirically determined that a circle with a
radius of 27μm encompasses the entire cell, without approaching another astrocyte's territory. The
results of this quantification is demonstrated in Fig 3. Analysis was conducted in ImageJ and
described in Figure S2. Experiments were conducted in triplicate. Cells counted are reported in the
legend for figure 3.

Synaptoneurosome (SN) preparation
SN fractions were isolated using a modified protocol(43). Three cortices (including
hippocampi) were pooled from Aldh1l1:eGFP/Rpl10a mice and homogenized in 3.5mL of: 5mM
Tris pH 7.5 and 250mM sucrose supplemented with 0.5mM DTT, 100μg/mL cycloheximide, 1mM
tetrodotoxin (Tocris #1069), 40units/mL rRNAsin (Promega #N2511), 20units/mL SUPERaseIn
(Ambion #AM2694) and 1 tablet/10mL cOmplete Mini EDTA-free protease inhibitor cocktail
(Roche #4693159001). Homogenization was performed in a glass dounce homogenizer (7mL max
volume) with 10 strokes each of the loose and tight pestles. The homogenate was first spun at
1000xg in a Sorvall RT7 plus centrifuge for 10 minutes at 4°C without brake. 2mL of the 2.5mL
supernatant was layered onto a sucrose percoll gradient as described (43) in Seton open-top tubes
(#5042) and spun at 32,500xg for 5minutes in a Sorvall RC-5C without brake. The SN band was
collected by puncturing the bottom of the tube, allowing the bottom milliliter to drip out and

collecting the next 3.5mL, which are the combined SN fractions as described(43). 200μl of the SN
fraction was used as the SN input sample. The remaining homogenate supernatant (~500μl) was
spun at 20,000xg for 15mins at 4C after incubation with 1% NP-40(Sigma) and 30mM DHPC
(Avanti). 60ul of the supernatant was added to 140μl of 0.15M KCl supplemented as the
homogenization buffer, and was set aside for use as the Cortex input sample. The remaining
supernatant and the SN fraction were used for either TRAP or Western blotting.
353
<u>Translating ribosome affinity purification (TRAP)</u>
TRAP was performed on 3 replicates of SN fraction (PAP TRAP sample) and cortex
homogenate (cortex TRAP sample), modified from protocols as described(40, 44). Briefly,
biotinylated-Protein L coated streptavidin magnetic beads were coupled with 2 monoclonal mouse
Anti-EGFP antibodies(40). Samples were added to conjugated beads and incubated on a inverted
mixer for 3 hours at 4°C. Beads were then washed 3 times with a high-salt buffer (10 mM HEPES
at pH 7.4, 350 mM KCl, 5 mM $MgCl_2$, 1% NP-40, 0.5 mM dithiothreitol, 100 μg/mL
cycloheximide). RNA purification was performed on eluted beads and respective input samples in
parallel using Qiagen RNEasy MinElute kit. RNA quality and concentration was assessed using
PicoChips on the Agilent BioAnalyzer. As large subunit was tagged, representative traces (Fig
S6) containing both 18S(small subunit) and 28S(large subunit) ribosomal RNA confirm that 80S
ribosomes were harvested from the SN fraction and comparison groups. All samples used in this
experiment had RIN scores > 8 for RNAseq library preparation.
367
368 <u>Library preparation</u>
369 Double stranded cDNA was prepared from 2ng of RNA using the Nugen Ovation RNAseq
370 System V2. Illumina sequencing libraries were generated from 600 ng of amplified cDNA, after
371 shearing to ~200bp using a Covaris sonicator as described in the protocol (NEBNext Ultra DNA
372 Library Prep Kit for Illumina, #E7370S).
373
<u>Western Blots</u>
Three independent replicates of the cortex and SN input samples were obtained. Crude
lysate was run on a 4-12% polyacrylamide gel (BioRad) and transferred to PVDF membranes
using a semi-dry transblot apparatus (BioRad). Membranes were probed using the following
antibodies at the stated concentrations: Chicken GFP (Aves) 1:1000, Mouse Psd95 (Enzo) 1:1000,
Mouse Ezrin (Abcam)1:500, LaminB2 (Sigma), 1:500.
380
<u>RNA-sequencing</u>
Libraries were sequenced on an Illumina HiSeq 2500, for a total of 175M 50bp reads across
the 12 samples. Reads were analyzed as described(45). Briefly, reads were trimmed using
Trimmomatic (v0.32), rRNA reads were depleted with bowtie2. Surviving reads were aligned to
Ensembl release 75 of the mouse genome using STAR and then counted using HTSeq. Differential
expression analysis was performed using the edgeR package. CPMs of all 12 are in Table S1. Raw
and analyzed RNA-sequencing data are also available at GEO: GSE74456.
PAP enriched transcripts were determined by the intersect of: transcripts enriched in SN
input/Cortex input (FDR < 0.1, Log_2(Fold Change) > 1.5) and transcripts enriched in PAP
TRAP/SN input (FDR < 0.1, Log_2(Fold Change) > 1.5). These transcripts are listed in Table S2.
Soma enriched transcripts were determined by the intersect of: transcripts depleted in SN
input/Cortex input (FDR < 0.1, Log_2(Fold Change) < 1.5) and transcripts enriched in Cortex

TRAP/Cortex Input (FDR < 0.1, Log$_2$(Fold Change) > 1.5). These transcripts are listed in Table
S3.

<u>Gene Ontology Analysis</u>
PAP and Soma lists (see RNA-sequencing and Fig. 3C) were analyzed using the BiNGO
plugin(v3.0.3) in Cytoscape(v3.2.1). Mouse Genome Informatics (MGI) IDs and descriptions were
obtained from the biomaRt package from Bioconductor in R. Overrepresented biological
processes were identified using a hypergeometric test with Benjamini-Hochberg. Lists were
compared to a background list of genes detected at CPM > 2 in Cortex input sample.

<u>Sequence feature analysis</u>

For each gene, the sequence of the longest isoform was obtained using the biomaRt
package in R and used in subsequent analyses. We analyzed sequence length, GC content, and
individual base content of the 5'UTR, CDS, and 3'UTR, as well as complexity of secondary
structure in the 5'UTR and 3'UTR using the Vienna RNA package(46). All comparisons
between Soma or PAP lists were made with Wilcoxon rank-sum test with Benjamini-Hochberg
correction. To predict RNA structure complexity, we used the ViennaRNA RNAfold program,
which calculates the minimum free energy secondary structure of the input RNA sequence. Since
longer sequences are more likely by chance to have more complex secondary structures, we
normalized this free-energy value by the length of the sequence.

Author contributions statement:

RO developed and KS and RO refined the PAP-TRAP method. KS, AL, and JD designed and
conducted bioinformatic analysis. RK, KS and JD conducted ISH, IF, imaging, quantification,
and analysis. KS conducted and analyzed puromycylation studies. AD conducted STORM. MV
developed image analysis pipeline for quantification. JD and KS conceived of study, designed
experiments, and wrote manuscript.

Acknowledgments:

We thank Dr. Joshua Dearborn for help with the neonatal injections, Dr. Min-Yu Sun for guidance
in preparing acute slices, Drs Dennis Oakley and James Fitzpatrick for training and assistance with
imaging, Dr. Kelly Monk for comments on the manuscript, and the members of the Dougherty lab
for advice. This work was supported by the CDI(MD-II-2013-269, and a WUCCI Microgrant),
NIH(DA038458-01, MH099798-01), WUSTL Interface of Psychology Neuroscience and
Genetics Training Grant (5T32 GM081739) and WUSTL Neurosciences Program Training Grant
(T32 GM008151). Key technical resources were provided by the Hope Center and the Genome
Technology Resource Center at Washington University(supported by NIH grants P30 CA91842
and UL1TR000448), and the Washington University Center for Cellular Imaging(supported by
CDI, St. Louis Children's Hospital, the Foundation for Barnes-Jewish Hospital, and
NIH:NS086741).

References

1.   Araque A, et al. (1999) Tripartite synapses: glia, the unacknowledged partner. *Trends*
Neurosci 22(5):208–215.

2.   Ullian EM, Sapperstein SK, Christopherson KS, Barres BA (2001) Control of synapse
number by glia. *Science* 291(5504):657–661.

3.   Mauch DH, et al. (2001) CNS synaptogenesis promoted by glia-derived cholesterol.
Science 294(5545):1354–1357.

4.   Pfrieger FW, Barres BA (1997) Synaptic Efficacy Enhanced by Glial Cells in Vitro.
Science 277(5332):1684–1687.

5.   Jourdain P, et al. (2007) Glutamate exocytosis from astrocytes controls synaptic strength.
Nat Neurosci 10(3):331–339.

6.   Swanson RA, et al. (1997) Neuronal regulation of glutamate transporter subtype expression
in astrocytes. *J Neurosci Off J Soc Neurosci* 17(3):932–940.

7.   Kang H, Schuman EM (1996) A requirement for local protein synthesis in neurotrophin-
induced hippocampal synaptic plasticity. *Science* 273(5280):1402–1406.

8.   Kislauskis EH, Li Z, Singer RH, Taneja KL (1993) Isoform-specific 3'-untranslated
sequences sort alpha-cardiac and beta-cytoplasmic actin messenger RNAs to different
cytoplasmic compartments. *J Cell Biol* 123(1):165–172.

9.   Zhang HL, et al. (2001) Neurotrophin-induced transport of a beta-actin mRNP complex
increases beta-actin levels and stimulates growth cone motility. *Neuron* 31(2):261–275.

10. Steward O, Levy WB (1982) Preferential localization of polyribosomes under the base of
dendritic spines in granule cells of the dentate gyrus. *J Neurosci Off J Soc Neurosci*
2(3):284–291.

11. Burgin KE, et al. (1990) In situ hybridization histochemistry of Ca2+/calmodulin-
dependent protein kinase in developing rat brain. *J Neurosci* 10(6):1788–1798.

12. Colman DR, Kreibich G, Frey AB, Sabatini DD (1982) Synthesis and incorporation of
myelin polypeptides into CNS myelin. *J Cell Biol* 95(2 Pt 1):598–608.

13. Martin KC, Ephrussi A (2009) mRNA Localization: Gene Expression in the Spatial
Dimension. *Cell* 136(4):719–730.

14. Jung H, Gkogkas CG, Sonenberg N, Holt CE (2014) Remote Control of Gene Function by
Local Translation. *Cell* 157(1):26–40.

15. Oberheim NA, Wang X, Goldman S, Nedergaard M (2006) Astrocytic complexity
distinguishes the human brain. *Trends Neurosci* 29(10):547–553.

16.   Mennerick S, Zorumski CF (1994) Glial contributions to excitatory neurotransmission in
467   cultured hippocampal cells. *Nature* 368(6466):59–62.

468   17.   Špaček J (1982) Ribosome-associated membrane contacts between astrocytes in the anoxic
469   brain. *Acta Neuropathol (Berl)* 57(4):270–274.

470   18.   Doyle JP, et al. (2008) Application of a translational profiling approach for the comparative
471   analysis of CNS cell types. *Cell* 135(4):749–762.

472   19.   Nielsen S, et al. (1997) Specialized Membrane Domains for Water Transport in Glial Cells:
473   High-Resolution Immunogold Cytochemistry of Aquaporin-4 in Rat Brain. *J Neurosci*
474   17(1):171–180.

475   20.   Dani A, Huang B, Bergan J, Dulac C, Zhuang X (2010) Superresolution Imaging of
476   Chemical Synapses in the Brain. *Neuron* 68(5):843–856.

477   21.   Danbolt NC, Storm-Mathisen J, Kanner BI (1992) An [Na+ + K+]coupledl-glutamate
478   transporter purified from rat brain is located in glial cell processes. *Neuroscience*
479   51(2):295–310.

480   22.   Berger UV, DeSilva TM, Chen W, Rosenberg PA (2005) Cellular and Subcellular mRNA
481   Localization of Glutamate Transporter Isoforms GLT1a and GLT1b in Rat Brain by In Situ
482   Hybridization. *J Comp Neurol* 492(1):78–89.

483   23.   Schmidt EK, Clavarino G, Ceppi M, Pierre P (2009) SUnSET, a nonradioactive method to
484   monitor protein synthesis. *Nat Methods* 6(4):275–277.

485   24.   Oberheim NA, et al. (2009) Uniquely Hominid Features of Adult Human Astrocytes. *J
486   Neurosci* 29(10):3276–3287.

487   25.   Gerstner JR, et al. (2012) Time-of-day regulates subcellular trafficking, tripartite synaptic
488   localization and polyadenylation of the astrocytic Fabp7 mRNA. *J Neurosci Off J Soc
489   Neurosci* 32(4):1383–1394.

490   26.   Derouiche A, Frotscher M (2001) Peripheral astrocyte processes: monitoring by selective
491   immunostaining for the actin-binding ERM proteins. *Glia* 36(3):330–341.

492   27.   Rothstein JD, et al. (1996) Knockout of Glutamate Transporters Reveals a Major Role for
493   Astroglial Transport in Excitotoxicity and Clearance of Glutamate. *Neuron* 16(3):675–686.

494   28.   Chung W-S, et al. (2013) Astrocytes mediate synapse elimination through MEGF10 and
495   MERTK pathways. *Nature* 504(7480):394–400.

496   29.   Kucukdereli H, et al. (2011) Control of excitatory CNS synaptogenesis by astrocyte-
497   secreted proteins Hevin and SPARC. *Proc Natl Acad Sci U S A* 108(32):E440-449.

498   30.   Christopherson KS, et al. (2005) Thrombospondins Are Astrocyte-Secreted Proteins that
499   Promote CNS Synaptogenesis. *Cell* 120(3):421–433.

31. Mayford M, Baranes D, Podsypanina K, Kandel ER (1996) The 3'-untranslated region of
CaMKII alpha is a cis-acting signal for the localization and translation of mRNA in
dendrites. *Proc Natl Acad Sci U S A* 93(23):13250–13255.

32. Andreassi C, Riccio A (2009) To localize or not to localize: mRNA fate is in 3′UTR ends.
Trends Cell Biol 19(9):465–474.

33. Condeelis J, Singer RH (2005) How and why does beta-actin mRNA target? *Biol Cell*
97(1):97–110.

34. Han TW, et al. (2012) Cell-free formation of RNA granules: bound RNAs identify features
and components of cellular assemblies. *Cell* 149(4):768–779.

35. Gruber AR, Lorenz R, Bernhart SH, Neuböck R, Hofacker IL (2008) The Vienna RNA
Websuite. *Nucleic Acids Res* 36(suppl 2):W70–W74.

36. Sun S, et al. (2015) Translational profiling identifies a cascade of damage initiated in motor
neurons and spreading to glia in mutant SOD1-mediated ALS. *Proc Natl Acad Sci U S A*
112(50):E6993-7002.

37. Rothstein JD, Van Kammen M, Levey AI, Martin LJ, Kuncl RW (1995) Selective loss of
glial glutamate transporter GLT-1 in amyotrophic lateral sclerosis. *Ann Neurol* 38(1):73–
84.

38. Miller S, et al. (2002) Disruption of Dendritic Translation of CaMKIIα Impairs
Stabilization of Synaptic Plasticity and Memory Consolidation. *Neuron* 36(3):507–519.

39. Greco TM, Seeholzer SH, Mak A, Spruce L, Ischiropoulos H (2010) Quantitative Mass
Spectrometry-based Proteomics Reveals the Dynamic Range of Primary Mouse Astrocyte
Protein Secretion. *J Proteome Res* 9(5):2764–2774.

40. Doyle JP, et al. (2008) Application of a translational profiling approach for the comparative
analysis of CNS cell types. *Cell* 135(4):749–762.

41. Dani A, Huang B, Bergan J, Dulac C, Zhuang X (2010) Superresolution imaging of
chemical synapses in the brain. *Neuron* 68(5):843–856.

42. Reddy AS, et al. (2016) A Comprehensive Analysis of Cell Type-Specific Nuclear RNA
From Neurons and Glia of the Brain. *Biol Psychiatry*. doi:10.1016/j.biopsych.2016.02.021.

43. Westmark PR, Westmark CJ, Jeevananthan A, Malter JS (2011) Preparation of
synaptoneurosomes from mouse cortex using a discontinuous percoll-sucrose density
gradient. *J Vis Exp JoVE* (55). doi:10.3791/3196.

44. Heiman M, et al. (2008) A translational profiling approach for the molecular
characterization of CNS cell types. *Cell* 135(4):738–748.

45.   Reddy AS, et al. (2016) A Comprehensive Analysis of Cell Type-Specific Nuclear RNA
534   From Neurons and Glia of the Brain. *Biol Psychiatry*. doi:10.1016/j.biopsych.2016.02.021.

46.   Lorenz R, et al. (2011) ViennaRNA Package 2.0. *Algorithms Mol Biol AMB* 6:26.

Figure Legends

Figure 1: Subcellular localization in cortical astrocytes shows ribosomal proteins in
peripheral processes and in close proximity to synapses (A-B) Confocal IF of a cortical
astrocyte, stained with Gfap(A) or Aqp4(B)(white). EGFP/Rpl10a(green) shows ribosomal tag
extending throughout astrocyte, past Gfap+ processes (yellow dashed line) and within fine
processes highlighted by Aqp4. Scale bar=10μm. (C) STORM imaging showing an EGFP/Rpl10a
(green) filled astrocyte process proximal to synapses(as illustrated, these are defined by apposition
of Bassoon(red), and Homer(blue)). Inset of box on left, and side view is a 90 rotation of a second
synapses, again showing EGFP/Rpl10a puncta surrounding a synapse.

Figure 2: Method for sparse astrocyte labeling and FISH of *Slc1a2* reveals its peripheral
localization. (A) Early postnatal pups were injected transcranially with AAV9 expressing GFP,
resulting in sparse labeling of cortical astrocytes after 3 weeks. Scale bar=20μm. (B) Confocal IF
of a GFP-labeled astrocyte shows extension of *Slc1a2*(red) FISH into fine, Gfap(white) negative
peripheral processes, scale bar=10μm.

Figure 3: Peripheral ribosomes are actively translating in astrocytes (A) Acute slices(300 μM)
from P21 mice with sparse astrocyte labeling(generated as in Fig. 2), are treated for 10 minutes of
puromycin, with or without 30 minutes of anisomycin pretreatment, then fixed, sectioned, and
processed for IF against puromycin(red) and GFP(green). (B) Max projection super-resolution
Structured Illumination Microscopy(SIM) to detect puromyclation(red) of synthesizing proteins in
a GFP-labeled astrocyte shows translation occurs in peripheral processes, and is blocked by
pretreatment with anisomycin. Scale bar=10μm. (C) Quantification of puromycin intensity (only
puromycin pixels that were in astrocytes labeled with GFP were measured) at increasing radii from
the nucleus indicates robust translation occurs in PAPs. Repeated measures ANOVA revealed
main effects of condition $F(2, 138)=9.694$, $p=0.0001$ and distance $F(8, 1104) = 19.023$, $p < 2E-16$
with a significant interaction between condition and distance $F(16, 1104) = 2.019$, $p=0.01$. Data
represented as mean +/- SEM. Asterisks represent one-sided T-tests, *post hoc*. **** $p < 0.001$,
***$p < 0.005$, **$p < 0.01$. N(cells) = 54(puromycin), 48(anisomycin+puromycin).

Figure 4: Identification of peripherially enriched transcripts (A) Diagram of experimental
steps in PAP-TRAP and comparison samples for RNA-seq (B) Representative immunoblots for
input and SN fraction confirms enrichment of synaptic and PAP proteins, depletion of nuclear
proteins (LaminB2) and presence of EGFP/Rpl10a (C) Diagram of analytical strategy for defining
PAP and Soma enriched transcripts (D) Examples of PAP-enriched transcripts, including those
related to Glutamate metabolism(*Slc1a2, Slc1a3, Glul)*, fatty acid synthesis(*Fads, Scd)*, and
interesting signaling molecules(*Ptch1, Sparc, Ntsr2*).

Figure 5: *In vivo* **validation of PAP-TRAP candidates** (A) FISH on GFP-labeled astrocytes
confirms presence of PAP-TRAP identified mRNAs(Cpe, Clu, Glul and Sparc) in peripheral
processes. Cortical astrocytes were labeled as described in Fig 2. Images are representative of three
independent experiments with each probe.

Figure 6: Pathway and sequence analysis on PAP-enriched transcripts (A) Representative
significant GO terms for PAP-enriched transcripts, hypergeometric test with Benjamini-Hochberg
correction(B-H) (B) Quantification of expression of PAP-enriched and Soma-enriched transcripts
indicates Soma transcripts have lower median expression in cortical astrocytes(Wilcoxon test, B-
H corrected, *p<0.05) (C) Quantification of length of PAP and Soma transcripts indicates PAP-
enriched transcripts have longer 3'UTRs(Wilcoxon test, B-H corrected, ****p<0.0001). (D) RNA
structure-score (minimum free energy of most stable predicted structure, normalized to length),
indicates PAP-enriched transcripts have more stable 3'UTR secondary structures (Wilcoxon test,
B-H corrected, **p<0.01, lower values are more stable).

A eGFP/Rpl10a | Gfap | Merge/DAPI | Inset

B eGFP/Rpl10a | Aqp4 | Merge/DAPI | Inset

C STORM | Inset | Illustration | Side View

Bassoon: presynapse

Gfp/Rpl10a: Astrocyte

Homer: postsynapse

500nm | 200nm | 200nm

A P2 intracranial injections
AAV9:GFP

19 days →

Harvest Tissue
for ISH or slice
preparations at
P21

GFP

B

| AAV9:GFP | Gfap | Slc1a2 mRNA | Merge/DAPI |

A

3µM Puromycin

Puromycin

His
Lys
Val
Met

Puromycylated
nascent peptides

IF detection of
Puromycin *in situ*

P21 Acute mouse slices
(AAV9:GFP at P2)

B

	AAV9:GFP	Puromycin	GFAP	Merge/DAPI
3µM Puromycin				
Inset				
1mM Anisomycin 3µM Puromycin				

C

- Puromycin
- Anisomycin + Puromycin

*** **** *** * *** **** **** ****

A

Aldh1l1::eGFP/Rpl10a cortices

TRAP   SN isolation

3%
10%
15%
23%

TRAP

Cortex input   Cortex TRAP   SN input   PAP TRAP

RNA isolation and library prep

RNA sequencing

B

IB:

kDa:
100 — Psd95
75 — Ezrin
68 — Lamin B2
50 — eGFP/Rpl10a

Cortex input   SN input

C

Enriched in SN input vs Cortex input   **PAP enriched transcripts**   Enriched in PAP vs SN input

Depleted in SN input vs Cortex input   **Soma enriched transcripts**   Enriched in Cortex vs Cortex input

D

Symbol	Description
Apoe	apolipoprotein E
Cldn10	claudin 10
Clu	clusterin
Cpe	carboxypeptidase E
Csrp1	cysteine and glycine-rich protein 1
Cxcl14	chemokine (C-X-C motif) ligand 14
Fads1	fatty acid desaturase 1
Fads2	fatty acid desaturase 2
Glud1	glutamate dehydrogenase 1
Glul	glutamate-ammonia ligase (glutamine synthetase)
Kcnj10	potassium inwardly-rectifying channel, subfamily J, member 10
Kif1c	kinesin family member 1C
Mertk	c-mer proto-oncogene tyrosine kinase
Myo10	myosin X
Pla2g7	phospholipase A2, group VII (platelet-activating factor acetylhydrolase, plasma)
Ptch1	patched homolog 1
S100a1	S100 calcium binding protein A1
Scd1	stearoyl-Coenzyme A desaturase 1
Scd2	stearoyl-Coenzyme A desaturase 2
Slc1a2	solute carrier family 1 (glial high affinity glutamate transporter), member 2
Slc1a3	solute carrier family 1 (glial high affinity glutamate transporter), member 3
Sparc	secreted acidic cysteine rich glycoprotein
Ntsr2	neurotensin receptor 2

A

AAV9::GFP	mRNA	Merge/DAPI	inset

Clu

Cpe

Glul

Sparc

No probe

Supplementary Materials for:

Title: Astrocytes locally translate transcripts in their peripheral
processes

Authors: Kristina Sakers[1-3], Allison M. Lake[2,3], Rohan Khazanchi[2,3], Rebecca Ouwenga[1-
3,] Michael J. Vasek[2,3], Adish Dani[4], Joseph D. Dougherty[2,3]*

Affiliations:

[1]Division of Biology and Biomedical Sciences, Washington University School of
Medicine, 660 S. Euclid Ave, Saint Louis MO, 63108, USA.

[2]Department of Genetics, Washington University School of Medicine, 660 S. Euclid Ave,
Saint Louis MO, 63108, USA.

[3]Department of Psychiatry, Washington University School of Medicine, 660 S. Euclid Ave,
Saint Louis MO, 63108, USA.

[4]Department of Pathology and Immunology, Washington University School of Medicine,
660 S. Euclid Ave, Saint Louis MO, 63108, USA.

*Correspondence to:

Joseph Dougherty, PhD

Department of Genetics

Washington University School of Medicine

Campus Box 8232

660 S. Euclid Ave, Saint Louis MO, 63108, USA.

jdougherty@genetics.wustl.edu

(314)362-3774

25
26
27 **This PDF file includes:**

28
29   Tables legends for S1-S7

30
31 **Other Supplementary Materials for this manuscript includes the following:**

32
33   Supplementary Tables S1 to S4 as .csv

34
35
36 **Supplemental Figure Legends**

37
38

Fig. S1: EGFP/Rpl10a in Aldh1L1 bacTRAP mouse cortex extends beyond Gfap+ processes (A) Examination of localization of Rps16 by confocal reveals detectable Rps16(violet/white) in fine processes. Note Rps16 is also found robustly in adjacent neurons, as expected (*). Therefore, presence of GFP signal was used to mask all non-astrocyte pixels, and a binary image created showing GFP positive pixels that also contained Rps16 signal. This approach allows for identification of Rps16 labeling specifically in the astrocyte even in the presence of Rps16 labeling in adjacent cells. (B) A tissue section processed in parallel, but omitting Rps16 primary antibody, demonstrates absence of signal in the astrocyte.

Fig. S2: Puromycylation detection and quantification method. (A) Example of masking for quantification of peripheral puromyclation (Confocal IF). Note, as expected, robust translation also occurs in adjacent neurons(*).Therefore, presence of GFP signal was used to mask all non-astrocyte pixels, and a binary image created showing GFP positive pixels that also contained Puro-IF signal (GFP+Puro) for all downstream quantification. (B) Illustration of method for quantification of relative amount of translation at increasing distance from the astrocyte nucleus. A nucleus of a GFP+ astrocyte is selected and thresholded (middle left). A series of Sholl-like concentric rings are demarked for quantification (red rings). All Puro-IF outside of GFP+ area is masked, and Puro-IF positive area within each ring is measured (middle right). Intensity of Puro-IF within each GFP+ pixel is also measured (right panel).

Fig. S3: Pathway analysis using Gene Ontologies for PAP enriched transcripts. All enriched GO terms for PAP-enriched transcripts, $p < 0.005$, hypergeometric test with B-H correction.

Fig. S4: Pathway analysis using Gene Ontologies Soma enriched transcripts. All enriched GO terms for Soma-enriched transcripts, $p < 0.05$, hypergeometric test with B-H correction.

Fig. S5: Nucleotide composition does not differ between PAP and Soma-enriched transcripts. (A-D) Percent composition of each nucleotide for 5' UTR (white), coding sequence (CDs, green), and 3'UTR (purple) in PAP and Soma lists. $p > 0.05$, Wilcoxon test with B-H (E) GC percentage. $p > 0.05$, Wilcoxon test with B-H correction.

Fig. S6 Bioanalyzer confirms RNA quality from Cortex TRAP and PAP TRAP samples. (A) PicoChip electrophoresis for representative samples, marker is highlighted

in green. (B) Fluorescent trace visualization of PicoChip electrophoresis for Cortex TRAP
and PAP TRAP. Ribosomal RNAs from both subunits are captured.

Fig. S7 A small number of neuronal transcripts are bound to peripheral astrocyte
ribosomes: in vitro or in vivo mechanism? (A) Scatterplot of Cortex TRAP compared to
Cortex Input CPM for all transcripts (Log$_2$). We detect robust enrichment for sets of ~ 200
previously detected[1] astrocyte-enriched transcripts (green dots), and substantial, though
not complete, depletion of ~200 previously detected[1] neuron-enriched transcripts (blue
dots) in the current cortex TRAP experiment. This degree of enrichment and depletion is
typical for TRAP[2, 3] (B) While PAP-TRAP still enriches for most astrocyte transcripts
relative to SN samples, scatterplot of PAP-TRAP compared to SN input shows an
unexpected relative blunting of both enrichment and depletion relative to Cortex TRAP.
Note, significantly enriched transcripts are still detectable, as evidenced in Table S2.
Surprisingly, a handful of 'neuronal' transcripts now show relative enrichment following
capture of astrocyte ribosomes (arrows) from the SN fraction (PAP-TRAP). (C) These
include examples of many transcripts that are strong candidates to be locally translated in
neurons (*Camk2a, Shanks)*, or are known to be locally translated in oligodendrocytes (e.g.,
Mbp)[4]. (D) Roughly 15% of PAP-Enriched transcripts from Table S2, which show clear
enrichment by PAP-TRAP, are depleted when examining the Cortex TRAP vs Input
comparison, suggesting they are not present on most astrocyte ribosomes in the cell body,
as illustrated by Venn Diagram. (E) Illustrations of some of the possible alternate models
for these unexpected results. Left panel: interactions in solution between RNA granules
originally from neurons with captured granules from astrocytes *in vitro*, mediated by the
disordered nature of domains in many RBPs[5]. Right panels: low level exchange of
mRNA or ribosomal components between adjacent processes, or perhaps following
astrocyte phagocytosis of synapses[6], *in vivo*. Regardless, if needed, this 15% can be
removed from the PAP-Enriched list computationally (provided as Table S4), and all the
major conclusions regarding the remaining PAP-Enriched genes are still supported. (F)
77% of the original PAP GO terms survive at B-H corrected p<0.005. (G) Quantification
of expression of PAP-enriched and Soma-enriched transcripts indicates Soma transcripts
have lower median expression in cortical astrocytes(Wilcoxon test, B-H corrected,
*p<0.0001). (H) Quantification of length of PAP and Soma transcripts indicates PAP-
enriched transcripts are longer (Wilcoxon test, B-H corrected, ****p<0.0001, *p<0.05).
(I) RNA structure-score(minimum free energy of most stable predicted structure,
normalized to length), indicates PAP-enriched transcripts have more stable 3'UTR
secondary structures(Wilcoxon test, B-H corrected, *p<0.05, lower values are more stable)

Table S1. CPM
Counts per million for all biological replicates. Nomenclature: Cortex input = cortex
homogenate transcripts, Cortex TRAP = transcripts on GFP+ ribosomes in
Aldh1l1:eGFP/Rpl10a mouse cortex, SN input = transcripts in SN fraction, PAP TRAP =
transcripts on GFP+ ribosomes in Aldh1l1:eGFP/Rpl10a SN fraction.
118

Table S2. PAP enriched transcripts

PAP-enriched transcripts, as defined in Figure 4, with respective sample enrichment and FDR significance. Log fold changes (base 2) and FDR for three comparisons are included in the table. Abbreviations as in Table S1.

Table S3. Soma enriched transcripts

Soma-enriched transcripts, as defined in Figure 4, with respective sample enrichment and FDR significances, as in Table S2.

Table S4. PAP enriched transcripts, with putative neuron or oligodendrocyte derived transcripts removed.

PAP-enriched transcripts, as defined in S7, with 33 transcripts removed that were significantly depleted (FDR<.1, logFC< -1) in Cortex TRAP vs Cortex input comparison.

References for Supplemental Material

1. Dougherty JD, et al. (2012) Candidate pathways for promoting differentiation or quiescence of oligodendrocyte progenitor-like cells in glioma. *Cancer Res* 72(18):4856–4868.

2. Okaty BW, Sugino K, Nelson SB (2011) A Quantitative Comparison of Cell-Type-Specific Microarray Gene Expression Profiling Methods in the Mouse Brain. *PLOS ONE* 6(1):e16493.

3. Dougherty JD, Schmidt EF, Nakajima M, Heintz N (2010) Analytical approaches to RNA profiling data for the identification of genes enriched in specific cells. *Nucleic Acids Res* 38(13):4218–4230.

4. Li Z, Zhang Y, Li D, Feng Y (2000) Destabilization and Mislocalization of Myelin Basic Protein mRNAs in quaking Dysmyelination Lacking the QKI RNA-Binding Proteins. *J Neurosci* 20(13):4944–4953.

5. Calabretta S, Richard S (2015) Emerging Roles of Disordered Sequences in RNA-Binding Proteins. *Trends Biochem Sci* 40(11):662–672.

6. Chung W-S, et al. (2013) Astrocytes mediate synapse elimination through MEGF10 and MERTK pathways. *Nature* 504(7480):394–400.

C

logFC PAP-TRAP vs SN Input	logFC SN Input vs Cortex Input	logFC Cortex TRAP vs Cortex Input	symbol	description
0.75	1.35	-3.28	Bsn	bassoon
0.71	1.05	-3.64	Cacna1i	calcium channel, voltage-dependent, alpha 1I subunit
1.73	1.17	-3.10	Camk2a	calcium/calmodulin-dependent protein kinase II alpha
1.14	1.13	-1.98	Grin1	glutamate receptor, ionotropic, NMDA 1 (zeta 1)
1.30	0.99	-1.73	Homer2	homer scaffolding protein 2
1.77	1.50	-2.61	Mbp	myelin basic protein
0.70	1.37	-1.65	Pld3	phospholipase D family, member 3
1.21	2.03	-2.11	Shank1	SH3/ankyrin domain gene 1
1.64	1.02	-2.76	Shank2	SH3/ankyrin domain gene 2
2.28	1.07	-1.79	Shank3	SH3/ankyrin domain gene 3
1.08	1.66	-3.74	Slit1	slit homolog 1 (Drosophila)
1.55	1.41	-2.91	Slit3	slit homolog 3 (Drosophila)