An immuno-suppressive aphid saliva protein is delivered into the cytosol of plant

mesophyll cells during feeding.

Sam T. Mugford[1], Elaine Barclay[1], Claire Drurey[1,2], Kim C. Findlay[1] and Saskia

A. Hogenhout[1*].

[1]Cell and Developmental Biology, John Innes Centre, Norwich, NR4 7UH, UK.

[2]Current address: Institute of Infection, Immunity and Inflammation, Glasgow

University, G12 8TA, UK.

[*]corresponding author: saskia.hogenhout@jic.ac.uk

Abstract

Herbivore selection of plant hosts and plant responses to insect colonization have

been subjects of intense investigations. A growing body of evidence suggests that for

successful colonization to occur, (effector/virulence) proteins in insect saliva must

modulate plant defense responses to the benefit of the insect. A range of insect saliva

proteins that modulate plant defense responses have been identified, but there is no

direct evidence that these proteins are delivered into specific plant tissues and enter

plant cells. Aphids and other sap-sucking insects of the order Hemiptera use their

specialized mouthparts (stylets) to probe plant mesophyll cells, until they reach the

phloem cells for long-term feeding. Here we show by immunogold-labeling of

ultrathin sections of aphid feeding sites that an immuno-suppressive aphid effector

localizes in the cytoplasm of mesophyll cells near aphid stylets, but not in cells further

away from aphid feeding sites. In contrast, another aphid effector protein localizes in

the sheaths composed of gelling saliva that surround the aphid stylets. Thus, insects

deliver effectors directly into plant tissue. Moreover, different aphid effectors locate

extracellularly in the sheath saliva or are introduced into the cytoplasm of plant cells.

Introduction:

Insect herbivores likely modulate a range of plant processes in order to successfully

colonize their host plants. A growing body of evidence suggests that the saliva of

herbivorous insects contains virulence factors (effectors) that interfere with host plant

defences and facilitate colonisation (Musser et al. 2002; Will et al. 2007; Mutti et al

2008; Bos et al. 2010; Stuart et al. 2012; Atamian et al. 2013; Pitino and Hogenhout

2013; Chaudhary et al., 2014; Elzinga et al. 2014; Guo et al. 2014; Rodriguez et al.

2014; Naessens et al. 2015; Wang et al. 2015) Sap-sucking insects of the order

Hemiptera, such as aphids, whiteflies, leafhoppers, psyllids and planthoppers, are

stealthy feeders. These insects possess piercing-sucking mouthparts, consisting of a

pair of stylets, which navigate to vascular elements for long-term feeding.

Hemipterans cause direct feeding damage to plants and also transmit the majority of

plant viruses and several plant pathogenic bacteria (Hogenhout et al 2008; Guerrieri

and Digilio 2008; Orlovskis et al. 2015). The green peach aphid (GPA) *Myzus*

persicae alone transmits over a 100 plant viruses.

The feeding behaviour of aphids has been studied extensively. While navigating to the

phloem, in the pathway phase, aphid stylets probe various plant cells of the mesophyll

(Tjalingi 2006). During a probe the aphid stylets penetrate cells, deposit saliva inside,

and acquire cell contents (Martín et al. 1997). The pathway phase continues until the

aphid reaches a phloem sieve-element cell, where they remain to feed often for many

hours. The phloem feeding phase starts with release of saliva followed by extended

periods of phloem sap acquisition that are occasionally interrupted with short

salivation periods. Whereas it has been shown that this feeding behaviour is

responsible for the transmission of a diversity of viruses by aphids, the biological

significance of this behaviour to the aphids themselves is not fully understood.

Aphids produce different types of saliva: 'gelling' saliva that forms a protective

sheath around the stylets, and 'watery' saliva that is thought to be injected into plant

cells during the pathway and phloem feeding phases (Tjallingii and Esch 1993;

Martín et al. 1997). A number of proteins are detected in aphid saliva (Ramsey et al.

2007; Harmel et al. 2008; Carolan et al. 2009; Cooper et a. 2010; Nicholson et al.

2012, 2014; Rao et al. 2013; Vandermoten et al. 2014; Chaudhary et al. 2015). Aphid

saliva proteins have also been detected in lysate of plant tissues previously exposed

to aphids (Mutti et al. 2008; Naessens et al. 2015). Some components of aphid saliva

trigger plant defence responses that are characteristic of pattern-triggered immunity

(PTI), including reactive oxygen species (ROS) bursts and callose deposition, that

require the plant membrane-associated receptor-like kinase (RLK)

BRASSINOSTEROID INSENSITIVE 1-ASSOCIATED KINASE (BAK1)

(Chaudhary et al. 2014; Prince et al. 2014). BAK1 is a key regulator of several cell

membrane-localized pattern recognition receptors (PRRs), which mediate the first

step of the plant defence response via the recognition of pathogen-associated

molecular patterns, such as flg22 (Chinchilla et al. 2007; Chaudhary et al. 2014).

Aphid saliva contains effectors that suppress plant defence responses (Will et al.

2007; Mutti et al 2008; Bos et al. 2010; Atamian et al. 2013; Pitino and Hogenhout

2013; Chaudhary et al., 2014; Elzinga et al. 2014; Guo et al. 2014; Rodriguez et al.

2014; Naessens et al. 2015; Wang et al. 2015). Mp10 is one of a number of candidate

effector proteins that have been identified and was found to suppress the plant ROS

burst in response to flg22 (Bos et al. 2010; Rodriguez et al. 2014). Mp10 and another

candidate effector protein expressed in GPA salivary glands, MpOS-D1, belong to the

chemosensory protein (CSP) family (Bos et al. 2010; Zhou et al. 2010). Other aphid

effector candidates, such as MpPIntO1 and MpC002, promote aphid colonization

(Mutti et al 2008; Pitino and Hogenhout 2013; Pitino et al. 2011; Coleman et al. 2015)

and are found in aphid saliva and/or in extracts of aphid-exposed leaves (Mutti et al

2008; Harmel et al. 2008). However, evidence that insects deliver effectors directly

into plant cells is currently lacking.

How pathogens and pests deliver effectors to the appropriate site of action has been a

major research topic of many research groups (Dodds and Rathjen 2010). Gram-

negative bacterial pathogens deliver their effectors via specialized mechanisms, such

as type III secretion systems (Gopalan et al. 1998). For fungi and oomycetes it was

discovered that effectors require conserved sequences, such as an RXLR motif, at

their N-termini for entering the plant host, but the mechanisms underlying their

delivery is not fully understood (Petre and Kamoun 2014). It is as yet unclear how,

where and when aphids deposit their effectors in the plant; they may do so via the

'gelling' saliva to embed the proteins within the sheaths surrounding the stylets in the

apoplastic space of plant tissue, or via 'watery' saliva to introduce them into plant

cells. In addition, aphids may secrete different effectors during the pathway and

phloem feeding phases.

Here we studied ultra-thin sections of plant tissues containing tracks of aphid stylets

and used antisera raised to the aphid effectors Mp10, MpOS-D1, MpC002 and

MpPIntO1 for immuno-gold labelling of the sections. Antisera of Mp10, but not of

MpOS-D1, labelled the cytosol and chloroplasts of plant mesophyll cells adjacent to

aphid stylet tracks, whereas antisera to MpPIntO1 and MpC002 labelled the aphid

stylet sheaths. These data indicate that aphid effectors are delivered into the cytosol of

plant cells during probing in the pathway phase, whereas other effectors are

embedded within the sheaths that surround the stylets in the apoplastic space of

mesophyll tissue.

Results

Candidate effector proteins are detected in total protein extracts of aphid-

exposed leaves

Affinity purified antisera were raised against recombinant GPA candidate effectors

and examined for specificity and background levels in aphid and plant extracts by

protein immuno-blotting. The antisera to Mp10, MpOSD1, MpPIntO1 and MpC002

detected bands of the predicted sizes in total protein extracted from whole aphids

(Fig. 1). Antisera to the CSPs Mp10 and MpOS-D1 detected Mp10 and MpOS-D1,

respectively, down to the ng level (Supplementary Fig. S1), indicating that the

antisera to these proteins are specific and sensitive. The antisera to all four effectors

detected proteins of the predicted sizes in extracts of *A. thaliana* leaves previously

exposed to GPA (Fig.1), but not in unexposed leaves. These bands were not detected

in blots probed with corresponding pre-immune sera (Supplementary Fig. S2). The

Mp10 and MpOS-D1 antibodies and pre-immune sera reacted with some non-specific

bands of different sizes in the aphid-infested and uninfested plant tissues, but the

bands corresponding to the Mp10 and MpOS-D1 proteins were only detected in

infested tissues and not with the pre-immune sera (Fig. 1, Supplementary Fig. S2).

Thus, GPA delivers all four effectors into plant tissues during feeding.

The aphid effector protein Mp10 localizes to the cytoplasm and chloroplasts of

mesophyll cells adjacent to aphid feeding sites

To investigate if GPA delivers effectors in or near feeding sites, semi-thin sections of

GPA-infested leaf samples were stained with toluidine blue to localize GPA stylets

tracks (Fig. 2A). Ultra-thin sections of the same sample were then used for immuno-

gold-labelling (IGL) with the Mp10 antiserum (Fig. 2B-D). Controls included the

GPA-infested *A. thaliana* leaf samples sections incubated with pre-immune sera and

sections of uninfested *A. thaliana* leaf samples incubated with antisera (Fig, 3, 4 and

Supplementary Fig. S2).

IGL with Mp10 antiserum (dilutions 1:10 and 1:100) showed high labelling density in

cytoplasm and chloroplasts of mesophyll cells adjacent to the aphid stylets tracks in

two independently prepared GPA-infested leaf samples (Fig. 2B-D, 3A-B), whereas

only low density background labelling was observed with pre-immune sera (dilution

1:50) in the sections of aphid-infested leaves (Fig. 3C) or with Mp10 antisera

(dilution 1:10) in sections of uninfested control samples (Fig. 3D). Thus, despite some

aspecific labelling of plant proteins on immuno-blots (Fig. 1), the labelling of plant

tissue adjacent to the aphid feeding site is specific. Quantification of the labelling

seen on micrographs revealed consistently more labelling with Mp10 antisera of the

cytosol and chloroplasts with Mp10 antisera in mesophyll cells adjacent to aphid

feeding sites compared to the two control treatments (Fig. 3G, Supplementary Fig.

S3). Labelling of mesophyll cells distal to the site of feeding was not detected (Fig.

3E; Supplementary Fig. S3). Chloroplasts are known to be prone to high levels of

non-specific background labelling in IGL experiments, however we do not see similar

levels of labelling of plastids in the uninfested or pre-immune controls, or of plastids

distal to the site of feeding. Therefore, these data suggest that Mp10 localizes in the

cytosol and chloroplasts of mesophyll cells adjacent to the stylets tracks.

Occasionally we found that cells adjacent to the stylets' track contained unusual

features (Fig. 3 A, C). The cytoplasm of these cells showed reduced granularity and a

higher number of vesicle-like structures that may be endoplasmic reticulum (ER)-

derived. These were not seen in all aphid-infested samples (e.g. Fig. 3 B), but may

represent a response of the cell to aphid probing.

Because GPA feeds from the phloem sieve cells, we quantified Mp10 antisera

labelling of the vascular bundles closest to the aphid feeding site (Fig. 2 A; Fig. 3F).

We did not detect specific labelling of any cell type in the vascular tissue in

experimental compared to the control samples (Supplementary Fig. S2B) suggesting

that Mp10 is not delivered into the vasculature or is not present at high enough

concentrations for detection in these tissues.

We also investigated labelling with antisera to MpOS-D1, a family member of Mp10,

in and near stylets tracks. We did not detect specific labelling of mesophyll or

vascular tissues with MpOS-D1 antisera in or near GPA feeding sites compared to

controls (Supplemental Fig. S2C, D). While the Mp-OSD1 antisera detected

denatured MpOS-D1 protein on immuno-blots (Fig. 1), it is possible that these

antisera do not detect this effector in resin sections. Alternatively, GPA may not

deliver Mp-OSD1 or delivers this effector at low concentrations or to different

locations inside plant tissues compared to Mp10.

Aphid effector MpPIntO1 localizes to the sheaths surrounding aphid stylets at

feeding sites.

To compare localizations of aphid effectors, we also conducted IGL with antisera to

MpPIntO1 and MpC002 on ultra-thin sections of tissues containing aphid stylets

tracks. We observed very strong and specific labelling of the stylets sheaths by the

MpPIntO1 antisera (Fig. 4B; Suppl. Fig. 2G), but not by the corresponding pre-

immune serum (Fig. 4C; Suppl. Fig. 2G).The MpC002 antisera also labelled the

sheaths more strongly than other tissues in the plant and this labelling was stronger

than detected with the corresponding pre-immune sera (Fig. 4D-E; Suppl. Fig. 2E).

However, the background labelling of uninfested plant tissue by the MpC002 antisera

was much higher than for MpPIntO1, suggesting that anti-MpC002 may be less

specific (Suppl. Fig. 2E-H). These data suggest that MpPIntO1, and possibly also

MpC002, are found in the sheaths that surround the stylets during feeding.

Discussion

In this study, we generated specific antisera to the GPA proteins Mp10, MpOS-D1,

MpC002 and MpPIntO1, which were previously shown to have activities

characteristic of effector proteins (Mutti et al 2008; Bos et al. 2010; Pitino et al. 2011;

Pitino and Hogenhout 2013), and used these antisera in immuno-gold labelling

experiments to examine if GPA secretes effector proteins into plants during feeding.

We found that the Mp10 antisera specifically labelled the cytoplasm and chloroplasts

of mesophyll cells adjacent to the aphid stylet tracks. No such labelling was observed

in cells further removed from the stylets tracks. In addition, immuno-gold labelling

experiments with the antisera to the three other effectors did not label these

compartments. Finally, we observed no or much reduced labelling densities in the

pre-immune sera and uninfested controls. Taken together, these experiments provide

strong evidence for Mp10 delivery by GPA into mesophyll cells, where Mp10 resides

in the cytosol and the chloroplasts.

We previously found that Mp10 suppresses the flg22-induced ROS burst (Bos et al.

2010) which is dependent on BAK1 (Chinchilla et al. 2007). The ROS burst seen in

response to aphid elicitors is also dependent on BAK1 (Chaudhary et al. 2014; Prince

et al. 2014). BAK1 is a key regulator of several plasma membrane-localized pattern

recognition receptors (PRRs), which sense potential invaders early on and trigger

downstream immune signalling. Thus, given that we found Mp10 in the cytoplasm of

mesophyll cells adjacent to stylets tracks, this effector may suppress the ROS burst

induced by aphids shortly after initial probing by the aphid in the pathway phase. The

additional detection of Mp10 in chloroplasts is of interest because chloroplasts are a

major sources of ROS and have a critical role in PTI (Shapiguzov et al. 2012; Nomura

et al. 2012; Caplan et al. 2015).

In addition to the suppression of PTI, heterologous expression of Mp10 in plants

elicits an effector-triggered immunity (ETI)-type response, leading to chlorosis and

activation of JA- and SA- signalling pathways (Bos et al. 2010; Rodriguez et al.

2014). However, in these expression experiments Mp10 is present at high levels in

many cells, whereas our data indicate that Mp10 is delivered in only some cells near

the aphid stylets in the presence of other effectors (such as MpPIntO1). Hence, it

remains to be investigated how Mp10 modulates cell defense responses during aphid

feeding.

Our IGL data show different localizations of MpPIntO1 and MpC002 compared to

Mp10. MpPIntO1 and MpC002 are abundant proteins in aphid saliva and can be

readily detected by protein immuno-blotting in artificial diets fed upon by aphids and

extracts of aphid-exposed leaves (Mutti et al. 2008; Harmel et al. 2008). Our IGL data

provide evidence that MpPIntO1, and possibly also MpC002, are present in the stylets

sheaths. We did not detect MpPIntO1 inside plant mesophyll and vascular cells

indicating that aphids do not deliver this effector inside these cells or else they are

present at or below detection-level concentrations. Our data are in agreement with

previous findings showing that MpPIntO1 is abundant in the gelling saliva, which

generate the sheaths (Harmel et al. 2008). We previously found that aphids reproduce

better on transgenic *A. thaliana* which produce MpPIntO1 and MpC002 under control

of 35S promoters (Pitino and Hogenhout 2013), though the effect of MpPIntO1 on

aphid performance is variable (Pitino and Hogenhout 2013; Elzinga et al. 2014). In

addition, RNA interference (RNAi)-mediated knock down of MpC002 in aphids

reduced aphid performance on plants, whereas MpPIntO1-RNAi aphids were not

affected (Pitino et al. 2011). Several pathogens deliver effectors in the apoplastic

space to modulate defence responses (Tian et al. 2004, 2007; De Wit 2016). Our IGL

data open up the possibility that the aphid MpPIntO1 and MpC002 effectors act in the

plant apoplast from within sheath saliva.

In some cases, changes in subcellular structures occurs in response to aphid feeding

(Fig. 3, A, C). It is known that pathogen attack can induce dramatic reorganisations of

subcellular structures, including organelles (Hardham et al. 2008; Ben Khaled et al.

2015). Aphids can also cause significant tissue damage and disruption of plant tissue

during feeding (Saheed et al. 2007). Moreover, probing of cells by aphid stylets

induces a rapid subcellular relocalization of vesicle-associated *Cauliflower mosaic*

virus (CaMV) particles that is essential for the acquisition and transmission of this

virus by aphids (Martiniere et al. 2013). The increase in abundance of vesicular

structures we observed might be associated with the activation of plant defence

responses, such as the delivery of defence compounds via vesicles to deter attack.

Alternatively, aphid effectors may modulate plant cell processes, such as the

reprogramming of vesicle trafficking, as has been shown to occur for pathogens (Ben

Khaled et al. 2015; Bozkurt et al. 2015).

Materials and methods

Antibody generation

Coding sequences corresponding to the predicted mature effector proteins (minus

predicted secretory signal peptide) were expressed in *Escherichia coli* as *N*-terminal

6Xhis tag fusions (Mp10, MpPIntO1 and MpOSD1) or as a *N*-terminal dC, and 6XHis

tag fusion (MpC002), purified and checked for purity by protein immuno-blotting

using anti-His-tag antibody (BacPower and FoldArt technologies, Genscript). Proteins

were injected into chicken (Mp10) or rabbit (MpOSD1, MpPIntO1, and MpC002) by

Genscript. Specific antisera were affinity purified using immobilized recombinant

protein. Aliquots of pre-immune serum were collected from the animals before

injection.

Western blotting

Arabidopsis thaliana Col-0 plants were exposed to GPA (*Myzus persicae* clone O) for

24h (1000 aphids per 3-week old plant), aphids were carefully removed, and the plant

rosette was rinsed in distilled water to remove any remaining aphid material from the

surface. Total protein was extracted from plant and aphids in sodium dodecyl sulfate

(SDS) buffer (5 µl per mg plant tissue, or 200 µl per mg aphid tissue), and 10 µl was

size-separated by SDS-polyacrylamide gel electrophoresis and blotted onto

nitrocellulose membrane using standard methods. Blots were probed overnight with

1:100 or 1:1000 antibody dilutions (or with pre-immune sera at double the

concentrations), and a secondary antibody-horseradish peroxidase-conjugate.

Immuno-localisation

Samples were harvested from aphid-infested (or uninfested) *A. thaliana* plants and

immediately fixed in 4% formaldehyde / 0.5% glutaraldehyde in phosphate-buffered

saline (PBS) at 4 degrees overnight. The samples were then embedded in LR White

resin (Agar Scientific) using the progressive lowering of temperature (PLT) method

using the Leica EM AFS2 (Automatic Freeze Substitution) machine (Caillaud et al.

2014). Sections were prepared on a Leica UC6 ultramicrotome. Sections of 0.5 μm

were stained with toluidine blue for light microscopy (Nikon Microphot-SA) and

ultrathin sections of approximately 90 nm, were picked up on to pyroxylin- and

carbon-coated gold grids (Agar Scientific) for immuno-labelling.

The ultrathin sections were immuno-gold labelled according to the following

protocol: 50mM glycine/PBS for 15 mins; Aurion blocking buffer (Aurion, The

Netherlands for 30 mins; 0.1% acetylated bovine serum albumin (BSA-C) (Aurion,

The Netherlands) in PBS for 2 X 5 mins; primary antibody of choice at 1:10 to 1:100

dilutions in 0.1% BSA-C/PBS for 90 mins; 0.1% BSA-C/PBS for 6 X 5 mins;

secondary antibody conjugated to 10nm gold particles (1/40 dilution in 0.1% BSA-

C/PBS) for 90 mins; 0.1% BSA-C/PBS for 6 X 5 mins; PBS for 2 X 5 mins; and

water for 2 X 3 mins.

Grids were viewed in a FEI Tecnai 20 transmission electron microscope (FEI UK Ltd,

Cambridge, UK) at 200 kV and imaged using an AMT XR60 digital camera (Deben,

Bury St Edmunds, UK) to record TIF files. Images were analysed using ImageJ. The

areas and the numbers of gold particles detected in each subcellular compartment

were measured across a series of images across each sample. The density across

different compartments in infested/uninfested samples were analysed by general

linear model (GLM, Genstat). Where differences in labelling intensity were detected

between aphid-infested and uninfested samples, replicate sections were probed with

the corresponding pre-immune serum, at a higher concentration that the antibody was

used at, to demonstrate that labelling was specific.

Acknowledgements

We thanks the members of the Hogenhout lab for useful discussions and feedback,

and the BBSRC for funding (BB/J0045531/1).

Figure legends

Figure 1. Aphid salivary proteins are detected in extracts of aphid-infested host plant tissue.

Total protein extracts from whole aphids (GPA) (lane 8) and *A. thaliana* rosettes

uninfested (lanes 1-2, two different plants) or infested with GPA (lanes 3-5, three

different plants) were separated by SDS-PAGE, transferred to nitrocellulose

membranes and probed with antisera raised against the aphid proteins Mp10, MpOS-

D1, MpPIntO1 and MpC002 as shown at the left of the blots. The bottom panel shows

an amido-black stained loading control. Arrows at right point to bands of predicted

sizes for the effectors, as indicated. Arrowheads indicate non-specific bands detected

by antibodies.

Figure 2. Mp10 protein is detected in the cytoplasm of mesophyll cells near an aphid stylets track.

(A): Semi-thin section (0.5 μm) from an aphid-infested *A. thaliana* leaf stained with

toluidine blue. Aphid stylets tracks (ST) are visible between mesophyll cells (Mc) and

a vascular bundle (VB). (B-D): Immuno-gold labeling of ultra-thin sections (90 nm)

of the tissue shown in (A) with anti-Mp10 (1:10 dilution) revealed a high density of

gold particles (arrows in D) in the cytoplasm (Cy) of mesophyll cells (Mc). No

obvious labeling was observed of aphid stylets (St), cell wall (CW), intercellular

space (IS), mesophyll cell vacuole (V) and inside vesicles (Ve). Images shown in B to

D are sequentially higher magnifications of the same area of a cell adjacent to the

stylets track. Scale bars, 500 nm.

Figure 3. Mp10 detection is specific and restricted to the cells adjacent to stylets

tracks.

Ultra-thin sections from *A. thaliana* leaves infested with GPA, and labeled with anti-

Mp10 (1:100 dilution) show labelling (examples indicated with arrows) in the cytosol

"Cy" (A-B, G) and plastid "P" (B, G) of cells adjacent to the stylets track, but not

when labelled with pre-immune serum (1:50) (C, G). Samples from uninfested plants

are not labelled with anti-Mp10 above background levels (D, G). Labeling with anti-

Mp10 was not detected in mesophyll cells distal to the feeding site (E), or an area of

the vascular bundle closest to the feeding site in GPA-infested tissues (F).

Quantification of labelling density (G) based on measurements of compartment area

and gold particle count across multiple images of different areas from the samples.

Values show mean (+/- SEM) density (infested, antiMp10 (at 1:100 dilution) n=13;

infested, pre-immune (1:50) n=4, uninfested anti-Mp10 (1:100) n=5). Asterisks

indicate significant differences in labelling both between infested vs. uninfested

samples, and between anti-Mp10 and pre-immune labelled samples for that

compartment ($p<0.01$, GLM). Abbreviations: "IS" intercellular space; "Cy"

cytoplasm; "V" vacuole; "CW" cell wall; "Ve" vesicles; "P" plastid; "U" unidentified

compartment. In (F): "CC" is a phloem companion cell, and "SE" are phloem sieve

elements. Insets (A, C, D and F) are magnified regions of the image, all scale bars are

500 nm.

Figure 4. MpPIntO1 and MpC002 antisera label the stylets sheaths

Ultra-thin sections from aphid-infested samples showing the gelling saliva that forms

the stylets sheath ("ST"), immuno-gold labelled with anti-MpPIntO1 (A-B at 1:100

dilution) or with the corresponding pre-immune serum (C, 1:50); or with anti-

MpC002 (D, 1:100) or the corresponding pre-immune serum (E, 1:50). Examples of

gold-labelling, where present, are indicated with arrows. Compartments are indicated

as: "ST" stylet track; "IS" intercellular space; "Cy" cytoplasm; "V" vacuole; "CW"

cell wall; "Ve" vesicles; "P" plastid; "M" mitochondria. Scale bars all indicate 500

nm. Quantification of labelling density can be found in Supplemental Fig. 3.

Supplementary Figure 1. Mp10 and MpOSD1 antisera are specific and do not

cross react.

Immuno-blots of pure recombinant Mp10 and MpOSD1 proteins probed with antisera

raised against the proteins show the sensitivity and specificity of these antisera. (A)

Replicate blots of a range of quantity of Mp10 protein (1-25ng) and 25ng OSD1

protein probed with (from top) decreasing concentration of the anti-Mp10 serum

(from 1:500 to 1:10,000) or with corresponding pre-immune serum (at 1:1000

dilution). (B) Replicate blots of a range of quantities of OSD1 protein (1-25ng) and

25ng Mp10 protein probed with (from top) decreasing concentration of the anti-OSD1

serum (from 1:500 to 1:10,000) or with corresponding pre-immune serum (at 1:1000

dilution).

Supplementary Figure 2. Immuno-blots of aphid-exposed plant tissue probed

with antisera raised against candidate effector proteins and also with

corresponding pre-immune serum.

Immuno-blots of total protein extracts from Arabidopsis rosettes; uninfested (lanes 1-

2) or infested with GPA (lanes 3-5), and extract from GPA aphids (lane 9), were

probed with antisera raised against the aphid salivary proteins (as shown in Fig. 1),

and with the corresponding pre-immune sera (PI); (from the top) Mp10, Mp10-PI,

OSD1, OSD1-PI, MpPIntO1, MpPIntO1-PI, MpC002, and MpC002-PI. The bottom

panel shows loading of the plant extract samples by amido-black staining of the

membrane. Arrowheads indicate bands of the expected sizes detected in aphid-

exposed plants and aphids, but not unexposed plants. Arrows indicate presumably

non-specific bands of other sizes. Pre-immune sera were used at higher concentrations

than all corresponding antibodies to ensure that specific antibody-labelled bands were

not also detected by pre-immune serum.

Supplementary Figure 3. Quantification of immuno-gold labelling over different

cellular compartments and tissues. Graphs show the density of gold particles in

different subcellular compartments (mean +/- SEM (n= from 3 to 13)) for (left) cells

adjacent to the aphid stylet track (or equivalent mesophyll cells from uninfested

tissues), and (right) cells in the vascular bundle. Samples were probed with affinity-

purified antibody (all at 1:100 dilution) raised against Mp10 (A-B); MpOSD1 (C-D);

MpC002 (E-F) or MpPIntO1 (G-H), or (where indicated) with the corresponding pre-

immune serum (all at 1:50 dilution). Asterisks indicate significant difference in

labelling both between infested and uninfested samples, and between antibody and

pre-immune labelled samples for that compartment ($p < 0.01$, GLM). Plus signs

indicate significantly higher labelling of stylet track relative to other compartments in

infested tissue (for which no uninfested control plants could be tested) and for which

the signal was significantly higher than for infested samples probed with the

corresponding pre-immune serum ($p < 0.01$, GLM). "ND": Not determined.

LITERATURE CITED

Atamian, H.S., Chaudhary, R., Cin, V.D., Bao, E., Girke, T., and Kaloshian, I.

2013. In planta expression or delivery of potato aphid Macrosiphum euphorbiae

effectors Me10 and Me23 enhances aphid fecundity. Mol Plant Microbe Interact.

26:67-74.

Ben Khaled, S.B., Postma, J., and Robatzek, S. 2015. A moving view: subcellular

trafficking processes in pattern recognition receptor–triggered plant immunity.

Annual Review of Phytopathology. *53*:379-402.

Bos, J.I.B., Prince, D., Pitino, M., Maffei, M.E., Win, J., and Hogenhout, S.A.

2010. A functional genomics approach identifies candidate effectors from the aphid

species *Myzus persicae* (green peach aphid). PLoS Genet. *6:*e1001216

Bozkurt, T.O., Belhaj, K., Dagdas, Y.F., Chaparro-Garcia, A., Wu, C.H., Cano,

L.M., and Kamoun, S. 2015. Rerouting of plant late endocytic trafficking toward a

pathogen interface. Traffic. *16*:204-26.

Caillaud, M-C., Wirthmueller, L., Sklenar, J., Findlay, K., Piquerez, S.J.M.,

Jones, A.M., Robatzek, S., Jones, J.D., and Faulkner, C. 2014. The plasmodesmal

protein PDLP1 localises to haustoria-associated membranes during downy mildew

infection and regulates callose deposition. PLoS Pathog *10*:e1004496.

Caplan, J.L., Kumar, A.S., Park, E., Padmanabhan, M.S., Hoban, K., Modla, S.,

Czymmek, K., and Dinesh-Kumar S.P. 2015. Chloroplast stromules function during

innate immunity. Developmental Cell. *34* 45-57.

Carolan, J.C., Fitzroy, C.I.J., Ashton, P.D., Douglas, A.E., and Wilkinson, T.L.

2009. The secreted salivary proteome of the pea aphid *Acyrthosiphon pisum*

characterised by mass spectrometry. Proteomics. *9*:2457-2467.

Chaudhary, R., Atamian, H.S., Shen, Z., Briggs, S.P., and Kaloshian, I. 2014.

GroEL from the endosymbiont *Buchnera aphidicola* betrays the aphid by triggering

plant defence. Proc Natl Acad Sci USA. *111*:8919-8924.

Chaudhary, R., Atamian, H.S., Shen, Z., Briggs, S.P., and Kaloshian, I. 2015.

Potato aphid salivary proteome: Enhanced salivation using resorcinol and

identification of aphid phosphoproteins. J Proteome Res. *14*:1762-1778.

Chinchilla, D., Zipfel, C., Robatzek, S., Kemmerling, B., Nürnberger, T., Jones,

J.D.G., Felix, G., and Boller, T. 2007. A flagellin-induced complex of the receptor

FLS2 and BAK1 initiates plant defence. Nature. *448*:497-500.

Coleman, A.D., Wouters, R.H., Mugford, S.T., and Hogenhout, S.A. 2015.

Persistence and transgenerational effect of plant-mediated RNAi in aphids. J Exp Bot.

66, 541-8.

Cooper, W.R., Dillwith, J.W. and Puterka, G.J. 2010. Salivary proteins of Russian

wheat aphid (Hemiptera: Aphididae). Environ Entomol. *39*:223-231.

De Wit P.J. 2016. Apoplastic fungal effectors in historic perspective; a personal

view. New Phytol. 2016. doi: 10.1111/nph.14144

Dodds, P.N., and Rathjen, J.P. 2010 Plant immunity: towards an integrated view of

plant-pathogen interactions. Nat Rev Genet. *11*:539-548.

Elzinga, D., De Vos, M., and Jander, G. 2014. Suppression of plant defenses by a

Myzus persicae (green peach aphid) salivary effector protein. Mol Plant Microbe

Interact. *27*:747-756.

Guerrieri, E., and Digilio, M.C. 2008. Aphid-plant interactions: a review. Journal of

Plant Interactions. *3*:223-232.

Guo, K., Wang, W., Luo, L., Chen, J., Guo, Y., and Cui, F. 2014. Characterization

of an aphid-specific, cysteine-rich protein enriched in salivary glands. Biophys Chem.

189:25-32.

Hardham, A.R., Takemoto, D., and White, R.G. 2008. Rapid and dynamic

subcellular reorganization following mechanical stimulation of *Arabidopsis* epidermal

cells mimics responses to fungal and oomycete attack. BMC Plant Biology. *8*:63.

Harmel, N., Létocart, E., Cherqui, A., Giordanengo, P., Mazzucchelli, G.,

Guillonneau, F., De Pauw, E., Haubruge, E., and Francis, F. 2008. Identification

of aphid salivary proteins: A proteomic investigation of *Myzus persicae.* Insect Mol

Biol. *17*:165-174.

Hogenhout, S.A., Ammar,, E-D., Whitfield, A.E. and Redinbaugh, M.G. 2008.

Insect vector interactions with persistently transmitted viruses. Annual Review of

Phytopathology. *46*:327-359.

Martín, B., Collar, J.L., Tjallingii, W.F., and Fereres, A. 1997. Intracellular

ingestion and salivation by aphids may cause the acquisition and inoculation of non-

persistently transmitted plant viruses. J Gen Virol. *78*:2701-2705.

Martiniere, A., Bak, A., Macia, J.L., Lautredou, N., Gargani, D., Doumayrou, J.,

Garzo, E., Moreno, A., Fereres, A., Blanc, S., and Drucker, M. 2013. A virus

responds instantly to the presence of the vector on the host and forms transmission

morphs. Elife. *2013*:1-26.

Musser, R.O., Hum-Musser, S.M., Eichenseer, H., Peiffer, M., Ervin, G.,

Murphy, J.B., and. Felton G. W. 2002. Herbivory: Caterpillar saliva beats plant

defences. Nature *416*:599-600.

Mutti, N.S., Louis, J., Pappan, L.K., Pappan, K., Begum, K., Chen, M.S., Park,

Y., Dittmer, N., Marshall, J., Reese, J.C., Reeck, G.R. 2008. A protein from the

salivary glands of the pea aphid, *Acyrthosiphon pisum,* is essential in feeding on a

host plant. Proc Natl Acad Sci USA. *105*:9965-9969.

Naessens, E., Dubreuil, G., Giordanengo, P., Baron, O.L., Minet-Kebdani, N.,

Keller, H., and Coustau, C. 2015. A secreted MIF cytokine enables aphid feeding

and represses plant immune responses. Curr Biol. *25*:1898-1903.

Nicholson, S.J., Hartson, S.D., and Puterka, G.J. 2012. Proteomic analysis of

secreted saliva from Russian Wheat Aphid (*Diuraphis noxia* Kurd.) biotypes that

differ in virulence to wheat. J Proteomics. *75*:2252-2268.

Nicholson, S.J., and Puterka, G.J. 2014. Variation in the salivary proteomes of

differentially virulent greenbug (*Schizaphis graminum* Rondani) biotypes*. J

Proteomics. *105:*186-203.

Nomura, H., Komori, T., Uemura, S., Kanda, Y., Shimotani, K., Nakai, K.,

Furuichi, T., Takebayashi, K., Sugimoto, T., Sano, S., Suwastika, I.N., Fukusaki,

E., Yoshioka, H., Nakahira, Y., and Shiina, T. 2012. Chloroplast-mediated

activation of plant immune signalling in Arabidopsis. Nat Commun. *26*:926.

Orlovskis Z., Canale M. C., Thole V., Pecher P., Lopes J. R. S., Hogenhout S. A.

2015. Insect-borne plant pathogenic bacteria: getting a ride goes beyond physical

contact. Current Opinion in Insect Science. *9*:16–23.

Petre, B., and Kamoun, S. 2014. How do filamentous pathogens deliver effector

proteins into plant cells? PLoS Biol. *12*:e1001801.

Pitino, M., Coleman, A.D., Maffei, M.E., Ridout, C.J., and Hogenhout, S.A. 2011.

Silencing of aphid genes by dsRNA feeding from plants. PloS one *6*:e25709.

Pitino, M., and Hogenhout, S.A. 2013. Aphid protein effectors promote aphid

colonization in a plant species-specific manner. Mol Plant Microbe Interact. *26*:130-

139.

Prince, D.C., Drurey, C., Zipfel, C., and Hogenhout, S.A. 2014. The Leucine-Rich

Repeat Receptor-Like Kinase BRASSINOSTEROID INSENSITIVE1-

ASSOCIATED KINASE1 and the Cytochrome P450 PHYTOALEXIN DEFICIENT3

Contribute to Innate Immunity to Aphids in Arabidopsis. Plant Physiol. 164:2207-

2219.

Ramsey, J.S., Wilson, A.C., de Vos, M., Sun, Q., Tamborindeguy, C., Winfield,

A., Malloch, G., Smith, D.M., Fenton, B., Gray, S.M., and Jander, G. 2007.

Genomic resources for Myzus persicae: EST sequencing, SNP identification, and

microarray design. BMC Genomics 16:423.

Rao, S.A.K., Carolan, J.C., and Wilkinson, T.L. 2013. Proteomic profiling of

cereal aphid saliva reveals both ubiquitous and adaptive secreted proteins. PLoS One.

8:e57413.

541 **Rodriguez, P., Stam, R., Warbroek, T., and Bos, J.I.B.** 2014. Mp10 and Mp42

542 from the aphid species *Myzus persicae* trigger plant defenses in *Nicotiana*

543 *benthamiana* through different activities. Mol Plant Microbe Interact. *27*, 30-39.

544 **Saheed, S.A., Liu, L., Jonsson, L., and Botha, C.E.J.** 2007. Xylem – as well as

545 phloem – sustains severe damage due to feeding by the Russian wheat aphid. South

546 African Journal of Botany. *73*:593–599.

547 **Shapiguzov, A., Vainonen, J.P., Wrzaczek, M., and Kangasjärvi, J.** 2012. ROS-

548 talk – how the apoplast, the chloroplast, and the nucleus get the message through.

549 Frontiers in Plant Science. *3*:292.

550 **Stuart J.J., Chen M.S., Shukle R., and Harris M.O.** 2012. Gall midges (hessian

551 flies) as plant pathogens. Ann Rev Phytopathol. *50*:339–357.

552 **Tian, M., Huitema, E., da Cunha, L., Torto-Alalibo, T., and Kamoun, S.** 2004. A

553 kazal-like extracellular serine protease inhibitor from *Phytophthora infestans* targets

554 the tomato pathogenesis-related protease P69B*. The Journal of Biological

555 Chemistry. *279*:26370–26377.

556 **Tian, M., Win, J., Song, J., van der Hoorn, R., van der Knaap, E., and Kamoun,**

557 **S.** 2007. A *Phytophthora infestans* cystatin-like protein targets a novel tomato papain-

558 like apoplastic protease. Plant Physiology. *143*:364–377.

559 **Tjallingii, W.F., and Esch, T.H.** 1993. Fine structure of aphid stylet routes in plant

560 tissues in correlation with EPG signals. Physiol Entomol. *18*:317-328.

561 **Tjalingi, W.F.** 2006. Salivary secretions by aphids interacting with proteins of

562 phloem wound responses. Journal of Experimental Botany. *57*:739–745.

Vandermoten, S., Harmel, N., Mazzucchelli, G,, De Pauw, E., Haubruge, and E.,

Francis, F. 2014. Comparative analyses of salivary proteins from three aphid species.

Insect Mol Biol. *23*:67-77.

Wang, W., Dai, H., Zhang, Y., Chandrasekar, R., Luo, L., Hiromasa, Y., Sheng,

C., Peng, G., Chen, S., Tomich, J.M., Reese, J., Edwards, O., Kang, L., Reeck,

G., and Cui, F. 2015. Armet is an effector protein mediating aphid-plant interactions.

FASEB J. *29*:2032-2045.

Will, T., Tjallingii, W.F., Thönnessen, A., and van Bel, A.J.E. 2007. Molecular

sabotage of plant defence by aphid saliva. Proc Natl Acad Sci USA. *104*:10536-

10541.

Zhou, J.J., Vieira, F.G., He, X.L., Smadja, C., Liu, R., Rozas, J., and Field, L.M.

2010. Genome annotation and comparative analyses of the odorant-binding proteins

and chemosensory proteins in the pea aphid *Acyrthosiphon pisum*. Insect Mol Biol.

Suppl 2:113-22.

Figure 1. Aphid salivary proteins are detected in extracts of aphid-infested host plant tissue.
Total protein extracts from whole aphids (GPA) (lane 8) and *A. thaliana* rosettes uninfested (lanes 1-2, two different plants) or infested with GPA (lanes 3-5, three different plants) were separated by SDS-PAGE, transferred to nitrocellulose membranes and probed with antisera raised against the aphid proteins Mp10, MpOS-D1, MpPIntO1 and MpC002 as shown at the left of the blots. The bottom panel shows an amido-black stained loading control. Arrows at right point to bands of predicted sizes for the effectors, as indicated. Arrowheads indicate non-specific bands detected by antibodies.

Figure 2. Mp10 protein is detected in the cytoplasm of mesophyll cells near an aphid stylets track.

(A): Semi-thin section (0.5 µm) from an aphid-infested *A. thaliana* leaf stained with toluidine blue. Aphid stylets tracks (ST) are visible between mesophyll cells (Mc) and a vascular bundle (VB). (B-D): Immuno-gold labeling of ultra-thin sections (90 nm) of the tissue shown in (A) with anti-Mp10 (1:10 dilution) revealed a high density of gold particles (arrows in D) in the cytoplasm (Cy) of mesophyll cells (Mc). No obvious labeling was observed of aphid stylets (St), cell wall (CW), intercellular space (IS), mesophyll cell vacuole (V) and inside vesicles (Ve). Images shown in B to D are sequentially higher magnifications of the same area of a cell adjacent to the stylets track. Scale bars, 500 nm.

Figure 3. Mp10 detection is specific and restricted to the cells adjacent to stylets tracks.

Ultra-thin sections from *A. thaliana* leaves infested with GPA, and labeled with anti-Mp10 (1:100 dilution) show labelling (examples indicated with arrows) in the cytosol "Cy" (A-B, G) and plastid "P" (B, G) of cells adjacent to the stylets track, but not when labelled with pre-immune serum (1:50) (C, G). Samples from uninfested plants are not labelled with anti-Mp10 above background levels (D, G). Labeling with anti-Mp10 was not detected in mesophyll cells distal to the feeding site (E), or an area of the vascular bundle closest to the feeding site in GPA-infested tissues (F).

Quantification of labelling density (G) based on measurements of compartment area and gold particle count across multiple images of different areas from the samples. Values show mean (+/- SEM) density (infested, antiMp10 (at 1:100 dilution) n=13; infested, pre-immune (1:50) n=4, uninfested anti-Mp10 (1:100) n=5). Asterisks indicate significant differences in labelling both between infested vs. uninfested samples, and between anti-Mp10 and pre-immune labelled samples for that compartment (*p*<0.01, GLM). Abbreviations: "IS" intercellular space; "Cy" cytoplasm; "V" vacuole; "CW" cell wall; "Ve" vesicles; "P" plastid; "U" unidentified compartment. In (F): "CC" is a phloem companion cell, and "SE" are phloem sieve elements. Insets (A, C, D and F) are magnified regions of the image, all scale bars are 500 nm.

Figure 4. MpPIntO1 and MpC002 antisera label the stylets sheaths
Ultra-thin sections from aphid-infested samples showing the gelling saliva that forms the stylets sheath ("ST"), immuno-gold labelled with anti-MpPIntO1 (A-B at 1:100 dilution) or with the corresponding pre-immune serum (C, 1:50); or with anti-MpC002 (D, 1:100) or the corresponding pre-immune serum (E, 1:50). Examples of gold-labelling, where present, are indicated with arrows. Compartments are indicated as: "ST" stylet track; "IS" intercellular space; "Cy" cytoplasm; "V" vacuole; "CW" cell wall; "Ve" vesicles; "P" plastid; "M" mitochondria. Scale bars all indicate 500 nm. Quantification of labelling density can be found in Supplemental Fig. 3.

Supplementary Figure 1. Mp10 and MpOSD1 antisera are specific and do not cross react.
Immuno-blots of pure recombinant Mp10 and MpOSD1 proteins probed with antisera raised
against the proteins show the sensitivity and specificity of these antisera. (A) Replicate blots of a
range of quantity of Mp10 protein (1-25ng) and 25ng OSD1 protein probed with (from top)
decreasing concentration of the anti-Mp10 serum (from 1:500 to 1:10,000) or with corresponding
pre-immune serum (at 1:1000 dilution). (B) Replicate blots of a range of quantities of OSD1
protein (1-25ng) and 25ng Mp10 protein probed with (from top) decreasing concentration of the
anti-OSD1 serum (from 1:500 to 1:10,000) or with corresponding pre-immune serum (at 1:1000
dilution).

Supplementary Figure 2. Immuno-blots of aphid-exposed plant tissue probed with antisera raised against candidate effector proteins and also with corresponding pre-immune serum. Immuno-blots of total protein extracts from Arabidopsis rosettes; uninfested (lanes 1-2) or infested with GPA (lanes 3-5), and extract from GPA aphids (lane 9), were probed with antisera raised against the aphid salivary proteins (as shown in Fig. 1), and with the corresponding pre-immune sera (PI); (from the top) Mp10, Mp10-PI, OSD1, OSD1-PI, MpPIntO1, MpPIntO1-PI, MpC002, and MpC002-PI. The bottom panel shows loading of the plant extract samples by amido-black staining of the membrane. Arrowheads indicate bands of the expected sizes detected in aphid-exposed plants and aphids, but not unexposed plants. Arrows indicate presumably non-specific bands of other sizes. Pre-immune sera were used at higher concentrations than all corresponding antibodies to ensure that specific antibody-labelled bands were not also detected by pre-immune serum.

Supplementary Figure 3. Quantification of immuno-gold labelling over different cellular compartments and tissues. Graphs show the density of gold particles in different subcellular compartments (mean +/- SEM (n= from 3 to 13)) for (left) cells adjacent to the aphid stylet track (or equivalent mesophyll cells from uninfested tissues), and (right) cells in the vascular bundle. Samples were probed with affinity-purified antibody (all at 1:100 dilution) raised against Mp10 (A-B); MpOSD1 (C-D); MpC002 (E-F) or MpPIntO1 (G-H), or (where indicated) with the corresponding pre-immune serum (all at 1:50 dilution). Asterisks indicate significant difference in labelling both between infested and uninfested samples, and between antibody and pre-immune labelled samples for that compartment ($p < 0.01$, GLM). Plus signs indicate significantly higher labelling of stylet track relative to other compartments in infested tissue (for which no uninfested control plants could be tested) and for which the signal was significantly higher than for infested samples probed with the corresponding pre-immune serum ($p < 0.01$, GLM). "ND": Not determined.