**Pollinator species traits do not predict either response to**

**agricultural intensification or functional contribution**

Ignasi Bartomeus[1,*] Daniel P. Cariveau [2,3], Tina Harrison[2], Rachael Winfree[2]

[1] Estación Biológica de Doñana (EBD-CSIC), Sevilla, Spain.

[2] Rutgers University, The State University of New Jersey, New Brunswick, NJ

08901, USA

[3] Department of Entomology, University of Minnesota, Saint Paul, MN 55113,

USA

[*] Correspondence author: nacho.bartomeus@gmail.com

**Running headline:** Pollinator traits are not predictive

**Summary:**

1. The response and effect trait framework, if supported empirically, would

provide for powerful and general predictions about how biodiversity loss will

lead to loss in ecosystem function.

2. This framework proposes that species traits will explain how different

species respond to disturbance (i.e. response traits) as well as their

contribution to ecosystem function (i.e. effect traits). However, predictive

response and effect traits remain elusive for most systems.

3. Here, we present detailed data on crop pollination services provided by

native, wild bees to explore the role of six commonly used species traits in

determining how crop pollination is affected by increasing agricultural

intensification. Analyses were conducted in parallel for three crop systems

(watermelon, cranberry, and blueberry) located within the same geographical

region (mid-Atlantic USA).

4. Bee species traits did not strongly predict species' response to any

agricultural intensification process, and the few traits that were weakly

predictive were not consistent across crops. Similarly, no trait predicted

species' overall functional contribution in any of the three crop systems,

although body size was a good predictor of per capita efficiency in two

systems.

5. Pollinator traits may be useful for understanding ecological processes in

some systems, but thus far we are unable to make generalizable predictions

regarding species responses to land-use change and its effect on the delivery

of ecosystem services.

**Keywords:** Biodiversity, bees, ecosystem services, ecosystem function,

response traits, effect traits, body size, diet specialism.

**Introduction**

Land-use change, along with other human-induced global change drivers, is

accelerating the rates of extinction of most taxa (Ellis et al. 2010). At the same

time, humanity relies on ecosystem services that wild species deliver, such as

pollination and pest control by insects, and nutrient cycling by microorganisms

(Cardinale *et al.* 2012). Thus, it is important to understand the relationship

between biodiversity loss and ecosystem service delivery (Schwartz et al.

2000). In particular, making generalizable predictions regarding how the

decline or local extinction of taxa will affect ecosystem services will allow for

targeted conservation actions to ameliorate negative impacts of land-use

change.

One avenue for predicting the functional consequences of biodiversity loss is

the response and effect trait framework (Lavorel & Garnier 2002, Naeem &

Wright 2003, McGill et al. 2006). Local extinction does not occur at random

because extinction risk is dependent on the species' characteristics.

Identifying which traits govern species responses to particular threats

('response traits') would provide the first step for predicting future species

loss. Furthermore, the magnitude by which ecosystem function declines when

a species is lost depends on that species' functional contribution. This, too, is

likely to be mediated by the species' traits ('effect traits'). Therefore, the

relationship between response and effect traits will mediate the magnitude of

the impact of human disturbance on ecosystem services (Schleuning, Fründ &

García 2015). For example, if the same species traits that are associated with

high function are also most sensitive to disturbance, ecosystem function

would be predicted to decline rapidly (Larsen, Williams & Kremen 2005).

However, for the response-effect trait framework to be generalizable, it is first

necessary to identify response and effect traits that are both explanatory and

possible to measure in the field (Cadotte, Carscadden & Mirotchnick 2011).

While a few generalities have emerged as to which traits make animal species

at greater risk of local decline, including dietary or habitat specialization and

body size (Fisher & Owens 2004; Ockinger *et al.* 2010), the correlation

between these response traits and extinction risk has been found to be weak,

variable, or context-dependent (Devictor, Julliard & Jiguet 2008; Powney *et al.*

2014; Fritz, Bininda-Emonds & Purviis 2009). Similarly, although some effect

traits have been identified, they are often weakly predictive, and their identity

varies by function and taxonomic group (Gagic *et al.* 2015). Lastly, within the

functional trait field as a whole, most progress has been made in identifying

functional traits for plants (Diaz *et al.* 2016), while little is known for animals

(Didham et al. 2016).

Here, we seek to identify response and effect traits for wild bee species

providing a key ecosystem service, crop pollination. The yield of most crop

plants increases with animal pollination (Klein *et al.* 2007). While managed

honey bees are a leading crop pollinator, wild insects contribute more than

half of pollinator visits to crop flowers across more than 40 crop systems

worldwide (Rader *et al.* 2016). A major threat to pollinators is habitat

destruction, primarily conversion to agriculture (Garibaldi *et al.* 2011), which is

also a leading cause of species loss worldwide (Pereira *et al.* 2010). Thus

agricultural land use has the potential to affect the ecosystem service upon

which agriculture itself depends (Deguines et al. 2014).

Our data sets were collected and analyzed in parallel and come from three

crop systems (watermelon, cranberry and blueberry) located within the same

geographical region (mid-Atlantic USA), but pollinated by distinct bee

communities. We determined whether six commonly-used species traits can

predict 1) species' responses to agricultural intensification (response traits)

and/or 2) species' contributions to crop pollination (effect traits).

**Material and methods:**

*Study system*

We selected 49 sites across three study systems that were located throughout

New Jersey and eastern Pennsylvania (USA). Watermelon sites (N = 17)

were located in 90 x 60 km region central New Jersey and Eastern

Pennsylvania, where the main types of land use are agriculture and suburban

development, interspersed with highly fragmented deciduous forest.

Cranberry and blueberry sites (N = 16 each) were both located within a 35 x

55 km area in southern New Jersey, where the main land cover types are

pine-oak ericaceous heath and agriculture. All sites in all systems were

separated by at least 1 km (range, watermelon: 2-90 km, cranberry: 1-32 km,

blueberry 1-38 km).

All three of crops are highly dependent upon bee pollination for marketable

fruit production (Klein *et al.* 2007). Commercial honey bees are used in most

of our study fields. However, honey bees are primarily managed hives, moved

throughout the region, and only found on sites during bloom. Therefore, honey

bees are not influenced by land cover in the same manner as wild bees and

are not used in our analyses. Wild bees are important pollinators in all three

systems (mean percentage of wild bee visits: 73% watermelon, 25%

cranberry, and 14% blueberry).

*Data collection:*

At all sites on all three crops, we used hand-netting to measure overall bee

abundance and species richness. To collect bees, we walked along fixed

transects at standard times of day and collected all bees observed to be

visiting flowers. In watermelon and blueberry, bees were netted three times

throughout the day for 20 minutes per transect (60 minutes per date per site)

and twice each day in cranberry for 30 minutes per transect (120 minutes per

date per site). Data were collected during the peak bloom in 2010

(watermelon: July, cranberry: late-May-early July, blueberry: April-early May).

Data were collected on three days per site for watermelon and blueberry and

two days per site for cranberry. Detailed methods can be found in Benjamin

Reilly & Winfree (2014), Winfree *et al.* (2015), Cariveau *et al.* (2013).

*Land cover characteristics of sites*

To relate pollinator response traits to land use, we required high-quality land

cover data for each pollinator collection site. For the cranberry and blueberry

sites in New Jersey, we used a continuous polygon layer classified by visual

photograph interpretation into 60 categories, at a minimum mapping unit of

4047 $m^2$ (1 acre; GIS Data provided by the New Jersey Department of

Environmental Protection). For watermelon sites that extend from central New

Jersey into Pennsylvania, we created a similar land cover data layer by

manually digitizing Google Earth imagery and visually classifying 15

categories, at a minimum mapping unit of 5,000 $m^2$ (1.24 acres). As each crop

was analyzed separately, our results are robust to using different land cover

data. However, to simplify the interpretation of results for the three crops, we

reclassified all land cover data into the following 7 broad categories:

agriculture, open managed (for example, mowed grass), open natural or semi-

natural (for example, old fields), semi-urban (<30% impervious surface), urban

(>30% impervious surface), wooded, and open water.

Prior to doing our spatial analyses, we explored several different landscape

characteristics, because agricultural land-use change can occur at multiple

scales and affect multiple landscape attributes. For each data collection site

we calculated four land cover variables: a) percent agriculture, b) percent

natural and semi-natural open habitat, c) forest edge length, and d) habitat

heterogeneity. We used agricultural land cover as our primary land use

change variable, as it is the dominant anthropogenic habitat type in all three

study systems. Percent open natural/semi-natural habitat makes up a small

proportion of the total land cover (mean 6.8% cover at 1500 m radius) but is

likely to be disproportionately important as forage and nesting habitat for bees

(Kleijn *et al.* 2006). Forest edge length may represent important habitat for

bees that nest in woods but forage in open areas (Kells & Goulson 2003).

Diverse patchy landscapes may be beneficial to bees as they may nest and

forage in different habitats (Kremen *et al.* 2007); thus we include Shannon H

diversity of the seven cover categories as a measure of habitat heterogeneity.

We calculated values for all four land cover variables at both a small scale

(300 m radius) and a large scale (1500 m radius), which correspond to typical

flight distances of small- and large-bodied bees, respectively (Greenleaf et al.

2007).

*Pollinator function*

To estimate the pollination services provided per bee species, we measured

two variables in the field, flower visitation frequency and per visit efficiency. As

variation in visitation frequency may be a function of land use at individual

farms, we use species abundances for each species at the site with its highest

abundance for each crop. Hence, we assess visitation frequency at its

maximum, which represents the optimal visitation frequency for each species.

To measure the pollination efficiency we quantified single-visit pollen

deposition by presenting virgin flowers to individual bees foraging on the

target crop. After visitation, we counted the number of pollen grains deposited

per flower visit (watermelon) or the number of pollen tetrads with pollen tubes

per flower visit (cranberry and blueberry). Because species identification in the

field is not possible for most bees and net collecting immediately after visits is

generally not possible, for the measurement of pollination efficiency we

grouped bees in species groups. Each group consisted of between one and

27 species, with the median number of species per group being 4 species

(Supporting Information, Table S1). Control flowers were left bagged until the

end of the field day, and contained few pollen grains (watermelon mean = 3

grains, N = 40 stigmas; cranberry mean = 0 tetrads, N = 82 stigmas; blueberry

mean = 2 tetrads, N= 734 stigmas). For detailed methods see Benjamin *et al.*

*(*2014), Winfree *et al.* (2015), Cariveau *et al.* (2013).

*Species traits*

Bee species vary in a number of traits that are associated with their response

to land-use change (Williams et al. 2010). Moreover, these traits will likely

affect the pollinator contribution to function, either by modifying its abundance

or because they are related to its per capita effectiveness. We obtained

detailed natural history data on 6 traits for the 90 bee species in our study: a)

sociality (solitary, facultative social, eusocial), b) nesting placement (hole,

cavity, stem, wood, ground), c) brood parasite (yes, no), d) body size, e) diet

breadth (level of specialization) and f) tongue length.

We obtained the trait data as follows. Species sociality level, nesting behavior

and brood parasite status were extracted from the literature (Bartomeus *et al.*

2013a). Body size (estimated from intertegular span, IT; Cane 1987) was

measured in the lab using collected specimens that had been identified to the

species level by professional taxonomists. Multiple specimens were measured

per species (mean = 6.6 specimens ± 3 S.E.) and the mean across the

measured specimens was used as the value for the species. Bee body size

also correlates strongly with foraging distance (Greenleaf *et al.* 2007), and

thus is ecologically related to mobility. Tongue length was measured in the lab

for 7.7 ± 1.2 SE specimens per species, and the mean across the measured

specimens is used. For the 40 specimens for which we cannot obtain a

tongue measure, we estimated tongue length from the species' body size and

phylogeny using an allometric equation (Cariveau et al. 2016).

Diet breadth was calculated using six independent datasets previously

collected at 139 sites throughout the study region by the Winfree laboratory

group. Each data set consists of individual pollinator specimens that were net-

collected while foraging on a flowering plant species; both pollinator and plant

were then identified to the species level. Those datasets comprise overall 393

pollinator species, and 392 plant species, with 3890 plant-pollinator

interactions (Supporting Information. Text S2). Prior to calculating diet

breadth, we rarefied the data to 20 visitation records per bee species, to avoid

confounding rarity with specialization (Blüthgen *et al.* 2008; Winfree *et al.*

2014). Nine species had fewer than 20 records and we were unable to

estimate diet breadth in the manner described above. Five of these species

are known to be specialized and we simulated the diet breadth index of 20

individuals visiting the known host plants. The four other species are known to

be generalists and we therefore used the mean diet breadth of its genus.

These four species were extremely rare (< 5 records each) in our analyzed

dataset.

To calculate diet breadth for each bee species, we considered the number of

plants species as well as the phylogenetic breadth that the bees fed upon by

using a rarefied phylogenetic diversity index(Nipperess & Matsen 2013). To

determine phylogenetic distances among plants, we first constructed a

general phylogenetic tree using the PHYLOMATIC "megatree" (version

R201120829, Chamberlain & Szocs 2013) which defines relationships

between higher plants (Webb, Ackerly & Kembel 2008). We then dated nodes

across this tree according to Wikström et al. (2001) and used the branch-

length adjustment algorithm BLADJ to estimate the age of all remaining,

undated nodes. Though this procedure implies that ages within our

phylogenies should be treated as approximations (Beaulieu *et al.* 2007),

previous analysis indicates marked improvements of phylogenetic analyses

when even a limited number of nodes are properly dated (Webb 2000).

*Statistical analysis*

**Response traits:** In order to investigate which traits are associated with

environmental variables related to agricultural intensification, we used fourth-

corner analysis (Legendre Galzin & Harmelin-Vivien 1997). This analysis is

specifically designed to directly analyze the relationship between the

biological characteristics of species and the characteristics of the habitat

where they are found.

The fourth-corner approach links an R matrix of environmental variables to a

Q matrix of species traits through an L matrix of species abundance (Dray &

Legendre 2008). A correlation coefficient is computed for each pair of species

traits and environmental variables. We tested the significance of the trait-

environment correlation by combining two permutation models (Dray &

Legendre 2008). The first permutation randomizes site rows (or equivalently,

environmental variables across sites) to test if communities in different

environments have different trait values. The second permutation randomizes

species columns (or equivalently, trait values across species) to test if trait-

environment relationships are independent of community composition (Dray &

Legendre 2008). Under the second null model, a consistent trait-environment

relationship will not be significant if communities in similar environments have

similar species composition. Thus even if the first permutation is significant

but the second is not, we cannot know if the observed trait-environment

correlation simply reflects random changes in trait values due to the shift in

species composition. Hence, as recommended by Dray & Legendre (2008),

we report the highest p-value of the two models. We the ran the fourth-corner

analysis for each crop independently using 9999 permutations for each null

model (R package "ade4"; Dray & Dufour 2007). No adjustments for p-values

were made, but due to the high number of tests performed in the analyses,

the possibility of significance by chance increases. Therefore, we mainly

discuss effect sizes and only highlight results with significance level $\leq 0.01$,

while we consider $0.01 > \alpha \leq 0.05$ to be relatively weakly supported

associations.

**Effect trait analysis:** To determine which traits influenced functional

contribution of each species we ran separate generalized linear models with

either visitation or per capita efficiency as response variables. Species traits

were predictors. The best model based on AICc was selected. When

differences between the best models were less than 2 we selected the simpler

model. The analysis for efficiency was done at the species group level (see

above: pollination function section). In order to obtain traits at the species

group level we calculated the mean values over species belonging to the

same group, weighted by the species mean abundance within the group. For

categorical variables we chose the dominant level.

All statistical analyses were performed in R software, version 3.0.3 (R Core

Team 2014).

**Results**

**Response traits**: Overall, we did not find a strong correlation between any

ecological traits and the environmental variables analyzed. Some traits

exhibited weak responses but they were not consistent across crops. For

watermelon, small bees and those with short tongues tended to decline with

increasing percentage of agriculture at 300m radius (Body size: r = 0.23, p =

0.03, Tongue length: r = 0.23, p = 0.03). Species nesting in stems were

positively associated with heterogeneous landscapes at 1500m radius (Stem

nesters: r = 0.12, p = 0.04). For cranberry and blueberry, we did not find any

trait significantly associated with any of the landscape characteristics. A

complete list of all comparisons is presented in Supporting Information (Table

S3).

**Effect traits:** As for response traits, no traits were highly predictive of either

visitation frequency or per visit efficiency across crops. For watermelon, no

trait was selected on the best model for visitation frequency. However,

pollination efficiency was positively correlated with body size and tongue

length ($R^2 = 0.75$, $F_{2,9} = 17.07$, $p < 0.001$, Fig 2A). For cranberry, visitation

frequency was positively related to cavity nesters ($R^2 = 0.38$, $F_{4,36} = 7.1$, $p <$

$0.0001$, Fig 2B). This result was driven by *Bombus* species, which are the only

cavity nesters in this data set. In contrast, efficiency per visit was not related

to any trait. For blueberry, visitation frequency was positively related to diet

specialism ($R^2 = 0.37$, $F_{1,20} = 13.5$, $p = 0.001$, Fig 2C), while efficiency per

visit is positively related to tongue length ($R^2 = 0.70$, $F_{1,5} = 14.9$, $p = 0.01$,

Fig 2D). Model selection, can be found in Supporting Information (Table S4).

**Discussion:**

Identifying traits that characterize which species are more sensitive to land-

use change or those that are functionally important is complex. We found

evidence for response and effect traits but they differed among crop species

as well as landscape variable used. Therefore, while some traits may be

important in some contexts, no traits were generalizable enough to be used to

predict how land-use change will influence the delivery of pollination services

in other systems. Further, the relationships identified were weak, especially for

response traits. This does not negate the importance of traits for

understanding which mechanisms underlie species responses to land use

change or pollination effectiveness, but it does suggest the commonly used

traits we studied here are not suitable for predictive purposes.

Being able to identify strong response traits would be a key tool for

understanding extinction risk, and an asset for conservation planning.

However, characterizing extinction risk based on traits is challenging. Despite

some generalities that emerge across taxa, with rare species, big species,

specialists, and higher trophic levels being in general more sensitive to

disturbances (Fisher & Owens 2004), there is a large variation in the response

of the species with those traits (Fritz, Bininda-Emonds & Purvis 2009; Seguin

*et al.* 2014). Work specifically on native bees has found that traits such as

specialization, body size, and sociality may predict responses to land use

(Winfree *et al.* 2009; Bommarco *et al.* 2010; Williams *et al.* 2010; Bartomeus

*et al.* 2013b; De Palma *et al.* 2015). However, studies often find contrasting

results. For example, De Palma et al. (2015) analyzed over 70,000 wild bee

records and found that small species were most sensitive to agricultural land

use, while others have found that larger species are more sensitive to

agricultural land use and/or environmental change generally (Larsen, Williams

& Kremen 2005; Bartomeus *et al.* 2013b; Rader *et al.* 2014), and some have

found little effect of body size (Williams et al. 2010). Here, we only found a

weak trend for big species to be more abundant in locally intense agricultural

areas in watermelon, but this trend disappears when land use is measured at

larger scales. Another trait, dietary specialization, is one of the few traits that

has been generally linked to increased species sensitivity to environmental

change (Williams *et al.* 2010; Scheper *et al.* 2014; De Palma *et al.* 2015), but

here we found that floral specialist bees did not decline with intensifying

agriculture. If anything, specialization is positively associated with agricultural

intensification in the blueberry system (Table S3), largely because the most

abundant bee species in this system (*Andrena bradleyi*) is a specialist on the

blueberry genus *Vaccinium*. Specialist bees observed in crop systems are

likely to be specialized on the crop plant family as was the case in our data

(e.g. *Habropoda sp.* , *Andrena bradleyi* in blueberry, but also *Peoponapis sp.*

in watermelon and *Mellita americana* in cranberry). We would expect different

responses from study designs that include natural habitat and a larger range

of specialist host plants (Forrest et al. 2015, Bartomeus & Winfree 2013).

Effect traits have been even harder to identify for pollinators. The limited data

published on particular plants suggests insects with larger bodies tend to

deposit more pollen per flower visit, but this pollen was not well distributed on

the stigma (Hoehn *et al.* 2008), or that the correlation between body size and

per visit pollination function is low (Larsen, Williams & Kremen 2005). Our

study supports the positive correlation between body size and per-visit pollen

deposition in both watermelon and blueberry (although note that tongue

length is correlated with body size in blueberry r = 0.76), but not for cranberry.

Hence, generality is difficult to achieve because a single pollinator trait, like

big body size, may not lead to high pollination function in all contexts. Rather it

seems likely that the most efficient trait will depend on the crop (Garibaldi *et*

*al.* 2015). Moreover, the total pollination provided by a pollinator species is the

product of visitation frequency and per capita efficiency (Kremen *et al.* 2005),

two processes that may be governed by different traits.

If generalizable response and effect traits can be found, the final step will be

to link response and effects to predict changes in ecosystem services. A

positive association between the response and effect traits (Naeem & Wright

2003) such that species with the strongest response to environmental change

also had the strongest effect on function, indicates the land-use change has

the potential for dramatic effects on ecosystem function. Whether response

and effect traits are in general positively, negatively, or uncorrelated is an

important question that has not yet been answered (Larsen, Williams &

Kremen 2005). Despite the conceptual elegance of the response-effect trait

framework, it is only effective if it is predictive, and strong evidence for the

generality of traits has not yet been found. For example, even the very

thorough and rigorously analyzed study of response-effect relationships by

Larsen, Williams & Kremen (2005) is based on a non-significant weak

relationship between pollinator per visit efficiency and body size. In our study,

even the strongest correlations find for watermelon, where big species are

less sensitive to local agricultural intensification and more efficient per visit,

but not more frequent flower visitors than smaller species are too weak to be

useful for predictive purposes.

Predictive response and/or effect traits are often assumed in the larger

literature as well. For example, recent re-evaluations of community stability in

food webs shows that using body size as proxy of extinction risk changes the

outcome of the stability simulations (Brose *et al.* 2016). However, the

assumption that body size is a good predictor of extinction risk is not directly

validated. Given the correlation showing that bigger species are more

sensitive is usually weak (Fisher & Owens 2004), these kind of approaches

could produce misleading outcomes.

Currently trait data may be too coarse to reveal ubiquitous response and

effect traits for four reasons. First, some traits may simply reflect identity of

genera or higher taxonomic groups. For example, some bumble bee species

in our three systems (especially *B. impatiens*) are common, functionally

dominant, and robust to extinction (Cariveau *et al.* 2013, Winfree *et al.* 2015).

Some of the important response and effect traits that we found, such as cavity

nesting and body size, may simply be proxies for bumble bees. Bumble bee

species also share other traits (e.g. sociality) that are commonly used in trait

analyses. Therefore, studies that don't include phylogenetic correlations may

be simply characterizing the general relationship between disturbance and the

functionally dominant bumble bees, or other dominant taxa. As there is a great

variability in the responses to disturbance among bumble bee species

(Cameron *et al.* 2011; Bartomeus *et al.* 2013b, Persson *et al.* 2015) this may

also explain why some studies find big species to be more sensitive to land

use change (Larsen, Williams & Kremen 2005) and other studies find the

opposite (Rader *et al.* 2014, this study). Second, traits may interact in complex

ways and single traits may be not able to capture responses and functional

contributions across species (e.g. Bommarco *et al.* 2010, De Palma *et al.*

2015). Third, phenotypical variability within species, usually ignored in trait-

based approaches, may play a more important role than previously though

(Bolnik et al. 2011). Finally, the most important traits may not have been

studied. Response traits such as dispersal ability, fecundity, and nest

microclimate/soil type, and effect traits like floral visitation behavior or

hairiness may be better predictors than the traits we have now. However, if

these traits are not easy to measure across bee species, they may be of little

use. Traits databases that include an increasing number of traits and agreed-

upon measurement techniques similar to those used in plant ecology (Kattge

*et al.* 2011) but that are also open-access may lead to significant

advancements in functional trait ecology in wild bees.

There is a call to be more predictive in ecology (Petchey *et al.* 2015). The use

of traits to predict species responses and subsequent changes in ecosystem

services is a potentially powerful approach. This is especially the case for

organisms such as insects where species identification is challenging and

detailed species-level natural history information is lacking. The ability to

effectively use a trait framework is becoming controversial because studies

thus far have not clearly related specific traits to specific threats or functions

(Didham et al. 2016; Shipley et al. 2016). A growing number of studies are

working to address the complexity and increase the predictability of this

framework (e.g. Laughlin & Messier 2015). However, until these approaches

yield consistent patterns across systems, site-specific species identity and

monitoring may at present be the best measure for predicting changes in

ecosystem services as a result of land-use change. A few dominant species

often drive ecosystem functioning (Kleijn *et al.* 2015; Winfree *et al.* 2015).

Identifying the sensitivity of the functionally dominant species may be the best

proxy thus far for predicting effects of species loss in ecosystem function.

**Acknowledgments:**

We thank Scott Chamberlain for help with BLADJ. IB was supported by

project Beefun (PCIG14-GA-2013-631653) funded by EU.

**Data Accessibility:** All data and code used in this manuscript is accessible in

github (https://github.com/ibartomeus/RE_traits) and will be archived on

acceptance in figshare.

**References:**

Bartomeus, I., Ascher, J.S., Gibbs, J., Danforth, B.N., Wagner, D.L., Hedtke, S.M. *et al.* (2013a) Data from: Historical changes in northeastern US bee pollinators related to shared ecological traits. Dryad Data Repository.

Bartomeus, I., Ascher, J.S., Gibbs, J., Danforth, B.N., Wagner, D.L., Hedtkee, S.M. *et al.* (2013b) Historical changes in northeastern US bee pollinators related to shared ecological traits. *Proceedings of the National Academy of Sciences, USA,* **110,** 4656–4660.

Bartomeus, I. & Winfree, R. (2013) Pollinator declines: reconciling scales and implications for ecosystem services. *F1000Res, 2,* 146.

Beaulieu, J.M., Leitch, I.J. & Knight, C.A. (2007) Genome size evolution in relation to leaf strategy and metabolic rates revisited. *Annals of Botany, 99,* 495-505.

Benjamin, F.E., Reilly, J.R. & Winfree, R. (2014) Pollinator body size mediates the scale at which land use drives crop pollination services. *Journal of Applied Ecology,* **51,** 440-449.

Blüthgen, N., Frund, J., Vazquez, D.P. & Menzel, F. (2008) What do interaction network metrics tell us about specialization and biological traits? *Ecology,* **89,** 3387-3399.

Bommarco, R., Biesmeijer, J.C., Meyer, B., Potts, S.G., Poyry, J., Roberts, S.P. *et al.* (2010) Dispersal capacity and diet breadth modify the response of wild bees to habitat loss. *Proceedings of the Royal Society of London, Series B: Biological Sciences,* **277,** 2075-2082.

Bolnick, D.I., Amarasekare, P., Araújo, M.S., Bürger, R., Levine, J.M., Novak, M., Rudolf, V.H., Schreiber, S.J., Urban, M.C., Vasseur, D.A. (2011) Why intraspecific trait variation matters in community ecology. *Trends in ecology & evolution.* **26**(4):183-92.

Brose, U., Blanchard, J.L., Eklof, A., Galiana, N., Hartvig, M., M, R.H. *et al.* (2016) Predicting the consequences of species loss using size-structured biodiversity approaches. *Biological Reviews of the Cambridge Philosophical Society,* Epub ahead of print.

Cadotte, M.W., Carscadden, K. & Mirotchnick, N. (2011) Beyond species: functional

diversity and the maintenance of ecological processes and services. *Journal*
*of Applied Ecology,* **48,** 1079-1087.

Cameron, S.A., Lozier, J.D., Strange, J.P., Koch, J.B., Cordes, N., Solter, L.F. *et al.*
(2011) Patterns of widespread decline in North American bumble bees.
*Proceedings of the National Academy of Sciences, USA,* **108,** 662-667.

Cane, J. (1987) Estimation of bee size using intertegular span (Apoidea). *Journal of*
*the Kansas Entomological Society,* **60,** 145-147.

Cardinale, B.J., Duffy, J.E., Gonzalez, A., Hooper, D.U., Perrings, C., Venail, P. *et al.*
(2012) Biodiversity loss and its impact on humanity. *Nature,* **486,** 59-67.

Cariveau, D.P., Nayak, G.K., Bartomeus, I., Zientek, J., Ascher, J.S., Gibbs, J. *et al.*
(2016) The allometry of bee proboscis length and its uses in ecology. *PLoS*
*One,* **11,** e0151482.

Cariveau, D.P., Williams, N.M., Benjamin, F.E. & Winfree, R. (2013) Response
diversity to land use occurs but does not consistently stabilise ecosystem
services provided by native pollinators. *Ecology Letters,* **16,** 903-911.

Chamberlain, S.A. & Szocs, E. (2013) taxize: taxonomic search and retrieval in R.
*F1000Res,* **2,** 191.

De Palma, A., Kuhlmann, M., Roberts, S.P.M., Potts, S.G., Börger, L., Hudson, L.N.
*et al.* (2015) Ecological traits affect the sensitivity of bees to land-use
pressures in European agricultural landscapes. *Journal of Applied Ecology,*
**52,** 1567–1577.

Deguines, N., Jono, C., Baude, M., Henry, M., Julliard, R. & Fontaine, C. (2014)
Large-scale trade-off between agricultural intensification and crop pollination
services. *Frontiers in Ecology and the Environment,* **12,** 212-217.

Devictor, V., Julliard, R. & Jiguet, F. (2008) Distribution of specialist and generalist
species along spatial gradients of habitat disturbance and fragmentation.
*Oikos,* **117,** 507-514.

Diaz, S., Kattge, J., Cornelissen, J.H., Wright, I.J., Lavorel, S., Dray, S. *et al.* (2016)
The global spectrum of plant form and function. *Nature,* **529,** 167-171.

Didham, R.K., Leather, S.R., Basset, Y. & Stewart, A. (2016) Circle the bandwagons
- challenges mount against the theoretical foundations of applied functional
trait and ecosystem service research. *Insect Conservation and Diversity,* **9,** 1-
3.

Dray, S. & Dufour, A.B. (2007) The ade4 package: Implementing the duality diagram
for ecologists. *Journal of Statistical Software,* **22,** 1-20.

Dray, S. & Legendre, P. (2008) Testing the species traits–environment relationships:
The fourth-corner problem revisited. *Ecology,* **89,** 3400-3412.

Ellis, E.C., Klein Goldewijk, K., Siebert, S., Lightman, D. & Ramankutty, N. (2010)
Anthropogenic transformation of the biomes, 1700 to 2000. *Global Ecology*
*and Biogeography,* **19,** 589-606.

Fisher, D.O. & Owens, I.P. (2004) The comparative method in conservation biology.
*Trends in Ecology & Evolution,* **19,** 391-398.

Fritz, S.A., Bininda-Emonds, O.R. & Purvis, A. (2009) Geographical variation in
predictors of mammalian extinction risk: big is bad, but only in the tropics.
*Ecology Letters,* **12,** 538-549.

Forrest, J.R.K., Thorp R.W., Kremen, C. & Williams, N.M. (2015) Contrasting
patterns in species and functional-trait diversity of bees in an agricultural
landscape. *The Journal of Applied Ecology,* **52**, 706–15.

Gagic, V., Bartomeus, I., Jonsson, T., Taylor, A., Winqvist, C., Fischer, C. *et al.*
        (2015) Functional identity and diversity of animals predict ecosystem
        functioning better than species-based indices. *Proceedings of the Royal
        Society of London, Series B: Biological Sciences,* **282**.

Garibaldi, L.A., Bartomeus, I., Bommarco, R., Klein, A.M., Cunningham, S.A., Aizen,
        M.A. *et al.* (2015) Trait matching of flower visitors and crops predicts fruit set
        better than trait diversity. *Journal of Applied Ecology,* **52,** 1436–1444.

Garibaldi, L.A., Steffan-Dewenter, I., Kremen, C., Morales, J.M., Bommarco, R.,
        Cunningham, S.A. *et al.* (2011) Stability of pollination services decreases with
        isolation from natural areas despite honey bee visits. *Ecology Letters,* **14,**
        1062-1072.

Greenleaf, S.S., Williams, N.M., Winfree, R. & Kremen, C. (2007) Bee foraging
        ranges and their relationship to body size. *Oecologia,* **153,** 589-596.

Hoehn, P., Tscharntke, T., Tylianakis, J.M. & Steffan-Dewenter, I. (2008) Functional
        group diversity of bee pollinators increases crop yield. *Proceedings of the
        Royal Society B-Biological Sciences,* **275,** 2283-2291.

Kattge, J., DÍaz, S., Lavorel, S., Prentice, I.C., Leadley, P., Bönisch, G. *et al.* (2011)
        TRY - a global database of plant traits. *Global Change Biology,* **17,** 2905-
        2935.

Kells, A.R. & Goulson, D. (2003) Preferred nesting sites of bumblebee queens
        (Hymenoptera: Apidae) in agroecosystems in the UK. *Biological Conservation,*
        **109,** 165-174.

Kleijn, D., Baquero, R.A., Clough, Y., Diaz, M., De Esteban, J., Fernandez, F. *et al.*
        (2006) Mixed biodiversity benefits of agri-environment schemes in five
        European countries. *Ecology Letters,* **9,** 243-254; discussion 254-247.

Kleijn, D., Winfree, R., Bartomeus, I., Carvalheiro, L.G., Henry, M., Isaacs, R. *et al.*
        (2015) Delivery of crop pollination services is an insufficient argument for wild
        pollinator conservation. *Nature Communications,* **6,** 7414.

Klein, A.M., Vaissiere, B.E., Cane, J.H., Steffan-Dewenter, I., Cunningham, S.A.,
        Kremen, C. *et al.* (2007) Importance of pollinators in changing landscapes for
        world crops. *Proceedings of the Royal Society of London, Series B: Biological
        Sciences,* **274,** 303-313.

Kremen, C. (2005) Managing ecosystem services: what do we need to know about
        their ecology? *Ecology Letters,* **8,** 468-479.

Kremen, C., Williams, N.M., Aizen, M.A., Gemmill-Herren, B., LeBuhn, G., Minckley,
        R. *et al.* (2007) Pollination and other ecosystem services produced by mobile
        organisms: a conceptual framework for the effects of land-use change.
        *Ecology Letters,* **10,** 299-314.

Larsen, T.H., Williams, N.W. & Kremen, C. (2005) Extinction order and altered
        community structure rapidly disrupt ecosystem functioning. *Ecology Letters,* **8,**
        538-547.

Laughlin, D.C. & Messier, J. (2015) Fitness of multidimensional phenotypes in
        dynamic adaptive landscapes. *Trends in Ecology & Evolution,* **30,** 487-496.

Lavorel, S. & Garnier, E. (2002) Predicting changes in community composition and
        ecosystem functioning from plant traits: revisiting the Holy Grail. *Functional
        Ecology,* **16,** 545-556.

Legendre, P., Galzin, R. & Harmelin-Vivien, M.L. (1997) Relating behavior to habitat:
        Solutions to the fourth-corner problem. *Ecology,* **78,** 547-562.

McGill, B.J., Enquist, B.J., Weiher, E. & Westoby, M. (2006) Rebuilding community
ecology from functional traits. *Trends in Ecology & Evolution,* **21,** 178-185.
Naeem, S. & Wright, J.P. (2003) Disentangling biodiversity effects on ecosystem
functioning: deriving solutions to a seemingly insurmountable problem.
*Ecology Letters,* **6,** 567-579.
Nipperess, D.A. & Matsen, F.A.t. (2013) The mean and variance of phylogenetic
diversity under rarefaction. *Methods in Ecology and Evolution,* **4,** 566-572.
Öckinger, E., Schweiger, O., Crist, T.O., Debinski, D.M., Krauss, J., Kuussaari, M. *et*
*al.* (2010) Life-history traits predict species responses to habitat area and
isolation: a cross-continental synthesis. *Ecology Letters,* **13,** 969-979.
Pereira, H.M., Leadley, P.W., Proenca, V., Alkemade, R., Scharlemann, J.P.,
Fernandez-Manjarres, J.F. *et al.* (2010) Scenarios for global biodiversity in the
21st century. *Science,* **330,** 1496-1501.
Persson, A.S., Rundlöf, M., Clough, Y. & Smith, H.G. (2015) Bumble bees show trait-
dependent vulnerability to landscape simplification. *Biodiversity and*
*Conservation,* **24,** 3469-3489.
Petchey, O.L., Pontarp, M., Massie, T.M., Kefi, S., Ozgul, A., Weilenmann, M. *et al.*
(2015) The ecological forecast horizon, and examples of its uses and
determinants. *Ecology Letters,* **18,** 597-611.
Powney, G.D., Preston, C.D., Purvis, A., Van Landuyt, W. & Roy, D.B. (2014) Can
trait-based analyses of changes in species distribution be transferred to new
geographic areas? *Global Ecology and Biogeography,* **23,** 1009-1018.
R Core Team (2014) R: A language and environment for statistical computing. *R*
*Foundation for Statistical Computing, Vienna, Austria.* R Foundation for
Statistical Computing, Vienna, Austria.
Rader, R., Bartomeus, I., Garibaldi, L.A., Garratt, M.P., Howlett, B.G., Winfree, R. *et*
*al.* (2016) Non-bee insects are important contributors to global crop
pollination. *Proceedings of the National Academy of Sciences USA,* **113,** 146-
151.
Rader, R., Bartomeus, I., Tylianakis, J.M., Laliberté, E. & van Kleunen, M. (2014)
The winners and losers of land use intensification: pollinator community
disassembly is non-random and alters functional diversity. *Diversity and*
*Distributions,* **20,** 908-917.
Scheper, J., Reemer, M., van Kats, R., Ozinga, W.A., van der Linden, G.T.,
Schaminee, J.H. *et al.* (2014) Museum specimens reveal loss of pollen host
plants as key factor driving wild bee decline in The Netherlands. *Proceedings*
*of the National Academy of the Sciences USA.*
Schleuning, M., Fründ, J. & García, D. (2015) Predicting ecosystem functions from
biodiversity and mutualistic networks: an extension of trait-based concepts to
plant-animal interactions. *Ecography,* **38,** 380-392.
Schwartz, M.W., Brigham, C.A., Hoeksema, J.D., Lyons, K.G., Mills, M.H. & van
Mantgem, P.J. (2000) Linking biodiversity to ecosystem function: implications
for conservation ecology. *Oecologia,* **122,** 297-305.
Seguin, A., Harvey, E., Archambault, P., Nozais, C. & Gravel, D. (2014) Body size as
a predictor of species loss effect on ecosystem functioning. *Scientific Reports,*
**4,** 4616.
Shipley, B., De Bello, F., Cornelissen, J.H., Laliberte, E., Laughlin, D.C. & Reich,
P.B. (2016) Reinforcing loose foundation stones in trait-based plant ecology.

*Oecologia,* **180,** 923-931.

Webb, C.O. (2000) Exploring the phylogenetic structure of ecological communities:
An example for rain forest trees. *American Naturalist,* **156,** 145-155.

Webb, C.O., Ackerly, D.D. & Kembel, S.W. (2008) Phylocom: software for the
analysis of phylogenetic community structure and trait evolution.
*Bioinformatics,* **24,** 2098-2100.

Wikström, N., Savolainen, V. & Chase, M.W. (2001) Evolution of the angiosperms:
calibrating the family tree. *Proc Biol Sci,* **268,** 2211-2220.

Williams, N.M., Crone, E.E., Roulston, T.a.H., Minckley, R.L., Packer, L. & Potts,
S.G. (2010) Ecological and life-history traits predict bee species responses to
environmental disturbances. *Biological Conservation,* **143,** 2280-2291.

Winfree, R., Aguilar, R., Vazquez, D.P., LeBuhn, G. & Aizen, M.A. (2009) A meta-
analysis of bees' responses to anthropogenic disturbance. *Ecology,* **90,** 2068-
2076.

Winfree, R., J, W.F., Williams, N.M., Reilly, J.R. & Cariveau, D.P. (2015)
Abundance of common species, not species richness, drives delivery of
a real-world ecosystem service. *Ecology Letters,* **18,** 626-635.

Winfree, R., Williams, N.M., Dushoff, J. & Kremen, C. (2014) Species
abundance, not diet breadth, drives the persistence of the most linked
pollinators as plant-pollinator networks disassemble. *American*
*Naturalist,* **183,** 600-611.

**Tables and Figures**

 **Fig. 1: Relationships between traits and environmental variables** for A)

 watermelon, B) blueberry and c) cranberry. Positive significant correlations

 are in red (p < 0.05). Note no correlation is significant despite all the

 combinations tested for both blueberry and cranberry. D) Detail of the

 relationship between body size (visualized as the community weighted means

 for each community) and percentage of agriculture at 300 meter radii.

640 **Fig. 2: Multipanel plot showing the relationships between species traits**

641 **and pollination function**, which is decomposed into efficiency (pollen

642 deposited per flower visit) and frequency of flower visits. A) watermelon, B)

643 cranberry, C-D) blueberry.

   **Supporting Information**

   Table S1: Equivalencies between species and groups used for single visit

   data.

| Crop | Species | Single Visit Group | Percentage within group |
|------|---------|--------------------|------------------------|
| Blueberry | Andrena_banksi | MED_AND | 0.00 |
| Blueberry | Andrena_barbara | MED_AND | 0.00 |
| Blueberry | Andrena_bradleyi | MED_AND | 0.93 |
| Blueberry | Andrena_carlini | LG_AND | 0.12 |
| Blueberry | Andrena_carolina | LG_AND | 0.01 |
| Blueberry | Andrena_cressonii | LG_AND | 0.01 |
| Blueberry | Andrena_fenningeri | MED_AND | 0.01 |
| Blueberry | Andrena_hilaris | MED_AND | 0.00 |
| Blueberry | Andrena_ilicis | MED_AND | 0.00 |
| Blueberry | Andrena_imitatrix | MED_AND | 0.01 |
| Blueberry | Andrena_mandibularis | MED_AND | 0.00 |
| Blueberry | Andrena_morrisonella | MED_AND | 0.01 |
| Blueberry | Andrena_screpteropsis | MED_AND | 0.00 |
| Blueberry | Andrena_vicina | LG_AND | 0.86 |
| Blueberry | Augochlora_pura | Green | 0.56 |
| Blueberry | Augochlorella_aurata | Green | 0.44 |
| Blueberry | Bombus_bimaculatus | Bom_Q | 0.24 |
| Blueberry | Bombus_griseocollis | Bom_Q | 0.49 |
| Blueberry | Bombus_impatiens | Bom_Q | 0.18 |
| Blueberry | Bombus_perplexus | Bom_Q | 0.08 |
| Blueberry | Ceratina_calcarata/dupla | Dialictus | 0.22 |
| Blueberry | Colletes_inaequalis | Coll | 0.26 |
| Blueberry | Colletes_thoracicus | Coll | 0.09 |
| Blueberry | Colletes_validus | Coll | 0.65 |
| Blueberry | Habropoda_laboriosa | HAB | 1.00 |
| Blueberry | Lasioglossum_acuminatum | Dialictus | 0.02 |
| Blueberry | Lasioglossum_coeruleum | Dialictus | 0.02 |
| Blueberry | Lasioglossum_fuscipenne | Dialictus | 0.02 |
| Blueberry | Lasioglossum_leucocomum | Dialictus | 0.02 |
| Blueberry | Lasioglossum_oblongum | Dialictus | 0.10 |
| Blueberry | Lasioglossum_pilosum | Dialictus | 0.15 |
| Blueberry | Lasioglossum_versatum | Dialictus | 0.02 |
| Blueberry | Lasioglossum_weemsi | Dialictus | 0.07 |
| Blueberry | Lasioglossum_zephyrum | Dialictus | 0.05 |

| | | | |
|---|---|---|---|
| Blueberry | Nomada_luteola | Dialictus | 0.02 |
| Blueberry | Osmia_taurus | Dialictus | 0.02 |
| Blueberry | Osmia_cornifrons | Dialictus | 0.02 |
| Blueberry | Osmia_pumila | Dialictus | 0.07 |
| Blueberry | Sphecodes_aroniae | Dialictus | 0.02 |
| Blueberry | Sphecodes_stygius | Dialictus | 0.02 |
| Blueberry | Xylocopa_virginica | XYL | 1.00 |
| Cranberry | Agapostemon_splendens | Green | 0.02 |
| Cranberry | Andrena_cressonii | Melitta | 0.01 |
| Cranberry | Andrena_imitatrix | Melitta | 0.01 |
| Cranberry | Andrena_morrisonella | Melitta | 0.01 |
| Cranberry | Andrena_spiraeana | Melitta | 0.01 |
| Cranberry | Andrena_vicina | Melitta | 0.02 |
| Cranberry | Augochlora_pura | Green | 0.16 |
| Cranberry | Augochlorella_aurata | Green | 0.63 |
| Cranberry | Augochloropsis_metallica | Green | 0.16 |
| Cranberry | Augochloropsis_sumptuosa | Green | 0.03 |
| Cranberry | Bombus_bimaculatus | Bombus_bimaculatus | 1.00 |
| Cranberry | Bombus_citrinus | Bombus_spp | 0.50 |
| Cranberry | Bombus_griseocollis | Bombus_griseocollis | 1.00 |
| Cranberry | Bombus_impatiens | Bombus_impatiens | 1.00 |
| Cranberry | Bombus_perplexus | Bom_pervag | 0.85 |
| Cranberry | Bombus_sandersoni | Bombus_spp | 0.50 |
| Cranberry | Bombus_vagans | Bom_pervag | 0.15 |
| Cranberry | Ceratina_calcarata/dupla | Small_black | 0.12 |
| Cranberry | Coelioxys_immaculata | Megachile | 0.02 |
| Cranberry | Coelioxys_porterae | Megachile | 0.04 |
| Cranberry | Coelioxys_sayi | Megachile | 0.02 |
| Cranberry | Colletes_consors | Megachile | 0.02 |
| Cranberry | Halictus_rubicundus | Small_black | 0.02 |
| Cranberry | Heriades_carinatus | Osmia | 0.08 |
| Cranberry | Hoplitis_truncata | Osmia | 0.12 |
| Cranberry | Hylaeus_affinis | Small_black | 0.08 |
| Cranberry | Lasioglossum_apopkense | Small_black | 0.01 |
| Cranberry | Lasioglossum_coeruleum | Small_black | 0.01 |
| Cranberry | Lasioglossum_creberrimum | Small_black | 0.02 |
| Cranberry | Lasioglossum_fuscipenne | Small_black | 0.05 |
| Cranberry | Lasioglossum_georgeickworti | Small_black | 0.05 |
| Cranberry | Lasioglossum_lineatulum | Small_black | 0.01 |
| Cranberry | Lasioglossum_oblongum | Small_black | 0.15 |
| Cranberry | Lasioglossum_pilosum | Small_black | 0.02 |
| Cranberry | Lasioglossum_planatum | Small_black | 0.02 |
| Cranberry | Lasioglossum_subviridatum | Small_black | 0.17 |
| Cranberry | Lasioglossum_trigeminum | Small_black | 0.02 |

| Cranberry | Lasioglossum_versatum | Small_black | 0.04 |
|---|---|---|---|
| Cranberry | Megachile_addenda | Megachile | 0.22 |
| Cranberry | Megachile_gemula | Megachile | 0.32 |
| Cranberry | Megachile_mendica | Megachile | 0.28 |
| Cranberry | Megachile_texana | Megachile | 0.08 |
| Cranberry | Melitta_americana | Melitta | 0.95 |
| Cranberry | Nomada_bella/lepida | Small_black | 0.01 |
| Cranberry | Nomada_pygmaea | Small_black | 0.01 |
| Cranberry | Nomada_rodecki | Small_black | 0.07 |
| Cranberry | Osmia_inspergens | Osmia | 0.07 |
| Cranberry | Osmia_pumila | Osmia | 0.13 |
| Cranberry | Osmia_virga | Osmia | 0.60 |
| Cranberry | Panurginus_atramontensis | Small_black | 0.09 |
| Cranberry | Sphecodes_aroniae | Small_black | 0.03 |
| Cranberry | Sphecodes_fattigi | Small_black | 0.01 |
| Cranberry | Xylocopa_virginica | XYL | 1.00 |
| Watermelon | Agapostemon_sericeus | LG | 0.25 |
| Watermelon | Agapostemon_texanus | LG | 0.11 |
| Watermelon | Agapostemon_virescens | LG | 0.52 |
| Watermelon | Anthidium_oblongatum | LDS | 0.03 |
| Watermelon | Augochlora_pura | SG | 0.80 |
| Watermelon | Augochlorella_aurata | SG | 0.20 |
| Watermelon | Augochloropsis_metallica | LG | 0.11 |
| Watermelon | Bombus_bimaculatus | BOM | 0.01 |
| Watermelon | Bombus_fervidus | BOM | 0.00 |
| Watermelon | Bombus_griseocollis | BOM | 0.01 |
| Watermelon | Bombus_impatiens | BOM | 0.98 |
| Watermelon | Bombus_perplexus | BOM | 0.00 |
| Watermelon | Bombus_vagans | BOM | 0.00 |
| Watermelon | Calliopsis_andreniformis | SD | 0.03 |
| Watermelon | Ceratina_calcarata/dupla | CER | 0.81 |
| Watermelon | Ceratina_strenua | CER | 0.19 |
| Watermelon | Halictus_confusus | HAL_MDS | 0.83 |
| Watermelon | Halictus_ligatus | HAL_MDS | 0.16 |
| Watermelon | Halictus_parallelus | LDS | 0.03 |
| Watermelon | Halictus_rubicundus | LDS | 0.56 |
| Watermelon | Hoplitis_pilosifrons | HAL_MDS | 0.00 |
| Watermelon | Hoplitis_producta | HAL_MDS | 0.00 |
| Watermelon | Hylaeus_affinis | TD | 0.01 |
| Watermelon | Lasioglossum_admirandum | SD | 0.01 |
| Watermelon | Lasioglossum_albipenne | SD | 0.00 |
| Watermelon | Lasioglossum_atwoodi | SD | 0.00 |
| Watermelon | Lasioglossum_bruneri | SD | 0.01 |
| Watermelon | Lasioglossum_callidum | SD | 0.01 |

| | | | |
|---|---|---|---|
| Watermelon | Lasioglossum_cattellae | TD | 0.00 |
| Watermelon | Lasioglossum_cinctipes | SD | 0.00 |
| Watermelon | Lasioglossum_coreopsis | TD | 0.00 |
| Watermelon | Lasioglossum_coriaceum | SD | 0.00 |
| Watermelon | Lasioglossum_cressonii | SD | 0.01 |
| Watermelon | Lasioglossum_ellisiae | TD | 0.01 |
| Watermelon | Lasioglossum_ephialtum | SD | 0.02 |
| Watermelon | Lasioglossum_georgeickworti | SD | 0.00 |
| Watermelon | Lasioglossum_gotham | SD | 0.00 |
| Watermelon | Lasioglossum_illinoense | TD | 0.05 |
| Watermelon | Lasioglossum_imitatum | TD | 0.59 |
| Watermelon | Lasioglossum_laevissimum | SD | 0.00 |
| Watermelon | Lasioglossum_leucocomum | SD | 0.02 |
| Watermelon | Lasioglossum_leucozonium | LDS | 0.09 |
| Watermelon | Lasioglossum_lineatulum | SD | 0.00 |
| Watermelon | Lasioglossum_mitchelli | TD | 0.17 |
| Watermelon | Lasioglossum_nymphaearum | SD | 0.04 |
| Watermelon | Lasioglossum_oblongum | SD | 0.01 |
| Watermelon | Lasioglossum_obscurum | SD | 0.01 |
| Watermelon | Lasioglossum_paradmirandum | TD | 0.04 |
| Watermelon | Lasioglossum_pectinatum | SD | 0.00 |
| Watermelon | Lasioglossum_pectorale | SD | 0.02 |
| Watermelon | Lasioglossum_pilosum | SD | 0.28 |
| Watermelon | Lasioglossum_planatum | SD | 0.00 |
| Watermelon | Lasioglossum_platyparium | SD | 0.00 |
| Watermelon | Lasioglossum_rozeni | SD | 0.00 |
| Watermelon | Lasioglossum_smilacinae | SD | 0.00 |
| Watermelon | Lasioglossum_subviridatum | SD | 0.00 |
| Watermelon | Lasioglossum_tegulare | TD | 0.07 |
| Watermelon | Lasioglossum_trigeminum | SD | 0.02 |
| Watermelon | Lasioglossum_truncatum | SD | 0.01 |
| Watermelon | Lasioglossum_versatum | SD | 0.44 |
| Watermelon | Lasioglossum_viridatum | SD | 0.01 |
| Watermelon | Lasioglossum_weemsi | TD | 0.05 |
| Watermelon | Lasioglossum_zephyrum | SD | 0.04 |
| Watermelon | Megachile_brevis | LDS | 0.06 |
| Watermelon | Megachile_mendica | LDS | 0.18 |
| Watermelon | Megachile_rotundata | LDS | 0.03 |
| Watermelon | Megachile_sculpturalis | LDS | 0.03 |
| Watermelon | Melissodes_bimaculata | MEL | 0.99 |
| Watermelon | Melissodes_trinodis | MEL | 0.00 |
| Watermelon | Nomada_articulata | TRI | 0.01 |
| Watermelon | Peponapis_pruinosa | PEP | 1.00 |
| Watermelon | Ptilothrix_bombiformis | MEL | 0.01 |

| | | | |
|---|---|---|---|
| Watermelon | Triepeolus_lunatus | TRI | 0.01 |
| Watermelon | Triepeolus_remigatus | TRI | 0.98 |
| Watermelon | Xylocopa_virginica | XYL | 1.00 |

Text S2: Datasets used for calculating dietary specialization: Six datasets

were used to create the phylogenetic distance index. All data were collected in

the region of the crop study. Specimens were collected using a hand net and

the bee species and plant species were recorded. This resulted in a total of

18,733 bee x plant interactions for species that were also in the crop dataset.

The number of species, sites, and years of collection are as follows: 1) Pine

barrens in 2003: 280 bee x plant interactions. Habitat types were extensive

pine-oak forest (14 sites), forest fragments (14 sites), suburban back yards (7

sites), and agricultural field borders (5 sites) in New Jersey (Winfree et al.

2007). Bees were collected in temporally stratified sampling rounds between

April and September. 2) NJPA: 3906 bee x plant interactions. Data collected

on watermelon field margins at a total of 20 sites. Farm types included small-

scale mixed farming, both crops and field margins, both organic and low-

pesticide-input conventional. All bees were collected in three temporally

stratified sampling rounds in July, in each of 3 years. 3) NFWF 3906 bee x

plant interactions. Habitat types were old fields. Bees were collected in May

through Sept at 25 sites for two years. *Lasioglossum* species where not

included for this dataset due to recent changes in its taxonomy. 4) NSF 2006

666 bee x plant interactions. Habitat types were deciduous forest fragments

(13 sites), and suburban / urban yards (3 sites) and sites with extensive

forests with diverse wildflower communities (4 sites). All bees were collected

in sampling rounds between April and early June. 5) CIG 4600 bee x plant

interactions. Site were comprised of old fields as well as pollinator

enhancement sites. Bees using were collected using a hand net from a total of

a total 18 sites in 2011-2013. For each bee specimen, the plant species was

recorded. 6) Cape May 5858 bee x plant interactions. This study included only

one site. The habitat was an old field that had been planted in 20 species of

native perennial plants. Sampling took place over 3 years in sampling rounds

that occurred in May through September.

Winfree, R. Griswold, T. & Kremen, C. (2007). Effect of human disturbance on

bee communities in a forested ecosystem. *Conservation Biology*. 21: 213-223.

Table S3: Fourth corner results showing correlations and p-values for each

trait. Significant values in bold.

| Crop | Test | r | Standard deviation | P-value |
|------|------|---|--------------------|---------|
| watermelon | ag_300 / Nest_.cavity | 0.19 | 1.92 | 0.05 |
| watermelon | open_300 / Nest_.cavity | -0.09 | -0.83 | 0.43 |
| watermelon | forest_edge_300 / Nest_.cavity | -0.05 | -0.31 | 0.71 |
| watermelon | shannonH_300 / Nest_.cavity | -0.09 | -0.88 | 0.4 |
| watermelon | ag_1500 / Nest_.cavity | -0.11 | -1 | 0.33 |
| watermelon | open_1500 / Nest_.cavity | -0.15 | -1.4 | 0.18 |
| watermelon | forest_edge_1500 / Nest_.cavity | -0.02 | -0.2 | 0.86 |
| watermelon | shannonH_1500 / Nest_.cavity | -0.06 | -0.54 | 0.62 |
| watermelon | ag_300 / Nest_.hole | 0 | 0.07 | 0.93 |
| watermelon | open_300 / Nest_.hole | -0.03 | -0.85 | 0.32 |
| watermelon | forest_edge_300 / Nest_.hole | -0.06 | -0.76 | 0.34 |
| watermelon | shannonH_300 / Nest_.hole | -0.03 | -0.63 | 0.41 |
| watermelon | ag_1500 / Nest_.hole | -0.03 | -0.54 | 0.41 |
| watermelon | open_1500 / Nest_.hole | -0.05 | -0.94 | 0.22 |
| watermelon | forest_edge_1500 / Nest_.hole | -0.03 | -0.65 | 0.36 |
| watermelon | shannonH_1500 / Nest_.hole | -0.08 | -1.61 | 0.09 |
| watermelon | ag_300 / Nest_.soil | -0.12 | -1.2 | 0.27 |
| watermelon | open_300 / Nest_.soil | 0.11 | 1.16 | 0.27 |

| watermelon | forest_edge_300 / Nest_.soil | 0.01 | -0.07 | 0.95 |
|---|---|---|---|---|
| watermelon | shannonH_300 / Nest_.soil | 0.07 | 0.81 | 0.45 |
| watermelon | ag_1500 / Nest_.soil | 0.04 | 0.46 | 0.66 |
| watermelon | open_1500 / Nest_.soil | 0.07 | 0.73 | 0.5 |
| watermelon | forest_edge_1500 / Nest_.soil | -0.05 | -0.53 | 0.63 |
| watermelon | shannonH_1500 / Nest_.soil | 0.02 | 0.23 | 0.85 |
| watermelon | ag_300 / Nest_.stem | -0.08 | -1 | 0.26 |
| watermelon | open_300 / Nest_.stem | 0.03 | 0.44 | 0.66 |
| watermelon | forest_edge_300 / Nest_.stem | 0.01 | 0.11 | 0.93 |
| watermelon | shannonH_300 / Nest_.stem | 0.09 | 1.38 | 0.18 |
| watermelon | ag_1500 / Nest_.stem | 0 | 0.05 | 0.96 |
| watermelon | open_1500 / Nest_.stem | 0.07 | 0.87 | 0.27 |
| watermelon | forest_edge_1500 / Nest_.stem | 0.01 | 0.25 | 0.78 |
| **watermelon** | **shannonH_1500 / Nest_.stem** | **0.12** | **2.01** | **0.04** |
| watermelon | ag_300 / Nest_.wood | -0.03 | -0.22 | 0.76 |
| watermelon | open_300 / Nest_.wood | -0.06 | -0.84 | 0.43 |
| watermelon | forest_edge_300 / Nest_.wood | 0.05 | 0.66 | 0.52 |
| watermelon | shannonH_300 / Nest_.wood | -0.05 | -0.66 | 0.54 |
| watermelon | ag_1500 / Nest_.wood | 0.08 | 1.04 | 0.3 |
| watermelon | open_1500 / Nest_.wood | 0.04 | 0.51 | 0.59 |
| watermelon | forest_edge_1500 / Nest_.wood | 0.09 | 1.37 | 0.17 |
| watermelon | shannonH_1500 / Nest_.wood | -0.04 | -0.58 | 0.57 |
| watermelon | ag_300 / Socia.Eusocial | 0.02 | 0.18 | 0.91 |
| watermelon | open_300 / Socia.Eusocial | -0.01 | -0.09 | 0.93 |
| watermelon | forest_edge_300 / Socia.Eusocial | 0.04 | 0.27 | 0.86 |
| watermelon | shannonH_300 / Socia.Eusocial | 0.08 | 0.79 | 0.48 |
| watermelon | ag_1500 / Socia.Eusocial | -0.04 | -0.36 | 0.74 |
| watermelon | open_1500 / Socia.Eusocial | -0.16 | -1.61 | 0.11 |
| watermelon | forest_edge_1500 / Socia.Eusocial | -0.06 | -0.61 | 0.56 |
| watermelon | shannonH_1500 / Socia.Eusocial | 0.01 | 0.12 | 0.92 |
| watermelon | ag_300 / Socia.fac_social | -0.07 | -0.64 | 0.54 |
| watermelon | open_300 / Socia.fac_social | 0.03 | 0.45 | 0.72 |
| watermelon | forest_edge_300 / Socia.fac_social | -0.01 | 0.02 | 0.99 |
| watermelon | shannonH_300 / Socia.fac_social | 0.06 | 0.86 | 0.47 |
| watermelon | ag_1500 / Socia.fac_social | -0.01 | -0.15 | 0.86 |
| watermelon | open_1500 / Socia.fac_social | 0.07 | 0.73 | 0.47 |
| watermelon | forest_edge_1500 / Socia.fac_social | 0 | -0.02 | 0.99 |
| watermelon | shannonH_1500 / Socia.fac_social | 0.09 | 1.52 | 0.13 |
| watermelon | ag_300 / Socia.Solitary | 0.03 | 0.31 | 0.79 |
| watermelon | open_300 / Socia.Solitary | -0.01 | -0.15 | 0.89 |
| watermelon | forest_edge_300 / Socia.Solitary | -0.03 | -0.16 | 0.86 |

| | | | | |
|---|---|---|---|---|
| watermelon | shannonH_300 / Socia.Solitary | -0.14 | -1.46 | 0.14 |
| watermelon | ag_1500 / Socia.Solitary | 0.05 | 0.54 | 0.61 |
| watermelon | open_1500 / Socia.Solitary | 0.13 | 1.33 | 0.2 |
| watermelon | forest_edge_1500 / Socia.Solitary | 0.07 | 0.74 | 0.48 |
| watermelon | shannonH_1500 / Socia.Solitary | -0.08 | -0.83 | 0.44 |
| watermelon | ag_300 / Paras.No | -0.1 | -0.85 | 0.48 |
| watermelon | open_300 / Paras.No | -0.02 | -0.24 | 0.88 |
| watermelon | forest_edge_300 / Paras.No | -0.02 | -0.22 | 0.82 |
| watermelon | shannonH_300 / Paras.No | 0.18 | 1.67 | 0.08 |
| watermelon | ag_1500 / Paras.No | 0 | 0 | 1 |
| watermelon | open_1500 / Paras.No | -0.22 | -2.07 | 0.05 |
| watermelon | forest_edge_1500 / Paras.No | -0.12 | -1.13 | 0.23 |
| watermelon | shannonH_1500 / Paras.No | 0.06 | 0.57 | 0.59 |
| watermelon | ag_300 / Paras.Yes | 0.1 | 0.85 | 0.48 |
| watermelon | open_300 / Paras.Yes | 0.02 | 0.24 | 0.88 |
| watermelon | forest_edge_300 / Paras.Yes | 0.02 | 0.22 | 0.82 |
| watermelon | shannonH_300 / Paras.Yes | -0.18 | -1.67 | 0.08 |
| watermelon | ag_1500 / Paras.Yes | 0 | 0 | 1 |
| watermelon | open_1500 / Paras.Yes | 0.22 | 2.07 | 0.05 |
| watermelon | forest_edge_1500 / Paras.Yes | 0.12 | 1.13 | 0.23 |
| watermelon | shannonH_1500 / Paras.Yes | -0.06 | -0.57 | 0.59 |
| **watermelon** | **ag_300 / ITfam** | **0.23** | **2.19** | **0.02** |
| watermelon | open_300 / ITfam | -0.07 | -0.61 | 0.56 |
| watermelon | forest_edge_300 / ITfam | -0.1 | -0.82 | 0.41 |
| watermelon | shannonH_300 / ITfam | -0.15 | -1.39 | 0.18 |
| watermelon | ag_1500 / ITfam | -0.13 | -1.15 | 0.27 |
| watermelon | open_1500 / ITfam | -0.1 | -0.94 | 0.36 |
| watermelon | forest_edge_1500 / ITfam | -0.03 | -0.29 | 0.79 |
| watermelon | shannonH_1500 / ITfam | -0.09 | -0.78 | 0.45 |
| watermelon | ag_300 / PDrar20 | -0.08 | -0.76 | 0.52 |
| watermelon | open_300 / PDrar20 | -0.04 | -0.53 | 0.63 |
| watermelon | forest_edge_300 / PDrar20 | 0.02 | 0.18 | 0.88 |
| watermelon | shannonH_300 / PDrar20 | 0.08 | 1 | 0.35 |
| watermelon | ag_1500 / PDrar20 | 0.05 | 0.51 | 0.7 |
| watermelon | open_1500 / PDrar20 | -0.13 | -1.56 | 0.12 |
| watermelon | forest_edge_1500 / PDrar20 | -0.06 | -0.7 | 0.52 |
| watermelon | shannonH_1500 / PDrar20 | 0.02 | 0.22 | 0.86 |
| **watermelon** | **ag_300 / tongue** | **0.23** | **2.18** | **0.02** |
| watermelon | open_300 / tongue | -0.07 | -0.64 | 0.55 |
| watermelon | forest_edge_300 / tongue | -0.09 | -0.76 | 0.45 |
| watermelon | shannonH_300 / tongue | -0.14 | -1.38 | 0.19 |
| watermelon | ag_1500 / tongue | -0.12 | -1.11 | 0.29 |
| watermelon | open_1500 / tongue | -0.1 | -0.92 | 0.37 |
| watermelon | forest_edge_1500 / tongue | -0.03 | -0.26 | 0.82 |

| | | | | |
|---|---|---|---|---|
| watermelon | shannonH_1500 / tongue | -0.08 | -0.73 | 0.48 |
| cranberry | ag_300 / Nest_.cavity | 0.1 | 1.09 | 0.3 |
| cranberry | open_300 / Nest_.cavity | -0.15 | -1.62 | 0.11 |
| cranberry | forest_edge_300 / Nest_.cavity | 0 | 0.05 | 0.97 |
| cranberry | shannonH_300 / Nest_.cavity | -0.07 | -0.71 | 0.51 |
| cranberry | ag_1500 / Nest_.cavity | 0.05 | 0.48 | 0.65 |
| cranberry | open_1500 / Nest_.cavity | -0.08 | -0.69 | 0.5 |
| cranberry | forest_edge_1500 / Nest_.cavity | -0.07 | -0.75 | 0.47 |
| cranberry | shannonH_1500 / Nest_.cavity | 0.04 | 0.38 | 0.72 |
| cranberry | ag_300 / Nest_.hole | 0.01 | 0.12 | 0.91 |
| cranberry | open_300 / Nest_.hole | -0.05 | -0.56 | 0.58 |
| cranberry | forest_edge_300 / Nest_.hole | 0.01 | 0.18 | 0.88 |
| cranberry | shannonH_300 / Nest_.hole | -0.03 | -0.56 | 0.58 |
| cranberry | ag_1500 / Nest_.hole | -0.04 | -0.43 | 0.68 |
| cranberry | open_1500 / Nest_.hole | -0.08 | -0.64 | 0.56 |
| cranberry | forest_edge_1500 / Nest_.hole | -0.02 | -0.22 | 0.87 |
| cranberry | shannonH_1500 / Nest_.hole | -0.05 | -0.51 | 0.61 |
| cranberry | ag_300 / Nest_.soil | -0.09 | -0.89 | 0.41 |
| cranberry | open_300 / Nest_.soil | 0.09 | 0.97 | 0.36 |
| cranberry | forest_edge_300 / Nest_.soil | 0 | 0.01 | 1 |
| cranberry | shannonH_300 / Nest_.soil | 0.09 | 0.81 | 0.45 |
| cranberry | ag_1500 / Nest_.soil | -0.04 | -0.33 | 0.76 |
| cranberry | open_1500 / Nest_.soil | 0.13 | 1.08 | 0.3 |
| cranberry | forest_edge_1500 / Nest_.soil | 0.09 | 0.87 | 0.41 |
| cranberry | shannonH_1500 / Nest_.soil | -0.02 | -0.17 | 0.91 |
| cranberry | ag_300 / Nest_.stem | -0.03 | -0.52 | 0.36 |
| cranberry | open_300 / Nest_.stem | 0.02 | 0.21 | 0.84 |
| cranberry | forest_edge_300 / Nest_.stem | 0 | 0.07 | 0.94 |
| cranberry | shannonH_300 / Nest_.stem | 0.02 | 0.32 | 0.56 |
| cranberry | ag_1500 / Nest_.stem | -0.01 | -0.11 | 0.9 |
| cranberry | open_1500 / Nest_.stem | 0.03 | 0.51 | 0.44 |
| cranberry | forest_edge_1500 / Nest_.stem | 0.02 | 0.36 | 0.61 |
| cranberry | shannonH_1500 / Nest_.stem | 0 | 0.05 | 0.92 |
| cranberry | ag_300 / Nest_.wood | -0.05 | -0.55 | 0.6 |
| cranberry | open_300 / Nest_.wood | 0.17 | 1.77 | 0.07 |
| cranberry | forest_edge_300 / Nest_.wood | -0.02 | -0.3 | 0.76 |
| cranberry | shannonH_300 / Nest_.wood | 0.01 | 0.1 | 0.93 |
| cranberry | ag_1500 / Nest_.wood | 0 | 0.05 | 0.96 |
| cranberry | open_1500 / Nest_.wood | 0 | -0.04 | 0.97 |
| cranberry | forest_edge_1500 / Nest_.wood | 0 | 0 | 1 |
| cranberry | shannonH_1500 / Nest_.wood | -0.01 | -0.07 | 0.94 |
| cranberry | ag_300 / Socia.Eusocial | 0.02 | 0.28 | 0.82 |
| cranberry | open_300 / Socia.Eusocial | 0.03 | 0.19 | 0.88 |
| cranberry | forest_edge_300 / Socia.Eusocial | 0.01 | 0.09 | 0.93 |

| | | | | |
|---|---|---|---|---|
| cranberry | shannonH_300 / Socia.Eusocial | 0.02 | 0.28 | 0.8 |
| cranberry | ag_1500 / Socia.Eusocial | -0.01 | -0.08 | 0.95 |
| cranberry | open_1500 / Socia.Eusocial | 0.02 | 0.18 | 0.88 |
| cranberry | forest_edge_1500 / Socia.Eusocial | -0.01 | -0.14 | 0.92 |
| cranberry | shannonH_1500 / Socia.Eusocial | 0.01 | 0.12 | 0.91 |
| cranberry | ag_300 / Socia.fac_social | 0.03 | 0.75 | 0.47 |
| cranberry | open_300 / Socia.fac_social | 0.04 | 0.29 | 0.77 |
| cranberry | forest_edge_300 / Socia.fac_social | -0.05 | -1.15 | 0.29 |
| cranberry | shannonH_300 / Socia.fac_social | -0.04 | -0.71 | 0.32 |
| cranberry | ag_1500 / Socia.fac_social | 0.04 | 0.71 | 0.47 |
| cranberry | open_1500 / Socia.fac_social | 0.01 | 0.23 | 0.77 |
| cranberry | forest_edge_1500 / Socia.fac_social | 0.01 | 0.22 | 0.81 |
| cranberry | shannonH_1500 / Socia.fac_social | 0.03 | 0.56 | 0.5 |
| cranberry | ag_300 / Socia.Solitary | -0.03 | -0.41 | 0.74 |
| cranberry | open_300 / Socia.Solitary | -0.04 | -0.37 | 0.72 |
| cranberry | forest_edge_300 / Socia.Solitary | 0.01 | 0.15 | 0.9 |
| cranberry | shannonH_300 / Socia.Solitary | -0.01 | -0.09 | 0.94 |
| cranberry | ag_1500 / Socia.Solitary | 0 | -0.03 | 0.97 |
| cranberry | open_1500 / Socia.Solitary | -0.03 | -0.23 | 0.86 |
| cranberry | forest_edge_1500 / Socia.Solitary | 0.01 | 0.07 | 0.96 |
| cranberry | shannonH_1500 / Socia.Solitary | -0.02 | -0.32 | 0.79 |
| cranberry | ag_300 / Paras.No | 0 | -0.01 | 1 |
| cranberry | open_300 / Paras.No | 0.01 | 0.16 | 0.87 |
| cranberry | forest_edge_300 / Paras.No | 0 | -0.05 | 0.96 |
| cranberry | shannonH_300 / Paras.No | 0 | 0.01 | 0.97 |
| cranberry | ag_1500 / Paras.No | 0 | 0.04 | 0.97 |
| cranberry | open_1500 / Paras.No | 0.01 | -0.05 | 0.96 |
| cranberry | forest_edge_1500 / Paras.No | 0.02 | 0.14 | 0.88 |
| cranberry | shannonH_1500 / Paras.No | 0.03 | 0.24 | 0.74 |
| cranberry | ag_300 / Paras.Yes | 0 | 0.01 | 1 |
| cranberry | open_300 / Paras.Yes | -0.01 | -0.16 | 0.87 |
| cranberry | forest_edge_300 / Paras.Yes | 0 | 0.05 | 0.96 |
| cranberry | shannonH_300 / Paras.Yes | 0 | -0.01 | 0.97 |
| cranberry | ag_1500 / Paras.Yes | 0 | -0.04 | 0.97 |
| cranberry | open_1500 / Paras.Yes | -0.01 | 0.05 | 0.96 |
| cranberry | forest_edge_1500 / Paras.Yes | -0.02 | -0.14 | 0.88 |
| cranberry | shannonH_1500 / Paras.Yes | -0.03 | -0.24 | 0.74 |
| cranberry | ag_300 / ITfam | 0.13 | 1.37 | 0.19 |
| cranberry | open_300 / ITfam | -0.13 | -1.36 | 0.18 |
| cranberry | forest_edge_300 / ITfam | -0.05 | -0.54 | 0.62 |

| | | | | |
|---|---|---|---|---|
| cranberry | shannonH_300 / ITfam | -0.12 | -1.31 | 0.2 |
| cranberry | ag_1500 / ITfam | 0.14 | 1.45 | 0.15 |
| cranberry | open_1500 / ITfam | 0.01 | 0.02 | 0.99 |
| cranberry | forest_edge_1500 / ITfam | 0 | -0.08 | 0.93 |
| cranberry | shannonH_1500 / ITfam | 0.11 | 1.18 | 0.27 |
| cranberry | ag_300 / PDrar20 | -0.05 | -0.47 | 0.68 |
| cranberry | open_300 / PDrar20 | 0.18 | 1.87 | 0.08 |
| cranberry | forest_edge_300 / PDrar20 | 0.02 | 0.16 | 0.88 |
| cranberry | shannonH_300 / PDrar20 | 0.09 | 0.87 | 0.42 |
| cranberry | ag_1500 / PDrar20 | -0.06 | -0.66 | 0.55 |
| cranberry | open_1500 / PDrar20 | 0.06 | 0.46 | 0.71 |
| cranberry | forest_edge_1500 / PDrar20 | 0.03 | 0.27 | 0.82 |
| cranberry | shannonH_1500 / PDrar20 | -0.01 | -0.08 | 0.95 |
| cranberry | ag_300 / tongue | 0.11 | 1.19 | 0.25 |
| cranberry | open_300 / tongue | -0.15 | -1.54 | 0.13 |
| cranberry | forest_edge_300 / tongue | -0.01 | -0.13 | 0.91 |
| cranberry | shannonH_300 / tongue | -0.09 | -0.9 | 0.39 |
| cranberry | ag_1500 / tongue | 0.07 | 0.72 | 0.5 |
| cranberry | open_1500 / tongue | -0.06 | -0.5 | 0.66 |
| cranberry | forest_edge_1500 / tongue | -0.06 | -0.64 | 0.58 |
| cranberry | shannonH_1500 / tongue | 0.06 | 0.6 | 0.6 |
| blueberry | ag_300 / Nest_.cavity | -0.14 | -0.76 | 0.52 |
| blueberry | open_300 / Nest_.cavity | 0.11 | 0.68 | 0.52 |
| blueberry | forest_edge_300 / Nest_.cavity | 0.1 | 0.61 | 0.64 |
| blueberry | shannonH_300 / Nest_.cavity | 0.09 | 0.53 | 0.69 |
| blueberry | ag_1500 / Nest_.cavity | -0.34 | -1.13 | 0.21 |
| blueberry | open_1500 / Nest_.cavity | 0.2 | 1.33 | 0.19 |
| blueberry | forest_edge_1500 / Nest_.cavity | 0.24 | 0.94 | 0.31 |
| blueberry | shannonH_1500 / Nest_.cavity | -0.04 | -0.25 | 0.82 |
| blueberry | ag_300 / Nest_.hole | -0.03 | -0.02 | 0.93 |
| blueberry | open_300 / Nest_.hole | -0.02 | -0.35 | 0.82 |
| blueberry | forest_edge_300 / Nest_.hole | 0.05 | 0.23 | 0.81 |
| blueberry | shannonH_300 / Nest_.hole | 0.07 | 0.4 | 0.72 |
| blueberry | ag_1500 / Nest_.hole | 0.08 | 0.52 | 0.44 |
| blueberry | open_1500 / Nest_.hole | -0.05 | -0.55 | 0.6 |
| blueberry | forest_edge_1500 / Nest_.hole | 0.01 | 0 | 1 |
| blueberry | shannonH_1500 / Nest_.hole | 0.05 | 0.29 | 0.71 |
| blueberry | ag_300 / Nest_.soil | 0.12 | 0.75 | 0.5 |
| blueberry | open_300 / Nest_.soil | -0.12 | -0.78 | 0.51 |
| blueberry | forest_edge_300 / Nest_.soil | -0.05 | -0.3 | 0.77 |
| blueberry | shannonH_300 / Nest_.soil | -0.08 | -0.47 | 0.67 |
| blueberry | ag_1500 / Nest_.soil | 0.41 | 1.28 | 0.14 |
| blueberry | open_1500 / Nest_.soil | -0.09 | -0.55 | 0.65 |
| blueberry | forest_edge_1500 / Nest_.soil | -0.4 | -1.48 | 0.08 |

| | | | | |
|---|---|---|---|---|
| blueberry | shannonH_1500 / Nest_.soil | -0.08 | -0.48 | 0.72 |
| blueberry | ag_300 / Nest_.wood | 0.01 | 0.07 | 0.95 |
| blueberry | open_300 / Nest_.wood | 0.03 | 0.09 | 0.92 |
| blueberry | forest_edge_300 / Nest_.wood | -0.07 | -0.54 | 0.74 |
| blueberry | shannonH_300 / Nest_.wood | -0.02 | -0.19 | 0.92 |
| blueberry | ag_1500 / Nest_.wood | -0.15 | -0.48 | 0.52 |
| blueberry | open_1500 / Nest_.wood | -0.12 | -0.93 | 0.4 |
| blueberry | forest_edge_1500 / Nest_.wood | 0.25 | 1.07 | 0.18 |
| blueberry | shannonH_1500 / Nest_.wood | 0.15 | 0.85 | 0.33 |
| blueberry | ag_300 / Socia.Eusocial | -0.17 | -1.03 | 0.36 |
| blueberry | open_300 / Socia.Eusocial | 0.13 | 0.83 | 0.47 |
| blueberry | forest_edge_300 / Socia.Eusocial | 0.16 | 1.1 | 0.29 |
| blueberry | shannonH_300 / Socia.Eusocial | 0.13 | 0.84 | 0.42 |
| blueberry | ag_1500 / Socia.Eusocial | -0.32 | -1.04 | 0.38 |
| blueberry | open_1500 / Socia.Eusocial | 0.21 | 1.38 | 0.17 |
| blueberry | forest_edge_1500 / Socia.Eusocial | 0.2 | 0.77 | 0.67 |
| blueberry | shannonH_1500 / Socia.Eusocial | -0.01 | -0.02 | 0.98 |
| blueberry | ag_300 / Socia.fac_social | 0.08 | 0.7 | 0.5 |
| blueberry | open_300 / Socia.fac_social | -0.06 | -0.48 | 0.66 |
| blueberry | forest_edge_300 / Socia.fac_social | -0.12 | -1.09 | 0.29 |
| blueberry | shannonH_300 / Socia.fac_social | -0.09 | -0.76 | 0.46 |
| blueberry | ag_1500 / Socia.fac_social | -0.12 | -0.59 | 0.31 |
| blueberry | open_1500 / Socia.fac_social | -0.18 | -1.52 | 0.12 |
| blueberry | forest_edge_1500 / Socia.fac_social | 0.22 | 1.36 | 0.09 |
| blueberry | shannonH_1500 / Socia.fac_social | 0.13 | 1.19 | 0.26 |
| blueberry | ag_300 / Socia.Solitary | 0.11 | 0.67 | 0.55 |
| blueberry | open_300 / Socia.Solitary | -0.08 | -0.55 | 0.63 |
| blueberry | forest_edge_300 / Socia.Solitary | -0.07 | -0.44 | 0.67 |
| blueberry | shannonH_300 / Socia.Solitary | -0.06 | -0.38 | 0.74 |
| blueberry | ag_1500 / Socia.Solitary | 0.39 | 1.27 | 0.13 |
| blueberry | open_1500 / Socia.Solitary | -0.08 | -0.52 | 0.66 |
| blueberry | forest_edge_1500 / Socia.Solitary | -0.34 | -1.3 | 0.17 |
| blueberry | shannonH_1500 / Socia.Solitary | -0.08 | -0.51 | 0.73 |
| blueberry | ag_300 / Paras.No | 0 | -0.71 | 1 |
| blueberry | open_300 / Paras.No | 0 | 0.65 | 1 |
| blueberry | forest_edge_300 / Paras.No | 0 | -0.69 | 0.91 |
| blueberry | shannonH_300 / Paras.No | 0 | 0.43 | 0.88 |
| blueberry | ag_1500 / Paras.No | 0 | 0.69 | 0.99 |
| blueberry | open_1500 / Paras.No | 0 | -0.7 | 0.97 |
| blueberry | forest_edge_1500 / Paras.No | 0 | 0.96 | 0.34 |

| | | | | |
|---|---|---|---|---|
| blueberry | shannonH_1500 / Paras.No | 0 | -0.72 | 0.88 |
| blueberry | ag_300 / ITfam | -0.03 | -0.13 | 0.9 |
| blueberry | open_300 / ITfam | 0.01 | 0.02 | 0.99 |
| blueberry | forest_edge_300 / ITfam | -0.05 | -0.33 | 0.78 |
| blueberry | shannonH_300 / ITfam | -0.02 | -0.16 | 0.88 |
| blueberry | ag_1500 / ITfam | -0.42 | -1.33 | 0.16 |
| blueberry | open_1500 / ITfam | -0.02 | -0.09 | 0.95 |
| blueberry | forest_edge_1500 / ITfam | 0.42 | 1.59 | 0.08 |
| blueberry | shannonH_1500 / ITfam | 0.12 | 0.73 | 0.52 |
| blueberry | ag_300 / PDrar20 | -0.08 | -0.4 | 0.73 |
| blueberry | open_300 / PDrar20 | 0.13 | 0.66 | 0.53 |
| blueberry | forest_edge_300 / PDrar20 | 0 | -0.02 | 0.98 |
| blueberry | shannonH_300 / PDrar20 | 0.03 | 0.15 | 0.89 |
| blueberry | ag_1500 / PDrar20 | -0.42 | -1.39 | 0.17 |
| blueberry | open_1500 / PDrar20 | -0.04 | -0.19 | 0.87 |
| blueberry | forest_edge_1500 / PDrar20 | 0.35 | 1.37 | 0.17 |
| blueberry | shannonH_1500 / PDrar20 | 0.32 | 1.59 | 0.11 |
| blueberry | ag_300 / tongue | -0.17 | -0.91 | 0.41 |
| blueberry | open_300 / tongue | 0.05 | 0.29 | 0.82 |
| blueberry | forest_edge_300 / tongue | 0.08 | 0.41 | 0.72 |
| blueberry | shannonH_300 / tongue | 0.08 | 0.41 | 0.71 |
| blueberry | ag_1500 / tongue | -0.48 | -1.51 | 0.12 |
| blueberry | open_1500 / tongue | 0.07 | 0.35 | 0.73 |
| blueberry | forest_edge_1500 / tongue | 0.36 | 1.39 | 0.18 |
| blueberry | shannonH_1500 / tongue | -0.02 | -0.08 | 0.95 |

684 Table S4: Model selection procedure showing all models within 2 AICc values.

| Crop | Measure | Model | Delta AICc |
|---|---|---|---|
| Watermelon | Visitation Frequency | ~ body size + tongue | 0 |
| Watermelon | Visitation Frequency | ~ tongue | 0.06 |
| Watermelon | Visitation Frequency | ~ . | 0.28 |
| Watermelon | Visitation Frequency | ~ body size | 1.99 |
| Watermelon | Pollen deposition | ~ body size + tongue | 0 |
| Cranberry | Visitation Frequency | ~ sociality + tongue | 0 |
| Cranberry | Visitation Frequency | ~ nest place | 1.43 |
| Cranberry | Visitation Frequency | ~ sociality | 1.56 |
| Cranberry | Pollen deposition | ~ . | 0 |
| Blueberry | Visitation Frequency | ~ specialization | 0 |
| Blueberry | Pollen deposition | ~ tongue | 0 |