Title: A novel ENU-induced ankyrin-1 mutation impairs parasite invasion and increases erythrocyte
clearance during malaria infection in mice

Author list and affiliations

Hong Ming Huang[1], Denis C. Bauer[2], Patrick M. Lelliott[3], Andreas Greth[4], Brendan J. McMorran[1],
Simon J. Foote[1], Gaetan Burgio[1]*

[1] Department of Immunology and Infectious Disease, John Curtin School of Medical Research,
Australian National University, ACT, Australia.

[2] CSIRO, Sydney, NSW, Australia.

[3] IFReC Research Building, Osaka University, 3-1 Yamada-oka, Suita, Osaka 565-0871, Japan.

[4] synaps studios GmbH, Rebmoosweg 73A, CH-5200 Brugg, Switzerland.

Corresponding author*

Dr. Gaetan Burgio

The John Curtin School of Medical Research, Australian National University,

131 Garran Road, ACT 2601, Australia.

Gaetan.burgio@anu.edu.au

+61 2 612 59428

Abstract

Genetic defects in various red blood cell (RBC) cytoskeletal proteins have been long associated with changes in susceptibility towards malaria infection. In particular, while ankyrin (Ank-1) mutations account for approximately 50% of hereditary spherocytosis (HS) cases, an association with malaria is not well-established, and conflicting evidence has been reported. We describe a novel N-ethyl-N-nitrosourea (ENU)-induced ankyrin mutation MRI61689 that gives rise to two different ankyrin transcripts: one with an introduced splice acceptor site resulting a frameshift, the other with a skipped exon. *Ank-1*$^{(MRI61689/+)}$ mice exhibit an HS-like phenotype including reduction in mean corpuscular volume (MCV), increased osmotic fragility and reduced RBC deformability. They were also found to be resistant to rodent malaria *Plasmodium chabaudi* infection. Parasites in *Ank-1*$^{(MRI61689/+)}$ erythrocytes grew normally, but red cells showed resistance to merozoite invasion. Uninfected *Ank-1*$^{(MRI61689/+)}$ erythrocytes were also more likely to be cleared from circulation during infection; the "bystander effect". This increased clearance is a novel resistance mechanism which was not observed in previous ankyrin mouse models. We propose that this bystander effect is due to reduced deformability of *Ank-1*$^{(MRI61689/+)}$ erythrocytes. This paper highlights the complex roles ankyrin plays in mediating malaria resistance.

Introduction

Malaria is a mosquito-borne disease caused by the protozoan *Plasmodium*, responsible for many deaths every year, mostly children [1]. In endemic regions with limited healthcare access, host genetics is one of the major determinants of malaria susceptibility and survival [2-4]. This is evident from the distributions of various genetic polymorphisms in humans, such as Duffy antigen negativity and sickle cell trait, which coincide with malaria distribution [3,5,6]. It is thought that these genetic polymorphisms confer protection against malaria, thus providing a survival advantage in the face of malaria-induced mortality [7,8].

In addition, these polymorphisms also provide crucial insights into host-parasite interactions. *Plasmodium* relies on a favourable host environment in order to thrive, many erythrocyte-related polymorphisms have been discovered that interfere with parasite survival thus contributing to malaria resistance. These include polymorphisms that affect the cytoskeleton of erythrocytes, such as Southeast Asian Ovalocytosis (SAO), hereditary elliptocytosis (HE) and spherocytosis (HS) [9-13]. Several hypotheses have been proposed for the mechanisms by which they confer malaria protection, including reduced erythrocyte invasion, intra-erythrocytic growth and cytoadherence [8,13-19]. However, due to the heterogeneity of the manifestation of these disorders in the human population, contradicting evidences for resistance mechanisms has often been presented. A study done by Facer [13] showed that only patients carrying certain spectrin mutations have impaired parasite invasion of red blood cells (RBC), but not others that also exhibited HE symptoms. A similar observation was reported by Chishti, et al. [17], where individuals with defective protein 4.1 exhibited intra-erythrocytic growth inhibition, but not those with glycophorin C defects, despite the fact that both defects gave rise to HE. These differences in malaria resistance mechanisms remained largely unexplored, and further studies in this aspect would potentially provide useful insight into host-parasite interactions.

The RBC cytoskeletal protein ankyrin-1 (ANK-1), is a 210kDa protein responsible for connecting the

spectrin network with the RBC membrane through interactions with Band 3, protein 4.2 and the

Rhesus complex [20-22]. Spherocytosis is a genetic disorder where RBCs are abnormally small and are

known as spherocytes. ANK-1 mutations account for more than 50% of human HS cases [23]. However,

similar to SAO and HE, HS is a heterogeneous disorder where the symptoms vary greatly depending

on the mutations. The disorder can range from asymptomatic through to severe anaemia requiring

splenectomy [24]. Despite a possible association with malaria, HS is actually common in Northern

European and Japanese populations [25-28], and much rarer in other populations [29]. Nevertheless,

several *in vitro* and *in vivo* studies have repeatedly reported association between HS and malaria

resistance, and several mechanisms have been suggested. An *in vitro* study done by Schulman, et al.

[16] using RBCs from HS patients suggested that parasite invasion and growth in these erythrocytes

was impaired. This is further supported by studies done in mice with ankyrin mutations. Both *Ank-*

1$^{(nb/nb)}$ and *Ank-1 $^{(MRI23420/+)}$* have shown inhibited intra-erythrocytic growth and erythrocyte invasion

possibly due to spectrin and ankyrin deficiency, respectively [30,31]. On the other hand, another

mutation described by Rank, et al. [32], *Ank-1$^{1674/+}$*, parasite invasion appeared to be normal in these

erythrocytes. Instead, increased erythrocyte fragility was proposed as a contributing factor for

increased malaria resistance [32]. Taking these observations together, it is possible that disruption to

erythrocyte cytoskeletons can mediate multiple mechanisms of resistance.

In a large phenotypic N-ethyl-N-nitrosourea (ENU) mutagenesis screen, using either abnormal red

cells or resistance to malaria as to the screened phenotypes, we identified many novel mutations

that give rise to RBC abnormalities and consequently malaria resistance in mice. We report here an

ENU-induced mutation in the ankyrin-1 gene (*Ank-1$^{(MRI61689)}$*) which was found to exhibit a HS-like

phenotype, with significantly lower RBC volumes, increased osmotic fragility and decreased

deformability. *Ank-1$^{(MRI61689)}$* also confers resistance towards *Plasmodium chabaudi adami* infection

in mice, and *Ank-1$^{(MRI61689/+)}$* mice were shown to show both reduced merozoite invasion and

increased RBC clearance, possibly as a consequence of reduced red blood cell deformability.

Results

The MRI61689 mutation gives rise to an hereditary spherocytosis-like phenotype

The G1 mouse carrying the MRI61689 mutation was initially identified from an ENU suppressor

screen for the recessive mutation db/db. The G1 MRI61689 exhibited abnormal blood parameters on

an ADVIA haematological analyser, with reduced MCV of 48.6fL compared to the background of

53.3±0.5fL. This MCV values from the B6.BKS(D)-Lepr[db]/J background is comparable to C57BL/6 mice.

We crossed the G1 founder mouse with B6.BKS(D)-Lepr[db]/J to produce G2 mice where approximately

half of the animals exhibited an abnormal phenotype (Table 1). The affected G2 progeny, which

were obligate heterozygotes for the ENU-induced mutation, showed reduction in MCV (46.1±0.2fL)

compared to unaffected progeny (51.4±0.4fL), lower mean corpuscular haemoglobin (MCH)

(13.5±0.1pg compared to 14.6±0.1pg of wild-type), elevated RBC count ($11.1\pm0.1\times10^9$ cells/ml

compared to $10.5\pm0.1\times10^9$ cells/ml of wild-type) (Table 1). No differences were observed for total

haemoglobin (HB), mean corpuscular haemoglobin concentration (MCHC), white blood cell (WBC)

count, platelets count (PLT) or reticulocyte percentage (Table 1).

However, when two affected mice were intercrossed, a quarter of the pups were found to die within

1 week postnatally, suggesting homozygosity for MRI61689 might be incompatible with life. Blood

smears were taken from these pups and compared with the other affected and unaffected mice. The

heterozygotes have slightly smaller RBCs but no target cells or spherocytes were observed (Figure

1a). Conversely, homozygous mice had significantly smaller RBCs with anisocytosis, fragmented RBCs,

acanthocytes and reticulocytosis (Figure 1a). Under SEM, RBCs of heterozygous mice seemed to have

less distinct discoid shape, but otherwise no distinguishing features were observed. (Figure 1b). On

the other hand, the RBCs of homozygous mice appeared very deformed, acanthocytic and appeared

to lack the discoid shape (Figure 1b).

When subjected to osmotic stress, RBCs of heterozygous mice showed significantly increased

fragility compared to wild-type erythrocytes (with 50% haemolysis at approximately 5.6 g/L

compared to 4.5 g/L of wild-type) (Figure 1c). The RBC deformability was assessed using an *in vitro*

spleen retention assay by filtering RBCs through a layer of beads with varying sizes. This is thought to

model splenic filtration *in vivo*, and the retention is thought to indicate deformability. As shown in

Figure 1d, the heterozygous mice exhibited up to 70% of the RBCs retention compared to wild-type

of approximately 3.5% retention, thus suggesting a significantly reduced RBC deformability.

MRI61689 carries a splice site mutation in *Ank-1* gene resulting in an alternative transcript and exon

skipping

To identify the causative mutation responsible for this abnormal RBC count, we sequenced the

exomes of 2 heterozygous mice. Exome sequencing revealed a number of variants. These were

prioritised based on filters as shown in Table 2. Through further genotyping using Sanger sequencing,

a mutation in *Ank-1* gene was found to correlate with all the affected mice and was proven to

segregate perfectly with the reduced MCV for over 3 generations of mouse crosses. The mutation

was found in the 17-18 intron of *Ank-1* gene, with T to A transversion 11 base pair upstream of exon

18 (IVS17-11T>A) (Figure 2a). This is situated in the ankyrin-repeats domain involved in band 3

binding. We proposed that the mutation introduced a new acceptor splice site for exon 18,

potentially leading to a frameshift mutation.

To assess this hypothesis, transcript analysis was performed. Embryonic liver RNA was extracted,

cDNA was synthesized and PCR-amplified using primers listed in the experimental procedures. Figure

2b shows the PCR products of embryonic liver cDNA from non-mutant, *Ank-1*[(MRI61689/+)] and *Ank-*

1[(MRI61689/MRI61689)] when amplified using primer set 1, which covers exon 17 to 21. Bands of

approximately 400bp can be observed in all the genotypes, but *Ank-1*[(MRI61689/MRI61689)] also exhibited a

second smaller product of approximately 300bp length. We proposed that this second band resulted

from exon skipping. Sanger sequencing of these PCR products revealed that the 300bp product

lacked exon 18, confirming that the exon 18 was skipped, and exon 19 was directly connected to

exon 17 during transcription (Figure 2c). This transcript is predicted to produce a shortened, in-

frame 207kDa ANK-1 protein.

To examine the effect of MRI61689 mutation in heterozygous mice, we further designed a primer set

containing the predicted acceptor splice site (primer set 2). Figure 2d shows that the mutant

transcript is only present in $Ank-1^{(MRI61689/+)}$ and $Ank-1^{(MRI61689/MRI61689)}$ mice, as predicted. Further

Sanger sequencing revealed an insertion of 11bp into the transcript adding an additional donor

splicing site and causing a frameshift mutation in the exon that would result in a premature stop

codon at amino acid position 724, as illustrated in Figure 2e, thus giving rise to a truncated protein of

78.5kDa. Therefore the homozygous mice exhibit a mutation at 11 bp upstream of the exon 18

donor splicing site resulting in two alternative transcripts: the skipping of exon 18 and an 11bp

insertion and creation of an additional donor splicing site leading to frameshift mutation that would

result in a premature stop codon and a truncated protein.

We hypothesize that this mutation would reduce the Ank-1 expression levels. We assessed this

hypothesis by examining the gene expression levels of $Ank-1$ in embryonic liver using qPCR at mRNA

level and Western blotting at protein level in mature RBCs. As shown in Figure 3a, $Ank-1$ mRNA

levels in both $Ank-1^{(MRI61689/+)}$ and $Ank-1^{(MRI61689/MRI61689)}$ E14 embryonic livers were significantly

reduced, up to 60% and 80% reduction compared to the wild-type, respectively. However, no

significant reduction in the full length ANK-1 (210kDa) protein levels was observed in both

Coomassie and Western blotting (Figure 3b-d). No truncated form of ANK-1 (78.5kDa) was observed

in $Ank-1^{(MRI61689/+)}$ erythrocytes. Furthermore, no reduction was observed for the protein levels of

other cytoskeletal proteins, including Band 3, protein 4.2, alpha- and beta-spectrin (Figure 3b and 3c).

This suggested that erythrocyte protein levels might be compensated by the WT allele in Ank-

$1^{(MRI61689/+)}$ mice, and the reduction in $Ank-1$ mRNA levels did not seem to affect the protein levels.

Ank-1$^{(MRI61689/+)}$ mice are resistant to *Plasmodium chabaudi* infection

We hypothesized that the *Ank-1$^{(MRI61689)}$* mutation confers malaria resistance. The malaria

susceptibility of *Ank-1$^{(MRI61689/+)}$* mice was examined by injecting a lethal dose of *Plasmodium*

chabaudi adami DS, a murine strain of malaria that models the *Plasmodium falciparum* erythrocytic

stage [33]. The *Ank-1$^{(MRI61689/+)}$* mice exhibited significantly lower peak parasitemia, with only

approximately 13% parasitemia compared to 52% parasitemia of wild-type (Figure 4a) but no delay

in the appearance of parasites was observed. In addition, *Ank-1$^{(MRI61689/+)}$* have a significantly

increased survival rate, where all the *Ank-1$^{(MRI61689/+)}$* mice survived the infection (Figure 4b)

compared to the 16% survival of wild-type mice. Since *Ank-1$^{(MRI61689)}$* directly affects the red cell

(cytoskeletal protein), we hypothesized that the resistance was likely due to a RBC-autonomous

effect. Therefore, we postulated three mechanisms of *P. chabaudi* resistance in *Ank-1$^{(MRI61689/+)}$* mice.

Firstly, the maturation of parasite inside the *Ank-1$^{(MRI61689/+)}$* erythrocytes could be impaired leading

to reduced growth and death of parasites [31]. Secondly, the *Ank-1$^{(MRI61689/+)}$* erythrocytes might be

resistant to merozoite invasion, which resulted in reduced parasitemia and delayed course of

infection [15]. Finally, the infected *Ank-1$^{(MRI61689/+)}$* erythrocytes might more prone to destruction during

the course of infection (increased clearance), thus posing a challenge for the parasite to establish a

successful infection [34].

Ank-1$^{(MRI61689)}$ does not impair the intra-erythrocytic growth of *P. chabaudi*.

To elucidate the possible mechanisms of resistance, the effect of the *Ank-1$^{(MRI61689/+)}$* mutation on

parasite intra-erythrocytic growth was investigated using the TUNEL assay at 1-10% parasitemia.

TUNEL detects DNA fragmentation, a marker for apoptosis or necrosis. In conjunction with a DNA

fluorescent dye, DAPI, it is possible to detect dying parasites in the erythrocytes (Figure 4c) [31,35]. We

measured the TUNEL-positivity of *P. chabaudi* in *Ank-1*^(MRI61689/+) erythrocytes during the late

trophozoite stage of the infections, which was the portion of the parasite lifecycle affected by the

Ank1 mutation in the *Ank-1*^(MRI23420/+) line. As shown in Figure 4d, no differences was observed in the

percentage of TUNEL-positive parasites in both wild-type and *Ank-1*^(MRI61689/+) erythrocytes (24.7 ± 2.3%

in *Ank-1*^(MRI61689/+) mice compared to 29.5 ± 2.9% in wild-type). This indicated that *Ank-1*^(MRI61689) did

not impair parasite intra-erythrocytic growth.

Ank-1^(MRI61689/+) erythrocyte is resistant to *P. chabaudi* invasion, and have increased clearance from

circulation

Erythrocyte invasion and clearance were assessed via an *in vivo* erythrocyte tracking (IVET) assay.

Labelled blood of wild-type and *Ank-1*^(MRI61689/+) mice were injected into infected wild-type mice

during merozoite invasion to examine the ability of *Plasmodium chabaudi* to invade and grow within

erythrocytes of both genotypes. The result compared the percentage of parasitized cells of both

genotypes and expressed as a ratio of parasitemia in *Ank-1*^(MRI61689/+) RBC to wild-type RBC

populations. As shown in Figure 4e, lower parasitemia ratio (approximately 0.55) was observed from

30 minutes after injection with labelled blood, and was consistently lower in *Ank-1*^(MRI61689/+) blood

over 36 hours post injection, which indicates a lower invasion rate into the *Ank-1*^(MRI61689/+) RBCs.

Additionally, the remaining proportion of labelled RBCs was also monitored over the course of the

assay. A significant reduction of *Ank-1*^(MRI61689/+) erythrocytes in infected mice compared to wild-type

erythrocytes was observed, with up to a 45% reduction in *Ank-1*^(MRI61689/+) RBC number compared

with wild-type (Figure 4f). On the other hand, a smaller reduction was observed for *Ank-1*^(MRI61689/+)

erythrocytes compared to wild-type in uninfected mice, suggesting *Ank-1*^(MRI61689/+) RBCs are more

likely to get cleared from circulation during malaria infection. As the percentage of infected

erythrocytes was low (5-20%) (Supp. figure 1), this indicates that the majority of the RBCs getting

cleared were uninfected RBCs, possibly as a result of bystander effect. This experiment suggested

that two possible mechanisms of resistance are both operating to produce the lower parasitemia

and increased survival in $Ank-1^{(MRI61689/+)}$ mice, the reduction of parasite invasion and increased

clearance of $Ank-1^{(MRI61689/+)}$ RBCs.

Discussion

We report a novel mutation in $Ank-1$ gene, MRI61689, causing a hereditary spherocytosis-like

phenotype, with reduced MCV, increased osmotic fragility and reduced deformability. MRI61689 is

an intronic mutation between exon 17 and 18 where two possible splice variants could arise, one

has an introduced acceptor site resulting a frameshift, whereas the other consists of a skipped exon

18. The $Ank-1$ mRNA levels were reduced, but no reduction in protein levels were observed. The

predicted truncated form (78.5kDa) was also not observed. $Ank-1^{(MRI61689/+)}$ mice also have increased

resistance to *Plasmodium chabaudi* infections, and the erythrocyte invasion was impaired but the

intra-erythrocytic growth appeared normal. The $Ank-1^{(MRI61689/+)}$ RBCs were also more likely to be

cleared from circulation during infection, an observation independent of number of parasitized

erythrocytes.

In comparison of $Ank-1^{MRI61689}$ mice to other previously described $Ank-1$ mouse models, they

appeared comparable to $Ank-1^{MRI23420}$ mice, but more severe than $Ank-1^{1674}$ and $Ank-1^{nb}$ mice. More

specifically, most homozygous $Ank-1^{(MRI61689)}$ mice died within a week after birth, similar to

homozygous $Ank-1^{MRI23420}$ mice, while homozygous $Ank-1^{1674}$ and $Ank-1^{nb}$ mice were viable [30,32].

However, no notable differences were observed in heterozygous $Ank-1^{MRI61689}$ mice in terms of their

RBC microcytosis, morphology and susceptibility to osmotic stress compared to heterozygous $Ank-$

1^{1674} and $Ank-1^{MRI23420}$ mice [31,32]. However, similar to $Ank-1^{1674/+}$ mice and unlike $Ank-1^{(MRI23420/+)}$ mice,

$Ank-1^{(MRI61689/+)}$ mice exhibited similar levels of ankyrin and other RBC cytoskeletal protein to wild-

type, which might suggest compensation by the wild-type allele, and thus warrants further studies.

In humans, many *ANK-1* mutations that result in frameshift have been described, most of which

situated in the band 3 binding domain towards the N terminus [36,37]. While no *Ank-1*[(MRI61689)]

homologous mutation has been described in humans, a frameshift mutation has been reported to be

in the exon 17, called Ankyrin Osaka I, which gave rise to symptomatic HS [37]. Furthermore, exon

skipping in human *ANK-1* gene has also been documented. Edelman, et al. [38] reported a HS patient

exhibiting a severe ankyrin-deficient HS due to an introduction of a new splice acceptor site for exon

17, known as ankyrin[Ankara]. Under further examination, this mutation was found to give rise to

multiple splice forms, including insertions and skipped exons. Most splice forms with insertion were

expected to cause frameshift, potentially leading to premature termination of ankyrin [38]. This finding

is in agreement with what we observed for *Ank-1*[(MRI61689/MRI61689)] mice, where frameshift caused by

new splice acceptor site, leading to high mortality rate with severe HS-like phenotype in *Ank-*

1[(MRI61689/MRI61689)] mice. It is likely that the surviving *Ank-1*[(MRI61689/MRI61689)] mice relied on the exon-

skipping splice form to produce in-frame functional *Ank-1* proteins. However, further studies are

required for support this hypothesis.

In terms of the response to malaria infection, *Ank-1*[(MRI61689/+)] mice exhibited similar degree of

malaria resistance as *Ank-1*[(MRI23420/+)] and *Ank-1*[1674/+] mice, with at least 30-40% reduction in

parasitemia and increased survival [31,32], and unlike *Ank-1*[nb/+] mice with only 10% reduction [30].

Previous studies suggested the reduction in erythrocyte invasion and intra-erythrocytic growth to be

the major resistance mechanisms [30,31]. However, normal parasite invasion was reported in *Ank-1*[1674/+]

mice [32]. As a result, we have explored the potential mechanisms of *Ank-1*[MRI61689/+] mice in this study

in attempt to elucidate the complex roles ankyrin plays during malaria infections.

First, we report that *Ank-1*[(MRI61689/+)] mice exhibit normal parasite intra-erythrocytic growth (Figure

4d), in contrast to *Ank-1*[(MRI23420/+)] mice. TUNEL assay detects the presence of DNA fragmentation

which occurs during apoptosis and necrosis [39], which indicates dying parasites in RBCs [35]. McMorran,

et al. [35] reported TUNEL-positive parasites in C57BL/6 wild-type mice at a level consistent with the

observations in this study. On the other hand, Greth, et al. [31] reported a lower TUNEL-positive in

their SJL/J wild-type mice, which most likely due to differences in the genetic background of

experimental mice. Nevertheless, we did not observe abnormal parasite morphology under light

microscopy unlike *Ank-1*$^{(MRI23420/+)}$ mice, which support the deductions of parasite death from the

TUNEL assays.

In terms of parasite invasion, *Ank-1*$^{(MRI61689/+)}$ RBCs were found to be more resistant to merozoite

invasion as shown in the IVET assay (Figure 4e). The *Ank-1*$^{(MRI61689/+)}$ erythrocytes were less infected

compared to the wild-type 30 minutes after injection during merozoite invasion, and stayed

consistently lower compared to wild-type throughout the erythrocytic cycle. This reduction in

invasion has also been observed in *Ank-1*$^{(MRI23420/+)}$ mice [31]. In contrast, Rank, et al. [32] reported no

difference in parasite invasion for *Ank-1*$^{1674/+}$ mice, indicating a different effect mediated by *Ank-1*^{1674}

mutation compared to *Ank-1*$^{(MRI23420)}$ and *Ank-1*$^{(MRI61689)}$ mutations.

From these comparisons with other ankyrin mouse models, it is evident that RBC cytoskeleton plays

an important yet complex role during malaria infections. However the exact mechanism for each of

these different phenotypes for each ankyrin haplotype remains elusive, different ankyrin mutations

can exert different effects on the parasites depending on the location of the mutations, giving rise to

multiple resistance mechanisms. This hypothesis is also consistent with the heterogeneous HS

symptoms associated with ankyrin mutations, which highlights the complicated interactions

between RBC cytoskeletons and malaria parasites.

On the other hand, one important observation from the IVET assay is the rapid clearance of *Ank-*

1$^{(MRI61689/+)}$ erythrocytes from the circulation within 36 hours post-injection, up to 40% of the initial

RBC number (Figure 4f). However, at this timepoint the parasitemia of the host mice was only 5-20%

(Supp. Figure 1a), therefore the rate of RBC clearance cannot be explained by the clearance of

parasitized RBCs, instead, it is likely that the majority of the RBCs being cleared were uninfected. In

comparison, no loss was observed for wild-type erythrocytes in both infected and uninfected mice.

Since, in these experiments both wild-type and *Ank-1*[(MRI61689/+)] blood were subjected to the same

host environment simultaneously, it implies that the clearance of *Ank-1*[(MRI61689/+)] erythrocytes is cell

autonomous, rather than due to other effects of the host animal. This is the first observation of

increased, uninfected RBC clearance associated with an ankyrin mutation. Bystander clearance is

typically observed in inflammation, such as during sepsis [40], and is thought to cause severe malaria

anaemia during malaria infection in humans through the destruction of normal uninfected RBCs [41,42].

It is possible that *Ank-1*[(MRI61689)] causes a more exaggerated bystander effect during malaria infection,

leading to a rapid reduction of *Ank-1*[(MRI61689/+)] erythrocyte numbers.

Reduced RBC deformability was proposed to be one of the mechanisms of bystander clearance

during malaria anaemia through phagocytosis and splenic filtration [43,44]. From the *in vitro* spleen

retention assay (Figure 1d), *Ank-1*[(MRI61689/+)] erythrocytes exhibited reduced deformability, making

them more likely to be retained in the filter layer, which is likely to promote their destruction in *in*

vivo settings. Therefore, we proposed that *Ank-1*[MRI61689] mutation causes alteration to the

erythrocyte, which renders them more likely to be cleared from the circulation.

In summary, we report that ENU-induced *Ank-1*[MRI61689] mutation causes an HS-like phenotype in mice,

and confers significant resistance to *P. chabaudi* infection. We propose that *Ank-1*[(MRI61689/+)]

erythrocytes are significantly resistant to parasite invasion but are able to support normal parasite

intra-erythrocytic growth. We also described a novel observation of increased RBC clearance

associated with this ankyrin mutation. This study emphasizes the importance of RBC cytoskeletal

proteins in mediating multiple complex mechanisms of resistance towards malaria, which provide

further insights to the complex interaction between the host and parasites.

Methods and Materials

Mice and Ethics Statement

All mice used in this study were housed with 12 hour light-dark cycle under constant temperature at

21 °C. All procedures were conformed to the National Health and Medical Research Council (NHMRC)

Australian code of practice. Experiments were performed under ethics agreement AEEC A2014/54,

which were approved by the Australian National University animal ethics committees.

ENU Mutagenesis and Dominant Phenotype Screening

B6.BKS(D)-Lepr^{db}/J male mice were injected intraperitoneally with two dose of 100 mg/kg ENU

(Sigma-Aldrich, St Louis, MO) at one week interval for a recessive suppressor screen of db/db

mutation. The treated males (G0) were crossed to females from the isogenic background to produce

the first generation progeny (G1). The seven-week-old G1 progeny were bled and analysed on an

Advia 120 Automated Haematology Analyser (Siemens, Berlin, Germany) to identify abnormal red

blood cell count. Mouse carrying MRI61689 mutation was identified with a "mean corpuscular

volume" (MCV) value three standard deviations lower from other G1 progeny. It was crossed with

B6.BKS(D)-Lepr^{db}/J mice to produce G2 progeny to test the heritability of the mutations and the

dominance mode of inheritance. Mice that exhibited low MCV (<48fL) were selected for whole

exome sequencing and genotyping. This abnormal red blood cell count was unrelated to the obesity

phenotype.

Microscopy

For light microscopy, cells were briefly fixed in methanol for one minute and air-dried before being

stained in a 10% Giemsa solution (Sigma-Aldrich, St Louis, MO) at pH 7.4 for 10 minutes. For

scanning electron microscopy (SEM), fresh blood was first fixed immediately upon drawing in 3%

EM-grade glutaraldehyde (Sigma-Aldrich, St Louis, MO) overnight at 4°C. The samples were washed

with mouse tonicity phosphate buffered saline (MT-PBS) (150mM NaCl, 16mM Na$_2$HPO$_4$, 4mM

NaH$_2$PO$_4$, pH 7.4) 3 times, 10 minutes soak each. The cells were then adhered to the cover slips with

0.1% polyethyleneimine (PEI) for 10 minutes, before washing off with MT-PBS. The cells were then

dried serially using 30%, 50%, 70%, 80%, 90%, 100%, 100% ethanol, each with 10 minutes soak. The

cells were then soaked in 1:1 ethanol: hexamethyldisilazane solution for 10 minutes, followed by 2

washes with 100% hexamethyldisilazane, each 10 minutes. The coverslips were then air-dried

overnight and coated with gold and examined under JEOL JSM-6480LV scanning electron microscope.

Osmotic Fragility Measurement

To assess the susceptibility of RBC membrane to osmotic stress, 5μl of mouse whole blood was

diluted 100-fold with phosphate buffer (pH 7.4) containing 0 to 10g/L of sodium, and incubated for

at least 10 minutes at room temperature. The cells were centrifuged at 800g for 3 minutes, and the

supernatant, which contains free haemoglobin, was measured at 540nm to assess the degree of

haemolysis. The absorbance values were expressed as percentage of haemolysis, with haemolysis at

0g/L sodium considered as 100% lysis.

In vitro spleen retention assay

The RBC deformability of both wild-type and *Ank-1*$^{(MRI61689/+)}$ were assessed according to the protocol

described previous by Deplaine, et al. [45] with modifications. Briefly, RBCs from wild-type and *Ank-*

1$^{(MRI61689/+)}$ mice were stained with 10μg/ml of either hydroxysulfosuccinimide Atto 633 (Atto 633) or

hydroxysulfosuccinimide Atto 565 (Atto 565) (Sigma-Aldrich, St Louis, MO), followed by three

washes with in MTRC (154mM NaCl, 5.6mM KCl, 1mM MgCl$_2$, 2.2mM CaCl$_2$, 20mM HEPES, 10mM

glucose, 4mM EDTA, 0.5% BSA, pH 7.4, filter sterilized). The stained RBCs were mixed in equal

proportion and diluted with unstained wild-type RBCs to give approximately 10-20% of the sample

being labelled RBCs. The samples were further diluted to 1-2% haematocrit with MTRC, before

passing through spleen retention filter bed. The pre-filtered and post-filtered samples were analysed

on BD LSRFortessa (BD Biosciences, Franklin Lakes, NJ) flow cytometer to determine the proportion

being retained in the filter bed.

Whole exome sequencing

DNA from two G2 mice carrying the abnormal red blood cell parameters (MCV <48fL) were extracted

with Qiagen DNeasy blood and tissue kit (Qiagen, Venlo, Netherlands) for exome sequencing as

previous described [46]. Briefly, at least 10μg of DNA was prepared for exome enrichment with Agilent

Sure select kit paired-end genomic library from Illumina (San Diego, CA), followed by high

throughput sequencing using a HiSeq 2000 platform. The bioinformatics analysis was conducted

according to the variant filtering method previously described by Bauer, et al. [47]. Private variants

that were shared between the two mutants but not with other B6.BKS(D)-Lepr[db]/J, C57BL/6 mice or

previously described ENU mutants were annotated using ANNOVAR [48]. Private non-synonymous

exonic and intronic variants within 20 bp from the exon spicing sites were retained as potential

candidate ENU mutations.

PCR and Sanger sequencing

DNA from mutant mice were amplified through PCR with 35 cycles of 30 seconds of 95^0C

denaturation, 30 seconds of 56-58^{0}C annealing and 72^{0}C elongation for 40 seconds. The primers

used in the PCR are described as below. The PCR products were examined with agarose gel

electrophoresis before being sent to the Australian Genome Research Facility (AGRF) in Melbourne,

Australia, for Sanger sequencing. Logarithm of odds (LOD) score was calculated based on the

number of mice that segregated with the candidate mutations.

Primers:

Acp5-F: 5'-CAGAAGGATGCCTTTGGGTA-3'; *Acp5*-R: 5'-ACCAGCGCTTGGAGATCTTA-3'

Kcnk1-F: 5'-GGGCCTTTTCCTCCTTACAGA-3'; *Kcnk1*-R: 5'-CAGGAAACGGTGACAAATCC-3'

Epas1-F: 5'-GGAAGCCAGAACTTCGATGA-3'; *Epas1*-R: 5'-GTAGTGTTCCCTGGGGTGT-3'

Picalm-F: 5'-TCACTGAATGTAATTGGGATATCAT-3'; *Picalm*-R: 5'-CACCCTCTCTTCACTTTTGTG-3'

Socs6-F: 5'-CCGCTTTGTTATCCGTCAGT-3'; *Socs6*-R: 5'-TGGCAGCAAAGACTTCAATG-3'

Ank1-F: 5'-TCCCTGGCTTAAAGTTGGTG-3'; *Ank1*-R: 5'-CTCTCCCTTAGCTGCATTCC-3'

Quantitative PCR and cDNA sequencing

RNA was isolated from embryonic livers of E14 embryos using Qiagen RNeasy kit (Qiagen, Venlo,

Netherlands), followed by cDNA synthesis using Transcriptor High Fidelity cDNA Synthesis Kit (Roche,

Basel, Switzerland). Quantitative PCR was carried out on ViiA™ 7 Real-Time PCR System (Thermo

Scientific, Waltham, MA). The $\Delta\Delta C_T$ method [49] was used to determine the cDNA levels of *Ank-1* and

the housekeeping gene β-actin and expressed as a fold-change of the mutants to the wild-type. The

primers used for *Ank-1* gene spanned exon 2 to 4: *Ank-1*-F: 5'-TAACCAGAACGGGTTGAACG-3'; *Ank-1*-

R: 5'-TGTTCCCCTTCTTGGTTGTC-3'; β-Actin-F: 5'- TTCTTTGCAGCTCCTTCGTTGCCG-3'; β-Actin-R: 5'-

TGGATGCGTACGTACATGGCTGGG-3'.

To characterize the effect of MRI61689 mutation, cDNA were amplified through PCR using two

primers set were design as shown below: Primer set 1 was designed to amplify wild-type *Ank-1*

transcript, whereas primer set 2 was designed to only amplify the predicted mutant transcript with

11bp insertion. Amplified PCR product were analysed using agarose gel electrophoresis and each

band was purified and sequenced.

Primer set 1: Forward: ATGCAGAGTCGGTACAAGGC; Reverse: CCGTTCGAGCTGACCTCATT

Primer set 2: Forward: CCTGGGGAACAAGTTTCTTT; Reverse: GTGCAAGGGGCTGTATCCTA

SDS-PAGE, Coomassie staining and Western blot

RBC ghosts were prepared by lysing mouse RBCs with ice-cold 5mM phosphate buffer (ph7.4) and

centrifuged at 20,000g for 20 minutes followed by removal of the supernatant. The pellet was

further washed with the 5mM phosphate buffer until the supernatant became clear. The RBC ghosts

or whole blood lysates were denatured in SDS-PAGE loading buffer (0.0625M Tris pH 6.8, 2% SDS, 10%

glycerol, 0.1M DTT, 0.01% bromophenol blue) at 95°C for 5 minutes before loading onto a Mini-

PROTEAN® TGX™ Precast Gels (Bio-Rad, Hercules, CA). The gels were then either stained with

Coomassie blue solution (45% v/v methanol, 7% v/v acetic acid, 0.25% w/v Brilliant Blue G) overnight

or transferred to a nitrocellulose membrane. The western blot was carried out using these primary

antibodies: anti-alpha 1 spectrin (clone 17C7), anti-beta 1 spectrin (clone 4C3) (Abcam, Cambridge,

UK), anti-GAPDH (clone 6C5) (Merck Millipore, Darmstadt, Germany), anti-N-terminal *Ank-1* "p89",

anti-Band 3 and anti-protein 4.2 (kind gifts from Connie Birkenmeier, Jackson Laboratory, US).

Malaria infection

250μl of thawed *P. chabaudi adami* infected blood was injected into the intraperitoneal cavity of a

C57BL/6 donor mouse. When the donor mouse reached 1-10% parasite load (parasitemia), blood

was collected through cardiac puncture. The parasitized blood was diluted with Krebs' buffered

saline with 0.2% glucose as described previously [50]. Each experimental mouse was infected with

1×10^4 parasites intraperitoneally. The parasitemia of these mice were monitored either using light

microscopy or flow cytometry.

Terminal deoxynucleotidyl transferase dUTP nick end labelling (TUNEL) staining

3μl of infected blood containing 1-10% parasitemia were collected during trophozoite stage and

fixed in 1 in 4 diluted BD Cytofix™ Fixation Buffer (BD Biosciences, Franklin Lakes, NJ) for at least day

until they were needed. Each sample was then washed twice with MT-PBS, and adhered to a glass

slide pre-coated with 0.1% polyethylenimine (PEI) for 10 minutes at room temperature. The excess

cells were washed off with the wash solution from APO-BrdU TUNEL assay kit (Thermo Scientific,

Waltham, MA) and incubated overnight at room temperature with TUNEL labelling solution (1mM

Cobalt Chloride, 25mM Tris-HCl pH 6.6, 200mM sodium cacodylate, 0.25mg/ml BSA, 60uM BrdUTP,

15U Terminal transferase). The slides were washed three times with rinse buffer from APO-BrdU

TUNEL assay kit, followed by staining with 50μg/ml of anti-BrdU-Biotin antibody (Novus Biologicals,

Littleton, CO) in MT-PBT (MT-PBS, 0.5% BSA, 0.05% Triton X-100) for 1 hour. The slides were then

washed three times with MT-PBT, followed by probing with 2μg/ml Alexa Fluor® 594 conjugated

streptavidin (Thermo Scientific, Waltham, MA). Next, they were washed three times with MT-PBS

and mounted with SlowFade® Gold antifade reagent with DAPI (Thermo Scientific, Waltham, MA)

and sealed. When the slides were dried, they were examined using Axioplan 2 fluorescence light

microscope (Carl Zeiss, Oberkochen, Germany) between 600x to 1000x magnification. At least 100

DAPI-positive cells were counted, and each was graded as either positive or negative for TUNEL

staining, as an indication of DNA fragmentation.

In vivo erythrocyte tracking (IVET) assays

The IVET assay was carried out as previously described by Lelliott, et al. [51,52]. Briefly, 1.5ml to 2ml of

whole blood was collected from wild-type and *Ank-1*[(MRI61689/+)] mice via cardiac puncture. Both wild-

type and *Ank-1*[(MRI61689/+)] blood were either stained with 10μg/ml of Atto 633 or 125μg/ml of EZ-

Link™ Sulfo-NHS-LC-Biotin (Biotin) (Thermo Scientific, Waltham, MA) for 45 minutes at room

temperature, followed by washing three times with MT-PBS. The *Ank-1*[(MRI61689/+)] blood was mixed

with wild-type blood in two different dye combinations to correct for any dye effects. 1×10^9

erythrocytes were injected intravenously into infected wild-type mice at 1-5% parasitemia during

schizogony stage, usually 8-10 days post-infection with 1×10^{4} parasites. Blood samples were

collected at 30 minutes, 3hours, 12 hours, 20 hours and 36 hours after injection. The ratio of

infected $Ank\text{-}1^{(MRI61689/+)}$ to wild-type erythrocytes was determined on flow cytometry, as an

indication of the relative susceptibility to malaria infections between wild-type and $Ank\text{-}1^{(MRI61689/+)}$

mice. The proportion of labelled blood populations were also tracked over time to determine the

clearance of these RBCs from the circulation.

Flow cytometry analysis of blood samples

For both malaria infections and IVET assay, 2µl of whole blood samples were stained with 2µg/ml

streptavidin-PE-Cy7 (only for experiments with biotinylated erythrocytes), 1µg/ml anti-CD45–

allophycocyanin (APC)–eFluor 780 (clone 30-F11), 1µg/ml anti-CD71 (TFR1)–PerCP–eFluor 710 (clone

R17217) (eBioscience, San Diego, CA), 4µM Hoechst 33342 (Sigma-Aldrich, St Louis, MO) and 12µM

JC-1 (Thermo Scientific, Waltham, MA) in MTRC. All samples analysed through flow cytometry were

performed on BD LSRFortessa (BD Biosciences, Franklin Lakes, NJ), where 100,000 to 2,000,000

events were collected and visualized on FACSDiva^{TM} and FlowJo software. The RBCs and leukocytes

were first selected on forward scatter and side scatter channels (FSC/SSC) signals, followed by gating

of single cells based on FS area to height ratio. RBCs were further isolated by gating on CD71

negative and CD45 negative population, followed by gating on Atto-labelled and Biotin-labelled

erythrocytes on appropriate channels (APC for Atto-633, PE for Atto-565 and PE-Cy7 for Biotin). The

parasitemia of each labelled erythrocyte population was determined by gating on Hoechst 33342

positive and JC-1 positive population.

Statistical analysis

The LOD score method coupled with Bonferroni correction was used to determine the causative

mutation for MRI61689. The statistical significance of the malaria survival was tested using the Log-

Rank test. The statistical significance of parasite infection was determined via the statmod software

package   for   R   (http://bioinf.wehi.edu.au/software/compareCurves)   using   the

'compareGrowthCurves' function with 10,000 permutation, followed by adjustments for multiple

testing. The statistical significance for the ratios of IVET assays was determined using the one sample

t-test with hypothetical mean of 1. For the rest of the results, statistical significance was determined

using two-tailed Students t-tests.

References

1   W.H.O. *World Malaria Report 2015*. (World Health Organisation (WHO), 2015).

2   Kwiatkowski, D. P. How malaria has affected the human genome and what human genetics
can teach us about malaria. *Am J Hum Genet* **77**, 171-192 (2005).

3   Piel, F. B. *et al.* Global distribution of the sickle cell gene and geographical confirmation of
the malaria hypothesis. *Nature communications* **1**, 104, doi:10.1038/ncomms1104 (2010).

4   Carter, R. & Mendis, K. N. Evolutionary and historical aspects of the burden of malaria. *Clin*
Microbiol Rev **15**, 564-594 (2002).

5   Zimmerman, P. A. *et al.* Emergence of FY*A(null) in a Plasmodium vivax-endemic region of
Papua New Guinea. *Proc Natl Acad Sci U S A* **96**, 13973-13977 (1999).

6   Miller , L. H., Mason , S. J., Clyde , D. F. & McGinniss , M. H. The Resistance Factor to
Plasmodium vivax in Blacks. *New England Journal of Medicine* **295**, 302-304,
doi:doi:10.1056/NEJM197608052950602 (1976).

7   Cyrklaff, M. *et al.* Hemoglobins S and C interfere with actin remodeling in Plasmodium
falciparum-infected erythrocytes. *Science* **334**, 1283-1286, doi:10.1126/science.1213775
(2011).

8   Allen, S. J. *et al.* Prevention of cerebral malaria in children in Papua New Guinea by
southeast Asian ovalocytosis band 3. *Am J Trop Med Hyg* **60**, 1056-1060 (1999).

9   Jarolim, P. *et al.* Deletion in erythrocyte band 3 gene in malaria-resistant Southeast Asian
ovalocytosis. *Proceedings of the National Academy of Sciences of the United States of*
America **88**, 11022-11026 (1991).

10   Mgone, C. S. *et al.* Occurrence of the erythrocyte band 3 (AE1) gene deletion in relation to
malaria endemicity in Papua New Guinea. *Trans R Soc Trop Med Hyg* **90**, 228-231 (1996).

11   Genton, B. *et al.* Ovalocytosis and cerebral malaria. *Nature* **378**, 564-565,
doi:10.1038/378564a0 [doi] (1995).

12   Lecomte, M. C. *et al.* [Hereditary elliptocytosis in West Africa: frequency and repartition of
spectrin variants]. *C R Acad Sci III* **306**, 43-46 (1988).

13   Facer, C. A. Erythrocytes carrying mutations in spectrin and protein 4.1 show differing
sensitivities to invasion by Plasmodium falciparum. *Parasitol Res* **81**, 52-57 (1995).

14   Kidson, C., Lamont, G., Saul, A. & Nurse, G. T. Ovalocytic erythrocytes from Melanesians are
resistant to invasion by malaria parasites in culture. *Proceedings of the National Academy of*
Sciences of the United States of America **78**, 5829-5832 (1981).

15   Hadley, T. *et al.* Resistance of Melanesian elliptocytes (ovalocytes) to invasion by
Plasmodium knowlesi and Plasmodium falciparum malaria parasites in vitro. *J Clin Invest* **71**,
780-782 (1983).

16   Schulman, S. *et al.* Growth of Plasmodium falciparum in human erythrocytes containing
abnormal membrane proteins. *Proc Natl Acad Sci U S A* **87**, 7339-7343 (1990).

17   Chishti, A. H., Palek, J., Fisher, D., Maalouf, G. J. & Liu, S. C. Reduced invasion and growth of
Plasmodium falciparum into elliptocytic red blood cells with a combined deficiency of
protein 4.1, glycophorin C, and p55. *Blood* **87**, 3462-3469 (1996).

18   Cortes, A., Benet, A., Cooke, B. M., Barnwell, J. W. & Reeder, J. C. Ability of Plasmodium
falciparum to invade Southeast Asian ovalocytes varies between parasite lines. *Blood* **104**,
2961-2966, doi:10.1182/blood-2004-06-2136 2004-06-2136 [pii] (2004).

19   Dua, M., Raphael, P., Sijwali, P. S., Rosenthal, P. J. & Hanspal, M. Recombinant falcipain-2
cleaves erythrocyte membrane ankyrin and protein 4.1. *Mol Biochem Parasitol* **116**, 95-99
(2001).

20   Nicolas, V. *et al.* Rh-RhAG/Ankyrin-R, a New Interaction Site between the Membrane Bilayer
and the Red Cell Skeleton, Is Impaired by Rhnull-associated Mutation. *Journal of Biological*
Chemistry **278**, 25526-25533, doi:10.1074/jbc.M302816200 (2003).

21   Grey, J. L., Kodippili, G. C., Simon, K. & Low, P. S. Identification of contact sites between
ankyrin and band 3 in the human erythrocyte membrane. *Biochemistry* **51**, 6838-6846,
doi:10.1021/bi300693k (2012).

22   Su, Y. *et al.* Associations of protein 4.2 with band 3 and ankyrin. *Molecular and cellular*
biochemistry **289**, 159-166, doi:10.1007/s11010-006-9159-x (2006).

23   Eber, S. W. *et al.* Ankyrin-1 mutations are a major cause of dominant and recessive
hereditary spherocytosis. *Nat Genet* **13**, 214-218, doi:10.1038/ng0696-214 (1996).

24   Casale, M. & Perrotta, S. Splenectomy for hereditary spherocytosis: complete, partial or not
at all? *Expert review of hematology* **4**, 627-635, doi:10.1586/ehm.11.51 (2011).

25   Godal, H. C. & Heisto, H. High prevalence of increased osmotic fragility of red blood cells
among Norwegian blood donors. *Scandinavian journal of haematology* **27**, 30-34 (1981).

26   Eber, S. W., Pekrun, A., Neufeldt, A. & Schroter, W. Prevalence of increased osmotic fragility
of erythrocytes in German blood donors: screening using a modified glycerol lysis test.
Annals of hematology **64**, 88-92 (1992).

27   Jensson, O., Jonasson, J. L. & Magnusson, S. Studies on herediatary spherocytosis in Iceland.
Acta medica Scandinavica **201**, 187-195 (1977).

28   Yawata, Y. *et al.* Characteristic features of the genotype and phenotype of hereditary
spherocytosis in the Japanese population. *Int J Hematol* **71**, 118-135 (2000).

29   Glele-Kakai, C. *et al.* Epidemiological studies of spectrin mutations related to hereditary
elliptocytosis and spectrin polymorphisms in Benin. *Br J Haematol* **95**, 57-66 (1996).

30   Shear, H. L., Roth, E. F., Jr., Ng, C. & Nagel, R. L. Resistance to malaria in ankyrin and spectrin
deficient mice. *Br J Haematol* **78**, 555-560 (1991).

31   Greth, A. *et al.* A novel ENU-mutation in ankyrin-1 disrupts malaria parasite maturation in
red blood cells of mice. *PLoS One* **7**, e38999, doi:10.1371/journal.pone.0038999 (2012).

32   Rank, G. *et al.* Novel roles for erythroid Ankyrin-1 revealed through an ENU-induced null
mouse mutant. *Blood* **113**, 3352-3362, doi:blood-2008-08-172841 [pii] 10.1182/blood-2008-
08-172841 (2009).

33   Stephens, R., Culleton, R. L. & Lamb, T. J. The contribution of *Plasmodium chabaudi* to our
understanding of malaria. *Trends in Parasitology* **28**, 73-82, doi:10.1016/j.pt.2011.10.006
(2012).

34 Ayi, K., Turrini, F., Piga, A. & Arese, P. Enhanced phagocytosis of ring-parasitized mutant
erythrocytes: a common mechanism that may explain protection against falciparum malaria
in sickle trait and beta-thalassemia trait. *Blood* **104**, 3364-3371, doi:10.1182/blood-2003-11-
3820 (2004).
35 McMorran, B. J. *et al.* Platelets Kill Intraerythrocytic Malarial Parasites and Mediate Survival
to Infection. *Science* **323**, 797-800, doi:10.1126/science.1166296 (2009).
36 Gallagher, P. G. Hematologically important mutations: ankyrin variants in hereditary
spherocytosis. *Blood Cells Mol Dis* **35**, 345-347, doi:10.1016/j.bcmd.2005.08.008 (2005).
37 Nakanishi, H., Kanzaki, A., Yawata, A., Yamada, O. & Yawata, Y. Ankyrin gene mutations in
japanese patients with hereditary spherocytosis. *Int J Hematol* **73**, 54-63 (2001).
38 Edelman, E. J., Maksimova, Y., Duru, F., Altay, C. & Gallagher, P. G. A complex splicing defect
associated with homozygous ankyrin-deficient hereditary spherocytosis. *Blood* **109**, 5491-
5493, doi:10.1182/blood-2006-09-046573 (2007).
39 Kyrylkova, K., Kyryachenko, S., Leid, M. & Kioussi, C. Detection of apoptosis by TUNEL assay.
Methods Mol Biol **887**, 41-47, doi:10.1007/978-1-61779-860-3_5 (2012).
40 Straat, M., van Bruggen, R., de Korte, D. & Juffermans, N. P. Red blood cell clearance in
inflammation. *Transfusion medicine and hemotherapy : offizielles Organ der Deutschen
Gesellschaft fur Transfusionsmedizin und Immunhamatologie* **39**, 353-361,
doi:10.1159/000342229 (2012).
41 Jakeman, G., Saul, A., Hogarth, W. & Collins, W. Anaemia of acute malaria infections in non-
immune patients primarily results from destruction of uninfected erythrocytes. *Parasitology*
119, 127-133 (1999).
42 Lamikanra, A. A. *et al.* Malarial anemia: of mice and men. *Blood* **110**, 18-28 (2007).
43 Dondorp, A. M. *et al.* Prognostic significance of reduced red blood cell deformability in
severe falciparum malaria. *Am J Trop Med Hyg* **57**, 507-511 (1997).
44 Dondorp, A. *et al.* The role of reduced red cell deformability in the pathogenesis of severe
falciparum malaria and its restoration by blood transfusion. *Transactions of the Royal Society
of Tropical Medicine and Hygiene* **96**, 282-286 (2002).
45 Deplaine, G. *et al.* The sensing of poorly deformable red blood cells by the human spleen can
be mimicked in vitro. *Blood* **117**, e88-95, doi:10.1182/blood-2010-10-312801 (2011).
46 Hortle, E. *et al.* Adenosine monophosphate deaminase 3 activation shortens erythrocyte
half-life and provides malaria resistance in mice. *Blood*, doi:10.1182/blood-2015-09-666834
(2016).
47 Bauer, D. C., McMorran, B. J., Foote, S. J. & Burgio, G. Genome-wide analysis of chemically
induced mutations in mouse in phenotype-driven screens. *BMC Genomics* **16**, 1-8,
doi:10.1186/s12864-015-2073-4 (2015).
48 Wang, K., Li, M. & Hakonarson, H. ANNOVAR: functional annotation of genetic variants from
high-throughput sequencing data. *Nucleic Acids Research* **38**, e164-e164,
doi:10.1093/nar/gkq603 (2010).
49 Livak, K. J. & Schmittgen, T. D. Analysis of relative gene expression data using real-time
quantitative PCR and the 2− ΔΔCT method. *methods* **25**, 402-408 (2001).
50 Jarra, W. & Brown, K. Protective immunity to malaria: studies with cloned lines of
Plasmodium chabaudi and P. berghei in CBA/Ca mice. I. The effectiveness and inter‐and
intra‐species specificity of immunity induced by infection. *Parasite immunology* **7**, 595-606
(1985).
51 Lelliott, P. M., Lampkin, S., McMorran, B. J., Foote, S. J. & Burgio, G. A flow cytometric assay
to quantify invasion of red blood cells by rodent Plasmodium parasites in vivo. *Malar J* **13**,
100, doi:10.1186/1475-2875-13-100 (2014).
52 Lelliott, P. M., McMorran, B. J., Foote, S. J. & Burgio, G. In vivo assessment of rodent
Plasmodium parasitemia and merozoite invasion by flow cytometry. *JoVE (Journal of
Visualized Experiments)*, e52736-e52736 (2015).

Acknowledgement

We would like to acknowledge Shelley Lampkin and Australian Phenomics Facility (APF) for the

maintenance of the mouse colonies. This study was funded by the National Health and Medical

Research Council of Australia (Program Grant 490037, and Project Grants 605524 and APP1047090),

Australian Society of Parasitology (ASP), OzEMalaR, National Collaborative Research Infrastructure

Strategy (NCRIS), the Education Investment Fund from the Department of Education and Training,

the Australian Phenomics Network, Howard Hughes Medical Institute and the Bill and Melinda Gates

Foundation.

   Figure Legends

Figure 1. **The phenotypic characterisation of Ank-1$^{(MRI61689/+)}$ mice.** The morphology of Ank-1$^{(MRI61689/+)}$

and Ank-1$^{(MRI61689/MRI61689)}$ erythrocytes under light microscopy with Giemsa stain (a) and scanning

electron microscopy (b). The osmotic fragility curve of wild-type and Ank-1$^{(MRI61689/+)}$ erythrocytes

when subjected to osmotic stress (c) (n=5-7 per group). The *in-vitro* spleen retention rate of wild-

type and Ank-1$^{(MRI61689/+)}$ erythrocytes when passing through filter beds (d) (n=3 per group). **

indicates P<0.01, *** indicates P<0.001, and all error bars are standard error of mean (SEM).

Figure 2. **The identification of Ank-1$^{(MRI61689)}$ mutation and its effects on transcription.** The

sequencing of Ank-1$^{(MRI61689)}$ mutation, showing a T to A transversion (a). Gel electrophoresis of

amplified cDNA product from wild-type, Ank-1$^{(MRI61689/+)}$ and Ank-1$^{(MRI61689/MRI61689)}$ embryonic livers

with primers that spanned the exon 17 to 21 of ankyrin-1 cDNA (primer set 1) (b). The sequencing

results of both bands, showing exon skipping in the abnormal transcript of Ank-1$^{(MRI61689/MRI61689)}$

embryonic liver (c). Gel electrophoresis of amplified cDNA product from wild-type, Ank-1$^{(MRI61689/+)}$

and Ank-1$^{(MRI61689/MRI61689)}$ embryonic livers with primers that contained the predicted 11bp insertion

(primer set 2) (d). Sequencing result showing an 11bp insertion between exon 17 and 18 cDNA of the

Ank-1$^{(MRI61689)}$ transcript (e). The predicted effect of the insertion on the translation of ankyrin-1,

showing a frameshift and a premature chain termination (f).

Figure 3. **The effect of Ank-1$^{(MRI61689)}$ mutation on the Ank-1 expression.** Quantitative PCR showing

the ankyrin-1 mRNA levels in both Ank-1$^{(MRI61689/+)}$ and Ank-1$^{(MRI61689/MRI61689)}$ embryonic liver (a). The

protein levels of various cytoskeletal proteins examined with both Coomassie (b) and Western blot

(c). The relative protein levels of ankyrin-1 calculated from the western blot (d) (n=3 per group).

Error bars indicate SEM.

Figure 4. **The response of *Ank-1*^(MRI61689/+) mice to malaria infection.** The parasite load of wild-type

and *Ank-1*^(MRI61689/+) mice when infected with 1x10^4 *P. chabaudi* (a) and the associated survival curve

(b). To determine the mechanisms of resistance, parasite intra-erythrocytic growth was assessed

through TUNEL assay at 1-10% parasitemia during late trophozoite stage, as visualised from

immunofluorescent images showing presence of parasites and TUNEL-positive parasites (c). The

number of TUNEL-positive parasites in both wild-type and *Ank-1*^(MRI61689/+) mice were counted and

expressed as a percentage (d) (n=3). For parasite invasion and RBC clearance, The IVET assay was

done showing the ratio of infected *Ank-1*^(MRI61689/+) to wild-type erythrocytes over 36 hours (e) and

the relative number *Ank-1*^(MRI61689/+) and wild-type erythrocyte in both infected and uninfected mice

(f) during *P. chabaudi* infection (n=7). ** indicates P<0.01, *** indicates P<0.001, ## and ###

indicates P<0.01 and P<0.001 respectively when compared to wild-type RBC number in infected mice,

whereas ^^ indicates P<0.01 when compared to *Ank-1*^(MRI61689/+) RBC number in uninfected mice.

Error bars indicate SEM.

Tables

	WBC (x10^6/ml)	RBC (x10^9/ml)	HGB (g/L)	MCV (fL)	MCH (pg)	MCHC (g/L)	PLT (x10^6/ml)	%Retics
Wild type	8.7±0.5	10.5±0.1	153.2±2.2	51.4±0.4	14.6±0.1	283.6±2.8	1151±57	2.69±0.26
Ank-1$^{(MRI61689/+)}$	11.9±4.5	11.1±0.1	151.2±1.4	46.1±0.2	13.5±0.1	281.4±9.3	1151±61	2.24±0.21
p-values	NS	P<0.01	NS	P<0.001	P<0.001	NS	NS	NS

668   **Table 1**. **The complete blood count of *Ank-1*$^{(MRI61689/+)}$ mice.** The haematological parameters of *Ank-*

669   *1*$^{(MRI61689/+)}$ compared to wild-type mice (n=19-25). WBC = white blood cell count; RBC = red blood cell

670   count; HGB = haemoglobin; MCV = mean corpuscular volume; MCH = mean corpuscular

671   haemoglobin; MCHC = mean corpuscular haemoglobin concentration; PLT = platelet

672   concentration; %Retics = percentage of reticulocytes.

Chromosome	Gene name	Location	Reference base	Variant base	Number of mutant mice with mutation	LOD score (Threshold: 2.08)
9	Acp5	22129643	C	A	(3/8)	-0.32
8	Kcnk1	126025024	C	T	(4/8)	0
17	Epas1	86825679	A	T	(0/8)	-2.33
7	Picalm	90165538	G	T	(0/8)	-2.33
18	Socs6	88869240	A	G	(5/8)	0.32
8	Ank1	23106019	T	A	(8/8)	2.33

675 **Table 2: The identification of MRI61689 mutation.** Six variants were selected from the exome

676 sequencing, each mutation was sequenced in *Ank-1*[(MRI61689/+)] mice and the number of mutant mice

677 carrying each mutation was determined and LOD score was calculated, with LOD threshold being

678 2.08 (n=8).

Figure 1

a)

b)

Wild-type

Ank-1(MRI61689/+)

Ank-1(MRI61689/MRI61689)

c)

d)

Figure 2

a)

T→A

Wild-type

Ank-1(MRI61689/+)

Ank-1(MRI61689/MRI61689)

b)

c)

Exon 17   Exon 18

Wild-type

Exon 17   Exon 19

Skipped

d)

e)

Wild-type

Ank-1(MRI61689)

f)

Wild-type   gca cat ctg ttt ctt ttt ag AGT GGA CTC ACT CCC
             ------------------------------ Ser   Gly   Leu Thr Pro

Ank-1(MRI61689) gca cat cag TTT CTT TTT AGA GTG GAC TCA CTC CC……TGA
                 ------------- Phe Leu Phe Arg Val Asp Ser Leu…………STOP

Figure 3

a)

Ank1 gene expression in embryonic liver

b)

c)

d)

Figure 4