1
2  TITLE
3  Containing Emerging Epidemics: a Quantitative Comparison of Quarantine and Symptom
4  Monitoring
5

6  AUTHORS
7  Corey M Peak[1], Lauren M Childs[1], Yonatan H Grad[2,3], Caroline O Buckee[1*]
8
AFFILIATIONS
1. Center for Communicable Disease Dynamics, Department of Epidemiology, Harvard T.H.
Chan School of Public Health, 677 Huntington Avenue, Boston MA 02115 USA
2. Department of Immunology and Infectious Diseases, Harvard TH Chan School of Public
Health, Boston, MA 02115
3. Division of Infectious Diseases, Brigham and Women's Hospital, Harvard Medical School,
Boston, MA 02115
16
17  *To whom correspondence should be addressed: cbuckee@hsph.harvard.edu
18

**ABSTRACT**

19
Strategies for containing an emerging infectious disease outbreak must be non-
pharmaceutical when drugs or vaccines for the pathogens do not yet exist or are
unavailable. The success of these non-pharmaceutical strategies will not only depend on
the effectiveness of quarantine or other isolation measures, but also on the epidemiological
characteristics of the infection. There is currently no systematic framework to assess the
relationship between different containment strategies and the natural history and
epidemiological dynamics of the pathogen, however. Here we compare the effectiveness of
quarantine and symptom monitoring, implemented via contact tracing, in controlling
epidemics using an agent-based branching model. We examine the relationship between
epidemic containment and the disease dynamics of symptoms and infectiousness for seven
case study diseases with diverse natural histories including Ebola, Influenza A, and Severe
Acute Respiratory Syndrome (SARS). We show that the comparative effectiveness of
symptom monitoring and quarantine depends critically on the natural history of the
infectious disease, its inherent transmissibility, and the intervention feasibility in the
particular healthcare setting. The benefit of quarantine over symptom monitoring is
generally maximized for fast-course diseases, but we show the conditions under which
symptom monitoring alone can control certain outbreaks. This quantitative framework can
guide policy-makers on how best to use non-pharmaceutical interventions to contain
emerging outbreaks and prioritize research during an outbreak of a novel pathogen.
39

## SIGNIFICANCE

Quarantine and symptom monitoring of contacts with suspected exposure to an infectious disease are key interventions for the control of emerging epidemics; however, there does not yet exist a quantitative framework for comparing the control performance of each. Here, we use a mathematical model of seven case study diseases to show how the choice of intervention is influenced by the natural history of the infectious disease, its inherent transmissibility, and the intervention feasibility in the particular healthcare setting. We use this information to identify the most important characteristics of the disease and setting that need to be characterized for an emerging pathogen in order to make an informed decision between quarantine and symptom monitoring.

## INTRODUCTION

The global burden of emerging infectious diseases is growing and prompts the need for effective containment policies [1-3]. In many cases, strategies must be non-pharmaceutical, as targeted drugs or vaccines for the pathogens are unavailable. Among the various containment strategies, isolation of ill and potentially infectious patients is one of the most intuitive, and relies on targeting individuals by tracing the contacts of known cases. Contacts with symptoms can then be hospitalized or isolated, but policy makers must also decide how best to handle contacts who do not meet the case definition for infection. Two strategies have historically been used in the case of a potentially infected but symptom-free contact: quarantine and symptom monitoring.

Quarantine of potentially infected contacts is a highly conservative approach to epidemic containment. However, there are substantial costs associated with quarantine policies, ranging from direct costs, like implementation expenses and the restriction of personal liberties, to indirect costs, including stigmatization of health workers and sometimes interruption of financial and trade markets [4-8]. A less conservative but substantially cheaper and more socially palatable approach is active symptom monitoring of contacts. In this strategy, health workers check on contacts one or two times a day, moving the contacts to isolation if symptoms occur (see definitions in *Methods*). Given the importance of rapid decision making in the event of novel emerging pathogens such as Ebola, and the potentially devastating consequences of poor containment strategies, quantitative guidelines are urgently needed.

The recent Ebola epidemic in West Africa highlights the uncertainties surrounding the relationship between non-pharmaceutical interventions and both disease biological

dynamics and the intervention setting [7]. Recommendations by the United States Centers for

Disease Control and Prevention (CDC) implicitly recognized the importance of the

implementation setting by differentiating its international response, where quarantine was

prioritized [9], and its domestic response, where symptom monitoring was prioritized [10,11].

During the Severe Acute Respiratory (SARS) epidemic, broad quarantine interventions

were applied in Taiwan and subsequently abandoned [12]. These recent epidemics exemplify

the urgent need for evidence-based guidelines on how to decide whether quarantine of an

infectious disease is, according to Gates et al, at worst "counterproductive" or at best "one

of the few tactics that can reduce its spread" [13].

The success of these approaches is not simply a reflection of the efficiency of their

implementation, however, but crucially depend on the biology and natural history of the

pathogen in question.  Disease stages are typically divided into those describing the

symptom history of the individual (e.g., the incubation period and duration of symptoms)

and the temporality of that individual's infectiousness to others (e.g., the latent period and

duration of infectiousness). Studies of epidemic dynamics typically focus solely on the

latter, but the interaction between the two has important consequences for how diseases

spread and the impact of interventions such as ring vaccination [14], travel screening [15], and

symptom monitoring. To examine the relationship between epidemiological dynamics of a

pathogen and non-pharmaceutical containment strategies, we develop an agent-based

branching model that accommodates dynamics of symptoms and infectiousness and allows

interventions to be targeted via contact tracing. To assess diseases with a wide range of

natural histories that have the potential for causing sudden, severe epidemics, we consider

case studies of seven known pathogens: Ebola; hepatitis A; influenza A; Middle East

Respiratory Syndrome (MERS); pertussis; SARS; and smallpox. We identify which disease

characteristics and intervention attributes are most critical in deciding between quarantine

and symptom monitoring, and provide a clear, general framework for understanding the

consequences of isolation policies during an epidemic of known or novel pathogens.

**RESULTS**

**Intervention Effectiveness Depends on Disease Epidemiological Dynamics.**

103           To assess the impact of quarantine and symptom monitoring, we developed a

104    general mathematical model of disease transmission and interventions targeted via contact

105    tracing (**Fig 1**). The model structure accommodates five key metrics of intervention

106    performance in a given setting (**Table 1**). We used particle filtering to parameterize seven

107    case studies of outbreak-prone pathogens (see *Methods* and **Table S1**).

108           **Fig 2** illustrates model dynamics for two infections with different epidemiological

characteristics, Ebola and influenza A, showing the range of epidemic outcomes resulting

from different interventions for a given $R_0$ value. Unimpeded exponential epidemic growth

in our branching model (red) can be reduced by the increasingly conservative

interventions of health-seeking behavior (teal), symptom monitoring (gold), and

quarantine (blue) (**Figs 2A-B**). Under a given intervention policy, we estimate the effective

reproductive number ($R_e$) as the average number of infections generated by an infectious

individual in the population. The effective reproductive number under symptom

monitoring ($R_S$), quarantine ($R_Q$), and the absolute difference between the two ($R_S - R_Q$)

increases with $R_0$ and differs by disease (**Figs 2C-D**).

118           We find the effectiveness of symptom monitoring and quarantine in controlling a

disease in a particular setting depends critically on its biological dynamics (e.g. latent and

infectious periods) and transmissibility ($R_0$) (**Fig 3a**). Holding transmissibility constant ($R_0$

arbitrarily set to 2.75±0.25), biological dynamics alone strongly influence the effectiveness

of quarantine and especially symptom monitoring, as seen by the wide spread in $R_S$ (**Fig

3b**).

In simulations with high intervention performance settings (i.e., 90% isolation

effectiveness, 90% of contacts traced within one day, and symptoms monitored daily),

diseases such as MERS and Ebola could be controlled (i.e., $R_e < 1$) with either quarantine or

symptom monitoring; diseases such as hepatitis A with only quarantine; but diseases such

as pertussis require additional interventions in order to reduce the effective reproductive

number below one, due in large part to pre-symptomatic infectiousness (**Fig 3c**). The

absolute comparative effectiveness ($R_S - R_Q$) varies widely by disease, as demonstrated by

the line length in **Fig 3c.** The relative comparative effectiveness ($\frac{R_S - R_Q}{R_S}$) also varies widely,

with quarantine reducing $R_S$ by over 65% for influenza A and hepatitis A and by less than

10% for pertussis (**Fig S1**). The reader can explore results from landscapes with different

intervention performance settings and disease transmissibility in the **interactive**

**supplement** (https://coreypeak.shinyapps.io/InteractiveQuarantine).

**Categorizing Disease Control Frontiers**

In order to compare the effectiveness of symptom monitoring and quarantine, one

must select an appropriate metric to compare $R_S$ and $R_Q$. We therefore categorized

intervention response heterogeneity into four control quadrants (**Fig 3a**). In quadrant I,

where neither intervention is sufficient to prevent epidemic growth, the relative difference

$\frac{R_S - R_Q}{R_S}$ can distinguish whether quarantine is merited or could be paired with other

strategies to achieve control.  Because quarantine is by definition the more conservative

intervention, simulation results in quadrant II occur only stochastically. In quadrant III,

where both interventions are sufficient and the number of prevented cases can be more

directly estimated, the distinguishing metric was the absolute difference $R_S - R_Q$ and its

inverse ($\frac{1}{R_S - R_Q}$), which can be interpreted as the number of contacts that must be

quarantined in order to prevent one additional case over symptom monitoring (an analog

of "number needed to treat"). For the example of SARS, Day et al. propose that mass

quarantine may be unnecessary because effective symptomatic isolation alone would

sufficiently control the disease (hence placing the disease in quadrant III) [8]. In quadrant IV,

where quarantine but not symptom monitoring can control the disease, quarantine would

be strongly considered as the minimum sufficient strategy to prevent exponential epidemic

growth.

The following two sections aim to identify which disease characteristics and

intervention performance metrics most strongly influence these differences in response to

quarantine and symptom monitoring.

**Ranking of epidemiological characteristics by importance for containment feasibility**

The comparative effectiveness of quarantine and symptom monitoring is strongly

influenced by differences in the infection's natural history. We measured partial rank

correlation coefficients to examine which biological characteristics in particular are most

influential after controlling for the other characteristics (*Methods*). As demonstrated by

strongly negative partial rank correlation coefficients in **Fig 4**, increasing the duration of

infectiousness ($d_{INF}$) and elongating the latent period ($T_{OFFSET}$) reduced the differences

between quarantine and symptom monitoring, thereby making the interventions more

similar. Other factors, such as overdispersed heterogeneity of the basic reproductive

number ($\kappa$), did not influence the average effect of symptom monitoring and quarantine, as

reflected by a coefficient of nearly zero. Longer incubation periods ($T_{INC}$) increased the

preference for quarantine, as seen by the positive partial rank correlation coefficient for

both absolute and relative comparative effectiveness. However, the length of the incubation

period does not generally influence comparative cost-effectiveness because the number of

days in quarantine ($d_Q$) increases as the incubation period lengthens (**Fig S2**).

Frequently, the most politically and economically pressing concerns are whether

control (i.e. $R_e < 1$) is logistically achievable and what would be the least invasive

intervention to achieve control. **Fig 5** shows frontiers where control of an Ebola-like

disease requires increasingly invasive interventions, moving from health-seeking behavior

(teal), to symptom monitoring (gold), to quarantine (blue), the most invasive. **Fig 5a** shows

how this frontier is influenced by the inherent transmissibility ($R_0$) and timing of the latent

period relative to the incubation period ($T_{OFFSET}$), with all other characteristics similar to

Ebola. When $R_0$ is large and symptoms emerge long after infectiousness (e.g., $T_{OFFSET} > 0$),

even quarantine is insufficient to control the disease with optimal intervention

performance. However, we observe that when transmissibility is relatively low (e.g.,

$R_0 < 2.5$), control of this hypothetical disease can be achieved even if infectiousness

precedes symptoms by several days (**Fig 5a**) or if a substantial fraction of transmission

events occur before symptom onset (adapting the framework of [16]) (**Fig 5b**).

**Ranking of intervention performance metrics by importance for containment**

**feasibility**

Policy makers facing an epidemic must also consider the expected performance of

interventions, since the effectiveness of targeted control policies will depend on their

feasibility within a particular healthcare system. Generally, we found the benefit of

quarantine over symptom monitoring increases with better intervention performance (i.e.

larger fraction of contacts traced ($P_{CT}$), better isolation effectiveness ($\gamma$), and shorter delays

in tracing a contact ($D_{CT}$) (**Fig 4**). However, the effectiveness of symptom monitoring

approached that of quarantine when the delay between symptom onset and isolation ($D_{SM}$)

is shortened, due either to more frequent symptom monitoring or more sensitive detection

of symptoms followed by prompt isolation.

While these patterns were highly consistent across the case study diseases, some

intervention performance metrics were particularly influential in the presence of certain

disease characteristics. For example, diseases with short incubation periods ($T_{INC}$) such as

influenza A were strongly influenced by delays in tracing a contact ($D_{CT}$) (**Fig S2**).

**DISCUSSION**

A key strategy to controlling the spread of infectious diseases focuses on tracing the contacts of infected individuals, with the goal of limiting subsequent spread should those contacts become infected and infectious. Here we present the first study comparing the effectiveness of the two primary non-pharmaceutical interventions: symptom monitoring and quarantine. We show that the interventions are not equivalent and that the choice of which intervention to implement to achieve optimal control depends on the natural history of the infectious disease, its inherent transmissibility, and the intervention feasibility in the particular healthcare setting.

Our results show that the benefit of quarantine over symptom monitoring is maximized for fast-course diseases (short duration of infectiousness and a short latent period compared to the incubation period), and in settings where isolation is highly effective, a large fraction of contacts are traced, or when there is a long delay between symptom onset and isolation. This delay ($D_{SM}$) not only captures ineffective symptom monitoring, but also the potential for symptoms to be masked for a period of time through biological (e.g., natural disease progression or self-medication with anti-pyretics) or behavioral (e.g., avoidance) mechanisms. In contrast, the widely-discussed "super spreading" disease characteristic did not independently impact the comparative effectiveness of interventions after holding the mean $R_0$ constant. However, this characteristic could remain important to understand disease control during early, highly stochastic stages of emergence [17]. Our findings are consistent with Fraser, Riley, et al. [16] that both inherent transmissibility and the proportion of transmission from

asymptomatically infected individuals are key epidemiological parameters for the

feasibility of control via quarantine.

In addition to identifying parameters that differentiate quarantine and symptom

monitoring, our results also characterize parameter spaces where symptom monitoring,

not just quarantine, is sufficient for containment of an emerging epidemic. Given the high

costs and poor scalability of quarantine, symptom monitoring is likely to be a key

intervention for future epidemic containment.

Although our results focus on the early stages of an outbreak, contact tracing,

symptom monitoring, and quarantine are often key tools for end-stage epidemic control

and elimination. As the effective reproductive number decreases below one (e.g. via

depletion of susceptible individuals, complementary interventions, seasonality, etc.), our

results suggest the preference for quarantine also decreases (**Fig 4**). However, one must

consider aspects such as geographic containment, public compliance, and, if the availability

of resources lags the epidemic curve, a possible resource-per-case surplus that may enable

the more conservative and costly approach of quarantine.

Our results suggest that symptom monitoring could effectively control an outbreak

of a new Ebola-like disease, even when infectiousness precedes symptoms and

interventions are not perfectly implemented. Because perfect interventions are not always

necessary, these results support the conclusion of Cetron et al. [18] that the optimal

containment strategy may allow "partial or leaky quarantine" in order to increase the

fraction of contacts who participate.

The case study diseases were selected to demonstrate a broad range of natural

histories and are not intended to represent all candidates for quarantine and symptom

monitoring. Furthermore, the selected diseases have a diverse set of transmission routes,

including bodily fluid, fecal-oral, and airborne, which will influence the contact networks

and traceability of contacts. In our framework, a reduction in the fraction of contacts who

are ultimately traced will decrease the preference for quarantine over symptom

monitoring, therefore supporting the previous findings that quarantine was inefficient for a

respiratory disease like SARS [19].

We do not consider group quarantines or other non-pharmaceutical interventions

implemented *en masse*. For such interventions as targeting all airplane passengers

returning from an affected region, the time at which symptom monitoring or quarantine

are initiated is not necessarily linked to the timing of their infector, so does not require the

correlation structure of our branching model. The effectiveness and efficiency of

quarantine, like any intervention, improves with better targeting, hence the case for mass

quarantine would require additional considerations for multisectoral cost-effectiveness [4].

We propose that the most influential parameters should be prioritized for early

characterization during an outbreak [20] and should be modeled with conservative

consideration of parameter uncertainty, including both real diversity and measurement

error. Our framework identifies the key infection-related parameters to define and can

form the basis of cost-benefit analyses. Such data-driven decision-making will be critical to

determining the optimal public health strategies for the inevitable next epidemic.

**METHODS**

**Definitions**

"Contact Tracing" is the process of identifying and assessing people who have been exposed to a disease [30]. Contacts who are symptomatic when traced are immediately placed in isolation; those who are not symptomatic are placed under either quarantine or symptom monitoring. Here, we model "forward" contact tracing whereby an infected individual names contacts they may have infected [31].

"Isolation" is the separation of a symptomatic individual believed to be infected [30]. By reducing the number of risky contact events, isolation reduces disease transmission when infectiousness coincides at least partly with symptoms.

"Quarantine" is the separation of an individual who is believed to be exposed, but is currently not ill [30]. If an individual becomes symptomatic, they will be isolated and receive healthcare.

"Symptom Monitoring" is the assessment of symptoms at regular intervals of an individual believed to be exposed, but not ill. If symptoms are detected, the individual is placed in isolation [30]. Although they may be encouraged to avoid mass-gatherings or other specific events, an individual under symptom monitoring is not separated from others and therefore does not experience a reduction in risky contacts until symptoms are detected.

"Health Seeking Behavior" is the act of seeking healthcare during the presentation of symptoms, leading to isolation. Practically, this intervention could be a health education campaign that prompts individuals to self-identify illness and seek effective isolation. This intervention, which accelerates isolation in a manner separate from contract tracing, provides a comparative care standard for our analysis.

**Model**

Individuals in our branching model progressed through a Susceptible-Exposed-
Infectious-Recovered (SEIR) disease process. We focus our analysis on the early epidemic
phase of an emerging infectious disease, assuming no changes to herd immunity within the
first few generations of transmission.

Following infection, the number of days before onset of infectiousness and onset of
symptoms are the latent period and incubation period, respectively (**Fig 1**). Because
clinical symptoms, pathogen concentration, and behavior of the patient can change
throughout the course of disease [21], we allowed relative infectiousness to vary with time $\tau$
since onset of infectiousness ($\beta_\tau$). The effective reproductive number in the presence of
health seeking behavior, symptom monitoring, and quarantine are respectively $R_{HSB}$, $R_S$, and
$R_Q$.

The recent SARS and Ebola epidemics highlighted that hospital isolation does not
always contain transmission, and we therefore allowed isolation effectiveness ($\gamma$) to vary to
reflect different settings [16,22,23]. The fraction of contacts who are traced ($P_{CT}$) can be less
than 1, encompassing symptomatic infectors who fail to recall contacts, asymptomatic
"silent" infection events, reluctance to report contacts, and challenges in identifying
contacts, especially for airborne transmission routes. Imperfections and uncertainties in
risk profiling can reduce the fraction of traced contacts that are truly infected ($P_{INF}$) [11,19].
Delays in tracing a contact ($D_{CT}$) can arise for numerous reasons, including intractable
roads, low mobile phone penetration, and personnel limitations. The delay between
symptom onset and isolation ($D_{SM}$) specifically applies to individuals under symptom

monitoring and is influenced by the frequency of monitoring, delays in recognizing

sometimes unreliable clinical features, and delays in prompt isolation upon symptom

detection.

**Simulation**

We drew disease characteristics for each simulated individual from disease-specific

input distributions. During each hour $\tau$ of infectiousness, an individual infected a number of

new individuals drawn from a Poisson distribution (or, if super-spreading factor $\kappa \neq 1$, a

negative binomial distribution [17]) with mean equal to the product of the expected number

of onward infections for the individual ($R_0$) and the relative infectiousness $\beta_\tau$ where

$\sum_{\tau=1}^{d_{INF}} \beta_\tau = 1$. We assumed time-varying relative infectiousness follows a triangular

distribution with time of peak infectiousness ($\tau_\beta$) occurring anywhere between the onset

and end of infectiousness, inclusively.

We recorded both the day of transmission and the source of each new infection for

each new transmission event, and drew disease characteristics for each newly infected

individual. An individual was identified by contact tracing with probability $P_{CT}$ at the earlier

time of either when their infector is isolated or the time of the transmission event if

infection occurred while the infector was isolated. After an operational lag time of $D_{CT}$ days,

a contact was placed under quarantine, symptom monitoring or, if already symptomatic,

isolation. An individual in isolation or quarantine had their infectiousness reduced by a

factor γ for the remainder of their disease. An individual under symptom monitoring was

isolated $D_{SM}$ days after symptom onset. A full description of the model process can be found

in **S1 Appendix**.

**Parameterization**

As compared to characteristics related to the natural history of symptoms and illness, key aspects of the natural history of infectiousness tend to be harder to observe and measure [24]. Therefore, we used a Sequential Monte Carlo particle filtering algorithm [25,26] to create a joint probability space of the time offset between the latent period and incubation period ($T_{OFFSET} = T_{LAT} - T_{INC}$), time of peak infectiousness ($\tau_\beta$), and duration of infectiousness ($d_{INF}$). From an uninformative prior distribution of each parameter bounded by published observations, we simulated five infection generations of 500 initial individuals and recorded the simulated serial interval (i.e., the time between symptom onset in infector-infectee pairs). Parameter sets were resampled with importance weights determined by the degree to which the distribution of simulated serial intervals matched published serial interval distributions, using the Kolmogorov-Smirnov test of the difference between cumulative distribution functions (**Table S1**) [27,28]. After perturbation, the process was repeated until convergence, which we defined to be when the median Kolmogorov-Smirnov statistic was within 10% of the previous two iterations. This cutoff criterion was chosen to balance the objectives of finding a stationary posterior set of particles while preserving some of the heterogeneity in input parameters.

Holding the incubation period distribution constant, we fit an offset for the latent period ($T_{OFFSET}$) for several reasons, including consistency with CDC methods for disease characterization [29], the biological expectation of these timings both being linked to pathogen load, and to parsimoniously limit each characteristic to one interpretable parameter. For the duration of infectiousness ($d_{INF}$), we fit the upper bound of a uniform

distribution with a lower bound of 1 day. To allow for variable infectiousness during this

duration, we assume a triangular distribution of relative infectiousness $\beta_\tau$ and fit the time

of peak infectiousness ($\tau_\beta$). A full description of the model parameterization can be found

in the **S2 Appendix.**

**Analysis**

Partial rank correlation coefficients were calculated to identify the most influential

disease characteristics (e.g. duration of infectiousness) and intervention performance

metrics (e.g. isolation effectiveness). In order to remove dependence between the

parameters jointly fit through the particle filtering method above, we used Latin Hypercube

Sampling to draw 5,000 sets from each marginal posterior parameter distribution

independently. To maximize coverage of the parameter space we allowed fractional

parameters ($\gamma$, $P_{CT}$, $P_{INF}$, $k$) to range from 0 to 1, delays ($D_{CT}$, $D_{SM}$) to range from 0 to 7 days,

$R_0$ to range from 1 to 5, and the incubation period ($T_{INC}$) to be shrunk by up to 50% or

stretched by up to 150%.

Using 1,000 samples drawn from the joint-parameter space from the particle

filtering method, we measured $R_0$, $R_Q$, $R_S$, and $R_{HSB}$ for each disease. We compared the

effectiveness of symptom monitoring and quarantine by the absolute difference $R_S - R_Q$

and the relative difference $\frac{R_S - R_Q}{R_S}$. We calculated the number of days an infected individual

was in quarantine but not yet infectious ($d_Q$) as surrogate for the marginal cost of

quarantine over symptom monitoring. As surrogates for cost-effectiveness, we calculate the

absolute difference per quarantine day $(R_S - R_Q)/d_Q$ and relative difference per

quarantine day $\left(\frac{R_S - R_Q}{R_S}\right)/d_Q$ and present these results in Fig S1 and S2.

When risk profiling was imperfect (i.e. $P_{INF}$ < 1), individuals who were not infected

may have been mistakenly traced as contacts and placed under symptom monitoring or

quarantine. Such events may be conceptualized as false positives and will decrease $P_{INF}$;

conversely, individuals who were infected but not traced are false negatives and will

decrease $P_{CT}$. We assumed that non-infected contacts were followed for a length of time set

up the 95th percentile incubation period ($T_{INC}^{95}$), at which point health authorities may

conclude the contact was not infected after all. This would change the number of days in

quarantine to $\widehat{d_Q} = \left(d_Q + T_{inc}^{95}\left(1/P_{inf} - 1\right)\right)$.

**ACKNOWLEDGEMENTS**

CMP, LMC, and COB were supported by Cooperative Agreement U54GM088558 from the

National Institute Of General Medical Sciences. CMP was also supported by National

Research Service Award T32AI007535-16A1. YHG was supported by the National Institutes

of Health (K08-AI104767) and the Smith Family Foundation.  The content is solely the

responsibility of the authors and does not necessarily represent the 433 official views of

the National Institute Of General Medical Sciences or the National Institutes of 434 Health.

The funders had no role in study design, data collection and analysis, decision to publish, or

preparation of the manuscript. The authors thank Colin Worby for helpful statistical

discussions.

**REFERENCES**

McMichael AJ. Environmental and social influences on emerging infectious diseases: past, present and future. *Philos Trans R Soc B Biol Sci* 2004; **359**: 1049–58.

Morens DM, Folkers GK, Fauci AS. The challenge of emerging and re-emerging infectious diseases. *Nature* 2004; **430**.

Jones KE, Patel NG, Levy MA. Global trends in emerging infectious diseases. *Nature* 2008; **451**: 990–3.

Barbera J, Macintyre A, Gostin L, *et al.* Large-Scale Quarantine Following Biological Terrorism in the United States. *Jama* 2001; **286**: 2711–7.

Union ACL, Partnership YG health J. Fear, Politics, and Ebola. How Quarantines Hurt the Fight Against Ebola and Violate the Constitution. 2015 http://media.wix.com/ugd/148599_3c2a77dfa3c84942be9b40efacda1876.pdf.

Ebola: Quarantine can undermine efforts to curb epidemic. http://www.msf.org/article/ebola-quarantine-can-undermine-efforts-curb-epidemic.

Drazen JM, Kanapathipillai R, Campion EW, *et al.* Ebola and Quarantine. *N Engl J Med* 2014; **371**: 2029–30.

Day T, Park A, Madras N, Gumel A, Wu J. When Is Quarantine a Useful Control Strategy for Emerging Infectious Diseases? *Am J Epidemiol* 2006; **163**: 479–85.

Nyenswah T, Fallah M, Sieh S, *et al.* Controlling the Last Known Cluster of Ebola Virus Disease — Liberia , January – February 2015. 2015; **64**: 500–4.

CDC. Interim U.S. Guidance for Monitoring and Movement of Persons with Potential Ebola Virus Exposure. Atlanta, Georgia, 2014 http://www.cdc.gov/vhf/ebola/exposure/monitoring-and-movement-of-persons-with-exposure.html.

Chung WM, Smith JC, Weil LM, *et al.* Active Tracing and Monitoring of Contacts Associated With the First Cluster of Ebola in the United States. *Ann Intern Med* 2015; : 1–10.

Rothstein MA, Alcalde MG, Elster NR, *et al.* Quarantine and isolation: Lessons learned from SARS. 2003 http://www.iaclea.org/members/pdfs/SARS REPORT.Rothstein.pdf.

Gates B. The Next Epidemic — Lessons from Ebola. 2015; : 1–4.

Kretzschmar M, van den Hof S, Wallinga J, van Wijngaarden J. Ring vaccination and smallpox control. *Emerg Infect Dis* 2004; **10**: 832–41.

Gostic KM, Kucharski AJ, Lloyd-Smith JO. Effectiveness of traveller screening for emerging pathogens is shaped by epidemiology and natural history of infection. *Elife* 2015; **2015**: 1–16.

Fraser C, Riley S, Anderson RM, Ferguson NM. Factors that make an infectious disease outbreak controllable. *Proc Natl Acad Sci U S A* 2004; **101**: 6146–51.

Lloyd-Smith JO, Schreiber SJ, Kopp PE, Getz WM. Superspreading and the effect of individual variation on disease emergence. *Nature* 2005; **438**: 355–9.

Cetron M, Maloney S, Koppaka R, Simone P. Isolation and Quarantine: Containment Strategies for Sars 2003. *Learn from SARS Prep Next Dis Outbreak Work Summ* 2004; : 71–83.

Glasser JW, Hupert N, McCauley MM, Hatchett R. Modeling and public health

emergency responses: Lessons from SARS. *Epidemics* 2011; **3**: 32–7.

20    Lessler J, Edmunds WJ, Halloran ME, Hollingsworth TD, Lloyd AL. Seven challenges
for model-driven data collection in experimental and observational studies.
*Epidemics* 2015; **10**: 78–82.

21    Fine P, Eames K, Heymann DL. 'Herd immunity': A rough guide. *Clin Infect Dis* 2011;
**52**: 911–6.

22    Klinkenberg D, Fraser C, Heesterbeek H. The effectiveness of contact tracing in
emerging epidemics. *PLoS One* 2006; **1**: e12.

23    Lloyd-Smith JO, Galvani AP, Getz WM. Curtailing transmission of severe acute
respiratory syndrome within a community and its hospital. *Proc Biol Sci* 2003; **270**:
1979–89.

24    Richardson M, Elliman D, Maguire H, Simpson J, Nicoll A. Evidence base of incubation
periods, periods of infectiousness and exclusion policies for the control of
communicable diseases in schools and preschools. *Pediatr Infect Dis J* 2001; **20**: 380–
91.

25    Andrieu C, Doucet A, Singh SS, Tadic VB. Particle methods for change detection,
system identification, and control. *Proc IEEE* 2004; **92**: 423–38.

26    Doucet A, de Freitas N, Gordon N. An Introduction to Sequential Monte Carlo
Methods. *Seq Monte Carlo Methods Pract* 2001; : 3–14.

27    Djuric PM, Miguez J. Assessment of Nonlinear Dynamic Models by Kolmogorov-
Smirnov Statistics. *IEEE Trans Signal Process* 2010; **58**: 5069–79.

28    Marsaglia G, Tsang WW, Wang J. Evaluating Kolmogorov's Distribution. *J Stat Softw*
2003; **8**: 1–4.

29    Centers for Disease Control and Prevention. Epidemiology and Prevention of Vaccine
Preventable Diseases, 13th ed. Washington D.C.: Public Health Foundation, 2015.

30    Centers for Disease Control and Prevention. CDC Methods for Implementing and
Managing Contact Tracing for Ebola Virus Disease in Less-Affected Countries. 2014.

31    Muller J, Kretzschmar M, Dietz K. Contact tracing in stochastic and deterministic
epidemic models. *Math Biosci* 2000; **164**: 39–64.

32    Wu JT, Riley S, Fraser C, Leung GM. Reducing the Impact of the Next Influenza
Pandemic Using Household-Based Public Health Interventions. *PLoS Med* 2006; **3**.
DOI:10.1371/journal.pmed.0030361.

33    WHO Ebola Response Team. Ebola Virus Disease in West Africa - The First 9 Months
of the Epidemic and Forward Projections. SUPPLEMENT. *N Engl J Med* 2014;
published online Sept 22. DOI:10.1056/NEJMoa1411100.

34    Althaus CL. Ebola superspreading. *Lancet Infect Dis* 2015; **15**: 507–8.

35    Vink MA, Bootsma MCJ, Wallinga J. Serial Intervals of Respiratory Infectious Diseases:
A Systematic Review and Analysis. *Am J Epidemiol* 2014; **180**: 865–75.

36    Brodribb HS. Infective hepatitis in a boarding-school. *Lancet* 1952; : 339–42.

37    Pickles W. Epidemic Catarrhal Jaundice. *Br Med J* 1930; : 944–6.

38    Biggerstaff M, Cauchemez S, Reed C, Gambhir M, Finelli L. Estimates of the
reproduction number for seasonal, pandemic, and zoonotic influenza: a systematic
review of the literature. *BMC Infect Dis* 2014; **14**: 480.

39    Viboud C, Boëlle P, Cauchemez S, *et al.* Risk factors of influenza transmission in
households. 2004; : 684–9.

40    Lessler J, Reich NG, Brookmeyer R, Perl TM, Nelson KE, Cummings DA. Incubation

periods of acute respiratory viral infections: a systematic review. *Lancet Infect Dis*
2009; **9**: 291–300.
41 Cauchemez S, Fraser C, Van Kerkhove MD, *et al.* Middle East respiratory syndrome
coronavirus: Quantification of the extent of the epidemic, surveillance biases, and
transmissibility. *Lancet Infect Dis* 2014; **14**: 50–6.
42 Assiri A, McGeer A, Perl TM, *et al.* Hospital outbreak of Middle East respiratory
syndrome coronavirus. *N Engl J Med* 2013; **369**: 407–16.
43 te Beest DE, Henderson D, van der Maas N a T, *et al.* Estimation of the serial interval
of pertussis in Dutch households. *Epidemics* 2014; **7**: 1–6.
44 Lipsitch M, Cohen T, Cooper B, *et al.* Transmission dynamics and control of severe
acute respiratory syndrome. *Science (80- )* 2003; **300**: 1966–70.
45 Nishiura H, Brockmann SO, Eichner M. Extracting key information from historical
data to quantify the transmission dynamics of smallpox. *Theor Biol Med Model* 2008;
**5**: 20.
509
510

511
**Fig 1. Schematic of the natural history of disease and the timing of interventions.**

Beginning on the left with the infection event, one progress through a latent period ($T_{LAT}$)

before becoming infectious for $d_{INF}$ days with late peak infectiousness $\tau_\beta$. For diseases A, B,

and C, symptoms are respectively shown to emerge before, concurrent with, and after

onset of infectiousness. We show here an individual who is traced shortly after infection

and is placed under symptom monitoring or quarantine after a short delay $D_{CT}$.

**Fig 2. Model dynamics and output for two exemplar diseases: Ebola and influenza A.**

Each line in the panels (A) and (B) designates one model run initiated with 100 infectious

individuals in Generation 1 and submitted to either no intervention (red), health seeking

behavior (teal), symptom monitoring every day (gold), or quarantine (blue) at Generation

3. Each point in panels C and D designates the simulated effective reproductive number

from one model run with input reproductive number (x-axis) between 1 and 5, with the

526    asterisk denoting the input R for panels (A) and (B) (1.83 and 1.46, respectively). Loess

527    curves are shown as heavier lines. Here, symptom monitoring performs similarly to

528    quarantine for Ebola control, but not for influenza A. Note the independent y-axes.

**531** **Fig 3. Infection control performance depends on disease biological dynamics and**

**532** **inherent transmissibility (R$_0$).**

**533** (A) The effective reproductive number under symptom monitoring (x-axis) and quarantine

**534** (y-axis) for 100 simulations of each disease when the basic reproductive number is set to

**535** published values [See Panel C, diamonds]. Quadrants indicate regions of control with (I)

**536** neither quarantine nor symptom monitoring, (II) only symptom monitoring, (III) either

**537** quarantine or symptom monitoring, and (IV) only quarantine. (B) As in (A), but the basic

**538** reproductive number ($R_0$) is set to for all diseases to 2.75 (+/- 0.25) to isolate inherent

**539** differences in biological dynamics. (C) Disease-specific mean basic reproductive number

**540** (diamond) and the mean effective reproductive numbers under symptom monitoring

541    (triangle) and quarantine (circle). The length of the horizontal line therefore equals the

542    absolute comparative effectiveness $R_S - R_Q$.

**Fig 4. Influence of disease characteristics and intervention performance metrics.**

Partial rank correlation coefficients (x-axis) measuring the influence of disease

characteristics and intervention performance metric (rows) on the absolute (red) and

relative (green) comparative effectiveness of quarantine and symptom monitoring, pooled

for all case study diseases. The 95% confidence intervals from 100 bootstrapped samples

are represented by error bars.

**Fig 5. Minimally invasive interventions sufficient to control a hypothetical disease.**

(A) Disease characteristics drawn from Ebola except symptoms are assumed to either:

precede infectiousness by up to 10 days (X = -10 days); coincide with infectiousness onset

(X = 0 days); or emerge up to 10 days after infectiousness onset  (X = +10 days). Points

represent simulations where health-seeking behavior (teal), symptom monitoring (gold),

or quarantine (blue) were the minimally sufficient intervention to bring $R_e$ below 1. (B) As

in (A), but the x-axis is transformed to represent the proportion of infections that occur

prior to symptoms in a analogous way to Fraser, Riley, et al 2004 [16]. Interventions are in an

optimal setting with all contacts being traced immediately, no infections occur during

isolation, and symptom monitoring is performed twice per day.

564
**Fig S1. Relative comparative effectiveness and cost-effectiveness.** The relative

comparative effectiveness varies widely by disease, with quarantine reducing $R_S$ by >65%

for influenza A and hepatitis A and by <10% for pertussis. However, due to a much shorter

incubation period of influenza A versus hepatitis A (**Table S1**), the relative cost-

effectiveness measured by the reduction per day of quarantine (outlined bars) is

substantially higher for influenza A than hepatitis A.

**Fig S2. Partial rank correlation coefficients for all outcomes.** Partial rank correlation

coefficients (x-axis) measuring the influence of disease characteristics and intervention

performance metrics (rows) on the impact, comparative effectiveness, and comparative

cost-effectiveness of the interventions under study. Disease-specific estimates are shown

with colored bars and pooled estimates with larger grey bars. For example, increasing the

delay in tracing a named contact $D_{CT}$ has a generally small effect negative effect on $R_S$-$R_Q$

when pooled across diseases (large grey bar), but for influenza A specifically (purple bar),

$D_{CT}$ has a rather large negative effect on $R_S$-$R_Q$. Note that pooled estimates for comparative

cost-effectiveness are not available due to non-monotonic relationships across diseases.

**Table S1.** Disease parameters

| | Inputs | | | Parameters fit via Sequential Monte-Carlo method | | |
|---|---|---|---|---|---|---|
| | Basic Reproductive Number $R_0$ | Serial Interval (days) | Incubation Period $T_{INC}$ (days) | Latent period offset $T_{OFFSET}$ $= T_{LAT} - T_{INC}$ (days) | Mean duration of infectiousness $(1 + d_{INF})/2$ (days) | Time of peak Infectiousness $\tau_\beta$ |
| Ebola | 1.83 (1.72, 1.94) [33] | 13.36 (2.66, 38.8) [34] | 7.87 (0.93, 28.2) [33] | 0.33 (0**, 1.01) | 6.53 (1.28, 13.7) | 0.10 (0, 0.37) |
| Hepatitis A | 2.25 (2, 2.5) * | 26.72 (20.7, 33.8) [35,36] | 29.11 (24.6, 34.1) [37] | -5.33 (-7.57, -3.26) | 6.23 (1.22, 15.8) | 0.35 (0, 0.98) |
| Influenza A | 1.54 (1.28, 1.80) [38] | 2.20 (0.63, 3.76) [35,39] | 1.40 (0.63, 3.10) [40] | -0.23 (-0.76, 0.29) | 1.88 (1.04, 3.84) | 0.49 (0.02, 0.98) |
| MERS | 0.95 (0.6, 1.3) [41] | 7.62 (2.48, 23.3) [42] | 5.20 (1.83, 14.7) [42] | -1.55 (-3.14, 0.02) | 8.35 (1.37, 19.9) | 0.37 (0.01 , 0.96) |
| Pertussis | 4.75 (4.5, 5) * | 19.26 (3.61, 57.2) [43] | 7.00 (4.00, 10.0) [29] | -2.14 (-5.39, 0.78) | 34.38 (2.67, 76.7) | 0.45 (0.11, 0.88) |
| SARS | 2.9 (2.2, 3.6) [44] | 8.32 (1.59, 19.2) [44] | 4.01 (1.25, 12.8) [40] | 0.16 (0**, 0.67) | 10.94 (1.50, 23.0) | 0.10 (0, 0.46) |
| Smallpox | 4.75 (4.5, 5) * | 15.54 (9.98, 24.2) [45] | 11.83 (8.47, 16.5) [45] | 0.03 (-1.80, 1.68) | 8.45 (1.37, 20.0) | 0.32 (0, 0.97) |

Median (95% CI)

* Assumed

** Sequential Monte-Carlo boundary condition reached

**S1 Appendix. Disease model**

The model simulates a branching network of infected individuals only. An individual
$i$ is assigned characteristics sampled from distributions defined for each disease. The
incubation period ($T_{INC}$), i.e. the time from infection to symptom onset, is drawn from
published distributions (**Table S1**). The duration of infectiousness ($d_{INF}$), time of peak
infectiousness ($\tau_\beta$), and time offset between the latent and incubation periods ($T_{OFFSET}$) are
drawn from the joint posterior distribution generated by the sequential Monte-Carlo (SMC)
particle filtering method described in **S2 Appendix**. For clarity, we describe the method for
an individual $i$, but the following process is repeated for an initial population of 1,000
individuals who initiate distinct trees.
The expected number of onward infections by $i$, $R_{0i}$, is distributed over each hour $\tau$
of disease $R_{\tau i} = \beta_{\tau i} R_{0i}$, where $\beta_{\tau i}$ is the relative infectiousness of $i$ on hour $\tau$ such that
$\sum_{\tau=0}^{d_{INF}} \beta_\tau = 1$. For parsimony and ease of interpretation, we assume $\beta_\tau$ follows a triangle
distribution with a peak value at time $\tau_\beta$ drawn from the SMC posterior.

$$\beta_{\tau i} = \begin{cases} \sqrt{\tilde{u}\, d_{INF}\, \tau_\beta} & , \ if \ \tilde{u} < {}^{\tau_\beta}/d_{INF} \\ d_{INF} - \sqrt{(1-\tilde{u})(d_{INF} - \tau_\beta)(d_{INF})} & , \ otherwise \end{cases}$$

where $\tilde{u} \sim unif(0,1)$
The number of infections ($N_{\tau i}$) generated by individual $i$ at hour $\tau$ is drawn from a
negative binomial distribution with mean equal to $R_{\tau i}$ and dispersion factor $\kappa$ to capture
super spreading tendencies. If $\kappa = 1$, the negative binomial distribution reduces to a
Poisson distribution with rate $\lambda = R_{\tau i}$.
Each individual $i$ then has a vector of $\langle N_{0i}, N_{1i}, \dots, N_{d_{inf_i}} \rangle$ of onward infections that
occur during each hour $\tau \epsilon \langle 0,1, \dots, d_{INF_i} \rangle$. A new individual $j$ is generated for each onward
infection $N_{t_i} \geq 1$. Disease characteristics for $j$ are drawn as above, adding the time of
infection to the latent and incubation periods for $j$.
Contact $j$ of infector $i$ will be traced with probability $P_{CT}$. If traced, $j$ is placed under
symptom monitoring or quarantine with an operational lag time of $D_{CT}$ days. The lag time
occurs after the earlier of isolation or removal from the disease system upon recovery or
death for individual $i$. We assume individuals granted access to the quarantine or isolation
room will be logged and given the same attention as contacts traced through
epidemiological interview and will therefore will begin monitoring or quarantine at the
time of the infection control breach or transmission event.
Next we determine the time of isolation for $j$. If time of symptom onset for $j$ occurs
before $j$ is traced, $j$ is immediately isolated (**Panel 1**). Otherwise, time of isolation for $j$
depends on whether symptom monitoring or quarantine is used. Under symptom
monitoring, isolation of $j$ occurs a delay $D_{SM}$ days after symptom onset. Note that for
contacts checked twice-daily, $D_{SM} \sim unif(0, 0.5)$. Upon isolation, the hourly expected number
of onward infections is reduced to $R_{t_j} = (1 - \gamma)\beta_{t_j} R_{0j}$ where $\gamma$ is effectiveness of isolation
with support $[0, 1]$. If $j$ is under quarantine, then $R_{t_j}$ is reduced by $(1 - \gamma)$ beginning at the
time $j$ is traced.

## S2 Appendix. Parameterization via Sequential Monte Carlo

The following parameterization method was repeated for each case study disease. Data-informed incubation period and serial interval distributions were collected through a literature review. Sequential Monte-Carlo (also known as particle filtering) methods were used to estimate the joint distribution of three disease parameters ($T_{OFFSET}$, $d_{INF}$, and $\tau_\beta$) using knowledge of the incubation period and serial interval distributions [28,29].

Here we assume that the latent period for an individual is some time ($T_{OFFSET}$) before ($T_{OFFSET} < 0$) or after ($T_{OFFSET} > 0$) the onset of symptoms. Therefore, $T_{OFFSET}$ is a translation of the incubation period distribution. We assume a uniform distribution of duration of infectiousness from 1 day to $d_{INF}$, allowing for partial days as well. We assume the distribution of relative infectiousness to follow a triangle distribution with a max at time $\tau_\beta$, which ranges from 0, indicating infectiousness is linearly decreasing, to 1, indicating infectiousness is linearly increasing.

The steps are as follows:

    i.    Draw initial parameter set Θ consisting of $T_{OFFSET}$, $d_{INF}$, and $\tau_\beta$ from a Latin Hypercube sample bounded by the range of parameter values found in the literature (Note: $d_{INF}$ is loosely bounded by symptom observations).

    ii.    Run the branching epidemic model under a situation with no interventions and measure the simulated serial intervals.

    iii.    Compare the simulated serial intervals to the published serial interval(s) in the literature using the Kolmogorov-Smirnov test statistic (KS).

    iv.    Draw a bootstrapped sample of Θ_candidate from Θ weighted by the KS statistic and restricted by an adaptive Θ threshold of (max KS in trial)*0.80

    v.    Perturb each Θ_candidate by up to 2% of the initial parameter value range

    vi.    Repeat steps (ii)-(v) until the median KS is within 10% of each of the previous two rounds.

    vii.    Generate an unweighted sample from final Θ joint probability space with replacement as the input parameter set for th1e case studies.

    viii.    To calculate partial rank correlation coefficients generate an unweighted sample with replacement from the parameter space of each of for $T_{OFFSET}$, $d_{INF}$, and $\tau_\beta$ independent distributions from Θ.