1   TITLE
2   Containing Emerging Epidemics: a Quantitative Comparison of Quarantine and Symptom
3   Monitoring
4

5   AUTHORS
6   Corey M Peak[1], Lauren M Childs[1], Yonatan H Grad[2,3], Caroline O Buckee[1*]
7
AFFILIATIONS
1. Center for Communicable Disease Dynamics, Department of Epidemiology, Harvard T.H.
Chan School of Public Health, 677 Huntington Avenue, Boston MA 02115 USA
2. Department of Immunology and Infectious Diseases, Harvard TH Chan School of Public
Health, Boston, MA 02115
3. Division of Infectious Diseases, Brigham and Women's Hospital, Harvard Medical School,
Boston, MA 02115
15
16   *To whom correspondence should be addressed: cbuckee@hsph.harvard.edu

18   ABSTRACT
19   Background
20   Strategies for containing an emerging infectious disease outbreak must be non-
21   pharmaceutical when drugs or vaccines for the pathogens do not yet exist or are
22   unavailable. However, little work exists to guide decisions between competing non-
23   pharmaceutical strategies, as exemplified by the confusion about whether to employ
24   quarantine or symptom monitoring during the recent Ebola epidemic in West Africa.
25

26   Methods
We compared the effectiveness of quarantine and symptom monitoring in controlling
epidemics using an agent-based branching model that accommodates non-pharmaceutical
interventions. We used Sequential Monte Carlo particle filtering methods to parameterize
disease dynamics of symptoms and infectiousness for seven case study diseases with
diverse natural histories including Ebola, Influenza A, and Severe Acute Respiratory
Syndrome (SARS). We quantify the key characteristics of an emerging disease that are most
influential for determining the optimal intervention, given varying feasibility of its
implementation.
35

36   Findings
The comparative effectiveness of symptom monitoring and quarantine depends critically
on the natural history of the infectious disease, its inherent transmissibility, and the
intervention feasibility in the particular healthcare setting. The benefit of quarantine over
symptom monitoring is generally maximized for fast-course diseases, or when there is a
long delay between symptom onset and isolation. We find that symptom monitoring could
effectively control an outbreak of a new Ebola-like disease, even when infectiousness
precedes symptoms and interventions are not perfectly implemented.
44

45   Interpretation
46   We establish a quantitative framework for guiding policy-makers in their decisions about
47   how best to use non-pharmacological interventions to contain emerging outbreaks. Our
48   method also provides guidelines for prioritizing research during an outbreak of a novel
49   pathogen, by highlighting which aspects of the disease determine the epidemic potential of
50   emerging pathogens.
51

52   Funding
53   NIH-MIDAS, Smith Family Foundation.
54

INTRODUCTION

The global burden of emerging infectious diseases is growing and prompts the need

for effective containment policies[1–3]. In many cases, strategies must be non-pharmaceutical,

as targeted drugs or vaccines for the pathogens are unavailable. Among the various

containment strategies, isolation of ill and potentially infectious patients is one of the most

obvious, and relies on targeting individuals by tracing the contacts of those infected.

Contacts with symptoms can then be hospitalized or isolated, but policy makers must also

decide how best to handle contacts who do not meet the case definition for infection. Two

strategies have historically been used in the case of a potentially infected but healthy

contact: quarantine and symptom monitoring. Precise definitions of these interventions

can be found in **Panel 1**.

The recent Ebola epidemic in West Africa highlighted the confusion about whether

quarantine or symptom monitoring should be employed and under what circumstances [4].

For example, the US Centers for Disease Control and Prevention (CDC) recommendations

differed between its international response, where quarantine was prioritized [5], and its

domestic response, where symptom monitoring was prioritized [6,7]. Similar confusion was

recorded during the Severe Acute Respiratory (SARS) epidemic, where broad quarantine

interventions were applied in Taiwan and subsequently abandoned [8]. These recent

epidemics highlight the urgent need for evidence-based guidelines on how to decide

whether quarantine of an infectious disease is, according to Gates et al, at worst

"counterproductive" or at best "one of the few tactics that can reduce its spread" [9].

Quarantine of potentially infected contacts is a highly conservative approach to

epidemic containment. However, there are substantial costs associated with quarantine

policies, ranging from direct costs, like implementation expenses and the restriction of

personal liberties, to indirect costs, including stigmatization of health workers and

sometimes interruption of financial and trade markets [4,10–13]. A less conservative but

substantially cheaper and more socially palatable approach is active symptom monitoring

of contacts. In this strategy, health workers check on contacts one or two times a day,

moving the contacts to isolation if symptoms occur (see **Panel 1**). Given the importance of

rapid decision making in the event of novel emerging pathogens such as Ebola, and the

potentially devastating consequences of poor containment strategies, quantitative

guidelines are urgently needed.

Here, we develop a mathematical model to compare the performance of symptom

monitoring and quarantine of traced contacts in containing an emerging infectious disease.

We consider case studies of seven known pathogens with a wide range of natural histories

that have the potential for causing sudden, severe epidemics. We use these case studies to

examine the impact of isolation policies for diseases with diverse known epidemiological

characteristics, and to provide a generalizable approach to decision making in the event of

future epidemics. We identify which disease characteristics and intervention attributes are

most critical in deciding between quarantine and symptom monitoring, and provide a clear,

general framework for understanding the consequences of isolation policies during an

epidemic.

METHODS

Model Structure

We developed an agent-based branching model, which accommodates dynamics of

symptoms and infectiousness. Specifically, we considered the epidemiological

characteristics of infections including Ebola, hepatitis A, influenza A, Middle East

Respiratory Syndrome (MERS), pertussis, SARS, and smallpox. Individuals in our branching

model progressed through a Susceptible-Exposed-Infectious-Recovered (SEIR) disease

process. We focused our analysis on the early epidemic phase of an emerging infectious

disease, assuming no changes to herd immunity within the first few generations of

transmission.

Model Inputs

Following infection, the number of days before onset of infectiousness and onset of

symptoms are the latent period and incubation period, respectively (**Fig 1**). Because

clinical symptoms, pathogen concentration, and behavior of the patient can change

throughout the course of disease [20], we allowed relative infectiousness to vary with time τ

since onset of infectiousness (β_τ). The basic reproductive number (R_0) is defined as the

average number of infections caused by one infectious individual who is not isolated over

the course of disease in a susceptible population. To incorporate the idea that symptomatic

individuals are likely to seek care, regardless of contact tracing policies in place, we

included a "health seeking behavior" (HSB) intervention condition (Panel 1), where

individuals were assumed to seek care at a random time while symptomatic. The effective

reproductive number in the presence of health seeking behavior, symptom monitoring, and

quarantine are respectively R_{HSB}, R_S, and R_Q.

The definitions of the five key intervention policy performance metrics are found in

Table 1. The recent SARS and Ebola epidemics highlighted that hospital isolation does not

always contain transmission, and we therefore allowed isolation effectiveness (γ) to vary to

reflect different settings [21-23]. The fraction of contacts traced (P_{CT}) can be less than 1,

encompassing symptomatic infectors who fail to recall contacts, asymptomatic "silent"

infection events, or reluctance to report contacts. Imperfections in risk profiling can reduce

the fraction of traced contacts that are truly infected (P_{INF}). For example, only 2 Ebola

infections were recorded among 179 contacts traced in the United States [7]. Delays in

tracing a contact (D_{CT}) can arise for numerous reasons, including intractable roads, low

mobile phone penetration, and personnel limitations. The delay between symptom onset

and isolation (D_{SM}) applies only to individuals under symptom monitoring and is influenced

by the frequency of monitoring and delays in prompt isolation upon symptom detection.

Simulation

We drew disease characteristics for each simulated individual from disease-specific

input distributions. During each hour τ of infectiousness, an individual infected a number of

new individuals drawn from a Poisson distribution (or, if super-spreading factor $\kappa \neq 1$, a

negative binomial distribution[24]) with mean equal to the product of the expected number

of onward infections for the individual (R_0) and the relative infectiousness β_τ where

$\sum_{\tau=1}^{d_{INF}} \beta_\tau = 1$. We assumed time-varying relative infectiousness follows a triangular

distribution with time of peak infectiousness (τ_β) occurring anywhere between the onset

and end of infectiousness, inclusively.

We recorded both the day of transmission and the infector for each new

transmission event, and drew disease characteristics for each newly infected individual. An

individual was identified by contact tracing with probability P_{CT} at the earliest time of the

following: (a) their infector was isolated; (b) their infector recovered/died from disease; or

(c) at the time of infection if the infector was isolated when the transmission event

occurred. After an operational lag time of D_{CT} days, a contact was placed under quarantine,

symptom monitoring or, if already symptomatic, isolation. An individual in isolation or

quarantine had their infectiousness reduced by a factor γ for the remainder of their

disease. An individual under symptom monitoring was isolated D_{SM} days after symptom

onset. A full description of the model process can be found in **S1 Appendix**. **Fig 2** shows

sample outputs for five simulated generations of transmission.

Parameterization

There is a general lack of published data on key characteristics including latent

period, relative infectiousness, and duration of infectiousness, while some other

characteristics, including the incubation period and serial interval, are generally estimable

[25]. Therefore, we used a Sequential Monte Carlo particle filtering algorithm [26,27] to create a

joint probability space of the time offset between the latent period and incubation period

($T_{OFFSET} = T_{LAT} - T_{INC}$), time of peak infectiousness (τ_β), and duration of infectiousness

(d_{INF}). From an uninformative prior distribution of each parameter bounded by published

observations, we simulated five infection generations of 500 initial individuals and

recorded the simulated serial interval (i.e., the time between symptom onset in infector-

infectee pairs). Parameter sets were resampled with importance weights determined by

the degree to which the distribution of simulated serial intervals matched published serial

interval distributions, using the Kolmogorov-Smirnov test of the difference between

cumulative distribution functions (**Table 2**) [28,29]. After perturbation, the process was

repeated until convergence, which we defined to be when the median Kolmogorov-Smirnov

statistic was within 10% of the previous two iterations. This cutoff criterion was chosen to

balance the objectives of finding a stationary posterior set of particles while preserving

some of the heterogeneity in input parameters.

Holding the incubation period distribution constant, we fit an offset for the latent

period (T_{OFFSET}) for several reasons, including consistency with CDC methods for disease

characterization [30], the biological expectation of these timings both being linked to

pathogen load, and to parsimoniously limit each characteristic to one interpretable

parameter. For the duration of infectiousness (d_{INF}), we fit the upper bound of a uniform

distribution with a lower bound of 1 day. To allow for variable infectiousness during this

duration, we assume a triangular distribution of relative infectiousness β_τ and fit the time

of peak infectiousness (τ_β). A full description of the model parameterization can be found

in the **S2 Appendix.**

Analysis

Partial rank correlation coefficients were calculated to identify the most influential

disease characteristics (e.g. duration of infectiousness) and intervention performance

metrics (e.g. isolation effectiveness). In order to remove dependence between the

parameters jointly fit through the particle filtering method above, we used Latin Hypercube

Sampling to draw 5,000 sets from each marginal posterior parameter distribution

independently. To maximize coverage of the parameter space we allowed fractional

parameters (γ, P_{CT}, P_{INF}, k) to range from 0 to 1, delays (D_{CT}, D_{SM}) to range from 0 to 7 days,

R_0 to range from 1 to 5, and the incubation period (T_{INC}) to be shrunk by up to 50% or

stretched by up to 150%.

193   Using 1,000 samples drawn from the joint-parameter space from the particle

filtering method, we measured R_0, R_Q, R_S, and R_{HSB} for each disease. We compared the

effectiveness of symptom monitoring and quarantine by the absolute difference $R_S - R_Q$

and the relative difference $\frac{R_S - R_Q}{R_S}$. We calculated the number of days an infected individual

was in quarantine but not yet infectious (d_Q) as surrogate for the marginal cost of

quarantine over symptom monitoring. As surrogates for cost-effectiveness, we calculate the

absolute difference per quarantine day $\left(R_S - R_Q\right)/d_Q$ and relative difference per

quarantine day $\left(\frac{R_S - R_Q}{R_S}\right)/d_Q$.

201   When risk profiling was imperfect (i.e. P_{INF} < 1), individuals who were not infected

may have been mistakenly traced as contacts and placed under symptom monitoring or

quarantine. We assumed that non-infected contacts were followed for a length of time set

up the 95th percentile incubation period (T_{INC}^{95}), at which point health authorities may

conclude the contact was not infected after all. This would change the number of days in

quarantine to $\widehat{d_Q} = \left(d_Q + T_{inc}^{95}\left(1/P_{inf} - 1\right)\right)$.

Role of the Funding Source

RESULTS

The effectiveness of symptom monitoring and quarantine in controlling a disease in

a particular setting depends critically on its transmissibility (R_0) and on its biological

dynamics (e.g. latent and infectious periods) (**Fig 3a**). By holding transmissibility constant

(R_0 arbitrarily set to 2·75±0·25), **Fig 3b** shows how biological dynamics alone strongly

influence the effectiveness of quarantine and especially symptom monitoring, as seen by

the wide spread in R_S.

When intervention performance is "high" (see **Table 1**), diseases such as MERS and

Ebola can be controlled with either quarantine or symptom monitoring; diseases such as

hepatitis A with only quarantine; and diseases such as pertussis require additional

interventions, e.g. vaccination (**Fig 3c**). The absolute comparative effectiveness ($R_S - R_Q$)

varies widely by disease, as demonstrated by the line length in **Fig 3c.** The relative

comparative effectiveness ($\frac{R_S - R_Q}{R_S}$) also varies widely, with quarantine reducing R$_S$ by over

65% for influenza A and hepatitis A and by less than 10% for pertussis (**Fig S1**). The reader

can explore results from landscapes with different intervention performance settings and

disease transmissibility in the **interactive supplement**

(https://coreypeak.shinyapps.io/InteractiveQuarantine).

When choosing between symptom monitoring and quarantine, one must consider

the effective reproductive numbers under each intervention and choose an appropriate

metric to compare these two values. We categorized intervention response heterogeneity

into four control quadrants (**Fig 3a**). In quadrant I, where neither intervention is sufficient

to prevent epidemic growth, the relative difference $\frac{R_S - R_Q}{R_S}$ can distinguish whether

quarantine is merited or could be paired with other strategies to achieve control. Because

quarantine is by definition the more conservative intervention, simulation results in

quadrant II occur only stochastically. In quadrant III, where both interventions are

sufficient and the number of prevented cases can be more directly estimated, the

distinguishing metric was the absolute difference $R_S - R_Q$ and its inverse ($\frac{1}{R_S - R_Q}$), which

can be interpreted as the number of contacts that must be quarantined in order to prevent

one additional case over symptom monitoring (an analog of "number needed to treat"). For

the example of SARS, Day et al. propose that mass quarantine may be unnecessary because

effective symptomatic isolation alone would sufficiently control the disease (hence placing

the disease in quadrant III) [13]. In quadrant IV, where quarantine but not symptom

monitoring can control the disease, quarantine would be strongly considered as the

minimum sufficient strategy to prevent exponential epidemic growth.

The following two sections aim to identify which disease characteristics and

intervention performance metrics most strongly influence these differences in response to

quarantine and symptom monitoring.

Influential Disease Characteristics

The comparative effectiveness of quarantine and symptom monitoring is strongly

influenced by differences in the components of the infection's natural history. We examined

which biological characteristics in particular are most influential. As demonstrated by

strongly negative partial rank correlation coefficients in **Fig 4**, increasing the duration of

infectiousness (d_{INF}) and elongating the latent period (T_{OFFSET}) reduced the differences

between quarantine and symptom monitoring, thereby making the interventions more

similar. Other factors, such as overdispersed heterogeneity of the basic reproductive

number (κ), did not influence the average effect of symptom monitoring and quarantine, as

reflected by a partial rank correlation coefficient of nearly zero. Longer incubation periods

(T_{INC}) increased the preference for quarantine, as seen by the positive partial rank

correlation coefficient for both absolute and relative comparative effectiveness. However,

the length of the incubation period does not generally influence comparative cost-

effectiveness because the number of days in quarantine (d_Q) increases as the incubation

period lengthens (**Fig S2**).

Frequently, the most pressing concerns are whether control (i.e. $R_e < 1$) is

achievable and what would be the least invasive intervention to achieve control. **Fig 5**

shows frontiers where control of an Ebola-like disease requires increasingly invasive

interventions, namely health-seeking behavior (teal), symptom monitoring (gold), or

quarantine (blue). **Fig 5a** shows how this frontier is influenced by the inherent

transmissibility (R_0) and timing of the latent period relative to the incubation period

(T_{OFFSET}), with all other characteristics similar to Ebola. When R_0 is large and symptoms

emerge long after infectiousness (e.g., $T_{OFFSET} > 0$), even quarantine is insufficient to

control the disease with optimal intervention performance [see **Table 1**]. However, we

observe that when transmissibility is relatively low (e.g., $R_0 < 2 \cdot 5$), control of this

hypothetical disease can be achieved even if infectiousness precedes symptoms by several

days (**Fig 5a**) or if a substantial fraction of transmission events occur before symptom

onset (adapting the framework of [22]) (**Fig 5b**).

Intervention Performance Metrics

Policy makers facing an epidemic must also consider the expected performance of

interventions, since the effectiveness of targeted control policies will depend on their

feasibility within a particular healthcare system. Generally, we found the benefit of

quarantine over symptom monitoring increases with better intervention performance (i.e.

larger fraction of contacts traced (P_{CT}), better isolation effectiveness (γ), and shorter delays

in tracing a contact (D_{CT}) (**Fig 4**). However, the effectiveness of symptom monitoring

approached that of quarantine when the delay between symptom onset and isolation (D_{SM})

is shortened, due either to more frequent symptom monitoring or more sensitive detection

of symptoms followed by prompt isolation.

While these patterns were highly consistent across the case study diseases, some

intervention performance metrics were particularly influential in the presence of certain

disease characteristics. For example, diseases with short incubation periods (T_{INC}) such as

influenza A were strongly influenced by delays in tracing a contact (D_{CT}) (**Fig S2**).

DISCUSSION

A key strategy to controlling the spread of infectious diseases focuses on tracing the

contacts of infected individuals, with the goal of limiting subsequent spread should those

contacts become infected and infectious. Here we present the first study comparing the

effectiveness of the two primary non-pharmacological interventions for a directly-

transmitted infection, symptom monitoring and quarantine. We show that the

interventions are not equivalent and that the choice of which intervention to implement to

achieve optimal control depends on the natural history of the infectious disease, its

inherent transmissibility, and the intervention feasibility in the particular healthcare

setting.

Our results show that the benefit of quarantine over symptom monitoring is

maximized for fast-course diseases (short duration of infectiousness and a short latent

period compared to the incubation period), and in settings where isolation is highly

effective, a large fraction of contacts are traced, or when there is a long delay between

symptom onset and isolation. This delay (D_{SM}) not only captures ineffective symptom

monitoring, but also the potential for symptoms to be masked for a period of time through

biological (e.g., natural disease progression or self-medication with anti-pyretics) or

behavioral (e.g., avoidance) mechanisms. In contrast, the widely-discussed "super

spreading" disease characteristic did not impact the comparative effectiveness of

interventions, although this characteristic could remain important to understand disease

control during early, highly stochastic stages of emergence [24]. Our findings are consistent

with Fraser, Riley, et al. [22] that both inherent transmissibility and the proportion of

transmission from asymptomatically infected individuals are key epidemiological

parameters for the feasibility of control via quarantine.

Our results uniquely identify parameter spaces where symptom monitoring, not just

quarantine, is sufficient for containment of an emerging epidemic. Given the high costs and

poor scalability of quarantine, symptom monitoring is likely to be a key intervention for

future epidemic containment. Our results suggest that symptom monitoring could

effectively control an outbreak of a new Ebola-like disease, even when infectiousness

precedes symptoms and interventions are not perfectly implemented. Because perfect

interventions are not always necessary, these results support the conclusion of Cetron et al.

[31] that the optimal containment strategy may allow "partial or leaky quarantine" in order to

increase the fraction of contacts who participate.

We do not consider mass quarantines or other non-pharmaceutical interventions

done on a population level. For mass interventions, such as targeting all airplane

passengers returning from an affected region, the time at which symptom monitoring or

quarantine are initiated is not necessarily linked to the timing of their infector, so does not

require the correlation structure of our branching model. The effectiveness and efficiency

of quarantine, like any intervention, improves with better targeting, hence the case for

mass quarantine would require additional considerations for multisectoral cost-

effectiveness [10].

We propose that the most influential parameters should be prioritized for early

characterization during an outbreak and should be modeled with conservative

consideration of parameter uncertainty, including both real diversity and measurement

error. Our framework identifies the key infection-related parameters to define and can

form the basis of cost-benefit analyses. Such data-driven decision-making will be critical to

determining the optimal public health strategies for the inevitable next epidemic.

ACKNOWLEDGEMENTS
343
344   The authors thank Colin Worby for helpful statistical discussions. CMP, LMC, and COB were

345   supported by Cooperative Agreement U54GM088558 from the National Institute Of

346   General Medical Sciences. CMP was also supported by National Research Service Award

347   T32AI007535-16A1. YHG was supported by the National Institutes of Health (K08-

348   AI104767) and the Smith Family Foundation.

Panel 1: Definitions of interventions

Contact Tracing is the process of identifying and assessing people who have been exposed to a disease [33]. Contacts who are symptomatic when traced are immediately placed in **isolation**; those who are not symptomatic are placed under either **quarantine** or **symptom monitoring**. Here, we model "forward" contact tracing whereby an infected individual names contacts they may have infected [34].

Isolation is the separation of a *symptomatic* individual believed to be infected [33]. By reducing the number of risky contact events, isolation reduces disease transmission when infectiousness coincides at least partly with symptoms.

Quarantine is the separation of an individual who is believed to be exposed, but is currently not ill [33]. As soon as an individual is placed in quarantine, we assume they have the same reduction in risky contacts as in isolation. If an individual becomes symptomatic, they will be isolated and receive healthcare.

Symptom Monitoring is the assessment of symptoms at regular intervals of an individual believed to be exposed, but not ill. If symptoms are detected, the individual is placed in isolation [33]. Although they may be encouraged to avoid mass-gatherings or other specific events, an individual under symptom monitoring is not separated from others and therefore does not experience a reduction in risky contacts until symptoms are detected.

Health Seeking Behavior is the act of seeking healthcare during the presentation of symptoms, leading to isolation. Practically, this intervention could be a communications campaign that prompts individuals to self-identify their illness and seek effective isolation. This intervention, which accelerates isolation in a manner separate from contract tracing, provides a comparative care standard for our analysis.

Panel 2. Research in context

Evidence before this study
We searched PubMed and Google Scholar on July 17, 2016, with the terms "quarantine" and "symptom monitoring" and "contact tracing". We did not find any studies that compare the impact on disease spread of quarantine and symptom monitoring of contacts. However, previous work has demonstrated how disease characteristics can strongly influence the effectiveness of non-pharmaceutical interventions such as isolation and quarantine [13,22,35] and traveler screening [36].

Added value of this study
We provide a formal mathematical framework for comparing the relative merits of quarantine and symptom monitoring targeted by contact tracing. We use this framework and a range of case study diseases to show how the natural history of the infectious disease, its inherent transmissibility, and the intervention feasibility in the particular healthcare setting all influence the comparative effectiveness of quarantine and symptom monitoring.

Implications of all available evidence
In certain circumstances, quarantine and/or symptom monitoring can be powerful non-pharmaceutical interventions for epidemic control. When choosing between these interventions, some aspects of the disease and setting are more influential than others and should therefore be prioritized when studying an emerging disease and modeled with a range of uncertainty for re-emerging diseases.

404 **Table 1.** Intervention Parameters

	Variable	Definition	Example Intervention Performance	
			Optimal	High
Isolation Effectiveness	γ	Proportion of infections prevented by isolation as compared to an individual never in isolation (R_0)	1	0·9
Fraction of contacts traced	P_{CT}	Proportion of infected contacts identified by infector	1	0·9
Fraction of traced contacts who are truly infected	P_{INF}	Proportion of individuals under quarantine or symptom monitoring who were truly infected before they were traced	1	0·5
Delay in tracing a named contact	D_{CT}	Number of days between nomination of a contact and monitoring or quarantine	0·25 days ± 0·25	0·5 days ± 0·5
Delay from symptom onset to isolation	D_{SM}	Number of days between symptom onset and isolation of an individual being symptom monitored	0.25 days ± 0.25	0.5 days ± 0.5

406   **Table 2.** Disease parameters

	Inputs			Parameters fit via Sequential Monte-Carlo method		
	Basic Reproductive Number R_0	Serial Interval (days)	Incubation Period T_{INC} (days)	Latent period offset T_{OFFSET} $= T_{LAT} - T_{INC}$ (days)	Mean duration of infectiousness $(1 + d_{INF})/2$ (days)	Time of peak Infectiousness τ_β
Ebola	1·83 (1·72, 1·94) [37]	13·36 (2·66, 38·8) [38]	7·87 (0·93, 28·2) [37]	0·33 (0**, 1·01)	6·53 (1·28, 13·7)	0·10 (0, 0·37)
Hepatitis A	2·25 (2, 2·5) *	26·72 (20·7, 33·8) [39,40]	29·11 (24·6, 34·1) [41]	-5·33 (-7·57, -3·26)	6·23 (1·22, 15·8)	0·35 (0, 0·98)
Influenza A	1·54 (1·28, 1·80) [42]	2·20 (0·63, 3·76) [39,43]	1·40 (0·63, 3·10) [44]	-0·23 (-0·76, 0·29)	1·88 (1·04, 3·84)	0·49 (0·02, 0·98)
MERS	0·95 (0·6, 1·3) [45]	7·62 (2·48, 23·3) [46]	5·20 (1·83, 14·7) [46]	-1·55 (-3·14, 0·02)	8·35 (1·37, 19·9)	0·37 (0·01 , 0·96)
Pertussis	4·75 (4·5, 5) *	19·26 (3·61, 57·2) [47]	7·00 (4·00, 10·0) [30]	-2·14 (-5·39, 0·78)	34·38 (2·67, 76·7)	0·45 (0·11, 0·88)
SARS	2·9 (2·2, 3·6) [15]	8·32 (1·59, 19·2) [15]	4·01 (1·25, 12·8) [44]	0·16 (0**, 0·67)	10·94 (1·50, 23·0)	0·10 (0, 0·46)
Smallpox	4·75 (4·5, 5) *	15·54 (9·98, 24·2) [48]	11.83 (8.47, 16.5) [48]	0.03 (-1.80, 1.68)	8.45 (1.37, 20.0)	0.32 (0, 0·97)

407   Median (95% CI)

408   * Assumed

409   ** Sequential Monte-Carlo boundary condition reached

Fig 1. Schematic of the natural history of disease and the timing of interventions.

Beginning on the left with the infection event, one progress through a latent period (T_{LAT})

before becoming infectious for d_{INF} days with a varying degree of infectiousness β_τ. For

diseases A, B, and C, symptoms are respectively shown to emerge before, concurrent with,

and after onset of infectiousness. We show here an individual who is traced shortly after

infection and is placed under symptom monitoring or quarantine after a short delay D_{CT}.

Fig 2. Model dynamics and output for two exemplar diseases: Ebola and Influenza A.

Each line in the panels (A) and (B) designates one model run initiated with 100 infectious

individuals in Generation 1 and submitted to either no intervention (red), health seeking

behavior (teal), symptom monitoring every day (gold), or quarantine (blue) at Generation

3. Each point in panels C and D designates the simulated effective reproductive number

from one model run with input reproductive number (x-axis) between 1 and 5, with the

asterisk denoting the input R for panels (A) and (B) (1·83 and 1·46, respectively). Loess

curves are shown as heavier lines. Here, symptom monitoring performs similarly to

quarantine for Ebola control, but not for influenza A. Note the independent y-axes.

429
430

431
Fig 3. Infection control performance depends on disease biological dynamics and

inherent transmissibility (R$_0$).

(A) The effective reproductive number under symptom monitoring (x-axis) and quarantine

(y-axis) for 100 simulations of each disease when the basic reproductive number is set to

published values [See Panel C, diamonds]. Quadrants indicate regions of control with (I)

neither quarantine nor symptom monitoring, (II) only symptom monitoring, (III) either

quarantine or symptom monitoring, and (IV) only quarantine. (B) As in (A), but the basic

reproductive number (R_0) is set to for all diseases to 2·75 (+/- 0·25) to isolate inherent

differences in biological dynamics. (C) Disease-specific mean basic reproductive number

(diamond) and the mean effective reproductive numbers under symptom monitoring

(triangle) and quarantine (circle). The length of the horizontal line therefore equals the

absolute comparative effectiveness $R_S - R_Q$.

444
Fig 4. Influence of disease characteristics and intervention performance metrics.

446 Partial rank correlation coefficients (x-axis) measuring the influence of disease

447 characteristics and intervention performance metric (rows) on the absolute (red) and

448 relative (green) comparative effectiveness of quarantine and symptom monitoring, pooled

449 for all case study diseases. The 95% confidence intervals from 100 bootstrapped samples

450 are represented by error bars.

451
452

Fig 5. Minimally invasive interventions sufficient to control a hypothetical disease.

(A) Disease characteristics drawn from Ebola except symptoms are assumed to either:

precede infectiousness by up to 10 days (X = -10 days); coincide with infectiousness onset

(X = 0 days); or emerge up to 10 days after infectiousness onset (X = +10 days). Points

represent simulations where health-seeking behavior (teal), symptom monitoring (gold),

or quarantine (blue) were the minimally sufficient intervention to bring R_e below 1. (B) As

in (A), but the x-axis is transformed to represent the proportion of infections that occur

prior to symptoms in a analogous way to Fraser, Riley, et al 2004 [22]. Interventions are in

the "optimal" setting with all contacts being traced immediately, no infections occur during

isolation, and symptom monitoring is performed twice per day.

465   REFERENCES

467   1   McMichael AJ. Environmental and social influences on emerging infectious diseases:
468   past, present and future. *Philos Trans R Soc B Biol Sci* 2004; **359**: 1049–58.

469   2   Morens DM, Folkers GK, Fauci AS. The challenge of emerging and re-emerging
470   infectious diseases. *Nature* 2004; **430**.

471   3   Jones KE, Patel NG, Levy MA. Global trends in emerging infectious diseases. *Nature*
472   2008; **451**: 990–3.

473   4   Drazen JM, Kanapathipillai R, Campion EW, *et al.* Ebola and Quarantine. *N Engl J Med*
474   2014; **371**: 2029–30.

475   5   Nyenswah T, Fallah M, Sieh S, *et al.* Controlling the Last Known Cluster of Ebola Virus
476   Disease — Liberia , January – February 2015. 2015; **64**: 500–4.

477   6   CDC. Interim U.S. Guidance for Monitoring and Movement of Persons with Potential
478   Ebola Virus Exposure. Atlanta, Georgia, 2014
479   http://www.cdc.gov/vhf/ebola/exposure/monitoring-and-movement-of-persons-
480   with-exposure.html.

481   7   Chung WM, Smith JC, Weil LM, *et al.* Active Tracing and Monitoring of Contacts
482   Associated With the First Cluster of Ebola in the United States. *Ann Intern Med* 2015; :
483   1–10.

484   8   Rothstein MA, Alcalde MG, Elster NR, *et al.* Quarantine and isolation: Lessons learned
485   from SARS. 2003 http://www.iaclea.org/members/pdfs/SARS
486   REPORT.Rothstein.pdf.

487   9   Gates B. The Next Epidemic — Lessons from Ebola. 2015; : 1–4.

488   10   Barbera J, Macintyre A, Gostin L, *et al.* Large-Scale Quarantine Following Biological
489   Terrorism in the United States. *Jama* 2001; **286**: 2711–7.

490   11   Union ACL, Partnership YG health J. Fear, Politics, and Ebola. How Quarantines Hurt
491   the Fight Against Ebola and Violate the Constitution. 2015
492   http://media.wix.com/ugd/148599_3c2a77dfa3c84942be9b40efacda1876.pdf.

493   12   Ebola: Quarantine can undermine efforts to curb epidemic.
494   http://www.msf.org/article/ebola-quarantine-can-undermine-efforts-curb-
495   epidemic.

496   13   Day T, Park A, Madras N, Gumel A, Wu J. When Is Quarantine a Useful Control
497   Strategy for Emerging Infectious Diseases? *Am J Epidemiol* 2006; **163**: 479–85.

498   14   Legrand J, Grais RF, Boelle PY, Valleron a J, Flahault a. Understanding the dynamics
499   of Ebola epidemics. *Epidemiol Infect* 2007; **135**: 610–21.

500   15   Lipsitch M, Cohen T, Cooper B, *et al.* Transmission dynamics and control of severe
501   acute respiratory syndrome. *Science (80-)* 2003; **300**: 1966–70.

502   16   Chowell G, Blumberg S, Simonsen L, Miller M a., Viboud C. Synthesizing data and
503   models for the spread of MERS-CoV, 2013: Key role of index cases and hospital
504   transmission. *Epidemics* 2014; **9**: 40–51.

505   17   Kretzschmar M, van den Hof S, Wallinga J, van Wijngaarden J. Ring vaccination and
506   smallpox control. *Emerg Infect Dis* 2004; **10**: 832–41.

507   18   Bauch CT, Rao ASRS, Pham BZ, *et al.* A dynamic model for assessing universal
508   Hepatitis A vaccination in Canada. *Vaccine* 2007; **25**: 1719–26.

509   19   Boelle P-Y, Ansart S, Cori A, Valleron A-J. Transmission parameters of the A/H1N1
510   (2009) influenza virus pandemic: a review. *Influenza Other Respi Viruses* 2011; **5**:

306–16.

20   Fine P, Eames K, Heymann DL. 'Herd immunity': A rough guide. *Clin Infect Dis* 2011;
52: 911–6.

21   Klinkenberg D, Fraser C, Heesterbeek H. The effectiveness of contact tracing in
emerging epidemics. *PLoS One* 2006; **1**: e12.

22   Fraser C, Riley S, Anderson RM, Ferguson NM. Factors that make an infectious
disease outbreak controllable. *Proc Natl Acad Sci U S A* 2004; **101**: 6146–51.

23   Lloyd-Smith JO, Galvani AP, Getz WM. Curtailing transmission of severe acute
respiratory syndrome within a community and its hospital. *Proc Biol Sci* 2003; **270**:
1979–89.

24   Lloyd-Smith JO, Schreiber SJ, Kopp PE, Getz WM. Superspreading and the effect of
individual variation on disease emergence. *Nature* 2005; **438**: 355–9.

25   Richardson M, Elliman D, Maguire H, Simpson J, Nicoll A. Evidence base of incubation
periods, periods of infectiousness and exclusion policies for the control of
communicable diseases in schools and preschools. *Pediatr Infect Dis J* 2001; **20**: 380–
91.

26   Andrieu C, Doucet A, Singh SS, Tadic VB. Particle methods for change detection,
system identification, and control. *Proc IEEE* 2004; **92**: 423–38.

27   Doucet A, de Freitas N, Gordon N. An Introduction to Sequential Monte Carlo
Methods. *Seq Monte Carlo Methods Pract* 2001; : 3–14.

28   Djuric PM, Miguez J. Assessment of Nonlinear Dynamic Models by Kolmogorov-
Smirnov Statistics. *IEEE Trans Signal Process* 2010; **58**: 5069–79.

29   Marsaglia G, Tsang WW, Wang J. Evaluating Kolmogorov's Distribution. *J Stat Softw*
2003; **8**: 1–4.

30   Centers for Disease Control and Prevention. Epidemiology and Prevention of Vaccine
Preventable Diseases, 13th ed. Washington D.C.: Public Health Foundation, 2015.

31   Cetron M, Maloney S, Koppaka R, Simone P. Isolation and Quarantine: Containment
Strategies for Sars 2003. *Learn from SARS Prep Next Dis Outbreak Work Summ* 2004; :
71–83.

32   Ferguson NM, Cummings D a T, Cauchemez S, *et al.* Strategies for containing an
emerging influenza pandemic in Southeast Asia. *Nature* 2005; **437**: 209–14.

33   Centers for Disease Control and Prevention. CDC Methods for Implementing and
Managing Contact Tracing for Ebola Virus Disease in Less-Affected Countries. 2014.

34   Muller J, Kretzschmar M, Dietz K. Contact tracing in stochastic and deterministic
epidemic models. *Math Biosci* 2000; **164**: 39–64.

35   Wu JT, Riley S, Fraser C, Leung GM. Reducing the Impact of the Next Influenza
Pandemic Using Household-Based Public Health Interventions. 2006; **3**.
DOI:10.1371/journal.pmed.0030361.

36   Gostic KM, Kucharski AJ, Lloyd-Smith JO. Effectiveness of traveller screening for
emerging pathogens is shaped by epidemiology and natural history of infection. *Elife*
2015; **2015**: 1–16.

37   WHO Ebola Response Team. Ebola Virus Disease in West Africa - The First 9 Months
of the Epidemic and Forward Projections. SUPPLEMENT. *N Engl J Med* 2014;
published online Sept 22. DOI:10.1056/NEJMoa1411100.

38   Althaus CL. Ebola superspreading. *Lancet Infect Dis* 2015; **15**: 507–8.

39   Vink MA, Bootsma MCJ, Wallinga J. Serial Intervals of Respiratory Infectious Diseases:

A Systematic Review and Analysis. *Am J Epidemiol* 2014; **180**: 865–75.

40   Brodribb HS. Infective hepatitis in a boarding-school. *Lancet* 1952; : 339–42.

41   Pickles W. Epidemic Catarrhal Jaundice. *Br Med J* 1930; : 944–6.

42   Biggerstaff M, Cauchemez S, Reed C, Gambhir M, Finelli L. Estimates of the
reproduction number for seasonal, pandemic, and zoonotic influenza: a systematic
review of the literature. *BMC Infect Dis* 2014; **14**: 480.

43   Viboud C, Boëlle P, Cauchemez S, *et al.* Risk factors of influenza transmission in
households. 2004; : 684–9.

44   Lessler J, Reich NG, Brookmeyer R, Perl TM, Nelson KE, Cummings DA. Incubation
periods of acute respiratory viral infections: a systematic review. *Lancet Infect Dis*
2009; **9**: 291–300.

45   Cauchemez S, Fraser C, Van Kerkhove MD, *et al.* Middle East respiratory syndrome
coronavirus: Quantification of the extent of the epidemic, surveillance biases, and
transmissibility. *Lancet Infect Dis* 2014; **14**: 50–6.

46   Assiri A, McGeer A, Perl TM, *et al.* Hospital outbreak of Middle East respiratory
syndrome coronavirus. *N Engl J Med* 2013; **369**: 407–16.

47   te Beest DE, Henderson D, van der Maas N a T, *et al.* Estimation of the serial interval
of pertussis in Dutch households. *Epidemics* 2014; **7**: 1–6.

48   Nishiura H, Brockmann SO, Eichner M. Extracting key information from historical
data to quantify the transmission dynamics of smallpox. *Theor Biol Med Model* 2008;
5: 20.

581
582   **Fig S1. Relative comparative effectiveness and cost-effectiveness.** The relative

583   comparative effectiveness varies widely by disease, with quarantine reducing R_S by >65%

584   for influenza A and hepatitis A and by <10% for pertussis. However, due to a much shorter

585   incubation period of influenza A versus hepatitis A (**Table 2**), the relative cost-

586   effectiveness (outlined bars) is substantially higher for influenza A than hepatitis A.

587
588

589
Fig S2. Partial rank correlation coefficients for all outcomes. Partial rank correlation

coefficients (x-axis) measuring the influence of disease characteristics and intervention

performance metrics (rows) on the impact, comparative effectiveness, and comparative

cost-effectiveness of the interventions under study. Disease-specific estimates are shown

with colored bars and pooled estimates with grey bars. Note that pooled estimates for

comparative cost-effectiveness are not available due to non-monotonic relationships across

diseases.

S1 Appendix. Disease model

The model simulates a branching network of infected individuals only. An individual i is assigned characteristics sampled from distributions defined for each disease. The incubation period (T_{INC}), i.e. the time from infection to symptom onset, is drawn from published distributions (**Table 2**). The duration of infectiousness (d_{INF}), time of peak infectiousness (τ_β), and time offset between the latent and incubation periods (T_{OFFSET}) are drawn from the joint posterior distribution generated by the sequential Monte-Carlo (SMC) particle filtering method described in **S2 Appendix**. For clarity, we describe the method for an individual i, but the following process is repeated for an initial population of 1,000 individuals who initiate distinct trees.

The expected number of onward infections by i, R_{0_i}, is distributed over each hour τ of disease $R_{\tau_i} = \beta_{\tau_i} R_{0_i}$, where β_{τ_i} is the relative infectiousness of i on hour τ such that $\sum_{\tau=0}^{d_{INF}} \beta_\tau = 1$. For parsimony and ease of interpretation, we assume β_τ follows a triangle distribution with a peak value at time τ_β drawn from the SMC posterior.

$$\beta_{\tau_i} = \begin{cases} \sqrt{\tilde{u}\, d_{INF}\, \tau_\beta} & , \; if \; \tilde{u} < {}^{\tau_\beta}\!/_{d_{INF}} \\ d_{INF} - \sqrt{(1-\tilde{u})(d_{INF} - \tau_\beta)(d_{INF})} & , \; otherwise \end{cases}$$

where $\tilde{u} \sim unif(0,1)$

The number of infections (N_{τ_i}) generated by individual i at hour τ is drawn from a negative binomial distribution with mean equal to R_{τ_i} and dispersion factor κ to capture super spreading tendencies. If $\kappa = 1$, the negative binomial distribution reduces to a Poisson distribution with rate $\lambda = R_{\tau_i}$.

Each individual i then has a vector of $\langle N_{0_i}, N_{1_i}, \dots, N_{d_{inf_i}} \rangle$ of onward infections that occur during each hour $\tau \, \epsilon \langle 0,1, \dots, d_{INF_i} \rangle$. A new individual j is generated for each onward infection $N_{t_i} \geq 1$. Disease characteristics for j are drawn as above, adding the time of infection to the latent and incubation periods for j.

Contact j of infector i will be traced with probability P_{CT}. If traced, j is placed under symptom monitoring or quarantine with an operational lag time of D_{CT} days. The lag time occurs after the earlier of isolation or removal from the disease system upon recovery or death for individual i. We assume all contacts of an individual in hospital isolation are documented, so an individual j who was infected by an isolated individual i will be traced from the time of the transmission event.

Next we determine the time of isolation for j. If time of symptom onset for j occurs before j is traced, j is immediately isolated (**Panel 1**). Otherwise, time of isolation for j depends on whether symptom monitoring or quarantine is used. Under symptom monitoring, isolation of j occurs a delay D_{SM} days after symptom onset. Note that for contacts checked twice-daily, $D_{SM} \sim unif(0, 0.5)$. Upon isolation, the hourly expected number of onward infections is reduced to $R_{t_j} = (1 - \gamma)\beta_{t_j} R_{0_j}$ where γ is effectiveness of isolation with support [0, 1]. If j is under quarantine, then R_{t_j} is reduced by $(1 - \gamma)$ beginning at the time j is traced.

S2 Appendix. Parameterization via Sequential Monte Carlo

638
639
The following parameterization method was repeated for each case study disease.
Data-informed incubation period and serial interval distributions were collected through a
literature review. Sequential Monte-Carlo (also known as particle filtering) methods were
used to estimate the joint distribution of three disease parameters (T_{OFFSET}, d_{INF}, and τ_β)
using knowledge of the incubation period and serial interval distributions [28,29].
Here we assume that the latent period for an individual is some time (T_{OFFSET})
before ($T_{OFFSET} < 0$) or after ($T_{OFFSET} > 0$) the onset of symptoms. Therefore, T_{OFFSET} is a
translation of the incubation period distribution. We assume a uniform distribution of
duration of infectiousness from 1 day to d_{INF}, allowing for partial days as well. We assume
the distribution of relative infectiousness to follow a triangle distribution with a max at
time τ_β, which ranges from 0, indicating infectiousness is linearly decreasing, to 1,
indicating infectiousness is linearly increasing.
The steps are as follows:
653

    i.   Draw initial parameter set Θ consisting of T_{OFFSET}, d_{INF}, and τ_β from a
654
Latin Hypercube sample bounded by the range of parameter values
found in the literature (Note: d_{INF} is loosely bounded by symptom
observations).
ii.   Run the branching epidemic model under a situation with no
interventions and measure the simulated serial intervals.
iii. Compare the simulated serial intervals to the published serial
interval(s) in the literature using the Kolmogorov-Smirnov test statistic
(KS).
iv.   Draw a bootstrapped sample of Θ$_\text{candidate}$ from Θ weighted by the KS
statistic and restricted by an adaptive Θ threshold of (max KS in
trial)*0.80
v.   Perturb each Θ$_\text{candidate}$ by up to 2% of the initial parameter value range
vi.   Repeat steps (ii)-(v) until the median KS is within 10% of each of the
previous two rounds.
vii. Generate an unweighted sample from final Θ joint probability space
with replacement as the input parameter set for the case studies.
viii. To calculate partial rank correlation coefficients generate an
unweighted sample with replacement from the parameter space of each
of for T_{OFFSET}, d_{INF}, and τ_β independent distributions from Θ.
674
675