Activity of the *C. elegans* egg-laying behavior circuit is controlled by competing

activation and feedback inhibition

Kevin M. Collins[1,2,6], Addys Bode[1], Robert W. Fernandez[2], Jessica E. Tanis[2,4], Jacob Brewer[2],

Matt Creamer[3,5], and Michael R. Koelle[2,3]

[1] Department of Biology, University of Miami, Coral Gables, FL USA

[2] Department of Molecular Biophysics and Biochemistry, Yale University, New Haven, CT USA

[3] Interdepartmental Neuroscience Program, Yale University, New Haven, CT USA

[4] Present address: Department of Biological Sciences, University of Delaware, Newark, DE

USA

[5] Present address: Department of Molecular, Cellular, and Developmental Biology, Yale

University, New Haven, CT USA

[6]To whom correspondence should be addressed:

kcollins@bio.miami.edu

Department of Biology

University of Miami

1301 Memorial Drive

Coral Gables, FL 33143

Tel: (305) 284-9058

Abstract

Like many behaviors, *Caenorhabditis elegans* egg laying alternates between inactive and active states. To understand how the underlying neural circuit turns the behavior on and off, we optically recorded circuit activity in behaving animals while manipulating circuit function using mutations, optogenetics, and drugs. In the active state, the circuit shows rhythmic activity phased with the body bends of locomotion. The serotonergic HSN command neurons initiate the active state, but accumulation of unlaid eggs also promotes the active state independent of the HSNs. The cholinergic VC motor neurons slow locomotion during egg-laying muscle contraction and egg release. The uv1 neuroendocrine cells mechanically sense passage of eggs through the vulva and release tyramine to inhibit egg laying, in part via the LGC-55 tyramine-gated Cl⁻ channel on the HSNs. Our results identify discrete signals that entrain or detach the circuit from the locomotion central pattern generator to produce active and inactive states.

Introduction

Neural circuits are the functional units underlying all thoughts and behaviors, but there is as yet no neural circuit in any organism for which we know with precision how the many signaling events among its cells together generate its activity. This would require, among other things, understanding how neuromodulators shape activity at chemical and electrical synapses to alter the excitability of specific cells in the circuit to generate regulated, dynamic patterns of circuit responses and a coherent behavioral output. One approach to this problem is to analyze small neural circuits typical of invertebrate model organisms, so that both the simplicity of the circuits and the powerful experimental approaches uniquely possible within these systems can provide a penetrating analysis (Marder, 2012).

C. elegans egg-laying behavior is controlled by a small circuit that offers many experimental advantages for study (Schafer, 2006). This circuit, diagrammed in Figure 1A, consists of two serotonergic Hermaphrodite Specific Neurons (HSNs) and six cholinergic Ventral C neurons (VCs), each of which synapse onto a set of vulval muscles whose contraction expels eggs. Four neuroendocrine uv1 cells also regulate egg laying (Jose et al., 2007). Despite its anatomical simplicity, the egg-laying circuit produces a regulated, rhythmic behavior that alternates between quiescent periods of about 20 minutes during which no egg laying occurs, and active states lasting a few minutes during which ~5 eggs are laid (Waggoner et al., 1998). Active states appear to result when the HSNs release serotonin that signals through G protein coupled receptors on the vulval muscles to increase their excitability (Waggoner et al., 1998; Shyn et al., 2003; Hapiak et al., 2009; Emtage et al., 2012). Strong regulation by sensory stimuli is superimposed on the pattern of alternating behavioral states. For example, carbon dioxide regulates neuropeptide release from head sensory neurons that

signal through receptors on the HSNs to inhibit egg laying (Ringstad and Horvitz, 2008; Hallem

et al., 2011). Worms also halt egg laying in the absence of food and restart the behavior when

re-fed (Daniels et al., 2000; Dong et al., 2000).

C. elegans egg laying has been studied for decades, and dozens of genes have been

identified by mutations that either reduce (Desai et al., 1988) or increase (Bany et al., 2003)

egg laying. Some of the identified genes encode ion channels that regulate cell and synaptic

electrical excitability (Elkes et al., 1997; Johnstone et al., 1997; Lee et al., 1997; Weinshenker

et al., 1999; Jospin et al., 2002; Jose et al., 2007; Collins and Koelle, 2013). Other identified

genes encode components of G protein signaling pathways that act in specific cells of the

circuit (Brundage et al., 1996; Koelle and Horvitz, 1996; Hajdu-Cronin et al., 1999; Miller et al.,

1999; Williams et al., 2007; Porter et al., 2010). These results suggest neuromodulators,

including serotonin, signal through G proteins to regulate the excitability of specific cells in the

circuit to control circuit activity and egg laying.

The ability to leverage egg-laying mutants to understand the molecular mechanisms that

initiate, sustain, and terminate the active state of egg laying has been limited by the tools

available to analyze activity in the cells of the egg-laying circuit. Ca^{2+} activity in cells in the

circuit was initially optically recorded in immobilized animals, which have limited ability to

engage in egg-laying behavior (Shyn et al., 2003; Zhang et al., 2008). More recently we

developed methods to optically record Ca^{2+} activity of the vulval muscles in moving animals as

they engage in normal egg-laying behavior. Using ratiometric imaging to correct for movement

and focus artifacts, and recording at 20 frames/sec for periods of up to one hour, we were able

to measure activity of the vulval muscles as animals cycled through active and inactive states

(Collins and Koelle, 2013; Li et al., 2013). Our recordings showed that the vulval muscles are

excited rhythmically in phase with the body bends of animal locomotion, and we proposed this

was due to signaling from the cholinergic motor neurons, including the VCs, that are

rhythmically active during locomotion. We found that a conserved ERG K^+ channel depresses

response of the vulval muscles to excitation, thus maintaining the inactive behavior state, and

we proposed that serotonin signaling onto vulval muscle receptors enhances vulval muscle

excitability to allow the strong muscle contractions and egg release seen in the active state.

We have now generated tools to carry out Ca^{2+} recordings in moving animals for each neuron

type in the egg-laying circuit. Combining these recordings with precise manipulations of the

circuit using mutations, optogenetics, and drugs allows us to deduce how specific signals from

these cells drive specific features of circuit activity and behavior.

Results

Distinct patterns of activity in the egg-laying circuit accompany the active and inactive

behavior states

We carried out long-term Ca^{2+} recordings of HSN and VC neuron activity in behaving

animals using methods we developed previously for the vulval muscles (Collins and Koelle,

2013). We generated transgenic strains of *C. elegans* co-expressing the fluorescent Ca^{2+}

reporter GCaMP5 and the Ca^{2+}-independent fluorescent protein mCherry specifically in the

HSN or VC neurons, and used ratiometric imaging to record cell activity and egg-laying

behavior as animals moved through active and inactive egg-laying states (Figure 1). Because

of the large size of the egg-laying synapse, we were able to clearly observe Ca^{2+} transients

localized to the presynaptic terminus in HSN (Figure 1B; Movie 1) and VC (Figure 1C; Movie 2).

For comparison, we also carried out Ca^{2+} recordings of vulval muscles, and as in our earlier

studies of these muscles, were able to clearly distinguish the weak twitching contractions

(Figure 1D; Movie 3) from the strong contractions that drove egg release (Figure 1E).

Through long-term Ca^{2+} imaging, we observed a striking increase in rhythmic activity in

the cells of the circuit that began prior to the first egg-laying event and persisted beyond the

last egg-laying event of each active state (Figure 1F). We defined the active state as one

minute before the first egg-laying event and one minute after the last egg-laying event.

We found that HSNs displayed long (~4 seconds at half-maximum amplitude) Ca^{2+}

transients that were largely consistent in waveform and amplitude, but that varied in occurring

either as single events or trains (Figure 1F). There was also significant HSN activity outside of

the active state, but HSN transients were more frequent during the active state (Figure 1G).

Active state HSN transients typically occurred in bursts, within which HSN transients occurred

every ~20 seconds (Figure 1G). Only 11% of HSN transients were accompanied by an egg

release (Figure 1H), but almost every egg-laying event occurred during an HSN transient

(Figure 1F; out of 49 total egg-laying events observed, 48 occurred within an HSN transient).

VC transients were shorter than HSN transients (~2 seconds at half-max) and variable

in amplitude. We rarely observed VC transients during the inactive state, but we found a clear

induction of transients during the active state (Movie 2; Figure 1F, G), and these transients

occurred in bursts around egg-laying events with an inter-transient interval of ~10 seconds,

about twice as frequent as seen for the HSNs. Although Ca^{2+} transients were most intense in

the VC presynaptic varicosities and axonal processes, they were also visible in all six VC cell

bodies, especially those of the VC4 and VC5 neurons that most closely flank the vulva (Figure

1C and Movie 2). Vulval muscle twitching and egg-laying contractions caused significant

mechanical deformation of the VC neurons. 34% of VC transients were accompanied by an

egg release, a larger fraction than seen for HSN and vulval muscle Ca^{2+} transients (Figure 3H,

p <0.0001; Fisher's exact test). Every egg-release event was accompanied by a VC transient.

There was no correlation seen between the magnitudes of VC or HSN transients that did or did

not coincide with egg release, while the magnitude of muscle Ca^{2+} transients that resulted in

egg laying were ~4-fold larger than the muscle twitch transients that did not release eggs

(Figure 1H).

The close apposition of HSN, VC, and vulval muscle synaptic regions prevented us from

simultaneously imaging more than one cell type, but we were able to compare timing of activity

in the different cells using the moment of egg release as an objective landmark. HSN Ca^{2+}

transients peaked ~2 seconds prior to egg release (Figure 1I) and decayed slowly. Shorter VC

Ca^{2+} transients followed HSN activity, peaking ~0.1 seconds before egg release. Vulval muscle

Ca^{2+} transients peaked with egg laying. We also tracked vulval muscle contraction by

measuring the size and fluorescence intensity of the mCherry labeled muscles, and saw that

muscle relaxation and decay of vulval muscle Ca^{2+} immediately followed egg release (Figure

1I).

HSN, VC, and vulval muscle activity are phased with animal locomotion

C. elegans moves with sinusoidal body bends consisting of waves of body-wall muscle

contraction driven by the Ventral A and B (VA/VB) ventral nerve cord motor neurons. We

previously observed that vulval muscle Ca^{2+} transients occurred rhythmically when the vulva

was at a particular phase of the locomotion body bend (Collins and Koelle, 2013). Using our

video recordings of Ca^{2+} activity in moving animals, we examined whether HSN and VC activity

are similarly phased with locomotion. Thus we determined timing of the peaks of HSN, VC, and

vulval muscle Ca^{2+} transients relative to when the vulva passed through the previous and

subsequent most contracted or relaxed state of a body bend. The results were plotted on

circular histograms with ventral contraction at 0/360°, ventral relaxation at 180°, and

intermediate phases of body bends at the angles in between (Figure 2A). We found that the

HSN, VC, and vulval muscle Ca^{2+} transients occurred when the vulva was at particular phases

of the locomotor body bend. HSN transients just prior to egg release reached their maximum at

~180°, when the vulva and adjacent body wall muscles were at their most relaxed state (Figure

2B). VC and vulval muscle transients that accompanied egg laying occurred later, starting at

~225° and 270°, respectively (Figure 2C and 2D). HSN transients that did not result in egg

laying showed no clear evidence of phasing (Figure 2B). In contrast, VC transients and vulval

muscle twitch transients that did not lead to egg release were still phased, but delayed by ~45°

toward the most ventral contracted state. These results confirm our previous observations that

vulval muscle twitching and egg-laying activity are phased with locomotion (Collins and Koelle,

2013). These results extend that analysis to show that HSN, VC, and vulval muscle activity

during egg laying events occur earlier in the body bend cycle, when the adjacent body wall

muscles are in a more relaxed state.

Optogenetic activation of HSNs initiates the egg-laying active state

To directly test how neurotransmitter signaling from the HSNs regulates egg-laying

circuit activity, we used the *egl-6* promoter to express Channelrhodopsin-2 in the HSNs

(Emtage et al., 2012), allowing us to drive neurotransmitter release specifically from the HSNs

with blue light. We repeated the observation of Emtage et al. (2012) that optogenetic activation of HSNs stimulates egg-laying behavior (Movie 4), but we activated the HSNs while simultaneously recording behavior and Ca^{2+} activity in the VCs (Figure 3A) or vulval muscles (Figure 3B). We found that activation of HSNs resulted in circuit activity reminiscent of a spontaneous active state, including rhythmic Ca^{2+} activity of both VCs and vulval muscles, and egg-laying events that accompanied a subset of these Ca^{2+} transients. Some differences from a spontaneous active state were notable. Almost every VC transient after optogenetic HSN activation was accompanied by an egg-laying event, whereas only one third of VC transients during spontaneous active states were accompanied by egg-laying events. These results suggest that the high level of HSN activity after optogenetic activation induces strong coupling of VC and vulval muscle excitation.

During both spontaneous and optogenetically-induced active states, rhythmic vulval muscle twitches occur far more frequently than do VC transients, suggesting these twitches are excited by something other than the VCs. Careful observation shows that the most dramatic Ca^{2+} changes during vulval muscle twitching occur at the ventral tips of the vm1 muscles (Figure 1D). The VCs innervate vm2 muscles but not the vm1 muscles. The vm1 muscles are instead innervated by the VB6 and VA7 neurons that also release acetylcholine to drive body wall muscle contraction for locomotion (White, J.G. et al., 1986). Consistent with this, vm1 twitching occurs at the same time the adjacent ventral body wall muscles are contracting during locomotion (Figure 2D). Increased electrical excitability of the vulval muscles during a spontaneous or optogenetically-induced active state may allow for vm1 excitation by VB6/VA7 with each body bend.

VC neuron activity helps coordinate egg laying and locomotion

Our observation that VC activity always accompanies egg release (Figure 1I and 3A) is consistent with a proposal that the VCs release acetylcholine that signals through vulval muscle nicotinic receptors to drive egg laying (Waggoner et al., 2000; Kim et al., 2001). While VC4 and VC5 make extensive synapses onto the vm2 muscles, VC1-3 and VC6 make extensive, distributed synapses onto the ventral body wall muscles (White et al., 1976). How VC activity during egg laying might affect body wall muscle contraction has not been examined. We optogenetically activated the HSNs to induce active states and recorded animal speed around egg-laying events (Figure 3C). We found a clear and significant reduction of locomotion speed beginning ~1 s before egg release, a time when VC activity is increasing (Figure 1I). Locomotion speed recovered immediately after egg release (Figure 3C). These results are consistent with the hypothesis that acetylcholine released from the VCs slows locomotion during egg laying.

To test directly how VC activity regulates locomotion and egg laying, we used a *lin-11* promoter fragment to express Channelrhodopsin-2 specifically in the VCs (Bany et al., 2003). Optogenetic activation of the VCs was not sufficient to drive egg laying (Figure 3F), but did lead to a rapid shortening of the body (Movie 5 and Figure 3D) and cessation of locomotion (Figure 3E). The shortening phenotype seen in VC ChR2 animals after blue light exposure closely resembles that seen for optogenetic activation of the body wall muscles (Nagel et al., 2005), except the head and tail of VC-activated animals were still mobile and active, consistent with the fact that VC neurons innervate only the central body wall muscles (White et al., 1976). Our results indicate that while VC activity immediately precedes egg-laying events, it is not sufficient to drive egg release. Optogenetic VC activation is sufficient to produce a slowing of locomotion as seen during egg-laying events. In contrast, HSN activity is sufficient to induce all

217 of the hallmarks of the active state, including rhythmic vulval muscle twitching, VC transients,

218 and the coordinated vulval muscle contractions that produce egg-laying events.

Animals lacking HSNs still enter the egg-laying active state

We next tested whether HSN activity was necessary for circuit activity by recording

vulval muscle activity in *egl-1(dm)* mutants, which lack HSN neurons (Trent et al., 1983).

Animals lacking HSNs have a substantial lengthening of the inactive state (Waggoner et al.,

1998) and are strongly egg-laying defective, accumulating ~50 eggs in the uterus rather than

the 12-15 seen in the wild type (Trent et al., 1983). We expected to find that loss of HSNs led

to a dramatic reduction in vulval muscle activity. Surprisingly, we found the opposite; animals

lacking HSNs had more frequent vulval muscle Ca^{2+} transients during the inactive state and

robust activity during infrequent active states that led to multiple eggs being laid in a short

period (compare Figures 4A and 4B; quantitation in Figure 4C). There was a significant

reduction in the amplitude of vulval muscle Ca^{2+} transients during egg laying in *egl-1(dm)*

mutants while the amplitude of twitch Ca^{2+} transients was unaffected (Fig. 4D). Vulval muscle

activity in *egl-1(dm)* mutants lacking HSNs resembled the frequent yet uncoordinated vulval

muscle activity found in animals missing the vm2 vulval muscle arms onto which the HSNs

normally make synapses (Li et al., 2013). Such activity involves asynchronous contraction of

the anterior and posterior vulval muscles, which is ineffective at allowing egg release. One

quantitative measure of this asynchrony is that overall vulval muscle Ca^{2+} transient peaks in

egl-1(dm) mutants were longer in duration, likely resulting in individual muscle cells beginning

their transients at different times (Figure 4E). Vulval muscle Ca^{2+} transients in animals lacking

HSNs remained phased with locomotion, including the ~45° advance in phase of vulval muscle

activity that results in egg-laying events (compare Figures 4F and 4G). These results suggest

that the HSNs are necessary to coordinate contraction of anterior and posterior vulval muscles,

and that signals from other cells besides the HSNs can also induce the egg-laying active state.

We hypothesized that accumulating unlaid eggs could stretch the uterus or body,

leading to signals responsible for the increased activity of the vulval muscles independent of

neurotransmitter signaling from the HSNs. To test if egg accumulation contributed to the onset

of the active phase, we sterilized animals with FUDR (Mitchell et al., 1979), a DNA synthesis

inhibitor that blocks germ line cell division and thus egg production. FUDR-treated animals had

a dramatic reduction of vulval muscle activity and appeared to be stuck in the inactive state,

with no egg-laying transients observed (Figures 4A and 4B). The inhibition of vulval muscle

activity by FUDR occurred in both wild-type animals and HSN-deficient *egl-1(dm)* mutants

(Figures 4C and 4D). The residual vulval muscle Ca^{2+} twitch transients in FUDR-treated

animals had phasing identical to those seen in the inactive state of untreated animals (Figures

4F and 4G). These results suggest that germ line activity and/or the accumulation of eggs in

the uterus is required to initiate the egg-laying active state.

The uv1 neuroendocrine cells are mechanically deformed and activated by egg-laying

events

We previously identified a role for the uv1 neuroendocrine cells in inhibiting egg-laying

behavior (Jose et al., 2007). The four uv1 cells sit at the junction between the uterine lumen

and the vulval canal (Newman and Sternberg, 1996), and as such are positioned to

mechanically sense the accumulation of unlaid eggs in the uterus and/or passage of eggs

through the vulval canal as they are laid. To record uv1 Ca^{2+} activity, we co-expressed

GCaMP5 and mCherry in uv1 from the *ocr-2* promoter and made recordings in behaving

animals. We found that the uv1 cells were mechanically deformed as each egg passed through

the vulva, and then showed strong Ca^{2+} transients after egg release (Figure 5A-B; Movie 5).

Egg-laying events did not always trigger Ca^{2+} transients in uv1, with uv1-silent events usually

being the first or last within an active state (data not shown). We also observed weaker uv1

Ca^{2+} transients a few minutes before or after the egg-laying active state (Figure 5B-D).

Because these were not closely associated with egg-laying events their functional importance

remains unclear. Typical active state uv1 Ca^{2+} transients began immediately with or slightly

before egg release, coinciding with the initial mechanical deformation of uv1 caused by the egg,

with the Ca^{2+} transient peak occurring ~2 seconds later (Figure 5E). Based on the position of

the uv1 cells, their morphological deformations during passage of eggs, and the timing of uv1

Ca^{2+} transients, our data suggest the uv1 cells are mechanically activated by eggs passing

through the vulva during egg laying.

Tyramine signals through LGC-55 receptors to inhibit HSN activity and egg laying

The uv1 cells make the biogenic amine neurotransmitter tyramine, and *tdc-1* mutations

that disrupt tyramine biosynthesis lead to increased egg-laying behaviors (Alkema et al., 2005).

We found that exogenous tyramine reduced egg laying in wild-type animals (Figure 6A). To

identify targets of tyramine inhibition, we tested response to exogenous tyramine in strains

mutant for three known *C. elegans* tyramine receptors. Animals lacking *lgc-55*, which encodes

a tyramine-gated Cl⁻ channel, showed resistance to inhibition by exogenous tyramine (Figure

6A), although tyramine at high levels still had a residual ability to reduce egg laying in *lgc-55*

mutants (Figure 6B). We transformed into *C. elegans* a fosmid transgene containing the full-

length *lgc-55* gene fused to GFP coding sequences, and found LGC-55 is expressed in the

HSNs, uv1, and vulval muscles (Figure 6C and Pirri et al., 2009). Re-expression of LGC-55

specifically in HSN from the *tph-1* promoter restored tyramine sensitivity to *lgc-55* mutants

(Figure 6D). We have previously shown that two Cl^- extruding transporters, KCC-2 and ABTS-1,

are expressed in the HSNs where they promote the development of inhibitory ligand-gated Cl^-

channel signaling (Tanis et al., 2009; Bellemer et al., 2011). These data suggest that tyramine

signaling through LGC-55 would hyperpolarize the HSN and inhibit activity. To test this directly,

we compared HSN activity in wild-type and *lgc-55* mutant animals. We observed a significant

increase in the frequency of Ca^{2+} transients in HSNs of *lgc-55* mutant animals (Figures 6E and

6F) in both the inactive and active states of egg-laying behavior. Mean HSN inter-transient

intervals in wild-type animals were 41±5 seconds in the inactive state and 17±2 seconds during

the active state, while intervals in *lgc-55* mutants were reduced to 22±2 seconds in the inactive

state and 13±1 seconds during the active state. Thus, the absence of inhibitory feedback by

tyramine signaling onto the HSNs leads to increased activity in both the active and inactive

egg-laying behavior states.

Discussion

Using a combination of optogenetic, molecular genetic, and Ca^{2+} imaging approaches,

we show an unexpected level of coordination of *C. elegans* egg-laying circuit activity by the

rhythm of locomotion and by egg production. Using recordings of cell activity in behaving

animals as they transition from the inactive to active egg-laying states, we were able to show

that circuit activity is diminished during the inactive state, strongly increased and rhythmic

during the active state, and this rhythm is phased with sinusoidal locomotion. Sterilization

eliminated the active state rhythm, locking animals in the inactive state. Our results suggest

that activity in the egg-laying circuit, like that of many other modulated motor circuits, responds

to rhythmic input from central pattern generators (Marder et al., 2015).

A working model shown in Figure 7 explains how different inputs into the egg-laying

circuit are responsible for the inactive state and the active state rhythms observed during egg-

laying behavior. Active state vulval muscle Ca^{2+} transients peak rhythmically each time the

vulva approaches a ventral body bend, the point at which the cholinergic VB and VA motor

neurons release acetylcholine to rhythmically excite both the vm1 muscles and adjacent

ventral body wall muscles during locomotion. During the inactive state, K^+ channels including

ERG depress vulval muscle excitability below threshold, explaining why we fail to observe

vulval muscle activity with each body bend during this state (Collins and Koelle, 2013). While

we observe rhythmic activity in the HSNs in the active phase, the periodicity and phasing of

HSN activity does not correlate with that of locomotion. Consistent with this, previous work

found changes in osmolarity can drive rhythmic HSN activity even in immobilized animals

(Zhang et al., 2008). Thus HSN rhythmic activity is independent of locomotion, although we

found HSN Ca^{2+} transients that happen to fall at a specific phase of locomotor body bends are

more likely to be followed by egg release. We propose that release of serotonin from the HSNs,

along with an unknown signal resulting from mechanical stretch of the uterus by accumulating

eggs, together raise vulval muscle electrical excitability to result in the active state during which

the vulval muscles undergo rhythmic twitching or egg laying contractions in response to the

same cholinergic VA and VB neurons that drive locomotor body bends. This model accounts

for the phasing of vm1 transients with body bends, although egg-laying events require

additional activity in the vm2 muscles that results in the coordinated contraction of all vm muscles that results in egg release. We found that vm transients that lead to egg release are on average slightly phase advanced relative to those that do not. This phase advance could from the excitatory input onto the vm2 muscles occurring earlier in the body bend, and candidates for this input are the HSNs and VCs, both of which synapse onto the vm2 muscles and have phased activity during egg-laying events. The functional purpose of this phase advance may be mechanical. Passage of each 25x40μm egg requires contraction of vm1 and vm2 muscles on both the anterior and posterior sides of the vulva (Li et al., 2013). There may not be sufficient room for vulva to fully open and release egg unless the adjacent body wall muscles are in a contracted state.

The HSNs are command neurons that integrate diverse sensory inputs and initiate the ~2 minute active state of the egg-laying circuit. Optogenetic activation of HSNs recapitulates features of the active state including phased VC and vulval muscle activity and multiple egg-laying events. However, stimulation of the HSNs fails to drive tetanic contraction of the vulval muscles as would be predicted for a motor neuron releasing a fast-acting neurotransmitter. Instead, HSN makes and releases the neuromodulator serotonin which signals through G protein coupled serotonin receptors to activate the vulval muscles (Desai et al., 1988; Shyn et al., 2003; Carnell et al., 2005; Hobson et al., 2006; Hapiak et al., 2009). The HSNs also release serotonin onto the AVF interneurons in the nerve ring to promote a state of locomotion arousal that accompanies the active state (Hardaker et al., 2001). Animals defective in serotonin signaling have reduced egg laying (Tanis et al., 2008), but this defect is not as severe as that of *egl-1(dm)* animals missing HSNs (Trent et al., 1983), suggesting HSN releases additional neurotransmitters. We found that optogenetic activation of HSNs leads to

rhythmic activity of VC neurons, which are not known to express serotonin receptors, so an unknown signal derived from HSN could directly activate VCs, as depicted in the Figure 7 model.

How is HSN activity controlled? The HSNs receive little synaptic input near the vulval except from the PLM mechanosensory neurons (White, J.G. et al., 1986). The PLMs inhibit HSN activity and egg laying in response to gentle touch (Zhang et al., 2008), but recent work indicates the PLM neurons are also deformed during normal patterns of locomotion that could contribute to rhythmic activity seen in HSN during egg laying (Topalidou et al., 2012). Neuroendocrine signaling also regulates HSN activity. *flp-10* and *flp-17* neuropeptides are released from the CO_2-responsive BAG neurons in the head, and these neuropeptides signal through EGL-6 receptor and $G\alpha_o$ to activate IRK channels that inhibit HSN activity and egg laying (Ringstad and Horvitz, 2008; Hallem et al., 2011; Emtage et al., 2012). Because loss of EGL-6 or IRK channels do not increase egg laying as much as cell-specific inhibition of $G\alpha_o$, there must be additional receptors and effectors of $G\alpha_o$ that inhibit HSN activity. $G\alpha_q$ signaling opposes $G\alpha_o$ to increase serotonin release from the HSNs (Tanis et al., 2008), but the ligands and receptors that activate $G\alpha_q$ have not yet been identified. Because we find that HSN Ca^{2+} transients vary more in frequency than amplitude, we propose that these diverse sensory signals increase or reduce the probability that HSN reaches a threshold potential that can initiate, sustain, or terminate HSN activity and thus the active behavior state of the egg-laying circuit.

We have also identified an HSN-independent mechanism that induces the active state. Mutants lacking HSNs are strongly egg-laying defective, but eventually enter active states with properly phased vulval muscle twitching and egg-laying transients. The active state was

eliminated in animals sterilized by FUDR, reducing vulval muscle activity and phasing to that seen in the inactive state. Recent work has identified the gonad as an important site of regulation for adult-specific behaviors in *C. elegans* (Fujiwara et al., 2016). We propose that the accumulation of unlaid eggs in the uterus, perhaps leading to mechanical stretch, provides an essential, permissive signal for entry into the active state. Several steps during embryo production and maturation could regulate entry into the active state to result in the ~20 period of the egg-laying inactive state.. Animals grown on food generate one new oocyte from each of two gonad arms every ~10 minutes, balancing out the 3-5 eggs laid in the ~2 minute active state that occurs every ~20 minutes, to result in a ~constant number of unlaid eggs in the uterus (McCarter et al., 1999). New embryos are moved by gonadal sheath contractions into the spermatheca for fertilization and deposition into the uterus. Excitable cells drive each of these steps, providing potential sources of regulation by cell signaling and sensory feedback (Kariya et al., 2004). Eight uterine muscles line the uterine lumen and form gap junctions with the vm2 muscles (White, J.G. et al., 1986), and the addition of unlaid eggs into the uterus could induce a mechanical stretch that cross-excites the vm2 muscles. Future work will be required to understand how the accumulation of unlaid eggs promotes the active state.

Our results strongly imply that acetylcholine released from the VC motor neurons slows locomotion during vulval muscle contraction. We found that animal locomotion slows during egg release, and that optogenetic activation of the VC neurons halts movement and results in body wall muscle contraction and animal shortening. This function of the VCs is likely via acetylcholine signaling through nicotinic receptors expressed on ventral body wall muscles and neurons that regulate locomotion (White, J.G. et al., 1986; Nagel et al., 2005). The effect of VC activity to slow locomotion and contract body wall muscles may help compress the body and

uterus to encourage expulsion of eggs, analogous to the role of body wall muscle contraction in

facilitating defecation (Liu and Thomas, 1994).

Our results provide more complex evidence as to whether VC acetylcholine also directly

triggers egg release. We found that VC activity always accompanies egg-laying contractions of

the vulval muscles. The vulval muscles also express nicotinic acetylcholine receptors, and

receptor agonists including levamisole and nicotine stimulate egg laying (Weinshenker et al.,

1995; Waggoner et al., 2000). However, we found little correlation between the strength of the

VC activity and whether an egg is laid (Figure 1H), and that optogenetic activation of the VC

neurons fails to drive egg laying. Further, VC-defective and acetylcholine-deficient mutants

have *increased* egg-laying behavior (Bany et al., 2003; Ringstad and Horvitz, 2008). These

data together are consistent with a model in which acetylcholine released from the VA/VB

motor neurons signal through nicotinic receptors on the vm1 muscles to stimulate rhythmic

vulval muscle twitching, which combined with acetylcholine released from the VC neurons to

similarly excite the vm2 muscles results in egg laying. High levels of VC activity, such as

occurs during optogenetic activation, may drive release of additional signals, such as

neuropeptides, that inhibit egg-laying circuit activity (Bany et al., 2003; Banerjee et al., 2016),

or by freezing locomotion may feedback to halt the VA/VB activity required to drive vm1

contraction.

How is VC activity controlled? The VC neurons are unusual in that they are largely

presynaptic, only receiving input from each other and the HSNs. As noted above, optogenetic

activation of HSNs results in VC activity, suggesting a currently unknown signal from HSN

directly activates VCs. A second possible input onto VCs is suggested by the fact that the VC

neurons extend processes devoid of synapses dorsally along the vulval hypodermis that could

be mechanosensory (White, J.G. et al., 1986). Thus, rhythmic VC activity could be stimulated mechanically by body bends or by a humoral signal released from the locomotor circuit (Hu et al., 2011). Further, VC activity could be mechanically stimulated by vulval muscle contraction (Zhang et al., 2010), particularly as we observed mechanical deformation of VC presynaptic termini during strong twitching and egg-laying contractions. In this way, the VCs could act like baroreceptors–mechanically activated by vulval muscle contraction to drive contraction of the body wall muscles that results in animal slowing until the feedback of egg release. In this model, the coordination of locomotion and egg laying by tying both behaviors to the same central pattern generator allows productive egg release.

The uv1 neuroendocrine cells inform the circuit that eggs have been successfully laid. Passage of eggs through the vulva mechanically deforms the uv1 cells and triggers a strong Ca^{2+} transient. Our results suggest this activates uv1 to release tyramine that inhibits egg-laying, at least in part via the LGC-55 receptor on the HSN neurons. Active states last ~2 minutes, and inhibitory tyramine release from uv1 could provide a feedback signal that terminates activity in the circuit. The four uv1 cells have a unique structural position in the vulva that may explain how they become activated during egg laying (Newman et al., 1996). uv1 cells extend processes that contact and make adherens junctions with the ventral vulF vulval cells. uv1 cells also makes dorsal attachments to utse, a large, H-shaped multinucleate uterine seam cell that attaches the uterus to the lateral epidermis and that functions as a hymen broken by the first egg-laying event. The acute mechanical strain on vulF, uv1, and/or utse during egg passage could mechanically activate ion channels. Previous work has shown uv1 expresses several TRPV channels that may be mechanosensory and contribute to this response (Jose et al., 2007). A dominant-negative OCR-2 mutant expressed in uv1 leads to

hyperactive egg-laying behavior phenotype. We still observe Ca^{2+} transients in uv1 after egg-

laying events in this *ocr-2(vs29)* mutant (data not shown), suggesting other channels contribute

to the uv1 activity we see after egg laying. Although our results indicate that uv1 releases

tyramine to inhibit the HSNs via LGC-55, this signaling event does not fully explain the ability of

uv1 cells to inhibit the egg-laying circuit. We found that inhibition of egg laying by exogenous

tyramine is reduced but not eliminated in *lgc-55* null mutants, suggesting other tyramine

receptors help terminate the active state. The uv1 cells also make *nlp-7* and *flp-11*, two

neuropeptides that inhibit HSN activity and egg-laying behavior (Banerjee et al., submitted),

and the identification of the receptors and signaling pathways through which they inhibit

serotonin release will help to more fully understand how activity of the egg-laying circuit is

terminated.

Every neural circuit may require specific signals to initiate activity when its function is

required, signals to coordinate the dynamic activity of its various cells, and mechanisms to

determine when the circuit has successfully completed its task and signal to turn off activity.

Many circuits also must respond to specific signals from central pattern generators that

coordinate the activity of multiple circuits. A great challenge is to reduce these abstract

concepts to concrete, specific mechanisms in the case of a real circuit. For the *C. elegans* egg-

laying circuit, signals in each of the categories outlined above are now identified, albeit in

varying levels of detail and completeness. Continuing analysis of the *C. elegans* egg-laying

circuit using recordings of Ca^{2+} activity in behaving animals, combined with genetic,

optogenetic, and pharmacological manipulation, provides a path towards detailed

understanding the mechanisms that control activity of this model circuit.

Materials and Methods

Nematode Culture, Strains, and Maintenance

Caenorhabditis elegans strains were maintained as hermaphrodites at 20°C on Nematode Growth Medium (NGM) agar plates with *E. coli* OP50 as a source of food as described (Brenner, 1974). All strains are derived from the Bristol N2 wild-type strain, and all assays were performed using age-matched adult hermaphrodites ~24 hours past the late L4 stage. Animals were treated with 0.1 mg/ml floxuridine (FUDR) at a late L4 stage when all non-germline cell divisions are complete. A list of strains, mutants, and transgenes used in this study can be found in Table 1.

Molecular Biology and Transgenes

Vulval muscles: GCaMP5G (Akerboom et al., 2012) and mCherry were expressed in the vulval muscles from the *unc-103e* promoter as previously described (Collins and Koelle, 2013). Briefly, pKMC274 (GCaMP5, 20ng/*µ*l) and pKMC257 (mCherry, 2ng/*µ*l) were co-injected along with the pL15EK *lin-15* rescue plasmid (50ng/*µ*l) into LX1832 *lite-1(ce314), lin-15(n765ts)* X animals. The GCaMP5/mCherry extrachromosomal transgene produced was chromosomally integrated using UV/TMP mutagenesis, creating the integrated transgene *vsIs164*, and the resulting transgenic animals were backcrossed to the LX1832 parental line six times.

HSN: GCaMP5G and mCherry were expressed in the HSN motor neurons using the *nlp-3* promoter. mCherry and GCaMP5G coding sequences were ligated into pJB9 bearing the *nlp-3* promoter and *nlp-3* 3' untranslated region, and the resulting plasmids (GCaMP5, pKMC300, 80ng/*µ*l; mCherry, pKMC299, 20ng/*µ*l) were co-injected with pL15EK (50ng/*µ*l) into LX1832 worms. The extrachromosomal transgene produced was integrated with UV/TMP, creating the

integrated transgene *vsIs183,* and the resulting transgenic animals were backcrossed to the LX1832 parental line six times. The *wzIs30* transgene expressing Channelrhodopsin-2 from the *egl-6* promoter has been described (Emtage et al., 2012).

VC: GCaMP5G and mCherry were expressed in the VC motor neurons using pKMC145, a pDM4-based plasmid bearing a *lin-11* enhancer region fused to the *pes-10* basal promoter with a modified multiple cloning site (Bany et al., 2003; Tanis et al., 2008). Derivatives expressing GCaMP5 (pKMC273, 80ng/μl) and mCherry (pKMC275, 30ng/μl) were co-injected with pL15EK (50ng/μl) into LX1832 worms. The extrachromosomal transgene produced was integrated with UV/TMP, creating the transgene *vsIs172*, and transgenic animals were backcrossed to the LX1832 parental line six times. To activate the VCs optogenetically, coding sequences for ChR2-YFP bearing the H134R, T159C mutations (Erbguth et al., 2012) were amplified by PCR, ligated into pKMC145 to generate pKMC283, injected at 80ng/μl into LX1832 along with pL15EK at 50ng/μl. The extrachromosomal transgene produced was integrated with UV/TMP creating the transgene *keyIs3,* and the resulting animals were backcrossed to LX1832 parental line six times.

uv1: GCaMP5G and mCherry were expressed in uv1 using the *ocr-2* promoter. GCaMP5 and mCherry coding sequences were ligated into pJT79 between the *ocr-2* promoter and 3' untranslated region (Jose et al., 2007) to generate plasmids pKMC281 and pKMC284, respectively. Long-range PCR products amplified from these plasmids were injected into worms at 80ng/μl each along with pL15EK (50ng/μl) as previously described (Jose et al., 2007). The extrachromosomal transgene produced was integrated with UV/TMP creating the transgene *vsIs177*, and the resulting animals were backcrossed to the LX1832 parental line six times.

lgc-55: A ~40 kb genomic fosmid clone (WRM063D_A12; Perkins, 2011) recombineered to

express LGC-55 tagged with GFP at the C-terminus (Sarov et al., 2012) was injected at

100ng/μl into LX2081 *unc-119(ed3)* mutant animals expressing TagRFP pan-neuronally to

generate the strain LX2126. These animals were then crossed with LX2096 animals to

exchange pan-neural mCherry transgene for one that expresses in the vulval muscles,

generating LX2126. Fluorescence was visualized on immobilized animals with a 63x water

objective on a Zeiss LSM 710 Duo confocal microscope. For cell-specific rescue, LGC-55 was

expressed in the HSN neurons of *lgc-55* mutant animals using the *tph-1* promoter. Briefly, a

lgc-55 cDNA (Pirri et al., 2009) was inserted into pJM66A plasmid (Moresco and Koelle, 2004)

to generate pJT105. pJT105, or a GFP-expressing control plasmid (pJM60A), was injected at

100ng/μl along with pL15EK rescue plasmid (50ng/μl) into *lgc-55(tm2913); lin-15(n765ts)*

double mutant animals. Egg-laying behavior was analyzed in five independent transgenic lines

expressing LGC-55 or GFP, and one strain each were kept for long-term storage.

Ratiometric Ca^{2+} Imaging

Animals were mounted between glass coverslips and chunked sections of NGM plates for

imaging as described (Collins and Koelle, 2013; Li et al., 2013). The stage and focus were

adjusted manually to keep the egg-laying system in view during recording periods. Three

different imaging systems were used to collect recordings, and each gave quantitatively similar

results. First, single, two-channel confocal slices (18 μm thick) of GCaMP5 and mCherry

fluorescence were collected through a 20x Plan Apochromat objective (0.8NA) using a Zeiss

Axio Observer microscope and recorded at 20 fps at 256x256 pixel resolution, 16-bit depth

using a 710 Duo LIVE scanner as described previously (Collins and Koelle, 2013; Li et al.,

2013). Second, widefield GCaMP5 and mCherry fluorescence was imaged through a 20x Plan

Apochromat objective (0.8NA) using a Zeiss Axio Observer microscope, and the GCaMP5 and

mCherry fluorescence was separated using a W-View Gemini image splitter at 20 fps at

256x256 pixel resolution, 16-bit depth onto a single, 4x4 binned Hamamatsu ORCA Flash 4.0

V2 sCMOS sensor after excitation with 470 nm and 590 nm LEDs (Zeiss Colibri.2). Third,

single, two-channel confocal slices (20 μm thick) of GCaMP5 and mCherry fluorescence were

collected using an inverted Leica TCS SP5 confocal microscope with the resonant scanner

running at 8,000Hz and collected at 20 fps at 256x256 pixel resolution, 12-bit depth after 2x

digital zoom.

Quantitative ratiometric analysis was performed after importing image sequences into

Volocity (Perkin Elmer). GCaMP5/mCherry ratios were calculated, and pixels with mCherry

fluorescence intensities ~2 standard deviations above background were selected as objects for

mean ratio measurement. While the Volocity algorithm set with these parameters typically

identified the vulval muscles and uv1 cells as a single object, the complex morphology of the

HSN and VC neurons often led to the cells being found as two or more objects within each

video frame. These were rejoined using a custom script in MATLAB (Mathworks). The ratio of

the joined object was scaled in proportion to the area of each component object to generate an

area-adjusted GCaMP5/mCherry ratio. The ratio data were then smoothed using a 150 msec

(three timepoint) rolling average, and the lowest 10% of the GCaMP5/mCherry ratio values

over the entire recording period were averaged to establish a ΔR/R baseline. Ca^{2+} transients

and their features (peak amplitude and width at half-maximum) were identified from ratio traces

using the *findpeaks* algorithm in MATLAB, and peaks were confirmed by visual inspection of

GCaMP5/mCherry recordings and ratio traces. Egg-laying events were observed by allowing a

small amount of bright field light to transmit into the mCherry channel whose gain was

increased for egg visualization. The egg-laying active state was defined in recordings to

include those transients that occurred one minute before and one minute after each egg-laying

event. Those transients not found within an active state were considered to be in the inactive

state. To facilitate comparisons of ΔR/R between different reporters and recording conditions

(Figure 1H and 5D), the maximum peak amplitude was normalized to 1. For phasing analysis,

the timing of individual Ca^{2+} transient peaks relative to ventral relaxation and contraction was

determined independent of whether the animal was engaged in forward or reverse locomotion,

as described previously (Collins and Koelle, 2013).

Optogenetics

Channelrhodopsin-2 (ChR2) expressing strains were grown in the presence or absence of the

ChR2 cofactor all-*trans* retinal (ATR). ATR was prepared at 100mM in 100% ethanol and

stored at -20°C. To prepare NGM plates for behavior analysis, ATR was diluted to 0.4 mM with

warmed cultures of OP50 bacteria in B Broth, and 200 μl of culture was seeded onto each 60

mm NGM plate. The plates were allowed to grow for 24 hours at 25-37°C, after which late L4

worms were staged onto prepared plates for behavioral assays 24 hours later. During Ca^{2+}

imaging experiments, ChR2 was activated with the same light used for GCaMP5 excitation.

For timed behavior recording experiments on plates, a OTPG_4 TTL Pulse Generator (Doric

Optics) was used to trigger image capture (Grasshopper 3, 4.1 Megapixel, USB3 CMOS

camera, Point Grey Research) and shutter opening on a EL6000 metal halide light source

generating 8-16 mW/cm^2 (depending on the magnification used) of ~470±20nm blue light via a

EGFP filter set mounted on a Leica M165FC stereomicroscope. Worm size, speed, and body

bends were determined from image sequences using Volocity.

Tyramine behavior assays

Three adult animals were placed onto each of three 35 mm agar plates prepared with

tyramine-HCl (Sigma) and acetic acid seeded with OP50 bacteria as described (Pirri et al.,

2009). The average number of eggs laid was counted after 30 minutes from at least 11 trials

for each genotype and tyramine concentration.

Statistical analyses

Data were analyzed using Prism 6 (GraphPad). Average worm speed, worm length, body

bends, or eggs laid were compared using one-way ANOVA with Bonferroni correction for

multiple comparisons, and error bars indicate 95% confidence intervals for the mean. Ca^{2+}

transient peak amplitudes, widths, and inter-transient intervals were pooled from multiple

animals (n≥5 animals per experiment), compared using the Mann-Whitney test (for pair-wise

comparisons) or the Kruskal-Wallis test with Dunn's correction (for multiple comparisons), and

error bars indicate quartile intervals for the indicated median. $p<0.05$ interactions were deemed

significant and are labeled with an asterisk.

Acknowledgments

This work was funded by a postdoctoral fellowship from the American Heart Association to

KMC (POST4990016) and by grants from the NINDS to MRK (NS036918) and to MRK and

KMC (NS086932). Confocal instrumentation was supported by a grant to the Yale Liver Center

(DK34989). Some strains were provided by the *C. elegans* Genetics Center, which is funded

by NIH Office of Research Infrastructure Programs (P40 OD010440). We thank members of

the Koelle and Collins labs, and Drs. James Baker, Julia Dallman, and Mason Klein for

comments and helpful discussions on the manuscript.

Competing Interests

607 The authors declare no financial and non-financial competing interests.

References

Akerboom J et al. (2012) Optimization of a GCaMP calcium indicator for neural activity imaging.
J Neurosci Off J Soc Neurosci 32:13819–13840.

Alkema MJ, Hunter-Ensor M, Ringstad N, Horvitz HR (2005) Tyramine Functions
independently of octopamine in the Caenorhabditis elegans nervous system. Neuron
46:247–260.

Bany IA, Dong M-Q, Koelle MR (2003) Genetic and cellular basis for acetylcholine inhibition of
Caenorhabditis elegans egg-laying behavior. J Neurosci Off J Soc Neurosci 23:8060–
8069.

Bellemer A, Hirata T, Romero MF, Koelle MR (2011) Two types of chloride transporters are
required for GABA(A) receptor-mediated inhibition in C. elegans. EMBO J 30:1852–
1863.

Brenner S (1974) The genetics of Caenorhabditis elegans. Genetics 77:71–94.

Brundage L, Avery L, Katz A, Kim UJ, Mendel JE, Sternberg PW, Simon MI (1996) Mutations
in a C. elegans Gqalpha gene disrupt movement, egg laying, and viability. Neuron
16:999–1009.

Carnell L, Illi J, Hong SW, McIntire SL (2005) The G-protein-coupled serotonin receptor SER-1
regulates egg laying and male mating behaviors in Caenorhabditis elegans. J Neurosci
Off J Soc Neurosci 25:10671–10681.

Collins KM, Koelle MR (2013) Postsynaptic ERG Potassium Channels Limit Muscle Excitability
to Allow Distinct Egg-Laying Behavior States in Caenorhabditis elegans. J Neurosci Off
J Soc Neurosci 33:761–775.

Daniels SA, Ailion M, Thomas JH, Sengupta P (2000) egl-4 acts through a transforming growth
factor-beta/SMAD pathway in Caenorhabditis elegans to regulate multiple neuronal
circuits in response to sensory cues. Genetics 156:123–141.

Desai C, Garriga G, McIntire SL, Horvitz HR (1988) A genetic pathway for the development of
the Caenorhabditis elegans HSN motor neurons. Nature 336:638–646.

Dong MQ, Chase D, Patikoglou GA, Koelle MR (2000) Multiple RGS proteins alter neural G
protein signaling to allow C. elegans to rapidly change behavior when fed. Genes Dev
14:2003–2014.

Elkes DA, Cardozo DL, Madison J, Kaplan JM (1997) EGL-36 Shaw channels regulate C.
elegans egg-laying muscle activity. Neuron 19:165–174.

Emtage L, Aziz-Zaman S, Padovan-Merhar O, Horvitz HR, Fang-Yen C, Ringstad N (2012)
IRK-1 Potassium Channels Mediate Peptidergic Inhibition of Caenorhabditis elegans
Serotonin Neurons via a Go Signaling Pathway. J Neurosci Off J Soc Neurosci
32:16285–16295.

Erbguth K, Prigge M, Schneider F, Hegemann P, Gottschalk A (2012) Bimodal activation of different neuron classes with the spectrally red-shifted channelrhodopsin chimera C1V1 in Caenorhabditis elegans. PloS One 7:e46827.

Fujiwara M, Aoyama I, Hino T, Teramoto T, Ishihara T (2016) Gonadal Maturation Changes Chemotaxis Behavior and Neural Processing in the Olfactory Circuit of Caenorhabditis elegans. Curr Biol CB 26:1522–1531.

Hajdu-Cronin YM, Chen WJ, Patikoglou G, Koelle MR, Sternberg PW (1999) Antagonism between G(o)alpha and G(q)alpha in Caenorhabditis elegans: the RGS protein EAT-16 is necessary for G(o)alpha signaling and regulates G(q)alpha activity. Genes Dev 13:1780–1793.

Hallem EA, Spencer WC, McWhirter RD, Zeller G, Henz SR, Rätsch G, Miller DM, Horvitz HR, Sternberg PW, Ringstad N (2011) Receptor-type guanylate cyclase is required for carbon dioxide sensation by Caenorhabditis elegans. Proc Natl Acad Sci U S A 108:254–259.

Hapiak VM, Hobson RJ, Hughes L, Smith K, Harris G, Condon C, Komuniecki P, Komuniecki RW (2009) Dual excitatory and inhibitory serotonergic inputs modulate egg laying in Caenorhabditis elegans. Genetics 181:153–163.

Hardaker LA, Singer E, Kerr R, Zhou G, Schafer WR (2001) Serotonin modulates locomotory behavior and coordinates egg-laying and movement in Caenorhabditis elegans. J Neurobiol 49:303–313.

Hobson RJ, Hapiak VM, Xiao H, Buehrer KL, Komuniecki PR, Komuniecki RW (2006) SER-7, a Caenorhabditis elegans 5-HT7-like receptor, is essential for the 5-HT stimulation of pharyngeal pumping and egg laying. Genetics 172:159–169.

Hu Z, Pym ECG, Babu K, Vashlishan Murray AB, Kaplan JM (2011) A neuropeptide-mediated stretch response links muscle contraction to changes in neurotransmitter release. Neuron 71:92–102.

Johnstone DB, Wei A, Butler A, Salkoff L, Thomas JH (1997) Behavioral defects in C. elegans egl-36 mutants result from potassium channels shifted in voltage-dependence of activation. Neuron 19:151–164.

Jose AM, Bany IA, Chase DL, Koelle MR (2007) A specific subset of transient receptor potential vanilloid-type channel subunits in Caenorhabditis elegans endocrine cells function as mixed heteromers to promote neurotransmitter release. Genetics 175:93–105.

Jospin M, Jacquemond V, Mariol M-C, Segalat L, Allard B (2002) The L-type voltage-dependent Ca2+ channel EGL-19 controls body wall muscle function in Caenorhabditis elegans. J Cell Biol 159:337–348.

Kariya K-I, Bui YK, Gao X, Sternberg PW, Kataoka T (2004) Phospholipase Cepsilon regulates ovulation in Caenorhabditis elegans. Dev Biol 274:201–210.

683   Kim J, Poole DS, Waggoner LE, Kempf A, Ramirez DS, Treschow PA, Schafer WR (2001)
684   Genes affecting the activity of nicotinic receptors involved in Caenorhabditis elegans
685   egg-laying behavior. Genetics 157:1599–1610.

686   Koelle MR, Horvitz HR (1996) EGL-10 regulates G protein signaling in the C. elegans nervous
687   system and shares a conserved domain with many mammalian proteins. Cell 84:115–
688   125.

689   Lee RY, Lobel L, Hengartner M, Horvitz HR, Avery L (1997) Mutations in the alpha1 subunit of
690   an L-type voltage-activated Ca2+ channel cause myotonia in Caenorhabditis elegans.
691   EMBO J 16:6066–6076.

692   Li P, Collins KM, Koelle MR, Shen K (2013) LIN-12/Notch signaling instructs postsynaptic
693   muscle arm development by regulating UNC-40/DCC and MADD-2 in Caenorhabditis
694   elegans. eLife 2:e00378.

695   Liu DW, Thomas JH (1994) Regulation of a periodic motor program in C. elegans. J Neurosci
696   Off J Soc Neurosci 14:1953–1962.

697   Marder E (2012) Neuromodulation of neuronal circuits: back to the future. Neuron 76:1–11.

698   Marder E, Goeritz ML, Otopalik AG (2015) Robust circuit rhythms in small circuits arise from
699   variable circuit components and mechanisms. Curr Opin Neurobiol 31:156–163.

700   McCarter J, Bartlett B, Dang T, Schedl T (1999) On the control of oocyte meiotic maturation
701   and ovulation in Caenorhabditis elegans. Dev Biol 205:111–128.

702   Miller KG, Emerson MD, Rand JB (1999) Goalpha and diacylglycerol kinase negatively
703   regulate the Gqalpha pathway in C. elegans. Neuron 24:323–333.

704   Mitchell DH, Stiles JW, Santelli J, Sanadi DR (1979) Synchronous growth and aging of
705   Caenorhabditis elegans in the presence of fluorodeoxyuridine. J Gerontol 34:28–36.

706   Moresco JJ, Koelle MR (2004) Activation of EGL-47, a Galpha(o)-coupled receptor, inhibits
707   function of hermaphrodite-specific motor neurons to regulate Caenorhabditis elegans
708   egg-laying behavior. J Neurosci Off J Soc Neurosci 24:8522–8530.

709   Nagel G, Brauner M, Liewald JF, Adeishvili N, Bamberg E, Gottschalk A (2005) Light activation
710   of channelrhodopsin-2 in excitable cells of Caenorhabditis elegans triggers rapid
711   behavioral responses. Curr Biol CB 15:2279–2284.

712   Newman AP, Sternberg PW (1996) Coordinated morphogenesis of epithelia during
713   development of the Caenorhabditis elegans uterine-vulval connection. Proc Natl Acad
714   Sci 93:9329–9333.

715   Newman AP, White JG, Sternberg PW (1996) Morphogenesis of the C. elegans hermaphrodite
716   uterus. Dev Camb Engl 122:3617–3626.

Perkins, Jaryn Daniel (2011) Comparison of fosmid libraries made from two geographic
isolates of Caenorhabditis elegans. Available at: http://dx.doi.org/10.14288/1.0071587
[Accessed August 6, 2016].

Pirri JK, McPherson AD, Donnelly JL, Francis MM, Alkema MJ (2009) A tyramine-gated
chloride channel coordinates distinct motor programs of a Caenorhabditis elegans
escape response. Neuron 62:526–538.

Porter MY, Xie K, Pozharski E, Koelle MR, Martemyanov KA (2010) A Conserved Protein
Interaction Interface on the Type 5 G Protein β Subunit Controls Proteolytic Stability and
Activity of R7 Family Regulator of G Protein Signaling Proteins. J Biol Chem
285:41100–41112.

Ringstad N, Horvitz HR (2008) FMRFamide neuropeptides and acetylcholine synergistically
inhibit egg-laying by C. elegans. Nat Neurosci 11:1168–1176.

Sarov M et al. (2012) A genome-scale resource for in vivo tag-based protein function
exploration in C. elegans. Cell 150:855–866.

Schafer WF (2006) Genetics of egg-laying in worms. Annu Rev Genet 40:487–509.

Shyn SI, Kerr R, Schafer WR (2003) Serotonin and Go modulate functional states of neurons
and muscles controlling C. elegans egg-laying behavior. Curr Biol CB 13:1910–1915.

Tanis JE, Bellemer A, Moresco JJ, Forbush B, Koelle MR (2009) The potassium chloride
cotransporter KCC-2 coordinates development of inhibitory neurotransmission and
synapse structure in Caenorhabditis elegans. J Neurosci Off J Soc Neurosci 29:9943–
9954.

Tanis JE, Moresco JJ, Lindquist RA, Koelle MR (2008) Regulation of serotonin biosynthesis by
the G proteins Galphao and Galphaq controls serotonin signaling in Caenorhabditis
elegans. Genetics 178:157–169.

Topalidou I, Keller C, Kalebic N, Nguyen KCQ, Somhegyi H, Politi KA, Heppenstall P, Hall DH,
Chalfie M (2012) Genetically separable functions of the MEC-17 tubulin
acetyltransferase affect microtubule organization. Curr Biol CB 22:1057–1065.

Trent C, Tsuing N, Horvitz HR (1983) Egg-laying defective mutants of the nematode
Caenorhabditis elegans. Genetics 104:619–647.

Waggoner LE, Dickinson KA, Poole DS, Tabuse Y, Miwa J, Schafer WR (2000) Long-term
nicotine adaptation in Caenorhabditis elegans involves PKC-dependent changes in
nicotinic receptor abundance. J Neurosci Off J Soc Neurosci 20:8802–8811.

Waggoner LE, Zhou GT, Schafer RW, Schafer WR (1998) Control of alternative behavioral
states by serotonin in Caenorhabditis elegans. Neuron 21:203–214.

Weinshenker D, Garriga G, Thomas JH (1995) Genetic and pharmacological analysis of
neurotransmitters controlling egg laying in C. elegans. J Neurosci Off J Soc Neurosci
15:6975–6985.

Weinshenker D, Wei A, Salkoff L, Thomas JH (1999) Block of an ether-a-go-go-like K(+)
channel by imipramine rescues egl-2 excitation defects in Caenorhabditis elegans. J
Neurosci Off J Soc Neurosci 19:9831–9840.

White JG, Southgate E, Thomson JN, Brenner S (1976) The structure of the ventral nerve cord
of Caenorhabditis elegans. Philos Trans R Soc Lond B Biol Sci 275:327–348.

White, J.G., Southgate, E., Thomson, J.N., Brenner, S. (1986) The structure of the nervous
system of the nematode Caenorhabditis elegans. Phil Trans R Soc 314:1–340.

Williams SL, Lutz S, Charlie NK, Vettel C, Ailion M, Coco C, Tesmer JJG, Jorgensen EM,
Wieland T, Miller KG (2007) Trio's Rho-specific GEF domain is the missing Galpha q
effector in C. elegans. Genes Dev 21:2731–2746.

Zhang M, Chung SH, Fang-Yen C, Craig C, Kerr RA, Suzuki H, Samuel ADT, Mazur E,
Schafer WR (2008) A self-regulating feed-forward circuit controlling C. elegans egg-
laying behavior. Curr Biol CB 18:1445–1455.

Zhang M, Schafer WR, Breitling R (2010) A circuit model of the temporal pattern generator of
Caenorhabditis egg-laying behavior. BMC Syst Biol 4:81.

Table 1: Strains used in this study

Strain	Feature	Genotype	Figures
LX1832	Strain for transgene production	*lite-1(ce314) lin-15(n765ts)* X	1-6
LX1836	HSN Channelrhodopsin	*wzIs30* IV; *lite-1(ce314) lin-15(n765ts)* X	3
LX1918	vulval muscle GCaMP5, mCherry	*lite-1(ce314) vsIs164 lin-15(n765ts)* X	1, 2, 3
LX1932	HSN Channelrhodopsin, vulval muscle GCaMP5, mCherry	*wzIs30* IV; *lite-1(ce314) vsIs164 lin-15(n765ts)* X	3
LX1938	No HSNs, vulval muscle GCaMP5, mCherry	*egl-1(n986dm)* V; *lite-1(ce314) vsIs164 lin-15(n765ts)* X	4
LX1986	uv1 GCaMP5, mCherry	*vsIs177*; *lite-1(ce314) lin-15(n765ts)* X	5
LX1960	VC GCaMP5, mCherry	*vsIs172*; *lite-1(ce314) lin-15(n765ts)* X	1, 2
LX1970	HSN Channelrhodopsin, VC GCaMP5, mCherry	*wzIs30* IV; *vsIs172*; *lite-1(ce314) lin-15(n765ts)* X	3
LX2004	HSN GCaMP5, mCherry	*lite-1(ce314), vsIs183 lin-15(n765ts)* X	1, 2, 6
LX2038	*lgc-55* null mutant HSN GCaMP5, mCherry	*lgc-55(tm2913)* V *lite-1(ce314) vsIs183 lin-15(n765ts)* X	6
MIA3	VC Channelrhodopsin	*keyIs3*; *lite-1(ce314) lin-15(n765ts)* X	3
MT13113	*tdc-1* null mutant; no tyramine biosynthesis	*tdc-1(n3419)* II	6
N2	Bristol strain	wild type	6
OH313	*ser-2* null mutant	*ser-2(pk1357)* X	6
VC125	*tyra-3* null mutant	*tyra-3(ok325)* X	6
QW89	*lgc-55* null mutant	*lgc-55(tm2913)* V	6
LX2096	vulval muscle mCherry	*vsIs191; lin-15(n765ts)* X	
LX2081	pan-neuronal TagRFP	*unc-119(ed3)* I; *otIs356*	
LX2137	vulval muscle mCherry, *lgc-55::gfp*	*vsIs191*; *vsEx791*	6
LX1330	*lgc-55* mutant expressing GFP in HSN from *tph-1* promoter	*lgc-55(tm2913)* V; *lin-15(n765ts)* X *vsEx557*	6
LX1329	*lgc-55* mutant expressing LGC-55 in HSN from *tph-1* promoter	*lgc-55(thm2913)* V; *lin-15(n765ts)* X *vsEx558*	6

Figure 1. Cell-specific reporters of activity in the *C. elegans* egg-laying behavior circuit.

(**A**) Schematic of the circuit. HSN (green) and VC (blue) motor neurons synapse onto the vm2 muscle postsynaptic termini (center of schematic). The uv1 neuroendocrine cells (pink) extend processes (grey) along the vulval slit and vm2 postsynaptic terminus. (**B-E**) Individual video

frames of the GCaMP5:mCherry fluorescence ratio showing active state Ca^{2+} transients in

HSNs (**B**), VCs (**C**), and vulval muscles during twitching (**D**) and egg-laying behaviors (**E**).

Arrowheads, HSN and VC presynaptic termini; asterisks, cell bodies; scale bar, 10 μm. (**F**) 30

minute recordings of HSN, VC, and vulval muscle activity (left panel), showing distinct active

(yellow) and inactive (grey) egg-laying behavior states, with expanded timescale of one active

state at right. Arrowheads show egg-laying events. (**G**) Scatter plots and median HSN, VC, and

vulval muscle (vm) inter-transient intervals during egg-laying inactive (–, filled circles) and

active (+, open circles) states. Asterisks indicate significant differences (p<0.0001). (**H**)

Relationship between Ca^{2+} transient amplitude and egg release. Scatter plots and medians of

normalized amplitude with (+; open circles) and without (–; closed circles) egg release. Also

shown is the percent of total transients that accompanied egg release. (**I**) Timing of HSN, VC,

and vulval muscle Ca^{2+} transients and egg release. Shown at top is a curve of the median of

Ca^{2+} from HSN (green), VC (blue), and vulval muscles (orange) from normalized ΔR/R traces

(with the peak Ca^{2+} set to 100%) synchronized to the moment of egg release (0 s, arrowhead

and dotted line). Bars indicate 20% change in median GCaMP/mCherry ratio. The timing of the

HSN Ca^{2+} peak is significantly different from that of the VCs and vulval muscles (p<0.0001;

n≥31 egg-laying events). Shown at bottom is a trace of median vulval muscle size. Bar shows

a 5% change in median object size based on mCherry fluorescence.

Figure 2. HSN, VC, and vulval muscle activity is phased with animal locomotion.

(**A**) The position of the vulva within a sinusoidal locomotor body bend at the moment a Ca^{2+} transient peaked was used to assign a body bend "phase", in units of degrees, for each Ca^{2+} transient observed in HSN (**B**), VC (**C**), and vulval muscle (**D**) for the recordings analyzed in Figure 1. The fraction of transients observed in each of eight 45° bins was plotted on a circular histogram with ventral relaxation at 180°, and ventral contraction at 0/360°. Ca^{2+} transients that resulted in egg release had significantly different phasing from those that did not (HSN, p=0.0058; VC, p=0.001; vulval muscles, p=0.0071; Mann-Whitney test). The phasing of HSN, VC, and vulval muscle transients is significantly different from a random distribution (R, dashed line at 0.125; p<0.05) except for those HSN transients that did not result in egg release.

Figure 3. Optogenetic activation of HSN neurons initiates the egg-laying active state, and optogenetic activation of VC neurons slows locomotion.

(**A**) Animals expressing Channelrhodopsin-2 (ChR2) in the HSNs and GCaMP5/mCherry in the VC neurons were grown in the presence (+ATR; top, blue) or absence (–ATR; bottom, black) of all-*trans* retinal. 489 nm laser light was used to simultaneously stimulate ChR2 activity and excite GCaMP5 fluorescence during the entire recording. Arrowheads indicate egg-laying events. (**B**) The same experiment as (A) except that GCaMP5/mCherry were expressed in the

vulval muscles rather than VCs. (**C**) Brief reduction of animal speed during optogenetically-

induced egg laying. The egg-laying active state in animals expressing ChR2 in the HSN

neurons was induced with 30 s exposure to blue light. Average speed (black trace) about each

egg-laying event (0 s) was calculated from the worm centroid position, and the grey area

shows the 95% confidence interval. Average speed at the moment of egg laying release (0 s)

is reduced compared to 1 s before or after (p <0.001). See also Movie 4. (**D**) Activation of the

VC neurons slows locomotion but fails to induce egg laying. Animals expressing ChR2 in the

VCs were grown in the presence (+ATR; red) or absence (–ATR; black/grey) of all-*trans* retinal,

and worm length (upper photos and graph) was determined. Micrographs are still images from

a representative animal at indicated time points (see Movie 5). Inset arrows indicate regions of

the head and tail that remain mobile after VC activation. Scatter plots of body bends (**E**) and

egg laying (**F**) from the same recordings shown in D. There was a significant change in worm

length and body bends, but not egg laying, during blue light in animals grown on ATR (red;

p<0.0001; n=17 animals). Positive control animals expressing ChR2 in the HSNs (green)

showed a significant increase in egg laying during blue light (p<0.0001; n≥14 animals).

Figure 4. Active states require egg production but not the HSNs.

(**A**) Vulval muscle activity in wild-type animals either untreated or after sterilization with floxuridine (FUDR). Left panel shows 30-minute recordings with the grey underlined regions

expanded at right. (**B**) Vulval muscle activity in *egl-1(n986dm)* mutants lacking HSNs. (**C-D**)

Scatter plots of vulval muscle inter-transient intervals (**C**) and amplitudes (**D**). Line indicates

median, error bars indicate quartiles, and asterisks indicate significant differences ($p<0.01$). (**E**)

Vulval muscle transients are wider in animals lacking HSNs. Cumulative distribution plots of

Ca^{2+} transient peak widths; asterisk indicates $p<0.0001$ (Mann-Whitney test). **(F, G)** Vulval

muscle Ca^{2+} transient phasing after sterilization resembles that of the inactive state. The

relative timing of peak vulval muscle Ca^{2+} transients during twitching (red), egg-laying (yellow),

and after FUDR treatment (blue) are plotted as in Figure 2. Twitch, egg-laying, and FUDR Ca^{2+}

transient timings were significantly different from each other and from a random distribution (R,

dashed line at 0.125; $p<0.05$). (**F**) Transients from 14 untreated and 5 FUDR-treated wild-type

animals. (**G**) Transients from 23 untreated and 10 FUDR-treated *egl-1(n986dm)* animals.

Figure 5. The tyraminergic uv1 neuroendocrine cells are mechanically deformed and activated by egg laying.

(**A**) Fluorescence micrographs of uv1 showing GCaMP5, mCherry, and the GCaMP5/mCherry ratio before egg laying, during egg passage through the vulva, and after egg release. Times of movie frames in seconds are at top, and white scale bar is 10 μm. Arrowheads in mCherry micrographs indicate position of uv1 cells as they are deformed by egg passage, the dotted line indicates position and opening of the vulva. (**B**) GCaMP5/mCherry ratio ($\Delta R/R$) trace of uv1 activity, including an egg-laying active state. Egg-laying events are indicated by arrowheads. (**C**) Scatter plots of uv1 inter-transient intervals during the inactive (closed circles)

and active (open circles) egg-laying behavior states. Line indicates median, error bars indicate

quartiles, and asterisks indicate significant differences (p<0.0001). (**D**) Scatter plots and

medians of normalized amplitude with (+; open circles) and without (−; closed circles) an

accompanying egg release. Error bars indicate quartiles, and asterisks indicate significant

differences (p<0.0001, Mann-Whitney test). (**E**) uv1 Ca^{2+} transients follow egg-laying events.

Normalized traces of median Ca^{2+} in HSN (green) and VC (blue) from Figure 1 are compared

to uv1 (purple), synchronized to the moment of egg release (0 s, arrowhead and dotted line).

Bars indicates 20% change in median GCaMP/mCherry ratio. Asterisk indicates the uv1 Ca^{2+}

peak is significantly later than the HSN and VC peaks (p <0.0001; n=27 egg-laying events from

13 animals).

Figure 6. Tyramine inhibits egg laying, in part, through LGC-55 receptors on the HSNs.

(**A**) Exogenous tyramine inhibits egg-laying behavior and requires the LGC-55 receptor.

Scatter plots and means showing the number of eggs laid after 30 minutes by wild type (grey),

tdc-1(n3419) tyramine biosynthetic enzyme mutant (blue), and *lgc-55(tm2913), ser-2(pk1357),*

and *tyra-3(ok325)* tyramine receptor mutant animals (orange, red, and purple, respectively) on

plates without (-, closed circles) or with (+, open circles) 15 mM tyramine. Error bars indicate

95% confidence intervals, and asterisks indicate significant responses (*p*<0.01) while n.s. is

not significant (p>0.05). (**B**) Dose-response curve of tyramine inhibition on egg laying in wild-

type and *lgc-55* mutant animals. Asterisks indicate significant differences at 10 mM (p<0.0001)

and 20 mM tyramine (p=0.0164). (**C**) LGC-55 is expressed in HSN, uv1, and vulval muscles.

Expression of LGC-55 tagged with GFP compared to vulval muscle mCherry was visualized

using confocal microscopy; scale bar is 10 μm. GFP is localized to the HSN and uv1 cell

bodies (arrows) as well as the ventral tips of the vulval muscles. (**D**) Re-expression of LGC-55

in HSN restores tyramine inhibition of egg laying. Scatter plots of eggs laid by *lgc-55* mutants

expressing either GFP or *lgc-55* in the HSN from the *tph-1* promoter. Asterisk indicates

significant differences in egg laying (p=0.0029). (**E**) Loss of LGC-55 increases HSN activity.

Representative Ca^{2+} ratio traces show HSN activity in wild type (top, grey) and *lgc-55* mutant

animals (bottom, blue) during the active state. (**F**) Scatter plots and medians of HSN inter-

transient intervals in wild type and *lgc-55* mutant animals during the egg-laying inactive (–,

closed circles) and active (+, open circles) states. Asterisks indicate statistically significant

differences (p<0.0001; n≥7 animals per genotype).

Figure 7. Working model of how circuit connectivity, signaling, and activity contribute to the observed rhythms that accompany the active and inactive egg-laying behavior states. VA and VB motor neuron synapses release acetylcholine (ACh) that rhythmically excites the body wall muscles (bwm) and vm1 vulval muscles during each locomotion body bend, every ~10 seconds. Eggs are produced from each gonad arm every 10 minutes, and we propose the accumulation of 2-3 eggs mechanically excites the uterine muscles (um) facilitating exit from the inactive state. Sustained bursts of HSN command motor neuron activity trigger serotonin release that drives ~2 minute active states. HSN also makes synapses and gap junctions with the cholinergic VC motor neurons, and VC makes synapses and gap junctions with the vm2 muscles whose contraction allows egg release. Passage of eggs mechanically activates the uv1 cells triggering release of tyramine and neuropeptides, which inhibit HSN activity to terminate the active state. See text for further details.

Movie 1. Ratio recording of HSN Ca^{2+} transients during the egg-laying active state. High Ca^{2+} is indicated in red while low calcium is in blue. Large panel is an expanded view showing the freely-moving animal that has been contrast enhanced to make the worm and its laid eggs visible. Inset is cropped to a small area containing the HSN, and stabilized to remove movement. Text labels indicate when Ca^{2+} transients and egg release occur.

Movie 2. Ratio recording of VC Ca^{2+} transients during the egg-laying active state. High Ca^{2+} is indicated in red while low calcium is in blue. Contrast is enhanced to make the worm visible, although the laid eggs are not easily visible. Text labels indicate when vulval muscle (vm) twitches or egg-laying contractions occur, which result in small or large displacements of the VC4 and VC5 cell bodies, respectively.

Movie 3. Ratio recording of vulval muscle Ca^{2+} transients during the egg-laying active state. High Ca^{2+} is indicated in red while low calcium is in blue. Large panel is an expanded view showing the freely-moving animal that has been contrast enhanced to make the worm and its laid eggs visible. Inset is cropped to a small area containing the vulval muscles, and stabilized to remove movement. Text labels indicate when Ca^{2+} transients and egg release occur.

Movie 4. Activation of the HSNs using Channelrhodopsin-2 induces the egg-laying active state. Animals were recorded for a total of 90 seconds with continuous blue light stimulation beginning at 30 s and ending at 60s, during which five eggs are laid. Recording is sped up 3-fold.

Movie 5. Activation of the VCs using Channelrhodopsin-2 induces animal paralysis and shortening of body length. Animals were recorded for a total of 90 seconds with continuous blue light stimulation beginning at 30 s and ending at 60s. Recording is sped up 3-fold.

Movie 6. Ratio recording of uv1 Ca^{2+} transients during the egg-laying active state. High Ca^{2+}

is indicated in red while low calcium is in blue. Large panel is an expanded view showing the

freely-moving animal. Inset is cropped to a small area containing the uv1s, and stabilized to

remove movement. Microscope is focused to image the two uv1 cells on the right side of this

animal, while the two additional uv1 cells on the left side are out of focus and not visible. Text

labels indicate when egg releases occur, each of which is followed by a uv1 Ca^{2+} transient.