Investigation of the sexual reproduction strategy of the Périgord black truffle (*Tuber*

***melanosporum* Vittad.) revealed trioecy**

Herminia De la Varga[1], François Le Tacon[1], Mélanie Lagoguet[1], Flora Todesco[1], Torda

Varga[1*], Igor Miquel[1], Dominique Barry-Etienne[2], Christophe Robin[3], Fabien Halkett[1],

Francis Martin[1], Claude Murat[1,4]

[1]INRA, Université de Lorraine, UMR1136 Interactions Arbres-Microorganismes, Laboratoire

d'Excellence ARBRE, F-54280 Champenoux, France

[2]ALCINA sarl, 10 rue des Amaryllis, F-34070 Montpellier, France

[3] INRA, Université de Lorraine, UMR 1121 Laboratoire Agronomie-Environnement, Nancy-

Colmar, Vandoeuvre-les-Nancy 54518, France

[4]Corresponding author:

Claude Murat

UMR1136 INRA/Université de Lorraine. "Interactions Arbres/Micro-organismes"

Laboratoire d'Excellence ARBRE. INRA Nancy. 54280 Champenoux. France

Tel: +33 (0)3 83 39 41 27; Fax: +33 (0)3 83 39 40 69; Email: claude.murat@nancy.inra.fr

Running title: Black truffle sexual reproduction strategy

[*] Current address: Synthetic and Systems Biology Unit, Biological Research Centre,

Hungarian Academy of Sciences, Temesvári krt. 62, H-6726 Szeged, Hungary.

Summary

Fungi display a rich variety of sexual reproduction systems and are therefore good models to

investigate sex evolution. Moreover, understanding the investment in sexual reproduction of

edible fungi is a critical challenge for improving their cultivation. The Périgord black truffle

(*Tuber melanosporum* Vittad.) is a heterothallic ascomycete associated with trees through

ectomycorrhizas and forming highly prized ascocarps. The aim of this study was to unravel

the sexual reproduction strategy (hermaphroditic versus male/female specialization) of *T.*

melanosporum in a truffle orchard by disentangling the contribution of female and male

genotypes to the formation of ascocarps in a five-year investigation. Few genotypes

were hermaphrodites, co-occurring with numerous genotypes behaving only as female or

male, revealing trioecy. The genetic diversity of the male genotypes was higher than female

diversity, suggesting for male elements a higher recruitment from ascospores. Most of the

female and male genotypes were transitory (present only one year), whereas some genotypes

persisted for several years: female-fertile genotypes as mycorrhizas, and male-fertile

genotypes as soil free-living mycelium. Contrary to other ascomycetes, a high number of

female-fertile only genotypes was found. We hypothesized that the mycorrhizal life style

favours female-fertile strains and therefore that the life strategy influences fungal sexual

strategy.

Introduction

In fungi, sexual lifestyles are rich with species displaying typical mating between compatible strains (i.e. heterothallism) whereas others are able to reproduce sexually in the absence of a compatible partner (i.e. homothallism). The life cycle of the ascomycetes species is mainly haploid and heterothallic species form male elements (e.g. conidia, sexual spores or hyphae) (Glass and Kuldau, 1992; Leslie and Klein, 1996), and female elements (ascogonia) are usually much more complex. In theory, one strain of a heterothallic ascomycete is hermaphrodite or monoecious (Esser and Kuenen, 1967), meaning that it can form either male or female elements (Glass and Kuldau, 1992). A haploid (hence self-sterile) hermaphrodite propagating by asexually producing spores or reproducing sexually was proposed to be an idealized heterothallic ascomycete (Leslie and Klein, 1996). Hermaphroditism is not always maintained as demonstrated for *Hypomyces solani* (*Fusarium solani*) (Hansen and Snyder, 1943), and the occurrence of strains forming only male or female elements (i.e. dioecy) was observed in ascomycetes (Benjamin, 1950). Trioecy, the co-occurrence of hermaphrodite and dioecy, was observed for *Triceromyces* species (Benjamin, 1986) and *Fusarium fujikuroi* (Leslie, 1995). Few examples of trioecy have been described in fungi, although it is widespread in animals and plants (Weeks et al., 2006; Chaundhuri et al., 2011; Armova-Zvuloni et al., 2014; Joseph and Murthy, 2014; Mirski and Brzosko, 2015). Moreover, the relative abundance of hermaphrodite, male/female producing strains in the field for mycorrhizal (plant symbiotic) ascomycete species has never been investigated.

Beside the theoretical and evolutionary interest of investigating fungal sexual reproduction, it is also critical for overcoming the cultivation of species forming edible mushrooms. This is particularly true for the most famous edible fungi (such as Boletes, Chanterelles, and truffles), whose growth and sexual reproduction are strictly linked to host plants through a symbiotic

association called ectomycorrhiza (Murat et al., 2008). Cultivation of the black Perigord

truffle (*Tuber melanosporum* Vittad.) started forty years ago thanks to inoculated seedlings

(Chevalier and Grente, 1979); however, the control of its sexual reproduction cycle is not

achieved (Le Tacon et al., 2015). *Tuber melanosporum* is a heterothallic ascomycete species

(Martin et al., 2010; Rubini et al., 2011b) with the female mycelium found in the root system

as ectomycorrhiza and forming the structure (i.e. peridium and gleba) of the ascocarps (Fig.

1). The origin of male elements is unknown, but it was suggested to be conidia, ascospores, or

soil mycelium (Fig. 1; Le Tacon et al., 2015). The small-scale genetic structure of female

genotypes was characterized (Rubini et al., 2011a; Murat et al., 2013), although male

genotypes were never analysed under field conditions. Female genotypes were detected in the

root system as this is where they tended to aggregate according to their mating type (Murat et

al., 2013). The same result was found recently in one natural production site of *T. aestivum* in

Germany (Molinier et al., 2016). Interestingly, a high density of ascocarps was not observed

for either species in the contact zone of the opposite mating types suggesting a more complex

system of sexual reproduction.

The aim of this study was to unravel the sexual reproduction strategy (trioecious versus

dioecious or hermaphroditic) of *T. melanosporum* in the field. We used a five-year sampling

of *T. melanosporum* ascocarps, ectomycorrhizas and soil in one orchard and established the

genotypes of both female and male elements. The investigation of sex strategy in truffles

provides new insights into the sex specialisation in fungi, but more importantly, it improves

our understanding of sexual reproduction and its impact on the eukaryotic tree of life.

Results

Ascocarps and ectomycorrhiza samplings

A total of 241 ascocarps were harvested during five consecutive seasons of truffle production

(from November to March of the following year) from 2010/2011 to 2014/2015 (Table 1; Fig.

S1). The ascocarps were randomly distributed around the productive trees and could be also

found near the trunk, at 4 to 5 m from it, in the zone of extension of the roots, and between it

(Fig. S1). The productive area under each tree extended slowly over five years.

A total of 475 ectomycorrhizal root tips were collected (205 in 41 root samples, with 5 *T.*

melanosporum morphotype each, in 2011 and 270 in 45 root samples, with 6 *T.*

melanosporum morphotype each, in November 2014) (Fig. S1). Only 37 ectomycorrhiza

samples did not amplified with the *T. melanosporum* specific primers, and the presence of *T.*

melanosporum ectomycorrhizas was not confirmed only in five root samples (two in 2011 and

three in 2014). For genotyping, only one *T. melanosporum* ectomycorrhiza per root sample

was selected.

Genotyping of female elements (gleba and ectomycorrhiza)

The genotypes of ascocarp female elements (i.e. gleba) were successfully obtained for 238 out

of the 241 ascocarps (Table 1 and Table S1). A total of 73 female multilocus genotypes

(called hereafter genotypes) with significant Psex value were found during the five years

(Table 1, Tables S1 and Table S2). Twelve genotypes were present in more than one season

(Table S1). These genotypes accounted for 26% to 94% of the ascocarps for seasons 3 and 1

respectively, for a total of 154 truffles (64 % of the total harvested ascocarps over the five

seasons). One persistent genotype (R002) was found throughout the season under the same

tree (F11, area 3), representing 15% of the total ascocarps harvested (Fig. S2 and Table S1).

Another genotype (R021) fructified under the A11 tree (in area 1) for three seasons and

represented 16 % of the total ascocarp harvested (Fig. 2 and Fig. S1). The maximum size

(clonal subrange) of the female genotypes was 22.36 m, corresponding to the R002 genotype

found in areas 1 and 3 of the truffle orchard (Table 1 and Fig. S2). Most female genotypes

were found in only one ascocarp (Table S1).

A total of 117 ectomycorrhizas were genotyped, representing 9 and 14 genotypes for 2010

and 2014 samplings, respectively (Table 1 and Table S1). Five genotypes (R001, R002, R003,

R004 and R007) were found in both ectomycorrhizal samplings in 2010 and 2014 (Table S1).

The maximum ectomycorrhiza genotype size was 19.02 m for R021 (Table 1 and Fig. 2).

Eighty-six percent of ectomycorrhiza genotypes were detected also as female elements in

ascocarps, represented in 16 genotypes with significant Psex value (Table S1). Similarly, the

female element of the ascocarps shared the same genotype with that of the nearby

ectomycorrhizas.

When considering only the mating type locus, the aggregation index (Ac) varied from 0.42 to

0.57 for ascocarp female element (gleba) and from 0.40 to 0.96 for ectomycorrhizas (Table 1).

In all cases, the Ac was significant (P-value < 0.05), indicating that genotype of the same

mating type tend to aggregate. In the orchard, large patches from 5 to 20 m^2 of different

genotypes of the same mating type for the root colonization were observed (Fig. S3). A single

tree can be colonized by one patch (E10 and B11) or by two contiguous patches of opposite

mating types (F10, F11, D11, A11, and A12). Tree D11 harboured maternal strains and

ectomycorrhizas of *MAT1-2* with the exception of one ectomycorrhiza sampled in square F5

formed by *MAT1-1* mycelium (Fig. S3 and Table S1). This genotype was not detected in male

strains (see below). At the contact zone between patches of opposite mating types, no

differences in the ascocarp density were observed when compared to density within the patch

(Fig. S1 and Fig. S3).

Zygote analysis and genotyping of male elements

We successfully genotyped 206 zygotes out of the 241 ascocarps (86.5 %; Table S1). By

subtracting the female haploid contribution, we were able to reconstruct all male genotypes. A

total of 138 male genotypes were found (Table 1 and Table S1). A total of 9 zygotes (4%)

were homozygous for all microsatellite loci (i.e. only mating type locus is heterozygous). In

zygotes, a significant deficit of heterozygosity was observed with F_{IS} values of 0.02 and 0.28

for seasons 1 and 3, respectively (Table 1). The relationship of female and male genotypes in

zygotes was investigated by kinship coefficient calculation. The kinship coefficient varied

from -0.5 to 1.5 (Fig. S4).

Most of the male genotypes (75%) were transitory, i.e. found only in one ascocarp (Table S1).

Only 8 persistent male genotypes with significant Psex values were found. These persistent

genotypes produced 21 % to 37 % of the ascocarps (Table 1). Sixteen male genotypes were

found during several seasons and 5 (only one genotype with significant Psex) were found

under different trees (Table S1). One genotype (R102) was detected for four seasons under

A11 where it fertilized 8 % of the ascocarps (Fig. S5). The maximum male genotype size was

22.64 m for genotype R102 with detection of most of them in one single ascocarp (Table 1).

Identification of hermaphrodite strains

The hermaphrodite strains are those detected as both female and male elements. In our

dataset, 8 genotypes (3.9%) were detected as female and male (detected in the gleba and/or

ectomycorrhizal tissues and in the zygotes). However, only 3 genotypes (1.5 % of the total

number of female and male genotypes) were supported by significant Psex values and thus

considered true hermaphrodites (Table 1 and Fig. 2). The R021 genotype was the most

frequent female genotype (see above), and it has been also detected in 5 ascocarps as male

element and once as ectomycorrhiza (Fig. 2A). The R060 genotype was found 5 times as

female genotype, once as male strains and once as ECM (Fig. 2B). Finally, the R068 was

detected 3 times in female strains, once in male strains, and once as ECM (Fig. 2C).

Conversely, 135 genotypes were detected only as male-fertile (64%) and 70 genotypes were

only female-fertile (33%).

Small-scale genetic structure for both female and male elements

According to the season, the Simpson's diversity varied from 0.81 to 0.98, 0.98 to 0.99, and

0.79 to 0.99 for female, male and ectomycorrhizal samples, respectively (Table 1). The inter-

individual diversity varied from 0.52 to 0.91 and 0.83 to 0.97 for female and male elements,

respectively (Table 1). Both indices were similar but they were slightly higher for male

elements compared to female ones and ectomycorrhizas. The genetic structure was assessed

using isolation by distance (IBD), and the slope was used to investigate dissemination

capacities (Fig. 3). A significant genetic structure was detected for female and male elements

when all samples were used together or when using a culled dataset. When all samples were

considered, a higher slope was found for female versus male strains (0.125 versus 0.049),

indicating a higher dissemination capacity or higher density of male elements (Fig. 3). This

analysis, however, could be biased due to the repeated sampling of the same genotype. Using

the culled dataset the slope was even higher for female versus male strains but it was not

significant (Fig. 3).

Quantity of mating types mycelium in soil samples

In the 20 soil cores harvested in May 2015 (Fig. S1), both mating type mycelium were

detected in 16 soil samples (Table 2). In one soil sample, any of both mating types of

mycelium were detected while in 2 soil samples only *MAT1-1* was found, and in one soil

sample only *MAT1-2* was detected. The quantity of mycelium ranged from 0 to 7.4 mg of

mycelium per gram of soil and 0 to 40.49 mg of mycelium per grams of soil for *MAT1-1* and

MAT1-2, respectively. With two exceptions (squares E4 and E8 in area 1), the most frequent

mycelium corresponded to those of maternal strains and ectomycorrhiza.

Discussion

In this study, we characterized *in situ the* genotypes of female and male elements for *T.*

melanosporum by analysing both maternal and zygotic tissues of ascocarps. This involved

disentangling female and male contributions of all fruiting bodies collected in an orchard for

five years. We revealed a strong genetic structure for both elements and trioecy through the

coexistence in this population of hermaphrodite, female-fertile only and male-fertile only

strains. Moreover, equilibrium between vegetative propagation and sexual reproduction was

observed, but the factors conditioning the shift between both modes are still unknown. More

broadly, the maintenance of sexual reproduction in a natural population is an intriguing

question always discussed among evolutionists (Lehtonen and Kokko, 2014). Fungi are good

models to investigate sex evolution since they are particularly rich in their mode of sexual

reproduction. The recent progress in fungal genomics has allowed investigation of sexual

reproduction in species where it had not been previously described such as the cheese

ascomycete *Penicillium roqueforti* (Ropars et al., 2012). For mycorrhizal fungal species, the

mating type locus was described from genomic data for arbuscular endomycorrhiza

(*Rhizophagus irregularis,* Ropars et al., 2016), ectomycorrhizal basidiomycetes (*Laccaria*

bicolor, Niculita et al. 2008) and ascomycetes (*T. borchii*, Belfiori et al., 2015; *T. indicum*,

Belfiori et al., 2013 and *T. melanosporum*, Rubini et al., 2011b). However, the distribution of

mating type strains and sexual reproduction strategies (hermaphrodite versus dioecy) has not

yet been investigated, although mycorrhizal fungi play key roles in ecosystem functioning and

biogeochemical cycles (Smith and Read, 2008).

In *T. melanosporum*, the female elements are provided by the ectomycorrhizas that play the

role of gametophytes (Rubini et al., 2011a; Murat et al. 2013). Conversely, the origin of the

male elements was unknown (Le Tacon et al., 2015). For both female and male elements, we

observed similar features with the co-occurrence of small size genotypes, often detected

transitorily in only one ascocarp and large perennial genotypes. These results suggest that, for

both female and male elements, there is a mix of new genotype recruitment from ascospores

and perennial strains disseminated by vegetative propagation.

A strong genetic structure was found for both female and male elements, as evidenced by IBD

analysis. The IBD is influenced by spore dispersal. In fungi, two main strategies of dispersal

exist: 1) wind for epigeous and 2) animals for hypogeous species. This strong genetic

structure is characteristic of hypogeous fungi that are expected to have reduced spore

dispersal than epigeous (Kretzer et al., 2005). When considering culled data, similar slopes

for both male and female elements (0.048 and 0.051, respectively), suggest similar capacities

of dissemination for both sexes. Focusing on male elements, 16 perennial genotypes were

found. Among them, genotype R102 was found in 18 ascocarps (8.7%) over 4 years (Fig. S5).

Only the three male genotypes corresponding to the true hermaphrodite genotypes were

present on ectomycorrhizas. This can be due to a poor ability to form associations with the

host of male-fertile genotypes that probably survive as free-living mycelium in the soil or are

linked to non-ectomycorrhizal plants. Indeed, herbaceous plants can host truffle mycelium,

but the nature of this interaction (i.e. surface contact only, endophytism, endomycorrhiza) is

unknown (Gryndler et al., 2014). Moreover, we detected the presence of both mating types in

the soil mycelium in 16 out of 20 soil cores. This result demonstrated that mycelium of the

opposite mating type is present close to female strains in ectomycorrhiza. We can therefore

hypothesize that genotypes producing only the male elements (the so-called male strain)

survive in the soil through vegetative propagation. In many ascomycetes, conidia (i.e. asexual

spores) serve for vegetative propagation or as male elements (Nelson, 1996; Maheshwari,

1999). Urban *et al.* (2004) described the existence of anamorphous structures producing

mitotic conidia in soils where *T. borchii* and *T. oligospermum* ascocarps were present. Healy

et al. (2013) suggested that Pezizales mitospores, including *Tuber* mitospores, which failed to

form ectomycorrhizas, could act as spermatia. However, the question of whether conidia of

Tuber species act as spermatia remains unanswered and mitotic conidia have not yet been

observed in *T. melanosporum* orchards.

Male genotypes presented a higher genetic diversity and were less perennial than female ones,

suggesting that ascospore recruitment is more important for male than for female elements. It

is therefore tempting to hypothesize that most of the male elements originate from

germinating ascospores whose mycelium does not survive after sexual reproduction. Indeed,

sexual spores have been proposed as male elements for ascomycetes[2] and for basidiomycetes

(Nieuwenhuis et al., 2011). The inoculation of ascospores in order to improve the production

of ascocarps is a common practice (Olivier et al., 2012) but is totally empirical, as scientific

background was lacking. This practice is not recent since Ciccarello (1564), Bradley (1726)

and Buffon (1749) proposed methods to produce ascocarps by inoculating pieces of truffles

under mature trees. The role of ascospores was not directly demonstrated in this study, but the

possibility to trap truffle production in a few cm^2, as recently demonstrated (Bonneau, 2016),

open new perspectives to confirm the role of ascospores as male elements.

Hermaphroditism is the common rule in heterothallic ascomycetes ((Glass and Kuldau, 1992;

Nieuwenhuis and Aanen, 2012), and haploid self-sterile hermaphrodite has been proposed as

the optimal situation (Leslie and Klein, 1996). For *T. melanosporum* Rubini et al., 2011a;

Murat et al., 2013) and *T. aestivum* (Molinier et al., 2016) female strains are not randomly

distributed according to their mating types since they form large patches of different

genotypes in the root system, all sharing the same mating type. In our study, we confirmed

this spatial aggregation of genotypes of the same mating type in the root system. This

aggregation was stable over five years despite genotype turnover (Fig. S3). Interestingly, few

ascocarps were harvested in the contact zone of both mating types suggesting that 1) these

strains were not hermaphrodites; 2) all opposite mating type strains could not mate; or 3) they

had very limited dispersal capacities. We found three genotypes as true hermaphrodites since

they were identified as both female and male elements. On the contrary, most genotypes were

identified as female-fertile or male-fertile only, suggesting a specialisation of the partners in

one sex. In the literature, the terms male-sterile and female-sterile are commonly used, but in

our study based on field observations, we cannot definitively rule out that the female-fertile

and male-fertile strains are sterile for the opposite sex. Indeed, two hypotheses can be

proposed: 1) the capacity to produce male or female elements is lost due to mutations or 2)

genetic or epigenetic regulation impedes forming male or female elements. If mutations drive

the sex specialisation in *Tuber*, female-fertile and male-fertile strains are effectively sterile for

the opposite sex. In case of a regulation mechanism, sex specialization may be reversible.

In our population, female-fertile strains were relatively frequent (33% versus 64% of male-

fertile strains). This finding contradicts the model developed by Leslie and Klein[2] for

pathogenic and saprotrophic fungi. They hypothesised that male-sterile (i.e. female-fertile)

strains have reduced fitness for vegetative propagation, and therefore, they are not frequent. In

T. melanosporum the female-fertile strains were found in ectomycorrhizas and 17 % of these

were persistent, suggesting they dispersed by vegetative propagation. The female structures

are more complex than male structures and they require more energy. The necessity of

female-fertile strains to form ectomycorrhiza for carbon acquisition from the host tree could

favour their fitness in vegetative propagation explaining the difference with the Leslie and

Klein (1996) model. We may attempt to propose that the fungal life cycle (pathogen/

symbiotic/ saprotrophic) could influence the reproduction strategy.

To summarize, in our population a mix of few hermaphrodites with a majority of female-

fertile and male-fertile strains were identified suggesting trioecy. Trioecy is known in plants

(Joseph and Murthy, 2014; Mirski and Brzosko, 2015) and animals ((Weeks et al., 2006;

Chaudhuri et al., 2011). In fungi, trioecy was reported for the ascomycete Triceromyces

(Benjamin, 1986), and it can exist for *F. fujikuroi* (Leslie, 1995)*,* but it doesn't seem to be a

widespread situation. In animals, trioecy can occur when environmental conditions change or

when a species occupies a new habitat, leading to a transition from hermaphroditism to dioecy

or *vice versa* (Weeks et al., 2006). Trioecy is therefore a transitory status and in

Caenorhabditis elegans it seems evolutionarily unstable (Chaudhuri et al., 2011). In

heterothallic ascomycetes, hermaphroditism could be the ancestral status since it is expected

for most of the species (Glass and Kuldau, 1992). It is therefore tempting to hypothesize that

hermaphroditism has been lost in our *T. melanosporum* population in order to favour female-

fertile and male-fertile strains. In a theoretical analysis, Nauta and Hoekstra (1992) suggested

that in heterothallic populations, hermaphroditism could only be stable if the fitness of

ascospore formation is variable. For ascomycetes, 8 spores are expected in each asci but for *T.*

melanosporum the number of spores per asci varied from 1 to 4 (Chevalier et al., 2001). In

Neurospora crassa mutations leads to spore number reduction and some of these reduced the

female or male fertility (Leslie and Raju, 1985). There were no differences in the number of

spores per asci between hermaphrodite and female strains (data not shown) and currently it is

not possible to conclude that spore fitness influences the sexual reproduction strategy in

truffle. But the investigation of sexual reproduction of truffle species with 8 spores such as *T.*

panniferum or *T. regianum* (Chevalier et al., 2001) can allow addressing this hypothesis.

To conclude, trioecy with a low rate of hermaphrodite was observed in a *T. melanosporum*

population. Sexual dimorphism could therefore be more frequent in fungi than expected, and

progress in genome sequencing could allow its investigation. Indeed, in contrast to animals

and plants, dioecious fungi often are morphologically similar, and sexual dimorphism can be

detected only at genome level or at gene regulation level (Samils et al., 2013). Moreover, the

life strategy could influence the sexual reproduction mode, which has been poorly

investigated to date. The understanding of sexual reproduction of economically interesting

fungal species such as those producing edible mushrooms is a major issue for a better control

of their life cycle and cultivation.

Experimental procedures

Truffle orchard and sampling

Ascocarps and ectomycorrhizas were sampled at a long-term experimental site located at

Rollainville in north-eastern France. The study site is described in a study by Murat *et al.*

(2013). Samples and trees were identified by a letter and number and mapped on a grid of 1 m

x 1 m squares set up with camping pickets (Fig. S1). Three different grids were made to

identify 3 areas (areas 1, 2 and 3) that cover all the productive zones of the plantation (Fig.

S1).

As described in Murat *et al.* (2013)*,* the sampling started under trees F10, F11 and E10 (area

3) in 2010-2011 season and under the trees A11, A12, B11 (area 1) and D11 (area 2) in 2011-

2012 season (Fig. S1). The mature truffles were systematically harvested during the

production season with help from a well-trained dog, and at the time of harvest, they were

precisely mapped on the grid with 5 cm precision. The ascocarps were then washed to remove

soil particles and stored at -20°C for molecular analysis.

During this five-year interval, two ectomycorrhizal samplings were done. The first one was

done under F10/F11/E10 trees (area 3 in Fig. S1) in spring 2011, and results on that sampling

were published by Murat *et al.* (2013). The second one was done in November 2014, 45 root

samples were sampled all over the truffle orchard (Fig. S1). Root samples were harvested

from the first 10 cm of soil and mapped on the same grid used for the positioning of the

ascocarps. The ectomycorrhizas were carefully retrieved from the soil and washed in water

under a dissecting microscope. From each root sample, the *T. melanosporum* ectomycorrhizas

were identified as described by Zambonelli *et al.* (1993) and Rauscher *et al.* (1995), and

stored individually in microcentrifuge tubes at -20°C for molecular analyses.

In order to investigate the distribution and abundance of both mating types in the soil, 20 soil

cores were harvested in May 2015 (Fig. S1) at 10-15 cm depth. All plant debris, stones and

roots were discarded from the soil samples, and samples were kept at -20°C for DNA

extractions.

DNA extractions

Genomic DNA from the gleba (i.e. inner part of the ascocarps) and ectomycorrhizal tips were

extracted by using the DNeasy Plant Mini Kit (Qiagen SA, Courtaboeuf, France), following

the manufacturer's instructions.

From each ascocarp, a mixture of spores was isolated and their DNA was extracted as

described below. Thin slides of each ascocarp were put onto a layer of water in a Petri dish to

let spores be released into the water in order to isolate the pool of spores. The liquid was

collected in a 1.5 mL tube and centrifuged at 14,000 rpm for 5 minutes. The supernatant was

then discarded in order to obtain a mixture of asci and spores. DNA from the isolated mixture

of spores from each ascocarp was extracted as described in Rubini *et al.* (1999) with some

modifications. First, to each pool of spores, we added 300 mL of NTE buffer (Tris-HCl 200

mM, NaCl 250 mM EDTA 25 mM), two tungsten beads, and then disrupted the spores with a

Tissue Lyser (Qiagen) for 10 minutes at 30 J. The tubes were then centrifuged at 14,000 rpm

for 10 minutes, and the recovered supernatant was added to a new tube. Thirty microliters of

NaAc (3M) and 330 μL isopropanol were added, and those tubes were mixed and centrifuged

again at 14,000 rpm for 10 minutes. After discarding the supernatant and cleaning the pellet

with 200 μL of ethanol (70%), we recovered the pellet (DNA) in 50 μL of TE. DNA extracts

were stored at -20°C.

Total DNA from soil samples was extracted using the Power Soil® DNA extraction Kit

(MoBio, Laboratories, Carlsbad, CA) following manufacturer's protocol and stored at -20°C

for molecular analysis.

Molecular genotyping

All extracted DNA samples were amplified using species-specific *T. melanosporum* primers[48]

in order to check the species and DNA quality.

All of the samples in which *T. melanosporum* identity was confirmed were genotyped by

using primer pairs corresponding to ten microsatellite markers (Tm16_ATA12,

Tm241_TAA17, Tm2_TAT15, Tm98_TAT15, Tm112_TAT19, Tm9_ATCA12,

Tm1_ATTG18, Tm75_GAAA14, Tm22_CCTCAT17 and Tm269_TGTTGC15) as described

by Murat *et al*. (2011; 2013). The genotyping was done by the INRA plateform Gentyane

(Clermont-Ferrand). The *T. melanosporum* mating types of all the samples were analysed

using specific primers for either the *MAT1-1-1* or the *MAT1-2-1* genes using the PCR

conditions described by Rubini *et al*. (2011b). Hereafter, according to Rubini *et al*. (2011a),

the mating types are termed *MAT1-1* and *MAT1-2*.

From the mix of DNA spores, we obtained zygotic genotypes from which the male genotype

was deduced by subtraction of the corresponding female genotype.

The soil mycelium of both mating types was quantified by quantitative real time PCR (qPCR)

using a protocol developed from the international patent (EP2426215 A1) in a confidential

research program with ALCINA sarl (Montpellier, France). qPCR reactions were carried out

with a StepOne Plus[TM] Real-Time PCR System machine provided with the StepOne software

v. 2.3 (Life Technologies, Carlsbad, CA). Two standard curves (R^2=0.99; Eff = 96.99% and

R^2=0.99; Eff = 99.67% for *MAT 1-1* and *MAT 1-2,* respectively) were obtained, as described

in Parladé *et al.* (2013), by mixing a known amount of soil harvested in a cereal field close to

the truffle orchard (in which the absence of *T. melanosporum* was confirmed) with a known

amount of fresh immature ascocarp of *T. melanosporum* belonging to one or the other mating

type. DNA from the mixture was extracted as all the other soil samples and serial tenfold

dilutions were done to obtain a standard curve for each mating type. Absolute quantification

of mating types in soil samples was obtained by interpolating their threshold cycle (Ct) values

on the corresponding standard curve.

Data analyses

All the samples with null alleles were discarded from our analyses. The multilocus genotype

identification was achieved using the software MLGsim 2.0 (2003). For genotypes sampled

more than once, the probability of independent occurrence (Pgen) and the PSex probability of

arising by chance (Psex) were calculated. The P-value for testing the significance of the PSex

for each genotypes was estimated using 1000 simulations. When the PSex values fell below

the P-value (0.05), it was concluded that identical genotypes originated from the same genet

(clonal multiplication).

The clonal diversity (R), the Simpson's diversity index modified for finite sample sizes (D),

the clonal subrange, and the aggregation index (Ac) for mating types were estimated using the

RClone package (2016) for R software, based on the GenClone software package v2.0 (2007).

The inter-individual diversity (He=1-Qinter) and isolation by distance (IBD) analysis were

obtained with GenePop software v4.2 online (1995; 2008). The significance of IBD was

tested using Mantel test with 10,000 permutations. All values were calculated for all the data

and also separately for female, male, and ECM data. A culled dataset was constructed to

reduce the bias due to sampling the same genotype several times. For each season, the

isobarycentre was considered for samples sharing the same genotype and having significant

Psex. If two samples with the same genotypes were found in different areas of the truffle

orchard (i.e. areas 1, 2 or 3), they were separated. Similarly, we did not calculate the

isobarycentre with samples sharing the same genotype in different years.

All the maps were obtained with a dedicated python program developed for this study. The

mapping program can be downloaded using a Unix terminal and the following command: git

clone https://git.igzor.net/inra/iam_mapping.git

Acknowledgments

The French National Research Agency (ANR) as part of the "Investissements d'Avenir"

program (ANR-11-LABX-0002-01, Lab of Excellence ARBRE) and ANR SYSTERRA

SYSTRUF (ANR-09-STRA-10) financed this study. ALCINA also contributed by co-

financing the ClimaTruf project. We are grateful to Dr Francesco Paolocci and its team in

Perugia (Italy) for providing us the protocol of DNA extraction from mix of spores and for

the constructive discussion. Authors would also like to thank Prof Fabienne Malagnac for the

critical reading of the first version of the manuscript. We have no conflict of interest to

declare.

References

Armoza-Zvuloni, R., Kramarsky-Winter, E., Loya, Y., Schlesinger, A., and Rosenfeld, H.

(2014) Trioecy, a unique breeding strategy in the sea anemone *Aiptasia diaphana* and its

association with sex steroids. *Biol Reprod* **90**: 122.

Arnaud-Haond, S., and Belkhir, K. (2007) GENCLONE: a computer program to analyse

genotypic data, test for clonality and describe spatial clonal organization. *Mol Ecol Notes* **7**:

15-17.

Bailleul, D., Stoeckel, S., and Arnaud-Haond, S. (2016) RClone: a package to identify

MultiLocus Clonal Lineages and handle clonal data sets in R. *Methods Ecol Evol*

doi:10.1111/2041-210X.12550.

Belfiori, B., Riccioni, C., Paolocci, F., and Rubini, A. (2013) Mating Type Locus of Chinese

Black Truffles Reveals Heterothallism and the Presence of Cryptic Species within the *T.*

indicum Species Complex. *Plos One* **8:** e82353.

Belfiori, B., Riccioni, C., Paolocci, F., and Rubini, A. (2016) Characterization of the

reproductive mode and life cycle of the whitish truffle *T. borchii*. *Mycorrhiza* **26**: 515-527.

Benjamin, R.K. (1986) Laboulbeniales on semiaquatic Hemiptera. V. Triceromyces: with a

description of monoecious-dioecious dimorphism in the genus. *Aliso* **11:** 245-278.

Benjamin, R.K., and Shanor, L. (1950) Discovery of Dioecism in *Laboulbenia formicarum*.

Science **111**: 33-34.

Bonneau, L. (2016) A propos des apports de spores dans les brulés. L'historique et le vécu

"en temps reel" par l'expérimentateur. *Le Trufficulteur* **94**: 9-12.

Bradley, R. (1726) New Improvements of Planting and Gardening: Both Philosophical and

Practical: In three parts. London, UK: Printed for W. Mears.

Buffon, G., Sonnini, C.S., and Latreille, P. (1749) Histoire Naturelle Des Plantes. Tome

Troisième. Plantes Cryptogames, Des Champignons. In Histoire Naturelle Générale et

Particulière Avec La Description Du Cabinet Du Roi (1749 à 1789). Paris, France:

Imprimerie Royale.

Chaudhuri, J., Kache, V., and Pires-daSilva, A. (2011) Regulation of Sexual Plasticity in a

Nematode that Produces Males, Females, and Hermaphrodites. *Cur Biol* **21**: 1548-1551.

Chevalier, G., and Grente, J. (1979) Application pratique de la symbiose ectomycorhizienne:

production à grande echelle de plants mycorhizes par la truffe (*Tuber melanosporum* Vitt.). In

International Congress on the Science and Cultivation of Edible Fungi. France, pp. 483-505.

Chevalier, G., Gregori, G., Frochot, H., and Zambonelli, A. (2001) The cultivation of the

Burgundy truffle. In Proceedings of the Second International Conference on Edible

Mycorrhizal Mushrooms. Bencivenga, M., and Granetti, B. (eds). Spoleto, Italy: Comunità

Montana dei Monti Martani e del Serano.

Ciccarello, A. (1564) Opusculum De Tuberibus. Ludouici Bozetti (eds.). Pavia, Italy.

Esser, K., and Kuenen, R. (1967) Genetics of Fungi. Berlin-Heidelberg-New York.

Glass, N.L., and Kuldau, G.A. (1992) Mating Type and Vegetative Incompatibility in

Filamentous Ascomycetes. *Ann Rev Phytopathol* **30**: 201-224.

Gryndler, M., Černá, L., Bukovská, P., Hršelová, H., and Jansa, J. (2014) *Tuber aestivum*

association with non-host roots. *Mycorrhiza* **24**: 603-610.

Hansen H. N. & Snyder W. C. The Dual Phenomenon and Sex in *Hypomyces solani* F.

curcurbitae. *Am J Bot* **30**: 419-422.

Healy, R.A., Smith, M.E., Bonito, G.M., Pfister, D.H., Ge, Z.W., Guevara, G.G. et al. (2013)

High diversity and widespread occurrence of mitotic spore mats in ectomycorrhizal Pezizales.

Mol Ecol **22**: 1717-1732.

Joseph, K.S., and Murthy, H.N. (2014) Sexual system of *Garcinia indica* Choisy: geographic

variation in trioecy and sexual dimorphism in floral traits. *Plant Syst Evol* **301**: 1065-1071.

Kretzer, A.M., Dunham, S., Molina, R., and Spatafora, J.W. (2005) Patterns of vegetative

growth and gene flow in *Rhizopogon vinicolor* and *R. vesiculosus* (Boletales, Basidiomycota).

Mol Ecol **14**: 2259-2268.

Le Tacon, F., Rubini, A., Murat, C., Riccioni, C., Robin, C., Belfiori, B. et al. (2015)

Certainties and uncertainties about the life cycle of the Périgord black truffle (Tuber

melanosporum Vittad.). *Ann For Sci* **73**: 105-117.

Lehtonen, J., and Kokko, H. (2014) Sex. *Curr Biol* **24**: R305-R306.

Leslie, J.F. (1995) *Gibberella fujikuroi*: available populations and variable traits. *Can J Bot*

73: 282-291.

Leslie, J.F., and Raju, N.B. (1985) Recessive Mutations from Natural Populations of

Neurospora crassa That Are Expressed in the Sexual Diplophase. *Genetics* **111**: 759-777.

Leslie, J.F., and Klein, K.K. (1996) Female Fertility and Mating Type Effects on Effective

Population Size and Evolution in Filamentous Fungi. *Genetics* **144**: 557-567.

Maheshwari, R. (1999) Microconidia of *Neurospora crassa*. *Fungal Genet Biol* **26**: 1-18.

Martin, F., Kohler, A., Murat, C., Balestrini, R., Coutinho, P.M., Jaillon, O. et al. (2010)

Périgord black truffle genome uncovers evolutionary origins and mechanisms of symbiosis.

Nature **464**: 1033–1038.

Mirski, P., and Brzosko, E. (2015) Are hermaphrodites better adapted to the colonization

process in trioecious populations of *Salix myrsinifolia*? *Acta Soc Bot Polo* **84**: 167-175.

Molinier, V., Murat, C., Baltensweiler, A., Büntgen, U., Martin, F., Meier, B. et al. (2016)

Fine-scale genetic structure of natural *Tuber aestivum* sites in southern Germany. *Mycorrhiza*

doi:10.1007/s00572-016-0719-y.

Murat, C., Mello, A., Abbà, S., Vizzini, A., and Bonfante, P. (2008) Edible Mycorrhizal

Fungi: Identification, Life Cycle and Morphogenesis. In *Mycorrhiza*. Varma, P.D.A. (ed):

Springer edn. Berlin Heidelberg, Germany, pp. 707-732.

Murat, C., Riccioni, C., Belfiori, B., Cichocki, N., Labbé, J., Morin, E. et al. (2011)

Distribution and localization of microsatellites in the Perigord black truffle genome and

identification of new molecular markers. *Fungal Genet Biol* **48**: 592-601.

Murat, C., Rubini, A., Riccioni, C., De la Varga, H., Akroume, E., Belfiori, B. et al. (2013)

Fine-scale spatial genetic structure of the black truffle (*Tuber melanosporum*) investigated

with neutral microsatellites and functional mating type genes. *New Phytol* **199**: 176–187.

Nauta, M.J., and Hoekstra, R.F. (1992) Evolution of reproductive systems in filamentous

ascomycetes. II. Evolution of hermaphroditism and other reproductive strategies. *Heredity*

68: 537-546.

Nelson, M.A. (1996) Mating systems in ascomycetes: a romp in the sac. *Trends Genet* **12**: 69-

74.

Niculita-Hirzel, H., Labbé, J., Kohler, A., Le Tacon, F., Martin, F., Sanders, I.R., and Kües,

U. (2008) Gene organization of the mating type regions in the ectomycorrhizal fungus

Laccaria bicolor reveals distinct evolution between the two mating type loci. *New Phytol*

180: 329-342.

Nieuwenhuis, B.P.S., and Aanen, D.K. (2012) Sexual selection in fungi. *J Evol Biol* **25**: 2397-

2411.

Nieuwenhuis, B.P.S., Debets, A.J.M., and Aanen, D.K. (2011) Sexual selection in mushroom-

forming basidiomycetes. *Proc R Soc Lond B Biol Sci* **278**: 152-157.

Olivier, J., Savignac, J., and Sourzat, P. (2012) *Truffe et trufficulture*. Périgueux, France:

FANLAC Editions.

Paolocci, F., Rubini, A., Granetti, B., and Arcioni, S. (1999) Rapid molecular approach for a

reliable identification of *Tuber* spp. ectomycorrhizae. *FEMS Microbiol Ecol* **28**: 23-30.

Parladé, J., De la Varga, H., De Miguel, A., Sáez, R., and Pera, J. (2013) Quantification of

extraradical mycelium of *Tuber melanosporum* in soils from truffle orchards in northern

Spain. *Mycorrhiza* **23**: 99-103.

Rauscher, T., Agerer, R., and Chevalier, G. (1995) Ektomykorrhizen von *Tuber*

melanosporum, Tuber mesentericum und *Tuber rufum* (Tuberales) an *Corylus avellana*. *Nova*

Hedwigia **61**: 281-322.

Raymond, M., and Rousset, F. (1995) GENEPOP (Version 1.2): Population Genetics

Software for Exact Tests and Ecumenicism. *J Hered* **86**: 248-249.

Ropars, J., Dupont, J., Fontanillas, E., Rodríguez de la Vega, R.C., Malagnac, F., Coton, M. et

al. (2012) Sex in Cheese: Evidence for Sexuality in the Fungus *Penicillium roqueforti*. *PLoS*

ONE **7**: e49665.

Ropars, J., Lo, Y.-C., Dumas, E., Snirc, A., Begerow, D., Rollnik, T. et al. (2016) Fertility

depression among cheese-making Penicillium roqueforti strains suggests degeneration during

domestication. *Evolution* doi:10.1111/evo.13015.

Rousset, F. (2008) GENEPOP'007: a complete re-implementation of the genepop software

for Windows and Linux. *Mol Ecol Resour* **8**: 103-106.

Rubini, A., Belfiori, B., Riccioni, C., Arcioni, S., Martin, F., and Paolocci, F. (2011a) *Tuber*

melanosporum: mating type distribution in a natural plantation and dynamics of strains of

different mating types on the roots of nursery-inoculated host plants. *New Phytol* **189**: 723-

735.

Rubini, A., Belfiori, B., Riccioni, C., Tisserant, E., Arcioni, S., Martin, F., and Paolocci, F.

(2011b) Isolation and characterization of MAT genes in the symbiotic ascomycete *Tuber*

melanosporum. *New Phytol* **189**: 710-722.

Samils, N., Gioti, A., Karlsson, M., Sun, Y., Kasuga, T., Bastiaans, E. et al. (2013) Sex-linked

transcriptional divergence in the hermaphrodite fungus *Neurospora tetrasperma*. *P Roy Soc*

Lond B Biol Sci **280**: 20130862.

Smith, S.E., and Read, D.J. (2008) Mycorrhizal symbiosis. 3rd edn. San Diego, USA:

Elsevier Academic Press Inc

Stenberg, P., Lundmark, M., and Saura, A. (2003) MLGsim: a program for detecting clones

using a simulation approach. *Mol Ecol Notes* **3**: 329-331.

Urban, A., Neuner-Plattner, I., Krisai-Greilhuber, I., and Haselwandter, K. (2004) Molecular

studies on terricolous microfungi reveal novel anamorphs of two Tuber species. *Mycol Res*

108: 749-758.

Weeks, S.C., Sanderson, T.F., Reed, S.K., Zofkova, M., Knott, B., Balaraman, U. et al. (2006)

Ancient androdioecy in the freshwater crustacean Eulimnadia. *P Roy Soc Lond B Biol Sci*

273: 725-734.

Zambonelli, A., Salomoni, S., and Pisi, A. (1993) Caratterizzazione anatomo-morfologica

delle micorrize di *Tuber* spp. su *Quercus pubescens* Will. *Micol Ital* **3**: 73-90.

Tables and Figure legends

Table 1. Sampling details, genotypic and genetic diversity for female, male elements and ectomycorrhiza for five years in the Rollainville orchard.

Table 2. Quantity of soil mycelium of both mating types in mg of mycelium/g of soil. The position in the truffle orchard is indicated as well as the maternal mating type in the sampling square. For square without ascocarps the most frequent mating type of maternal tissue in the surrounding square is indicated (identified by an asterisk).

Fig. 1. Schematic representation of the sexual reproduction for the heterothallic ascomycete *T. melanosporum*. In order that sexual reproduction occurs female and male elements needs to mate. In nature, the female element is found in the host tree root system as ectomycorrhiza, although the male elements could have different origins: germinating ascospores, soil mycelium (i.e. free-living or associated to ectomycorrhiza for hermaphrodite strains) or conidia. Conidia have never been observed for *T. melanosporum,* but they have been described in other *Tuber* species. After the fertilization, a diploid transitory phase occurs (that cannot be detected in mature ascocarps), followed by a meiosis phase that will end in the formation of a mature ascocarp. The structures of ascocarps (peridium, gleba and ascospores) are represented by photographs.

Fig. 2. Map of distribution of the hermaphrodite genotypes R021, R060 and R068 in the Rollainville orchard. Representation of the distribution of the hermaphrodite genotypes along the different seasons and in the different areas of the orchard. a. Distribution of genotype R021. b. Distribution of genotype R060. c. Distribution of genotype R068. Dots represent female ascocarp genotypes, crosses represent ectomycorrhiza genotypes and triangles represent paternal genotypes. Each season is represented by a different colour.

Fig. 3. Isolation by distance analysis (IBD) representing genetic *vs* geographic distance for

female and male genotypes.

Isolation by distance analyses obtained with female (a and c) and male (b and d) genotypes. a

and b were obtained using all the samples while c and d were obtained using culled data to

avoid oversampling of the same genotype. The slope with 95 % interval confidence is

indicated for each graph.

Table 1. Sampling details, genotypic and genetic diversity for female, male elements and ectomycorrhiza for five years in the Rollainville

orchard.

	2010-2011	2011-2012	2012-2013	2013-2014	2014-2015	All seasons
Number of sampled ascocarps	17	42	30	101	51	238
Female elements						
Number of samples genotyped	17	42	30	98	51	238
Number of genotypes (no. of genotypes with Psex < 0.05)	7 (6)	21 (8)	23 (7)	31 (12)	13 (10)	73 (21)
Maximum number of ascocarps per genotype	4	6	3	18	20	40
Number of genotype found in a single ascocarp	2	11	15	12	3	42
Number of persistent genotype s (found in 2; 3; 4 or 5 seasons)						12 (6; 3; 2; 1)
Percent of ascocarps produced by persistent genotypes	94.12%	45.24%	26.67%	63.34%	88.24%	63.87%
Clonal Subrange (m)	3.74	3.23	4.06	20.46	22.36	22.71
Simpson diversity index (D)	0.88	0.94	0.98	0.92	0.81	0.99
Inter-individual diversity (1-Qinter = He)	0.52	0.84	0.91	0.90	0.77	0.88
Aggregation index (Ac) (p-value)	0.46 (0.002)	0.57 (0)	0.28 (0)	0.54 (0)	0.43 (0)	0.46 (0)
Male elements						
Number genotyped zygotes	14	35	28	78	51	206
Number of homozygous zygotes (% genotyped)	1 (7.14%)	1 (2.85%)	1 (3.57%)	2 (2.56%)	4 (7.84%)	9 (4.37%)
Zygotes Fis	0.02	0.18	0.28	0.18	0.15	0.21
Number genotypes (no. of genotypes with Psex < 0.05)	12 (1)	33 (4)	24 (3)	56 (11)	36 (10)	138 (15)
Max number of ascocarps per genotype	2	3	3	10	4	18
Number of male genotypes found in a single ascocarp	8	26	18	39	19	110
Number of persistent genotypes (found in 2; 3; 4 or 5 seasons)						16 (12; 1; 3; 0)

Ectomycorrhiza (ECM) samples						
Number of genotyped ECM	81				36	117
Number of genotypes (no. of genotypes with Psex < 0.05)	9(9)				23(14)	27 (18)
Clonal Subrange (m)	4.58				19.02	19.02
Simpson diversity index (D)	0.79				0.99	0.87
Aggregation index (Ac) - ECM (p-value)	1 (0)				0.22 (0)	0.85 (0)
Number of ECM genotype occurring as female/male/both strains						16/3/3
Aggregation index (Ac) - ECM and female (p-value)	0.96 (0)				0.40 (0)	0.77 (0)
Number of hermaphrodite* genotypes (no. genotypes with Psex < 0.05)	1 (1)	2 (2)	2 (1)	6 (2)	4 (2)	8 (3)

610
611 *Hermaphrodites are genotypes found as female and male elements*

612   Table 2. Quantity of soil mycelium of both mating types in mg of mycelium/g of soil.

613   The position in the truffle orchard is indicated as well as the maternal mating type in the

614   sampling square. For square without ascocarps the most frequent mating type of

615   maternal tissue in the surrounding square is indicated (identified by an asterisk).

Area	Square	Maternal MAT	*MAT 1-1*	*MAT 1-2*
1	A4	2	0.428	16.37
1	A8	2*	0.402	21.58
1	C3	2*	0.444	18.22
1	C6	2	0.408	23.03
1	E4	1	0.753	40.49
1	E8	1*	0	0.021
1	G3	1	6.14	0
1	G6	1*	7.4	0.04
2	B4	2	0.535	25.39
2	D2	2	0	0
2	D6	2*	0.11	5.6
2	F4	2*	0.022	1.13
3	A2	2	0.171	9.51
3	B7	2*	0.179	9.009
3	C4	2	0.567	31.56
3	D5/6	2	0.112	4.32
3	D9	1	3.55	0.04
3	E2	2	0.464	38.58
3	F7	1	1.87	0
3	G4	2*	0.224	10.79

617
618

619   **Fig. 1**

Female elements   **Male elements**

Ascogones issued from ectomycorrhizas   Soil mycelium   Germinating   Antherozoïds issued from   Conidies *
formed by female fertile genotypes and   (n)   ascospores   ectomycorrhizas formed by   (n)
hermaphrodite genotypes   (n)   hermaphrodite genotypes
(n)   (n)

or/and   or/and   or/and   **?**

Fertilization

Transitory ascogenous tissue (2n)

Meiosis

Mature ascomata

Peridium,
female origin (n)

White sterile veins,   Ascospores (n) in dark
female origin (n)   fertile veins

Gleba,
female and male
origin**
Dark fertile veins,
female (n) and
male** (n) origin

* Never described for *T. melanosporum*
** After ascospore formation male genotypes are not detected
anymore unless breaking ascospores cell walls.

620
621

Fig. 2

625 **Fig. 3**

626
627