1

Evapotranspiration is resilient in the face of land cover and climate change in a

humid temperate catchment

S. K. Hamilton[a,b,c*], M. Z. Hussain[a,b], C. Lowrie[b,d], B. Basso[a,e], and G. P. Robertson[a,b,f]

[a] W.K. Kellogg Biological Station, Michigan State University, Hickory Corners, MI
49060

[b] Great Lakes Bioenergy Research Center, Michigan State University, East Lansing, MI
48824

[c] Department of Integrative Biology, Michigan State University, East Lansing, MI 48824

[d] Department of Mathematics, Michigan State University, East Lansing, MI 48824

[e] Department of Earth and Environmental Sciences, Michigan State University, East
Lansing, MI 48824

[f] Department of Plant, Soil, and Microbial Sciences, Michigan State University, East
Lansing, MI 48824

* corresponding author: hamilton@msu.edu

Tel: 1-269-671-2231; Fax: 1-269-671-2104

Abstract

In temperate humid catchments, evapotranspiration returns more than half of the annual

precipitation to the atmosphere, thereby determining the balance available to recharge

groundwaters and support stream flow and lake levels. Changes in evapotranspiration

rates and therefore catchment hydrology could be driven by changes in land use or

climate. Here we examine the catchment water balance over the past 50 y for a catchment

in southwest Michigan covered by cropland, grassland, forest, and wetlands. Over the

study period about 27% of the catchment has been abandoned from row-crop agriculture

to perennial vegetation and about 20% of the catchment has reverted to deciduous forest,

and the climate has warmed by 1.14°C. Despite these changes in land use, precipitation

and stream discharge, and by inference catchment-scale evapotranspiration, have been

stable over the study period. The remarkably stable rates of evapotranspirative water loss

from the catchment across a period of significant land cover change suggest that rainfed

annual crops and perennial vegetation do not differ greatly in evapotranspiration rates,

and this is supported by measurements of evapotranspiration from various vegetation

types based on soil water monitoring in the same catchment. Compensating changes in

the other meteorological drivers of evaporative water demand besides air temperature—

wind speed, atmospheric humidity, and net radiation—are also possible, but cannot be

evaluated due to insufficient local data across the 50-y period. Regardless of the

explanation, this study shows that the water balance of this landscape has been resilient in

the face of both land cover and climate change over the past 50 y.

Keywords: evapotranspiration, evaporation, crops, forest, crops, land use, climate

change

1. Introduction

In temperate humid catchments, evapotranspiration (ET) returns more than half of the annual precipitation to the atmosphere (Hanson, 1991; Williams *et al.,* 2012; Zhang *et al.,* 2016), mainly during the growing season by plant transpiration. The balance between precipitation and ET recharges groundwaters and supports stream flow and lake levels. Paired catchment studies often have shown that changes in the nature of the vegetation cover, especially deforestation or afforestation, alter ecosystem ET rates and thereby change stream flows (Bosch and Hewlett, 1982; Hornbeck *et al.,* 1993; Zhang *et al.,* 2001; Price, 2011; Brown *et al.,* 2005 and 2013). However, these studies are often conducted in small experimental catchments and generally compare stream water yields between two kinds of perennial vegetation (woody and herbaceous).

There have been fewer catchment-scale comparisons of water yield from annual vegetation such as maize (*Zea mays*, known as corn in the U.S.) and soybean (*Glycine max*) vs. perennial vegetation such as forest or grasslands (Price, 2011), yet land cover change from perennial vegetation to cropland and vice versa has occurred throughout the world as a result of agricultural expansion and contraction. In eastern North America, the original forests and grasslands were largely converted to agricultural lands by European settlers, but since the mid 1900s a substantial fraction of the converted land has reverted back to successional fields and forests as the more marginal agricultural lands were abandoned due to low profitability, poor suitability to mechanized cultivation, and concerns about soil erosion and degradation (Houghton and Hackler 2000; Ramankutty *et al.,* 2010).

Land cover in agricultural regions is expected to continue to change in the future.

As grain crops have become more profitable over the past decade due to global demand

for food and US policies that support ethanol production from maize, more land in

grasslands (including CRP land) is being converted to grow maize and soybean (Lark *et*

al., 2015). Meanwhile, successional ecosystems are becoming mature forests in many

locations (Pugh, 2015). Climate change and invasive plant species will increasingly drive

changes in the nature and phenology of vegetation communities (Simberloff, 2000;

Parmesan and Hanley 2015). Further changes to the nature of vegetation in agricultural

landscapes may occur if cellulosic biofuel crops are increasingly grown in the future

(Gelfand *et al.,* 2013).

Recently we reported ET measurements in candidate cellulosic cropping systems

at a location in southwest Michigan, USA using two distinct approaches: 1) by

monitoring soil water content with time domain reflectometry in annual crops (maize) as

well as perennial grasslands and hybrid poplar stands (Hamilton *et al.,* 2015); and 2) by

monitoring energy and water vapor fluxes using eddy covariance in maize, switchgrass,

and prairie at a nearby site (Abraha *et al.,* 2015). Results suggest strikingly similar

growing-season ET among these diverse plant systems, raising the question of whether

land cover changes would significantly affect ET in the Midwest U.S., as suggested in

some modeling studies (e.g., Le *et al.,* 2011; VanLoocke *et al.,* 2012; Zhuang *et al.,*

2013).

The objective of this study is to examine trends in ET over 50 y in a particularly

well-characterized, temperate humid catchment that has experienced significant land

cover change, but without the complications of urbanization, dams, and stormwater

management changes that are typical of larger catchments. We infer ET from the balance

between precipitation and discharge, and the results are compared with our independent

measurements of ET made on annual and perennial vegetation in the same catchment.

2. Methods

2.1 Study site

Augusta Creek is a 3[rd]-order stream in southwest Michigan (Kalamazoo and Barry

counties) that drains a predominantly rural landscape (95 km^2) composed of a mosaic of

forest, fallow fields, annual crops, wetlands, lakes, light residential development, and golf

courses (Figure 1). There are no impervious surfaces or storm drainage systems that drain

into the stream above the discharge measurement point, and urban land use covers just

2.4% of the catchment (land cover proportions over the 50-y period are presented later).

The stream is groundwater-fed, gaining water along most of its length. Its tributaries

emanate from wetlands or small lakes, and prairie fen wetlands line much of the stream

channels.

The stream runs through deep glacial deposits that lie well above the bedrock. The

most common soils in upland areas are well-drained Typic Hapludalfs developed on

postglacial terrain (Thoen, 1990), and there is little to no overland flow from upland areas

to the stream due to the high permeability of these coarse-textured soils (Rheaume,

1990). Irrigation of crops was rare in the area until very recently; some expansion has

taken place since 2005, supplied by groundwater wells.

Augusta Creek is in the vicinity of the W.K. Kellogg Biological Station (KBS),

where we conduct agricultural experiments under the aegis of the Great Lakes Bioenergy

Research Center (GLBRC) and KBS Long Term Ecological Research site

(www.lter.kbs.msu.edu; 42.3956° N, 85.3749° W and 288 m asl). Mean annual air

temperature is 10.1 °C and annual precipitation is 1005 mm, 511 mm of which falls as

rain during the May-Sep growing season (1981–2010) (NCDC 2013). In this region ET is

normally water–limited during the warmer part of the year (i.e., during at least part of the

growing season), and energy–limited during the cooler months (McVicar *et al.*, 2012a).

2.2 Land cover changes

Land cover for 1960 was estimated from georectified and mosaicked aerial photographs

in a geographic information system (ArcGIS). The catchment boundaries above the

discharge measurement point (US Geological Survey; Hydrologic Unit Codes

04050003040060 plus 04050003040070) and wetlands and lakes (National Wetlands

Inventory: http://www.fws.gov/wetlands/) were overlain on the aerial photo mosaic and

land cover was examined in the upland portions of the catchment. Based on the National

Wetlands Inventory, wetlands and lakes contiguous with the stream system amount to

15.9 km^2, or 16.6 % of the catchment above the discharge measurement point. Isolated

wetlands and small lakes also occur throughout the upland catchment, covering 5.2% of

its area. Wetland areas were assumed to be constant over the study period; there has been

no wetland drainage or creation in the catchment since 1960, and within the area mapped,

wetland boundaries generally include intermittently wet soils with high water tables as

well as areas with surface water.

Land cover for 2014 was estimated from the Cropland Data Layer

(http://nassgeodata.gmu.edu/CropScape/). For this purpose, we combined all field crops

(primarily maize, soybean and small grains) into the annual crop category, and all forests

(deciduous and coniferous) into the forest category. Conifers are not native to the upland

landscape here, but have been planted throughout the catchment; their total area as of

2014 amounts to ~3% of the total forest area and 1% of the upland catchment. The

grassland and pasture category includes hay as well as fallow fields (no native grassland

remains). The width of rural roads was exaggerated 3–5 fold in the Cropland Data Layer,

presumably due to automated classification of mixed pixels, so vegetated edges of

roadways were manually reclassified as grasslands. Land cover for an intermediate date

(1978), based on aerial photo interpretation, was available from the Michigan Resource

Inventory System (MIRIS) (http://www.ciesin.org/IC/mdnr/mrip.html); this data set was

comparable for forest but combines annual crops with some kinds of pasture, and was

therefore not compared for those categories.

2.3 Discharge and climate records

The US Geological Survey has monitored the discharge of Augusta Creek below the

lowermost tributary inflow since 1964 (station 04105700). The long-term mean discharge

at this point, which drains 95.3 km^2, is 1.28 m^3 s^{-1}. Daily discharge measurements for

October 1964 through September 2014 were partitioned into baseflow and stormflow

using the Web-based Hydrograph Analysis Tool (WHAT) described by Lim *et al.*,

(2005). Mean annual baseflow and stormflow discharges were calculated on a standard

U.S. water-year basis beginning on 1 October of each year, representing the transition

between warm and cool seasons, and water years are labeled by the starting year (i.e.,

water year 1964 is 1 October 1964–30 September 1965).

Climate data were drawn from several sources and compiled on a water-year

basis. Precipitation observations are from at least three stations (except 1992 which has

two) distributed across the catchment from north to south (Figure 1). Air temperature,

saturated vapor pressure, and drought index data were obtained from the Midwest

Regional Climate Center (http://mrcc.isws.illinois.edu/).

2.4 Estimation of evapotranspiration from water balances

Evapotranspiration has often been estimated from catchment water balances (e.g., Zhang

et al., 2016). For Augusta Creek, the water balance for the upland portion of the

catchment was determined as the difference between annual totals of precipitation falling

on the uplands (i.e., the catchment excluding wetlands and lakes contiguous with the

stream channels) and the annual stream baseflow discharge. Isolated lakes and wetlands

were included in the upland catchment area. The difference between precipitation inputs

on the uplands and stream baseflow outputs is therefore considered to represent the ET of

the upland catchment.

This approach to ET estimation assumes that stormflow represents direct capture

of precipitation from the wetlands and lakes contiguous with the stream system, whereas

baseflow represents infiltration and percolation of precipitation falling on the upland

catchment. The validity of this assumption is supported by the water balance calculations

(see Results below) as well as the high permeability of the soils in the uplands (Rheaume,

1990). Other assumptions that are reasonable in this case include no inter-basin transfers

of water, which is true in this catchment, and no significant trend in water storage in the

aquifer or surface water bodies over the study period. Although there are no continuous

water table measurements spanning this study period, water levels of local lakes that are

connected to the groundwater have shown no unidirectional trend since the late 1960s

(see Figure S1 for an example of water level data for a lake in the Augusta Creek

catchment). Additional evidence for no interannual trend in groundwater levels is

provided by a compilation of static water level measurements that are made when

residential water supply wells are constructed, which shows no trend over the study

period (Figure S2).

2.5 Evapotranspiration estimation from soil water content measurements

Since 2009, soil water profiles throughout the root zone and below were monitored

hourly using permanently installed, horizontally inserted time domain reflectometry

(TDR) probes at depths of 20, 35, 50, 65, 90 and 125 cm as well as a vertically inserted

probe at 0–10 cm depth. Our methods for estimating ET from soil water profiles are

described by Hamilton *et al.* (2015), who presented data on six biofuel cropping systems

harvested each fall. The TDR measurements provide an estimate of ET when daily

drawdowns in soil water can be measured and the soil water content is below its drained

upper limit, which is typical of most of the growing season. The sum of the daily

drawdowns in soil water content over the entire profile (0-150 cm) across the growing

season provides an estimate of ET; on days when new infiltration of rain water prevented

a measurable soil water drawdown, we estimated ET using a crop growth model (Basso

and Ritchie, 2012).

Here we present the mean ET rates for three of those systems that resemble

vegetation found on the broader landscape: 1) continuous no-till maize; 2) a restored

native prairie planted with 18 species of forbs and grasses; and 3) a hybrid poplar

plantation (*Populus nigra* × *P. maximowiczii 'NM6'*). In addition, we present comparable

water use measurements for three other systems in the same vicinity: 1) a fallow field

abandoned from row-crop agriculture in 2008 and harvested each fall; 2) a mature

deciduous forest (>50 y old) dominated by sugar maple (*Acer saccharum*), red oak

(*Quercus rubra*) and hickory (*Carya* spp.) trees; and 3) an early successional forest (ca.

25 years old) dominated by shrubs including autumn olive (*Elaeagnus umbellata*) and

honeysuckle (*Lonicera* sp.*) as well as a few medium-sized sugar maple and black cherry

(*Prunus serotina*) trees.

3. Results

3.1 Land use and climate changes

Maize has been the dominant agricultural crop over the 50-y study period with the

balance of harvested crops shifting increasingly to soybean since the 1970s, as in the

greater Midwest US region (Gage *et al.,* 2015). Data on Kalamazoo County from the

annual Census of Agriculture (U.S. Department of Agriculture:

http://www.agcensus.usda.gov/) indicate that in 1964 maize accounted for 69% of

harvested cropland, soybean for 5.7%, and the balance was mostly oats with some barley

and wheat. By 1987 maize was 58% and soybean 28% of harvested cropland, and by

2007 these two crops accounted for 64% and 32% of the harvested cropland.

Land cover in the upland catchment changed significantly between 1960 and 2014

(Figure 2). The proportion of the upland catchment in annual crops decreased from 57 to

30%, while forest increased from 15 to 35%. The proportion of grassland remained

similar, although only 20% of the 1960 upland grassland was still grassland in 2014;

most of the 1960 grassland became forest (43%) or cropland (22%), while some newly

abandoned cropland became grassland. The 1978 MIRIS land cover data (not shown; see

Methods) indicate that 94% of the forest present in 2014 existed by 1978, so most

reforestation began between 1960–78. Urban and residential development represents a

small fraction of the catchment (<2.4%), not including golf courses created during the

study period that covered 4.5% of the upland catchment by 2014 (the golf courses

occasionally irrigate during dry summers but are not significant water users at the

catchment scale). Similar changes in land cover occurred in adjacent catchments.

Annual precipitation for the Augusta Creek catchment over the 50 y averaged 948

$\pm$ 118 mm y^{-1} with no linear temporal trend (p = 0.93) (Figure 3a). No linear trend exists

in mean annual values for either the Palmer Drought Severity Index or the Palmer

Hydrological Drought Index (p = 0.34 and 0.67, respectively; Figure S3).

One or more of the four meteorological variables that control atmospheric

evaporative demand—wind speed, atmospheric humidity, net radiation, and air

temperature—could have changed over the 50 y, as global- and continental-scale analyses

have indicated significant changes in these variables in recent decades (Wild, 2009;

Willett *et al.,* 2008; McVicar *et al.,* 2012b). The effects of changes in these variables on

atmospheric evaporative demand could be to enhance or counteract each other, and the

resultant effect on ET is particularly important where evaporation is limited by energy

rather than water (e.g., decreasing wind speeds tend to counteract the effect of increasing

temperatures: McVicar *et al.,* 2012a, b). The region has experienced a 1.14°C increase in

mean annual air temperature (50-y mean = 8.95°C) which in turn equates to a 0.90

millibar (mb) increase in saturated vapor pressure (50-y mean = 13.5 mb) over the 50-y

period (Figure 4). Consistent data across the study period for wind speed, atmospheric

humidity, and net radiation are not available for this locale.

3.2 Catchment hydrology

Stream discharge partitioned into stormflow and baseflow shows how

groundwater dominates the total flow of Augusta Creek; baseflow averaged 78% of the

total discharge (Figure 3b). There is no linear trend in total ($p = 0.14$), stormflow ($p =$

0.91), or baseflow ($p = 0.83$) discharge over the 50 y. In this catchment, stormflow likely

reflects mainly precipitation falling on lakes and wetlands that are contiguous with the

stream channels because upland soils are highly permeable and there are few impervious

surfaces and little overland runoff from uplands to the streams. This is supported by the

comparison of annual stormflow volumes to the annual precipitation falling on

contiguous lakes and wetlands: on average, stormflow amounts to 57% (range, 44-73%)

of the precipitation with no linear trend over the 50 y ($p = 0.09$, data not shown). The

balance, which equates to a mean of 408 mm y^{-1}, could largely be explained by

evapotranspirative losses from the lakes and wetlands. If stormflow originating as

overland flow from the uplands were important, the total stormflow volume would

exceed the precipitation on lakes and wetlands.

Our annual water balances for Augusta Creek resemble earlier estimates

calculated by Rheaume (1990) over three representative years (1971, 1977 and 1985),

which indicated that 62, 65 and 59%, respectively, of the annual precipitation was

returned to the atmosphere as ET, mainly during the growing season (May–Sep),

although those estimates included ET from contiguous lakes and wetlands as well as

uplands. That study also employed hydrograph separation to estimate that about 75% of

the annual stream flow in those years was supported by groundwater discharge; our

estimate of mean baseflow contribution over the 50-y period is 78%.

Our estimate of ET, based on the difference between precipitation on the upland

catchment and baseflow discharge out of the catchment, averaged 563 ± 103 mm y^{-1}

(Figure 3c), with no linear trend (p = 0.98). Expressed as a percentage of annual

precipitation, ET averaged $59 \pm 6\%$ (Figure 3d), also with no trend over the 50 y (p =

0.88). Therefore these data show that ET from upland areas of the Augusta Creek

catchment has remained remarkably stable over the past 50 y in spite of large changes in

land cover towards less area in annual crops and more in deciduous forest.

3.3 ET rates from representative vegetation types

We estimated ET in annual crops and perennial vegetation over the 2009–2014

period from high-resolution changes in soil water profiles (Figure 5). Except for 2012,

which was a drought year, mean growing season ET rates were 495 ± 48 mm y^{-1} for

maize, 524 ± 79 for grasslands (fallow and prairie), and 532 ± 47 for woody vegetation

(deciduous forest, shrubland, and poplar). These rates are statistically indistinguishable

among vegetation types (p>0.05), further supporting the hypothesis that ET rates are

similar among annual crops, perennial grasslands, and forests in the Augusta Creek

catchment. These ET observations span years of varying warmth (Figure 4) but show no

relationship with mean growing-season temperature.

While soil water-based ET rates, excluding the 2012 drought year, are lower than

the catchment-based ET rates of 600 ± 59 mm y^{-1} in those years (2009, 2010, 2011, 2013,

and 2014 in Figure 3c), the soil water-based ET estimates reflect only the growing

seasons. Year-round eddy covariance measurements of water fluxes in maize and

grasslands at KBS indicate that about 30% of ET occurs outside the May-Sep growing

season (Abraha *et al.,* 2015). Adding 30% to the soil water-based ET rates brings rates

for maize, grasslands, and woody vegetation to 643, 681, and 692 mm y^{-1}, respectively,

all higher but within 15% of the catchment-based ET measurements over those years.

4. Discussion

4.1 Possible explanations for the stability of ET

There are several possible explanations for the long-term stability of catchment

ET that we believe are unlikely. One is that there may not have been sufficient time for

hydrologic responses to be detected. While the mean transit time for groundwater

movement in this kind of catchment is likely greater than a decade (e.g., Saad, 2008),

groundwater discharge rates from an unconfined and connected aquifer system would

respond to changing recharge at far faster time scales (McDonnell and Beven, 2014).

Succession from grassland to forest can be protracted, but the MIRIS forest cover data

indicate that most of the reforestation occurred in the first 14 years of the study period

(i.e., 1964–78). Many long-term paired catchment studies have shown that water yield

after regrowth of harvested forest tends to approach a stable rate within about 10–25

years (Hornbeck *et al.,* 1993; Brown *et al.,* 2013).

Another possibility is that the degree of land cover change over the study period

(27% of the upland catchment abandoned from annual crops and 20% of it becoming

reforested; Figure 2) may not be sufficiently large to signal a change in water yield, even

if annual crops and perennial vegetation had large differences in ET rates. Again, this is

unlikely because long-term paired catchment studies have shown significant change with

as little as 20% of the catchment either deforested or afforested (Brown *et al.,* 2005).

Also possible is that there are offsetting effects exerted by different land covers in

the vicinity (Albertson *et al.,* 2001; van Dijk *et al.*, 2012), but this does not seem likely

because adjacent catchments have similar mosaics of land cover, and the entire region has

experienced similar changes in vegetation over this time period. Compensating land use

changes that result in no net change in ET are also a possibility, such as the changes in

crops grown as noted above. However the ET rate of oats that were commonly grown in

the 1960s and 1970s is unlikely to differ much from the maize that replaced them (Allen

et al., 1998).

Over the past 50 y the mean annual air temperature has increased by about 1.14

°C (Figure 4), and the frost-free season has become longer by about 9 days (Kunkle,

2015). Evapotranspiration could increase with warming if available water were not

limiting, other meteorological changes did not offset the temperature effect (McVicar *et*

al., 2012b), and the vegetation could remain active over the longer growing season.

However, during the growing season when most (~70%) of the ET occurs, available soil

water typically becomes limiting to ET (Hamilton *et al.,* 2015). Also, most annual crops

and many grasses would senesce before the end of the potential growing season because

their development is regulated by degree-days (Parmesan and Hanley, 2015).

Extrapolation of observations from small catchments that are entirely covered by

one kind of vegetation to complex mixtures of vegetation may not be as straightforward

as it would seem. Models of ET and discharge from catchments with mixed land covers

has often proven challenging to validate, and a variety of possible reasons have been

considered (van Dijk *et al.,* 2012). Methodological issues identified by those authors

include uncertainties in land cover, precipitation and discharge data; in the case of the

Augusta Creek catchment, however, the precipitation and land cover data are likely to be

quite accurate. It is also possible that other catchment climate characteristics that we have

not considered are more influential to ET than land cover (e.g., Wilcox and Huang,

2010). Physical explanations noted by van Dijk *et al.,* (2012) for poor model performance

include recirculation of intercepted rainfall, which tends to be more important in forests,

and lateral water redistribution between vegetation types; identifying the potential

importance of these physical explanations in the Augusta Creek catchment is beyond the

scope of this study.

We cannot rule out the possibility that changes in the meteorological drivers of

atmospheric water demand (i.e., temperature, net radiation, wind speed, and atmospheric

humidity; McVicar *et al.*, 2012a, b) could have offset the effects of land cover changes on

ET. The steadily increasing partial pressure of atmospheric carbon dioxide could also

have reduced plant transpiration rates, although its effect on ET is most pronounced in

warm, highly water-limited (i.e., arid and subarid) regions (Donohue *et al.*, 2013; Yang *et*

al., 2016; Trancoso *et al.*, 2017). In any case the offset of land cover effects on ET by

these atmospheric changes would be a regional phenomenon contributing to the resilience

of catchment ET and discharge.

4.2 Conclusion

Evapotranspirative water loss in the upland portion of the Augusta Creek

catchment has been remarkably resilient across a 50-y period of decreasing cropland,

increasing perennial vegetation cover, and warming temperatures, leaving a relatively

consistent proportion of precipitation for groundwater recharge and streamflow. Our ET

estimates based on catchment water balances compare well with direct measurements in

the same catchment since 2009 based on soil water monitoring by time-domain

reflectometry for grasslands, annual crops, and perennial bioenergy crops and forest.

These observations suggest that water use by rainfed annual crops and perennial

vegetation is similar in this setting, and that in humid catchments with soil permeability

little affected by land cover, catchment water balances are not likely to be very sensitive

to near-term future changes in land cover and climate as long as the land is vegetated, and

crops are not irrigated. One such land cover change could be an increase in the cultivation

of perennial herbaceous crops for biofuel production, which, based on our findings, does

not seem likely to alter catchment water balances in this kind of setting.

Figures:

Figure 1. Location of Augusta Creek in Michigan (inset) and catchment boundaries

(shaded). Precipitation measurement sites are shown by triangles. The catchment

boundaries can be viewed in Google Earth: (catchment boundary kml file here).

Figure 2. Land cover change in the Augusta Creek catchment. Estimates are based on

analysis of aerial photos (1960) or satellite imagery (2014 Cropland Data Layer:

http://nassgeodata.gmu.edu/CropScape/).

Figure 3. Precipitation, stream discharge, and evapotranspiration (ET). Panels show

annual (Oct-Sep) values of (a) precipitation measured at 3-6 stations (mean = blue

line); (b) stream discharge partitioned into baseflow and stormflow; (c)

evapotranspiration (ET) estimated as the difference between precipitation and

baseflow discharge; and (d) ET as a percentage of annual precipitation. Horizontal

lines show the means.

Figure 4. Air temperature for the Augusta Creek catchment, derived from the Midwest

Regional Climate Center database (http://mrcc.isws.illinois.edu/). The positive

change is significant ($P = 0.005$) and amounts to 1.14°C over the 50 y.

Figure 5. Rainfall (blue) and evapotranspiration over the growing season (2009-14) from

annual maize and herbaceous and perennial vegetation, estimated from continuous

observations of plant water uptake in soil profiles.

Figure 1. Location of Augusta Creek in Michigan (inset) and watershed boundaries (shaded). Precipitation measurement sites are shown by triangles.

Figure 2. Land cover change in the Augusta Creek watershed. Estimates are based on analysis of aerial photos (1960) or satellite imagery (2014 Cropland Data Layer: http://nassgeodata.gmu.edu/CropScape/).

Figure 3. Precipitation, stream discharge, and evapotranspiration (ET). Panels show annual (Oct-Sep) values of (a) precipitation measured at 3-6 stations (mean = blue line); (b) stream discharge partitioned into baseflow and stormflow; (c) evapotranspiration (ET) estimated as the difference between precipitation and baseflow discharge; and (d) ET as a percentage of annual precipitation.

Figure 4. Air temperature (a) and saturated vapor pressure (b) for the Augusta Creek watershed, derived from the Midwest Regional Climate Center database (http://mrcc.isws.illinois.edu/). The positive changes are significant (P = 0.005 and 0.003 for temperature and vapor pressure, respectively) and amount to 1.14°C and 0.90 millibars over the 50 years.

Figure 5. Rainfall (blue) and evapotranspiration over the growing season (2009-14) from annual maize and herbaceous and perennial vegetation, estimated from continuous observations of plant water uptake in soil profiles.

Supplementary figures:

Supplementary Figure S1. Water levels in Fair Lake, one of two lakes forming the

headwaters of the Augusta Creek system. Note that these data extend back to well

before the start of the study period on 1 Oct 1964, and encompass a series of

drought years in the early 1960s. Since 1967 there has not been a unidirectional

trend across years that would suggest large changes in groundwater or surface

water storage. No other local lakes, whether draining to streams or isolated, are

known to have had their water levels change unidirectionally over the study

period. Data are from http://lter.kbs.msu.edu/datatables/381.

Supplementary Figure S2. Static water levels measured upon installation of residential

water supply wells in the vicinity of the Augusta Creek catchment. Data compiled

from public records by Shu-Guang Li of Michigan State University.

Supplementary Figure S3. The Palmer Drought Severity Index (a) and the Palmer

Hydrological Drought Index (b) for the region encompassing the Augusta Creek

watershed, derived from the Midwest Regional Climate Center database

(http://mrcc.isws.illinois.edu/). There is no significant linear trend in either index

($p = 0.34$ and 067, respectively).

Evapotranspiration is resilient in the face of land cover and climate change in a

humid temperate catchment

Supplementary Information

Supplementary Figure S1. Water levels in Fair Lake, one of two lakes forming the
headwaters of the Augusta Creek system. Note that these data extend back to well before
the start of the study period on 1 Oct 1964, and encompass a series of drought years in
the early 1960s. Since 1967 there has not been a unidirectional trend across years that
would suggest large changes in groundwater or surface water storage. No other local
lakes, whether draining to streams or isolated, are known to have changed
unidirectionally over the study period. Data are available at
http://lter.kbs.msu.edu/datatables/381.

Supplementary Figure S2. Static water levels measured upon installation of residential water supply wells in the vicinity of the Augusta Creek watershed. Data compiled from public records by Shu-Guang Li of Michigan State University.

Supplementary Figure S3. The Palmer Drought Severity Index (a) and the Palmer Hydrological Drought Index (b) for the region encompassing the Augusta Creek watershed, derived from the Midwest Regional Climate Center database (http://mrcc.isws.illinois.edu/). There is no significant linear trend in either index (p = 0.34 and 067, respectively).

References

Abraha, M., Chen, J., Chu, H., Zenone, T., John, R., Su, Y.-J., Hamilton, S.K., and

Robertson, G.P. (2015). Evapotranspiration of annual and perennial biofuel crops

in a variable climate. *Global Change Biology Bioenergy*,

doi:10.1111/gcbb.12239.

Albertson, J.D., Katul, G.G., Wiberg, P. (2001). Relative importance of local and

regional controls on coupled water, carbon, and energy fluxes. *Advances in Water*

Resources, 24, 1101-1118.

Allen, R.G., Pereira, L.S., Raes, D., Smith, M. (1998). Crop Evapotranspiration—

Guidelines for Computing Crop Water Requirements (FAO Irrigation and

Drainage Paper 56, Food and Agricultural Organization, Rome).

Basso, B., Ritchie, J.T. (2012). Assessing the impact of management strategies on water

use efficiency using soil-plant-atmosphere models. *Vadose Zone Journal*, 11(3),

doi: 10.2136/vzj2011.0173.

Bosch, J.M., Hewlett, J.D. (1982). A review of catchment experiments to determine the

effect of vegetation changes on water yield and evapotranspiration. *Journal of*

Hydrology, 55 (1/4), 3–23.

Brown, A.E., Zhang, L., McMahon, T.A., Westem, A.W., Vertessy, R.A. (2005). A

review of paired catchment studies for determining changes in water yield

resulting from alterations in vegetation. *Journal of Hydrology*, 310, 28-61.

Brown, A.E., Westem, A.W., McMahon, T.A., Zhang, L. (2013). Impact of forest cover

changes on annual streamflow and flow duration curves. *Journal of Hydrology*,

483, 39-50.

Donohue, R.J., Roderick, M.L., McVicar, T.R., Farquhar, G.D. (2013). Impact of CO_2

fertilization on maximum foliage cover across the globe's warm, arid

environments, *Geophysical Research Letters,* 40, 3031–3035, doi:

10.1002/grl.50563.

Gage, S.H., Doll, J.E., Safir, G.R. (2015). A crop stress index to predict climatic effects

on row-crop agriculture in the U.S. North Central Region. Pages 77-103 in

Hamilton S K, Doll J E, Robertson G P, editors. The Ecology of Agricultural

Landscapes: Long-Term Research on the Path to Sustainability. Oxford

University Press, New York, New York, USA.

Gelfand, I., Sahajpal, R., Zhang, X., Izaurralde, R.C., Gross, K.L., and Robertson, G.P.

(2013). Sustainable bioenergy production from marginal lands in the US Midwest

Nature, 493, 514–517.

Hamilton, S.K., Hussain, M.Z., Bhardwaj, A.K., Basso, B., Robertson, G.P. (2015).

Comparative water use by maize, perennial crops, restored prairie, and poplar

trees in the US Midwest. *Environmental Research Letters*, 10 065015, doi:

10.1088/1748-9326/10/6/064015.

Hanson, R.L. (1991). Evapotranspiration and Droughts, in Paulson, R.W., Chase, E.B.,

Roberts, R.S., Moody, D.W., Compilers, National Water Summary 1988-89--

Hydrologic Events and Floods and Droughts: U.S. Geological Survey Water-

Supply Paper 2375, 99-104.

Hornbeck, J.W., Adams, M.B., Corbett, E.S., Verry, E.S., Lynch, J.A. (1993). Long-term

impacts of forest treatments on water yield: a summary for northeastern USA

Journal of Hydrology, 150 (2/4), 323–344.

Houghton, R.A., Hackler, J.L. (2000). Changes in terrestrial carbon storage in the United

States. I: The roles of agriculture and forestry. *Global Ecology & Biogeography*,

9, 125-144.

Kunkle, K. (2015). Change in Length of Growing Season by State, 1895-2014. Data from

http://www3.epa.gov/climatechange/science/indicators/health-society/growing-

season.html; downloaded 11 March 2016.

Lark, T.J., Salmon, J.M., Gibbs, H.K. (2015). Cropland expansion outpaces agricultural

and biofuel policies in the United States. *Environmental Research Letters*, 10

044003 http://dx.doi.org/10.1088/1748-9326/10/4/044003.

Le, P.V.V., Kumar, P., Drewry, D.T. (2011). Implications for the hydrologic cycle under

climate change due to the expansion of bioenergy crops in the Midwestern United

States. *Proceedings of the National Academy of Sciences of the USA*, 108, 15085–

15090.

Lim, K.J., Engel, B.A., Tang, Z., Choi, J., Kim, K., Muthukrishnan, S., Tripathy, D.

(2005). Web GIS-based Hydrograph Analysis Tool, WHAT. *Journal of the

American Water Resources Association*, 41, 1407-1416.

McDonnell, J.J., Beven, K. (2014). Debates—The future of hydrological sciences: A

(common) path forward? A call to action aimed at understanding velocities,

celerities and residence time distributions of the headwater hydrograph. *Water

Resources Research*, 50, doi:10.1002/2013WR015141.

McVicar, T.R., Roderick, M.L., Donohue, R.J. and Van Niel, T.G. (2012a) Less bluster

ahead? Ecohydrological implications of global trends of terrestrial near-surface

wind speeds. *Ecohydrology*, 5, 381–388. doi: 10.1002/eco.1298

McVicar, T.R., Roderick, M.L., Donohue, R.J., Li, L.T., Van Niel, T.G., Thomas, A.,

Grieser, J., Jhajharia, D., Himri, Y., Mahowald, N.M., Mescherskaya, A.V.,

Kruger, A.C., Rehman, S., and Dinpashoh, Y. (2012b) Global review and

synthesis of trends in observed terrestrial near-surface wind speeds: Implications

for evaporation. *Journal of Hydrology*, 416-417, 182-205.

doi:10.1016/j.jhydrol.2011.10.024

NCDC (National Climate Data Center). (2013). Summary of Monthly Normal 1981–

2010. Gull Lake Biology Station, MI, USA. <http://www.ncdc.noaa.gov/cdo-

web/search> Accessed January 17, 2013.

Parmesan, C., Hanley, M.E. (2015). Plants and climate change: complexities and

surprises *Annals of Botany*, 116, 849–864.

Price, K. (2011). Effects of watershed topography, soils, land use, and climate on

baseflow hydrology in humid regions: A review. *Progress in Physical

Geography*, 35, 465-492.

Pugh, S.A. (2015). Forests of Michigan, 2014. Resource Update FS-35. Newtown

Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research

Station. 4 p.

Ramankutty, N., Heller, E., Rhemtulla, J. (2010). Prevailing myths about agricultural

abandonment and forest regrowth in the United States. *Annals of the Association

of American Geographers*, 100, 502-512.

Rheaume, S.J. (1990). Geohydrology and water quality of Kalamazoo County, Michigan,

1986-88. U.S. Geological Survey Water-Resources Investigations Report 90-

4028, 102 p.

Saad, D.A. (2008). Agriculture-related trends in groundwater quality of the glacial

deposits aquifer, central Wisconsin. *Journal of Environmental Quality*, 37, S240-

S248.

Simberloff, D. (2000). Global climate change and introduced species in United States

forests. *Science of the Total Environment*, 262, 253-262.

Thoen, G. (1990). Soil Survey of Barry County, Michigan (USDA Soil Conservation

Service and Michigan Agricultural Experiment Station, East Lansing, Michigan,

Washington, DC), 286 p.

Trancoso, R., Larsen, J.R., McVicar, T.R., Phinn, S.R., McAlpine, C.A. (2017). CO_2-

vegetation feedbacks and other climate changes implicated in reducing base flow.

Geophysical Research Letters, 44, 2310–2318, doi:10.1002/2017GL072759

Van Dijk, A. I. J. M., Peña-Arancibia, J.L., Bruijnzeel, L. A. (Sampurno). (2012). Land

cover and water yield: inference problems when comparing catchments with

mixed land cover. *Hydrology and Earth System Sciences*, 16, 3461–3473, doi:

10.5194/hess-16-3461-2012

VanLoocke, A., Twine, T., Zeri, M., and Bernacchi, C.J. (2012). A regional comparison

of water use efficiency for miscanthus, switchgrass and maize. *Agricultural and*

Forest Meteorology, 164, 82–95.

Wilcox, B.P., Huang, Y. (2010). Woody plant encroachment paradox: Rivers rebound as

degraded grasslands convert to woodlands, *Geophysical Research Letters*, 37,

L07402, doi: 10.1029/2009GL041929

Wild, M. (2009). Global dimming and brightening: a review. *Journal of Geophysical*

Research-Atmospheres, 114: D00D16, doi: 10.1029/2008JD011470.

Willett, K.M., Jones, P.D., Gillett, N.P., Thorne, P.W. (2008). Recent Changes in surface

humidity: development of the HadCRUH dataset. *Journal of Climate*, 21, 5364-

5383.

Williams, C.A., Reichstein, M., Buchmann, N., Baldocchi, D., Beer, C., Schwalm, C.,

Wohlfahrt, G., Hasler, N., Bernhofer, C., Foken, T., Papale, D., Schymanski S.,

Schaefer, K. (2012). Climate and vegetation controls on the surface water

balance: Synthesis of evapotranspiration measured across a global network of

flux towers *Water Resources Research*, 48, W06523, doi:

10.1029/2011WR011586.

Yang, Y., Donohue, R.J., McVicar, T.R., Roderick, M.L., Beck, H.E. (2016). Long-term

CO_2 fertilization increases vegetation productivity and has little effect on

hydrological partitioning in tropical rainforests, *Journal of Geophysical Research-*

-Biogeosciences, 121, 2125–2140, doi:10.1002/2016JG003475

Zhang, L., Dawes, W.R., Walker, G.R. (2001). Response of mean annual

evapotranspiration to vegetation changes at catchment scale. *Water Resources*

Research, 37, 701–708.

Zhang, Y., Peña-Arancibia, J.L., McVicar, T.R., Chiew, F.H.S., Vaze, J., Liu, C., Lu, X.,

Zheng, H., Wang, Y., Liu, Y.Y., Miralles, D.G., Pan, M. **(**2016). Multi-decadal

trends in global terrestrial evapotranspiration and its components. *Scientific*

Reports, 6, 19124, doi:10.1038/srep19124

Zhuang, Q., Qin, Z., Chen, M. (2013). Biofuel, land and water: switchgrass or

Miscanthus? *Environmental Research Letters*, 8 015020, doi:10.1088/1748-

9326/8/1/015020.

Acknowledgments

We thank A.K. Bhardwaj, S. Bohm, K. Kahmark, and S.-G. Li for instrumentation and

data assistance, local citizens W. Shafer, T. Smith, and W. Knollenberg for precipitation

data supplemental to that from our research and National Weather Service stations, and

the numerous people at Michigan State University and the U.S. Geological Survey who

helped maintain the precipitation and stream discharge records since 1964. T. McVicar,

J.J. McDonnell and T. Dunne read earlier versions and provided helpful advice on data

interpretation. Financial support for this work was provided by the U.S. Department of

Energy through the Great Lakes Bioenergy Research Center (DOE BER Office of

Science DE-FC02-07ER64494 and DOE OBP Office of Energy Efficiency and

Renewable Energy DE-AC05-76RL01830), the U.S. National Science Foundation

(LTER program, DEB 1027253), and the Michigan Agricultural Experiment Station.