

# Comparison of African, Asian, and American Zika Viruses in Swiss Webster mice: Virulence, neutralizing antibodies, and serotypes

Charles B. Stauff<sup>a,b</sup>, Oleksandr Gorbatshevych<sup>a</sup>, Jeronimo Cello<sup>a</sup>, Eckard Wimmer<sup>a,b</sup>, and Bruce Futcher<sup>a</sup>

<sup>a</sup>Dept. of Molecular Genetics and Microbiology, Stony Brook University, 11794-5222; <sup>b</sup>Codagenix Inc., Stony Brook, NY 11794

This manuscript was compiled on September 16, 2016

The sequence of Zika virus has evolved as it has spread out of Africa and into the Americas. It is unclear whether American strains of the virus define a new serotype. Here, we have tested the virulence and immunogenicity of three wild-type ZIKV strains in neonatal Swiss Webster mice. We found that all three ZIKV strains (African MR766, 1947; Asian FSS13025, 2010; and American, PRVABC59, 2015) are capable of killing neonatal mice after intracranial injection. Intraperitoneal injection with these viruses did not kill, but produced neutralizing antibodies as measured by a PRNT<sub>50</sub> assay. Sera from mice infected with each virus were tested for neutralizing activity against the infecting virus and also the other two viruses by a PRNT<sub>50</sub> assay. In general, the antibodies induced by each virus were good at neutralizing that virus (the homologous virus), but somewhat poorer at neutralizing the other two viruses (heterologous viruses). Antibodies induced by the African strain MR766 were about 4-fold worse at neutralizing the American strain PRVABC59 than the homologous strain, while antibodies induced by the American strain were about 10-fold worse at neutralizing the African strain than the homologous strain. Because the antibodies are cross-neutralizing at some level, the viruses do not form separate serotypes. Nevertheless, these results raise concern that the immunity conferred by the African virus may protect only relatively poorly against the new American strains. This has implications for the possible spread of the American ZIKV strains to Africa and Southeast Asia, and also for the development of vaccines.

## Introduction

Zika virus has migrated out of Africa where it was discovered in 1947[1] into Southeast Asia[2--4], into Micronesia [5, 6] and French Polynesia [7, 8] and most recently into South and Central America [9--13]. Since 2015, mosquito-borne Zika virus transmission has been reported in 64 countries and territories, with 50 countries and territories reporting their first ever Zika virus outbreaks (WHO). Alarmingly, Zika virus infection causes birth defects such as microcephaly in infants [14--16] and Guillain-Barre syndrome in adults [17--19]. Zika virus has also reached the United States, with 4,729 locally acquired cases in US Territories and 1,657 travel-associated cases reported within the United States itself[20]. Zika virus infection within the United States and its territories are already associated with 22 cases of Guillain-Barre syndrome. 885 infections are of pregnant women [21] whose children are now at risk for developing birth defects.

Zika virus was first discovered in Uganda in 1947. Subsequence serological studies suggested that, at least in some regions of Africa, a very large proportion of the adult population is seropositive for Zika virus, and so is presumably immune to new infections. However, as Zika virus has spread across the globe, its primary genome sequence has changed. Because of the significant amino acid changes from the ancestral MR766 to the new American

strains, a question arises as to whether the American strains of virus form a novel serotype. That is, will individuals with immunity against the ancestral African virus also have cross-immunity against the new American strains? If not, there is a concern that the novel strains may cause a wave of infection across Africa and Southeast Asia, notwithstanding the fact that older forms of Zika virus are already endemic there. Furthermore, there is an issue of whether a vaccine for Zika virus should be based on the MR766 amino acid sequence.

Zika virus, like dengue virus, infects primates [1, 22]. However, neonatal wild-type mice are also sensitive to Zika [23] and dengue [24] viruses. Previously, we have used a proxy challenge model using immune competent mice to gauge the effectiveness of dengue virus vaccine candidates to protect neonatal mice against lethal challenge [25]. We feel that this model will be equally applicable to research with ZIKV, a closely related flavivirus. Here, we infect neonatal mice with Zika. After intracranial injection, these infections are lethal, while after intraperitoneal injection, infections induce neutralizing antibodies. We compare the African, Asian, and American viruses for virulence in this model, and ask whether the antibodies induced by one strain differ in potency against the other strains.

## Materials/Methods

**Cells and Viruses.** Vero cells were grown in Dulbecco's Modified Eagle's Medium (DMEM) supplemented with 10% bovine calf serum (BCS; Gemclone) and Penicillin/Streptomycin (CellGro).

ZIKV strain MR766-SM150 (LC002520) and FSS13025 (JN860855) were acquired from Dr. Robert Tesh (UTMB, Galveston TX). ZIKV strain PRVABC59 (KU501215) was provided by BEI Resources (NIAID, NIH: Zika Virus, PRV-ABC59, NR-50240). MR766 [1] was isolated in 1947 from a primate in Uganda, FSS13025 [25] was isolated in Cambodia from a human patient in 2010, and PRVABC59 [26, 27] is a recent isolate (2015) from a human patient in Puerto Rico. Before infection in animals, stocks of each virus were generated by infecting confluent Vero cells at an MOI of 0.1 and harvesting cell culture supernatant at 3 days post infection.

**Mouse Infection.** To prepare immune serum, African (MR766 P3) or Asian (FSS13025) or American (PRVABC59) Zika viruses were used to infect 4 week old Swiss Webster (Charles River) mice at a dose of  $1 \times 10^6$  PFU delivered intraperitoneally in a volume of 100  $\mu$ L DMEM. Six mice were infected with each virus, but one of the mice infected with FSS13025 failed to develop any antibody titer against any virus (perhaps due to an injection failure), and this serum was not included in analysis. At 28 days' post-infection, mice were euthanized and terminally bled through cardiac puncture. Whole blood was allowed to clot for 30 minutes at room temperature, centrifuged at 10,000 rpm for 15 minutes, serum collected and frozen at  $-80^{\circ}\text{C}$ .


To determine  $\text{LD}_{50}$  in newborn Swiss Webster mice, litters ( $n=7-12$ ) were intracranially injected with 20  $\mu$ L of ZIKV MR766, FSS13025, or PRVABC59 diluted in Dulbecco's Modified Eagle's Medium (DMEM). Mice were checked daily for morbidity and mortality for 21 days post infection. All animal experiments were conducted under approval by the Institutional Animal Care and Use Committee at Stony Brook University.

**PRNT<sub>50</sub> Assay.** Standard working solutions of MR766 (Monkey / Uganda / 1947), FSS13025 (Human / Cambodia / 2010), and PRVABC59 (Human / Puerto Rico / 2015) were prepared at a concentration of 500 PFU/mL. Mouse serum was heat treated for 30 minutes at  $56^{\circ}\text{C}$  and then mixed with 100 PFU of each virus separately with 4-fold serum dilutions of 1:10, 1:40, 1:160, and 1:640 diluted in Vero infection medium (5% BCS DMEM). Each serum/virus

mixture was incubated at 37°C for 2 hours and then added to confluent Vero cell monolayers in a 12-well format with a volume of 250  $\mu$ L. Plates were rocked at room temperature for 30 minutes, then 250  $\mu$ L of additional Vero infection medium was added and the plates incubated at 37°C, 5% CO<sub>2</sub> for 1 hour. Medium was then aspirated from each plate, the wells were washed once with sterile DPBS, and 1 mL of 0.6% Tragacanth Gum, 5% BCS DMEM was added to each well. The plates were incubated at 37°C, 5% CO<sub>2</sub> for 4 days before being stained with crystal violet and read. The PRNT<sub>50</sub> titer was determined by taking the reciprocal of the highest dilution that reduced the plaque count by 50% or more. PRNT<sub>50</sub> values were log<sub>2</sub> transformed and compared using a paired t-test for MR766 and PRVABC59 infected mice, for which the virus-to-virus differences in PRNT<sub>50</sub> values were distributed normally. There was not a normal distribution for the mice infected with FSS13025, so these results were not analyzed with the paired t-test.

## Results and Discussion

Alignment of the sequence from the African strain MR766 (Uganda, 1947) with the American strain PRVABC59 (Puerto Rico, 2015) shows about 10% nucleotide change. A pairwise alignment of the polyprotein amino acid sequences for MR766 and either PRVABC59, or FSS13025 reveal a similar mutation profile where both the American and Asian strains contain about 125 amino acid differences from the African strain polyprotein (Fig. 1 B). Interestingly, both PRVABC59 and FSS13025 contain a four amino acid insertion in the envelope E protein that is responsible for host cell receptor binding (Fig. 1 B in red). FSS13025 is closely related to PRVABC59, containing only one amino-acid substitution in the E protein. Swiss Webster neonatal mice are highly susceptible to DENV[28] and ZIKV[23] when delivered intracranially. In preparation for future vaccine studies, we established the virulence of wild-type Zika viruses using this model. Individual litters were infected and mice were weighed and checked daily for signs of infection. The most commonly observed sign of infection was piloerection and the Swiss Webster neonates were susceptible to infection by all three strains of ZIKV (Fig. 2). Our strain of MR766 was passaged 150 times in suckling Swiss Webster mice [23] and was very lethal in this model with an LD<sub>50</sub> of 1 PFU, possibly because of this passaging. FSS13025 and PRVABC59 were also lethal in newborn mice, with LD<sub>50</sub> values of 250 and 2.7, respectively (Table 1) without recorded passaging in newborn mice. Thus, PRVABC59 is almost as virulent in mice as MR766, despite the fact that the latter has been mouse-adapted by passaging, and the former has not. Mean time to death (MTD) was also calculated for the highest dose administered in each group (Table 1). Similar in pattern to the LD<sub>50</sub> results, MR766 infected mice succumbed more quickly to infection (MTD=6.91 days) compared to FSS13025 (MTD= 13.0 days) or PRVABC59 (MTD= 11.4 days).




**Figure 1.** Mutation profiles of FSS13025 and PRVABC59 when aligned to MR766. Single amino acid differences are denoted by black and red bars relative to the polyprotein sequence of MR766 (B). The four red bars highlight a four amino acid insertion in FSS13025 and PRVABC59. Mutations map to the polyprotein products outlined in (A).

Six mice were infected intraperitoneally with the African and American viruses, and five mice with the Asian virus, and immune sera were harvested. All sera were tested by PRNT<sub>50</sub> assay for neutralizing antibody titers against all three viruses (Fig. 3). Titers from mice injected with the African strain

were 4-fold higher against the homologous African strain than against the heterologous Asian or American strains (Fig. 3A, Tables S1-S3). These differences were statistically significant ( $p = 0.042$  for each comparison) by a paired t-test. Conversely, titers from mice injected with the American strain were 10-fold higher against the homologous American strain than against the heterologous African strain (Fig. 3C, Tables S1-S3), and about 1.6-fold higher than against the Asian strain (Fig. 3C, Tables S1-S3). The 10-fold difference between the American and African strain titers was highly significant ( $p = 0.014$ ) by a paired t-test. Results with mice injected with the Asian strain were intermediate, but appeared qualitatively more similar to results with the American strain than the African strain (Fig. 3B). Because titer differences with the Asian virus were not normally distributed, significance could not be evaluated by the paired t-test.

These results show that there is significant strain specificity in the neutralizing antibodies induced by African and American strains: Antibodies raised against the African strain are more potent (~4-fold) against the African than the Asian or American strains, while antibodies raised against the American strain are more potent (~10-fold) against the American strain than the African strain (Tables S1-S3). This is consistent with the sequence divergence between these strains. The fact that these strains can be distinguished serologically needs to be taken into account in vaccine design: a vaccine against the American strains is more likely to be effective if it is based on an American strain than if it is based on an African strain.




**Figure 2.** Neonatal Swiss Webster mice were infected with MR766 (A), FSS13025 (B) or PRVABC59 (C) injected intracranially.

Our serological results so far are based on the in vitro PRNT<sub>50</sub> assay. An obvious next step would be to look at in vivo protection. Neonatal mice that have acquired antibodies induced by one virus (either by direct injection of serum, or maternal antibodies)<sup>24</sup> could be challenged with the homologous and heterologous viruses.

A recent paper correlated neutralizing antibodies<sup>[27]</sup> against Zika virus (specifically anti E glycoprotein antibodies) with protection against challenge. Using tests for neutralizing antibodies in infected mice is therefore essential to testing vaccine candidates. Larocca et al<sup>[27]</sup> noted that a microneutralization titer (essentially a PRNT<sub>100</sub> value) of 10 was sufficient to protect mice against challenge. The limit of detection (blue line) for the PRNT<sub>50</sub> used in our study was a 10-fold dilution in serum. It may be that the PRNT<sub>50</sub> titers we observed, while low, are still sufficient to protect mice against heterologous challenge with other strains of ZIKV.

Two recent papers have reached conclusions somewhat different from ours. First, Aliota et al. <sup>[29]</sup> infected rhesus macaques with the African Zika strain MR766, and challenged these macaques 70 days later with a French Polynesian strain. The macaques were fully protected against the second infection, showing no viremia. Aliota et al. concluded that "Immunogen selection is unlikely to adversely affect the breadth of vaccine protection" and "ZIKV strain selection is unlikely to compromise vaccine effectiveness". However, protection after 70 days (while certainly a good thing!) is not a quantitative measure of vaccine effectiveness. At 70 days after the primary infection, antibody titers are likely very high, and likely



**Figure 3.** 4-week old, Swiss Webster mice were infected with  $1 \times 10^6$  PFU of either the MR766, FSS13025, or PRVABC59 strains of ZIKV. Serum was collected from each mouse at 28 days post infection and tested by PRNT<sub>50</sub> against homologous and heterologous strains of ZIKV. In each panel (A, B, or C), results from serum from an individual mouse are marked with a different symbol; i.e., an open circle with no fill represents a single mouse in that panel. Yellow shading in symbols indicates the homologous virus.

to afford protection even if those antibodies are not perfectly matched against the new antigens. We found a 4 to 10-fold strain-dependent difference in the neutralizing activity of our sera, which does not immediately conflict with Aliota et al as these titers may still be protective in vivo[27]. Regardless of the difference we feel that for an optimal vaccine, immunogen selection is likely important.

**Table 1.** Neonatal, 1-2 day old, Swiss Webster mice were infected intracranially with serial dilutions of MR766, FSS13025, or PRVABC59 and mortality assessed using Kaplan-Meier curves. LD<sub>50</sub> and mean time to death (MTD) values were calculated for the highest dose used in the infections.

Virus	LD <sub>50</sub>	MTD
MR766 <sup>a</sup>	1	6.91
FSS13025 <sup>b</sup>	250	13
PRVABC59 <sup>c</sup>	2.7	11.4

<sup>a</sup>  $1 \times 10^3$  PFU dose

<sup>b</sup>  $1 \times 10^3$  PFU dose

<sup>c</sup>  $5 \times 10^3$  PFU dose

Second, Dowd et al.[30] did mouse experiments broadly similar to ours, with several differences in technical detail, but unlike us did not find statistically-significant differences in cross neutralization. Possibly relevant technical differences are that we compared a recent American virus, PRVABC59, to the African virus MR766, while Dowd et al. compared an Asian virus from 2013 (H/FP/2013). In addition, we infected juvenile wild-type mice to obtain sera, while Dowd et al. infected mutant *irf3/irf3*<sup>-/-</sup> mice. However, it is not clear that either of these technical differences should matter. We saw moderate differences in PRNT<sub>50</sub> (i.e., neutralizing antibody titer) between viruses of four-fold (when MR766 was the infecting virus) to ten-fold (when PRVABC59 was the infecting virus). We judged these differences to be statistically significant using a paired t-test, which is an appropriate and relatively powerful test under these circumstances. In contrast, Dowd et al. saw essentially no differences when MR766 was the infecting virus, and differences of perhaps two-fold when the Asian virus H/FP/2013 was the infecting virus (Fig. 4A of Dowd et al.), and they judged these differences to be not statistically

significant using ANOVA. We believe ANOVA is less powerful than a paired t-test under these circumstances, particularly since the number of mice used both by us and by Dowd et al. was rather small. However regardless of the statistical test, the fact remains that the fold-differences in neutralizing titres were at best small (~2-fold) for Dowd et al., but moderate (~4-fold to ~10-fold) for us, and the reasons for these differences are unknown.

In summary, our results suggest that the new American strains of Zika may be relatively virulent, at least in juvenile mice. The American strain PRVABC59 is almost as virulent in mice as MR766, despite the fact that MR766 has been repeatedly passaged in mice, while PRVABC59 has not. Furthermore, we detect moderate serological differences

between the ancestral African virus and the modern American Zika virus, with a four to ten-fold difference in neutralization titer when comparing homologous to heterologous viruses. On the one hand, a four to ten-fold difference is relatively modest. On the other hand, given that immunity can wane with time, and that individual responses vary, even a four-fold difference could be functionally important for a vaccine. In our studies, the number of mice used was small, and the serological results are based solely on the in vitro PRNT<sub>50</sub> assay, so of course further and larger studies of various types are required. Nevertheless, since vaccines against Zika virus are already being used in human volunteers (NCT02840487, NCT02887482, NCT02809443), it is important to consider that for maximum effectiveness against the American virus, vaccines should be based on an immunogen from an American virus.

**Acknowledgments.** This study was supported by a grant from the NIH (NIH 1R01AI110792 – 01A1) and also funded by the Center for Biotechnology, a New York State Center for Advanced Technology; New York State Department of Economic Development Contract #C150127; and corporate support.

**Competing Interests.** CS is an employee of, and EW is a founder of Codagenix, Inc. which has potential commercial interests in anti-Zika vaccines. EW, JC and BF have interests in patents held by the Research Foundation of Stony Brook University which may be applicable to the manufacture of anti-Zika vaccines.

## Bibliography

1. Dick GWA, Kitchen SF, Haddow AJ (1952) Zika Virus (I). Isolations and serological specificity. *Transactions of The Royal Society of Tropical Medicine and Hygiene* 46(5):509–520.
2. Buathong R et al. (2015) Detection of Zika Virus Infection in Thailand, 2012–2014. *The American Journal of Tropical Medicine and Hygiene* 93(2):380–383.
3. Haddow AD et al. (2012) Genetic Characterization of Zika Virus Strains: Geographic Expansion of the Asian Lineage. *PLoS Neglected Tropical Diseases* 6(2):e1477.
4. Olson JG, Ksiazek TG, Suhandiman n, Triwibowo n (1981) Zika virus, a cause of fever in Central Java, Indonesia. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 75(3):389–393.
5. Duffy MR et al. (2009) Zika Virus Outbreak on Yap Island, Federated States of Micronesia. *New England Journal of Medicine* 360(24):2536–2543.
6. Lanciotti RS et al. (2008) Genetic and Serologic Properties of Zika Virus Associated with an Epidemic, Yap State, Micronesia, 2007. *Emerging Infectious Diseases* 14(8):1232–1239.
7. Baronti C et al. (2014) Complete Coding Sequence of Zika Virus from a French Polynesia Outbreak in 2013. *Genome Announcements* 2(3):e00500–14–e00500–14.
8. Cao-Lormeau VM (2014) Zika Virus, French Polynesia, South Pacific, 2013. *Emerging Infectious Diseases* 20(11):1960–1960.
9. Campos GS, Bandeira AC, Sardi SI (2015) Zika Virus Outbreak, Bahia, Brazil. *Emerging Infectious Diseases* 21(10):1885–1886.
10. Petersen E et al. (2016) Unexpected and Rapid Spread of Zika Virus in The Americas - Implications for Public Health Preparedness for Mass Gatherings at the 2016 Brazil Olympic Games. *International Journal of Infectious Diseases*.
11. Zanluca C et al. (2015) First report of autochthonous transmission of Zika virus in Brazil. *Memórias do Instituto Oswaldo Cruz* 110(4):569–572.
12. Guerbois M et al. (2016) Outbreak of Zika virus infection, Chiapas State, Mexico, 2015, and first confirmed transmission by *Aedes aegypti* mosquitoes in the Americas. *Journal of Infectious Diseases* p. jiw302.
13. Roa M (2016) Zika virus outbreak: reproductive health and rights in Latin America. *The Lancet* 387(10021):843.
14. Brasil P et al. (2016) Zika Virus Infection in Pregnant Women in Rio de Janeiro — Preliminary Report. *New England Journal of Medicine*.
15. Cauchemez S et al. (2016) Association between Zika virus and microcephaly in French Polynesia, 2013–15: a retrospective study. *The Lancet* 387(10033):2125–2132.

16. Mlakar J et al. (2016) Zika virus associated with microcephaly. *New England Journal of Medicine*.
17. Broutet N et al. (2016) Zika Virus as a Cause of Neurologic Disorders. *New England Journal of Medicine*.
18. Carteaux G et al. (2016) Zika Virus Associated with Meningoencephalitis. *New England Journal of Medicine*.
19. Oehler E et al. (2014) Zika virus infection complicated by Guillain-Barre syndrome—case report, French Polynesia, December 2013. *Euro Surveill* 19(9):20720.
20. (2014) *Zika Virus*. <http://www.cdc.gov/zika/geo/pregwomen-uscases.html>.
21. (2014) *Zika Virus*. <http://www.cdc.gov/zika/geo/united-states.html>.
22. Abbink P et al. (2016) Protective efficacy of multiple vaccine platforms against Zika virus challenge in rhesus monkeys. *Science* p. aah6157.
23. Dick GWA (1952) Zika virus (II). Pathogenicity and physical properties. *Transactions of The Royal Society of Tropical Medicine and Hygiene* 46(5):521–534.
24. Shen SH et al. (2015) Large-scale recoding of an arbovirus genome to rebalance its insect versus mammalian preference. *Proceedings of the National Academy of Sciences of the United States of America* 112(15):4749–4754.
25. Shan C et al. (2016) An Infectious cDNA Clone of Zika Virus to Study Viral Virulence, Mosquito Transmission, and Antiviral Inhibitors. *Cell Host & Microbe* 19(6):891–900.
26. Faria NR et al. (2016) Zika virus in the Americas: Early epidemiological and genetic findings. *Science* 352(6283):345–349.
27. Larocca RA et al. (2016) Vaccine protection against Zika virus from Brazil. *Nature*.
28. Chen W, Kawano H, Men R, Clark D, Lai CJ (1995) Construction of intertypic chimeric dengue viruses exhibiting type 3 antigenicity and neurovirulence for mice. *Journal of Virology* 69(8):5186–5190.
29. Aliota MT et al. (2016) Heterologous protection against Asian Zika virus challenge in rhesus macaques. *bioRxiv* p. 059592.
30. Dowd KA et al. (2016) Broadly Neutralizing Activity of Zika Virus-Immune Sera Identifies a Single Viral Serotype. *Cell Reports* 16(6):1485–1491.