Quorum sensing integrates environmental cues, cell density and cell

history to control bacterial competence

Stefany Moreno-Gámez[1,2], Robin A. Sorg[1], Arnau Domenech[1,4], Morten Kjos[1,3],

Franz J. Weissing[2], G. Sander van Doorn[2,5] and Jan-Willem Veening[1,4,5]

1. Molecular Genetics Group, Groningen Biomolecular Sciences and Biotechnology

Institute, Centre for Synthetic Biology, University of Groningen, Nijenborgh 7, 9747

AG Groningen, the Netherlands

2. Groningen Institute for Evolutionary Life Sciences, University of Groningen,

Groningen, P.O. Box 11103, 9700 CC, The Netherlands

3. Present address: Department of Chemistry, Biotechnology and Food Science,

Norwegian University of Life Sciences, N-1432 Ås, Norway

4. Present address: Department of Fundamental Microbiology, Faculty of Biology and

Medicine, University of Lausanne, Biophore Building, CH-1015 Lausanne,

Switzerland

5. Correspondence to G. Sander van Doorn: g.s.van.doorn@rug.nl, tel: +31 (0)50 363

8097 and Jan-Willem Veening: Jan-Willem.Veening@unil.ch, tel: +41 (0)21 6925625,

Twitter: @JWVeening

Keywords: Quorum sensing, *Streptococcus pneumoniae*, antibiotics, competence,

transformation, bacterial stress response, mathematical modeling

Abstract

Streptococcus pneumoniae becomes competent for genetic transformation when exposed to an autoinducer peptide named CSP. This peptide was originally described as a quorum-sensing (QS) signal, enabling individual cells to regulate competence in response to population density. However, recent studies suggest that CSP may instead serve as a probe for sensing environmental cues, such as antibiotic stress or environmental diffusion. Here, we show that competence induction can be simultaneously influenced by cell density, external pH, antibiotic-induced stress and cell history. Our experimental data is explained by a mathematical model where the environment and cell history modify how cells produce or sense CSP. Taken together, model and experiments indicate that autoinducer concentration can function as an indicator of cell density across environmental conditions, while also incorporating information on environmental factors or cell history, allowing cells to integrate cues such as antibiotic stress into their QS response. This unifying perspective may apply to other debated QS systems.

Introduction

Bacteria release small diffusible molecules in the extracellular medium known as autoinducers. These molecules induce the expression of particular functions including biofilm formation, luminescence and genetic competence as well as their own production [1,2]. The most prevalent functional interpretation of the production and response to autoinducers is known as quorum sensing (QS). According to this view, the concentration of autoinducer molecules is a proxy for cell density, allowing bacteria to regulate the expression of those phenotypes that are only beneficial when

expressed by many cells [1,2]. However, it is likely that the concentration of autoinducer

molecules does not only reflect cell density, but that it is also affected by

environmental factors, such as the diffusivity of the medium. In fact, alternative

hypotheses state that bacteria release autoinducers to sense these environmental factors

rather than to monitor cell density. The most well-known hypothesis proposed by

Redfield is that the function of autoinducers is diffusion sensing, allowing cells to

avoid the secretion of costly molecules under conditions where they would quickly

diffuse away [3]. Other potential roles suggested for autoinducer production are sensing

local cell density together with diffusion [4], the positioning of other cells during biofilm

formation [5] and temporal variations in pH [6].

We study pneumococcal competence, a system classically used as an example of

QS. However, whether competence is actually controlled by QS has been recently

debated. Competence is a transient physiological state that is developed by

Streptococcus pneumoniae, as well as other bacteria. Upon entry into competence,

pneumococci upregulate the expression of genes required for uptake of exogenous

DNA as well as bacteriocins and various genes involved in stress response [7]. In *S.*

pneumoniae, competence is regulated by an autoinducer molecule known as the

competence-stimulating peptide (CSP) in a two-component regulatory system formed

by the histidine kinase ComD and the response regulator ComE [8,9] (Figure 1). Despite

the detailed understanding of the regulatory network of competence induction, little is

known about why competence is controlled by an autoinducer peptide like CSP. CSP

has been classically thought to be a QS signal[10], whose function could be to monitor

the density of potential DNA donors[14]. However, competence can be induced in

response to environmental factors like pH, oxygen, phosphate and antibiotic stress [11–

14]. Based on this evidence and the finding that competence initiates at the same time in

pneumococcal cultures inoculated at different initial densities, it was suggested that

CSP acts as a timing device that allows cells to mount a timed response to

environmental stress independently of cell density [15,16]. Since then, this hypothesis has

established in the field as an alternative to the QS view of competence [13,16–18].

Recently, Prudhomme et al. renamed the timing device mechanism as a growth-time

dependent mechanism and proposed that a subpopulation of competent cells that

originates stochastically spreads the competent state to the rest of the population by

cell-cell contact [19]. Another alternative to QS is that pneumococcal competence is an

instance of diffusion sensing. This was suggested by Yang et al. based on the

observation that the quorum for competence induction is not fixed but decreases with

more restrictive diffusion [20].

Here, we study the regulation of pneumococcal competence by cell density and two

environmental factors, antibiotic stress and pH. Using batch-culture experiments,

single cell analyses and mathematical modeling we show that these factors

simultaneously regulate competence development because they all affect the CSP

concentration: cell density sets the amount of cells producing CSP, whereas the

environment and cell history modify the rate at which individual cells produce or sense

CSP. Since there is density regulation and we show that CSP is exported

extracellularly, we advocate to keep using the term 'quorum sensing' in the context of

pneumococcal competence but with a broader meaning to acknowledge that in addition

to cell density, multiple factors are integrated into this QS response.

Results

A mathematical model of pneumococcal competence development

We developed a mathematical model of pneumococcal competence based on the

network of protein interactions known to regulate competence development (Figure 1)

during growth in a well-mixed liquid medium. Briefly, the precursor of CSP, ComC, is

cleaved and exported to the extracellular space by the membrane protein complex

ComAB [9,21]. Upon binding to CSP, ComD phosphorylates the response regulator

ComE, which in its phosphorylated form upregulates transcription of the operons

comAB, *comCDE* and *comX* [9,22,23]. The latter encodes the sigma factor ComX, which

controls transcription of genes required for uptake and processing of exogenous DNA

[7,24]. Our model uses ordinary differential equations (ODEs) and consists of two

components. At the population-level it keeps track of the population density and the

extracellular concentration of CSP; at the cell-level it keeps track of the intracellular

concentrations of the proteins involved in competence regulation (Figure 1). All the

cells export CSP to the medium at a rate determined by the intracellular concentrations

of ComC and ComAB. The concentration of CSP then feeds back into the intracellular

concentrations of all the proteins involved in competence since their transcription rates

depend on the ratio of ComE to ComE~P and thus on the rate at which ComD

phosphorylates ComE. Different environmental scenarios are simulated by changing

model parameters according to the known effects of such environmental factors in the

competence regulatory network. Since the model is primarily concerned with

competence initiation we purposely left out genes crucial for other aspects of

competence development (e.g. the stabilizing factor ComW and the immunity gene

comM) [17] and genes involved in competence shut-off such as DprA [25]. A detailed

description of the model and the choice of parameter values is provided in

Supplementary Note 1.

We use the model to determine the effect of environmental factors and cell

history (i.e. environments experienced in the past) on the relationship between cell

density and CSP concentration. Crucially, the model assumes that all cells are

homogeneous and that competence is only regulated by CSP, whose production

increases with cell density since cells release all CSP they produce to a common

extracellular pool. We are interested in determining whether these assumptions are

sufficient to explain our experimental results in well-mixed cultures or if additional

mechanisms need to be incorporated (e.g. density-independent competence induction

and cell-cell contact dependent competence transmission [15,16,19]).

Competence develops at a critical CSP concentration

It has been reported that competence develops at a fixed time after inoculation from

acid to alkaline conditions (pH 6.8 → 7.9) regardless of the inoculum size [15,19]. This

observation has motivated the view that competence develops independently of cell

density and rather acts as a timed response at the single cell level to the pH shift

occurring at the moment of inoculation. We extended previous studies by exploring a

wider range of inoculation densities (OD$_{595nm}$: 10^{-1} to 10^{-7}) (approximately 10^8 to 10^2

cells ml^{-1}) and preculturing conditions. We used the encapsulated serotype 2 strain *S.*

pneumoniae D39 [26] and cells were washed before inoculation to remove CSP produced

during the preculture. Importantly, we verified that CSP is actually present in the

supernatant of competent cultures of strain D39 (Supplementary Fig. 1). To monitor

competence development, the ComX-dependent promoter of the late competence gene

ssbB was fused to the firefly *luc* gene and inserted at the non-essential *bgaA* locus.

Activation and expression of *ssbB* is a good reporter for competence development

since SsbB expression strongly correlates with actual transformation with externally

added DNA (e.g. refs. [13,27]).

As shown in Figure 2a, we find that the inoculation density in strain D39 does have

an effect on the time of competence development, with competence initiating later for

lower inoculum sizes. For instance, for the lowest inoculation density, competence

initiates more than 4 hours later than for the highest inoculation densities (Figures 2a

and left panel of 2c). Note that our luminometer can detect light from competent cells

at an OD_{595} of 1.56×10^{-3} or higher (Supplementary Fig. 2) and therefore we cannot

exclude the possibility that a very small subpopulation of cells initiates competence

before we can detect it. Nevertheless, in all cases our estimates of the density of

competence initiation are far higher than the detection threshold, indicating that

competence in the majority of the population had not developed before crossing the

density threshold (Figure 2c, right panel).

Importantly, we observe that the population density at competence initiation is not

constant but positively related to the inoculation density. Hence, the dependency of the

time of competence initiation on the inoculation density is not a consequence of

competence developing at a fixed critical cell density for every condition. Instead, our

results are consistent with the mathematical model, which predicts that competence

develops when the CSP concentration has reached a critical threshold. The model

shows that competence will start faster for higher inoculation densities because the

CSP concentration reaches the critical threshold for competence activation earlier if

more cells are producing CSP (Figure 2d, left panel). Moreover, the model shows that

populations inoculated at low densities initiate competence at a lower density than

populations inoculated at high densities consistent with the experimental data (right

panels of Figures 2c and 2d). This is because cells inoculated at low cell densities

already had time to start transcribing competence regulatory genes and accumulate

some CSP once they reached the same cell density of cultures freshly inoculated at a higher cell density (Figure 2e). Thus, the critical CSP threshold is reached sooner for low-density inoculated cultures. Notably, a common misconception in the field is that in a QS system the critical concentration of autoinducer should always be attained at the same fixed cell density [15,16,19,20].

It is well known that the pH of the medium affects competence development, with natural competence being inhibited under acid conditions [11,28]. Under our experimental conditions, competence only naturally develops in alkaline growth medium with a pH > 7.4. So far, we have studied competence with cells precultured in a non-permissive pH for competence development (pH 6.8). These preculture conditions were reflected in the model simulations by assuming that cells initially were in the competence-off state. We also simulated the alternative scenario that cells are already competent at inoculation. For this cell history, the model predicts that the time of competence initiation is lower, but only for high inoculation densities (Figure 2d, left panel). This happens because when cells are competent initially and are inoculated at high density they can produce enough CSP to remain competent (Supplementary Fig. 3). However, when inoculation density is low, cells cannot produce enough CSP and initial competence switches off. The timing of the subsequent competence initiation is then the same as if cells were not competent when inoculated.

To verify the predicted effect of cell-history on the timing of competence initiation, we controlled the competence state of cells at inoculation by manipulating the pH during preculture. Specifically, we compared the time of competence initiation for cells coming from a non-permissive (pH 6.8) and a permissive (pH 7.9) pH history for competence development. For inoculation densities below OD_{595nm} of 0.01, the pH of the preculture did not have an effect on the timing of competence initiation as

predicted by the model (Figures 2b and c). On the other hand, for inoculation densities

above OD_{595nm} 0.01, there was a time delay in competence initiation for cells with an

acid history whereas cells with a non-acid history were competent when inoculated and

remained competent afterwards. This suggests that when the inoculation density is

high, there are enough cells to take the CSP concentration above the threshold for

competence activation if they are already producing CSP – as is the case of cells

coming from a non-acid history. By contrast, if cells come from a non-permissive pH

for competence development, the machinery for CSP production needs to be activated.

This causes a delay in competence initiation at high cell densities, which, at least for

our strain, would not result from regulation by a cell-density independent timing

device[15,16,19].

To further study competence in conditions closer to what *S. pneumoniae*

experiences in nature, we monitored competence development at the single-cell level

in microcolonies growing on a semi-solid surface. To this end, *ssbB* was fused to *gfp*

and competence initiation was followed using automated fluorescence time-lapse

microscopy. We observed that competence synchronizes in neighboring microcolonies

that are not in direct physical contact with each other (Figures 3A, 3B and

Supplementary Movie 1). In addition, more than 95% of the population became

competent in sharp contrast with competence development in other species such as

Bacillus subtilis where less than 20% of the population enters the competent state [29,30].

Both observations are consistent with the view that CSP diffuses extracellularly and

drives competence development across the population. We provide experimental

evidence for this claim by studying mixed populations of our wildtype D39 strain and

a *comC*⁻ mutant, which is unable to produce and export CSP, and therefore only

develops competence in the presence of external CSP. We followed competence

development in these mixed populations and found that competence can propagate

from the wildtype to the $comC^-$ mutant without the need of cell-to-cell contact (Figure

3C and Supplementary Movie 2). Note that the $comC$- mutant does not activate

competence when grown alone (Supplementary Fig. 4 and Supplementary Movies 3

and 4). This implies that CSP can diffuse extracellularly which confirms our previous

finding in liquid culture that CSP can be recovered from the supernatant of a

competent culture (Supplementary Fig. 1). It is worth noting that although cells release

CSP to an extracellular pool, a fraction might remain attached to them or in close

proximity due to diffusivity on the polyacrylamide surface. This can introduce

variation in the microenvironment that different cells experience and therefore in the

extent of synchronization of competence initiation.

In the experiments discussed so far, we used an encapsulated strain, in contrast to

the studies of Claverys et al. [15] and Prudhomme et al.[19], which were mainly based on

unencapsulated strains. We explored whether this could explain our different

observations regarding the effect of inoculation density on competence development

by studying additional strains: an unencapsulated version of our strain D39, the clinical

isolate PMEN14 together with its unencapsulated version and S. mitis, which is

naturally unencapsulated. Although we still observe that competence develops later for

smaller inoculums, the slope of the RLU signal decreases with inoculation density for

the two capsule knockouts and S. mitis, in agreement with the results of Claverys et

al.[15] and Prudhomme et al.[19] (Supplementary Fig. 5). A decreasing slope of the RLU

signal likely indicates decreased synchronization in competence development at lower

inoculation density. Since cells are synchronized by the common extracellular CSP

pool, a possible scenario is that the absence of a capsule impedes cells from exporting

all the CSP they produce. This would translate into less synchronization especially at

low inoculation densities. In the extreme scenario where cells would not share any

CSP with other cells, competence regulation would be in fact independent from cell

density. Note, however, that for the capsule knockout of our strain we can still detect

CSP in the cell-free supernatant and also observe competence propagation in the

absence of cell-cell contact (Supplementary Fig. 6 and Supplementary Movie 5).

pH and competence development

In order to understand how environmental factors affect competence, we quantified the

effect of external pH on natural competence development. We studied competence at a

fine-grained range of pH values from 6.8 to 8.5 and found a clear-cut value that

separated permissive from non-permissive external pH values for natural competence

development as reported before [11,27]. For our media this was pH 7.4 (Figure 4a).

However, not only competence always developed at pH higher than 7.4 but the critical

cell density for competence initiation decreased with increasing pH (Figure 4a and b).

Therefore, pH does not relate to competence as a binary permissive/non-permissive

condition but competence development is more efficient in more alkaline media. The

data suggests that for non-permissive pH conditions the cell density at which

competence would initiate is above the carrying capacity of the medium, which was

also previously proposed by Chen and Morrison[11].

To study the effect of pH on competence we deleted *comC* from the *comCDE*

operon and put it under the control of an IPTG-inducible promoter [31] at an ectopic

locus. We then tested competence development in this strain at different pH values

(7.2, 7.4, 7.6) and at varying IPTG concentrations. For any given IPTG concentration

competence always developed earlier at higher pH. This indicates that cells need to

express more *comC* to reach the critical CSP threshold for competence activation the

lower the pH (Supplementary Fig. 8). In fact, at low IPTG concentrations, competence

only develops above pH 7.4. The same pattern is observed in an IPTG-inducible

comCDE strain: for a fixed level of *comCDE* expression, the time of competence

initiation still decreases with the pH (Supplementary Fig. 8). Remarkably, in this

genetic background competence hardly develops at pH 7.2 even when *comCDE* is fully

induced. Cells might need to express more *comC* at lower pH because CSP export

and/or detection reduces with decreasing pH. This would also explain why competence

barely develops at low pH in the IPTG-inducible *comCDE* strain since for a fixed level

of *comE* expression, reduced CSP export and/or detection would bias the

ComE~P/ComE ratio towards no competence development. We tested whether pH

affects CSP detection by using a *comC*⁻ mutant. We performed experiments with

medium at different initial pH values where we added various concentrations of

synthetic CSP, using the *comC*⁻ mutant. We found that competence was mainly

dependent on the CSP concentration and only minor differences were found among

media with different pH (Supplementary Fig. 8). This suggests that competence

development is not mainly mediated by pH-dependent CSP detection so possibly it is

mediated by pH-dependent export. As peptidase-containing ATP-binding cassette

transporters such as ComAB require ATP to transport substrates [32], it might be that the

proton motive force influences its activity. Therefore, we incorporated the effect of pH

in our model by changing the rate at which cells export CSP. In agreement with the

experimental results, the modified model confirms that the density of competence

initiation decreases with the rate of export of CSP and thus with higher pH. Also, the

model predicts that for rates of CSP export below a certain threshold competence does

not develop any more since cells never manage to accumulate enough CSP for

competence to initiate (Figure 4c). Note however that this is a simplification of the

effect of pH in competence regulation since pH might also affect ComD and/or the

stability of CSP (as in other QS systems [6]) and as our data suggest it might also be

involved in the shutdown of competence (Supplementary Fig. 8). However, regardless

of the exact mechanism, as long as higher pH increases the rate at which single cells

produce and/or sense CSP, our model predicts that the density at which the critical

CSP concentration for competence activation is attained will decrease with increasing

pH (see Section C of the Supplementary Note 1 for a simple mathematical argument).

Finally, we assessed the joint effect of pH and cell density on competence

regulation. We did this by studying competence initiation for cultures inoculated at

different cell densities in media with different pH both experimentally and using the

model. The model predicts that competence will initiate earlier both for higher

inoculation densities and more alkaline pH (Figure 5a left panel): While higher

inoculation densities mean that more cells will start producing CSP after inoculation,

higher pH increases the rate at which individual cells produce CSP. The experimental

data is consistent with this prediction (Figure 5b left panel). Therefore, the observation

that pH affects competence development is not conflicting with regulation by cell

density because the CSP concentration depends on both of these factors.

Induction of competence by antibiotics

The induction of pneumococcal competence is affected by the presence of certain

classes of antibiotics [13,27], which has been considered additional evidence for the

hypothesis that competence can be regulated independently of cell density [13,15]. We

evaluated this claim by studying the role of HPUra and streptomycin on competence

regulation. We chose these antibiotics since the mechanisms by which they induce

competence at the molecular level have been elucidated to some extent: HPUra stalls

replication forks during DNA replication while initiation of DNA replication

continues, thereby increasing the copy number of genes near the origin of replication

(*oriC*). As a consequence, it up-regulates transcription of *comAB, comCDE* and *comX*

as these operons are located proximal to *oriC* [27]. Streptomycin causes mistranslation

and is thought to regulate competence via the membrane protease HtrA which targets

misfolded proteins and also represses competence possibly by degrading CSP [33] (also

see Supplementary Fig. 9). By increasing the amount of misfolded proteins,

streptomycin could reduce the rate at which CSP is degraded by HtrA leading to

competence induction.

We reproduced the effect of HPUra and streptomycin on competence regulation in

our model by increasing the transcription rate of *comAB, comCDE* and *comX* and by

reducing the rate at which CSP degrades, respectively. Our model predicts that the

presence of antibiotics lowers the pH threshold for competence development (Figure

5a) since antibiotics can counteract the effect of acidic pH to the point that cells can

still accumulate enough CSP to become competent. They do this by increasing the rate

at which single cells produce CSP (reducing the number of cells needed to reach the

critical CSP concentration for competence initiation) or by increasing the rate at which

they sense CSP (reducing the critical CSP concentration for competence initiation).

Also, it predicts that for pH values where competence is already induced without

antibiotics, it will develop faster in the presence of antibiotics (Figure 5a). We tested

these predictions experimentally using antibiotics at concentrations that have a

minimal effect on growth since we did not incorporate growth reduction due to

antibiotic stress in the model. In agreement with previous studies [13,27,33] and with the

model predictions, we find that antibiotics can induce competence at pH values that are

repressive for natural competence development (Figure 5b and c). We also find support

for the second prediction of the model since for permissive pH values for natural

competence development (above 7.4), competence is induced earlier in the presence of

antibiotics (Figure 5b and d). Remarkably, both the model and the experiments show

that the combined effect of pH and cell density in the presence of antibiotics remains

the same as when no antibiotics are added (compare left panel with middle and right

panels in Figures 5a and 5b): Competence induction still occurs earlier for high

densities of inoculation and more alkaline pH values. In the case of HPUra at pH 7.3 it

is even possible to see that competence does not develop for the highest inoculation

density as the population probably reaches carrying capacity before enough CSP is

produced. Finally, note that there might be alternative mechanisms from the ones

incorporated in the model by which antibiotic stress affects competence. For instance,

antibiotics reduce growth and can induce stress responses that lead to global changes

on transcription and translation.

Bistable region for competence development and cell history

An important feature of the competence regulatory network is the presence of a

positive feedback that couples CSP detection to CSP production (Figure 1). Signaling

systems that contain positive feedback loops often exhibit switch-like responses

resulting in the occurrence of alternative stable states [34]. We varied the strength of the

positive feedback loop in the model by changing the rate of CSP export and found that

the competence regulatory network exhibits bistability for a range of intermediate CSP

export rates. In this range, the model predicts the existence of two alternative states

where competence switches 'ON' or 'OFF' depending on the initial conditions (Figure

6a).

Since in the model the rate of CSP export is positively correlated to the pH, we

expected to find a region of pH values exhibiting similar bistability as an additional

experimental corroboration of the model. Indeed, we found support for the existence of

a bistable region at pH 7.4 where the wild type developed competence if CSP was

externally added in concentrations above 4 ng mL^{-1} (Figure 6b). Thus, whereas

competence always switched on for pH values above 7.4 regardless of the initial CSP

concentration, for pH 7.4 both 'ON' and 'OFF' states were observed depending on the

initial CSP concentration. Moreover, at pH 7.4 competence developed with 4 ng mL^{-1}

of CSP a concentration that would not induce competence in the *comC*$^{-}$ mutant

(Supplementary Fig. 8), which indicates that CSP production in the wild type was

kick-started by the initial addition of CSP resulting in enough overall CSP for

competence induction.

Bistable systems usually exhibit hysteresis. For this reason, we expected that at pH

7.4 where both the 'ON' and 'OFF' states are attainable, cell history, would influence

competence induction. From our previous experiments we determined that cells

coming from acid preculture inoculated at pH 7.4 do not develop competence at any

density of inoculation (Figure 5b left panel, second column). We then studied whether

there is history-dependence by inoculating cells coming from non-acid preculture at

pH 7.4. We found that cells coming from a non-acid preculture became competent

when inoculated at densities above OD$_{595nm}$ of 2.4×10^{-4}, which demonstrates that cell

history can influence competence development at this pH (Figure 6c). Past history has

an effect on competence because it determines the state of the machinery for CSP

production, which is 'OFF' when cells come from acid preculture but 'ON' when they

come from non-acid conditions. This explains why the effect of non-acid cell history

appears from a minimum inoculation density, since enough cells need to be inoculated

in order for them to produce the amount of CSP necessary for the system to remain

'ON' (Figure 6a inset). We then hypothesized that at pH 7.3 the critical inoculation

density of cells coming from non-acid history would have to be even higher than the

one at pH 7.4 as the model predicted that a higher initial concentration of CSP would

be necessary for the system to remain 'ON' at lower pH. We confirmed this prediction

experimentally by showing that at pH 7.3 competence does not develop for an

inoculation density of OD_{595nm} 2.4 x 10^{-4} (as for pH 7.4) but from 7.4 x 10^{-4} upwards

(Supplementary Fig. 10). Thus, our results show that, as a consequence of the positive

feedback involved in CSP production, previous exposure to different environmental

conditions can determine whether competence is induced or not by modifying the state

of the machinery for CSP production and/or sensing.

Discussion

Recently, the view that bacteria use autoinducers as QS signals has been debated since

autoinducer concentration can change in response to the environment. Here, we show

experimentally that cell density, pH and antibiotic stress simultaneously regulate

competence development in *S. pneumoniae* (Figures 2-5), a system classically framed

in the paradigm of QS. Using a mathematical model, we show that this occurs because

pH and antibiotics modify the rates at which single cells produce and sense CSP and

therefore the strength of the positive feedback loop coupling CSP detection to CSP

production (Figures 4 and 5). This environmental dependency does not override

regulation by cell density but rather modulates the relationship between the number of

cells and the CSP concentration. A fundamental aspect to the dependency on cell

density is that cells share CSP with others. Importantly, here we provide evidence both

in liquid culture (Supplementary Fig. 1) and through single-cell observations (Figure

3) that CSP is exported to the extracellular space. Finally, we show that competence

development is history-dependent since past environmental conditions can modify the

status of the machinery to produce and respond to CSP determining whether

competence switches on or not (Figure 6). Hysteresis in the competence response

might be especially important in the natural niche of the pneumococcus, the human

nasopharynx. In particular it is consistent with the observation that there is constitutive

upregulation of competence in pneumococcal biofilms during nasopharyngeal

colonization [35]. In this context, once competence is triggered for the first time, cells

would be primed to rapidly initiate another round of competence.

Why is competence controlled by CSP? CSP does not act as a timing device in our

encapsulated strain since competence develops in a cell-density dependent manner

without the necessity of cell-cell contact [15,16,19] (Figures 2 and 3). Regarding the

hypothesis that CSP is a probe to test diffusion [20], our results suggest that focusing on

diffusion alone oversimplifies the information and functionality that cells can gather

through CSP production. We hypothesize that by using an autoinducer peptide,

bacteria can coordinate the development of competence and in particular the

expression of fratricins and bacteriocins, which are under the control of the competent

state. These proteins can lyse or inhibit the growth of surrounding cells that are not

competent, increasing the efficiency of genetic transformation and mediating

competition with other bacteria [36–39]. By coordinating competence expression via CSP,

an isogenic bacterial population can increase the total concentration of secreted

fratricins and bacteriocins in times where population density is high, which likely

translates into a higher amount of lysed cells and therefore potential DNA donors.

Importantly, coordinating competence expression can also prevent the killing of clonal

siblings since immunity to these proteins comes with the competent state. Note

however that the extent to which cells synchronize competence development may vary

depending on the strain, genotype and growth conditions. In particular, our results

suggest that unencapsulated strains may synchronize less, which would explain the

difference between the findings reported by Prudhomme et al.[19] and our study.

Decreased synchronization may result from cells exporting less CSP to the

extracellular space and keeping more to themselves. In fact, in other species like *S.*

thermophilus where competence is controlled by ComS, a peptide that rapidly gets

imported back into the cell, the rate of competence development decreases with the

inoculation density [40] as observed for the unencapsulated pneumococcal strains.

Finally, note that *S. pneumoniae* grows primarily in biofilms where there is

heterogeneity in the physiological status and microenvironment that different cells

experience. This can certainly influence the degree of synchronization in competence

initiation across a population especially in the light of our findings that both current

and past environmental conditions affect the competence regulatory network. Indeed,

recent work in *S. mutans* showed heterogeneous competence activation of cells within

biofilms and upon different environmental pH ranges [41,42]. While studies of well-mixed

cultures give insight into the response mechanism shared by all cells in a population,

additional work is needed in the future to study how CSP production and detection by

individual cells is shaped by their spatial context and history and to unravel how

individual responses translate into patterns of population synchronization across

different genotypes and strains.

What is the relevance of the information carried by CSP? Alkaline pH and

antibiotic stress can induce competence by increasing the rate at which single cells

produce and sense CSP. We expect this to be a general mechanism by which sources

of stress that are alleviated through competence induce this state (e.g. mobile genetic

elements as hypothesized by Croucher et al.[43]). Upregulating competence in the

presence of antibiotics can increase survival by activating the expression of stress

response genes [7,18], facilitating repair of damaged DNA and mediating acquisition of

resistance [18,44]. Our findings suggest that strategies to prevent competence

development in response to antibiotics can focus on counteracting the effect of

antibiotics on the rate at which cells produce or sense CSP. Regarding the benefits of

upregulating competence with alkaline pH, these are less clear and could be an

example of a non-adaptive response resulting from the inherent biochemical properties

of ComAB and possibly ComD. Importantly, CSP can integrate additional

environmental cues like oxygen availability through the CiaRH two component

system, which represses *comC* post-transcriptionally and is required for virulence

expression and host colonization[14,45]. In fact, CiaRH is key for the regulation of

competence by multiple environmental signals in other streptococci like *S. mutans* [46].

Our findings support the view that functional hypotheses stressing individual

factors like diffusion or population density underplay the complexity of information

integrated by QS systems [4,47–51]. Although the term 'quorum sensing' overemphasizes

the role of population density, we advocate for keeping it due to its widespread use and

the fact that density will modify autoinducer concentration in any autoinducer

production system. Crucially, QS should be used in a broad sense that acknowledges

that bacteria integrate past and current environmental factors in addition to population

density into their QS responses. This view might be very useful for other autoinducer

production systems like competence in *Vibrio cholerae*, where the synthesis of the

autoinducer, CAI-1, depends on the intracellular levels of cAMP-CRP and therefore

might incorporate information on the metabolic status of the cell [52,53]. Also in other

systems, clear links between signal production, quorum threshold and environmental

conditions have been shown to affect QS [54–58].

Given that many biotic and abiotic factors can modify autoinducer concentrations

[59], future work should aim to study the relevance of such factors in the natural context

where bacteria secrete autoinducers. Such work is crucial to assess whether

upregulating QS in response to a particular factor provides a benefit for bacteria or is

merely a result of the biochemical properties of the QS regulatory network. An

interesting possibility is that, as in other biological systems [60], bacteria could perform

collective sensing of the environment through social interactions. In this context, by

secreting autoinducers cells could share individual estimates of environmental

conditions (e.g. antibiotic stress) for which upregulating QS is beneficial. Then,

autoinducer secretion would provide a way to get a more reliable estimate of the

environmental conditions by allowing a population to pool estimates made by

individual cells. Importantly, such a role for autoinducer secretion would explain the

dependency of QS on both cell density and the environment.

Methods

Bacterial strains and growth conditions

All pneumococcal strains used in this study are derivatives of the clinical isolate *S.*

pneumoniae D39 [26] unless specified otherwise. To monitor competence development,

strains either contain a transcriptional fusion of the firefly *luc* and the *gfp* gene with the

late competence gene *ssbB* or a full translational *ssbB-gfp* fusion. Cells were grown in

in C+Y complex medium at 37ºC. C+Y was adapted from Adams and Roe [61] and

contained the following components: adenosine (65 µm), uridine (107 µM), L-

asparagine (331 µM), L-cysteine (71 µM), L-glutamine (150 µM), L-tryptophane

(29.4) µM, casein hydrolysate (5 g L^{-1}), BSA (8 mg L^{-1}), biotin (2.46 µM), nicotinic

acid (4.87 µM), pyridoxine (3.4 µM), calcium pantothenate (5.04 µM), thiamin (1.9

µM), riboflavin (0.744 µM), choline (48 µM), $CaCl_2$ (113 µM), K_2HPO_4 (48.8 mM),

$MgCl_2$ (2.46 mM), $FeSO_4$ (1.8 µM), $CuSO_4$ (2 µM), $ZnSO_4$ (1.74 µM), $MnCl_2$ (40

µM), glucose (11.1 mM), sodium pyruvate (2.7 mM), saccharose (944 µM), sodium

acetate (24.4 mM) and yeast extract (2.5 g L^{-1}).

Construction of recombinant strains

To transform *S. pneumoniae*, cells were grown in C+Y medium (pH 6.8) at 37ºC to an

OD_{595} of 0.1. Then, cells were treated for 12 minutes at 37ºC with synthetic CSP-1

(100 ng mL^{-1}) and incubated for 20 minutes at 30ºC with the transforming DNA. After

incubation with the transforming DNA, cells were grown in C+Y medium (pH 6.8) at

37ºC for 90 minutes. *S. pneumoniae* transformants were selected by plating inside

Columbia agar supplemented with 3% of defibrinated sheep blood (Johnny Rottier,

Kloosterzande, The Netherlands) and the appropriate antibiotics. Working stocks of

cells were prepared by growing cells in C+Y (pH 6.8) until an OD_{595} of 0.4. Cells were

collected by centrifugation (1595 x *g* for 10 minutes) and resuspended in fresh C+Y

medium (pH 6.8) with 15% glycerol and stored at -80ºC. See the details on the

construction of the strains below and see Supplementary Table 1 for a list of all the

strains used in this study.

Strain DSM2

To follow competence development during pneumococcal growth, a transcriptional

fusion of two reporter genes, *luc* (firefly luciferase) and *gfp*, to the late competence

promoter P$_{ssbB}$ was used. The plasmid pLA18 [27], containing the P$_{ssbB}$-*luc-gfp* construct

was transformed into *S. pneumoniae* D-PEP1 [62] and transformants were selected on

Columbia blood agar supplemented with 1 μg mL^{-1} tetracycline. The P$_{ssbB}$-*luc-gfp*

construct integrates into the *bgaA* locus, and correct transformants were verified by

PCR.

Strains ADP243 and ADP62

To prevent the production of CSP, the original *comC* gene was replaced by an

erythromycin resistance marker, leaving the promoter and therefore the polycistronic

nature of *comDE* intact. The upstream region was amplified using primers LAG11

(GGCGGATCCGGCAGTTTGTGTAATAGTAC) and ADP2/48+AscI

(ACGTGGCGCGCCGTTCCAATTTAACTGTGTTTTTCAT), the downstream

region with primers ADP2/47+BamHI

(ACGTGGATCCGAAATAAGGGGAAAGAGTAATGGATTTATTTG) and LAG54

(AATCGCCATCTTCCAATCCC), and the erythromycin resistance marker with

0292_ery_R+BamHI (GCGGATCCTGTCTTTGACCCAATCATTC) and

sPG19+AscI (ACGTGGCGCGCCCGGAGGAATTTTCATATGAAC). All three

fragments were digested with the proper restriction enzymes (AscI and/or BamHI) and

ligated. The Δ*comC::ery* fragment containing the erythromycin resistance marker

flanked by the sequence up- and downstream of *comC* without altering the natural

transcription of *comDE*- was transformed into D39 and DLA3[27] resulting in strains

ADP243 (Δ*comC::ery*) and ADP62 (Δ*bgaA*:: P$_{ssbB}$-*luc*, Δ*comC::ery*), respectively.

Transformants were selected on Columbia blood agar containing 0.25 μg mL^{-1}

erythromycin. Correct deletion of *comC* was verified by PCR and sequencing.

Strains ADP49 and ADP51

To monitor natural competence in two clinical isolates, the PCR product of the

fragment *ssbB-luc-kan* from strain MK134[27] was transformed into the pneumococcal

strain PMEN14 and the *Streptococcus mitis* strain NTCC10712. Transformants were

selected on Columbia blood agar containing 250 μg mL^{-1} kanamycin resulting in

strains ADP49 (PMEN14, *ssbB-luc*) and ADP51 (*S. mitis ssbB-luc*), respectively.

Unencapsulated strains ADP25, ADP26, ADP92

To delete the polysaccharide capsule, the whole *cps* operon was replaced with a

chloramphenicol   resistance   cassette.   The   primers   used   were: ADP1/51

(CGGTCTTCAGTATCAGGAAGGTCAG)   and   ADP1/52+AscI

(CGATGGCGCGCCCTTCTTTCTCCTTAATAGTGG) to amplify the upstream

region;   primers   ADP1/53+NotI

(CACGGCGGCCGCGAGAAAGTTTTAAAGGAGAAAATG)   and   ADP1/54

(GATAGAGACGAGCTGCTGTAAGGC) to amplify the downstream region;

sPG11+AscI   (ACGTGGCGCGCCAGGAGGCATATCAAATGAAC)   and

sPG12+NotI (ACGTGCGGCCGCTTATAAAAGCCAGTCATTAG) to amplify the

chloramphenicol cassette. Products were digested with the appropriate enzymes (AscI

and/or NotI) and ligated. Strains D39, DLA3 and ADP49 were transformed with the

ligation product and transformants were selected on Columbia blood agar containing

4.5 μg mL^{-1} chloramphenicol resulting in strains ADP25 (Δ*cps::chl*), ADP26 (Δ*bgaA*::

P$_{ssbB}$-*luc*, Δ*cps::chl*), ADP92 (PMEN14, *ssbB-luc*, Δ*cps::chl*).

Strains ADP249 and ADP151

To follow competence development of micro-colonies during time-lapse microscopy,

the PCR product of the C-terminal fusion of *ssbB* with *gfp* and a downstream

kanamycin cassette were amplified using chromosomal DNA of strain RA42 as a

template [63] and transformed into the D39 strain and into its unencapsulated variant

ADP25 resulting in strains ADP249 (P_{ssbB}-*ssbB-gfp*) and ADP151 (P_{ssbB}-*ssbB-gfp*,

Δ*cps::chl*), respectively. Transformants were selected on Columbia blood agar

containing 250 μg mL^{-1} kanamycin.

Strains ADP247 and ADP248

Plasmid pPEP43 [62] containing a constitutive promoter driving the expression of the red

fluorescent protein mKate2 (cytoplasmic localization) was transformed into strain D39

and its unencapsulated derivative ADP25. pPEP43 integrates into the *cep*-locus, and

transformants were selected on Columbia blood agar containing 125 μg mL^{-1}

spectinomycin. The resulting strains, ADP235 (Δ*cep::p3-mkate2*) and ADP244

(Δ*cep::p3-mkate2*, Δ*cps::chl*), were transformed with the PCR product P_{ssbB}-*ssbB-gfp*

including the kanamycin cassette (same as for strain ADP249). Transformants were

selected on Columbia blood agar containing 250 μg mL^{-1} kanamycin. Finally, the

resulting strains, ADP245 (Δ*cep::p3-mkate2*, P_{ssbB}-*ssbB-gfp*) and its unencapsulated

variant ADP246 (Δ*cep::p3-mkate2*, Δ*cps::chl*, P_{ssbB}-*ssbB-gfp*), were transformed with

the *comC::ery* fragment described above. This resulted in strains ADP247 (Δ*cep::p3-*

mkate2, P_{ssbB}-*ssbB-gfp*, Δ*comC::ery*) and ADP248 (Δ*cep::p3-mkate2*, Δ*cps::chl*, P_{ssbB}-

ssbB-gfp, Δ*comC::ery*). Transformants were selected on Columbia blood agar

containing 0.25 μg mL^{-1} erythromycin.

Strain ADP95

To use an IPTG-inducible system, we first transformed the codon optimized *lacI* gene

[31] into strain DLA3 [27]. For that, we PCR-ed the fragment including *lacI* integrated into

the *prsA*-locus together with a gentamycin resistance cassette from chromosomal DNA

of strain DCI23 [31] using primers OLI40 (CCATGGCATCAGCGAGAAGGTGATAC)

and OLI41 (GCGGCCGCAGGATAGAAAGGCGAGAG) and transformed the

resulting PCR product to strain DLA3 while selecting for gentamycin (20 μg mL^{-1})

resulting in strain ADP95 (D39, Δ$bgaA$::P$_{ssbB}$-luc, Δ $prsA$::$lacI$).

Strains ADP112 and ADP107

To control the production of $comC$, and thereby CSP, a strain was constructed with an

ectopic copy of $comC$ (at the cep locus) under control of the IPTG-inducible promoter

P$_{lac}$ [31], and the subsequent deletion of the original $comC$ (replaced by an erythromycin

resistance cassette). The inducible system was created using BglFusion cloning [62]. To

amplify   the $comC$ fragment,   primers   ADP2/38

(CAGTGGATCCGGTTTTTGTAAGTTAGCTTACAAG)   and   ADP3/31

(CAGTCTCGAGCCCAAATCCAAATAAATCCATTAC) were used using D39

chromosomal DNA as a template. To control the production of $comCDE$, a strain with

an IPTG-inducible $comCDE$ was created, with the deletion of the original $comCDE$

locus. The inducible system was created as explained above. To amplify the $comCDE$

fragment,   primers   ADP2/38

(CAGTGGATCCGGTTTTTGTAAGTTAGCTTACAAG)   and   ADP2/58

(ACGTCTCGAGGCGGCCGCCAATTTCTTGCTAATTGTC) were used using

chromosomal DNA of D39 as a template. PCR products were digested with BglII and

XhoI and were ligated with similarly digested pPEP1 plasmid containing the promoter

P$_{lac}$ [31]. The ligations were transformed into strain ADP95 and transformants were

selected on Columbia blood agar containing 125 μg mL^{-1} spectinomycin. To these

strains, either the PCR product Δ$comC$::ery (see above) or a PCR product containing

Δ$comCDE$::chl from strain MK135 [27] was transformed and transformants were

selected on Columbia blood agar containing either 0.25 μg mL^{-1} erythromycin or 4.5

µg mL^{-1} chloramphenicol resulting in strains ADP112 (ΔbgaA::P$_{ssbB}$-luc, Δcep::P$_{lac}$-

comC, ΔcomC::ery) and ADP107 (ΔbgaA::P$_{ssbB}$-luc, Δcep::P$_{lac}$-comCDE,

ΔcomCDE::chl), respectively.

Strain MK356

To construct MK356 (ΔbgaA:: P$_{ssbB}$-luc-gfp, ΔhtrA::ery), htrA was replaced with an

erythromycin resistance gene. A region upstream of htrA was amplified from genomic

DNA of S. pneumoniae D39 using primers htrA-up-F (5'-

GAACCTGCGACCGTTCGCTTAGAAGG-3') and htrA-up-R+BamHI (5'-

GCGCGGATCCTCCATATGTTTGAATTACTG-3') and a region downstream of htrA

was amplified with primers htrA-Dwn-F-EcoRI (5'-

CGCGGAATTCGACATCTATGTAAAGAAAGC-3') and htrA-Dwn-R (5'-

GCTGTTGATAATTCTACTATATTCTTC-3'). An erythromycin resistance gene was

amplified from plasmid pORI28 [64] using primers EmR-F+BamHI (5'-

GCGCGGATCCTATGAACGAGAAAAATATAAAACAC-3') and EmR-R+EcoRI

(5'-CGCGGAATTCGCAGTTTATGCATCCCTTAACTTAC-3'). The three fragments

were digested with appropriate restriction enzymes (BamHI and/or EcoRI) and ligated.

The ΔhtrA::ery fragment, containing the an erythromycin resistance gene flanked by

the sequence up- and downstream of htrA, was transformed into strain DSM2.

Transformants were selected on Columbia blood agar containing 0.25 µg mL^{-1}

erythromycin. Correct deletion of htrA was verified by PCR.

Density and luminescence assays

Cells were pre-cultured in acid C+Y (pH 6.8) or in non-acid C+Y (pH 7.9) at 37ºC to

an OD$_{595nm}$ of 0.1. Right before inoculation, they were collected by centrifugation

(8000 rpm for 3 minutes) and resuspended in fresh C+Y (pH 7.9). Unless indicated

otherwise all experiments were started with an inoculation density of OD_{595} 0.002 with

cells from an acid preculture. Luciferase assays were performed in 96-wells plates with

a Tecan Infinite 200 PRO luminometer at 37ºC as described before [62]. Luciferin was

added at a concentration of 0.5 mg mL^{-1} to monitor competence by means of luciferase

activity. Optical density (OD_{595nm}) and luminescence (relative luminescence units

[RLU]) were measured every 10 minutes. The time and density of competence

initiation correspond to the first time point where the RLU signal is equal or above 200

units. RLU is used instead of RLU/OD for Figure 2 because 1) when competence

develops the rate at which the RLU signal increases is faster than the growth rate and

2) due to the very low inoculation densities used for Figure 2 the RLU/OD can be very

high at the start (clearly before competence has developed). The value of 200 units was

chosen because once this value is reached competence always developed. The effect of

pH on competence development was studied by inoculating cells in C+Y at a range of

pH values from 6.8 to 8.5. pH was adjusted by adding HCl and NaOH. The effect of

antibiotics was studied by adding streptomycin (3 µg mL^{-1}) and HPUra (0.075 µg mL^{-}

^{1}) to C+Y.

Detection of CSP in cell free supernatant

S. *pneumoniae* D39 wild-type and its *comC* deficient version, ADP243, were grown in

non-acid C+Y (pH 7.9) at 37ºC to an OD_{595nm} of 0.1. Cells were spun down by

centrifugation at 20000 x g for 5 min. The supernatant was sterilized by filtering twice

through 0.2 µm filters. The supernatant was plated on Columbia blood agar to confirm

that no viable cells were present. For both the wild-type and the *comC*-deficient strain,

the cell-free supernatant was diluted 1:1 with 2x concentrated C+Y medium containing

luciferin, and pH was adjusted to the indicated value by addition of HCl. The indicator

strain DSM2 pre-grown in C+Y pH 6.8 was then inoculated, and growth and luciferase

activity was monitored as described before.

Time-lapse fluorescence microscopy

A polyacrylamide slide was used as a semi-solid growth surface to spot the cells for

time-lapse microscopy. This slide was prepared with C+Y (pH 7.9) and 10%

acrylamide as reported previously [65]. Cells were pre-cultured in acid C+Y (pH 6.8) and

right before inoculation on the slide they were resuspended in fresh C+Y (pH 7.9) as

explained before. Phase contrast, GFP and RFP images were obtained using a

Deltavision Elite microscope (GE Healthcare, USA). Time-lapse videos were recorded

by taking images every 10 minutes after inoculation unless specified otherwise. File

conversions were done using Fiji and analysis of the resulting images was done using

Oufti [66].

Mathematical model

A mathematical model of the network of competence regulation (Figure 1) was

developed as a system of ODEs. The model incorporates the protein interactions

involved in sensing CSP via the two-component system formed by ComD and ComE

and exporting it via ComC and ComAB. Additionally, it explicitly models the

interaction of ComE and ComE~P with the gene promoters of *comAB*, *comCDE* and

comX. This is important since ComE~P binds these promoters as a dimer introducing

non-linearity into the system, which underlies the observed bistability. Population

growth is logistic and it is assumed that cells are homogeneous and that they export all

CSP they produce to a common extracellular pool. See Supplementary Note 1 for the

equations and further description.

Data availability

The authors declare that the data supporting the findings of the study are available in

this article and its Supporting Information files, or from the corresponding authors

upon request.

Author Contributions

SMG, AD, RAS, MK, GSvD, and JWV designed research; SMG, AD, RAS and MK performed experiments; SMG and GSvD developed the model; SM, AD, RAS, MK, FJW, GSvD, and JWV analyzed data; and SMG, FJW, GSvD and JWV wrote the paper.

Acknowledgments

We thank Martin Ackermann, Melanie Blokesch and two anonymous reviewers for helpful comments on an earlier version of this manuscript before submission to this journal, and Katrin Beilharz for the *htrA::ery* construct. SMG and GSvD were supported by Starting Independent Researcher Grant 309555 of the European Research Council and a VIDI fellowship (864.11.012) of the Netherlands Organization for Scientific Research (NWO). Work in the Veening lab is supported by the EMBO Young Investigator Program, a VIDI fellowship (864.12.001) and ERC starting grant 337399-PneumoCell.

Conflicts of interest

The authors state no conflicts of interest.

References

1. Fuqua, W. C., Winans, S. C. & Greenberg, E. P. Quorum sensing in bacteria: the

LuxR-LuxI family of cell density-responsive transcriptional regulators. *J.*

Bacteriol. **176,** 269–275 (1994).

2. Waters, C. M. & Bassler, B. L. Quorum sensing: cell-to-cell communication in

bacteria. *Annu. Rev. Cell Dev. Biol*. **21,** 319–346 (2005).

3. Redfield, R. J. Is quorum sensing a side effect of diffusion sensing? *Trends*

Microbiol **10,** 365–370 (2002).

4. Hense, B. A. *et al*. Does efficiency sensing unify diffusion and quorum sensing?

Nat. Rev. Microbiol. **5,** 230–239 (2007).

5. Alberghini, S. *et al*. Consequences of relative cellular positioning on quorum

sensing and bacterial cell-to-cell communication. *FEMS Microbiol. Lett*. **292,** 149–

161 (2009).

6. Decho, A. W. *et al*. Autoinducers extracted from microbial mats reveal a surprising

diversity of N-acylhomoserine lactones (AHLs) and abundance changes that may

relate to diel pH. *Environ. Microbiol*. **11,** 409–420 (2009).

7. Peterson, S. N. *et al*. Identification of competence pheromone responsive genes in

Streptococcus pneumoniae by use of DNA microarrays. *Mol. Microbiol*. **51,** 1051–

1070 (2004).

8. Håvarstein, L. S., Coomaraswamy, G. & Morrison, D. A. An unmodified

heptadecapeptide pheromone induces competence for genetic transformation in

Streptococcus pneumoniae. Proc. Natl. Acad. Sci. U. S. A. **92,** 11140–11144

(1995).

9. Pestova, E. V., Håvarstein, L. S. & Morrison, D. A. Regulation of competence for

genetic transformation in *Streptococcus pneumoniae* by an auto-induced peptide

pheromone and a two-component regulatory system. *Mol. Microbiol.* **21,** 853–862

(1996).

10. Håvarstein, L. & Morrison, D. Quorum sensing and peptide pheromones in

streptococcal competence for genetic transformation. in *Cell-cell signaling in*

bacteria 9–26 (ASM Press, 1999).

11. Chen, J. D. & Morrison, D. A. Modulation of competence for genetic

transformation in *Streptococcus pneumoniae. J. Gen. Microbiol.* **133,** 1959–1967

(1987).

12. Claverys, J.-P. & Havarstein, L. S. Extracellular-peptide control of competence for

genetic transformation in *Streptococcus pneumoniae. Front. Biosci. J. Virtual Libr.*

7, d1798-1814 (2002).

13. Prudhomme, M., Attaiech, L., Sanchez, G., Martin, B. & Claverys, J. P. Antibiotic

stress induces genetic transformability in the human pathogen *Streptococcus*

pneumoniae. Science **313,** 89–92 (2006).

14. Echenique, J. R., Chapuy-Regaud, S. & Trombe, M. C. Competence regulation by

oxygen in *Streptococcus pneumoniae*: involvement of *ciaRH* and *comCDE. Mol.*

Microbiol. **36,** 688–696 (2000).

15. Claverys, J.-P., Prudhomme, M. & Martin, B. Induction of competence regulons as

a general response to stress in gram-positive bacteria. *Annu. Rev. Microbiol.* **60,**

451–475 (2006).

16. Johnston, C., Martin, B., Fichant, G., Polard, P. & Claverys, J.-P. Bacterial

transformation: distribution, shared mechanisms and divergent control. *Nat. Rev.*

Microbiol. **12,** 181–196 (2014).

17. Johnsborg, O. & Havarstein, L. S. Regulation of natural genetic transformation and

acquisition of transforming DNA in *Streptococcus pneumoniae*. *FEMS*

MicrobiolRev **33,** 627–642 (2009).

18. Engelmoer, D. J. P. & Rozen, D. E. Competence increases survival during stress in

Streptococcus pneumoniae. *Evol*. *Int*. *J*. *Org*. *Evol*. **65,** 3475–3485 (2011).

19. Prudhomme, M., Berge, M., Martin, B. & Polard, P. Pneumococcal Competence

Coordination Relies on a Cell-Contact Sensing Mechanism. *PLoS Genet*. **12,**

e1006113 (2016).

20. Yang, J., Evans, B. A. & Rozen, D. E. Signal diffusion and the mitigation of social

exploitation in pneumococcal competence signalling. *Proc*. *Biol*. *Sci*. **277,** 2991–

2999 (2010).

21. Hui, F. M., Zhou, L. & Morrison, D. A. Competence for genetic transformation in

Streptococcus pneumoniae: organization of a regulatory locus with homology to

two lactococcin A secretion genes. *Gene* **153,** 25–31 (1995).

22. Ween, O., Gaustad, P. & Håvarstein, L. S. Identification of DNA binding sites for

ComE, a key regulator of natural competence in *Streptococcus pneumoniae*. *Mol*.

Microbiol. **33,** 817–827 (1999).

23. Martin, B. *et al*. ComE/ComE~P interplay dictates activation or extinction status of

pneumococcal X-state (competence). *Mol*. *Microbiol*. **87,** 394–411 (2013).

24. Martin, B., Garcia, P., Castanie, M. P. & Claverys, J. P. The *recA* gene of

Streptococcus pneumoniae is part of a competence-induced operon and controls

lysogenic induction. *Mol Microbiol* **15,** 367–379 (1995).

25. Mirouze, N. *et al*. Direct involvement of DprA, the transformation-dedicated RecA

loader, in the shut-off of pneumococcal competence. *Proc*. *Natl*. *Acad*. *Sci*. *U*. *S*. *A*.

110, E1035-1044 (2013).

26. Avery, A. T., MacLeod, C. M. & McCarty, M. Studies on the chemical nature of

the substance inducing transformation of pneumococcal types. Induction of

transformation by a desoxyribonucleic acid fraction isolated from Pneumococcus

type III. *J.Exp.Med*. **79,** 137–158 (1944).

27. Slager, J., Kjos, M., Attaiech, L. & Veening, J.-W. Antibiotic-induced replication

stress triggers bacterial competence by increasing gene dosage near the origin. *Cell*

157, 395–406 (2014).

28. Tomasz, A. & Mosser, J. L. On the nature of the pneumococcal activator

substance. *Proc Natl Acad Sci U A* **55,** (1966).

29. Maamar, H. & Dubnau, D. Bistability in the *Bacillus subtilis* K-state (competence)

system requires a positive feedback loop. *Mol.Microbiol*. **56,** 615–624 (2005).

30. Smits, W. K. *et al*. Stripping Bacillus: ComK auto-stimulation is responsible for

the bistable response in competence development. *Mol.Microbiol*. **56,** 604–614

(2005).

31. Liu, X. *et al*. High-throughput CRISPRi phenotyping identifies new essential

genes in *Streptococcus pneumoniae*. *Mol*. *Syst*. *Biol*. **13,** 931 (2017).

32. Lin, D. Y., Huang, S. & Chen, J. Crystal structures of a polypeptide processing and

secretion transporter. *Nature* **523,** 425–430 (2015).

33. Stevens, K. E., Chang, D., Zwack, E. E. & Sebert, M. E. Competence in

Streptococcus pneumoniae is regulated by the rate of ribosomal decoding errors.

mBio **2,** (2011).

34. Ferrell, J. E. Self-perpetuating states in signal transduction: positive feedback,

double-negative feedback and bistability. *CurrOpinCell Biol* **14,** 140–148 (2002).

35. Marks, L. R., Reddinger, R. M. & Hakansson, A. P. High levels of genetic

recombination during nasopharyngeal carriage and biofilm formation in

Streptococcus pneumoniae. *mBio* **3,** (2012).

36. Guiral, S., Mitchell, T. J., Martin, B. & Claverys, J. P. Competence-programmed

predation of noncompetent cells in the human pathogen *Streptococcus*

pneumoniae: genetic requirements. *Proc.Natl.Acad.Sci.U.S.A* **102,** 8710–8715

(2005).

37. Wei, H. & Håvarstein, L. S. Fratricide is essential for efficient gene transfer

between pneumococci in biofilms. *Appl. Environ. Microbiol.* **78,** 5897–5905

(2012).

38. Kjos, M. *et al*. Expression of *Streptococcus pneumoniae* Bacteriocins Is Induced

by Antibiotics via Regulatory Interplay with the Competence System. *PLoS*

Pathog. **12,** e1005422 (2016).

39. Wholey, W.-Y., Kochan, T. J., Storck, D. N. & Dawid, S. Coordinated Bacteriocin

Expression and Competence in *Streptococcus pneumoniae* Contributes to Genetic

Adaptation through Neighbor Predation. *PLoS Pathog.* **12,** e1005413 (2016).

40. Gardan, R. *et al*. Extracellular life cycle of ComS, the competence-stimulating

peptide of *Streptococcus thermophilus*. *J. Bacteriol.* **195,** 1845–1855 (2013).

41. Son, M., Ghoreishi, D., Ahn, S.-J., Burne, R. A. & Hagen, S. J. Sharply Tuned pH

Response of Genetic Competence Regulation in *Streptococcus mutan*s: a

Microfluidic Study of the Environmental Sensitivity of comX. *Appl. Environ.*

Microbiol. **81,** 5622–5631 (2015).

42. Shields, R. C. & Burne, R. A. Growth of *Streptococcus mutans* in Biofilms Alters

Peptide Signaling at the Sub-population Level. *Front. Microbiol.* **7,** 1075 (2016).

43. Croucher, N. J. *et al*. Horizontal DNA Transfer Mechanisms of Bacteria as

Weapons of Intragenomic Conflict. *PLoS Biol*. **14,** e1002394 (2016).

44. Cornick, J. E. & Bentley, S. D. *Streptococcus pneumoniae*: the evolution of

antimicrobial resistance to beta-lactams, fluoroquinolones and macrolides.

Microbes Infect. *Inst*. *Pasteur* **14,** 573–583 (2012).

45. Halfmann, A., Kovacs, M., Hakenbeck, R. & Bruckner, R. Identification of the

genes directly controlled by the response regulator CiaR in *Streptococcus*

pneumoniae: five out of 15 promoters drive expression of small non-coding RNAs.

Mol Microbiol **66,** 110–126 (2007).

46. Ahn, S.-J., Wen, Z. T. & Burne, R. A. Multilevel control of competence

development and stress tolerance in *Streptococcus mutans* UA159. *Infect*. *Immun*.

74, 1631–1642 (2006).

47. Lazazzera, B. A. Quorum sensing and starvation: signals for entry into stationary

phase. *Curr*.*Opin*.*Microbiol*. **3,** 177–182 (2000).

48. Platt, T. G. & Fuqua, C. What's in a name? The semantics of quorum sensing.

Trends Microbiol. **18,** 383–387 (2010).

49. West, S. A., Winzer, K., Gardner, A. & Diggle, S. P. Quorum sensing and the

confusion about diffusion. *Trends Microbiol*. **20,** 586–594 (2012).

50. Williams, P. & Cámara, M. Quorum sensing and environmental adaptation in

Pseudomonas aeruginosa: a tale of regulatory networks and multifunctional signal

molecules. *Curr*. *Opin*. *Microbiol*. **12,** 182–191 (2009).

51. Hense, B. A. & Schuster, M. Core principles of bacterial autoinducer systems.

Microbiol. *Mol*. *Biol*. *Rev*. *MMBR* **79,** 153–169 (2015).

52. Liang, W., Pascual-Montano, A., Silva, A. J. & Benitez, J. A. The cyclic AMP

receptor protein modulates quorum sensing, motility and multiple genes that affect

intestinal colonization in *Vibrio cholerae*. *Microbiol*. *Read*. *Engl*. **153,** 2964–2975

(2007).

53. Suckow, G., Seitz, P. & Blokesch, M. Quorum sensing contributes to natural

transformation of *Vibrio cholerae* in a species-specific manner. *J*. *Bacteriol*. **193,**

4914–4924 (2011).

54. Pai, A., Tanouchi, Y. & You, L. Optimality and robustness in quorum sensing

(QS)-mediated regulation of a costly public good enzyme. *Proc*. *Natl*. *Acad*. *Sci*.

U. *S*. *A*. **109,** 19810–19815 (2012).

55. Lee, J. *et al*. A cell-cell communication signal integrates quorum sensing and stress

response. *Nat*. *Chem*. *Biol*. **9,** 339–343 (2013).

56. Xavier, J. B., Kim, W. & Foster, K. R. A molecular mechanism that stabilizes

cooperative secretions in *Pseudomonas aeruginosa*. *Mol*. *Microbiol*. **79,** 166–179

(2011).

57. van Delden, C., Comte, R. & Bally, A. M. Stringent response activates quorum

sensing and modulates cell density-dependent gene expression in *Pseudomonas*

aeruginosa. *J*. *Bacteriol*. **183,** 5376–5384 (2001).

58. Dunlap, P. V. & Kuo, A. Cell density-dependent modulation of the *Vibrio fischeri*

luminescence system in the absence of autoinducer and LuxR protein. *J*. *Bacteriol*.

174, 2440–2448 (1992).

59. Boyer, M. & Wisniewski-Dyé, F. Cell-cell signalling in bacteria: not simply a

matter of quorum. *FEMS Microbiol*. *Ecol*. **70,** 1–19 (2009).

60. Berdahl, A., Torney, C. J., Ioannou, C. C., Faria, J. J. & Couzin, I. D. Emergent

sensing of complex environments by mobile animal groups. *Science* **339,** 574–576

(2013).

61. Adams, M. H. & Roe, A. S. A Partially Defined Medium for Cultivation of
Pneumococcus. *J. Bacteriol*. **49,** 401–409 (1945).

62. Sorg, R. A., Kuipers, O. P. & Veening, J.-W. Gene Expression Platform for
Synthetic Biology in the Human Pathogen *Streptococcus pneumonia*e. *ACS Synth*.
Biol. **4,** 228–239 (2015).

63. Aprianto, R., Slager, J., Holsappel, S. & Veening, J.-W. Time-resolved dual RNA-
seq reveals extensive rewiring of lung epithelial and pneumococcal transcriptomes
during early infection. *Genome Biol*. **17,** 198 (2016).

64. Leenhouts, K. *et al*. A general system for generating unlabelled gene replacements
in bacterial chromosomes. *Mol GenGenet* **253,** 217–224 (1996).

65. Sorg, R. A. & Veening, J.-W. Microscale insights into pneumococcal antibiotic
mutant selection windows. *Nat*. *Commun*. **6,** 8773 (2015).

66. Paintdakhi, A. *et al*. Oufti: an integrated software package for high-accuracy, high-
throughput quantitative microscopy analysis. *Mol*. *Microbiol*. **99,** 767–777 (2016).

67. Guiral, S., Hénard, V., Granadel, C., Martin, B. & Claverys, J.-P. Inhibition of
competence development in *Streptococcus pneumoniae* by increased basal-level
expression of the ComDE two-component regulatory system. *Microbiol*. *Read*.
Engl. **152,** 323–331 (2006).

 Figure legends

 Figure 1. Network of competence regulation in *S. pneumoniae*. ComC (C) binds the

 membrane protein complex ComAB, and it is processed and exported as CSP to the

 extracellular space. CSP binds to the histidine kinase ComD, which is located in the

 membrane as a dimer. Upon CSP binding, ComD autophosphorylates and transfers the

 phosphate group to the response regulator ComE [9,23]. The phosphorylated form of

 ComE (ComE~P) dimerizes and activates transcription of *comAB*, *comCDE* and *comX*

 by binding to their promoters [8,9]. Unphosphorylated ComE can also bind these

 promoters, repressing their transcription [23,67]. Synthesis of the alternative sigma factor

 ComX directs transcription of genes required for genetic transformation as well as

 other functions [7,24]. Two key features of this network are the presence of a positive

 feedback loop (since increasing CSP detection leads to increasing CSP production) and

 of non-linearity (since ComE~P interacts with the gene promoters as a dimer).

 Figure 2. Competence is regulated by cell density. (**a,b**) Growth curves (OD_{595nm})

 and competence expression measured as relative luminescence units (RLU) expressed

 from the promoter of the late competence gene *ssbB* for populations inoculated at a

 range of densities and grown in C+Y medium with initial pH 7.9. In **a**, cells were

 precultured in acid conditions (pH 6.8), while in **b** cells were precultured in non-acid

 conditions (pH 7.9). Four replicates are shown for each of seven inoculation densities

 (OD_{595nm}): 0.1 (Green), 0.05 (Red), 0.01 (Blue), 10^{-3} (Purple), 10^{-4} (Light green), 10^{-5}

 (Yellow) and 10^{-6} (Brown). Competence does not develop in cells coming from acid

 preculture and inoculated at a density of 0.1. (**c**) Effect of inoculation density on the

time until competence initiation (left panel) and the population density at which

competence was initiated (right panel). Competence initiation was defined as the time

where the RLU signal exceeded 200 units. Note that our luminometer has enough

sensitivity to detect light from competent cells at a density of 1.56×10^{-3} or higher even

if they correspond to a subpopulation (Supplementary Fig. 2). (**d**) Predictions of the

mathematical model concerning the effect of inoculation density on the timing of

competence initiation (left panel) and the density at which competence initiates (right

panel). In the model, competence initiation was defined as the time where the total

concentration of ComX times the population density exceeds 2000 units. Non-acid

preculture is simulated in the model by setting the initial amount of all proteins in the

competence regulatory network to the value they attain when cells are competent. (**e**)

The model predicts that populations inoculated at lower densities will reach a threshold

CSP concentration (dotted line) at a lower density than populations inoculated at

higher densities.

Figure 3. Competence can propagate by CSP diffusion without the necessity of

cell-to-cell contact. (**a**) Phase contrast and GFP images (left panel) of a set of

neighboring microcolonies of D39 with a fusion of the late competence gene *ssbB* to

gfp (P*ssbB*-*ssbB*-*gfp*). The first images (t = 0 min) were taken right after inoculation on

the microscopy slide and the second ones correspond to the moment of competence

initiation (t = 65 min). Images were taken every 5 minutes. Crucially, when

competence starts the microcolonies are not in direct physical contact with each other.

Analysis of the fluorescence signal across the entire population (right panel) shows

that once competence starts, after 60 minutes from inoculation, the distribution of

fluorescence intensity moves to higher intensity values through time. We set as a

threshold for counting competent cells a fluorescence signal 50% above background.

After 85 minutes, we counted 97 out of 99 cells as competent cells. (b) Distribution of

fluorescence signal within each microcolony in the window of competence initiation.

The microcolony numbers correspond to the ones indicated in the phase contrast image

in panel a). Competence initiates in all microcolonies within a window of 5 minutes.

(c) Time-lapse fluorescence microscopy tracking competence development in two

colonies with a fusion of the late competence gene *ssbB* to *gfp*: one formed by cells of

D39 (ADP249) and one formed by cells of a *comC* deficient D39 (ADP247). The two

strains are distinguishable because ADP247 constitutively expresses a red fluorescent

protein. Competence developed first in D39 and then it propagated to the *comC*[-]

mutant after 20 minutes without the necessity of cell-cell contact. Note that we

checked that the *comC* deficient D39 (ADP247) does not become competent by itself

when grown without D39 (Supplementary Fig. 4 and Supplementary Movies 3 and 4).

The scale bar is 5 µm in all images.

Figure 4. **Competence is upregulated by higher pH.** (**a**) Effect of initial medium pH

on growth curves (left panel) and the dynamics of competence expression (middle

panel). Competence expression was quantified as relative luminescence units (RLU)

normalized by cell density. In the right panel, competence expression is plotted in

relation to cell density. All populations were grown at the indicated initial pH and

inoculated at a density of OD_{595nm} 0.002. Three replicates are shown for each initial

pH. **b**) Effect of initial medium pH on the population density at which competence

was initiated (the density at which RLU exceeded 200 units). Competence did not

develop at pH 7.4 and below. Note that the indicated pH is the initial pH of the

medium, which does not stay constant due to acidification by growth (Supplementary

Fig. 7). Although the pH drops considerably in fully-grown cultures, acidification is

still minor at the density where competence develops. **c**) Predictions of the model on

the effect of the rate of CSP export, r_e, and thus the pH, on the density of competence

initiation. Competence does not develop anymore below a threshold rate of CSP

export.

Figure 5. Competence is simultaneously regulated by cell density, pH and

antibiotic stress. (**a,b**) Predictions of the mathematical model (**a**) and experimental

data (**b**) on the dependency of the time of competence initiation on inoculation density,

initial pH and antibiotic stress. The x-axis in **a** corresponds to the rate of CSP export in

the model, r_e, which is a proxy for pH. The color scales with the time of competence

initiation with more intense red corresponding to faster development of competence.

Black represents no competence development. In **b,** each box corresponds to the

average initiation time of three replicates. Both the model and the experimental data

show that competence develops faster at higher pH and higher inoculation densities. **c**)

Antibiotics induce competence at pH values that repress natural competence

development. **d**) At pH values that are not repressive for competence development (pH

> 7.4), competence develops faster in the presence of antibiotics. The stars indicate

which conditions are plotted in **c** and **d**. The concentrations of streptomycin and

HPUra are 3 μg mL^{-1} and 0.075 μg mL^{-1}, respectively. We chose these sub-MIC

concentrations to minimize the effect of antibiotic stress on growth.

1015

Figure 6. A bistable regime for competence development. (**a**) Extracellular

concentration of CSP in response to the rate of CSP export. The model predicts the

existence of a region where competence always switches on regardless of the initial

conditions (which would correspond to pH > 7.4) and of a bistable region (bordered by

the dashed lines). In the latter, the initial conditions can either switch on or not CSP

production and subsequently competence development. (Inset) In particular, the model

predicts that in this region non-acid cell history can allow competence development if

enough cells are inoculated since they can produce enough CSP to remain competent.

The inoculation densities are 0.1 (Blue) and 0.01 (Brown) and both the number of

ComX molecules (solid line) and the CSP concentration (dashed line) are shown. **b**)

Growth curves and competence expression measured as RLU units normalized by

density for cells coming from acid preculture (pH 6.8) and inoculated in medium at pH

7.4 with different initial concentrations of CSP. Three replicates are shown per

treatment and all the cultures are inoculated at OD_{595} 0.002. **c**) Growth curves and

competence expression measured as RLU units normalized by density for cells coming

from non-acid preculture (pH 7.9) and inoculated in medium at pH 7.4 at different

initial densities. Three replicates are shown per inoculation density. Competence does

not develop for cells inoculated at the same densities but coming from acid preculture

(pH 6.8) (Figure 5b, left panel, second column).

1035

a Acid preculture (pH 6.8 → pH 7.9)

b Non−acid preculture (pH 7.9 → pH 7.9)

c Experimental data

d Mathematical model

e

a Mathematical model

No antibiotic   Streptomycin   HPUra

Inoculation density

pH ($\times 10^{-5}$, r_e, export rate of CSP in s^{-1})

Initiation time (min)

b Experimental data

No antibiotic   Streptomycin   HPUra

Inoculation density

pH

Initiation time (min)

c Antibiotics induce competence at pH 7.4 (✳)

No antibiotic
Streptomycin
HPUra

pH: 7.4
I.D.: 2.2×10^{-3}

d Antibiotics speed up competence development (✳)

No antibiotic
Streptomycin
HPUra

pH: 7.8
I.D.: 0.02

a

'ON'

'OFF'

pH >7.4

[CSP] (y-axis): 0, 50, 100, 150

pH(×10⁻⁵, export rate of CSP, r_e) (x-axis): 0, 1, 2, 3, 4, 5

Inset: # of ComX (×10³, –): 0, 12.5, 25; [CSP] (--): 0, 50, 100; Time (min): 0, 400, 800; labels 0.1, 0.01

b Initial CSP concentration

Left: OD_{595} (0.01, 0.1, 1.0) vs Time (min) (0, 200, 400, 600, 800, 1000)

[CSP] (ng/mL): 20, 4, 0.8, None

Right: RLU/OD (10, 10^2, 10^3, 10^4, 10^5) vs Time (min) (0, 200, 400, 600, 800, 1000); 'ON', 'OFF'

c Cell history

Left: OD_{595} (0.01, 0.1, 1.0) vs Time (min) (0, 200, 400, 600, 800, 1000)

Inoculation density: 2.2×10^{-3}, 7.4×10^{-4}, 2.4×10^{-4}

Right: RLU/OD (10, 10^2, 10^3, 10^4, 10^5) vs Time (min) (0, 200, 400, 600, 800, 1000); 'ON', 'OFF'