Multiple factors influence population sex ratios in the Mojave Desert moss *Syntrichia*

caninervis[1]

Jenna T. Baughman[2,4], Adam C. Payton[3], Amber E. Paasch[2], Kirsten M. Fisher[2], and

Stuart F. McDaniel[3]

[2]Department of Biological Sciences, California State University, 5151 State University

Drive, Los Angeles, CA 90032, USA, [3]Department of Biology, University of Florida, 876

Newell Drive, Gainesville, FL 32611, USA

[4]Author for correspondence: Department of Integrative Biology and University and

Jepson Herbaria, University of California, 3040 Valley Life Sciences Building, Berkeley,

CA 94720, USA; e-mail: jbaughman@berkeley.edu; phone: 219-393-0179.

Female biased *Syntrichia caninervis* genetic sex ratio

[1]Manuscript received ________; revision accepted ________.

Acknowledgements

This work was supported by the California State University Department of

Biological Sciences Faculty-Student Achievement Fund to KMF, a Research Exchange

Grant to JTB from the NSF EDEN RCN (IOS 0955517), the NSF Louis Stokes Alliances

for Minority Participation (HRD 1363399) to AEP, the NIH-NIGMS Minority

Biomedical Research Support-Research Initiative for Scientific Enhancement program

(GM61331) to JTB, and an NSF grant (DEB 1541005) to SFM.

ABSTRACT

• *Premise of research:* Natural populations of many mosses appear highly female-

biased based on the presence of reproductive structures. This bias could be caused

by increased male mortality, lower male growth rate, or a higher threshold for

achieving sexual maturity in males. Here we test these hypotheses using samples

from two populations of the Mojave Desert moss *Syntrichia caninervis*.

• *Methods:* We used double digest restriction-site associated DNA (RAD)

sequencing to identify candidate sex-associated loci in panel of sex-expressing

plants. Next, we used putative sex-associated markers to identify the sex of

individuals without sex structures.

• *Key results:* We found an 18:1 phenotypic female:male sex ratio in the higher

elevation, moister site, and no sex expression at the drier, low elevation site. In

contrast, based on genetic data we found a 2:1 female bias in the low stress site

and only females in the high stress site. The area occupied by male and female

genets was indistinguishable.

• *Conclusions:* These data suggest that both differential mortality and sexual

dimorphism in thresholds for sex expression contribute to population sex ratio

biases, and that sex-specific life history and survival interact with environmental

stress to determine the frequency of sexual reproduction in *S. caninervis*.

Key words: bryophyte; Mojave Desert; moss; Pottiaceae; reproductive strategy; sex

expression; sex ratio; *Syntrichia caninervis*

INTRODUCTION

Female biased sex ratios are a common yet unexplained phenomenon in bryophyte populations (Bisang and Hedenäs, 2005). Most of the female biases documented in mosses to date have been based on counting the number of sexually mature male and female ramets in a population (but see McLetchie et al., 2001; Korpelainen et al., 2008; Bisang et al., 2010; Hedenäs et al., 2010; Bisang and Hedenäs, 2013; Bisang et al., 2014; Bisang et al., 2015; Hedenäs et al., 2016). In dioicous bryophytes, sex determination happens at meiosis with the production of spores (Bachtrog et al., 2011). A spore carrying the female sex-determining locus (U) will form a gametophyte that produces archegonia and ultimately bears the offspring embryo (a sporophyte). Spores with the male sex-determining locus (V) form a gametophyte that produces antheridia. Because the diploid sporophyte is produced by the union of a U-bearing egg with a V-bearing sperm, the sporophyte is always heterozygous at the sex-determining locus. As a consequence of meiotic segregation, therefore, the null expectation is a 1:1 sex ratio. Identifying the underlying mechanisms that generate the persistently female-biased phenotypic sex ratios in moss populations is an essential first step toward understanding the evolutionary causes and consequences of sex ratio bias in this ecologically important group of plants.

A female bias in the production of sexually mature ramets in a natural population could be caused by at least three distinct processes. First, an apparently female biased population sex ratio could simply be a product of faster female growth, as has been found in several species (Shaw and Beer, 1999; Stark et al., 2005a; McDaniel et al., 2008). In this case, female genotypes on average would produce more ramets and occupy larger

areas than male genotypes. Second, males might exhibit a lower frequency of sex

expression than females. If this were true, males would constitute a disproportionately

large fraction of the sterile plants in a population (termed 'the shy male hypothesis';

Mishler and Oliver, 1991; Stark et al., 2005a). Finally, a female bias could be causd by

elevated male mortality during spore production (McDaniel et al., 2007; Norrell et al.,

2014), establishment, or at some later point in the life-cycle. Regardless of the proximate

cause, elevated male mortality would decrease the amount of genetic diversity in males

relative to females.

One of the most extreme cases of sex ratio variation in mosses is in Mojave

Desert populations of *Syntrichia caninervis*. Previous data indicate that the phenotypic

sex ratio in *S. caninervis* populations ranges from 7:1 (Paasch et al., 2015) to 14 F:1 M

(Bowker et al., 2000; Bisang and Hedenäs, 2005), and that some populations lack male

sex expression entirely (Stark et al., 2001). Mojave Desert *S. caninervis* is extremely

desiccation tolerant and spends much of its life in an air-dried state, limiting all biological

functions to infrequent post-rainfall periods, primarily in cool winter months (Stark,

1997; Stark et al., 1998). Environmental variation that corresponds with changes in

timing and duration of this biologically active period appears to affect overall levels of

sex expression in a population. A survey of 890 *S. caninervis* individuals from a 10

hectare elevation gradient (Bowker et al., 2000) found that total percentage of expressing

individuals increased with elevation, and that male sex expression occured only at the

higher elevations, while lower elevation populations contained only a few expressing

females. In parallel with low levels of sex expression, sexual fertilization and production

of sporophytes is relatively rare, and established desert *S. caninervis* populations seem to

persist through vegetative cloning (Paasch et al., 2015).

Here we utilize restriction site-associated DNA genome sequencing (ddRADseq)

to identify the sex of sterile ramets and study the patterns of genetic variation in males

and females in two Mohave Desert populatoins of *S. caninervis*. We found extreme

female biased phenotypic sex ratios, consistent with previous studies, but less extreme

genetic biases, suggesting that males produce sex structures less frequently than females.

We also found lower genetic diversity in males than females, suggesting that the

remaining genetic sex ratio bias is best explained by elevated male mortality. Patterns of

genet size indicate that sexual dimorphism in growth rate is insufficient to explain the

female biased sex ratios in this species.

MATERIALS AND METHODS

Sample Collection— We collected male, female, and sterile *S. caninervis* samples

at a low-stress and a high-stress site in the Mojave Desert. The low-stress site is located

at an elevation of 1800 m in Sheep Creek Wash near Wrightwood, CA (34° 22' 33.85" N,

117° 36' 34.59" W), at the west edge of the Mojave Desert and the northern base of the

San Gabriel Mountains. The average high and low annual temperature is 16.8 °C and 1.7

°C, with an average annual precipitation of 49.4 cm (2007-2011, Wrightwood Weather

Station, NOAA National Climatic Data Center). This site experiences little foot traffic.

The high-stress site is at an elevation of 1257 m near Phelan, CA (34° 25' 29.80" N, 117°

36' 30.91" W), about nine miles northeast of the low-stress site. The average high and

low annual temperature is 27 °C and 10 °C, while average annual precipitation is 28.2 cm

(2005-2009, Phelan, CA, NOAA National Weather Service). This site is also highly

disturbed by foot traffic and erosion.

To establish a panel of plants of known sex to use to identify sex-linked molecular

markers, we isolated 11 *S. caninervis* female and 10 male ramets from the low stress site

in March and April of 2013. Field collections of *S. caninervis* were identified by leaf

morphology, color, and hair points then later confirmed with leaf cross-sections. In the

laboratory under a dissecting microscope patches were hydrated and screened for

presence of current or past antheridia and archegonia.

To estimate the phenotypic sex ratio of the low stress population, in May 2014 we

collected in 3-5 cm patches in three parallel 20 m linear north-south (N-S) transects, 10 m

apart from one another, collecting one patch every 2 m from a variety of shaded and

exposed microhabitats. In June 2014 we made collections from the high stress site in a

similar manner in two parallel 20 m N-S transects. Additionally, due to the highly

irregular distribution of the species in the high stres site, a third mini-transect was

sampled (beginning at about the 23 m mark of transect 2 and extending approximately 2

m), more densely (ca. every 500 cm) selecting samples in a northwest-southeast (NW-

SE) direction from a variety of microhabitats. To estimate the genetic sex ratio and

perform diversity analyses, a maximum of three individual sterile ramets were isolated

from each patch using a dissecting microscope, resulting in a total of 99 ramets from the

low stress site and, due to lower frequency of this species at this site, 42 ramets from the

high stress site, for a total of n = 141 individual ramets.

DNA Extraction and RADseq library preparation— Total genomic DNA was

extracted and isolated from 162 total ramets (21 of known sex, and 141 of unknown sex)

using a modified cetyl trimethyl ammonium bromide-beta mercaptoethanol (CTAB)

procedure(McDaniel et al., 2007). Prior to extraction, samples were ground dry to a fine

powder using a GenoGrinder 2010 bead shaker (SPEX CertiPrep, Metuchen, NJ). DNA

quality was evaluated for each sample by electrophoresis and quantified using a Qubit

Fluorometer (Invitrogen, Carlsbad, CA, USA) and samples' DNAs were normalized prior

to library preparation.

Illumina libraries were prepared for sequencing using a modified version of the

Peterson et al. (2012) protocol using the endonucleases EcoRI and MseI (New England

Biolabs, Ipswich, MA). Following double enzyme digestion, unique barcoded adaptors

were ligated to the resulting EcoRI cutsites and a non barcoded universal adaptor to the

MseI cutsite. Variable length barcodes of 8, 9, and 10 basepairs (bp) were used with each

barcode, differing by at least 4 bp. Illumina flowcell binding and sequencing primer sites

were added to the adaptor ligated fragments throught 10 cycles of PCR using NEB Q5

PCR mastermix (New England Biolabs, Ipswich, MA). Success of library construction

was evaluated throught agarose gel electrophoresis, after which 5 μL of each sample's

final library was pooled into a single tube for the 141 sterile samples and a separate tube

for the 21 samples of known sex. Size selection and sequencing were performed at the

University of Florida's Interdisciplinary Center for Biotechnology Research. Pooled

libraries were fractioned using Pippin ELF precision electrophoresis (Sage Science,

Beverly, MA) with the resulting 250-500 bp fraction being used for Illumina sequencing.

The sterile ramet sample library was sequenced using an Illumina NextSeq 500 at mid

throughput, producing 150 bp single-end reads. The library consisting of the 21 known-

sex ramets was pooled with libraries constructed in an identical manner for a non-related

study and sequenced on one lane of a HiSeq 3000 producing 100 bp single end reads.

Data analysis— Raw ddRAD sequence reads were assessed for quality with

fastQC (Gordon and Hannon, 2010). All manipulation of raw sequence reads was

performed using tools from the FASTX-Toolkit (Gordon and Hannon, 2010)

implemented in Galaxy on the University of Florida's Research Computing Cluster. To

eliminate low quality bases at the 3' end of reads, all sequences were trimed to 100 bp.

Reads were filterd for a minimum Phred quality score of 20 on at least 90% of the read.

All reads were then demultiplexed followed by the removal of the 5' barcode and EcoRI

enzyme cutsite.

RAD loci were assembled de novo using STACKS, version 1.24, (Catchen et al.,

2011; Catchen et al., 2013). Pertinent parameters for the STACKS pipeline were as

follows: a minimum of 2 reads required for each allele within an indivdial (-m option

ustacks), a maximum allowable distance of 2 nucleotides between alleles within a locus (-

M option ustacks), a maximum of 2 mismatches allowed when aligning secondary reads

to primary stacks (-N option ustacks), and a maximum nucleotide distance of 2 allowed

between loci of different inviduals (-n option cstacks). A subset of 30 samples, 15 from

each site, was used to create the master catalog of loci. Using the STACKS 'populations'

program, loci containing any SNP categorized as heterozygous for an individual were

identified. Since all individuals sampled were haploid gametophytes, all loci were

expected to be homozygous; therefore, those loci identified as heterozygous were

discarded as they likely represented paralogous or over-merged loci. Sequence data were

further filtered to retain only bi-allelic loci with 1 or fewer SNPs, found in a minimum of

74% of samples, with a minor allele frequency lower limit of 0.1, resulting in 2,234 loci

used for subsequent genetic analyses.

Twenty-one *S. caninervis* ramets of known sex (11 females expressing archegonia

and 10 males expressing antheridia) were used to identify potential sex-associated loci

(fragments) that would allow us to infer sex of sterile ramets. Any 80 bp locus that was

present in at least five individuals of one sex and absent in all individuals of the opposite

sex was considered a potential sex-associated locus. Using these criteria we identified

over 1,000 candidate sex-associated loci. From this list, the 100 male-associated and 100

female-associated loci that were present at the highest frequency in each sex (present in 9

of the 10 males and all of the 11 females) were selected for further analyses. A custom

Perl script was used to search for these sequences in the alignments from the final 131

sterile ramets used in our analyses (see Results). Of the 200 candidate sex-associated loci,

we identified loci that were present in the 131 sterile ramet samples and were in complete

linkage with one another. We discarded any loci that were not in complete linkage,

because this observation suggested that they were not truly sex-linked, along with any

loci that were not recovered in the sterile ramet data set. We counted the number of

ramets with only male-associated loci and the number with only female-associated loci

and tested for deviations from a 1:1 sex ratio with a Chi-square test and a 2-tailed p-

value.

In order to determine the number of unique genotypes among the 131 sterile

ramets, clonal assignments were made using genetic distance under the infinite allele

model with a pairwise distance threshold of 0 between individuals. Higher mismatch

thresholds, up to 8, were evaluated but all resulted in the same clonal assignments. Figure

2 shows the clonal decay distribution of number of unique genets identified at different

genetic distance threshold levels. Furthermore, all ramets that were identified as clones of

the same genet were also genetically assigned to the same sex. Clonal assignments were

used to fill in missing data, when possible, by finding a consensus sequence among

ramets per MLG using GENEIOUS version 8.1.4 (Kearse et al., 2012). To test whether

the female biased sex ratio was generated by greater female growth rates, we compared

the mean number of sites occupied by female and male MLGs, testing for significance

using a one-tailed t-test.

To compare levels of genetic variation between the two study populations, and

between males and females in the low stress population, we calculated five measures of

clonal diversity using the software GENODIVE version 2.0b23 (Meirmans and Van

Tienderen, 2004). Clonal diversity (P_d) is a measure of unique genets or multi-locus

genotypes (MLGs) relative to (divided by) the number of ramets sampled (Ellstrand and

Roose, 1987). The effective number of genotypes (N_{eff}) is an index that accounts for

frequencies of each genet and describes the number of MLGs that have equal frequencies

while minimizing low frequency MLGs. This index is analogous to effective number of

alleles but instead counts whole MLGs. Clonal evenness (effective number of genotypes

divided by number of genotypes) indicates how evenly the MLGs are divided over the

population and would be equal to 1.00 if all MLGs were represented equally. We also

calculated Simpson's diversity index, corrected for sample size, and the corrected

Shannon's index (Chao and Shen, 2003). The latter is a measure of clonal variation that

accounts for singletons (genets or MLGs sampled just once) in the population for sample

sizes greater than approximately 50 sampling units (Arnaud-Haond et al., 2007). We used

bootstrap tests with 1,000 permutations and subsampling to match population sizes to test

for differences in the latter three measures of clonal diversity.

To evaluate the degree to which differences between the two populations could be

due to fixed genetic differences, we first calculated the population differentiation statistic

F_{ST} among the 131 sterile ramets from the high and low stress sites using GENODIVE.

Next we performed a principal components analysis (PCA) of genetic distance

covariation, with the missing data filled in first with clones, where possible, then with the

mean allele frequencies of all ramets. The eigenvectors (axis loadings) from the first two

PCA axes were plotted with the ggplot2 R package (R Core Team, 2013). We also

estimated the population structure using the fastSTRUCTURE (Raj et al., 2014) inference

algorithm with a simple logistic prior and $K = 1$ through $K = 4$. The dataset used for this

contained 2,234 SNPs from 131 sterile ramets from the high and low stress sites where

missing data was filled in with clones, when possible. Membership coefficients for $K = 2$

were plotted using DISTRUCT version 2.2 from the fastSTRUCTURE software package.

For another means to visualize the patterns of genetic distance among genets in

the two populations, we constructed a midpoint-rooted neighbor joining genetic distance

tree using CLEARCUT. The dataset used for this contained 2,234 SNPs from 51 genets

from the high and low stress sites where missing data were filled in with clones, when

possible. Two genets with greater than 80% missing data were excluded. Because the

RAD loci are not completely linked, this tree does not represent the genealogical

relationships among these individuals but rather genetic similarity.

RESULTS

Phenotypic sex ratios— Of the 49 patches collected in the low stress site, 31

contained no sex structures, 17 contained ramets expressing archegonia and were

classified as female, 1 contained ramets of both sexes, and no patches contained ramets

with only male gametangia. The resulting phenotypic sex ratio of 18 F:1 M differs

significantly from 1:1 meiotic expectations (Chi-squared test, two-tailed p-value <

0.0001).

Sequencing statistiscs—Approximately 150 million total raw sequencing reads

were generated for the 141 ramet samples sequenced resulting in roughly 1.06 million

reads per barcode. About 40% of the reads were discarded due to low quality, resulting in

88 million reads that passed initial quality filters. Ten ramets with fewer than 4.5

thousand reads were discarded, leaving 98 sterile ramets in the high evelavtion, low stress

site, 33 sterile ramets in the low elevation, high stress site, and 21 ramets of known sex

from the low stress site, ranging from 51 thousand to 2.6 million reads per ramet. The

data matrix of 2,234 SNPs from 131 sterile ramet samples was 80% complete when using

original reads but increased to 94% complete when missing data were filled in with data

from identical clonal MLGs. The data matrix of concatenated loci used for the RAxML

analysis was 83.3% complete with a mean depth of about 5.6 reads per locus.

Identifying sex linked loci and genetic sex ratio— Of the 100 candidate male-

associated loci, 63 co-segregated exclusively within 33 sterile ramet samples and were in

complete linkage with one another. These 63 loci had an average read depth of 6.3 and on

average 33.5 male-associated loci were found within each newly identified male ramet. In

parallel, 65 of the 100 candidate female-associated loci co-segregated exclusively in the

remaining 98 sterile ramets and were also in complete linkage. These loci had a mean

read depth of 5.7 and an average of 25.3 female-associated loci were detected in each

newly identified female ramet. Of the remaining loci, 52 that were found in the test set of

21 ramets of known sex but were absent in the experimental set of 131 sterile ramets.

These may have been lost due to subtle differences in the library size selection procedure,

or may have been missed due to stochastic sampling. The remaining 20 loci may have

been incorrectly categorized as sex-linked, based on the small sample size of the trial set,

but it is also possible that some females could have contained fertilized archegonia that

went unnoticed during selection of ramets for DNA extraction and, thus, male-associated

loci were detected in unobserved embryos (indeed, 13 putative male-associated markers

found in sterile females, while 7 putative female-associated markers found in sterile

males). Using these putative sex linked markers, we found 65 female ramets and 33 male

ramets in samples from the low stress site, equating to an approximate 2:1 female-to-male

ratio. This represents a significant deviation from both the 1:1 expectation (Chi-squared

test, two-tailed p-value = 0.0012) and from the observed phenotypic sex expression at the

patch level (Chi-squared test, two-tailed p-value < 0.0001). Genetic sex ratio on a clonal

lineage level in the low stress site was also approximately 2:1 (31 female clonal lineages

and 14 male clonal lineages). All 33 of the ramets from the high stress site were found to

be female.

__Distribution of genotypic variation__— We found 53 unique MLGs among the 131

sterile ramets, based on analysis of 2,234 biallelic SNPs. The low stress site contained 45

unique MLGs from 98 ramets and the high stress site had eight MLGs from 33 ramets. A

total of 53 MLGs were distinguishable at genetic distance thresholds of 0 through 7, after

which the number of MLGs began to drop incrementally without distinct breaks,

indicating robust clonal assignments based on the SNP data (Figure 1).

Most unique MLGs in the low stress site were restricted to the 3-5cm patch they

were collected from. Of the 45 unique geneotypes in the low stress site, three were found

within adjacent patches (approximately 2 m apart) and only one MLG extended through

three patches for a total of about 4 m (Figure 2). Female and male MLGs did not occupy

significantly different numbers of patches (Figure 3A) nor did they contain significantly

different numbers of ramets per MLG (Figure 3B).

Both the Simpson's diversity and the corrected Shannon indices were

significantly higher in the low stress site than in the high stress site (p-value = 0.001).

Evenness was also slightly greater in the low stress site, but not significantly so (Table 1).

The Simpson's diversity and evenness values were not significantly different between the

two sexes in the low stress site, but the corrected Shannon Index was significantly higher

for females than for males (Table 2, p-value = 0.001).

Population differentiation and structure— The F_{ST} for the high and low stress

sites was 0.102 when all ramets were included. When estimated with just unique genets,

however, F_{ST} dropped to 0.028. The neighbor joining tree recovered genets of the high

stress site nested within the low stress site (Figure 5). Population structure analysis in

fastSTRUCTURE estimated a minimum of one model component to explain structure in

the data. However, this estimate was not supported by the corresponding marginal

likelihoods, which increased with the K parameter with no apparent maximum.

Membership coefficients for K = 2, visualized on Figure 5 show some support for

structure among the two sites.

The first axis of the principle components of covariance analysis (PCA) explained

8.112% of the total variation in the 131 ramets and 6.051% was explained by the second

axis. A scatterplot of the first two PC axes (Figure 6) shows a tighter clustering of many

ramets from the low stress site and lower association of ramets from the high stress site,

mostly resolved along PC axis 1. Yet, a few ramets from the high stress site are

genetically similar, and therefore clustered near, many in the low stress site.

DISCUSSION

S. caninervis has long been a model for investigations regarding the evolution and

ecology of sex ratio variation in mosses (Stark et al., 1998; Stark et al., 2001; Stark et al.,

2005a; Stark et al., 2005b; Stark and McLetchie, 2006; Paasch et al., 2015). However, the

inferences regarding sex ratio variation in this species, as well as other bryophytes, has

been limited by the large number of sterile plants in most bryophyte populations. In

principle, this limitation is overcome by the use of sex linked molecular markers, which

provide a simple means of determining the sex of sterile plants. Identifying reliable sex-

linked markers is not trivial, though, requiring screening large numbers of polymorphic

loci in a large pedigree or mapping population. In addition, sex linkage ideally should be

confirmed using a physical map, or molecular evolutionary analyses, for example to test

for complete linkage disquilibrium (LD) among sex linked loci, as predicted based on the

observation that recombination does not occur on UV sex chromosomes (Bachtrog et al.,

2011). To date, within mosses this has only been accomplished in the model system

Ceratodon purpureus (McDaniel et al., 2007; McDaniel et al., 2013). A less rigorous

approach involves testing for an association between a molecular marker and sex in a

large panel of unrelated individuals (as has been done in three *Drepanocladus* species;

(Bisang et al., 2010; Bisang and Hedenäs, 2013; Bisang et al., 2015; Hedenäs et al.,

2016). Here, we have screened several thousand restriction-site associated sequenced

(RADseq) loci in a small number of individuals of known sex, but additionally retained

only those that exhibited complete LD among all loci in a larger sample. Beyond the

large numbers of polymorphisms, RADseq loci have the advantage of being defined by a

unique DNA sequence, which both allows us to be confident in the homology of our loci

(unlike gel band-length approaches), and ultimately provides a means of identifying sex-

linked genes in published transcriptomes (Gao et al., 2014; Wickett et al., 2014).

We use these putative sex-linked markers to show that the genetic sex ratios in

two Mojave Desert populations populations of *S. caninervis* are female biased (2 F:1 M

in the low stress site, and entirely female in the high stress site). Importantly, however,

the phenotypic sex ratios in this sample were far more biased (17 F:1 M in the low stress

site). While it is certainly possible that we have over-estimated the long-term female sex

expression rates, due to drought in study area in the years prior to collection (2013-2014),

collections from seasons with more typical winter weather patterns reported phenotypic

sex ratios of 7 F:1 M (Paasch et al., 2015), approximately three times more female biased

than the genetic sex ratio we found. These data indicate that males constitute a

disporportionately large fraction of the sterile plants, providing unequivocal support for

the shy male hypothesis.

The greater frequency of female ramets, however, indicates that factors beyond

thresholds for sex expression must also contribute to the population sex ratio variation in

S. caninervis. Experimental manipulations show that females regenerate more readily

from plant fragments than males do under both cool conditions and desiccation stress

(Stark et al., 2004; Stark et al., 2005b; Stark and McLetchie, 2006), and potentially may

exhibit greater clonal propagation. However, the fact that we observed no difference in

number of sites occupied between female and male MLGs (Figure 3A) indicates that the

genetic female bias in this species is unlikely to be due to faster female growth.

A female genotype bias could also be causd by elevated male mortality. The

hypothesis uniquely predicts a lower male haplotype diversity, relative to females, as we

found (corrected Shannon index, 1.596 in females, 1.208 in males, $p = 0.001$). Indeed,

when comparing the two populations, the high stress site is less clonally diverse

(Simpson's and corrected Shannon indices, $p = 0.001$; P_d, no significance test) and has a

a greater bias than the low stress site. Importantly, the relatively low F_{ST} indicates that

most surveyed polymorphisms are shared between the two populations, suggesting that

the difference between these two populations are unlikely to result from fixed genetic

differences between the two populations.

With our current data it is not possible to isolate when in the life cycle male and

female demography differ, nor whether it is locally adapted male genotypes that survive

or only those genotypes that by chance land in more permissive environmental

conditions. However, the association between stress and male mortality suggests that the

interaction between male physiology and extrinsic environmental factors are more likely

to govern population sex ratio than exclusively intrinsic factors like sex ratio distorters

(McDaniel et al., 2007; Norrell et al., 2014). Indeed, the available evidence suggests that

S. caninervis female biased sex ratios in Mojave Desert populations correlate with

precipitation and temperature (this study; Bowker et al., 2000). One potential cause of the

elevated male mortality in *S. caninervis* is differential resourse allocation to reproduction.

Although sexual reproduction is costly for both sexes, males experience a higher initial

cost in the production of antheridia, while fertilized females bear the cost of nurturing a

sporophyte (Rydgren et al., 2010). Although we observe sporophytes only rarely in *S.*

caninervis, the relatively high diversity evident in our sample, along with the limited

population structure and weak structure in the population genealogy all indicate that

sexual reproduction is relatively frequent. Thus, males may experience sexual selection to

produce more antheridia (McDaniel et al., submitted), although it may be opposed by

natural selection which favors investment in the maintenance of vegetative tissues.

Several bryophyte demographic models predict the eventual local extinction of

males, based on vegetative growth patterns similar to those found in *S. caninervis*

(McLetchie et al., 2001; Crowley et al., 2005; Rydgren et al., 2010), although these

models generally assume that migration is negligable. The lack of population structure

between our two study sites indicates that population sex ratios may be influenced by

migration of male spores from other populations in addition to local population

dynamics. The mixing of genotypes in the PCA and distance tree suggests that both sites

in Sheeps Creek Wash draw spores from the same metapopulation. Indeed, the overall

genetic structure of S. caninervis is likely to be governed by low rates of sexual

reproduction throughout the whole region – to which the low stress site contributes some

spores – and those spores that land where they can grow and reproduce end up

contributing to subsequent generations. Other sites, like the high stress site, end up being

genotype sinks. These inferences are consistent with phylogeographic studies in other

species that report limited population structure even at much larger spatial scales

(McDaniel and Shaw, 2005; Vanderpoorten et al., 2008; Korpelainen et al., 2012; Shaw

et al., 2014).

Conclusions—This study demonstrates that the highly female-biased sex ratios

observed in Mojave Desert *S. caninervis* are congruent with both the shy male hypothesis

(Mishler and Oliver, 1991; Stark et al., 2005a) and increased male mortality. These

results highlight the importance of environmental conditions and demographic history for

shaping sex ratios in this species, and may have important consequences for the

persistence of local populations in the presence of long-term shifts in climate.

Importantly, both mechanisms are grounded in the disproportionate pre-zygotic energetic

cost of sexual reproduction experienced by males relative to females (Mishler and Oliver,

1991; Stark et al., 2000; Stark et al., 2005a). Genomic approaches like those we have

used here are likely to shed light on sex ratio variation in other moss species where other

processes may predominate (Cronberg, 2003; Bisang and Hedenäs, 2005; Stark et al.,

2010; Horsley et al., 2011; Bisang et al., 2014; Norrell et al., 2014). A future challenge is

to determine the contributions of migration and environmental factors like water

availability, through its effects on determining sex expression and mortality, on the

trajectory of population sex ratio change through time.

LITERATURE CITED

ARNAUD-HAOND S., C. M. DUARTE, F. ALBERTO, AND E. A. SERRÃO. 2007.
Standardizing methods to address clonality in population studies. *Molecular Ecology* 16:
5115-5139.

BACHTROG D., M. KIRKPATRICK, J. E. MANK, S. F. MCDANIEL, J. C. PIRES, W. R.
RICE, AND N. VALENZUELA. 2011. Are all sex chromosomes created equal? *Trends in*
Genetics 27: 350-357.

BISANG I., J. EARLEN, C. PERSSON, AND L. HEDENÄS. 2014. Family affiliation, sex ratio
and sporophyte frequency in unisexual mosses. *Botanical Journal of the Linnaean Society*
174: 163-172.

BISANG I., J. EHRLEN, H. KORPELAINEN, AND L. HEDENÄS. 2015. No evidence of sexual
niche partitioning in a dioecious moss with rare sexual reproduction. *Annals of Botany* 116:
771-779.

BISANG I. AND L. HEDENÄS. 2005. Sex ratio patterns in dioicous bryophytes re-visited.
Journal of Bryology 27: 207-219.

BISANG I. AND L. HEDENÄS. 2013. Males are not shy in the wetland moss *Drepanocladus*
lycopodioides. *International Journal of Plant Sciences* 174: 733-739.

BISANG I., H. KORPELAINEN, AND L. HEDENÄS. 2010. Can the sex- specific molecular
marker of Drepanocladus trifarius uncover gender in related species? *Journal of Bryology*
32: 305-308.

BOWKER M. A., L. R. STARK, D. N. MCLETCHIE, AND B. D. MISHLER. 2000. Sex
expression, skewed sex ratios, and microhabitat distribution in the dioecious desert moss
Syntrichia caninervis (Pottiaceae). *American Journal of Botany* 87: 517-526.

CATCHEN J., A. AMORES, P. HOHENLOHE, W. CRESKO, AND J. POSTLETHWAIT.
2011. Stacks: building and genotyping loci de novo from short-read sequences. *G3: Genes,*
Genomes, Genetics 1: 171-182.

CATCHEN J., P. HOHENLOHE, S. BASSHAM, A. AMORES, AND W. CRESKO. 2013.
Stacks: an analysis tool set for population genomics. *Molecular Ecology* 22: 3124-3140.

CHAO A. AND T. SHEN. 2003. Nonparametric estimation of Shannon's index of diversity when
there are unseen species in sample. *Environmental and Ecological Statistics* 10: 429-443.

CRONBERG N. 2003. Clonal distribution, fertility and sex ratios of the moss *Plagiomnium affine*
(Bland.) T. Kop. in forests of contrasting age. *Journal of Bryology* 25: 155-162.

CROWLEY P. H., C. STIEHA, AND D. N. MCLETCHIE. 2005. Overgrowth competition,
fragmentation and sex ratio dynamics: a spatially explicit, sub-individual-based model.
Journal of Theoretical Biology 233: 25-42.

ELLSTRAND N. C. AND M. L. ROOSE. 1987. Patterns of genotypic diversity in clonal plant-
species. *American Journal of Botany* 74: 123-131.

GAO B., D. ZHANG, X. LI, H. YANG, AND A. J. WOOD. 2014. De novo assembly and
characterization of the transcriptome in the desiccation-tolerant moss *Syntrichia caninervis*.
BMC Research Notes 7: 1-12.

GORDON A. AND G. J. HANNON. 2010. Fastx-toolkit.
http://hannonlab.cshl.edu/fastx_toolkit/index.

HEDENÄS L., I. BISANG, H. KORPELAINEN, AND B. CRONHOLM. 2010. The true sex
ratio in European *Pseudocalliergon trifarium* (Bryophyta: Amblystegiaceae) revealed my a
novel molecular approach. *Biological Journal of the Linnean Society* 100: 132-140.

HEDENÄS L., H. KORPELAINEN, AND I. BISANG. 2016. Identifying sex in non-fertile
individuals of the moss *Drepanocladus turgescens* (Bryophyta: Amblystegiaceae) using a
novel molecular approach. *Journal of Plant Research* 129: 1005-1010.

HORSLEY K., L. R. STARK, AND D. N. MCLETCHIE. 2011. Does the silver moss Bryum
argenteum exhibit sex-specific patterns in vegetative growth rate, asexual fitness or
prezygotic reproductive investment? *Annals of Botany* 107: 897-907.

KEARSE M., R. MOIR, A. WILSON, S. STONES-HAVAS, M. CHEUNG, S. STURROCK, S.
BUXTON, A. COOPER, S. MARKOWITZ, C. DURAN, T. THIERER, B. ASHTON, P.
MENTJIES, AND A. DRUMMOND. 2012. Geneious Basic: an integrated and extendable
desktop software platform for the organization and analysis of sequence data. *Bioinformatics*
28: 1647-1649.

KORPELAINEN H., I. BISANG, L. HEDENÄS, AND J. KOLEHMAINEN. 2008. The first sex-
specific molecular marker discovered in the moss Pseudocalliergon trifarium. *Journal of*
Heredity 99: 581-587.

KORPELAINEN H., A. K. JÄGERBRAND, AND M. VON CRÄUTLEIN. 2012. Genetic
structure of mosses *Pleurozium schreberi* (Willd. ex Brid.) Mitt. and *Racomitrium*
lanuginosum (Hedw.) Brid. along altitude gradients in Hokkaido, Japan. *Journal of Bryology*
34: 309-312.

MCDANIEL S. F., J. ATWOOD, AND J. G. BURLEIGH. 2013. Recurring evolution of dioecy
in bryophytes. *Evolution* 67: 567-572.

MCDANIEL S. F., S. B. CAREY, C. HENRIQUEZ, A. P. POWELL, G. T. GENNIS, E. Y.
WALKER, K. S. JONES, AND A. C. PAYTON. Sexual selection on male maturation
driven by mating season precipitation. *Evolution*.

MCDANIEL S. F. AND A. J. SHAW. 2005. Selective sweeps and intercontinental migration in
the cosmopolitan moss *Ceratodon purpureus* (Hedw.) Brid. *Molecular Ecology* 14: 1121-
1132.

MCDANIEL S. F., J. H. WILLIS, AND A. J. SHAW. 2007. A linkage map reveals a complex
basis for segregation distortion in an interpopulational cross in the moss *Ceratodon*
purpureus. *Genetics* 176: 2489-2500.

MCDANIEL S. F., J. H. WILLIS, AND A. J. SHAW. 2008. The genetic basis of developmental
abnormalities in interpopulation hybrids of the moss *Ceratodon purpureus*. *Genetics Society*
of America 1425-1435.

MCLETCHIE D. N., G. GARCÍA-RAMOS, AND P. H. CROWLEY. 2001. Local sex-ratio
dynamics: a model for the dioecious liverwort *Marchantia inflexa*. *Evolutionary Ecology* 15:
231-254.

MEIRMANS P. G. AND P. H. VAN TIENDEREN. 2004. GENOTYPE and GENODIVE: two
programs for the analysis of genetic diversity of asexual organisms. *Molecular Ecology*
Notes 4: 792-794.

MISHLER B. D. AND M. J. OLIVER. 1991. Gametophytic phenology of *Tortula ruralis*, a
desiccation-tolerant moss, in the Organ Mountains of southern New Mexico. *The Bryologist*
94: 143-153.

NORRELL T. E., K. S. JONES, A. C. PAYTON, AND S. F. MCDANIEL. 2014. Meiotic sex
ratio variation in natural populations of *Ceratodon purpureus* (Ditrichaceae). *American*
Journal of Botany 101: 1572-1576.

PAASCH A. E., B. D. MISHLER, S. NOSRATINIA, L. R. STARK, AND K. M. FISHER. 2015.
Decoupling of sexual reproduction and genetic diversity in the female-biased Mojave Desert
moss *Syntrichia caninervis* (Pottiaceae). *International Journal of Plant Sciences*.

PETERSON B. K., J. N. WEBER, E. H. KAY, H. S. FISHER, AND H. E. HOEKSTRA. 2012.
Double digest RADseq: An inexpensive method for *de novo* SNP discovery and genotyping
in model and non-model species. *PLOS One* 7.

R CORE TEAM. 2013. R: A language and environment for statistical computing. Vienna,
Austria.

RAJ A., M. STEPHENS, AND J. K. PRITCHARD. 2014. fastSTRUCTURE: Variational
Inference of Population Structure in Large SNP Data Sets. *Genetics* 197: 573-589.

RYDGREN K., R. HALVORSEN, AND N. CRONBERG. 2010. Infrequent sporophyte
production maintains a female-biased sex ratio in the unisexual clonal moss *Hylocomium*
splendens. *Journal of Ecology* 98: 1224-1231.

SHAW A. J. AND S. C. BEER. 1999. Life history variation in gametophyte populations of the
moss *Ceratodon purpureus* (Ditrichaceae). *American Journal of Botany* 86: 512-521.

SHAW A. J., G. K. GOLINSKI, E. G. CLARK, B. SHAW, H. K. STENØIEN, AND K. I.
FLATBERG. 2014. Intercontinental genetic structure in the amphi-Pacific peatmoss
Sphagnum miyabeanum (Bryophyta: Sphagnaceae). *Biological Journal of the Linnean*
Society 111: 17-37.

STARK L. R. 1997. Phenology and reproductive biology of *Syntrichia inermis* (Bryopsida,
Pottiaceae) in the Mojave Desert. *The Bryologist* 100: 13-27.

STARK L. R., D. N. MCLETCHIE, AND S. M. EPPLEY. 2010. Sex ratios and the shy male
hypothesis in the moss *Bryum argenteum* (Bryaceae). *The Bryologist* 113: 788-797.

STARK L. R., D. N. MCLETCHIE, AND B. D. MISHLER. 2005a. Sex expression, plant size,
and spatial segregation of the sexes across a stress gradient in the desert moss *Syntrichia*
caninervis. *American Bryological and Lichenological Society* 108: 183-193.

STARK L. R. AND D. N. MCLETCHIE. 2006. Gender-specific heat-shock tolerance of hydrated
leaves in the desert moss *Syntrichia caninervis*. *Physiologia Plantarum* 126: 187-195.

STARK L. R., N. MCLETCHIE, AND B. D. MISHLER. 2001. Sex expression and sex
dimorphism in sporophytic populations of the desert moss *Syntrichia caninervis*. *Plant*
Ecology 157: 183-196.

STARK L. R., B. D. MISHLER, AND D. N. MCLETCHIE. 1998. Sex expression and growth
rates in natural populations of the desert soil crustal moss *Syntrichia caninervis*. *Journal of*
Arid Environments 40: 401-416.

STARK L. R., B. D. MISHLER, AND D. N. MCLETCHIE. 2000. The cost of realized sexual
reproduction: assessing patterns of reproductive allocation and sporophyte abortion in a
desert moss. *American Journal of Botany* 87: 1599-1608.

STARK L. R., L. NICHOLS II, D. N. MCLETCHIE, S. D. SMITH, AND C. ZUNDEL. 2004.
Age and sex-specific rates of leaf regeneration in the Mojave Desert moss *Syntrichia*
caninervis. *American Journal of Botany* 91: 1-9.

STARK L. R., L. I. NICHOLS, D. N. MCLETCHIE, AND M. L. BONINE. 2005b. Do the sexes
of the desert moss *Syntrichia caninervis* differ in desiccation tolerance? A leaf regeneration
assay. *International Journal of Plant Sciences* 166: 21-29.

VANDERPOORTEN A., N. DEVOS, B. GOFFINET, O. J. HARDY, AND A. J. SHAW. 2008.
The barriers to oceanic island radiation in bryophytes: insights from the phylogeography of
the moss *Grimmia montana*. *Journal of Biogeography* 35: 654-663.

WICKETT N. J., S. MIRARAB, N. NGUYEN, T. WARNOW, E. CARPENTER, N. MATASCI,
S. AYYAMPALAYAM, M. S. BARKER, J. G. BURLEIGH, M. A. GITZENDANNER, B.
R. RUHFEL, E. WAFULA, J. P. DER, S. W. GRAHAM, S. MATHEWS, M.
MELKONIAN, D. E. SOLTIS, P. S. SOLTIS, N. W. MILES, C. J. ROTHFELS, L.
POKORNY, A. J. SHAW, L. DEGIRONIMO, D. W. STEVENSON, B. SUREK, J. C.
VILLARREAL, B. ROURE, H. PHILIPPE, C. W. DEPAMPHILIS, T. CHEN, M. K.
DEYHOLOS, R. S. BAUCOM, T. M. KUTCHAN, M. M. AUGUSTIN, J. WANG, Y.
ZHANG, Z. TIAN, Z. YAN, X. WU, X. SUN, G. K. WONG, AND J. LEEBENS-MACK.
2014. Phylotranscriptomic analysis of the origin and early diversification of land plants.
Proceedings of the National Academy of Sciences 111: E4859-E4868.

TABLES

Site	N	G	P_d	N_{eff}	Simpson's	Evenness	Shannon (corrected)
Low stress	98	45	0.459	36.379	0.983*	0.808	1.753*
High stress	33	8	0.242	6.368	0.869*	0.796	0.874*

600   Table 1. Clonal diversity indices for high and low stress Sheep Creek Wash sites.

601   Note: *Statistical difference between high and low, p-value = 0.001, based on 1,000

602   permutations with subsampling to match population sizes. N = number of ramets

603   sampled, G = number of genets, P_d = clonal diversity, N_{eff} = effective number of

604   genotypes.

Sex	N	G	P_d	N_{eff}	Simpson's	Evenness	Shannon (corrected)
Female	65	31	0.477	25.000	0.975	0.806	1.596*
Male	33	14	0.424	11.463	0.941	0.819	1.208*

Table 2. Clonal diversity indices for the sexes in low stress Sheep Creek Wash site.

Note: *Statistical difference between the sexes in low stress site, p-value = 0.001, based

on 1,000 permutations with subsampling to match population sizes. N = number of

ramets sampled, G = number of genets, P_d = clonal diversity, N_{eff} = effective number of

genotypes.

APPENDIX 1

Specimen vouchers were deposited in the California State University, Los Angeles

Herbarium (CSLA).

FIGURE LEGENDS

Figure 1. Clonal lineage decay at different genetic distance thresholds. This figure shows

the number of genets that would be identified at each genetic distance parameter setting.

Figure 2. Spatial distribution of genets. The low stress site contained 53 unique genets in

98 ramets from 33 patches along three transects (T1, T2, and T3). Females are

represented as circles and males as squares. Genets or multi-locus genotypes (MLGs) are

outlined in dashed lines. In cases that genets are in different patches, a dashed line

connects different ramets of the same MLG.

Figure 3. Number of patches occupied by (A) and (B) number of ramets per male and

female MLG. The average number of patches occupied by and the number of ramets

male and female MLGs in the Sheep Creek Wash low stress site do not differ

significantly. Error bars are standard deviation.

Figure 4. Proportion of genets with 1-5 ramets in the high and low stress sites. Proportion

of each vertical bar represents proportion of total genets or multi-locus genotypes

(MLGs) that were represented by each number of ramets.

Figure 5. fastSTRUCTURE membership coefficients for K = 2. Vertical bars represents

an individual ramets while colors represent genetic clusters detected (K = 2). Bars on the

left represent ramets from the high stress site (n = 33) and bars on the right represent

ramets from the low stress site (n = 98). Membership coefficients are plotted for each

ramet and colored corresponding to the proportion of its MLG that most closely aligns

each of the two clusters. The dataset used for this contained 2,234 SNPs from 131 sterile

ramets from the high and low stress sites where missing data was filled in with clones,

when possible.

Figure 6. CLEARCUT neighbor joining genetic distance tree of ramets from the low and

the high stress site of Sheep Creek. Branches that lead to the low stress ('L') site are blue

and those that lead to the high stress site ('H') are red. Branches that lead to mixed-

population groups are purple. Symbols indicate inferred sex.

Figure 7. Genetic distance PCA of all MLGs in the high and low stress sites Sheep Creek

Wash. Ramets from the high stress site are red and from the low stress site are blue.

Males are indicated with triangles and females with circles, and shape size is scaled to the

number of ramets sharing the same MLG ('Clones'). Missing alleles were filled in first

with clones, when possible, then with the mean allele frequencies of all ramets. PC1

explains 8.112% of the total variation in the samples and 6.051% is explained by PC2

(Cumulative 14.164%).

Figure 3

High stress   Low stress

0.08

Figure 7