Multiple factors influence population sex ratios in the Mojave Desert moss *Syntrichia*

caninervis[1]

Jenna T. Baughman[2.4], Adam C. Payton[3], Amber E. Paasch[2], Kirsten M. Fisher[2], and

Stuart F. McDaniel[3]

[2]Department of Biological Sciences, California State University, 5151 State University

Drive, Los Angeles, CA 90032, USA, [3]Department of Biology, University of Florida, 876

Newell Drive, Gainesville, FL 32611, USA

[4]Author for correspondence: Department of Integrative Biology and University and

Jepson Herbaria, University of California, 3040 Valley Life Sciences Building, Berkeley,

CA 94720, USA; e-mail: jbaughman@berkeley.edu; phone: 219-393-0179.

Female biased *Syntrichia caninervis* genotypic sex ratio

[1]Manuscript received _________; revision accepted _________.

Acknowledgements

This work was supported by the California State University Department of

Biological Sciences Faculty-Student Achievement Fund to KMF, a Research Exchange

Grant to JTB from the NSF EDEN RCN (IOS 0955517), the NSF Louis Stokes Alliances

for Minority Participation (HRD-1363399) to AEP, the NIH-NIGMS Minority

Biomedical Research Support-Research Initiative for Scientific Enhancement program

(GM61331) to JTB, and an NSF grant (DEB 1541005) to SFM.

ABSTRACT

• *Premise of research:* Natural populations of many mosses appear highly female-

biased based on the presence of reproductive structures. This bias could be caused

by increased male mortality, lower male growth rate, or a higher threshold for

achieving sexual maturity in males. Here we test these hypotheses using samples

from two populations of the Mojave Desert moss *Syntrichia caninervis*.

• *Methods:* We used double digest restriction-site associated DNA (RAD)

sequencing to identify candidate sex-associated loci in a panel of sex-expressing

plants. Next, we used putative sex-associated markers to identify the sex of

individuals without sex structures.

• *Key results:* We found an 18:1 phenotypic female: male sex ratio in the higher

elevation site (Wrightwood), and no sex expression at the low elevation site

(Phelan). In contrast, based on genetic data we found a 2:1 female bias in the

Wrightwood site and only females in the Phelan site. The area occupied by male

and female genets was indistinguishable.

• *Conclusions:* These data suggest that both differential mortality and sexual

dimorphism in thresholds for sex expression likely contribute to population

genetic and phenotypic sex ratio biases, and that phenotypic sex expression alone

fails to capture the extent of actual sex ratio bias present in natural populations of

S. caninervis.

INTRODUCTION

The ratio of males to females in sexually reproducing, dioecious species is generally predicted to be close to 1:1, given the assumption of equal parental investment in offspring of both sexes (Fisher, 1930). Analyses of consistent sex ratio biases therefore have provided important insights into the genetic basis of sex determination or the importance of variation in the roles of males and females in the life cycle (Charnov, 1982; West, 2009). Sex allocation theory predicts that skewed sex ratios can evolve as a result of both competitive and facilitative interactions among relatives (Hamilton, 1967), such that competition for resources among related individuals can bias sex ratios toward the more dispersive sex (Clark, 1978; Gowaty, 1993; Hewison and Gaillard, 1996), while mating among siblings can bias the sex ratio toward females (Herre, 1985; West and Herre, 1998; Reece et al., 2008). The fact that the cytoplasm is generally maternally inherited means that male-killing cyto-nuclear interactions often generate female biases (Schnable and Wise, 1998; Hurst et al., 1999). Other forms of sex-biased mortality also may cause sex ratio biases in specific demographic classes. Thus, generating a clear understanding of consistent sex ratio biases can provide insights into the natural history of a group of organisms, or the factors that govern genetic transmission from one generation to the next. Moreover, because biased sex ratios can profoundly influence the patterns of genetic variation within populations, such biases can have major implications for developing conservation strategies and estimating long term population persistence.

Female biased population sex ratios are a common yet unexplained phenomenon in bryophyte populations (Bisang and Hedenäs, 2005). In dioicous bryophytes, sex is determined at meiosis by a UV chromosomal system (Bachtrog et al., 2011); a spore

carrying the female sex-determining locus (U) will form a gametophyte that produces

archegonia and ultimately bears the offspring embryo (a sporophyte), while spores with

the male sex-determining locus (V) form gametophytes that produce antheridia. Because

the diploid sporophyte is produced by the union of a U-bearing egg with a V-bearing

sperm, the sporophyte is always heterozygous at the sex-determining chromosome (i.e.,

UV). As a consequence of meiotic segregation, therefore, the null expectation is a 1:1 sex

ratio. Understanding if the persistently female-biased phenotypic sex ratios in moss

populations represent a deviation from this genetic expectation is an essential first step

toward understanding the ultimate evolutionary causes (e.g., local resource competition,

genetic conflict) and consequences of sex ratio bias in this ecologically important group

of plants.

Most of the female biases documented in mosses to date have been based on

counting the number of sexually mature male and female ramets, or branches, in a

population (but see McLetchie et al., 2001; Korpelainen et al., 2008; Bisang et al., 2010,

2014, 2015, Hedenäs et al., 2010, 2016; Bisang and Hedenäs, 2013). A female bias in the

production of sexually mature ramets in a natural population could be caused by at least

three distinct processes. First, an apparently female biased population sex ratio could

simply be a product of faster female growth, as has been found in several species (Shaw

and Beer, 1999; Stark, Nichols, et al., 2005; McDaniel et al., 2008). In this case, a

population would contain more female ramets than male ramets, and individual female

genets (i.e., unique haplotypes) would on average would occupy larger areas than male

genets, although the genetic diversity within males and females would be similar (but see

Bengtsson and Cronberg, 2009). Second, males might reach sexual maturity less

frequently than females. In other words, fewer male haplotypes might produce

gametangia than female haplotypes. If this were true, males would constitute a

disproportionately large fraction of the sterile plants in a population (termed 'the shy male

hypothesis'; Mishler and Oliver, 1991; Stark et al., 2005). In this case we would expect a

genotypic sex ratio that is closer to 1:1 than the sex ratio observed from counting fertile

males and females, although again the genetic diversity within males and females would

be similar. Finally, a female bias could be caused by elevated male mortality during spore

production (McDaniel et al., 2007; Norrell et al., 2014), establishment, or at some later

point in the life-cycle. This hypothesis predicts a genotypic female bias at both the ramet

and genet level. Regardless of the proximate cause, elevated male mortality would

decrease the amount of genetic diversity in males relative to females.

One of the most extreme cases of phenotypic sex ratio variation in mosses is in

Mojave Desert populations of *Syntrichia caninervis*. Previous studies based on counts of

sexually mature plants indicate that female ramets of *S. caninervis* outnumber males by

as much as 7:1 (Paasch et al., 2015) or 14:1 (Bowker et al., 2000; Bisang and Hedenäs,

2005), and that some populations lack sexually mature males entirely (Stark et al., 2001).

Mojave Desert *S. caninervis* is extremely desiccation tolerant and spends much of its life

in an air-dried state, limiting all biological functions to infrequent post-rainfall periods,

primarily in cool winter months (Stark, 1997; Stark et al., 1998). Differences in the

timing and duration of this biologically active period, such as precipitation differences

along elevation gradients, appear to affect frequency of sex expression (phenotypic sex)

in a population. A survey of 890 *S. caninervis* individuals from a 10 hectare elevation

gradient (Bowker et al., 2000) found that total percentage of sexually mature individuals

increased with elevation. Male sex expression occurred only at the higher elevations,

while lower elevation populations contained only a few females with archegonia. In

parallel with low levels of sex expression, sexual fertilization and sporophyte production

is relatively rare. Established desert *S. caninervis* populations apparently persist through

vegetative propagation (Paasch et al., 2015).

Here we expand upon this work using restriction site-associated DNA genome

sequencing (ddRADseq) to identify the sex of sterile ramets and study the patterns of

genetic variation in males and females in two Mojave Desert populations of *S. caninervis*.

We used these data to address four main questions. First, to test whether the phenotypic

female bias was caused by higher rates of gametangial production by females, we

estimated the phenotypic and genotypic sex ratio in two populations. Second, we

estimated genet size to test whether sexual dimorphism in growth rate was sufficient to

explain the female biased phenotypic sex ratios in this species. Third, to evaluate whether

a genetic sex ratio bias was explained by elevated male mortality, we calculated levels of

genetic diversity in male and female haplotypes in both populations. Finally, to evaluate

the potential for inbreeding or sex-specific dispersal patterns to generate conditions

favorable to local mate competition or local resource competition, we generated several

estimates of population differentiation. Collectively these data point to differential sex

expression and elevated male mortality as key causes of female-biased sex ratios in

Mojave Desert populations of *S. caninervis*.

MATERIALS AND METHODS

Sample Collection— In the Mojave Desert, *S. caninervis* grows in a semi-

continuous carpet in both shaded and exposed microsites. We identified collections in the

field by the characteristic leaf morphology, color, and hair points, although all

identifications were confirmed with leaf cross-sections in the laboratory (Mishler, 2007).

We collected male, female, and sterile *S. caninervis* samples at two sites from Sheep

Creek Wash. The first site ('Wrightwood') is located at an elevation of 1800 m in Sheep

Creek Wash near Wrightwood, CA (34° 22' 33.85" N, 117° 36' 34.59" W), at the west

edge of the Mojave Desert and the northern base of the San Gabriel Mountains. The

average high and low annual temperature is 16.8 °C and 1.7 °C, with an average annual

precipitation of 49.4 cm (2007-2011, Wrightwood Weather Station, NOAA National

Climatic Data Center). This site experiences little foot traffic. The second site in the

lower portion of Sheep Creek Wash ('Phelan') is at an elevation of 1257 m near Phelan,

CA (34° 25' 29.80" N, 117° 36' 30.91" W), about nine miles northeast of the Wrightwood

site. The average high and low annual temperature is 27 °C and 10 °C, while average

annual precipitation is 28.2 cm (2005-2009, Phelan, CA, NOAA National Weather

Service). This site is also highly disturbed by foot traffic and erosion.

173   To establish a panel of plants of known sex to use to identify sex-linked molecular

174   markers, we isolated 11 *S. caninervis* female and 10 male ramets from the Wrightwood

175   site in March and April of 2013. In the laboratory under a dissecting microscope, patches

176   were hydrated and screened for presence of current or past antheridia and archegonia.

177   To estimate the phenotypic sex ratio of the Wrightwood population, in May 2014

we collected in 3-5 cm patches in three parallel 20 m linear north-south (N-S) transects,

10 m apart from one another, collecting one patch every 2 m from a variety of shaded and

exposed microhabitats. Collecting at regular intervals along a transect was important for

calculating spatial extent of haplotypes and sexes, while collection from different habitat

types increased our chances of finding both sexes. In June 2014 we made collections

from the Phelan site in a similar manner in two parallel 20 m N-S transects. Additionally,

due to the highly irregular distribution of the species in the Phelan site, a third mini-

transect was sampled (beginning at about the 23 m mark of transect 2 and extending

approximately 2 m), more densely (ca. every 0.3 m) selecting samples in a northwest-

southeast (NW-SE) direction from a variety of microhabitats. To estimate the genetic sex

ratio and perform diversity analyses, a maximum of three individual sterile ramets were

isolated from each patch using a dissecting microscope, resulting in a total of 99 ramets

from the Wrightwood site and, due to lower frequency of this species at this site, 42

ramets from the Phelan site, for a total of n = 141 individual ramets.

DNA Extraction and RADseq library preparation— Total genomic DNA was

extracted and isolated from 162 total ramets (21 of known sex, and 141 of unknown sex)

using a modified cetyl trimethyl ammonium bromide-beta mercaptoethanol (CTAB)

procedure (McDaniel et al., 2007). Prior to extraction, samples were ground dry to a fine

powder using a GenoGrinder 2010 bead shaker (SPEX CertiPrep, Metuchen, NJ). DNA

quality was evaluated for each sample by electrophoresis and quantified using a Qubit

Fluorometer (Invitrogen, Carlsbad, CA, USA), samples' DNAs were normalized prior to

library preparation.

Illumina libraries were prepared for sequencing using a modified version of the

Peterson et al. (2012) protocol using the endonucleases EcoRI and MseI (New England

Biolabs, Ipswich, MA). Following double enzyme digestion, unique barcoded adaptors

were ligated to the resulting EcoRI cut sites and a non-barcoded universal adaptor to the

MseI cut site. Variable length barcodes of 8, 9, and 10 base pairs (bp) were used with

each barcode, differing by at least 4 bp. Illumina flow cell binding and sequencing primer

sites were added to the adaptor ligated fragments throughout 10 cycles of PCR using

NEB Q5 PCR master mix (New England Biolabs, Ipswich, MA). Success of library

construction was evaluated through agarose gel electrophoresis, after which 5 µL of each

sample's final library was pooled into a single tube for the 141 sterile samples and a

separate tube for the 21 samples of known sex. Size selection and sequencing were

performed at the University of Florida's Interdisciplinary Center for Biotechnology

Research. Pooled libraries were fractioned using Pippin ELF precision electrophoresis

(Sage Science, Beverly, MA) with the resulting 250-500 bp fraction being used for

Illumina sequencing. The sterile ramet sample library was sequenced using an Illumina

NextSeq 500 at mid throughput, producing 150 bp single-end reads. The library

consisting of the 21 known-sex ramets was pooled with libraries constructed in an

identical manner for a non-related study and sequenced on one lane of a HiSeq 3000

producing 100 bp single end reads.

Data analysis— Raw ddRAD sequence reads were assessed for quality with

fastQC (Gordon and Hannon, 2010). All manipulation of raw sequence reads was

performed using tools from the FASTX-Toolkit (Gordon and Hannon, 2010)

implemented in Galaxy on the University of Florida's Research Computing Cluster. To

eliminate low quality bases at the 3' end of reads, all sequences were trimmed to 100 bp.

Reads were filtered for a minimum Phred quality score of 20 on at least 90% of the read.

All reads were then demultiplexed followed by the removal of the 5' barcode and EcoRI

enzyme cut site.

RAD loci were assembled de novo using STACKS, version 1.24, (Catchen et al.,

2011, 2013). Pertinent parameters for the STACKS pipeline were as follows: a minimum

of 2 reads required for each allele within an individual (-m option ustacks), a maximum

allowable distance of 2 nucleotides between alleles within a locus (-M option ustacks), a

maximum of 2 mismatches allowed when aligning secondary reads to primary stacks (-N

option ustacks), and a maximum nucleotide distance of 2 allowed between loci of

different individuals (-n option cstacks). A subset of 30 samples, 15 from each site, was

used to create the master catalog of loci. Using the STACKS 'populations' program, loci

containing any SNP categorized as heterozygous for an individual were identified. Since

all individuals sampled were haploid gametophytes, all loci were expected to be

homozygous; therefore, those loci identified as heterozygous were discarded as they

likely represented paralogous or over-merged loci. Approximately 23% of loci had at

least one individual with a heterozygous SNP call. Although removing only the

heterozygous allele may be sufficient for removing errors, to be conservative we remove

the entire locus from the dataset. Sequence data were further filtered to retain only bi-

allelic loci with 1 or fewer SNPs, found in a minimum of 74% of samples, with a minor

allele frequency lower limit of 0.1, resulting in 2,234 loci used for subsequent genetic

analyses.

Twenty-one S. caninervis ramets of known sex (11 females expressing archegonia

and 10 males expressing antheridia) were used to identify potential sex-associated loci

(fragments) that would allow us to infer sex of sterile ramets. Any 80 bp locus that was

present in most individuals of one sex and absent in all individuals of the opposite sex

was considered a potential sex-associated locus. Using these criteria, we identified over

5,000 candidate sex-associated loci. Although one sex-linked locus would suffice for our

purposes, we selected the 100 male-associated and 100 female-associated loci with the

best coverage across all 21 individuals for further analyses. This is because a single

autosomal locus may show a spurious association with sex in a sample of that size, due to

chance or because it has a sex-biased allele frequency; the region of suppressed

recombination on the UV sex chromosomes, however, makes it likely that many sex-

linked markers will segregate in complete linkage disequilibrium, providing a signature

of true sex linkage. We used a custom Perl script to search for the candidate sex-linked

sequences in the alignments from the final 131 sterile ramets used in our analyses (see

Results). We then used the loci that were in complete linkage with one another in the 131

sterile ramet samples to determine the sex of each haplotype and tested for deviations

from a 1:1 sex ratio with a Chi-square test and a 2-tailed p-value.

In order to determine the number of unique haplotypes among the 131 sterile

ramets, clonal assignments were made using genetic distance under the infinite allele

model with a pairwise distance threshold of 0 between individuals. Higher mismatch

thresholds, up to 8, were evaluated but all resulted in the same clonal assignments. Figure

1 shows the clonal decay distribution of number of unique genets identified at different

genetic distance threshold levels. Furthermore, all ramets that were identified as clones of

the same genet were also genetically assigned to the same sex. Clonal assignments were

used to fill in missing data, when possible, by finding a consensus sequence among

ramets per haplotype using GENEIOUS version 8.1.4 (Kearse et al., 2012). To test

whether the female biased sex ratio was generated by greater female growth rates, we

compared the mean number of sites occupied by female and male haplotypes, as well as

the mean number of ramets per male and female haplotypes, testing for significance using

a one-tailed t-test.

To compare levels of genetic variation between the two study populations, and

between males and females in the Wrightwood population, we calculated five measures

of clonal diversity using the software GENODIVE version 2.0b23 (Meirmans and

Tienderen, 2004). Clonal diversity (P_d) is a measure of unique genets relative to (divided

by) the number of ramets sampled (Ellstrand and Roose, 1987). The effective number of

genotypes (N_{eff}) is an index that accounts for frequencies of each genet and describes the

number of haplotypes that have equal frequencies while minimizing low frequency

haplotypes. This index is analogous to effective number of alleles but instead counts

whole haplotypes. Clonal evenness (effective number of genotypes divided by number of

genotypes) indicates how evenly the haplotypes are divided over the population and

would be equal to 1.00 if all haplotypes were represented equally. We also calculated

Simpson's diversity index, corrected for sample size, and the corrected Shannon's index

(Chao and Shen, 2003). The latter is a measure of clonal variation that accounts for

singletons (genets or haplotypes sampled just once) in the population for sample sizes

greater than approximately 50 sampling units (Arnaud-Haond et al., 2007). We used

bootstrap tests with 1,000 permutations and subsampling to match population sizes to test

for differences in the latter three measures of clonal diversity.

To evaluate the degree to which differences between the two populations could be

due to allele frequency differences, we first calculated the population differentiation

statistic F_{ST} among the 131 sterile ramets from the Phelan and Wrightwood sites using

GENODIVE. Next, we estimated the population structure using the fastSTRUCTURE

(Raj et al., 2014) inference algorithm with a simple logistic prior and K=1 through K=4.

The dataset used for this contained 2,234 SNPs from 131 sterile ramets from the Phelan

and Wrightwood sites, where missing data were filled in with data from identical clonal

haplotypes, when possible. Membership coefficients for K=2 were plotted using

DISTRUCT version 2.2 from the fastSTRUCTURE software package.

For another means to visualize the patterns of genetic distance among genets in

the two populations, we constructed a midpoint-rooted neighbor joining genetic distance

tree using CLEARCUT. The dataset used for this contained 2,234 SNPs from 51 genets

from the Phelan and Wrightwood sites, where missing data were again filled in with

identical haplotypes, when possible. Two genets with greater than 80% missing data were

excluded. Because the RAD loci are not completely linked, this tree does not represent

the genealogical relationships among these individuals but rather genetic similarity.

RESULTS

Phenotypic sex ratios— Of the 49 patches collected in the Wrightwood site, 31

contained no sex structures, 17 contained ramets expressing archegonia and were

classified as female, one contained ramets of both sexes, and no patches contained ramets

with only male gametangia. The resulting phenotypic sex ratio of 18 F:1 M differs

significantly from 1:1 meiotic expectations (Chi-squared test, two-tailed p-value <

0.0001).

Sequencing statistics—Approximately 150 million total raw sequencing reads

were generated for the 141 ramet samples sequenced resulting in roughly 1.06 million

reads per barcode. About 40% of the reads were discarded due to low quality, resulting in

88 million reads that passed initial quality filters. Ten ramets with fewer than 4.5

thousand reads were discarded, leaving 98 sterile ramets in the Wrightwood site, 33

sterile ramets in the Phelan site, and 21 ramets of known sex from the Wrightwood site,

ranging from 51 thousand to 2.6 million reads per ramet. The data matrix of 2,234 SNPs

from 131 sterile ramet samples was 80% complete when using original reads but

increased to 94% complete when missing data were filled in with data from identical

clonal haplotypes. The data matrix of concatenated loci used for the RAxML analysis

was 83.3% complete with a mean depth of about 5.6 reads per locus.

Identifying sex linked loci and genetic sex ratio— We used the 20 individuals of

known sex to identify 100 candidate male and 100 candidate female diagnostic markers.

We chose this approach because our sample size was necessarily small, meaning that any

single locus could show a spurious or population specific linkage to sex. Indeed,

approximately 25% of the loci from our trial set (52 loci between males and females)

were absent from our individuals of unknown sex, and 10% of the loci were apparently

misclassified as sex-linked because they failed to show complete linkage disequilibrium

with the other candidate loci. It is possible that the missing loci were lost due to subtle

differences in the library size selection procedure, or due to stochastic sampling during

the sequencing process. The remaining 65% of the candidate sex-linked loci were in

complete linkage with one another, as expected for a UV sex chromosome system. The

63 putatively male-linked loci had an average read depth of 6.3 and on average 33.5

male-associated loci were found within each newly identified genotypic male ramet. The

65 putatively female-linked loci had a mean read depth of 5.7 and an average of 25.3

female-associated loci were detected in each newly identified genotypic female ramet.

Thus, although some of these loci may later be shown to be only sex-correlated, or

limited to these populations, and not truly sex-linked, these data give us high confidence

that collectively these loci are diagnostic markers for sex in this sample of individuals.

Using these putative sex-linked markers, we found 65 female ramets and 33 male

ramets in samples from the Wrightwood site, equating to an approximate 2:1 female-

biased genetic sex ratio. This represents a significant deviation from both the 1:1

expectation (Chi-squared test, two-tailed p-value = 0.0012) and from the observed 18:1

phenotypic sex expression based on counts of sexually mature ramets (Chi-squared test,

two-tailed p-value < 0.0001). All 33 of the ramets from the Phelan site were female.

Including all of the 2,234 biallelic SNPs, we found 53 distinct genetic individuals,

or haplotypes, among the 131 sterile ramets. The Wrightwood site contained 45 unique

haplotypes and the Phelan site had eight haplotypes. A total of 53 haplotypes were

distinguishable at genetic distance thresholds of 0 through 7, after which the number of

haplotypes began to drop incrementally without distinct breaks, indicating that the

number of genotypes is unlikely to be inflated as a result of genotyping errors (Figure 1).

The plants in the Wrightwood site were assignable to 31 female haplotypes and 14 male

haplotypes, corresponding to an approximately 2:1 female-biased sex ratio.

Distribution of genotypic variation— To test whether males and females

exhibited different patterns of vegetative expansion, we counted the number of sites that

each haplotype occupied. Most haplotypes in the Wrightwood site were restricted to a

single 3-5 cm patch. Of the 45 haplotypes in the Wrightwood site, three were found

within adjacent patches (approximately 2 m apart) and only one haplotype spanned three

patches across about 4 m (Figure 2). Female and male haplotypes did not occupy

significantly different numbers of patches (mean female patches = 1.097 ± 0.301, mean

male patches = 1.143 ± 0.535, one-tailed t-test p-value = 0.3565) nor did they contain

significantly different numbers of ramets per haplotype (mean female ramets = 2.097 ±

1.044, mean male ramets = 2.357 ± 1.151, one-tailed t-test p-value = 0.2286).

To test whether the sexes harbored different amounts of genetic diversity, we

calculated several indices. The Simpson's diversity and evenness values were not

significantly different between the two sexes in Wrightwood, but the corrected Shannon

Index was significantly higher for females than for males (Table 1, p-value = 0.001).

Both the Simpson's diversity and the corrected Shannon indices were significantly higher

in Wrightwood than in Phelan (p-value = 0.001). Evenness was also slightly greater in

the Wrightwood site, but not significantly so (Table 2).

Population differentiation and structure— To evaluate the potential for sex

biased dispersal or inbreeding to drive sex ratio bias, we examined several measures of

population structure. The F_{ST} between the Phelan and Wrightwood sites was 0.102 when

all ramets were included. When estimated with just unique haplotypes, however, F_{ST}

dropped to 0.028, indicating that the patterns of haplotype expansion differ in the two

populations. The neighbor joining tree recovered haplotypes collected from Phelan nested

within Wrightwood haplotypes (Figure 3). Population structure analysis in

fastSTRUCTURE estimated a minimum of one model component to explain structure in

the data. However, this estimate was not supported by the corresponding marginal

likelihoods, which increased with the K parameter with no apparent maximum.

Membership coefficients for K =2, visualized on Figure 4 show some support for

structure among the two sites, consistent with inferences based on F_{ST}.

DISCUSSION

S. caninervis has long been a model for investigations regarding the evolution and

ecology of sex ratio variation in mosses (Stark et al., 1998, 2001; Stark, McLetchie, et al.,

2005; Stark, Nichols, et al., 2005; Stark and McLetchie, 2006; Paasch et al., 2015).

However, our understanding of sex ratio variation in this species, as well as other

bryophytes, has been limited by the large number of sterile plants in most bryophyte

populations. In principle, this limitation is overcome by the use of sex-linked molecular

markers, which provide a simple means of determining the sex of sterile plants.

Identifying reliable sex-linked markers is not trivial, though, requiring screening large

numbers of polymorphic loci in a large pedigree or mapping population. In addition, sex

linkage ideally should be confirmed using a physical map, or molecular evolutionary

analyses, for example to test for complete linkage disequilibrium (LD) among sex linked

loci, as predicted based on the observation that recombination does not occur on UV sex

chromosomes (Bachtrog et al., 2011). To date, within mosses this has only been

accomplished in the model system *Ceratodon purpureus* (McDaniel et al., 2007, 2013). A

less rigorous approach involves testing for an association between a molecular marker

and sex in a large panel of unrelated individuals (as has been done in three

Drepanocladus species; (Bisang et al., 2010, 2015; Bisang and Hedenäs, 2013; Hedenäs

et al., 2016). Here, we have used the latter approach, and screened several thousand

restriction-site associated sequenced (RADseq) loci in a small number of individuals of

known sex. Although the number of individuals was small, the large number of loci gave

us additional confidence because we could retain only those that exhibited complete LD

among all loci in a larger sample. Beyond the large numbers of polymorphisms, RADseq

loci have the advantage of being defined by a unique DNA sequence, which both allows

us to be confident in the homology of our loci (unlike gel band-length approaches), and

ultimately provides a means of identifying sex-linked genes in published transcriptomes

(Gao et al., 2014; Wickett et al., 2014).

We use these putative sex-linked markers to show that the genotypic sex ratios in

two Mojave Desert populations populations of *S. caninervis* are female biased (2 F:1 M

in the Wrightwood site, and entirely female in the Phelan site). Importantly, however, the

phenotypic sex ratios in this sample were far more biased (18 F:1 M in Wrightwood).

While it is certainly possible that we have over-estimated the long-term female sex

expression bias, due to drought in the study area in the years prior to collection (2013-

2014), collections from seasons with more typical winter weather patterns reported

phenotypic sex ratios of 7 F:1 M (Paasch et al., 2015), approximately three times more

female biased than the genotypic sex ratio we found. These data indicate that males

constitute a disproportionately large fraction of the sterile plants, providing unequivocal

support for the shy male hypothesis in these Mojave Desert populations of *S. caninervis*.

The greater frequency of female ramets, however, indicates that factors beyond

thresholds for sex expression must also contribute to the population sex ratio variation in

S. caninervis. Experimental manipulations show that females regenerate more readily

from plant fragments than males do under both cool conditions and desiccation stress

(Stark et al., 2004; Stark, Nichols, et al., 2005; Stark and McLetchie, 2006), and

potentially may accumulate more biomass. However, the fact that we observed no

difference in number of sites occupied between female and male haplotypes indicates that

the genotypic female bias in these populations is unlikely to be due to faster female

growth.

We do, however, find some evidence that elevated male mortality contributes to

the overall female biased sex ratio. The male-mortality hypothesis uniquely predicts a

lower male haplotype diversity, relative to females, as we found (corrected Shannon

index, 1.596 in females, 1.208 in males, $p = 0.001$). When comparing the two

populations, the Phelan site is less clonally diverse (Simpson's and corrected Shannon

indices, $p = 0.001$) and has a greater genotypic female bias than the Wrightwood site, also

suggesting greater mortality. Importantly, the relatively low F_{ST} between these two

populations indicates that most surveyed polymorphisms are shared between Phelan and

Wrightwood, suggesting that sex ratio differences between them are unlikely to result

from long isolation.

With our current data it is not possible to isolate when in the life cycle male and

female demography differ, nor whether the two sexes exhibit different microhabitat

preferences. However, the observed pattern of lower male diversity (i.e., elevated male

mortality) suggests that the interaction between male physiology and extrinsic

environmental factors are more likely to govern population sex ratio than exclusively

intrinsic factors like sex ratio distorters (McDaniel et al., 2007; Norrell et al., 2014).

Indeed, the available evidence suggests that *S. caninervis* female biased phenotypic sex

ratios in Mojave Desert populations correlate with precipitation and temperature (Bowker

et al., 2000). One potential cause of the rarity of males observed in *S. caninervis* is the

difference in timing of resource allocation to reproduction between the sexes. Although

sexual reproduction is costly for both sexes, in the moss *Hylocomium splendens* males

make a higher initial investment in the production of antheridia, while fertilized females

only allocate energy to reproduction after fertilization through nurturing a sporophyte

(Rydgren et al., 2010). Although we observe sporophytes only rarely in *S. caninervis*, the

relatively high diversity evident in our sample, along with the limited population

structure and weak structure in the population genealogy all indicate that sexual

reproduction is or has been present in these populations. Thus, males may experience a

trade-off between sexual selection, to produce more reproductive tissue, and natural

selection, which may favor investment in the maintenance of vegetative tissues, as has

been seen in the moss *C. purpureus* (McDaniel, 2005).

Several bryophyte demographic models predict the eventual local extinction of

males, based on vegetative growth patterns similar to those found in *S. caninervis*

(McLetchie et al., 2001; Crowley et al., 2005; Rydgren et al., 2010), although these

models generally assume that migration is negligible. The lack of population structure

between our two study sites indicates that population sex ratios may be influenced by

migration of spores from other populations in addition to local population dynamics. The

mixing of genotypes in the distance tree suggests that both sites in Sheep Creek Wash

draw spores from the same metapopulation. Some populations, like the Wrightwood site,

both recruit and produce spores that contribute to subsequent generations, while other

sites, like Phelan, permit growth but no sexual reproduction and become genotype sinks.

Our sampling is insufficient to quantitatively evaluate the population structure of Mojave

Desert populations of *S. caninervis*, but these inferences are consistent with studies in

other mosses that report some population structure at the patch scale (Hutsemékers et al.,

2010; Leonardía et al., 2013; Rosengren et al., 2016) but limited structure at regional or

larger spatial scales (McDaniel and Shaw, 2005; Vanderpoorten et al., 2008; Korpelainen

et al., 2012; Shaw et al., 2014; Désamoré et al., 2016).

Conclusions—This study demonstrates that the highly female-biased sex ratios

observed in Mojave Desert *S. caninervis* are congruent with both the shy male hypothesis

(Mishler and Oliver, 1991; Stark, McLetchie, et al., 2005) and increased male mortality

(Stark et al., 2000). These results suggest the importance of interactions between

environmental conditions and demographic factors for shaping sex ratios in this species,

and may have important consequences for the persistence of local populations in the

presence of long-term shifts in climate. Importantly, both mechanisms are grounded in

the disproportionate pre-zygotic energetic allocation to sexual reproduction by males

relative to females (Mishler and Oliver, 1991; Stark et al., 2000; Stark, McLetchie, et al.,

2005), in essence representing a trade-off between natural and sexual selection. Our data

suggest that local mate competition and local resource competition are less likely to be

major drivers of sex ratio variation in Mojave Desert populations of *S. caninervis*, but we

should caution that studies of inbreeding and sex-biased dispersal are needed in

bryophytes. We suspect that the wider use of genomic approaches like those we have

used here is likely to uncover cases of sex ratio variation where other processes may

predominate (Cronberg, 2003; Bisang and Hedenäs, 2005; Stark et al., 2010; Horsley et

al., 2011; Bisang et al., 2014; Norrell et al., 2014). A future challenge is to determine the

contributions of migration and environmental factors like water availability, through its

effects on determining sex expression and mortality, on the trajectory of population sex

ratio change through time.

LITERATURE CITED

ARNAUD-HAOND, S., C.M. DUARTE, F. ALBERTO, and E.A. SERRÃO. 2007. Standardizing

methods to address clonality in population studies. *Molecular Ecology* 16: 5115–

5139.

BACHTROG, D., M. KIRKPATRICK, J.E. MANK, S.F. MCDANIEL, J.C. PIRES, W.R. RICE,

and N. VALENZUELA. 2011. Are all sex chromosomes created equal? *Trends in*

Genetics 27: 350–357.

BENGTSSON, B.O., and N. CRONBERG. 2009. The effective size of bryophyte populations.

Journal of Theoretical Biology 258: 121–126.

BISANG, I., J. EARLEN, C. PERSSON, and L. HEDENÄS. 2014. Family affiliation, sex ratio

and sporophyte frequency in unisexual mosses. *Botanical Journal of the Linnaean*

Society 174: 163–172.

BISANG, I., J. EHRLEN, H. KORPELAINEN, and L. HEDENÄS. 2015. No evidence of sexual

niche partitioning in a dioecious moss with rare sexual reproduction. *Annals of*

Botany 116: 771–779.

BISANG, I., and L. HEDENÄS. 2013. Males are not shy in the wetland moss Drepanocladus

lycopodioides. *International Journal of Plant Sciences* 174: 733–739.

BISANG, I., and L. HEDENÄS. 2005. Sex ratio patterns in dioicous bryophytes re-visited.

Journal of Bryology 27: 207–219.

BISANG, I., H. KORPELAINEN, and L. HEDENÄS. 2010. Can the sex- specific molecular

marker of Drepanocladus trifarius uncover gender in related species? *Journal of*

Bryology 32: 305–308.

BOWKER, M.A., L.R. STARK, D.N. MCLETCHIE, and B.D. MISHLER. 2000. Sex

expression, skewed sex ratios, and microhabitat distribution in the dioecious desert

moss Syntrichia caninervis (Pottiaceae). *American Journal of Botany* 87: 517–526.

CATCHEN, J., A. AMORES, P. HOHENLOHE, W. CRESKO, and J. POSTLETHWAIT. 2011.

Stacks: building and genotyping loci de novo from short-read sequences. *G3: Genes,*

Genomes, Genetics 1: 171–182.

CATCHEN, J., P. HOHENLOHE, S. BASSHAM, A. AMORES, and W. CRESKO. 2013. Stacks:

an analysis tool set for population genomics. *Molecular Ecology* 22: 3124–3140.

CHAO, A., and T.-J. SHEN. 2003. Nonparametric estimation of Shannon's index of

diversity when there are unseen species in sample. *Environmental and Ecological*

Statistics 10: 429–443.

CHARNOV, E.L. 1982. The theory of sex allocation. *Monographs in population biology*

18: 1–355.

CLARK, A.B. 1978. Sex ratio and local resource competition in a prosimian primate.

Science 201: 163–165. Available at:

http://science.sciencemag.org/content/201/4351/163.long.

CRONBERG, N. 2003. Clonal distribution, fertility and sex ratios of the moss Plagiomnium

affine (Bland.) T. Kop. in forests of contrasting age. *Journal of Bryology* 25: 155–

162.

CROWLEY, P.H., C. STIEHA, and D.N. MCLETCHIE. 2005. Overgrowth competition,

fragmentation and sex ratio dynamics: a spatially explicit, sub-individual-based

model. *Journal of Theoretical Biology* 233: 25–42.

DÉSAMORÉ, A., J. PATIÑO, P. MARDULYN, S.F. MCDANIEL, F. ZANATTA, B. LAENEN, and

A. VANDERPOORTEN. 2016. High migration rates shape the postglacial history of

amphi-Atlantic bryophytes. *Molecular Ecology* 25: 5568–5584.

ELLSTRAND, N.C., and M.L. ROOSE. 1987. Patterns of genotypic diversity in clonal plant-

species. *American Journal of Botany* 74: 123–131.

FISHER, R.F. 1930. THE GENETICAL THEORY OF NATURAL SELECTION.

GAO, B., D. ZHANG, X. LI, H. YANG, and A.J. WOOD. 2014. De novo assembly and

characterization of the transcriptome in the desiccation-tolerant moss Syntrichia

caninervis. *BMC Research Notes* 7: 1–12.

GORDON, A., and G.J. HANNON. 2010. Fastx-toolkit.

GOWATY, P.A. 1993. Differential dispersal, local resource competition, and sex ratio

variation in birds. *The American Naturalist* 141: 263–280. Available at:

http://www.jstor.org/stable/2462668%5Cnhttp://about.jstor.org/terms.

HAMILTON, W.D. 1967. Extraordinary sex ratios. A sex-ratio theory for sex linkage and

inbreeding has new implications in cytogenetics and entomology. *Science (New*

York, N.Y.) 156: 477–488.

HEDENÄS, L., I. BISANG, H. KORPELAINEN, and B. CRONHOLM. 2010. The true sex ratio in

European Pseudocalliergon trifarium (Bryophyta: Amblystegiaceae) revealed my a

novel molecular approach. *Biological Journal of the Linnean Society* 100: 132–140.

HEDENÄS, L., H. KORPELAINEN, and I. BISANG. 2016. Identifying sex in non-fertile

individuals of the moss Drepanocladus turgescens (Bryophyta: Amblystegiaceae)

using a novel molecular approach. *Journal of Plant Research* 129: 1005–1010.

HERRE, E.A. 1985. Sex Ratio Adjustment in Fig Wasps. *Source: Science, New Series*

228: 896–898. Available at:

http://www.jstor.org/stable/1694703%5Cnhttp://about.jstor.org/terms.

HEWISON, A.J.M., and J.M. GAILLARD. 1996. Birth-sex ratios and local resource

competition in roe deer, Capreolus capreolus. *Behav. Ecol.* 7: 461–464. Available at:

http://beheco.oxfordjournals.org/cgi/content/abstract/7/4/461.

HORSLEY, K., L.R. STARK, and D.N. MCLETCHIE. 2011. Does the silver moss Bryum

argenteum exhibit sex-specific patterns in vegetative growth rate, asexual fitness or

prezygotic reproductive investment? *Annals of Botany* 107: 897–907.

HURST, G.D.D., F.M. JIGGINS, J.H.G. VON DER SCHULENBURG, D. BERTRAND, S.A.

WEST, I.I. GORIACHEVA, I.A. ZAKHAROV, ET AL. 1999. Male-killing Wolbachia in

two species of insect. *Proceedings of the Royal Society B: Biological Sciences* 266:

735.

HUTSEMÉKERS, V., O.J. HARDY, P. MARDULYN, A.J. SHAW, and A. VANDERPOORTEN.

2010. Macroecological patterns of genetic structure and diversity in the aquatic moss

Platyhipnidium riparioides. *New Phytologist* 185: 852–864.

KEARSE, M., R. MOIR, A. WILSON, S. STONES-HAVAS, M. CHEUNG, S. STURROCK, S.

BUXTON, ET AL. 2012. Geneious Basic: an integrated and extendable desktop

software platform for the organization and analysis of sequence data. *Bioinformatics*

28: 1647–1649.

KORPELAINEN, H., I. BISANG, L. HEDENÄS, and J. KOLEHMAINEN. 2008. The first sex-

specific molecular marker discovered in the moss Pseudocalliergon trifarium.

Journal of Heredity 99: 581–587.

KORPELAINEN, H., A.K. JÄGERBRAND, and M. VON CRÄUTLEIN. 2012. Genetic structure

of mosses Pleurozium schreberi (Willd. ex Brid.) Mitt. and Racomitrium

lanuginosum (Hedw.) Brid. along altitude gradients in Hokkaido, Japan. *Journal of*

Bryology 34: 309–312.

LEONARDÍA, A.A.P., B.C. TAN, and P.P. KUMAR. 2013. Population genetic structure of

the tropical moss Acanthorrhynchium papillatum as measured with microsatellite

markers. *Plant Biology* 15: 384–394.

MCDANIEL, S.F. 2005. Genetic correlations do not constrain the evolution of sexual

dimporphism in the moss Ceratodon purpureus. *Evolution* 59: 2353–2361.

MCDANIEL, S.F., J. ATWOOD, and J.G. BURLEIGH. 2013. Recurring evolution of dioecy in

bryophytes. *Evolution* 67: 567–572.

MCDANIEL, S.F., and A.J. SHAW. 2005. Selective sweeps and intercontinental migration

in the cosmopolitan moss Ceratodon purpureus (Hedw.) Brid. *Molecular Ecology*

14: 1121–1132.

MCDANIEL, S.F., J.H. WILLIS, and A.J. SHAW. 2007. A linkage map reveals a complex

basis for segregation distortion in an interpopulational cross in the moss Ceratodon

purpureus. *Genetics* 176: 2489–2500.

MᶜDANIEL, S.F., J.H. WILLIS, and A.J. SHAW. 2008. The genetic basis of developmental

abnormalities in interpopulation hybrids of the moss Ceratodon purpureus. *Genetics*

*Society of America*1425–1435.

MᶜLETCHIE, D.N., G. GARCÍA-RAMOS, and P.H. CROWLEY. 2001. Local sex-ratio

dynamics: a model for the dioecious liverwort Marchantia inflexa. *Evolutionary*

Ecology 15: 231–254.

MEIRMANS, P.G., and P.H. VAN TIENDEREN. 2004. GENOTYPE and GENODIVE: two

programs for the analysis of genetic diversity of asexual organisms. *Molecular*

Ecology Notes 4: 792–794.

MISHLER, B.D. 2007. Syntrichia. *In* F. of N. A. E. Committee [ed.], Flora of North

America North of Mexico, Vol. 27, 618–627.

MISHLER, B.D., and M.J. OLIVER. 1991. Gametophytic phenology of Tortula ruralis, a

desiccation-tolerant moss, in the Organ Mountains of southern New Mexico. *The*

Bryologist 94: 143–153.

NORRELL, T.E., K.S. JONES, A.C. PAYTON, and S.F. MᶜDANIEL. 2014. Meiotic sex ratio

variation in natural populations of Ceratodon purpureus (Ditrichaceae). *American*

Journal of Botany 101: 1572–1576.

PAASCH, A.E., B.D. MISHLER, S. NOSRATINIA, L.R. STARK, and K.M. FISHER. 2015.

Decoupling of sexual reproduction and genetic diversity in the female-biased

Mojave Desert moss Syntrichia caninervis (Pottiaceae). *International Journal of*

Plant Sciences.

PETERSON, B.K., J.N. WEBER, E.H. KAY, H.S. FISHER, and H.E. HOEKSTRA. 2012. Double

digest RADseq: An inexpensive method for de novo SNP discovery and genotyping

in model and non-model species. *PLOS One* 7: .

RAJ, A., M. STEPHENS, and J.K. PRITCHARD. 2014. fastSTRUCTURE: Variational

Inference of Population Structure in Large SNP Data Sets. *Genetics* 197: 573–589.

REECE, S.E., D.R. DREW, and A. GARDNER. 2008. Sex ratio adjustment and kin

discrimination in malaria parasites. *Nature* 453: 609–614.

ROSENGREN, F., N. CRONBERG, and B. HANSSON. 2016. Balance between inbreeding and

outcrossing in a nannandrous species, the moss Homalothecium lutescens.

HEREDITY 116: 107–113.

RYDGREN, K., R. HALVORSEN, and N. CRONBERG. 2010. Infrequent sporophyte

production maintains a female-biased sex ratio in the unisexual clonal moss

Hylocomium splendens. *Journal of Ecology* 98: 1224–1231.

SCHNABLE, P.S., and R.P. WISE. 1998. The molecular basis of cytoplasmic male sterility

and fertility restoration. *Trends in Plant Science* 3: 175–180.

SHAW, A.J., and S.C. BEER. 1999. Life history variation in gametophyte populations of

the moss Ceratodon purpureus (Ditrichaceae). *American Journal of Botany* 86: 512–

521.

SHAW, A.J., G.K. GOLINSKI, E.G. CLARK, B. SHAW, H.K. STENØIEN, and K.I. FLATBERG.

2014. Intercontinental genetic structure in the amphi-Pacific peatmoss Sphagnum

miyabeanum (Bryophyta: Sphagnaceae). *Biological Journal of the Linnean Society*

111: 17–37.

STARK, L.R. 1997. Phenology and reproductive biology of Syntrichia inermis (Bryopsida,

Pottiaceae) in the Mojave Desert. *The Bryologist* 100: 13–27.

STARK, L.R., L.N. II, D.N. MCLETCHIE, S.D. SMITH, and C. ZUNDEL. 2004. Age and sex-

specific rates of leaf regeneration in the Mojave Desert moss Syntrichia caninervis.

American Journal of Botany 91: 1–9.

STARK, L.R., and D.N. MCLETCHIE. 2006. Gender-specific heat-shock tolerance of

hydrated leaves in the desert moss Syntrichia caninervis. *Physiologia Plantarum*

126: 187–195.

STARK, L.R., D.N. MCLETCHIE, and S.M. EPPLEY. 2010. Sex ratios and the shy male

hypothesis in the moss Bryum argenteum (Bryaceae). *The Bryologist* 113: 788–797.

STARK, L.R., D.N. MCLETCHIE, and B.D. MISHLER. 2005. Sex expression, plant size, and

spatial segregation of the sexes across a stress gradient in the desert moss Syntrichia

caninervis. *American Bryological and Lichenological Society* 108: 183–193.

STARK, L.R., N. MCLETCHIE, and B.D. MISHLER. 2001. Sex expression and sex

dimorphism in sporophytic populations of the desert moss Syntrichia caninervis.

Plant Ecology 157: 183–196.

STARK, L.R., B.D. MISHLER, and D.N. MCLETCHIE. 1998. Sex expression and growth

rates in natural populations of the desert soil crustal moss Syntrichia caninervis.

Journal of Arid Environments 40: 401–416.

STARK, L.R., B.D. MISHLER, and D.N. MCLETCHIE. 2000. The cost of realized sexual

reproduction: assessing patterns of reproductive allocation and sporophyte abortion

in a desert moss. *American Journal of Botany* 87: 1599–1608.

STARK, L.R., L.I.I. NICHOLS, D.N. MCLETCHIE, and M.L. BONINE. 2005. Do the sexes of

the desert moss Syntrichia caninervis differ in desiccation tolerance? A leaf

regeneration assay. *International Journal of Plant Sciences* 166: 21–29.

VANDERPOORTEN, A., N. DEVOS, B. GOFFINET, O.J. HARDY, and A.J. SHAW. 2008. The

barriers to oceanic island radiation in bryophytes: insights from the phylogeography

of the moss Grimmia montana. *Journal of Biogeography* 35: 654–663.

WEST, S.A. 2009. Sex Allocation. *Princeton University Press*. Available at:

http://books.google.com/books?id=HeCZBO8J9IYC&pgis=1.

WEST, S.A., and E.A. HERRE. 1998. Partial local mate competition and the sex ratio: A

study on non-pollinating fig wasps. *Journal of Evolutionary Biology* 11: 531–548.

WICKETT, N.J., S. MIRARAB, N. NGUYEN, T. WARNOW, E. CARPENTER, N. MATASCI, S.

AYYAMPALAYAM, ET AL. 2014. Phylotranscriptomic analysis of the origin and early

diversification of land plants. *Proceedings of the National Academy of Sciences* 111:

E4859–E4868.

698   TABLES

Sex	N	G	P_d	N_{eff}	Simpson's	Evenness	Shannon (corrected)
Female	65	31	0.477	25.000	0.975	0.806	1.596*
Male	33	14	0.424	11.463	0.941	0.819	1.208*

700   Table 1. Clonal diversity indices for the sexes in the Wrightwood site.

701   Note: *Statistical difference between the sexes in Wrightwood, p-value = 0.001, based on

702   1,000 permutations with subsampling to match population sizes. N = number of ramets

703   sampled, G = number of genets, P_d = clonal diversity, N_{eff} = effective number of

704   genotypes.

Site	N	G	P_d	N_{eff}	Simpson's	Evenness	Shannon (corrected)
Wrightwood	98	45	0.459	36.379	0.983*	0.808	1.753*
Phelan	33	8	0.242	6.368	0.869*	0.796	0.874*

Table 2. Clonal diversity indices for Wrightwood and Phelan sites.

Note: *Statistical difference between Wrightwood and Phelan, p-value = 0.001, based on

1,000 permutations with subsampling to match population sizes. N = number of ramets

sampled, G = number of genets, P_d = clonal diversity, N_{eff} = effective number of

genotypes.

APPENDIX 1

Specimen vouchers were deposited in the California State University, Los Angeles

Herbarium (CSLA).

FIGURE LEGENDS

Figure 1. Clonal lineage decay at different genetic distance thresholds. This figure shows

the number of genets that would be identified at each genetic distance parameter setting.

Figure 2. Spatial distribution of genets. Wrightwood contained 53 unique genets in 98

ramets from 33 patches along three transects (T1, T2, and T3). Genotypic females are

represented as circles and genotypic males as squares. Genets are outlined in dashed

lines. In cases that genets are in different patches, a dashed line connects different ramets

of the same MLG. Presence of phenotypic sex in each patch is indicated with ♀ for

phenotypic females and ♂ for phenotypic males.

Figure 3. CLEARCUT neighbor joining genetic distance tree of ramets from the

Wrightwood and Phelan sites. Branches that join samples from the Wrightwood site

('W') site are blue and those that join the Phelan site ('P') are red. Branches that support

mixed-population groups are purple. Symbols indicate inferred sex.

Figure 4. fastSTRUCTURE membership coefficients for K = 2. Vertical bars represent

individual ramets while colors represent genetic clusters detected (K = 2). Bars on the left

represent ramets from the Phelan site (n = 33) and bars on the right represent ramets from

the Wrightwood site (n = 98). Membership coefficients are plotted for each ramet and are

colored corresponding to the proportion of each ramet MLG that most closely aligns with

each of the two clusters. The dataset used for this analysis contained 2,234 SNPs from

131 sterile ramets from Wrightwood and Phelan where missing data were filled in with

clones, when possible.

Legend:
- ● Genotypic female ramet
- □ Genotypic male ramet
- ⬭ Haplotype
- ♀ Phenotypic female present in patch
- ♂ Phenotypic male present in patch

0.08

Phelan   Wrightwood