Mapping eQTLs With RNA-Seq Reveals Novel SLE Susceptibility Genes, Non-Coding

RNAs, and Alternative-Splicing Events That Are Concealed Using Microarrays

Christopher A. Odhams[1*], Andrea Cortini[1], Lingyan Chen[1], Amy L. Roberts[1], Alfonso Buil[2-3],

Kerrin S. Small[4], Emmanouil T. Dermitzakis[2-3], David L. Morris[1], Timothy J. Vyse[1, 5*],

Deborah S. Cunninghame Graham[*1, 5]

[1]Division of Genetics and Molecular Medicine, King's College London, London, SE1 9RT,

UK. [2]Department of Genetic Medicine and Development, University of Geneva Medical

School, Geneva, CH – 1211 Geneva 4, Switzerland.

[3]Institute of Genetics and Genomics in Geneva, University of Geneva, Geneva, CH – 1211

Geneva 4 Switzerland.

[4]Department of Twin Research and Genetic Epidemiology, King's College London, London,

SE1 7EH, UK.

[5]Division of Immunology, Infection and Inflammatory Disease, King's College London,

London, SE1 1UL, UK.

*Correspondence: Dr Deborah S. Cunninghame Graham

Fax: +44 (0)207 188 2585

Telephone: +44 (0)207 848 8504

Email: deborah.cunninghame-graham@kcl.ac.uk

Abstract

Studies attempting to functionally interpret complex-disease susceptibility loci by GWAS and eQTL integration have predominantly employed microarrays to quantify gene-expression. RNA-Seq has the potential to discover a more comprehensive set of eQTLs and illuminate the underlying molecular consequence. We examine the functional outcome of 39 variants associated with Systemic Lupus Erythematosus (SLE) through integration of GWAS and eQTL data from the TwinsUK microarray and RNA-Seq cohort in lymphoblastoid cell lines. We use conditional analysis and a Bayesian colocalisation method to provide evidence of a shared causal-variant, then compare the ability of each quantification type to detect disease relevant eQTLs and eGenes. We discovered a greater frequency of candidate-causal eQTLs using RNA-Seq, and identified novel SLE susceptibility genes that were concealed using microarrays (e.g. *NADSYN1*, *SKP1*, and *TCF7*). Many of these eQTLs were found to influence the expression of several genes, suggesting risk haplotypes may harbour multiple functional effects. We pinpointed eQTLs modulating expression of four non-coding RNAs; three of which were replicated in whole-blood. Novel SLE associated splicing events were identified in the T-reg restricted transcription factor, *IKZF2*, the autophagy-related gene *WDFY4*, and the redox coenzyme *NADSYN1*, through asQTL mapping using the Geuvadis cohort. We have significantly increased our understanding of the genetic control of gene-expression in SLE by maximising the leverage of RNA-Seq and performing integrative GWAS-eQTL analysis against gene, exon, and splice-junction quantifications. In doing so, we have identified novel SLE candidate genes and specific molecular mechanisms that will serve as the basis for targeted follow-up studies.

Introduction

Genome-Wide Association Studies (GWAS) have successfully identified a large number of genetic loci that contribute to complex-disease susceptibility in humans (1). Evidence suggests these variants are enriched within regulatory elements of the genome and their effects play a central role in modulation of intermediate quantitative phenotypes such gene expression (1–6). Many expression quantitative trait loci (eQTL) mapping studies have since been conducted across a wide-range of ethnicities (7, 8), cell-types (9–16), disease states (17–22) and in response to various environmental stimuli (23, 24) - with each contributing to our understanding of the architecture of human regulatory variation in complex-disease.

In spite of diverse study designs, a significant constraint on the majority of such investigations is the use of 3´-targeted microarrays to profile gene expression. The effects of splicing are less likely to be detected through quantification of pre-defined probes that target common exons of a gene (25) and may explain why only a limited number of susceptibility loci localize to causal eQTL signals (26, 27). Technical limitations of microarrays and noise from the small probe design of exon-arrays, further hinder the accuracy of expression measurements (25, 28–30). RNA-Seq based eQTL mapping studies are beginning to emerge (31, 32) and, although large-scale analysis pipelines are still being streamlined, such types of investigations will greatly increase the likelihood of capturing disease associated eQTLs as quantification of overall gene and independent exon expression, and relative transcript abundance (including novel isoforms and non-coding RNAs) is possible (33–39).

Integrative studies using RNA-Seq to functionally annotate complex-disease susceptibility loci however have been limited (35, 40–44). Moreover, numerous investigations have aimed to explain the functional relevance of susceptibility loci by interrogation of GWAS SNPs themselves in eQTL datasets and simply testing for association with gene expression (45–47). Such inferential observations should be treated with caution as they may possibly be the

result of coincidental overlap between disease association and eQTL signal due to local LD

and general ubiquity of regulatory variants (48). This has become particularly important as

statistical power in eQTL cohorts grow and availability of summary-level data accession

through eQTL data-browsers increases (49–51).

In this investigation, we integrate eQTL data derived from both microarray and RNA-Seq

experiments with our GWAS results in Systemic Lupus Erythematosus (SLE [MIM: 152700]);

a heritable autoimmune disease with undefined aetiology and over 50 genetically associated

loci (52–54). We use summary-level *cis*-eQTL results in lymphoblastoid cell lines (LCLs)

taken from the TwinsUK cohort to directly compare the microarray (9) and RNA-Seq (39)

results in detecting SLE associated eQTLs along with their accompanying eGenes. We

apply a rigorous two-step approach – a combination of conditional (55) and Bayesian

colocalisation (56) analysis – to test for a shared causal variant at each locus. We

demonstrate the benefits of using RNA-Seq over microarrays in eQTL analysis by identifying

not only novel SLE candidate-causal eGenes but also putative molecular mechanisms by

which SLE-associated SNPs may act; including differential exon usage, and expression

modulation of non-coding RNA. Our investigation was extended to include RNA-Seq

expression data in whole blood in order to validate the eQTL signals detected in LCLs and

uncover the differences in genetic control of expression between cell-types. Finally, we

interrogate the Geuvadis RNA-Seq cohort (35) to identify SLE associated alternative-splicing

quantitative trait loci (asQTLs) and highlight the advantages of profiling at various resolutions

to detect eQTLs that would otherwise remain concealed. Through functional annotation of

SLE associated loci using microarray and RNA-Seq derived expression data, we have

supplied comprehensive evidence of the need to use RNA-Seq to detect disease

contributing eQTLs and, in doing so, have suggested novel functional mechanisms that

serve as a basis for future targeted follow-up studies.

Results

Discovery and classification of SLE candidate-causal eQTLs and eGenes

The first part of this study integrated the 39 SLE associated SNPs taken from a recent GWAS in Europeans (**Table 1**) with eQTLs from the TwinsUK gene-expression cohort (n=856) profiled using microarray and RNA-Seq (at both gene-level and exon-level resolutions). To accomplish this, we implemented a two-step pipeline (**Fig. 1**), and subjected the genomic intervals within +/-1Mb of each of the 39 GWAS SNPs to eQTL association analysis against expression quantifications in LCLs (**Table 2**).

Full results of the conditional and colocalisation analysis for each significant association are presented in **S1 Table**, **S2 Table**, and **S3 Table** for microarray, RNA-Seq (gene-level), and RNA-Seq (exon-level), respectively. Statistically significant SLE-associated *cis*-eQTLs showing evidence of a shared causal variant or in very strong LD and close ranking between the disease and *cis*-eQTL signal following conditional and colocalisation analyses were classified as SLE candidate-causal eQTLs as stated in Methods. SLE candidate-causal eGenes were defined as genes whose expression is modulated by the eQTL. The final column of **S1-S3 Tables** indicates whether each GWAS SNP is deemed to be candidate-causal. These SLE candidate-causal eQTLs and eGenes are presented in separate tables contingent on the dataset from which they were generated: results from microarray assessment are listed in **Table 3**, from RNA-Seq (gene-level) in **Table 4**, and from RNA-Seq (exon-level) in **Table 5**. Effect sizes are with respect to the minor allele; risk alleles are highlighted in **Table 1**. Overall, exon-level analysis was the most effective quantification type for the discovery of eQTLs and eGenes compared with gene-level RNA-Seq or microarray analysis following an FDR cut-off of q < 0.05 and conditional and colocalisation thresholding as described.

Table 1

Independent allelic associations at SLE susceptibility loci following meta-analysis with replication study

GWAS SNP (Tag SNP)	GWAS SNP Position (hg19)	Variant Alleles	Odds Ratio (CI)	*P*-value	GENCODE v10 Annotation	Annotated eGenes from microarray LCLs
rs2476601	1:114377568	G/**A**	1.43 (1.34-1.53)	1.10E-28	*PTPN22* Missense	
rs1801274	1:161479745	**G**/A	1.16 (1.11-1.21)	1.04E-12	*FCGR2A* Missense	*FCGR2B*
rs10912578 (rs844663)	1:173251856	G/**A**	1.27 (1.22-1.33)	4.16E-19	*LOC100506023* Intronic	
rs10753074 (rs1935325)	1:173346343	**T**/C	1.21 (1.15-1.26)	5.82E-12	*LOC100506023* Intronic	
rs17849501	1:183542323	C/**T**	2.10 (1.95-2.26)	3.45E-88	*NCF2* Synonymous	*SMG7*
rs3024505	1:206939904	G/**A**	1.17 (1.11-1.24)	4.64E-09	1kb 3' of *IL10*	
rs9782955	1:236039877	**C**/T	1.16 (1.11-1.22)	1.25E-09	*LYST* Intronic	*LYST*
rs268134	2:65608363	**G**/A	1.21 (1.15-1.27)	1.14E-10	*SPRED* Intronic	*SPRED2*
rs2111485	2:163110536	**G**/A	1.15 (1.11-1.2)	1.27E-11	8.9kb 5' of *FAP*	
rs11889341	2:191943742	C/**T**	1.70 (1.64-1.75)	2.48E-75	*STAT4* Intronic	
rs6736175 (rs16833249)	2:191946322	T/**C**	1.24 (1.19-1.29)	9.17E-17	*STAT4* Intronic	
rs3768792	2:213871709	A/**G**	1.24 (1.17-1.31)	1.21E-13	*IKZF2* 3'-UTR	
rs9311676 (rs11714389)	3:58470351	**C**/T	1.17 (1.13-1.22)	3.06E-14	*LOC101929223* NC	*ABHD6*
rs564799	3:159728987	**C**/T	1.14 (1.09-1.18)	1.54E-09	*IL12A-AS1* Intronic	*IL12A*
rs10028805	4:102737250	**G**/A	1.20 (1.15-1.25)	4.31E-17	*BANK1* Intronic	*BANK1*
rs7726414 (rs17167273)	5:133431834	C/**T**	1.45 (1.32-1.58)	4.44E-16	19kb 5' of *TCF7*	
rs10036748	5:150458146	C/**T**	1.38 (1.32-1.45)	1.27E-45	*TNIP1* Intronic	
rs2431697	5:159879978	**T**/C	1.26 (1.21-1.31)	8.01E-28	15kb 5' of *hsa-mir-146a*	
rs6568431	6:106588806	C/**A**	1.21 (1.15-1.27)	5.04E-14	31kb 3' of *PRDM1*	
rs6932056	6:138242437	T/**C**	1.83 (1.65-2.02)	1.97E-31	22kb 3' of *RP11-10J5.1*	
rs849142	7:28185891	**T**/C	1.14 (1.1-1.19)	8.61E-11	*JAZF1* Intronic	
rs4917014	7:50305863	**T**/G	1.18 (1.13-1.24)	6.39E-14	8.5kb 3' of *AC020743.4*	
rs3757387 (rs4728142)	7:128576086	T/**C**	1.45 (1.4-1.5)	1.14E-48	1.6kb 5' of *IRF5*	
rs35000415 (rs10488631)	7:128585616	C/**T**	1.83 (1.76-1.9)	1.20E-60	*IRF5* Intronic	*IRF5, TNPO3*
rs2736340	8:11343973	C/**T**	1.29 (1.22-1.37)	6.28E-20	7.5kb 5' of *BLK*	*BLK*
rs2663052	10:50069395	**G**/A	1.16 (1.1-1.22)	5.25E-09	*WDFY4* Intronic	*WDFY4*
rs4948496	10:63805617	T/**C**	1.14 (1.1-1.19)	1.04E-10	*ARID5B* Intronic	
rs12802200 (rs2396545)	11:566936	**C**/A	1.23 (1.15-1.31)	8.81E-10	*MIR210HG* NC	
rs2732549	11:35088399	**A**/G	1.24 (1.19-1.29)	1.20E-23	46kb 3' of *PDHX*	
rs3794060	11:71187679	T/**C**	1.23 (1.18-1.29)	1.32E-20	*NADSYN1* Intronic	
rs7941765 (rs6590343)	11:128499000	**C**/T	1.14 (1.1-1.19)	1.35E-10	547bp 3' of *RP11-744N12.3*	
rs10774625	12:111910219	**A**/G	1.13 (1.08 - 1.18)	4.09E-09	*ATXN2* Intronic	
rs1059312	12:129278864	A/**G**	1.17 (1.12-1.21)	1.48E-13	*SLC15A4* Synonymous	
rs4902562	14:68731458	G/**A**	1.14 (1.09-1.19)	6.15E-10	*RAD51B* Intronic	
rs2289583	15:75311036	C/**A**	1.19 (1.14-1.24)	6.22E-15	*SCAMP5* Intronic	*CSK, ULK3, MPI*
rs9652601	16:11174365	**G**/A	1.21 (1.15-1.26)	7.42E-17	*CLEC16A* Intronic	
rs34572943 (rs9936831)	16:31272353	G/**A**	1.71 (1.61-1.81)	3.39E-76	*ITGAM* Intronic	
rs11644034	16:85972612	**G**/A	1.25 (1.19-1.32)	9.58E-18	16kb 3' of *IRF8*	
rs2286672	17:4712617	C/**T**	1.25 (1.16-1.35)	2.93E-09	*PLD2* Missense	*RNF167*
rs2941509	17:37921194	C/**T**	1.35 (1.22-1.49)	7.98E-09	*IKZF3* 3'-UTR	
rs2304256	19:10475652	**C**/A	1.24 (1.17-1.31)	3.50E-13	*TYK2* Missense	
rs7444	22:21976934	T/**C**	1.27 (1.21-1.33)	1.84E-22	*UBE2L3* 3'-UTR	*UBE2L3*

SLE GWAS SNPs taken forward for *cis*-eQTL association analysis. Results from post-replication meta-analysis as described (57). SNP with the lowest P-value post meta-analysis or the SNP with the greatest evidence of a missense effect as defined by a Bayes Factor reported. Autosomal, non-MHC SNPs with MAF > 0.05 included in analysis only (39 total). Risk alleles are highlighted in bold type – minor allele on right. Functional annotation from HaploReg v4.0 using GENCODE genes v10. Stated eGenes detected from microarray studies in LCLs listed as described (57).

Table 2
Details of genotype-expression (eQTL) cohorts used in study

Cohort Name	TwinsUK (MuTHER)				Geuvadis
Total subjects	856				373
Ethnicity	EUR (UK)				EUR (CEU, GBR, FIN, TSI)
Sex	F				M / F
Age	37–85				*NA*
Investigation	Comparison of candidate-causal eQTL and eGene detection between microarray and RNA-Seq			Validation and comparison of LCL RNA-Seq discoveries in whole blood	Identification of asQTLs using RNA-Seq
Citation	*Grundberg et. al* (9)	*Buil et. al* (39)	*Buil et. al* (39)	*Buil et. al* (39)	*Lappalainen et. al* (35)
Expression profile type	**Microarray**	**RNA-Seq**	**RNA-Seq**	**RNA-Seq**	**RNA-Seq**
Unit of expression	**Probe**	**Gene**	**Meta-exon**	**Meta-exon**	**Splice-junction**
Cell-type	LCL	LCL	LCL	Whole Blood	LCL
Subjects used in analysis	777	683	765	384	373
Data format	Genevar (summary results)	Read-count	Summary eQTL results	Summary eQTL results	Raw sequence alignments
RNA Platform	Illumina HT-12 V3	Illumina HiSeq2000		Illumina HiSeq2000	Illumina HiSeq2000
RNA-Seq mapper	*NA*	BWA v0.5.9 (GRCh37/hg19)		BWA v0.5.9 (GRCh37/hg19)	GEM v1.349 (GRCh37/hg19)
Reference transcriptome	*NA*	GENCODE V10		GENCODE V10	GENCODE V10
RNA-Seq read length	*NA*	49-bp PE		49-bp PE	75-bp PE

Breakdown of genotype-expression (eQTL) cohorts used in analysis. TwinsUK cohort in lymphoblastoid cell lines (LCLs) used for microarray and RNA-Seq comparison (profiled at gene and meta-exon resolution); meta-exons are described as non-redundant overlapping portions of exons generated flattening of the transcriptome annotation. All TwinsUK (MuTHER) samples used in analysis are derived from the original 856 individuals. Validation of LCL data in whole blood carried out at meta-exon level using 384 of the 856 individuals. Geuvadis cohort used for asQTL identification; splice-junction quantifications were generated by Altrans (57) from the raw sequence alignments. Summary eQTL results include only the eQTL association results per test (where full genotype and expression data were not obtainable).

Table 3
Candidate-causal eQTLs and associated eGenes detected using microarray (probe-level)

Risk Locus	GWAS SNP	eGene	Probe ID	β	*P*-Value	FDR (q)
1p13.2	rs2476601	*BCL2L15*	ILMN_1655722	-0.075	8.26E-04	2.73E-02
3q25.33	rs564799	*IL12A*	ILMN_1671353	0.113	4.91E-13	4.86E-11
3p14.3	rs11714389	*ABHD6*	ILMN_1706344	-0.130	1.41E-13	1.63E-11
		PDHB	ILMN_1739274	-0.055	2.55E-05	1.26E-03
4q24	rs10028805	*BANK1*	ILMN_1661646	0.195	7.98E-13	6.91E-11
8p23.1	rs2736340	*BLK*	ILMN_1668277	-0.407	7.69E-25	1.07E-22
		FAM167A	ILMN_1687213	0.412	1.48E-43	1.03E-40
15q24.2	rs2289583	*CSK*	ILMN_1754121	-0.059	9.56E-07	5.10E-05
		MPI	ILMN_1761262	0.058	1.45E-04	5.58E-03
		ULK3	ILMN_1679495	-0.071	1.78E-09	1.12E-07
17p13.2	rs2286672	*INCA1*	ILMN_1704380	0.035	7.69E-04	2.66E-02
22q11.21	rs7444	*UBE2L3*	ILMN_1677877	-0.183	4.97E-25	8.61E-23

GWAS SNPs deemed to be candidate-causal eQTLs using microarray expression data profiled from 777 individuals of the TwinsUK cohort in lymphoblastoid cell lines. 768 probes, corresponding to 559 genes, were tested against in *cis* to the 39 GWAS SNPs.

Table 4
Candidate-causal eQTLs and associated eGenes detected using RNA-Seq (Gene-level)

Risk Locus	GWAS SNP	eGene	β	*P*-Value	FDR (q)
1p13.2	rs2476601	DCLRE1B	-0.330	8.41E-04	1.91E-02
3q25.33	rs564799	IL12A	0.372	1.07E-11	9.70E-10
3p14.3	rs11714389	ABHD6	0.320	7.00E-09	2.28E-07
		RPP14	-0.285	4.15E-07	1.64E-05
4q24	rs10028805	BANK1	-0.316	2.37E-08	1.29E-06
5q31.1	rs17167273	SKP1	-0.564	1.57E-05	5.06E-04
		TCF7	-0.534	2.89E-05	8.27E-04
5q33.3	rs2431697	MIR146A	0.256	1.52E-06	6.89E-05
8p23.1	rs2736340	FAM167A	0.668	1.40E-33	7.62E-31
		BLK	-0.596	3.13E-28	8.51E-26
		RP11-148O21.2	-0.591	1.11E-26	2.01E-24
11p15.5	rs2396545	HRAS	-0.248	1.56E-04	4.04E-03
11q13.4	rs3794060	NADSYN1	-0.620	6.43E-24	8.74E-22
		DHCR7	-0.200	1.70E-03	3.43E-02
15q24.2	rs2289583	ULK3	0.303	3.62E-07	1.79E-05
		UBE2Q2	0.249	5.66E-05	1.54E-03
		FAM219B	-0.198	1.18E-03	2.57E-02
		CSK	0.194	1.64E-03	3.43E-02
22q11.21	rs7444	UBE2L3	0.345	2.28E-06	9.54E-05

GWAS SNPs deemed to be candidate causal eQTLs using RNA-Seq expression data profiled from 683 individuals of the TwinsUK cohort in lymphoblastoid cell lines at gene-level resolution. 520 genes were tested against in *cis* to the 39 GWAS SNPs.

Table 5
Candidate causal eQTLs and associated eGenes detected using RNA-Seq (Exon-level)

Risk Locus	GWAS SNP	eGene	Meta-exon ID (chr. start. end)	β	*P*-Value	FDR (q)
1p13.2	rs2476601	*BCL2L15*	1.114420790.114424619	0.226	2.71E-10	1.80E-08
			1.114429871.114430169	0.199	2.72E-08	1.28E-06
		DCLRE1B	1.114449618.114449783	0.110	2.23E-03	3.85E-02
		MAGI3	1.114225519.114228545	0.115	1.41E-03	2.62E-02
1q32.1	rs3024505	*FCAMR*	1.207133024.207134568	0.114	1.62E-03	2.94E-02
		IL10	1.206945616.206945839	0.174	1.26E-06	4.88E-05
		IL24	1.207076321.207077484	0.108	2.81E-03	4.63E-02
2q24.2	rs2111485	*IFIH1*	2.163123589.163123889	0.119	1.00E-03	1.98E-02
			2.163128736.163128897	0.108	2.69E-03	4.49E-02
			2.163130305.163130454	0.116	1.25E-03	2.37E-02
			2.163136506.163136622	0.138	1.31E-04	3.29E-03
			2.163137838.163138055	0.110	2.34E-03	4.01E-02
			2.163163219.163163680	0.127	4.45E-04	9.72E-03
3q25.33	rs564799	*IFT80*	3.159996981.159997152	0.109	2.65E-03	4.44E-02
		IL12A	3.159706537.159706961	0.202	1.66E-08	8.24E-07
			3.159710799.159711631	0.286	7.56E-16	8.33E-14
			3.159713191.159713806	0.281	2.23E-15	2.31E-13
		RP11-432B6.3	3.159996981.159997152	0.109	2.65E-03	4.44E-02
		SMC4	3.160129546.160129872	0.120	8.69E-04	1.74E-02
			3.160137146.160137331	0.112	1.97E-03	3.48E-02
			3.160141213.160141438	0.112	1.98E-03	3.48E-02
			3.160141549.160141628	0.110	2.36E-03	4.03E-02
			3.160149431.160149613	0.126	4.69E-04	1.02E-02
			3.160150071.160150303	0.122	7.51E-04	1.53E-02
			3.160150814.160150997	0.137	1.41E-04	3.47E-03
3p14.3	rs11714389	*ABHD6*	3.58223233.58223643	0.132	2.62E-04	6.10E-03
			3.58242289.58242432	0.173	1.57E-06	5.89E-05
			3.58252916.58253072	0.174	1.34E-06	5.11E-05
			3.58255048.58255161	0.207	7.18E-09	3.76E-07
			3.58256659.58256791	0.190	1.21E-07	5.27E-06
			3.58260385.58260542	0.200	2.47E-08	1.18E-06
			3.58270924.58271180	0.221	6.02E-10	3.75E-08
			3.58279316.58280461	0.345	7.89E-23	1.13E-20
4q24	rs10028805	*BANK1*	4.102946358.102946666	0.264	1.07E-13	9.91E-12
			4.102981368.102981546	0.253	1.19E-12	9.92E-11
5q31.1	rs17167273	*SKP1*	5.133492082.133494319	0.276	7.20E-15	7.17E-13
5q33.3	rs2431697	*MIR146A*	5.159912306.159914433	0.248	3.36E-12	2.52E-10
			5.159895275.159895447	0.151	2.74E-05	7.97E-04
8p23.1	rs2736340	*BLK*	8.11351510.11352100	0.462	9.76E-42	3.16E-39
			8.11366659.11367400	0.350	1.72E-23	2.55E-21
			8.11412252.11412398	0.469	5.06E-43	2.01E-40
			8.11414166.11414346	0.448	6.05E-39	1.65E-36
			8.11415471.11415547	0.445	2.13E-38	5.26E-36
			8.11417842.11418961	0.485	2.45E-46	2.11E-43
			8.11420488.11420619	0.440	1.49E-37	2.96E-35
			8.11421412.11422113	0.446	1.32E-38	3.43E-36
		FAM167A	8.11278972.11282145	0.469	3.48E-43	1.64E-40
			8.11301540.11302395	0.463	7.22E-42	2.49E-39
		RP11-148O21.2	8.11415975.11416256	0.454	3.63E-40	1.04E-37
			8.11416421.11416495	0.325	2.69E-20	3.67E-18
			8.11417293.11417529	0.395	5.49E-30	9.17E-28
		RP11-148O21.4	8.11413760.11414170	0.353	6.24E-24	9.50E-22
			8.11415399.11415531	0.391	2.15E-29	3.48E-27
11p15.5	rs2396545	*ANO9*	11.417933.418589	0.180	5.66E-07	2.32E-05
			11.418720.419415	0.207	7.70E-09	3.95E-07
			11.419582.420860	0.196	4.54E-08	2.06E-06
			11.420929.421042	0.163	6.11E-06	2.03E-04
			11.428088.428199	0.119	9.93E-04	1.97E-02
			11.428475.428639	0.181	4.99E-07	2.07E-05
			11.428722.428826	0.164	5.44E-06	1.83E-04
			11.429474.430179	0.203	1.52E-08	7.63E-07
			11.430269.430403	0.198	3.28E-08	1.52E-06
			11.431694.433459	0.144	6.20E-05	1.69E-03
			11.433815.433937	0.156	1.45E-05	4.48E-04
			11.434024.434098	0.127	4.24E-04	9.30E-03
		HRAS	11.533277.533358	0.136	1.64E-04	3.97E-03
			11.533766.533976	0.151	2.72E-05	7.97E-04
			11.534212.534375	0.138	1.29E-04	3.24E-03
		PHRF1	11.587259.588497	0.126	4.86E-04	1.05E-02
			11.597395.597570	0.118	1.03E-03	2.01E-02
			11.607066.609720	0.170	2.37E-06	8.56E-05
			11.611634.612222	0.142	8.00E-05	2.09E-03
		RASSF7	11.561547.561892	0.160	9.26E-06	2.98E-04

Region	SNP	Gene	Position			
		RNH1	11.498457.498627	0.152	2.58E-05	7.72E-04
			11.504824.505881	0.309	2.28E-18	2.82E-16
			11.507113.507300	0.126	4.55E-04	9.90E-03
		TMEM80	11.700615.701127	0.131	2.75E-04	6.37E-03
11q13.4	rs3794060	*DHCR7*	11.71145460.71147019	0.246	5.41E-12	4.00E-10
			11.71148858.71148989	0.172	1.66E-06	6.15E-05
			11.71149795.71150129	0.179	6.25E-07	2.51E-05
			11.71152273.71152486	0.163	5.77E-06	1.93E-04
			11.71153309.71153399	0.207	7.53E-09	3.90E-07
			11.71155003.71155299	0.175	1.14E-06	4.45E-05
			11.71155901.71156004	0.150	3.12E-05	9.03E-04
		NADSYN1	11.71175099.71175554	0.227	2.04E-10	1.37E-08
			11.71185441.71186668	0.545	1.79E-60	9.26E-57
			11.71187079.71188484	0.538	1.06E-58	2.75E-55
			11.71189441.71190128	0.339	5.32E-22	7.45E-20
			11.71190340.71191320	0.419	7.54E-34	1.30E-31
			11.71191801.71193071	0.245	6.27E-12	4.57E-10
			11.71195358.71196694	0.280	2.94E-15	2.98E-13
			11.71202880.71202949	0.166	3.68E-06	1.28E-04
			11.71207481.71208657	0.248	3.30E-12	2.51E-10
			11.71209398.71211081	0.196	4.82E-08	2.15E-06
			11.71214910.71216920	0.297	5.05E-17	5.81E-15
		RP11-660L16.2	11.71159720.71159931	0.482	7.76E-46	5.74E-43
			11.71162736.71163203	0.530	1.44E-56	2.49E-53
12q24.12	rs10774625	*HECTD4*	12.112601913.112602079	0.113	1.81E-03	3.25E-02
15q24.2	rs2289583	*CSK*	15.75090574.75090653	0.153	2.27E-05	6.82E-04
			15.75091613.75091832	0.140	9.68E-05	2.48E-03
			15.75092753.75092846	0.143	7.38E-05	1.97E-03
			15.75093164.75093263	0.117	1.15E-03	2.20E-02
			15.75093863.75093936	0.140	1.08E-04	2.76E-03
			15.75094036.75094231	0.167	3.20E-06	1.14E-04
			15.75094338.75094424	0.129	3.65E-04	8.10E-03
		FAM219B	15.75192329.75195127	0.148	4.08E-05	1.15E-03
		MPI	15.75182867.75182995	0.150	3.29E-05	9.41E-04
			15.75185002.75185143	0.250	2.32E-12	1.87E-10
			15.75185479.75185661	0.191	1.08E-07	4.75E-06
			15.75189352.75189560	0.212	3.48E-09	1.92E-07
		ULK3	15.75128459.75129585	0.286	7.98E-16	8.61E-14
			15.75130607.75130685	0.130	3.14E-04	7.11E-03
			15.75130984.75131086	0.142	7.90E-05	2.09E-03
			15.75131351.75131391	0.145	5.99E-05	1.64E-03
			15.75131609.75131714	0.182	4.02E-07	1.69E-05
			15.75131899.75132054	0.150	3.16E-05	9.08E-04
			15.75132575.75132657	0.120	8.45E-04	1.70E-02
			15.75132839.75132982	0.133	2.32E-04	5.48E-03
			15.75133746.75133850	0.116	1.35E-03	2.54E-02
			15.75134416.75134536	0.129	3.46E-04	7.79E-03
			15.75134621.75134761	0.130	3.00E-04	6.88E-03
22q11.21	rs7444	*UBE2L3*	22.21965146.21965332	0.216	1.65E-09	9.46E-08
			22.21975804.21978323	0.422	2.19E-34	4.20E-32

GWAS SNPs deemed to be candidate causal eQTLs using RNA-Seq expression data profiled from 765 individuals of the TwinsUK cohort in lymphoblastoid cell lines at exon-level resolution. 4,786 exons, corresponding to 716 genes were tested against in *cis* to the 39 GWAS SNPs.

RNA-Seq better annotates SLE risk loci by revealing known and novel candidate genes

Fig. 2 illustrates the clear improvement of RNA-Seq relative to microarray in the discovery of candidate-causal eQTLs and their corresponding eGenes when annotating complex-disease susceptibility loci. With RNA-Seq, we replicated known SLE associated eQTLs and eGenes that were previously discovered using microarray. These included rs564799 for *IL12A*, rs2736340 for *BLK*, rs9311676 *ABHD6*, and rs2289583 for *ULK*, *CSK*, and *MPI* (**Tables 4**, and **5**; **Table 1** for previously reported associations in LCLs). Several eQTLs for eGenes that have been extensively studied in terms of their role in SLE pathogenesis were consistent across all three platforms: for example, rs10028805 for *BANK1* (57) and rs7444 for *UBE2L3* (58). **Fig. 2** also shows how exon-level RNA-Seq analysis led to the greatest frequency of candidate-causal eQTLs and eGenes than with either gene-level RNA-Seq or microarray (**S1 Fig.** and **S2 Fig.**). A total of 14 eQTLs modulating expression of 34 eGenes were detected using exon-level RNA-Seq contrasted to 11 eQTLs and 19 eGenes at gene-level RNA-Seq and only 8 eQTLs with 12 eGenes identified using microarray (**S1 Fig.** and **S2 Fig.**). Only one eQTL (rs2286672) and two eGenes (*PDHB*, *INCA1*) were found by microarray exclusively. These associations were either not significant post multiple testing using either RNA-Seq method, or were not deemed candidate-causal (**S1 Table S1** and **S3 Table**). In total 8 eQTLs regulating expression of 27 eGenes were detected using RNA-Seq but missed using microarray (**S3 Fig. S3** and **S4 Fig.**). Interestingly, exon-level analysis led to the greatest frequency of non-candidate-causal associations. Only 14 of the 34 significant eQTLs, q < 0.05, showed evidence of a shared causal variant post conditional and colocalisation testing (**S1 Fig.** and **S2 Fig.**).

Several eGenes known to be involved in the pathogenesis of SLE were identified using RNA-Seq exclusively and not reported in previous microarray-based eQTL studies in LCLs (**Table 1**). These include *IL10*, *IFIH1*, and the microRNA *MIR146A*. We believe a handful of

other eGenes unique to RNA-Seq to be novel SLE candidate genes. *NADSYN1* (NAD

Synthetase), *HECTD4* (HECT Domain E3 Ubiquitin Protein Ligase 4), *SKP1* (S-Phase

Kinase-Associated Protein 1) and *TCF7* (T-cell specific Transcription Factor 7) are examples

of novel eGenes.

RNA-Seq eQTL analysis reveals eQTLs regulating multiple eGenes

Fig. 2 also illustrates that many of the eQTLs discovered using RNA-Seq regulate multiple

eGenes. Exon-level analysis generated the greatest ratio (2.42) of candidate-causal eGenes

to eQTLs (**S5 Fig.**); suggesting that disease-associated haplotypes may be more functionally

potent and harbour multiple gene regulatory effects than previously thought.

One example of this effect is at the *TCF7-SKP1* locus where the disease-association signal

encompasses both genes (**S6 Fig.**). Both *TCF7* and *SKP1* were classified as candidate-

causal against the GWAS SNP rs7726414 using gene-level RNA-Seq (**Table 5**), and *SKP1*

but not *TCF7* at exon-level (**Table 4**). Interestingly, a missense variant of *TCF7* has been

implicated in Type 1 Diabetes risk (59), but there is only weak LD (r^2<0.4) between this

missense variant and rs7726414 (19 kb upstream of *TCF7*) or any protein-coding variants of

TCF7, suggesting that in SLE the causal mechanism may be dysregulated gene expression

of *TCF7* rather than a missense change *per se*. *SKP1*, part of the ubiquitin ligase complex, is

thought to stabilize the conformation of E3 ligases and its expression has been shown to be

upregulated in lung-cancer (60). Neither of these effects were present in the microarray data

as there is another more significant *cis*-eQTL, rs17167273, for the *TCF7* probe

(ILMN_1676470), which is not correlated with the rs7726414[T] risk variant (r^2:0.02, **S1**

Table). In addition, rs7726414[T] was not a significant eQTL for the *SKP1* microarray probe

(ILMN_1790710) (*P*=0.16). Our RNA-Seq eQTL analyses indicated that rs7726414[T]

represents a novel eQTL for dysregulated gene expression of *SKP1* in SLE (**S6 Fig.**). We

believe both *TCF7* and *SKP1* to be highly plausible candidate genes as the literature

suggests that knockdown of *TCF7* results in impaired stem cell potency and gene

expression regulation of CD34+ cells (61), whilst the mouse knockout of *SKP1* develop

highly penetrant T-cell lymphomas (62).

Other examples where either gene- or exon-level RNA-Seq analysis identified multiple

eGenes are at rs3024505 (*IL10*, *IL24*, and *FCAMR*), rs12802200 and rs3794060 (**Fig. 2**). In

the immune-gene concentrated 11p15.5 region (**S7 Fig.**), the GWAS SNP rs12802200, is an

eQTL for six eGenes, (*HRAS, TMEM80*, *RNH1*, *ANO9*, *PHRF1* and *RASSF7*; **Tables 4 and**

5) following exon- and/or gene-level RNA-Seq analysis, supporting our recent observations

of rs12802200 being a *cis*-eQTL for six genes across multiple immune cell-types at this

locus (63) and earlier reports demonstrating that rs12802200 was an eQTL for *TMEM80*

(whole blood(64)) and for both *RNH1* and *TMEM80* (LCLs (35)). 11p15.5 is also an

important regulatory region in non-immune types, because a total of eight eGenes in this

region have been identified in multiple non-immune cell types (65), four of which (*HRAS*,

TMEM80, *RNH1*, *ANO9*) were captured by our RNA-Seq analysis in LCLs. Since SNPs

within this susceptibility locus have also been previously shown to be correlated with

increased autoantibody production and interferon-α activity in sufferers of SLE (66), further

investigation will be required to elucidate a potential mechanism by which dysregulation of

gene expression may contribute to these effects. In this gene-dense region (**S7 Fig.**), none

of the six candidate-causal eGenes had an annotated microarray probe that passed quality

control.

In the 11q13.4 region, the intronic GWAS SNP rs3794060[C], was classified using gene-

level quantification as being a candidate-causal eQTL for both *NADSYN1* (NAD Synthetase

1) and *DHCR7* (7-Dehydrocholesterol Reductase) (**Table 4, Fig. 3A** and **S8 Fig.**). At exon-

level resolution, *NADSYN1* was also deemed candidate-causal, as well as the non-coding

RNA *RP11-660L16.2* (**Table 5**; described in following section). The risk variant rs3794060[C]

has been correlated with reduced circulating 25-hydroxy vitamin D concentrations (67) with

reduced vitamin D levels being recently associated with increased disease activity of SLE

(68). Interestingly, although a mouse knockout of *DHCR7* showed reduced serum and tissue

cholesterol levels (69), no autoimmune phenotype has been described at this locus. Given

the role oxidative stress plays in promoting inflammation and triggering autoimmunity

through tissue damage (70), it will be interesting to elucidate the role that dysregulation of

NADSYN1 expression plays in this process. Visualization of exon-level association data of

NADSYN1 against rs3794060 suggests a potential splicing mechanism affecting meta-exons

11 and 12 (P=1.79^{E-60}, 1.06^{E-58} respectively; **Fig. 3B**). The probe for *DHCR7*

(ILMN_1815626) showed a weak, but not statistically significant, association with rs3794060

(P=2.9x10^{-03}), whilst *NADSYN1* had no annotated probe in the microarray dataset.

RNA-Seq uncovers the role of non-coding RNA modulation at SLE susceptibility loci

Quantification of non-coding polyadenylated RNAs in the TwinsUK LCL cohort through RNA-

Seq revealed three candidate-causal eQTLs influencing the expression of four non-coding

eGenes (**Table 6**); none of which were captured using microarray.

We validated the effect at rs2431697 (5q33.3) where it is documented that the protective

minor allele [C] is associated with expression upregulation of the miRNA *MIR146A*, a

negative regulator of the type I Interferon pathway (71) (**Fig. 4A**). The best eQTL for

MIR146A at gene-level was rs2431697 (P=1.5x10^{-06}) and also at exon level for both of its

exons (P=3.4x10^{-12} and 1.2x10^{-04}). The decrease in rs2431697[T]-dependent expression of

MIR146A reported in peripheral blood leukocytes of SLE patients disrupts binding of

transcription factor Ets-1, uncouples of the type-1 IFN response (71), thereby increasing the

inflammatory response.

Fig. 4B shows GWAS variant rs2736340, and other SNPs in tight LD (r^2>0.8), within the

FAM167A-BLK (8p23.1) bi-directional promoter region. In this study we detected eQTLs at

the known eGenes *BLK* (B lymphocyte kinase) and *FAM167A* (Family with Sequence

Similarity 167, Member A) by all quantification methods (**S9 Fig.**). The rs2736340[T] risk

allele causes decreased expression of *BLK* (P=3.2x10^{-28}) and increased expression of

FAM167A (P=1.4x10^{-33}) at RNA-Seq gene-level (**Table 4**). This effect has been previously

described in microarray studies (72) - with reduced promoter activity of *BLK* leading to

altered B-cell development (73). The eQTL lies within a region of the genome subject to

multiple regulatory effects, with a 24bp region immediately around the variant containing

strong chromatin marks (peak height>60) in LCLs for H3K9me3 (gene repression) and

H3K4me3 (low expression when present in the promoter of CpG genes) assayed as part of

the ENCODE project (**S9 Fig.**). Interestingly, exon-level RNA-Seq analysis revealed that

rs2736340 also appeared to modulate the expression of two non-coding RNAs antisense to

the 3' region of *BLK*. These are: *RP11-148O21.2* and *RP11-148O21.4* (**Fig. 4B**). The SNP,

rs2736340, significantly reduced the expression of both *BLK* and three exons of *RP11-*

148O21.2 and the two exons of *RP11-148O21.4* (**Table 6**). Our analyses indicated that

expression disruption of these antisense RNAs caused by SLE risk variants represent a

potential novel mechanism at the locus.

There is an rs3794060 allele-dependent expression modulation of both exons of another

non-coding eGene, *RP11-660L16.2I* (**Fig. 4C**), which is located in the bi-directional promoter

between *DHCR7* and *NADSYN1*. The best eQTL for those exons is highly correlated with

the GWAS SNP (rs2282621, r^2:0.99). Both *DHCR7* and *NADSYN1* are candidate-causal

eGenes regulated in the same downward direction at RNA-Seq gene-level with respect to

the risk allele rs3794060[T] (P=1.7x10^{-03} and P=6.4x10^{-24}, respectively) (**S8 Fig.** and **Table**

4).

Table 6
Candidate-causal non-coding eGenes discovered using RNA-Seq (exon-level)

Risk Locus	GWAS SNP	eGene	Exon ID (chr. start. end)	β	*P*-Value	FDR (q)
5q33.3	rs2431697	*MIR146A*	5.159912306.159914433	0.248	3.36E-12	2.52E-10
			5.159895275.159895447	0.151	2.74E-05	7.97E-04
8p23.1	rs2736340	*RP11-148O21.2*	8.11415975.11416256	0.454	3.63E-40	1.04E-37
			8.11416421.11416495	0.325	2.69E-20	3.67E-18
			8.11417293.11417529	0.395	5.49E-30	9.17E-28
		RP11-148O21.4	8.11413760.11414170	0.353	6.24E-24	9.50E-22
			8.11415399.11415531	0.391	2.15E-29	3.48E-27
11q13.4	rs3794060	*RP11-660L16.2*	11.71159720.71159931	0.482	7.76E-46	5.74E-43
			11.71162736.71163203	0.530	1.44E-56	2.49E-53

GWAS SNPs deemed to be candidate causal eQTLs for non-coding eGenes using RNA-Seq expression data profiled from 765 individuals of the TwinsUK cohort in lymphoblastoid cell lines at exon-level resolution.

Confirmation of LCL candidate-causal eQTLs and eGenes using whole-blood RNA-Seq

To validate our LCL findings in a primary tissue-type, we extended our analytical pipeline to include an exon-level RNA-Seq dataset in 384 whole-blood samples from the TwinsUK cohort (**Table 2**). The full results of these analyses are provided in **S4 Table**.

We observed good correlation between LCLs and whole-blood effect-sizes (β) of GWAS SNPs against all matched *cis* exon-level associations (R^2=0.74; **S10 Fig.**). Seven of the 39 GWAS SNPs were classified as candidate-causal eQTLs in whole-blood, modifying 19 candidate-causal eGenes (**Table 7**). All seven of the whole-blood eQTLs and 15 of the 19 eGenes were deemed candidate-causal in LCLs, suggesting strong conservation across whole-blood cell types (**S11 Fig.**). The remaining four eGenes specific to whole-blood were: *PXK* (rs9311676); *IRF7* and *TALDO1* (rs12802200); and *SCAMP2* (rs2289583) (**Table 7**). Interestingly, the eQTLs regulating these four eGenes in whole-blood also regulated multiple eGenes in LCLs (**S11 Fig.**), implying that they tag highly regulatory haplotypes that may cause cell-type specific gene-expression disruption across the entire locus (same eQTL regulating different eGenes across different cell-types). Three of the four candidate-causal non-coding eGenes from LCLs were found in whole-blood (*RP11-148O21.2*, *RP11-148O21.4*, and *RP11-660L16.2*.). The LCL-specific *MIR146A* eGene association with GWAS SNP rs2431697 was not deemed to be significant (*P*=0.32), which is likely to be a result of its lymphocyte-specific gene expression profile (71) that is diluted in the heterogeneous population of whole-blood cell-types.

Inspection of specific exons modulated by the GWAS SNPs in each cell-type revealed instances of variability in the genetic control of exon usage between cell-types. A known splicing event in B-cells caused by branch-point SNP rs17266594 results in the loss of exon 2 in susceptibility gene *BANK1* which subsequently leads to B-cell hyper-responsiveness (57) (**S12 Fig.**). In whole-blood the GWAS variant, rs10028805, is associated with altered

expression of exon 2 (P=8.4x10^{-05}), with the best *cis*-eQTL for this effect being in near-

perfect LD (rs4411998; r^2:0.98). Both rs10028805 and rs4411998 are in strong LD with the

branch-point SNP (r^2:0.9). In LCLs however, the best *cis*-eQTL for exon 2, rs4572885

(P=9.74 x10^{-23}), has a large effect but is less correlated with the GWAS SNP (r^2:0.65) and

conditional analysis judges the effect of the GWAS SNP to be independent to the best *cis*-

eQTL for exon 2 (**S3 Table**). Interestingly, there is low correlation between the branch-point

SNP rs17266594 and the best *cis*-eQTL for exon 2 in LCLs (r^2:0.42); suggesting the

regulatory mechanism of exon 2 splicing in *BANK1* may be under two separate genetic

influences between the two cell-groups (**S12 Fig.**).

We saw a near identical pattern of differential exon usage within eGene *NADSYN1* between

LCLs and whole-blood driven by the GWAS SNP rs37940460 or tightly correlated variants

(**S13 Fig.**, **Table 7**). Variation at rs37940460 appeared to drive extensive expression

disruption of two meta-exons (11 and 12) of *NADSYN1* located near the centre of the gene

(meta-exon 11: LCL P=1.79x10^{-60}; whole-blood P=1.28x10^{-27}; meta-exon 12: LCL

P=1.06x10^{-58}; whole-blood P=6.30x10^{-26}). These two meta-exons were deemed to be

candidate-causal for SLE across both cell-types. We believe the meta-exons in the 3′ end of

NADSYN1 that are candidate-causal in LCLs (**Fig. 3**), are not detected in whole-blood may

be because of the smaller sample size or the mixed cell-type composition of the whole blood

cohort. This novel instance of specific exon expression disruption found in a primary cell-

type at *NADSYN1* may help to resolve the functional consequence of this locus.

Table 7
Candidate-causal eQTLs and eGenes detected using RNA-Seq (exon-level) in whole blood

Risk Locus	GWAS SNP	eGene	Exon ID (chr. start. end)	β	*P*-Value	FDR (q)
3p14.3	rs9311676	PXK	3.58383332.58383449	0.272	1.15E-07	2.03E-05
		RPP14	3.58303167.58305816	0.207	6.18E-05	5.45E-03
			3.58303167.58305920	0.211	4.52E-05	4.07E-03
4q24	rs10028805	BANK1	4.102750965.102751363	0.204	8.41E-05	7.11E-03
8p23.1	rs2736340	BLK	8.11400733.11400856	0.297	6.20E-09	1.40E-06
			8.11403240.11403612	0.268	1.74E-07	2.94E-05
			8.11405541.11405634	0.267	2.01E-07	3.26E-05
			8.11412252.11412398	0.282	3.72E-08	6.86E-06
			8.11412841.11412993	0.301	3.80E-09	9.63E-07
			8.11414166.11414346	0.290	1.50E-08	3.04E-06
			8.11415471.11415547	0.264	2.75E-07	4.29E-05
			8.11417842.11418961	0.373	1.26E-13	4.65E-11
			8.11420488.11420619	0.334	4.58E-11	1.33E-08
			8.11421412.11422113	0.344	1.16E-11	3.62E-09
		FAM167A	8.11278972.11282145	0.523	2.96E-27	6.00E-24
		RP11-148O21.2	8.11417293.11417529	0.222	1.79E-05	1.96E-03
			8.11415975.11416256	0.203	9.13E-05	7.56E-03
		RP11-148O21.4	8.11413760.11414170	0.220	2.09E-05	2.23E-03
11p15.5	rs12802200	ANO9	11.419582.420860	0.180	5.20E-04	3.35E-02
		HRAS	11.532242.532755	0.179	5.75E-04	3.64E-02
		IRF7	11.613785.614534	0.263	3.02E-07	4.54E-05
			11.614783.615728	0.216	2.89E-05	2.86E-03
		RNH1	11.504824.505881	0.238	3.84E-06	5.02E-04
		TALDO1	11.758950.759057	0.186	3.44E-04	2.36E-02
11q13.4	rs3794060	NADSYN1	11.71185441.71186668	0.526	1.28E-27	5.19E-24
			11.71187079.71188484	0.512	6.30E-26	8.52E-23
		RP11-660L16.2	11.71159720.71159931	0.437	1.48E-18	8.58E-16
			11.71162736.71163203	0.508	1.54E-25	1.56E-22
15q24.2	rs2289583	CSK	15.75094672.75095539	0.204	8.17E-05	7.05E-03
		MPI	15.75189853.75191798	0.252	1.02E-06	1.43E-04
		SCAMP2	15.75142855.75143014	0.188	2.83E-04	2.05E-02
		ULK	15.75128459.75129585	0.259	4.62E-07	6.69E-05
			15.75132839.75132982	0.189	2.58E-04	1.94E-02
			15.75134621.75134761	0.194	1.85E-04	1.42E-02
22q11.21	rs7444	UBE2L3	22.21975804.21978323	0.505	3.68E-25	2.99E-22

GWAS SNPs deemed to be candidate causal eQTLs using RNA-Seq expression data profiled from 384 individuals of the TwinsUK cohort in whole blood at exon-level resolution. In total, 3,793 exons were tested against, corresponding to 654 genes.

Splice-junction quantification reveals asQTLs and additional candidate eGenes

We extended our investigation to determine whether the GWAS SNPs (**Table 1**) had a direct

influence on the alternative-splicing of transcripts (alternative-splicing quantitative trait loci;

asQTL), and whether expression quantification at this resolution would reveal any additional

candidate-genes or potential functional mechanisms. We undertook *cis*-asQTL analysis

within a +/-1Mb window around each GWAS SNP against 33,039 splice-junction

quantifications, corresponding to 817 genes, in the Geuvadis cohort (**Table 2**). We identified

nine asQTLs significantly associated with 62 splice-junctions, corresponding to 10 eGenes

(**S5 Table**). After testing for a shared causal variant between the GWAS and asQTL signal,

six SLE candidate-causal asQTLs (26 splice-junctions) for seven eGenes remained (**Table**

8). Four eGenes (*TCF7*, *SKP1*, *BLK*, and *NADSYN1*) had been previously associated

through either gene-level or exon-level eQTL mapping using the TwinsUK cohort. The

remaining three candidate-causal eGenes detected using asQTL mapping (*IKZF2*, *WDFY4*,

and *IRF5*), as well as the novel causal mechanism involving *NADSYN1*, are described

below.

IKZF2

IKZF2 (Ikaros Family Zinc Finger 2) is novel SLE candidate-causal eGene detected only by

asQTL analysis. The GWAS association signal around the 3′ end of *IKZF2* tagged by risk

variant rs3768792[G] drove an increase in the fraction of splicing between exon 6A and exon

6B (P=3.8x10^{-05}); a bridge that is unique to the truncated isoform (ENST00000413091, 239

amino-acids) of *IKZF2* (**Fig. 5A**). Interestingly, this isoform possesses a premature

termination codon found on exon 6B that is not found on the canonical isoform

(ENST00000457361, 526 amino-acids) as in this isoform, exon 6A is spliced to exon 7 (**Fig.**

5B). This effect results in the premature truncation of the full-length protein and the

subsequent loss of the two zinc-finger dimerization domains found on exon 8 (**Fig. 5B**).

IKZF2 is a transcription factor thought to play a key role in T-reg stabilisation in the presence

of inflammatory responses (74). Since the Ikaros transcription factor family primarily regulate

gene expression through homo-/hetero-dimerization and DNA binding/protein-protein

interactions, the rs3768792[G] dependent asQTL effect on exon 6A to 6B resulting in less

functional *IKZF2* could be highly deleterious. *IKZF2* is known to regulate T-reg associated

genes, including *IL-2* and *FoxP3* (75, 76), therefore a decrease in the amount of DNA

binding *IKZF2* may result in loss of T-reg stability and a decrease of suppressive capacity

with consequential autoimmune sequelae. Interestingly, we identified an additional asQTL

variant (rs2291241) in near-perfect LD with the rs3768792 GWAS variant (r^2:0.99), located 9

bp upstream of exon 6B in truncated isoform ENST00000413091 (**S5 Table, Fig. 5C**). This

second asQTL, located within the polypyrimidine tract in the exon 6A/exon 6B intron, is a

highly plausible driving variant and may act through promotion of the described splicing

event (**Fig. 5C**).

WDFY4

We also discovered a novel putative SLE-associated splicing mechanism involving *WDFY4*

(WDFY Family Member 4), a gene belonging to a family thought to function as master

conductors of aggregate clearance by autophagy (77). Risk variant rs2263052[G] or

correlated SNPs (**Fig. 6A**) greatly increased the fraction of link-counts between exon 34A

and exon 34B (P=3.3x10^{-19}) which are unique to the truncated isoform ENST00000374161

(**Fig. 6B**). This truncated isoform (552 amino-acids) lacks the two WD40 domains found in

the full length isoform (ENST00000325239, 3184 amino-acids) that are essential to

enzymatic activity (77). There is a consequential decrease in the fraction of link-counts

between exon 34A and exon 35 (P=3.0x10^{-06}) that are unique to the canonical isoform of

WDFY4 (**Fig. 6B**). Interestingly, a known missense variant found in exon 31 of *WDFY4* (**Fig.**

6B), rs7097397 (Arg1816Gln), in strong LD (r^2:0.7) with rs2263052, has also been

implicated in SLE through GWAS (78). SIFT and PolyPhen predict the amino-acid

substitution to be tolerated (0.38) or benign (0.11) respectively (79); thereby suggesting the

risk haplotype may harbour two functional mechanisms influencing *WDFY4* (amino-acid

change and upregulation of a shorter isoform) that are both involved in SLE pathogenesis.

IRF5

The GWAS SNP and known asQTL, rs3757387, causes differential promoter usage of *IRF5*

(Interferon regulatory factor 5); a molecular mechanism that has previously been reported in

predisposition of SLE (80). Alteration of a consensus splice-site causes upregulation of a

shorter isoform of *IRF5* which subsequently leads to erroneous activation of the type-1 IFN-

pathway and pro-inflammatory cytokines (81). We replicated this known effect by observing

an increased fraction of splicing of the shorter isoform of *IRF5*, ENST00000489702, with

respect to the risk allele rs3757387[C] (**Table 8**). The risk allele increases splicing from the

first exon to the penultimate exon of ENST00000489702 (P=2.2x10^{-08}); and from the first

exon to the final exon of ENST00000489702 (P=5.9 x10^{-07}).

NADSYN1

Finally, using splice-junction quantification, we were able to pinpoint the specific transcript of

NADSYN1 that drives the exon-level association previously described (**Table 5**, **Fig. 3**). The

GWAS SNP rs3794060 leads to substantial upregulation of the meta-exon 10 to meta-exon

12 splice-site (P=8.0x10^{-12}) which is unique the ENST00000528509 transcript of *NADSYN1*

(**Table 8**, **S14 Fig.**). As a consequence of this splicing event, it appears the meta-exon 11 to

meta-exon 12 splice-site is highly reduced (P=2.1x10^{-14}) with reference to the risk allele [C].

Meta-exons 11 and 12 were implicated in exon-level analysis but a specific transcript could

not be isolated as the gene annotation was not collapsed to the granularity used in the

asQTL analysis. Interestingly, transcript ENST00000528509 is translated to a 294 amino

acid residue protein where the canonical transcript of *NADSYN1*, ENST00000319023, is 706

amino acids. The shorter protein lacks the NAD(+) Synthetase domain (located in positions

339-602aa) found in the canonical protein (Pfam: PF02540); thus implicating loss of this

domain as a potential causal mechanism.

Table 8
Candidate-causal asQTLs and associated eGenes detected using RNA-Seq (Splice-junction level)

Risk Locus	GWAS SNP	eGene	Spice Junction (chr: donor-acceptor exon)	β	*P*-Value	FDR (q)
2q34	rs3768792	IKZF2	2:213872084.213872808-213886368.213886444	-0.04	3.64E-09	5.51E-06
			2:213878515.213880002-213886368.213886444	-0.03	1.30E-06	1.07E-03
			2:213881647.213881768-213886368.213886444	-0.04	1.48E-07	1.41E-04
			2:213886717.213886854-213886368.213886444	0.07	3.80E-05	1.93E-02
			2:213914437.213914604-213886368.213886444	-0.04	4.55E-06	3.15E-03
5q31.1	rs7726414	SKP1	5:133541305.133541822-133561451.133561762	0.02	1.42E-06	1.16E-03
			5:133541645.133541822-133561451.133561762	-0.03	1.28E-06	1.07E-03
		TCF7	5:133478412.133478791-133474642.133474729	0.01	1.90E-06	1.49E-03
7q32.1	rs3757387	IRF5	7:128585899.128586088-128577666.128577888	0.02	2.18E-08	2.49E-05
			7:128586555.128586616-128577666.128577888	0.05	9.51E-07	8.13E-04
8p23.1	rs2736340	BLK	8:11351510.11352100-11403561.11403612	0.10	4.16E-05	2.08E-02
			8:11400733.11400856-11403561.11403612	-0.04	6.24E-05	2.96E-02
			8:11407668.11407771-11403240.11403612	0.02	1.74E-09	3.14E-06
			8:11412252.11412398-11403240.11403612	0.02	2.77E-09	4.61E-06
			8:11414167.11414297-11407503.11407771	0.01	1.36E-08	1.64E-05
			8:11415471.11415547-11407503.11407771	0.01	2.52E-06	1.90E-03
			8:11415471.11415547-11414166.11414346	-0.04	3.24E-06	2.32E-03
			8:11420488.11420619-11414166.11414346	0.02	1.09E-13	7.06E-10
10q11.23	rs2663052	WDFY4	10:50031227.50032214-50030425.50030582	-0.02	3.31E-19	2.15E-14
			10:50034716.50034954-50030425.50030582	0.02	3.01E-06	2.17E-03
11q13.4	rs3794060	NADSYN1	11:71171087.71171259-71171768.71171993	-0.06	2.95E-05	1.55E-02
			11:71184324.71184412-71187247.71188271	0.17	8.00E-12	2.08E-08
			11:71184615.71184732-71187079.71188484	0.06	1.02E-04	4.51E-02
			11:71185441.71186668-71187247.71188271	-0.19	2.13E-14	1.73E-10
			11:71185441.71185572-71189441.71189515	0.02	1.06E-05	6.71E-03
			11:71185441.71186668-71187079.71188484	-0.06	1.02E-04	4.51E-02

GWAS SNPs deemed to be candidate causal asQTLs using RNA-Seq expression data profiled from 373 individuals of the Geuvadis cohort in lymphoblastoid cell lines at splice-junction resolution. 33,039 splice-junctions, corresponding to 817 genes were tested against in cis to the 39 GWAS SNPs.

Discussion

Detailed characterization of the functional effects of human regulatory genetic variation associated with complex-disease is paramount to our understanding of molecular aetiology and poised to make significant contributions to translational medicine (82). Use of eQTL mapping studies to interpret GWAS findings have proved fundamental in our progression towards this goal - through prioritization of candidate genes, refinement of causal variants, and illumination of mechanistic relationships between disease-associated genetic variants and gene expression (82, 83). However, there is often a disparity between disease-associated genetic variation and phenotypic alteration, which historically may be due to the use of microarray-based technologies to profile genome-wide gene expression. With the advent of RNA-Seq, we can achieve more accurate quantification of the mRNA output of genes, individual exons, and isoform abundance, as well as unannotated and non-coding transcripts. Detection of splicing variants at susceptibility loci using RNA-Seq has the potential to uncover the role of specific isoforms implicated in disease risk, which are likely to have remained concealed by microarray, as a largely independent subset of variants control alternative splicing of isoforms compared to overall gene abundance (35).

Our two major motivations for this study were firstly to directly compare the ability of RNA-Seq with microarrays to detect candidate-causal eQTLs and their associated eGenes from GWAS data, and then to assess each platform's effectiveness in explaining the potential causal genes and mechanisms implicated by SLE risk alleles. A previous investigation of the same SLE risk alleles used in this study with eQTL data from LCL microarray datasets revealed that only 13 of the 39 risk alleles were eQTLs for a total of 15 eGenes (**Table S6**). Therefore, in an attempt to increase our understanding of the extent to which eQTLs can explain the functional consequences of our risk alleles and to achieve both of our major aims for this investigation, we set up an analytical pipeline to compare the results of eQTL annotation for 39 SLE susceptibility loci with *cis*-eQTL data from both microarray and RNA-

Seq experiments from the TwinsUK cohort in LCLs. We incorporated steps to minimize

false-positive associations through conditional and colocalisation analysis, as approximately

34% of all genes have a second independently associated *cis*-eQTL for any of their exons

when conditioning on the best *cis*-eQTL (35). The analytical pipeline we present will be

applicable to the functional annotation of susceptibility loci from a wide range of human

diseases.

Fig. 7 summarises the data generated in this manuscript illustrating which of the GWAS

SNPs show evidence of a candidate-causal eQTL association across the quantification

types. Our data analysis revealed that RNA-Seq is a more powerful in the identification of

candidate-causal eQTLs and their accompanying eGenes than corresponding microarray

datasets (**S6 Table**). Many of the published SLE candidate eGenes and associated

mechanisms were well-replicated by performing *cis*-eQTL analysis of RNA-Seq datasets at

various resolutions for each of the GWAS variants. These eGenes included, among others,

the effect of risk variants at *BLK* and *FAM167A* (**S9 Fig.**), *MIR146A* (**Fig. 4A**), *BANK1* (**S12

Fig.**) and *IRF5*. Microarray studies were unable to detect the novel SLE candidate-causal

eGenes identified from RNA-Seq data including *NADSYN1, TCF7*, *SKP1*, *WDFY4*, *IKZF2*,

and the non-coding RNA genes: *RP11-148O21.2, RP11-148O21.4*, and *RP11-660L16.2*

(**Fig. 2**).

Our results also demonstrate that RNA-Seq analysis is much better than microarrays in

identifying multiple eGenes for a single SNP (that may tag multiple functional variants). An

increased ratio of eQTLs to eGenes (average number of eGenes per candidate-causal

eQTL) was observed using RNA-Seq at exon-level (2.42) compared with gene-level (1.72)

and which were both greater than microarray (1.5) (**S5 Fig.**). The ability of RNA-Seq exon-

level analysis to identify multiple target eGenes for a specific eQTL is supported by recent

observations from capture Hi-C (cHi-C) experiments to functionally annotate chromatin

interactions, such as enhancer–promoter interfaces (84, 85). It has been shown that

chromatin interactions can control transcription in both *cis* and *trans* in a largely sequence-
specific manner, thus it is likely that some GWAS variants may functionally act through the
disruption of chromatin dynamics resulting in perturbation of expression of multiple genes
(84, 86, 87). Specific instances of this type of effect are seen in colorectal cancer risk loci
where the risk locus 11q23 mapped to interactions with genes *C11orf53*, *C11orf92* and
C11orf93, and separately, the risk SNP rs6983267 within 8q24 disrupts a chromatin
regulatory network involving interactions between three genes *CCAT2*, *CCAT1* and *MYC*
(84). Our results support this notion of multiple perturbed genes at a single susceptibility
locus. At 1q32.1, for example, rs3024505 was found to be associated with three plausible
candidate-causal eGenes: *IL10*, *IL24*, and *FCAMR* (located 1 kb, 130 kb, and 191 kb away
from rs3024505 respectively). These chromatin capture data also support the argument of
using RNA-Seq and extending the *cis*-eQTL distance (typically +/-0.25–1Mb) to a larger
region (+/-5Mb) around the associated SNP to identify effects caused by chromatin
interactions over a larger distance than commonly designated as *trans* from eQTL-type
analyses (88). Integration of eQTL data with epigenetic regulation (promoter methylation,
histone modification and expression of non-coding RNA) will allow the identification of the
potential mechanism of action (disruption of epigenetic landscape) and disease biology of
associated variants.

We have also demonstrated the power of RNA-Seq compared with microarray in the
discovery of alternative-splicing events. This is of significant importance as approximately
80% of all human genes undergo alternative splicing and it is estimated that 20-30% of
disease-associated mutations modify the configuration of expressed isoforms (89, 90). A
recent study has concluded that regulatory variants controlling gene splicing are major
contributors to complex traits (91). Two examples of this taken from our analyses are the risk
alleles at the *IKZF2* and *WDFY4* loci which drive up-regulation of short isoforms. Neither of
these effects were captured by the microarray probes that targeted the 3′-UTR of the
canonical longer isoforms. At *WDFY4*, the splicing variant rs2263052[G] is in strong LD with

an (Arg1816Gln) missense variant rs7097397[G], which we showed to be associated with

SLE (**Fig. 6**). These two potentially causal signals may reinforce each other. The novel

association identified in our group's recent GWAS study (63) at the *IKZF2* locus implicated a

risk haplotype tagged by the risk allele of rs3768792[G]. We identified, by RNA-Seq splice-

junction quantification exclusively, that variation at the rs3768792[G] risk allele led to

increased production of a shorter isoform of *IKZF2* (**Fig. 5**). Interestingly, other members of

this gene family, *IKZF1* and *IKZF3*, are also associated with SLE (63). An associated variant

in the 3'-UTR of *IKZF1* has been associated with increased expression of genes of the type

1 IFN pathway and decreased expression of complement genes, both mediated in *trans*

(64). The functional effect of the *IKZF3* common variant is less well documented, although

IKZF3 knockout mice develop spontaneous autoantibodies and B-cell lymphoma (92). The

IKZF2 knockout mouse has not been characterised for immune-deficient phenotypes. We

hypothesize that upregulation of the shorter isoform of *IKZF2* caused by rs3768792[G],

which lacks the dimerization domain, reduces translocation of the protein into the nucleus

and regulation of transcription of target genes. We believe this aberrant mechanism may

result in loss of T-reg stability. The asQTL discoveries described in this manuscript are

examples of how RNA-Seq can suggest a potential causal mechanism that can be easily

validated experimentally.

Our study also demonstrates that RNA-Seq is able to identify disease-relevant non-coding

RNAs. These type of transcripts have long been known to be of relevance to human

disease, however their detection and functional importance may have been under-estimated.

We identified three novel non-coding anti-sense RNAs using RNA-Seq: two regulated by

rs2736340 (*RP11-148O21.2* and *RP11-148O21.4*) (**Fig. 4B**), and one regulated by

rs3794060 (*RP11-660L16.2*) (**Fig. 4C**). These findings were replicated in whole blood (**Table**

7).

The data we present in this manuscript demonstrate that a comprehensive integrated

approach for eQTL analysis should be undertaken at gene-, exon- and exon-junction level

quantification. Excluding one or more levels of analysis will mean that eQTLs and/or eGenes

may be missed. This is illustrated by a number of examples in **Fig. 7**. At some loci, there

was an exon-level effect, which was not observed at the gene-level. For example, the risk

allele rs3024505[A] was an eQTL for *IL10*, *IL24*, and *FCAMR* at exon-level resolution only

(**Table 5**), but none of these eGenes were deemed to be significant at gene-level. This

suggests that the exon-level effect is more targeted than gene-level quantification, where

multiple different effects across the gene may dilute out the signal. However, some eGenes

exhibit a probable whole gene-level effect (*UBE2L3*, *BLK*, and *FAM167A)* as every

expressed exon showed a candidate-causal association for the respective risk alleles.

Hierarchical clustering tests could be designed to distinguish between these genuine gene-

level and exon-level effects, such as at *NADSYN1* or *BANK1*, where the gene-level effect is

likely to be driven by only a subset of exons (**Fig. 3**, **S12 Fig.**). There were occurrences

where significant candidate-causal eGenes were detected but there was no effect at the

exon-level. At *TCF7*, variants in low LD with rs7726414 exhibited significant exon-level *cis*-

eQTLs (**S3 Table**) and were deemed to be independent of the disease-association.

However, gene-level analysis revealed that the risk rs7726414 variant was candidate-causal

for total *TCF7* expression (**S2 Table**). These results emphasise that for any given eGene

there may be multiple genetic effects at different resolutions of quantification.

We understand the limitation of LCLs for transcript profiling studies. There is an inherent

limitation analysing LCL expression datasets, such as those available for our study, because

although LCLs are a good surrogate model for primary B-cells, the effect of EBV

transformation is likely to disrupt their underlying epigenetic and transcriptomic background.

The percentage of asQTLs in LCLs will exhibit significantly less replication in primary cell

types because of cell-type variability in the genetic control of isoform usage, approximately

70% of *cis*-eQTL detected in LCLs can be replicated in a primary tissue type (33). The

significance of alternative-splicing in genomic medicine will become better understood once

large RNA-Seq based eQTL cohorts emerge across a multitude of disease-relevant cell-

types. A gold standard of candidate-causal eQTL mapping strategies using RNA-Seq across

datasets using an explicit set of quantification types (gene-, exon-, splice-junction, isoform)

and analytical pipelines, will accelerate this process.

In summary, we have demonstrated the effectiveness of eQTL analysis using RNA-Seq by

increasing the numbers of candidate eGenes regulated by SLE associated alleles (**Fig. 2**,

Fig. 7, and **S6 Table**). We have shown that the power of RNA-Seq in eQTL annotation of

GWAS loci lies not only the assessment of the variants regulating the expression of

candidate genes, but also in uncovering putative molecular mechanisms, which allow for

more refined targeted follow-up studies to assess the phenotypic consequence of the

disease-associated variant. These studies could include knocking down target gene or

introducing recombinant vectors *in vitro* to overexpress target genes to evaluate phenotypic

consequence of expression changes of specific loci. Site directed mutagenesis could be

used to introduce candidate causal splice-sites and over-express target isoforms. The

CRISPR/Cas9 system for targeted genome editing presents an exciting opportunity for

eQTL/RNA-Seq targeted follow-up studies and the investigation of the effect that specific

variants have on the expression profile across different cell types.

Materials and Methods

Selection of SLE associated SNPs

SLE associated SNPs were taken from our recent 2015 publication (63). The study comprised a primary GWAS, with validation through meta-analysis and replication study in an external cohort (7,219 cases, 15,991 controls in total). The independently-associated susceptibility loci taken forward for this investigation were those that passed either genome-wide significance ($P< 5x10^{-08}$) in the primary GWAS or meta-analysis and/or those that reached significance in the replication study (False Discovery Rate, q<0.01). We defined the 'GWAS SNP' at each locus as either being the SNP with the lowest P-value post meta-analysis or the SNP with the greatest evidence of a missense effect as defined by a Bayes Factor. We omitted non-autosomal associations and those within the Major Histocompatibility Complex (MHC), and SNPs with a MAF <0.05. In total, 39 GWAS SNPs were taken forward (**Table 1**).

TwinsUK cohort eQTL datasets

Expression profiling by microarray (9) and RNA-Seq (39) of individuals from the UK Adult Twin Registry (TwinsUK) was carried out in two separate studies on the MuTHER (Multiple Tissue Human Expression Resource) cohort (**Table 2**). The MuTHER cohort is composed of 856 healthy female individuals of European descent aged between 37-85 years. We considered expression quantification data from both resting LCLs and whole blood. Profiling by microarray was performed using the Illumina Human HT-12 V3 BeadChips. For RNA-Seq, samples were sequenced using the Illumina HiSeq2000 and the 49-bp paired-end reads mapped with BWA v0.5.9 to the GRCh37 reference genome. Exons ('meta-exons' created by merging all overlapping exonic portions of a gene into non-redundant units) were quantified using read-counts against the GENCODE v10 annotation; with gene quantification defined as the sum of all exon quantifications belonging to the same gene. Full quality control and normalization procedures are described in the respective articles. Data from

each of the TwinsUK eQTL studies (**Table 2**) were provided in different formats. In each

instance it was necessary to generate summary *cis*-eQTL statistics per GWAS SNP (SNP,

expression-unit, β, standard error of β, and *P*-value of association) for integration analysis.

Per quantification type (microarray, RNA-Seq gene-level, and exon-level), each GWAS SNP

was subject to *cis*-eQTL analysis against all expression-units within +/-1Mb using no *P*-value

threshold. If the GWAS SNP was not found in an eQTL dataset, the most highly correlated,

closest tag SNP with $r^2 \geq 0.7$, common to all datasets, was used as its proxy (**Table 1**).

Adjustment for multiple testing of *cis*-eQTL results per quantification type were undertaken

using FDR with q <0.05 deemed significant.

Microarray cis-eQTL mapping

We used the Genevar (GENe Expression VARiation) portal to generate summary-level *cis*-

eQTL results (50). We ran the association between normalized expression data of the 777

individuals and each GWAS SNP implementing the external algorithm option (two-step

mixed model–based score test). In total 768 probes (559) genes, were tested.

RNA-Seq (gene-level) cis-eQTL mapping

RNA-Seq gene-level quantification was provided as residualized read-counts (effect of family

structure and other covariates regressed out). We had full genetic data for 683 individuals

and performed the analysis of each GWAS SNP against the transformed residuals using the

linear-model function within the MatrixeQTL R package (93). 520 genes were tested against

in *cis*.

RNA-Seq (exon-level) cis-eQTL mapping

P-values from the association of all SNPs against exon-level quantifications for 765

individuals using linear-regression were provided. We generated the t-statistic using the

lower-tail quantile function t-distribution function in R with 763 degrees of freedom. The

standard error and β were derived from the t-statistic. We then extracted the summary *cis*-

eQTL results for each GWAS SNP. 4,786 exons, corresponding to 716 genes for testing.

SLE candidate-causal *cis*-eQTL classification

Conditional analysis

We used the COJO (conditional and joint genome-wide association analysis) function of the

GCTA (Genome-wide Complex Trait Analysis) application to determine whether the GWAS

SNP had an independent effect on expression from that of the best *cis*-eQTL (55). For each

significant association (q<0.05), we re-performed the analysis using all SNPs within +/-1Mb

of the expression-unit in hand. We used the available genotype information of the 683

TwinsUK individuals to extract allele coding along with the MAF, and integrated this with the

cis-eQTL summary data. We discarded SNPs with: MAF < 0.05, imputation call-rates < 0.8,

and HWE $P < 1 \times 10^{-04}$. We used these individuals as the reference panel to calculate local

pairwise linkage disequilibrium (LD) between variants. Per significant association, all *cis*-

eQTLs were conditioned on by the best *cis*-eQTL. We then extracted the conditional *P*-value

of the GWAS SNP and considered associations to be independent to the best *cis*-eQTL if

$P_{cond} < 0.05$.

Colocalisation Analysis

We employed the 'coloc' Bayesian statistical method using summary data implemented in R

to test for colocalisation between *cis*-eQTL and disease causal variants derived from the

GWAS (56). The method makes the assumption of there being a single causal variant for

each trait (disease association and gene-expression from two separate studies) per locus

and calculates the posterior probabilities under five different causal variant hypotheses:

association with neither trait (H0), association with one trait but not the other (H1, H2),

association with both traits but from independent signals, and association with both traits

with a shared causal signal (H4). We extracted the necessary SNP statistics for the disease-

associated regions from our own GWAS and applied the same SNP filters used in the

conditional analysis. We tested for colocalisation between the GWAS summary data and *cis*-

eQTL data for each significant association within a +/-1Mb window of the GWAS SNP. We

assigned the prior probabilities, p1 and p2 (SNP is associated with GWAS and gene

expression respectively), as 1 x 10^{-04} i.e. 1 in 10,000 SNPs are causal to either trait, with p12

(SNP is associated with both traits) as 1x10^{-06} or 1 in 100 SNPs associated with one trait are

also associated with the other. For each *cis*-eQTL association colocalisation test, if the

posterior probability PP3 (two distinct causal variants, one for each trait) is greater than PP4

(single causal variant common to both traits), then greater posterior support is given to the

hypothesis that independent causal variants exist in both traits and thus the eQTL is unlikely

to be attributed to SLE genetic association.

Definition of SLE candidate-causal cis-eQTL and eGene

We defined a GWAS SNP as an SLE candidate-causal *cis*-eQTL if it met the following

criteria: significant post-multiple testing adjustment (q < 0.05), not independent to the best

cis-eQTL from conditional analysis (P_{cond} > 0.05), and supporting evidence of a shared

causal variant between gene expression and the primary GWAS signal based on

colocalisation (PP3 < PP4). The gene whose expression is modulated by the candidate-

causal eQTL is defined as an SLE candidate-causal eGene (**Fig. 1**).

Validation of LCL SLE candidate-causal cis-eQTLs in whole blood

Cis-eQTL summary data from whole blood at RNA-Seq exon-level were made available for

384 individuals of the 856 TwinsUK cohort individuals (**Table 2**). Expression profiling and

genotyping were identical to that as described for LCLs. We applied the same methodology

to this dataset to generate full *cis*-eQTL summary statistics, perform conditional and

colocalisation analysis, and classify SLE candidate-causal eQTLs and associated eGenes

(**Fig. 1**). In total, 3,793 exons were tested against, corresponding to 654 genes.

Geuvadis SLE candidate-causal *cis*-asQTL analysis

We investigated SLE disease-associated alternative splicing QTLs (asQTLs) using

European samples from the raw alignment files of the Geuvadis (35) 1000 Genomes RNA-

Seq project profiled in LCLs (**Table 2**). Genotype data and read-alignments were

downloaded from ArrayExpress for the 373 Europeans (comprising 91 CEU, 95 FIN, 94

GBR, and 93 TSI). We performed PCA on chromosome 20 using the R/Bioconductor

package SNPRelate (94) and decided to include the first three principle components as

covariates in the eQTL model as well as the binary imputation status (mixture of Phase 1

and Phase 2 imputed individuals). We removed SNPs with MAF<0.05, imputation call-

rates<0.8, and HWE P<1×10^{-04}. We removed non-uniquely mapped, non-properly paired

reads, and reads with more than eight mismatches for read and mate using Samtools(95).

We used the Altrans (96) method against GENCODE v10 to generate relative quantifications

(link-counts) of splicing events; which in brief, utilizes split and paired-end reads to count

links between exon-boundaries, which themselves are created by flattening the annotation

into unique non-redundant exon-groups. Following PCA of the link-counts, we decided to

normalize all link-counts with the first 10 principle components then removed exon-

boundaries with zero links in more than 10% of individuals. Link-counts were converted to

link-fractions (coverage of the link over the sum of the coverage of all the links that the first

exon makes) and merged in both 5′-3′ and 3′-5′ directions. Per GWAS SNP we performed

cis-eQTL analysis against the normalized link-fractions in MatrixeQTL with a linear-model

(93). 33,039 link-fractions were tested against corresponding to 817 genes in total. After

FDR multiple-testing adjustment we considered associations with q<0.05 as significant. As

full genetic and expression data were available, we decided to use the Regulatory Trait

Concordance (RTC) method to assess the likelihood of a shared functional variant between

the GWAS SNP and the asQTL signal (48). For each significant asQTL association we

extracted the residuals of the linear-regression of the best *cis*-eQTL against normalized link-

fractions and re-performed the analysis using all SNPs within the defined hotspot interval

against this pseudo-phenotype. The RTC score was defined as (N_{SNPs}-$Rank_{GWAS \: SNP}$)/N_{SNPs}

where N_{SNPs} is the number of SNPs in the interval, and $Rank_{GWAS \: SNP}$ is the rank of the

GWAS SNP association *P*-value against all other SNPs in the interval. We classified an SLE

candidate-causal *cis*-asQTL as a GWAS SNP with a significant association (q < 0.05) with

link-fraction quantification and an RTC score > 0.9.

Statistical analysis and data visualisation

We performed statistical analysis, graphics and data handling in R version 3.2.0 and ggplot2.

Genetic plots were generated using LocusZoom v1.1 (97). Karyotype diagrams were

modified from Ensembl (98). GWAS association plots and gene annotation graphic

visualisations were created using the UCSC Genome Browser(99).

Acknowledgements

DSCG and TJV were awarded an Arthritis Research UK PhD Studentship (for CO) (20332) "SLE pathogenesis: the molecular basis for multiple association signals in *IKZF2*" to undertake the analyses described in this manuscript. DSCG, DLM and TJV were awarded an Arthritis Research UK Project Grant (20265) "Post GWAS: A Functional Genomic Analysis of the SLE Susceptibility Gene *IKZF1*", which supported AC. TJV received funds from China Scholarship Council, number 201406380127 for LC. ALR was supported by MRC project grant L002604/1 "Functional Genomics of SLE: A Trans-ancestral approach" and an ARUK project grant (20580) "Targeted DNA sequencing in Indian and European samples contributes to causal allele identification for lupus". KSS was awarded funding by the MRC (MR/L01999X/1 and MR/M004422/1).

The Geuvadis 1000 Genomes RNA-Seq data was downloaded from the EBI ArrayExpress Portal (accessions E-GEUV-1). The RNA-Seq data on the TwinsUK samples in both LCLs and whole blood was made available through the EUROBATS project (EGAS00001000805 from the European Genome-Phenome Archive). The expression microarray data on the TwinsUK samples was downloaded from the EBI ArrayExpress Portal (E-TABM_1140).

The TwinsUK study was funded by the Wellcome Trust; European Community's Seventh Framework Programme (FP7/2007-2013). The study also receives support from the National Institute for Health Research (NIHR)- funded BioResource, Clinical Research Facility and Biomedical Research Centre based at Guy's and St Thomas' NHS Foundation Trust in partnership with King's College London. SNP Genotyping was performed by The Wellcome Trust Sanger Institute and National Eye Institute via NIH/CIDR.

723   **Conflict of Interest Statement**

724   The authors declare that there are no conflicts of interest.

References

References

1. Hindorff,L.A., Sethupathy,P., Junkins,H.A., Ramos,E.M., Mehta,J.P., Collins,F.S. and

Manolio,T.A. (2009) Potential etiologic and functional implications of genome-wide

association loci for human diseases and traits. *Proc. Natl. Acad. Sci. U. S. A.*, **106**,

9362–7.

2. Maurano,M.T., Humbert,R., Rynes,E., Thurman,R.E., Haugen,E., Wang,H.,

Reynolds,A.P., Sandstrom,R., Qu,H., Brody,J., *et al.* (2012) Systematic Localization of

Common Disease-Associated Variation in Regulatory DNA. *Science (80-.).*, **337**,

1190–1195.

3. The ENCODE Project Consortium, Dunham,I., Kundaje,A., Aldred,S.F., Collins,P.J.,

Davis,C. a, Doyle,F., Epstein,C.B., Frietze,S., Harrow,J., *et al.* (2012) An integrated

encyclopedia of DNA elements in the human genome. *Nature*, **489**, 57–74.

4. Kundaje,A., Meuleman,W., Ernst,J., Bilenky,M., Yen,A., Heravi-Moussavi,A.,

Kheradpour,P., Zhang,Z., Wang,J., Ziller,M.J., *et al.* (2015) Integrative analysis of 111

reference human epigenomes. *Nature*, **518**, 317–330.

5. Forrest,A.R.R., Kawaji,H., Rehli,M., Kenneth Baillie,J., de Hoon,M.J.L., Haberle,V.,

Lassmann,T., Kulakovskiy,I. V., Lizio,M., Itoh,M., *et al.* (2014) A promoter-level

mammalian expression atlas. *Nature*, **507**, 462–470.

6. Nicolae,D.L., Gamazon,E., Zhang,W., Duan,S., Dolan,M.E. and Cox,N.J. (2010) Trait-

associated SNPs are more likely to be eQTLs: annotation to enhance discovery from

GWAS. *PLoS Genet.*, **6**, e1000888.

7. Stranger,B.E., Montgomery,S.B., Dimas,A.S., Parts,L., Stegle,O., Ingle,C.E.,

Sekowska,M., Smith,G.D., Evans,D., Gutierrez-Arcelus,M., *et al.* (2012) Patterns of cis

regulatory variation in diverse human populations. *PLoS Genet.*, **8**, e1002639.

8. Spielman,R.S., Bastone,L.A., Burdick,J.T., Morley,M., Ewens,W.J. and Cheung,V.G.

(2007) Common genetic variants account for differences in gene expression among

ethnic groups. *Nat. Genet.*, **39**, 226–231.

9. Grundberg,E., Small,K.S., Hedman,Å.K., Nica,A.C., Buil,A., Keildson,S., Bell,J.T.,

Yang,T.-P., Meduri,E., Barrett,A., *et al.* (2012) Mapping cis- and trans-regulatory effects

across multiple tissues in twins. *Nat. Genet.*, **44**, 1084–1089.

10. Fairfax,B.P., Makino,S., Radhakrishnan,J., Plant,K., Leslie,S., Dilthey,A., Ellis,P.,

Langford,C., Vannberg,F.O. and Knight,J.C. (2012) Genetics of gene expression in

primary immune cells identifies cell type–specific master regulators and roles of HLA

alleles. *Nat. Genet.*, **44**, 502–510.

11. Naranbhai,V., Fairfax,B.P., Makino,S., Humburg,P., Wong,D., Ng,E., Hill,A.V.S. and

Knight,J.C. (2015) Genomic modulators of gene expression in human neutrophils. *Nat.*

Commun., **6**, 7545.

12. Nica,A.C., Parts,L., Glass,D., Nisbet,J., Barrett,A., Sekowska,M., Travers,M., Potter,S.,

Grundberg,E., Small,K., *et al.* (2011) The architecture of gene regulatory variation

across multiple human tissues: the MuTHER study. *PLoS Genet.*, **7**, e1002003.

13. Myers,A.J., Gibbs,J.R., Webster,J. a, Rohrer,K., Zhao,A., Marlowe,L., Kaleem,M.,

Leung,D., Bryden,L., Nath,P., *et al.* (2007) A survey of genetic human cortical gene

expression. *Nat. Genet.*, **39**, 1494–9.

14. Field,J.M., Hazinski,M.F., Sayre,M.R., Chameides,L., Schexnayder,S.M., Hemphill,R.,

Samson,R. a., Kattwinkel,J., Berg,R. a., Bhanji,F., *et al.* (2010) Part 1: Executive

summary: 2010 American Heart Association Guidelines for Cardiopulmonary

Resuscitation and Emergency Cardiovascular Care. *Circulation*, **122**, 640–657.

15. Emilsson,V., Thorleifsson,G., Zhang,B., Leonardson,A.S., Zink,F., Zhu,J., Carlson,S.,

Helgason,A., Walters,G.B., Gunnarsdottir,S., *et al.* (2008) Genetics of gene expression

and its effect on disease. **452**.

16. Schadt,E.E., Molony,C., Chudin,E., Hao,K., Yang,X., Lum,P.Y., Kasarskis,A., Zhang,B.,

Wang,S., Suver,C., *et al.* (2008) Mapping the genetic architecture of gene expression in

human liver. *PLoS Biol.*, **6**, e107.

17. Hao,K., Bossé,Y., Nickle,D.C., Paré,P.D., Postma,D.S., Laviolette,M., Sandford,A.,

Hackett,T.L., Daley,D., Hogg,J.C., *et al.* (2012) Lung eQTLs to help reveal the

molecular underpinnings of asthma. *PLoS Genet.*, **8**, e1003029.

18. Li,Q., Seo,J.H., Stranger,B., McKenna,A., Pe'Er,I., Laframboise,T., Brown,M.,

Tyekucheva,S. and Freedman,M.L. (2013) Integrative eQTL-based analyses reveal the

biology of breast cancer risk loci. *Cell*, **152**, 633–641.

19. Ongen,H., Andersen,C.L., Bramsen,J.B., Oster,B., Rasmussen,M.H., Ferreira,P.G.,

Sandoval,J., Vidal,E., Whiffin,N., Planchon,A., *et al.* (2014) Putative cis-regulatory

drivers in colorectal cancer. *Nature*, **512**, 87–90.

20. Zou,F., Chai,H.S., Younkin,C.S., Allen,M., Crook,J., Pankratz,V.S., Carrasquillo,M.M.,

Rowley,C.N., Nair,A.A., Middha,S., *et al.* (2012) Brain expression genome-wide

association study (eGWAS) identifies human disease-associated variants. *PLoS*

Genet., **8**, e1002707.

21. Albert,F.W. and Kruglyak,L. (2015) The role of regulatory variation in complex traits and

disease. *Nat. Rev. Genet.*, **16**, 197–212.

22. Gibson,G., Powell,J.E. and Marigorta,U.M. (2015) Expression quantitative trait locus

analysis for translational medicine. *Genome Med.*, **7**, 60.

23. Barreiro,L., Tailleux,L., Pai,A., Gicquel,B., Marioni,J.C. and Gilad,Y. (2012) Deciphering

the genetic architecture of variation in the immune response to Mycobacterium

tuberculosis infection. *Proc. Natl. Acad. Sci.*, **109**, 1204–1209.

24. Fairfax,B.P., Humburg,P., Makino,S., Naranbhai,V., Wong,D., Lau,E., Jostins,L.,

Plant,K., Andrews,R., McGee,C., *et al.* (2014) Innate Immune Activity Conditions the

Effect of Regulatory Variants upon Monocyte Gene Expression. *Science (80-.).*, **343**,

1246949–1246949.

25. Marioni,J.C., Mason,C.E., Mane,S.M., Stephens,M. and Gilad,Y. (2008) RNA-seq: An

assessment of technical reproducibility and comparison with gene expression arrays.

10.1101/gr.079558.108.

26. Zhang,X., Johnson,A.D., Hendricks,A.E., Hwang,S.-J., Tanriverdi,K., Ganesh,S.K.,

Smith,N.L., Peyser,P.A., Freedman,J.E. and O'Donnell,C.J. (2014) Genetic

associations with expression for genes implicated in GWAS studies for atherosclerotic

cardiovascular disease and blood phenotypes. *Hum. Mol. Genet.*, **23**, 782–795.

27. Zhang,X., Joehanes,R., Chen,B.H., Huan,T., Ying,S., Munson,P.J., Johnson,A.D.,

Levy,D. and O'Donnell,C.J. (2015) Identification of common genetic variants controlling

transcript isoform variation in human whole blood. *Nat. Genet.*, **47**, 345–352.

28. Zhao,S., Fung-Leung,W.-P., Bittner,A., Ngo,K. and Liu,X. (2014) Comparison of RNA-

Seq and microarray in transcriptome profiling of activated T cells. *PLoS One*, **9**,

e78644.

29. Majewski,J. and Pastinen,T. (2011) The study of eQTL variations by RNA-seq: from

SNPs to phenotypes. *Trends Genet.*, **27**, 72–79.

30. Russo,G., Zegar,C. and Giordano,A. (2003) Advantages and limitations of microarray

technology in human cancer. *Oncogene*, **22**, 6497–6507.

31. Sun,W. and Hu,Y. (2013) eQTL Mapping Using RNA-seq Data. *Stat. Biosci.*, **5**, 198–219.

32. 't Hoen,P. a C., Friedländer,M.R., Almlöf,J., Sammeth,M., Pulyakhina,I., Anvar,S.Y.,

Laros,J.F.J., Buermans,H.P.J., Karlberg,O., Brännvall,M., *et al.* (2013) Reproducibility

of high-throughput mRNA and small RNA sequencing across laboratories. *Nat.*

Biotechnol., **31**, 1015–22.

33. Pickrell,J.K., Marioni,J.C., Pai,A.A., Degner,J.F., Engelhardt,B.E., Nkadori,E.,

Veyrieras,J.-B., Stephens,M., Gilad,Y. and Pritchard,J.K. (2010) Understanding

mechanisms underlying human gene expression variation with RNA sequencing.

Nature, **464**, 768–772.

34. Montgomery,S.B., Sammeth,M., Gutierrez-Arcelus,M., Lach,R.P., Ingle,C., Nisbett,J.,

Guigo,R. and Dermitzakis,E.T. (2010) Transcriptome genetics using second generation

sequencing in a Caucasian population. *Nature*, **464**, 773–777.

35. Lappalainen,T., Sammeth,M., Friedländer,M.R., 't Hoen,P. a C., Monlong,J., Rivas,M. a,

Gonzàlez-Porta,M., Kurbatova,N., Griebel,T., Ferreira,P.G., *et al.* (2013) Transcriptome

and genome sequencing uncovers functional variation in humans. *Nature*, **501**, 506–11.

36. Battle,A., Mostafavi,S., Zhu,X., Potash,J.B., Weissman,M.M., McCormick,C.,

Haudenschild,C.D., Beckman,K.B., Shi,J., Mei,R., *et al.* (2014) Characterizing the

genetic basis of transcriptome diversity through RNA-sequencing of 922 individuals.

Genome Res., **24**, 14–24.

37. Lalonde,E., Ha,K.C.H., Wang,Z., Bemmo,A., Kleinman,C.L., Kwan,T., Pastinen,T. and

Majewski,J. (2011) RNA sequencing reveals the role of splicing polymorphisms in

regulating human gene expression. *Genome Res*, 10.1101/gr.111211.110.21.

38. Kumar,V., Westra,H.-J., Karjalainen,J., Zhernakova,D. V., Esko,T., Hrdlickova,B.,

Almeida,R., Zhernakova,A., Reinmaa,E., Võsa,U., *et al.* (2013) Human Disease-

Associated Genetic Variation Impacts Large Intergenic Non-Coding RNA Expression.

PLoS Genet., **9**, e1003201.

39. Buil,A., Brown,A.A., Lappalainen,T., Viñuela,A., Davies,M.N., Zheng,H.-F.,

Richards,J.B., Glass,D., Small,K.S., Durbin,R., *et al.* (2014) Gene-gene and gene-

environment interactions detected by transcriptome sequence analysis in twins. *Nat.*

Genet., **47**, 88–91.

40. Conde,L., Bracci,P.M., Richardson,R., Montgomery,S.B. and Skibola,C.F. (2013)

Integrating GWAS and Expression Data for Functional Characterization of Disease-

Associated SNPs: An Application to Follicular Lymphoma. *Am. J. Hum. Genet.*, **92**,

126–130.

41. Simon,L.M., Chen,E.S., Edelstein,L.C., Kong,X., Bhatlekar,S., Rigoutsos,I., Bray,P.F.

and Shaw,C.A. (2016) Integrative Multi-omic Analysis of Human Platelet eQTLs

Reveals Alternative Start Site in Mitofusin 2. *Am. J. Hum. Genet.*,

10.1016/j.ajhg.2016.03.007.

42. Li,H., Pouladi,N., Achour,I., Gardeux,V., Li,J., Li,Q., Zhang,H.H., Martinez,F.D., Skip

Garcia,J.G.N. and Lussier,Y.A. (2015) eQTL networks unveil enriched mRNA master

integrators downstream of complex disease-associated SNPs. *J. Biomed. Inform.*, **58**,

226–234.

43. Huang,L., Xu,W., Yan,D., Dai,L. and Shi,X. (2016) Identification of expression

quantitative trait loci of RPTOR for susceptibility to glioma. *Tumor Biol.*, **37**, 2305–2311.

44. Jeong,S., Patel,N., Edlund,C.K., Hartiala,J., Hazelett,D.J., Itakura,T., Wu,P.C.,

Avery,R.L., Davis,J.L., Flynn,H.W., *et al.* (2015) Identification of a novel mucin gene

HCG22 associated with steroid-induced ocular hypertension. *Investig. Ophthalmol. Vis.*

Sci., **56**, 2737–2748.

45. Moffatt,M.F., Kabesch,M., Liang,L., Dixon,A.L., Strachan,D., Heath,S., Depner,M., von

Berg,A., Bufe,A., Rietschel,E., *et al.* (2007) Genetic variants regulating ORMDL3

expression contribute to the risk of childhood asthma. *Nature*, **448**, 470–473.

46. Barrett,J.C., Hansoul,S., Nicolae,D.L., Cho,J.H., Duerr,R.H., Rioux,J.D., Brant,S.R.,

Silverberg,M.S., Taylor,K.D., Barmada,M.M., *et al.* (2008) Genome-wide association

defines more than 30 distinct susceptibility loci for Crohn's disease. *Nat. Genet.*, **40**,

955–962.

47. Li,X., Hastie,A.T., Hawkins,G. a., Moore,W.C., Ampleford,E.J., Milosevic,J., Li,H.,

Busse,W.W., Erzurum,S.C., Kaminski,N., *et al.* (2015) eQTL of bronchial epithelial cells

and bronchial alveolar lavage deciphers GWAS-identified asthma genes. *Allergy*, **70**,

n/a–n/a.

48. Nica,A.C., Montgomery,S.B., Dimas,A.S., Stranger,B.E., Beazley,C., Barroso,I. and

Dermitzakis,E.T. (2010) Candidate causal regulatory effects by integration of

expression QTLs with complex trait genetic associations. *PLoS Genet.*, **6**, e1000895.

49. Veyrieras,J.-B., Kudaravalli,S., Kim,S.Y., Dermitzakis,E.T., Gilad,Y., Stephens,M. and

Pritchard,J.K. (2008) High-Resolution Mapping of Expression-QTLs Yields Insight into

Human Gene Regulation. *PLoS Genet.*, **4**, e1000214.

50. Yang,T.-P., Beazley,C., Montgomery,S.B., Dimas,A.S., Gutierrez-Arcelus,M.,

Stranger,B.E., Deloukas,P. and Dermitzakis,E.T. (2010) Genevar: a database and Java

application for the analysis and visualization of SNP-gene associations in eQTL studies.

Bioinformatics, **26**, 2474–2476.

51. The GTEx Consortium (2013) The Genotype-Tissue Expression (GTEx) project. *Nat.*

Genet., **45**, 580–5.

52. Harley,I.T.W., Kaufman,K.M., Langefeld,C.D., Harley,J.B. and Kelly,J.A. (2009) Genetic

susceptibility to SLE: new insights from fine mapping and genome-wide association

studies. *Nat. Rev. Genet.*, **10**, 285–90.

53. Cui,Y., Sheng,Y. and Zhang,X. (2013) Genetic susceptibility to SLE: Recent progress

from GWAS. *J. Autoimmun.*, **41**, 25–33.

54. Deng,Y. and Tsao,B.P. (2014) Advances in lupus genetics and epigenetics. *Curr. Opin.*

Rheumatol., **26**, 1–11.

55. Yang,J., Lee,S.H., Goddard,M.E. and Visscher,P.M. (2011) GCTA: A Tool for Genome-
wide Complex Trait Analysis. *Am. J. Hum. Genet.*, **88**, 76–82.

56. Giambartolomei,C., Vukcevic,D., Schadt,E.E., Franke,L., Hingorani,A.D., Wallace,C. and
Plagnol,V. (2014) Bayesian Test for Colocalisation between Pairs of Genetic
Association Studies Using Summary Statistics. *PLoS Genet.*, **10**, e1004383.

57. Kozyrev,S. V, Abelson,A.-K., Wojcik,J., Zaghlool,A., Linga Reddy,M.V.P., Sanchez,E.,
Gunnarsson,I., Svenungsson,E., Sturfelt,G., Jönsen,A., *et al.* (2008) Functional variants
in the B-cell gene BANK1 are associated with systemic lupus erythematosus. *Nat.*
Genet., **40**, 211–216.

58. Lewis,M.J., Vyse,S., Shields,A.M., Boeltz,S., Gordon,P.A., Spector,T.D., Lehner,P.J.,
Walczak,H. and Vyse,T.J. (2015) UBE2L3 polymorphism amplifies NF-κB activation
and promotes plasma cell development, linking linear ubiquitination to multiple
autoimmune diseases. *Am. J. Hum. Genet.*, **96**, 221–234.

59. Noble,J.A., White,A.M., Lazzeroni,L.C., Valdes,A.M., Mirel,D.B., Reynolds,R., Grupe,A.,
Aud,D., Peltz,G. and Erlich,H.A. Associated With Type 1 Diabetes. **06**, 1579–1582.

60. Liu,Y., Wang,X., Cheng,X., Lu,Y. and Wang,G. (2015) Skp1 in lung cancer□: Clinical
significance and therapeutic efficacy of its small molecule inhibitors. **6**.

61. Wu,J.Q., Seay,M., Schulz,V.P., Hariharan,M., Tuck,D., Lian,J., Du,J., Shi,M., Ye,Z.,
Gerstein,M., *et al.* (2012) Tcf7 is an important regulator of the switch of self-renewal
and differentiation in a multipotential hematopoietic cell line. *PLoS Genet.*, **8**.

62. Zhou,W., Wei,W. and Sun,Y. (2013) Genetically engineered mouse models for functional
studies of SKP1-CUL1-F-box-protein (SCF) E3 ubiquitin ligases. *Cell Res*, **23**, 599–
619.

63. Bentham,J., Morris,D.L., Cunninghame Graham,D.S., Pinder,C.L., Tombleson,P.,

Behrens,T.W., Martín,J., Fairfax,B.P., Knight,J.C., Chen,L., *et al.* (2015) Genetic association analyses implicate aberrant regulation of innate and adaptive immunity genes in the pathogenesis of systemic lupus erythematosus. *Nat. Genet.*, 10.1038/ng.3434.

64. Westra,H.-J., Peters,M.J., Esko,T., Yaghootkar,H., Schurmann,C., Kettunen,J., Christiansen,M.W., Fairfax,B.P., Schramm,K., Powell,J.E., *et al.* (2013) Systematic identification of trans eQTLs as putative drivers of known disease associations. *Nat. Genet.*, **45**, 1238–43.

65. Ward,L.D. and Kellis,M. (2012) HaploReg: A resource for exploring chromatin states, conservation, and regulatory motif alterations within sets of genetically linked variants. *Nucleic Acids Res.*, **40**, 930–934.

66. Salloum,R., Franek,B.S., Kariuki,S.N., Rhee,L., Mikolaitis,R. a., Jolly,M., Utset,T.O. and Niewold,T.B. (2010) Genetic variation at the IRF7/PHRF1 locus is associated with autoantibody profile and serum interferon-alpha activity in lupus patients. *Arthritis Rheum.*, **62**, 553–561.

67. Wang,T.J., Zhang,F., Richards,J.B., Kestenbaum,B., van Meurs,J.B., Berry,D., Kiel,D.P., Streeten,E.A., Ohlsson,C., Koller,D.L., *et al.* (2010) Common genetic determinants of vitamin D insufficiency: a genome-wide association study. *Lancet*, **376**, 180–188.

68. Yap,K.S., Northcott,M., Hoi, a. B.-Y., Morand,E. and Nikpour,M. (2015) Association of low vitamin D with high disease activity in an Australian systemic lupus erythematosus cohort. *Lupus Sci. Med.*, **2**, e000064–e000064.

69. Wassif,C. a, Zhu,P., Kratz,L., Krakowiak,P. a, Battaile,K.P., Weight,F.F., Grinberg,A., Steiner,R.D., Nwokoro,N.A., Kelley,R.I., *et al.* (2001) Biochemical, phenotypic and neurophysiological characterization of a genetic mouse model of RSH/Smith--Lemli-- Opitz syndrome. *Hum. Mol. Genet.*, **10**, 555–564.

70. Shah,D., Mahajan,N., Sah,S., Nath,S.K. and Paudyal,B. (2014) Oxidative stress and its

biomarkers in systemic lupus erythematosus. *J. Biomed. Sci.*, **21**, 23.

71. Luo,X., Yang,W., Ye,D.-Q., Cui,H., Zhang,Y., Hirankarn,N., Qian,X., Tang,Y., Lau,Y.L.,

de Vries,N., *et al.* (2011) A functional variant in microRNA-146a promoter modulates its

expression and confers disease risk for systemic lupus erythematosus. *PLoS Genet.*, **7**,

e1002128.

72. Hom,G., Ph,D., Graham,R.R., Modrek,B., Taylor,K.E., Ortmann,W., Garnier,S., Lee,A.T.,

Chung,S.A., Ferreira,R.C., *et al.* (2008) Association of Systemic Lupus Erythematosus

with. *Genetics*.

73. Guthridge,J.M., Lu,R., Sun,H., Sun,C., Wiley,G.B., Dominguez,N., MacWana,S.R.,

Lessard,C.J., Kim-Howard,X., Cobb,B.L., *et al.* (2014) Two functional lupus-associated

BLK promoter variants control cell-type- and developmental-stage-specific transcription.

Am. J. Hum. Genet., **94**, 586–598.

74. Kim,H., Barnitz,R.A., Kreslavsky,T., Brown,F.D., Moffett,H., Lemieux,M.E., Kaygusuz,Y.,

Meissner,T., Holderried,T.A.W., Chan,S., *et al.* (2009) egulatory T cells (CD4 and CD8

T.

75. Baine,I., Basu,S., Ames,R., Sellers,R.S. and Macian,F. (2013) Helios induces epigenetic

silencing of IL2 gene expression in regulatory T cells. *J. Immunol.*, **190**, 1008–16.

76. Getnet,D., Grosso,J.F., Goldberg,M. V., Harris,T.J., Yen,H.R., Bruno,T.C., Durham,N.M.,

Hipkiss,E.L., Pyle,K.J., Wada,S., *et al.* (2010) A role for the transcription factor Helios in

human CD4+CD25+ regulatory T cells. *Mol. Immunol.*, **47**, 1595–1600.

77. Isakson,P., Holland,P. and Simonsen,A. (2013) The role of ALFY in selective autophagy.

Cell Death Differ., **20**, 12–20.

78. Yang,W., Shen,N., Ye,D.Q., Liu,Q., Zhang,Y., Qian,X.X., Hirankarn,N., Ying,D.,

971 Pan,H.F., Mok,C.C., *et al.* (2010) Genome-wide association study in asian populations

972 identifies variants in ETS1 and WDFY4 associated with systemic lupus erythematosus.

973 *PLoS Genet.*, **6**.

974 79. McLaren,W., Pritchard,B., Rios,D., Chen,Y., Flicek,P. and Cunningham,F. (2010)

975 Deriving the consequences of genomic variants with the Ensembl API and SNP Effect

976 Predictor. *Bioinformatics*, **26**, 2069–2070.

977 80. Graham,R.R., Kozyrev,S. V, Baechler,E.C., Reddy,M.V.P.L., Plenge,R.M., Bauer,J.W.,

978 Ortmann,W. a, Koeuth,T., González Escribano,M.F., Pons-Estel,B., *et al.* (2006) A

979 common haplotype of interferon regulatory factor 5 (IRF5) regulates splicing and

980 expression and is associated with increased risk of systemic lupus erythematosus. *Nat.*

981 *Genet.*, **38**, 550–555.

982 81. Graham,R.R., Kyogoku,C., Sigurdsson,S., Vlasova,I.A., Davies,L.R.L., Baechler,E.C.,

983 Plenge,R.M., Koeuth,T., Ortmann,W.A., Hom,G., *et al.* (2007) Three functional variants

984 of IFN regulatory factor 5 (IRF5) define risk and protective haplotypes for human lupus.

985 *Proc. Natl. Acad. Sci. U. S. A.*, **104**, 6758–63.

986 82. Lappalainen,T. (2015) Functional genomics bridges the gap between quantitative

987 genetics and molecular biology. *Genome Res.*, **25**, 1427–1431.

988 83. Battle, a. and Montgomery,S.B. (2014) Determining causality and consequence of

989 expression quantitative trait loci. *Hum. Genet.*, **133**, 727–735.

990 84. Jäger,R., Migliorini,G., Henrion,M., Kandaswamy,R., Speedy,H.E., Heindl,A., Whiffin,N.,

991 Carnicer,M.J., Broome,L., Dryden,N., *et al.* (2015) Capture Hi-C identifies the chromatin

992 interactome of colorectal cancer risk loci. *Nat. Commun.*, **6**, 6178.

993 85. Zhao,Z., Tavoosidana,G., Sjölinder,M., Göndör,A., Mariano,P., Wang,S., Kanduri,C.,

994 Lezcano,M., Sandhu,K.S., Singh,U., *et al.* (2006) Circular chromosome conformation

995 capture (4C) uncovers extensive networks of epigenetically regulated intra- and

interchromosomal interactions. *Nat. Genet.*, **38**, 1341–1347.

86. Kelsell,D.P., Norgett,E.E., Unsworth,H., Teh,M.-T., Cullup,T., Mein,C.A., Dopping-

Hepenstal,P.J., Dale,B.A., Tadini,G., Fleckman,P., *et al.* (2005) Mutations in ABCA12

underlie the severe congenital skin disease harlequin ichthyosis. *Am. J. Hum. Genet.*,

76, 794–803.

87. Steidl,U., Steidl,C., Ebralidze,A., Chapuy,B., Han,H., Will,B., Rosenbauer,F., Becker,A.,

Wagner,K., Koschmieder,S., *et al.* (2007) A distal single nucleotide polymorphism alters

long-range regulation of the PU . 1 gene in acute myeloid leukemia. *J. Clin. Invest.*,

117, 2611–2620.

88. Freedman,M.L., Monteiro,A.N.A., Gayther,S.A., Coetzee,G.A., Risch,A., Plass,C.,

Casey,G., De Biasi,M., Carlson,C., Duggan,D., *et al.* (2011) Principles for the post-

GWAS functional characterization of cancer risk loci. *Nat. Genet.*, **43**, 513–518.

89. Graveley,B.R. (2008) The haplo-spliceo-transcriptome: common variations in alternative

splicing in the human population. *Trends Genet.*, **24**, 5–7.

90. Wang,G.-S. and Cooper,T.A. (2007) Splicing in disease: disruption of the splicing code

and the decoding machinery. *Nat. Rev. Genet.*, **8**, 749–761.

91. Li,Y.I., van de Geijn,B., Raj,A., Knowles,D.A., Petti,A.A., Golan,D., Gilad,Y. and

Pritchard,J.K. (2016) RNA splicing is a primary link between genetic variation and

disease. *Science*, **352**, 600–4.

92. Wang,J.H., Avitahl,N., Cariappa,A., Friedrich,C., Ikeda,T., Renold,A., Andrikopoulos,K.,

Liang,L., Pillai,S., Morgan,B.A., *et al.* (1998) Aiolos regulates B cell activation and

maturation to effector state. *Immunity*, **9**, 543–553.

93. Shabalin,A.A. (2012) Matrix eQTL: ultra fast eQTL analysis via large matrix operations.

Bioinformatics, **28**, 1353–1358.

94. Zheng,X., Levine,D., Shen,J., Gogarten,S.M., Laurie,C. and Weir,B.S. (2012) A high-

performance computing toolset for relatedness and principal component analysis of

SNP data. *Bioinformatics*, **28**, 3326–3328.

95. Li,H., Handsaker,B., Wysoker,A., Fennell,T., Ruan,J., Homer,N., Marth,G., Abecasis,G.,

Durbin,R. and Subgroup,1000 Genome Project Data Processing (2009) The Sequence

Alignment/Map format and SAMtools. *Bioinformatics*, **25**, 2078–2079.

96. Ongen,H. and Dermitzakis,E.T. (2015) Alternative Splicing QTLs in European and

African Populations. *Am. J. Hum. Genet.*, **97**, 567–575.

97. Pruim,R.J., Welch,R.P., Sanna,S., Teslovich,T.M., Chines,P.S., Gliedt,T.P., Boehnke,M.,

Abecasis,G.R. and Willer,C.J. (2010) LocusZoom: Regional visualization of genome-

wide association scan results. *Bioinformatics*, **26**, 2336–2337.

98. Cunningham,F., Amode,M.R., Barrell,D., Beal,K., Billis,K., Brent,S., Carvalho-Silva,D.,

Clapham,P., Coates,G., Fitzgerald,S., *et al.* (2015) Ensembl 2015. *Nucleic Acids Res.*,

43, D662–D669.

99. Kent,W.J., Sugnet,C.W., Furey,T.S. and Roskin,K.M. (1976) The Human Genome

Browser at UCSC W. *J. Med. Chem.*, **19**, 1228–31.

1036

Legends to figures

Fig. 1: Two-stage *cis*-eQTL annotation pipeline for SLE susceptibility loci.

SLE susceptibility variants (**Table 1**) were annotated using residualized expression or summary-level eQTL statistics from three expression datasets: microarray probe-level expression data, and both gene-level and exon-level RNA-Seq quantifications. Each expression dataset was generated from LCLs from individuals of the TwinsUK cohort. **A**) We undertook *cis*-eQTL analysis of +/-1Mb intervals around each SNP and associations with q<0.05 after FDR adjustment were taken forward. **B**) Summary-level data from significant *cis*-eQTLs were tested for evidence of a shared causal variant using firstly conditional analysis using the TwinsUK genetic data as a reference panel, then colocalisation analysis to test for a single causal variant common to both traits. Associations passing these thresholds (described fully in methods) were classified as candidate-causal eQTLs and eGenes. Summary results per quantification type for significant and candidate-causal associations are shown in **Table 3** for microarray, **Table 4** for RNA-Seq (gene-level), and **Table 5** for RNA-Seq (exon-level). Full summary results are available in **Table S1**, **S2**, and **S3** respectively.

Fig. 2: Candidate-causal eQTLs and eGenes detected across quantification types. Candidate-causal eQTLs and eGenes detected using microarray (probe-level), RNA-Seq (gene-level), and RNA-Seq (exon-level) quantifications from the TwinsUK datasets are represented in their genomic context (**Tables 3**, **4**, and **5** respectively). Only chromosomes harbouring one or more candidate-causal eQTL or eGene per quantification type are shown.

Fig. 3: RNA-Seq gene-level and exon-level analysis implicate *NADSYN1* as a novel candidate-causal eGene.

A) *Cis*-eQTL analysis of GWAS SNP rs3794060 reveals the risk variant [C] leads to

substantial down-regulation at the gene-level of novel susceptibility gene *NADSYN1* in the

TwinsUK cohort that was not detected using microarray. **B**) The same analysis using exon-

level quantification leads to the inference of the gene-level effect being driven by

considerable expression disruption of two meta-exons of *NADSYN1* (meta-exon 11 and

meta-exon 12). Association *P*-values of rs3794060 against exon quantifications are plotted

with reference to the specific exon in the collapsed-gene model of *NADSYN1* (all annotated

transcripts combined).

1070

Fig. 4: Non-coding candidate-causal eGenes detected using exon-level RNA-Seq.

The three figures denote the eQTLs and corresponding non-coding eGenes identified from

cis-eQTL analysis of GWAS SNPs against TwinsUK exon-level quantifications in LCLs. The

top panels display the signal from the GWAS association plotted as $-\log_{10}$ (*P*), below which

the exon-level eQTL *P*-values for the effects showing colocalisation with the GWAS signal

are illustrated. The bar across the middle of the figure denotes the boundaries of the eQTL,

below which there is a panel showing the association *P*-value of the GWAS SNP against the

candidate-causal non-coding exons. The bottom panel shows LCL RNA-Seq alignments

from ENCODE to show that the regions containing the candidate-causal eQTLs are

expressed. **A**) GWAS SNP rs2431697 is a candidate-causal eQTL for non-coding eGene

MIR146A. **B**) GWAS SNP rs2736340 is a candidate-causal eQTL for non-coding eGenes

RP11-148O21.4 and *RP11-148O21.2*. **C**) GWAS SNP rs3794060 is a candidate-causal

eQTL for non-coding eGene *RP11-660L16.2*.

1084

1085

1086

Fig. 5: Identification of novel eGene *IKZF2* and potential causal mechanism using

RNA-Seq splice-junction quantification.

Cis-asQTL analysis of GWAS SNP rs3768792 against splice-junction quantifications

classified *IKZF2* as a candidate-causal eGene with risk variant [G] causing upregulation of

the exon 6A–exon 6B junction that is unique to truncated isoform ENST00000413091. **A**)

GWAS association signal across the *IKZF2* locus (chr2q34), tagged by rs3768792 localised

in the 3'-UTR of *IKZF2*. *Cis*-asQTL association signal of rs3768792 against splice-junction

quantification of exon 6A–exon 6B shows significance and colocalisation with the GWAS

signal. **B**) The exon 6A–exon 6B junction is unique to truncated isoform ENST00000413091.

Exon 6B harbours a premature stop-codon and therefore is not translated into the full-length

protein that contains the dimerization domains in exon 8. **C**) Close-up of the exon 6A–exon

6B junction and association ($P=3.80\mathrm{E}^{-05}$) with GWAS SNP rs3768792. A potential causal

asQTL in near-perfect LD was identified that is located within the polypyrimidine tract of the

junction and may induce splicing (rs2291241, $P=1.70^{\mathrm{E-05}}$).

1101

Fig. 6: Identification of potential disease-associated mechanisms at the *WDFY4*

susceptibility locus using asQTL mapping.

A) Our SLE GWAS indicates *WDFY4* as the candidate gene at the chr10q11.23 locus

tagged by intronic variant rs2663052, as well as the missense coding variant rs7097397 in

exon 31 that is in strong LD. *Cis*-eQTL showed rs2663052 is correlated with upregulation of

the exon 34A–34B junction of *WDFY4* (signal is colocalised with GWAS) that is unique to the

short isoform (ENST00000374161). This isoform lacks the two enzymatic WD40 domains of

the full length isoform (ENST00000325239). **B**) Two potential functional mechanisms may

occur when harbouring the risk haplotype that carries both risk alleles. Firstly an Arg to Gln

amino-acid substitution by rs7097397 in exon 31 that is shared by both the canonical and

short isoforms of *WDFY4*, and secondly an upregulation of the short isoform ($P=3.31^{\mathrm{E-19}}$)

that lacks functional domains, caused by rs2663052 or correlated variants, with

corresponding down-regulation of the full-length isoform (P=3.01^{E-06}).

1115

Fig. 7: Heatmap of candidate-causal eQTLs and eGenes detected across all
expression-quantification types.

Heatmap of all candidate-causal *cis*-eQTL associations across the four quantification types

(microarray, RNA-Seq gene-level, RNA-Seq exon-level, and RNA-Seq splice-junction level).

The first column is the key showing the relative P value of the eQTLs within each platform.

For the platform-specific columns, if an eQTL-eGene association is candidate-causal in at

least one quantification type, the data is displayed across all platforms. Rows are or ordered

by decreasing cumulative significance across quantification types. To normalize across

quantification types, relative significance of each association per column was calculated as

the $-\log_2$ (P/P_{max}); where P_{max} is the most significant association per quantification type. If an

association is deemed to be candidate-causal within a particular profiling-type, it is

highlighted with an asterisk.

Supporting Information

S1 Table. All significant eQTLs (q < 0.05) and associated eGenes detected at microarray (probe-level) with conditional and colocalisation results.

S2 Table. All significant eQTLs (q < 0.05) and associated eGenes detected at RNA-Seq (gene-level) with conditional and colocalisation results.

S3 Table. All significant eQTLs (q < 0.05) and associated eGenes detected at RNA-Seq (exon-level) with conditional and colocalisation results.

S1 Fig. Number of eQTL discoveries per quantification type. Including significant associations (q<0.05), and candidate-causal associations (significant, and not-independent and colocalised with GWAS).

S2 Fig. Number of eGene discoveries per quantification type. Including significant associations (q<0.05), and candidate-causal associations (significant, and not-independent and colocalised with GWAS).

S3 Fig. Shared candidate-causal eQTLs per quantification type.

S4 Fig. Shared candidate-causal eGenes per quantification type.

S5 Fig. Ratio of eQTLs to candidate-causal eGenes per quantification type.

S6 Fig. Gene-level and exon-level candidate-causal associations with *TCF7* and *SKP1* against rs7726414. *Cis*-eQTL analysis at gene-level and exon-level using RNA-Seq implicate novel SLE-associated eGenes *TCF7* and *SKP1*.

S7 Fig. High gene-density over associated variants tagged by GWAS SNP rs12802200. The six candidate-causal eGenes against rs1280220 discovered using RNA-Seq at either quantification method are marked with an asterisk.

S8 Fig. Candidate-causal eGenes *DHCR7* and *NADSYN1* for rs3794060, and non-

coding eGene *RP11-660L16.2*. The GWAS SNP rs3794060 is a candidate-causal eGene for *DHCR7* and *NADSYN1*, and also the non-coding eGene *RP11-660L16.2* at exon-level; which is located between *DCHR7* and *NADSYN1*.

S9 Fig. Candidate-causal eGenes *FAM167A*, *BLK* and two non-coding RNAs (*RP11-138021.4* and *RP11-138021.2*) driven by rs2736340. Associated variant rs2736340 lies in a region of intense regulatory chromatin marks located in the bi-directional promoter of *BLK* and *FAM167A* which are both candidate-causal eGenes at RNA-Seq gene-level. At exon-level, the non-coding eGenes *RP11-138021.4* and *RP11-138021.2* are also candidate-causal.

S10 Fig. Effect-size correlation of GWAS SNP associations with matched *cis*-exons between LCL and whole-blood.

S4 Table. All significant eQTLs (q < 0.05) and associated eGenes detected at RNA-Seq (exon-level) I with conditional and colocalisation results in whole-blood.

S11 Fig. Candidate-causal eQTLs and eGenes in whole blood. Comparison with LCL associations.

S12 Fig. Whole-blood exon-level eQTL effect on *BANK1* exon 2. Correlation between GWAS SNP and the best whole-blood eQTL for *BANK1* exon 2. Both are highly correlated with known branch-point SNP. All are weakly correlated with best LCL eQTL for *BANK1* exon 2.

S13 Fig. Exon-level eQTL analysis of *NADSYN1* in whole-blood and LCLs reveal near-identical splicing effect. Meta-exons 11 and 12 are substantially disrupted with reference to GWAS SNP rs3794060 in both LCLs and whole-blood.

S5 Table. All significant asQTLs (q < 0.05) and associated eGenes detected at RNA-Seq (splice-junction level).

S14 Fig. Proposed splicing mechanism of *NADSYN1* caused by risk haplotype tagged

by rs3794060. *NADSYN1* Ensembl transcript annotation displayed. *Cis*-asQTL identified the

meta-exon 10 to meta-exon 12 junction is upregulated with risk allele [C] and consequently

the meta-exon 11 to meta-exon 12 junction is downregulated.

S6 Table. Comparison of eQTLs and eGenes for SLE risk alleles between previously

reported in microarray studies and from RNA-Seq in current study.

Abbreviations

GWAS: Genome-Wide Association Study

eQTL: expression Quantitative Trait Loci

RNA-Seq: RNA-Sequencing

SLE: Systemic Lupus Erythematosus

asQTL: alternative-splicing Quantitative Trait Loci

SNP: Single Nucleotide Polymorphism

LCL: Lymphoblastoid cell line

Fig 1.

Fig 2.

Chromosome	I	II	III	VI	V	VIII	XI	XII	XV	XVII	XXII

Fig 3.

Fig 4.

Fig 5.

Fig 6.

Fig 7.