A critical review of the use and performance of different function types for modeling

temperature-dependent development of arthropod larvae

Brady K. Quinn*

Department of Biological Sciences, University of New Brunswick, 100 Tucker Park Road, Saint

John, NB, Canada E2L 4L5

*Corresponding author: bk.quinn@unb.ca, 1-506-343-7676

Highlights:

• Temperature-dependent development functions of arthropod larvae were reviewed

• 79 published datasets were re-tested and fit with 33 different function types

• 91.1 % of published studies did not fit their data with the best function of those tested

• Performance differed among functions and was related to taxon and temperature range tested

• Function type impacted predicted development times, so using the best function matters

ABSTRACT

Temperature-dependent development influences production rates of arthropods, including

crustaceans important to fisheries and agricultural pests. Numerous candidate equation types

(development functions) exist to describe the effect of temperature on development time, yet

most studies use only a single type of equation and there is no consensus as to which, if any

model predicts development rates better than the others, nor what the consequences of selecting a

potentially incorrect model equation are on predicted development times. In this study, a

literature search was performed of studies fitting development functions to development of

arthropod larvae (99 species). The published data of most (79) of these species were then fit with

33 commonly-used development functions. Overall performance of each function type and

consequences of using a function other than the best one to model data were assessed.

Performance was also related to taxonomy and the range of temperatures examined. The majority

(91.1 %) of studies were found to not use the best function out of those tested. Using the

incorrect model lead to significantly less accurate (e.g., mean difference $\pm$ SE 85.9 $\pm$ 27.4 %,

range: -1.7 to 1725.5 %) predictions of development times than the best function. Overall, more

complex functions performed poorly relative to simpler ones. However, performance of some

complex functions improved when wide temperature ranges were tested, which tended to be

confined to studies of insects or arachnids compared with those of crustaceans. Results indicate

the biological significance of choosing the best-fitting model to describe temperature-dependent

development time data.

Key Words: Temperature; arthropod; larval development; development functions; molting rate

1. Introduction:

Temperature affects biota at all levels, ranging from effects at the fundamental biochemical and physiological levels (Bělehrádek, 1935; Coutant and Talmage 1976; Somero, 2004) to effects on individual organisms (Brière et al., 1999; MacKenzie, 1988), populations (Aiken and Waddy, 1986; Cooper et al., 2012; McLaren et al., 1969), communities, and ecosystems (Menge, 1978; McQuaid and Branch, 1985). Through its effects on the physical and chemical properties of biologically active molecules, such as enzymes, temperature affects the rate at which numerous life processes occur, including metabolism, oxygen consumption, photosynthesis, movement, survival, growth, and embryonic development (Bělehrádek, 1935; Brière et al. 1999; Corkett, 1972; Coutant and Talmage 1976; Du et al., 2007; Geffen and Nash, 2012; Herzig, 1983; McLaren et al., 1969). Temperature also significantly affects larval development rate of poikilothermic animals, including some vertebrate larvae (Lind and Johansson, 2007; Kang et al., 2009; Miller et al., 2006) and those of invertebrates (e.g., de Severyn et al., 2000; Jenkins et al. 2006; Singh and Sharma, 1994). Temperature has particularly strong impacts on moulting and development of arthropods (Anger, 1984; Corkett and McLaren, 1970; Easterbrook et al., 2003; Hamasaki et al., 2009; Koda and Nakamura, 2010; MacKenzie, 1988; Marchioro and Forester, 2011; McLaren, 1963).

Within certain tolerance limits (Bělehrádek, 1935; Brière et al., 1999; Campbell et al., 1974; Shi and Ge, 2010) rates of biological processes of poikilotherms, including larval development, are positively correlated with temperature; thus, higher temperatures generally result in more rapid development than lower temperatures. This has important ecological implications, as environmental temperatures can influence generation times, production cycles, and population dynamics of such organisms. Higher or lower temperatures could, for example,

lead to changes in the amount and/or timing of peak secondary marine production of copepods

(Huntley and López, 1992; McLaren, 1963), outbreaks of agricultural pests (Easterbrook et al.,

2003) or vector-borne diseases (Bayoh and Lindsay, 2003), or introduction and establishment of

invasive species into new areas (de Rivera et al., 2007). Water temperatures could also influence

patterns of recruitment to populations of marine invertebrates, including crustaceans such as

lobsters and crabs, on which human fisheries depend (Aiken and Waddy, 1986; Anger, 1984;

Caddy, 1986; MacKenzie, 1988; Rothlisberg, 1979).

When modeling ecology and population dynamics of arthropods, equations are used to

represent the functional relationship between environmental temperature and development rate or

time of larvae. These equations, hereafter referred to as development functions, are derived by

rearing larvae at different controlled temperatures in a lab or hatchery setting, observing

development times of multiple larvae at each temperature, and then using regression analyses to

fit an equation relating temperature to development time (or its inverse, rate) to the data obtained.

There are countless potential forms of equation that can be used to fit such data, including

various linear, simple curvilinear, and complex non-linear functions (e.g., see reviews by Anger,

2001; Angilleta Jr., 2006; Blanco et al. 1995; Guerrero et al., 1994; Heip, 1974; Kontodimas et

al., 2004; McLaren, 1995; Shi and Ge, 2010; Smits et al. 2003). These functions differ in form,

assumptions, procedures used to derive their parameters, and most importantly in terms of the

development times predicted. For example, development times of American lobster, *Homarus*

americanus (H. Milne Edwards, 1837), larvae predicted with 33 of these development function

types can differ from each other and the data used to derive them by ≥ 50 days at the same

temperatures (see Fig. 1; Table S1). However, this is no clear standard rule or consensus as to

what is the "best" type of development function to apply to these kinds of data. Researchers are

generally left to choose the type of development function to use on their own, and will often

select one or a very few forms that have the best apparent match to their characters of their data

or has been used by other studies on the same or related species (e.g., Edgar and Andrew, 1990;

McLaren et al., 1969). Given the potential for different functions to make very different

predictions of development times (e.g., Fig. 1), however, development function choice should be

given more consideration in studies on these species.

It is possible that certain function types may in general be better representations of the

relationship between temperature and development of arthropod larvae, or specific sub-groups

within the Arthropoda (e.g., arachnids vs. crustaceans vs. insects), for example because they

come closer to capturing thermal performance relations of enzymes and other biomolecules

mediating moulting and development cycles in these taxa (Brière et al., 1999; Huey and

Stevenson, 1979; Somero, 2004). As a result, such functions might also achieve better fit to

development data, be able to more-closely match real observed development times, and make

better predictions of development in nature. Differences in methodologies used in studies on

different taxa, for example the fact that the range of temperatures tested is generally wider for

insects and arachnids than crustaceans; (reviewed by Hartnoll, 1982; Quinn and Rochette, 2015),

might also lead to apparent taxonomic differences in function performance and should be

investigated. Several previous studies have compared the characteristics of different

development function types in general (Anger, 2001; Blanco et al., 1995; Guerrero et al., 1994;

McLaren, 1995). Others have examined performance of one or two specific functions on

multiple species (e.g., Logan et al., 1976), or attempted to fit multiple function types to data for

one or two specific species under study to select the best function for their data (e.g., Angilletta

Jr., 2006; Heip, 1974; Kontodimas et al., 2004; Shi and Ge, 2010; Smits et al., 2003). Many other

studies seem to choose one or very few function(s) semi-arbitrarily, without discussing

alternatives (e.g., de Oliveria et al., 2009; Thompson 1982; see also Results). However, no

previous study has attempted to assess the degree to which one versus multiple types of

development functions are used in published studies, compared performance of different types of

development functions across multiple species, or assessed the overall impact of function choice

to predictions made with such functions. Such a large-scale analysis is needed, though, because it

could potentially allow functions that tend to better represent development data in general to be

identified, which can then allow for more informed decisions by future studies on arthropod

larval development.

In this study, a critical literature review was conducted to assess whether and to what

extent studies of temperature-dependent development of arthropod larvae attempt to represent

their data with more than one development function type, and also which specific types of

functions tend to be used. Then, data from previously published studies were extracted and

retested to derive multiple different development functions for the same datasets. The best model

type for each dataset was determined, and whether or not published studies actually used the best

function type for their data was recorded. Overall performance of different function types were

assessed by comparing overall function rankings, proportion of variance explained, and

information loss across datasets. Any taxonomic patterns (e.g., whether particular function types

performed better for arachnids than for crustaceans and/or insects) and whether performance was

related to the range of temperatures tested were also noted. The consequences of using different

function types were then tested by comparing the difference of predicted versus observed

development times, fit, and information loss of the function type used in original studies versus

that of the best function for the data.

2. Materials and methods:

2.1. Literature review:

A literature search was conducted through Web of Science (Thompson Reuters, 2015) for

the terms "temperature" AND "development". An initial search was carried out on 12 September

2012, through which the majority of the data in the present study were obtained; this search

yielded 1,052 results. A second search was carried out on 19 November 2014, which returned 35

additional results not available or published online at the time of the initial search. These 1,087

total search results were them further examined, and several criteria were used to remove non-

relevant results. Accessible peer-reviewed studies that reported larval development rates or times

of arthropods at different temperatures and derived a regression equation(s) (i.e., development

function) from their data were sought out. Studies that looked exclusively at growth (size

increase), which is a distinct process from development (Forster et al., 2011), were excluded.

After applying these criteria, 81 studies of 96 different arthropod species were obtained,

which provided a total of 99 species datasets for subsequent examination and analyses (Table S2).

Several specific types of development function were frequently utilized in these studies (Table

S2); these functions are presented in Table 1 and discussed in the next section (2.2.1, below).

Studies were published between 1970 and 2014, and conducted in several different countries on

marine, freshwater, and terrestrial species (Table S2) of various taxa within the Arachnida

(Phylum Arthropoda: Subphylum Chelicerata), Crustacea, and Insecta (Table S2). To assess

whether published studies tested multiple development functions on their data, each study was

carefully read and the number and types of development functions used to fit the data for each

study species were noted. Even if results from multiple functions were not reported, if alternative

functions than reported were at least mentioned in the Methods sections of studies they were

counted as having considered > 1 function. Also if any study tested multiple development

functions on their data and concluded one of these to be the "best" function for their data this

was also noted. The percent (%) of the 99 species datasets from the literature search on which

one, two, or more functions were tested, and the % of datasets on which different types of

functions were tested, were then calculated.

2.2. Meta-analysis:

2.2.1. Development functions considered in this study:

In the present study, 33 development function types were examined (Table 1; Figure 1).

These functions were used because they were found in the literature review in the present study

(see section 2.1 and Results) to be used quite frequently in studies of arthropod larvae, and were

discussed in reviews and studies on the topic of arthropod temperature-development functions by

Anger (2001), Guerrero et al. (1994), Heip (1974), Kontodimas et al. (2004), and Shi and Ge

(2010). Three linear and 30 nonlinear functions were examined, with *k*-values (*k* = numbers of

parameters + 1; Anderson, 2008) ranging from 3 to 8 (Table 1). Eight of these functions are fit

directly to development time data, while the remaining 25 functions are typically fitted to

development rate data, which are the inverse of time (Table 1); in some cases, the same function

form (e.g., quadratic) is applied to either development rate (function #15, Table 1) or time (#16),

yielding distinct functions (Fig. 1). 12 of these functions included a nonzero minimum

temperature (T_{min}) for development at which development rate is zero and development time

becomes infinite (see Fig. 1), 15 included a similar maximum temperature for development

(T_{max}), and 8 functions included both of these thresholds (Table 1). Linear functions can be used

to derive starting estimates for the values of T_{min} and T_{max} (Campbell et al., 1974; Kontodimas et

al., 2004; Table 1) to be used in deriving more complicated nonlinear functions. Attempts were

also made to test three additional functions found in these reviews: the Exponentially Modified

Gaussian (Naish and Hartwell, 1988), Sharpe-Schoolfield-Ikemoto (SSI; Sharpe and DeMichele,

1977; Schoolfield et al., 1981), and Weibull (Angilletta Jr., 2006) functions. However, these

three functions have very complex structures requiring specialized fitting procedures not readily

applicable in many statistical software packages (see review by Shi and Ge, 2010), and in the

present study they could not be fit to the datasets used; as such, they were not considered further

in this study.

2.2.2. Re-analysis of species datasets from published studies:

To assess whether the function(s) used in published studies were actually the "best"

functions for these published datasets, data were extracted for re-analysis from as many of the

studies obtained through the literature search described above as possible. Studies from the

literature search that did not present their data in a way that allowed it to be extracted for

retesting (e.g., only mean development times resented, without any measure of error), had to be

excluded. Therefore, only 79 species datasets out of the 99 initially obtained from the literature

search could be retested (Table S2); these included 10 arachnids (9 mites in the Subclass Acari

and one spider in Subclass Aranea), 28 Crustaceans (9 copepods and 19 decapods), and 41

insects in 8 orders (8 Coleoptera, 6 Diptera, 9 Hemiptera, 1 Homoptera, 1 Heteroptera, 8

Hymenoptera, 6 Lepidoptera, and 2 Thysanoptera) (Table S2). Raw data or means ± error

(standard deviation (SD), standard error (SEM), etc.) and sample sizes were extracted from

tables or figures in published papers for each of these 79 species and used to generate datasets

for reanalysis.

Each study dataset was analyzed with linear and nonlinear regressions (Table 1) between

temperature and development time or rate, as appropriate. These regression were carried out

using IBM SPSS Statistics 22 (SPSS Inc., 2014). To simplify analyses, only total development

times or rates (i.e., summed across multiple larval stages) were examined and data for individual

stages were not. For development functions including thermal limits or other parameters with

biological meaning (e.g., T_{min} or T_{max}; see Table 1 and section 2.2.1), unconstrained regressions

were initially carried out, with starting values for these parameters set to values estimated from

linear functions (see Table 1 and section 2.2.1, above). However, results were not accepted if this

yielded biologically unrealistic estimates, such as $T_{min} < 0°C$ in a species not known to survive

and develop at sub-zero temperatures, unreasonably low T_{min} (e.g., -100°C) or high T_{max} (e.g.,

100°C), or T_{min} or T_{max} within the temperature range for which successful development was

reported in the original study. In this case, constrained regressions were carried out (e.g., $T_{min} \geq$

0°C, $T_{min} < $ minimum T with successful development, $T_{max} > $ maximum T with successful

development, etc.) until satisfactory values were obtained.

Once a regression equation corresponding to each development function was obtained for

each dataset, AIC_C values (Akaike's Information Criterion (AIC) corrected for finite sample size;

Akaike, 1973; Anderson, 2008) were calculated using the residual sum of squares (RSS) between

observed development times and those predicted by each function for each study (Anderson,

2008). Development functions were then ranked for each species dataset based on AIC_C-values.

The "best" possible rank was 1, corresponding to the lowest AIC_C value, and the "worst"

possible rank – if there were no ties – was 33, which corresponded to the highest AIC_C value

among the functions tested. The percentage of retested datasets for which each of the 33

functions was concluded to be "best" was recorded. For each species dataset, whether or not the

function used in its original published study and concluded to be the "best" function for the data

was the same as that determined to be the "best" function in this study was noted. If more than

one function was used in an original study, whether the actual best function was among all these

functions was also noted. RSS values were also used to calculate R^2-values for each function, on

each dataset. These values were used as a measure of function performance (see next section,

2.2.3), indicating the proportion of the variation in observed development times that was

explained by each temperature-dependent development function.

2.2.3. Assessing and comparing the overall performance of different functions:

To determine whether any particular type(s) of function tended to do "better" than others,

the overall performance of each development function across species datasets was assessed using

three measures: average ranking (determined using AIC_C), R^2, and Δ_i values. These measures of

performance were also compared across different taxonomic groups (arachnids, crustaceans, and

insects) as it was possible that certain functions might better represent development of animals in

a certain group(s) better than animals in others; this could be due to real biological differences

among taxa or to different experimental methodologies used for their rearing.

Calculation of function rankings and R^2-values for each function on each dataset was

described above (previous section). Δ_i values were calculated once the best function for a given

dataset was determined to assess the information potentially lost by using a function other than

the best one (Anderson, 2008); a lower Δ_i value is better, and indicates less information loss. The

Δ_i value for a given function, "i", is calculated by subtracting the AIC_C value of the best function

of those tested from its AIC_C value (Anderson 2008); therefore, the best function will have $\Delta_i =$

0 . A function with a lower Δ_i, higher R^2, or lower (better) ranking value on average than other

functions was considered to have performed better overall than other functions.

Separate two-way ANOVAs were carried out in IBM SPSS Statistics 22 (SPSS Inc.,

2014) to compare each of these three measures of performance (rankings, R^2, and Δ_i) across

different development functions (factor with 33 levels; Table 1), as well as among different

taxonomic groups (factor with 3 levels: Arachnida, Crustacea, and Insecta). R^2-values were

arcsine-square root transformed to meet the assumptions of parametric tests. If a statistically-

significant ($p \le 0.05$) interaction between function and taxon was found, the data for that

measure were split by taxon and then separate one-way ANOVAs comparing different functions

were carried out for each taxon. If significant differences among functions were found, Tukey's

Honestly Significant Difference (HSD) test was used to perform post-hoc comparisons among

specific function types.

2.2.4. Assessing the consequences of development function choice:

The consequences of choosing one function versus another to predict larval development

at different temperatures were assessed by comparing whether and how much the best or only

function originally used in the study from which each species dataset was obtained ("original

best" function) predicted observed development times relative to the best function identified in

this study ("actual best" function). Three measures, described in detail below, were calculated for

each of the 79 species datasets to assess consequences of function choice. These were mean error,

R^2 decrease, and Δ_i resulting from using the original study function instead of the actual best one.

To calculate the first of these, error, the absolute deviance (in days) between predicted

(using a development function) and observed development time was calculated for both the best

and original study functions, at each temperature tested in original studies. The absolute deviance

for the actual best function was then subtracted from that for the original best/used function at

each temperature, to determine how much predictions were worse (i.e., further from observed

values) when using the original versus best function. These differences were then averaged

across all temperatures and data points to calculate a mean absolute error (in days) per each

dataset that was due to using the original versus actual best functions. Mean error per dataset was

also expressed as a percent improvement by performing the aforementioned calculations, but

before averaging differences between deviance of original and actual best functions across

temperatures these differences were divided by the best function's deviance and multiplied by

100 %; this translated the error from a "raw" measure (in days) to a percentage (%).

R^2 decrease was simply calculated for each dataset by subtracting the R^2-value of that

dataset's actual best function from that of its original used function. Percent R^2 increase was also

calculated for each dataset by dividing the R^2 change by the best function's R^2, and then

multiplying by 100 %. A large R^2 decrease implied that a lower proportion of the variation in

development time was explained by the original function than the best one.

The Δ_i of the original function for each dataset was also examined to assess the extent of

potential information lost by using these, rather than the best functions, to fit the data. As

described above, a lower Δ_i (closer to $0 = \Delta_i$ of best function) is better, and implies the function

retains more useful information than a function with a higher Δ_i. Generally a function with $\Delta_i < 2$

contains some useful information, even if it is not the "best" function, whereas a function with Δ_i

> 14 is highly unlikely to be informative (Anderson 2008). A high original function Δ_i value

would imply that the original function was considerably less informative than the best function.

301   If the best function for a given dataset was the same as the original function, all of the

measures described above would have a value of zero; if any measure were not significantly

different from zero, then, it would imply that using the best versus original function did not result

in meaningfully different predicted development times. Therefore, the distributions of each

measure of the consequences of using a function other than the best one (mean errors and R^2

decrease (both raw and % versions), as well as original function Δ_i values) across all species

datasets were compared to zero (null hypothesis of no differences) using five one-sample t-tests.

If the null hypothesis comparing these data against zero-values could be rejected, then a

significant ($p < 0.05$) impact of function choice was concluded.

2.3. Potential relationship between thermal range and function performance:

312   One interesting pattern noted during the literature review in this study was that studies of

temperature-dependent development differed considerably in their methodology, particularly

regarding the range of temperatures tested. Thermal ranges varied considerably among studies in

general, from as narrow as 6°C to as wide as 38°C (Table S2). Differences also appeared to exist

between studies of different taxonomic groups, particularly between studies of crustaceans

versus those of arachnids and insects (Hartnoll, 1982; Quinn and Rochette, 2015; Table S2). To

confirm whether such taxonomic differences were significant, the thermal range for each species

dataset obtained in the initial literature search (n = 99 total) was calculated as the maximum

temperature tested in its original study minus the minimum temperature tested; this included any

temperatures at which successful development was not observed (i.e., survival = 0 %), as this

implies a developmental threshold (i.e., T_{min} and T_{max}; see Table 1). One-way ANOVA was then

used to compare thermal ranges among the three major taxonomic groups of arthropods.

More complex functions with more parameters have lower power to model smaller

datasets (Angilletta Jr., 2006; Shi and Ge, 2010). This is especially true for functions containing

threshold temperatures (T_{min} and T_{max}) if these are fit to data recorded over narrow thermal

ranges not approaching a species' real thermal limits (Shi and Ge, 2010). In such cases, thermal

thresholds have to be extrapolated too far beyond actual observations, resulting in excessively

extreme estimates for these parameters. One could thus expect that more complex functions may

perform better when wider thermal ranges are tested, potentially approaching or encompassing

real thermal limits. To test whether or not more complex functions would be selected as better

functions when wider thermal ranges were tested, Pearson's correlation coefficients (R) were

calculated between the thermal ranges calculated for each dataset and the ranking of each

function per dataset. Thus, 33 separate correlation analyses were carried out (one for each type of

development function), each consisting of 99 thermal range-ranking pairs (one pair per dataset).

Whether correlations were significant ($p \leq 0.05$), and if significant R-coefficients were positive

or negative was examined. As lower ranking values implied better function performance (see

above), a positive correlation for a given function implied that the function did worse when a

wider range of temperatures was tested, whereas a negative correlation meant that the function

did better when a wider thermal range was tested.

3. Results:

<u>3.1. Results of literature review – usage of different functions:</u>

Out of 99 different species datasets, over half (59.6 %) were reported to have been fit

with only one development function and the vast majority (96.0 %) were fit with five or fewer

functions (Fig. 2). In most cases studies that used 2-5 functions examined insects or arachnids

(Fig. 2) and used 1-3 more complex functions plus the linear rate (function #5) or Ikemoto and

Takai (#4) functions to derive starting values for T_{min} and T_{max} parameters in these complex

functions (Table S2; see also below and Fig. 3). The only taxonomic group for which > 5

different functions were tested was Insecta, for which 4.0 % of all species datasets (representing

7.8 % of insects) were tested with 5-17 different development functions (Fig. 2).

The most frequently-used development function overall, and particularly among studies

of the Insecta and Arachnida, was the Linear rate function (#5), which was used in 34.9 % of all

datasets and representing 28.4 and 15.2 % of insect and arachnid datasets, respectively (Fig. 3A;

Table S2). Seven other function types were used for arachnids, and for insects a wide range of 24

further functions types were used (Fig. 3A). Functions used on insects and arachnids ranged in

complexity from relatively simple ($k = 3$) to very complex (#33, $k = 8$), but with no particular

function aside from the linear rate one (#5) predominating (Fig. 3A). Studies of Crustacea used a

more limited set of 8 functions, all but one of which (function #16, $k = 4$) were relatively simple

(#1-3 and 5-8, $k = 3$) (Fig. 3A). The Heip power function (#2) was the most commonly-used

function for Crustacea (9.4 % of all datasets, or 35.1 % of crustaceans), followed by the

Bĕlehrádek (#8, 6.6 % of datasets or 24.6 % of crustaceans) and Tauti or exponential (#7, 4.7 %

of datasets, 17.5 % of crustaceans) functions (Fig. 3A). The distribution of best functions as

concluded in reviewed studies showed a similar pattern, with the linear rate function (#5)

dominating for insects and arachnids and the Heip power function (#2) most often being

concluded best for crustacean data (Fig. 3B). Among insects and arachnids, the Brière-1 function

(#9) showed a slight tendency to be selected as best more often than other nonlinear functions, as

it was for 7.0 % of all datasets, representing 6.7 % of insects and 18.2 % of arachnids (Fig. 3B).

3.2. Results of meta-analysis – did previous studies use the "best" function for their data?

The development function(s) used to fit species datasets and/or concluded to have been

the "best" function for these data in their original studies were found, in the vast majority of

cases, not to be the best function for the data out of the 33 functions examined (Fig. 4). When

AIC_C was used to rank functions, the best function for 91.1 % of 79 retested study datasets was

concluded to be a different one than that selected in previous studies (Fig. 4A), and for 86.1 % of

datasets the actual best function was not even included in the set of all functions used in original

studies (Fig. 4B). The actual best function was found to be different from that concluded to be

best in original studies for all (100 %) arachnid and insect studies examined, and for most

(75.0 %) crustacean studies (Fig. 4A). Also, for all taxonomic groups a considerable majority of

datasets (100 % of Arachnida, 67.9 % of Crustacea, and 95.1 % of Insecta) were found to be best

fit using a function that was not used in original studies of these species (Fig. 4B).

3.3. Overall performance of different functions:

No one function was found to always be the "best" or "worst" for all reanalyzed datasets,

but some functions did tend to perform better than others (Fig. 5). Those that were ranked as the

best function by AIC_C particularly often (i.e., for ≥ 10 % of datasets) were the Heip power (#2),

hyperbola (#3), Bĕlehrádek (#8), quadratic time (#16), and 3rd order polynomial time (#19)

functions (Fig. 5). There were no clear taxonomic patterns in terms of which functions tended to

be ranked best, although function #19 did perform particularly well for insect datasets, and more

complex functions with $k > 5$ were rarely concluded to be best (Fig. 5). Of the 33 functions tested,

16 were also never concluded to be the best of those tested (Fig. 5).

There were significant interactions between the effects of taxonomic group and

development function type on overall performance of the 33 different development functions as

assessed with R^2-values ($F_{64, 2508} = 1.377$, $p = 0.026$) and by ranking functions with AIC_C ($F_{64,}$

$_{2508} = 2.362$, $p < 0.001$). Therefore differences in R^2 and rank among functions were compared

for each taxonomic group separately. In all three taxonomic groups, R^2-values (Arachnida: $F_{32,}$

$_{297} = 1.783$, $p = 0.007$; Crustacea: $F_{32, 891} = 3.721$, $p < 0.001$; Insecta: $F_{32, 1320} = 2.868$, $p < 0.001$)

and ranks significantly differed among different development functions (Arachnida: $F_{32, 297} =$

5.724, $p < 0.001$; Crustacea: $F_{32, 891} = 14.707$, $p < 0.001$; Insecta: $F_{32, 1320} = 17.823$, $p < 0.001$).

However, overall Δ_i values were found to not differ significantly among functions ($F_{32, 2508} =$

0.278, $p > 0.999$) or taxonomic groups ($F_{2, 2508} = 1.0$, $p = 0.368$), nor was there a significant

interaction between the effects of these factors on Δ_i values ($F_{64, 2508} = 0.448$, $p > 0.999$).

Differences in fit (R^2) and rankings among functions were actually very similar across the

different taxonomic groups (Fig. 6A-F; Table 2). Most function had relatively high overall

average R^2-values between 0.7 and 0.9 or higher (Fig. 6A-C), so on average, all temperature-

dependent development functions tested were able to explain the majority of variation in

observed development times. Functions with notably lower fit compared to others (lowest mean

$= 0.476$) did occur, though, and included the linear time (#6) Logan-6 (function #25), Logan-10

(#30), and W-L-D (#33) functions on arachnid and insect data (Fig. 6A, C; Table 2), and the

Ratkowsky (#27), Brière-1 (#9) and Brière-2 (#20) functions on crustacean data (Fig. 6B; Table

2). The explanatory power of development functions therefore differed by as much as ca. 10-

50% on average depending on which was used (Fig. 6A-C). Functions with lower (better) overall

ranks for all taxonomic groups included the Heip (#2), Hyperbola (#3), Bělehrádek (#8),

quadratic time (#16), and 3rd order polynomial time (#19) functions (Fig. 6D-E; Table 2), and

those with high (poor) overall ranks included those with low R^2-values described above and the

Holling Type III (#21) functions (Fig. 6D-E; Table 2). All mean Δ_i values were high (mean $\Delta_i \geq$

22.7) because each function was not selected as the best function by AIC_C at least once, and in

many cases the difference between the AIC_C values of the best function and the 2nd best function

(Δ_i) were quite large (as evidenced by the variance in Fig. 6G-I). There were therefore large

differences in the amount of information attained depending on which function was used, but no

clear pattern among functions or taxa (Fig. 6G-I).

3.4. Consequences of function choice:

All measures of prediction error, decreased fit, and increased information loss resulting

from using the best original rather than the actual best functions on species datasets were

significantly different from zero (one-sample t-tests, p < 0.05; Table 2). Development times

predicted with original studies' functions disagreed with observed developmental duration by

about 4 days or 85.9 % on-average, but could be off by as much as 132 days or 1725.5 % (Table

3). Fit (R^2) of original functions to arthropod datasets was lower by 0.091 on-average compared

with best functions (Table 3), meaning that nearly 10 % of the variation in the data would remain

unexplained if the original rather than the actual best function was used. The difference in R^2-

values between originally-used and best functions could be much greater in many cases, though,

as percent decrease in R^2 by using the best function was as high as 100 % for certain study

datasets (Table 3; Table S2). Additionally the mean Δ_i of the originally-used functions was 225

and could be as high as 3958.7 for some datasets (Table 3), indicating a substantial loss of

information relative to the actual best function as determined in the present study.

3.5. Range of temperature tested in studies versus function performance:

Thermal ranges tested in original studies differed significantly among studies of different

taxonomic groups ($F_{2, 96} = 16.533$, $p < 0.001$; see also Table S2). Interestingly, studies of

crustaceans tended to be carried out over significantly smaller thermal ranges (mean ± 95% C.I.

$= 11.9 ± 0.8$ °C, range $= 6.0$-22.5°C) than those of arachnids ($20.0 ± 1.1$ °C, range $= 12.5$-

30.0°C; Tukey's HSD test, $p < 0.001$) or insects ($17.5 ± 1.2$ °C, range $= 8.0$-38.0; Tukey's HSD

test, $p < 0.001$) implying notable differences in the way thermal effects are investigated in these

taxa; thermal ranges studied for Insects and Arachnids did not differ, however (Tukey's HSD test,

$p = 0.328$).

The range of temperatures examined in previous studies was significantly correlated with

performance (ranking) of 13 of the 33 development functions tested in this study (Fig. 7). With

one exception (function # 30), all functions for which rankings by AIC_C were significantly and

positively correlated with temperature range were those (#1, 2, 5-7, 13, 15, and 16) that had

fewer parameters ($k ≤ 4$) (Fig. 7). This means that these relatively simpler functions tended to do

more poorly (higher value = poorer rank, further from 1) when larger thermal ranges were tested.

Conversely, rankings of more complex functions with $k = 5$ (#20, 24, and 27) or $k = 7$ (#32) were

negatively correlated with temperature range (Fig. 7), meaning that these functions performed

better (lower value = better rank, closer to 1) when studies tested a wider range of temperatures.

4. Discussion:

4.1. Use and performance of different development functions in previous studies:

     Modeling functional relationships between temperature and life history characters is an essential component of studying the biology of poikilothermic organisms (Angilletta Jr., 2006; Bělehrádek, 1935; Papnikolaou et al., 2013; Shi and Ge, 2010). Predictions of generation times (Huntley and Lopez,1992), timing of seasonal events (Bayoh and Lindsay, 2003), dispersal potential (de Rivera et al., 2007), and recruitment to adult populations of Arthropoda (Aiken and Waddy, 1986; Caddy, 1986) produced by such modeling efforts are thus sensitive to the types of temperature-dependent larval development functions incorporated in these. Development times of arthropod larvae predicted for the same species and temperatures by different function types can differ substantially, which has important impacts on predictions made. Using the best possible function to represent a given species' and/or study's dataset should thus be a crucial component of the study of temperature-dependent arthropod larval development, which should precede reporting and use of the results of such studies in models. However, in the present study this important step was found to be largely bypassed by the majority of studies. Particular function types tended to be used more often than others for particular taxonomic groups with little or no clear justification for the choice made, while consideration of alternative function types was rarely reported in published papers. In most cases the function used in original published studies was not actually the best one for the datasets presented. Further, fitting these data with the best model resulted in better fit, less disagreement between predicted and observed development times, decreased information loss, and presumably also better predictive ability. These results demonstrate that development function choice is an important but often-ignored

step in research on arthropod larval development, which should be given greater consideration in

future studies.

Choosing one particular development function might have some justification if any

function(s) could be said to be better overall than others. In the present study, no single function

type was found to be the best or worst, although some did tend to perform better or worse than

others (see Fig. 5, 6 in Results). Functions that performed well overall might be recommended as

good starting points for fitting development data, and those that did poorly overall could

conversely be used with caution. (Table S1). Also, the more complex functions with high k-

values and including T_{min} and T_{max} parameters, which performed poorly overall, did somewhat

better on insect data than for other taxa and actually was among the best models for some insect

species datasets (Table S2). Therefore, it is difficult to make general statements about which

function is always best to use; this must rather be assessed on a case-by-case basis, for each

species and study. Results in this study showed that not using the best function for a given

dataset can result in very different predicted development times, which could lead to very

different (and potentially erroneous) inferences and predictions of species biology by modeling

studies (Miller et al., 1998; Miller et al., 2006; Quinn, 2014; Reitzel et al., 2004). The practice

among many fields of study has been to fit data with a development function type that has been

used in previous studies on similar species; for example, the frequent use of the Bĕlehrádek

function on copepod crustaceans (Anger, 2001; Corkett and McLaren, 1970; Hamasaki et al.

2009) or linear rate + complex function(s) on insects and arachnids (Golizadeh and Zalucki,

2012; Shi and Ge, 2010; Smits et al., 2003; Table S2). However, based on results of the present

study this practice should be discontinued.

4.2. Importance of temperature range tested to function performance:

An interesting finding in the present study was that overall performance of several

function types was correlated with the range of temperatures tested in original published studies.

Specifically, as the range of temperatures tested increased performance (i.e., likelihood to be

ranked as the best model) of the simpler functions examined decreased while that of more

complex functions increased. This result does make sense, however, if one considers the "real"

nature of temperature-biological rate relationships. Because the actual performance of the

enzymes mediating larval development most certainly have upper and lower functional threshold

temperatures, beyond which development cannot progress (Brière et al., 1999; Quinn and

Rochette, 2015; Somero, 2004), one can assume that for most species the "true" relationship

between temperature and development time resembles the Brière-2 function, or similar complex

asymmetrical curves (e.g., Huey and Stevenson, 1979; Shi and Ge, 2010). A study carried out

over a very wide range of temperatures should be able to approach or exceed thermal thresholds

and therefore identify these limiting temperatures, and thus be best explained by a complex

function. However, if one carries out their study over a more narrow thermal range, they will

only be able to observe a certain section of the development curve, which could be located

relatively far from one or both threshold temperatures. This could result in the observed

temperature-development data having a distinctly linear, quadratic, or power function-like shape,

such that one of these alternative, simpler functions would be identified as the "best" function for

the data over this specific range. Indeed, this seems to have been the case in several of the

datasets examined, in which thermal ranges and/or sample sizes were relatively small (e.g.,

Quinn et al., 2013; Corkett and McLaren, 1970; Hamasaki et al., 2009; Carlotti et al., 2007;

Table S1, S2) and the best function was determined to be one of the simpler functions, such as

the linear time (#6) or Heip power (#2) function, even though these should be the least-realistic

(Bělehrádek, 1935; Brière et al., 1999; Somero, 2004). Importantly, when this occurs the best

function will be the one that provides the most informative description of development times

over a very specific range of temperatures, but its performance is likely to degrade if

extrapolation beyond this range is attempted. Ultimately the "best" function should be of a

complex form resembling functions with T_{min} and $Tmax$ parameters, but most studies, especially

of Crustacea, are not conducted over sufficiently wide temperature ranges to be allow good

estimates of such functions' parameters to be derived.

The vast majority of insects and arachnids have terrestrial and/or freshwater aquatic

habitats, in which temporal variability in air and water temperatures can be very large (Pakyari et

al., 2011; Sanchez-Ramos et al., 2007; Stavrinides et al. 2010). As a result, the likelihood of

these organisms and their larvae being exposed to extreme temperatures exceeding thresholds for

moulting, development, and/or survival can be high. Conversely, many crustaceans (and all of

those examined in the present review; Table S2) inhabit the marine realm as larvae and/or adults

(Paul and Paul, 1999; Roberts et al., 2012; Thompson, 1982). While it is not impossible that

marine crustacean larvae could encounter temperatures too high for development or survival (e.g.,

such warm extremes could occur in shallow coastal areas, highly-stratified water columns,

intertidal zones at low tide, or more generally due to future climate change; Caffara et al., 2012;

Quinn and Rochette, 2015), they are thought to be far more likely to encounter lower limiting

temperatures, especially in the deeper ocean or temperate regions (Hartnoll, 1982; MacKenzie,

1988; Quinn, 2016). This perceived difference in limiting temperatures appears to have lead

studies on temperature-dependent development in these groups along different paths. Studies of

insects and arachnids often use very wide temperature ranges with the intent of capturing lower

and upper limiting temperatures for development in their species because these physiological

limits are known to be essential to modeling these species in their natural environments (Shi and

Ge, 2010; Smits et al., 2003). Studies of crustaceans, to the contrary, tend to be limited to more

narrow thermal ranges deemed "ecologically-relevant" (i.e., likely to be encountered by the

species in nature); occasionally these include lower limits, but in general physiological limits,

especially upper ones, are rarely sought (Hartnoll, 1982; Quinn, 2016).

While there is certainly logic behind the use of narrower, more-relevant thermal ranges in

studies of Crustacea, this approach still has potential to result in errors for two main, related

reasons. First, the type of development function used changes predicted development times both

at and between (interpolation) observed temperatures and especially outside of these

(extrapolation) (Angilletta Jr., 2006; Campbell et al., 1974; Quinn and Rochette, 2015). Second,

thermal development limits actually change the shape of the "real" and estimated (i.e., fitted by

regression) development curve, for example by decreasing its curvature when lower and upper

limits are further apart and increasing curvature when these are closer together (Bĕlehrádek,

1935; Brière et al., 1999; Shi and Ge, 2010; personal observations by author). All else being

equal, these difference in curvature can result in very different development times at the same

temperatures. As a result, it is important to know the physiological limits of a given species when

modeling its development (Quinn, 2016). Even if a species rarely encounters temperatures close

to these limits, development times calculated at intermediate temperatures will be impacted by

the values of these limits; if one ignores these limits and uses a different function type, or

attempts to estimate limits by extrapolation from a narrow thermal range, there is great potential

for erroneous predictions of development times to be made.

4.3. Discussion of potential limitations and next steps:

In the present review, published studies on arthropod larvae were obtained through

literature searches through Web of Science (Thompson Reuters, 2015). These searches were by

no means comprehensive – many other studies of temperature-dependent development of

arthropod larvae exist that were not indexed in this search engine – but it was extensive and did

provide a good sample of such studies encompassing many different years, regions, and

arthropod taxa (Table S2). This sample of the relevant literature was thus appropriate and useful

for the purposes of the present review of development function usage and performance. An

expanded search using additional search tools in a future study could obtained data for other

taxonomic groups within the Arthropoda (e.g, myriapods, more arachnids, other orders of

Crustacea and Insecta, etc.); indeed, an Insect Developmental Database has been created by

Nietschke et al. (2007) that could be used to obtain considerably more insect data for reanalyses.

Performance of development functions on data from species outside of the arthropod phylum

(e.g., molluscs: de Severyn et al., 2000; nematodes: Jenkins et al., 2006; Singh and Sharma,

1994; urochorates: Kang et al., 2009; vertebrates: McLaren and Cooley, 1972; Miller et al. 2006)

may also be attempted, and reveal additional patterns in study design, taxonomy, and function

usage of interest. However, overall patterns and conclusions of the present study would likely

hold true. To simplify analyses, this study conducted analyses on total larval development time

data rather than on individual larval stages. However, in most species development time of each

larval stages has a distinct response to temperature, requiring different developmental equations

to be derived for each stage (Corkett and McLaren, 1970; Hartnoll, 1982). Often survival to and

through later larval stages is very low, so power to fit more complex functions to later-stage

development data can be limited. As a result, the best function can potentially differ among

larval stages of the same species, in the same study; indeed, this was noted for American lobster

data in the present study (data not shown). A future study should investigate stage-specific

changes in the "best" development function(s), to confirm whether such patterns could impact

the overall performance of different function types, prediction error, and so on. However, it

would make mathematical sense for similar patterns to be found through such a detailed review

to those noted in the present study, given that similar factors (e.g., sample sizes and thermal

ranges tested) would impact function performance. There are also countless other development

function types in existence which were not included in the present study (e.g., Angilletta Jr.,

2006; Schoolfied et al., 1981; Shi and Ge, 2010). It is possible that one or more of the functions

not examined herein could actually be the closest to "real" development relationships and/or

perform better overall than all others. However, findings of this review that one or few functions

are used by most studies, the best function was not the one used in most original published

studies, and that using the non-best function results in poorer predictions would not change

through consideration of such additional functions.

In this study function performance was assessed mainly in terms of fit (R^2 and observed

versus predicted values) and information loss (AIC_C and Δ_i). These gave good indications of how

appropriate each function and its parameter estimates were for particular datasets (e.g., how well

sample sizes supported estimation of more complex functions). However, future studies could

take more thorough approaches to assessing predictive ability of different functions. One

approach to be used in the future to assess function performance could be cross-validation

(Picard and Cook, 1984; Anderson, 2008). Even better would be actually testing development

functions on new data, for example by predicting development times for a particular species at

different temperatures using different functions derived in a prior study, and then measuring new

development times and comparing these to predictions. If future studies attempted this, very

thorough tests and new evidence in favour of one function or another might be obtained.

<u>4.4. Implications of development time predictions based on different functions:</u>

Differences in development times predicted for the same species and temperature among

development functions have potential to impact various types of predictions relevant to arthropod

biology and ecology. Larval survival is usually inversely related to larval duration, such that

slower development results in fewer potential recruits to adult populations (Reitzel et al., 2004;

Roberts et al., 2012). Most life history and bio-physical models account for this by reducing

larval numbers in simulated cohorts by a certain percentage at each model time step, resulting in

substantial, exponential losses per each additional step spend in larval development (e.g., Miller

et al., 1998; Quinn, 2014). In many crustaceans, the larval phase of the life cycle is the main

dispersive phase (Anger, 1984; de Rivera et al., 2007; MacKenzie, 1988). Lengthening the larval

developmental period of such larvae can dramatically alter how far and to where larvae drift in

simulations with ocean currents; for example, simulations by Quinn (2014) showed that slowing

larval development (and thus lengthening drift time) of American lobster larvae by 60 % could

result in increased drift distances of larvae by up to ca. 500 km. If drift were overestimated in

such models, for instance due to use of inappropriate development functions, then the degree of

population mixing would be overestimated and an incorrect estimate of population structure

obtained; underestimation by the same means could also lead to considerable errors. Likewise if

dispersal ability of larvae of an invasive species, such as the green crab *Carcinus maenas* (L.) (de

Rivera et al., 2007), were underestimated in this way potential invasions to new regions that

could be predicted may be missed. Using development functions to estimate the timing of

seasonal peaks in abundance of disease vectors (Bayoh and Lindsay, 2003), agricultural pests

(Campbell et al., 1994; Easterbrook et al., 2003; Stavrinides et al., 2010), or species that serve as

important food sources to others (e.g., copepod secondary productivity in the ocean: Carlotti et

al., 2007) also depends on being able to make good estimates of larval development. For many

species very small differences in development time similar to the difference in "errors" of best

and original studies' functions are enough to dramatically alter the nature and implications of

modeled predictions (e.g., ≤ 1-5 days; Gadino and Walton, 2012). The type of development

function used can thus have large impacts on predictions, so it is important that studies attempt to

find the best model for their data. Importantly, much research is now being initiated to assess

how future climate change will impact many species, including arthropods and their larvae

(Caffara et al., 2012; Quinn and Rochette, 2015). Use of non-best development functions within

such research clearly could result in erroneous predictions as well and so should be avoided.

4.5. Recommendations and conclusions:

Based on results of this study, it is recommended that future studies examining effects of

temperature on development of arthropod larvae consider and attempt to fit multiple alternative

development function types to their data to determine the best way to model their results and

report that this was attempted. No one function type is better or worse overall, but the range of

temperatures to be considered and potential use of results (e.g., for extrapolation to temperatures

not observed) can be used as a guide when deciding which functions are most likely to provide

good representations of data. In general simpler functions could provide better descriptions of

development observed over relatively narrow thermal ranges, but provide poor extrapolation

ability. Conversely, wider ranges encompassing lower and/or upper limiting temperatures for

development can support more complex functions, which potentially resemble more closely true

enzymatic and biological thermal performance curves (Brière et al., 1999; Somero, 2004) and

may allow modeling over all temperatures potentially encountered. Studies on crustaceans in

particular should be conducted over wider thermal ranges in the future so that limiting

temperatures of these species can be identified and more complex, presumably realistic

development functions reliably fit to data for these organisms. Considering different potential

function types to find the best for each dataset should lead to better predictions of larval

development times in support of subsequent research on these important species.

Acknowledgements:

Thanks are due to Rémy Rochette, Joël Chassé, Jeff Houlahan, and Heather Hunt for advice and

guidance during this project, and the University of New Brunswick, Saint John Campus, for

providing resources that made the review and analyses possible. The author also thanks two

anonymous reviewers for providing comments that improved the scope and quality of the

manuscript.

References:

Aiken, D.E., Waddy, S.L., 1986. Environmental influence on recruitment of the American

lobster, *Homarus americanus*: a perspective. Can. J. Fish. Aquat. Sci. 43, 2258-2270.

Akaike, H., 1973. Information theory as an extension of the maximum likelihood principle, in:

Petrov, B.N., Csaki, F. (Eds.), Second International Symposium on Information Theory.

Akademiai Kiado, Budapest, pp. 267-281.

Al-Saffar, Z.Y., Grainger, J.N.R., Aldrich, J., 1996. Temperature and humidity affecting

689   development, survival and weight loss of the pupal stage of *Drosophila melanogaster*,

690   and the influence of alternating temperature on the larvae. J. Therm. Biol. 21, 389-396.

691   Analytis, S., 1977. Über die relation zwischen biologischer entwicklung und temperatur bei

692   phytopathogenen Pilzen. Phytopath. Z. 90, 64-76.

693   Analytis, S., 1981. Relationship between temperature and development times in phytopathogenic

694   fungus and in plant pests: a mathematical model. Agric. Res. (Athens). 5, 133-159.

695   Anderson, D.R., 2008. Model Based Inference in the Life Sciences: A Primer on Evidence.

696   Springer Science+Business Media, LLC, New York.

697   Anger, K., 1983. Temperature and the larval development of *Hyas araneus* L. (Decapoda:

698   Majidae); extrapolation of laboratory data to field conditions. J. Exp. Mar. Biol. Ecol. 69,

699   2013-215.

700   Anger, K., 1984. Development and growth in larval and juvenile *Hyas coarctatus* (Decapoda,

701   Majidae) reared in the laboratory. Mar. Ecol. Prog. Ser. 19, 115-123.

702   Anger, K., 1991. Effects of temperature and salinity on the larval development of the Chinese

703   mitten crab *Eriocheir sinensis* (Decapoda: Grapsidae). Mar. Ecol. Progr. Ser. 72, 103-110.

704   Anger, K., 2001. Crustacean Issues 12: The Biology of Decapod Crustacean Larvae. A.A.

705   Balkema, Rotterdam.

706   Angilletta, Jr., M.J., 2006. Estimating and comparing thermal performance curves. J. Therm.

707   Biol. 31, 541-545.

708   Bayoh, M.N., Lindsay, S.W., 2003. Effect of temperature on the development of the aquatic

709   stages of *Anopheles gambiae* sensu stricto (Diptera: Culicidae). Bull. Entomol. Res. 93,

710   375-381.

711 Bělehrádek, J., 1935. Temperature and living matter. Protoplasma Monographia, No. 8.

712   Borntraeger, Berlin.

713 Blanckenhorn, W.U., 1997. Effects of temperature on growth, development and diapause in the

714   yellow dung fly – against all the rules? Oecologia 111, 318-324.

715 Blanco, J. M., Guerrero, F., Rodríguez, V., 1995. The fate of comparisons of models in

716   temperature-dependent growth of copepods: a reply to the comments by McLaren. J.

717   Plankton Res. 17, 1391-1394.

718 Brière, J.F., Pracross, P., Rioux, A.Y., Pierre, J.S., 1999. A novel rate model of temperature-

719   dependent development for arthropods. Environ. Entomol. 28, 22-29.

720 Caddy, J.F., 1986. Modelling stock-recruitment processes in Crustacea: some practical and

721   theoretical perspectives. Can. J. Fish. Aquat. Sci. 43, 2330-2344.

722 Caffara, A., Rinaldi, M., Eccel, E., Rossi, V., Pertot, I., 2012. Modelling the impact of climate

723   change on the interaction between grapevine and its pests and pathogens: European

724   grapevine moth and powdery mildew. Agr. Ecosyst. Environ. 148, 89-101.

725 Campbell, A., Frazer, B.D., Gilbert, N., Gutierrez, A.P., Mackauer, M., 1974. Temperature

726   requirements of some aphids and their parasites. J. Appl. Ecol. 11, 431-438.

727 Campolo, O., Malacrinò, A., Laudani, F., Maione, V., Zappalà, L., Palmeri, V., 2014. Population

728   dynamics and temperature-dependent development of *Chrysomphalus aonidum* (L.) to

729   aid sustainable pest management decisions. Neotrop. Entomol. 43, 453-464.

730 Carlotti, F., Bonnet, D., Halsband-Lenk, C., 2007. Development and growth rates of

731   *Centropages typicus*. Progress in Oceanography 72, 164-194.

Cave, R.D., Schiacchetano, C., Diaz, R., 2009. Temperature-dependent development of the

cycad Aulacapsis scale, *Aulacapsis yasumatsui* (Hemiptera: Diaspididae). Florida

Entomol. 92, 578-581.

Choy, S., 1991. Embryonic and larval biology of *Liocarcinus holsatus* and *Necora puber*

(Crustacea: Brachyura: Portunidae). J. Exp. Mar. Biol. Ecol. 148, 77-92.

Cooper, B.S., Tharp II, J.M., Jernberg, I.I., Angilletta Jr., M.J., 2012. Developmental plasticity of

thermal tolerances in temperate and subtropical populations of *Drosophila melanogaster*.

J. Therm. Biol. 37, 211-216.

Corkett, C.J., 1972. Development rate of copepod eggs of the genus *Calanus*. J. Exp. Mar. Biol.

Ecol. 10, 171-175.

Corkett, C.J., McLaren, I.A., 1970. Relationships between development rate of eggs and older

stages of copepods. J. Mar. Biol. Assoc. U.K. 50, 161-168.

Coutant, C.C., Talmage, S.S., 1976. Thermal effects. J. Wat. Poll. Contr. Fed. 48, 1486-1544.

Criales, M.M., Anger, K., 1986. Experimental studies on the larval development of the shrimps

Crangon crangon and *C. allmanni*. Helgoländer Meeresun. 40, 241-265.Cummings, G.,

Fiddler, F., Vaux, D.L., 2007. Error bars in experimental biology. J. Cell Biol. 177, 7–11.

Davidson, J., 1942. On the speed of development of insect eggs at constant temperatures. Ayst. J.

Exp. Biol. Med. Sci. 20, 233-239.

Dawirs, R.R., 1985. Temperature and larval development of *Carcinus maenas* (Decapoda) in the

laboratory; predictions of larval dynamics in the sea. Mar. Ecol. Progr. Ser. 24, 297-302.

de Oliveira, S.A., Souza, B., Auad, A.M., da Silva, D.M., Souza, L.S., Carvalho, C.A., 2009.

Desenvolvimento e reprodução de *Sipha flava* (Forbes) (Hemiptera: Aphididae) em

diferentes temperaturas. Neotrop. Entomol. 38, 311-316.

de Rivera, C.E., Hitchcock, N.G., Teck, S.J., Steves, B.P., Hines, A.H., Ruiz, G.M., 2007. Larval

development rate predicts range expansion of an introduced crab. Mar. Biol. 150, 1275-

1288.

de Severeyn, Y. G., Severeyn, H., Grant, W., Reverol, Y., 2000. Effect of water temperature on

larval development of the bivalve mollusk *Tivela mactroides*: evaluation in the laboratory

and via simulation. Ecol. Model. 129, 143-151.

Du, W.-G., Hu, L.-J., Lu, J.-L., Zhu, L.-J., 2007. Effects of incubation temperature on embryonic

development rate, sex ratio and post-hatching growth in the Chinese three-keeled pond

turtle, *Chinemys reevesii*. Aquaculture 272, 747-753.

Easterbrook, M.A., Fitzgerald, J.D., Pinch, C., Tooley, J., Xu, X.-M., 2003. Development time

and fecundity of three important arthropod pests of strawberry in the United Kingdom.

Ann. Appl. Biol. 143, 325-331.

Edgar, A.J., Andrew, T.E., 1990. A simple method for fitting Bĕlehrádek's equation to

embryonic development data of zooplankton. Hydrobiologia 194, 177-181.

Eliopoulos, P.A., Kontodimas, D.C., Stathas, G.J., 2010. Temperature-dependent development of

Chilocorus bipustulatus (Coleoptera: Coccinellidae). Environ. Entomol. 39, 1352-1358.

Emmert, C.J., Mitzell III, R.F., Andersen, P.C., Frank, J.H., Stimac, J.L., 2008. Effects of

contrasting diets and temperatures on reproduction and prey consumption by

Proprioseiopsis asetus (Acari: Phytoseiidae). Exp. Appl. Acarol. 44, 11-26.

Escribano, R., Rodriguez, L., Irribarren, C., 1998. Temperature-dependent development and

growth of *Calanus chilensis* Brodsky from Northern Chile. J. Exp. Mar. Biol. Ecol. 229,

19-34.

Forster, J., Hirst, A.G., 2012. The temperature-size rule emerges from ontogenetic difference

between growth and development rates. Functional Ecol. 26, 483-492.

Forster, J., Hirst, A.G., Woodward, G., 2011. Growth and development rates have different

thermal responses. Am. Nat. 178, 668-678.

Gadino, A.N., Walton, V.M., 2012. Temperature-related development and population parameters

for *Typhlodromus pyri* (Acari: Phytoseiidae) found in Oregon vineyards. Exp. Appl.

Acarol. 58, 1-10.

Gao, G.-Z., Perkins, L.E., Zalucki, M.P., Lu, Z.-Z., Ma, J.-H., 2013. Effect of temperature on the

biology of *Acyrthosiphon gossypii* Mordvilko (Homoptera: Aphididae) on cotton. J. Pest.

Sci. 86, 167-172.

Geffen, A.J., Nash, R.D.M., 2012. Egg development rates for use in egg production methods

(EPMs) and beyond. Fish. Res. 117-118, 48-62.

Golizadeh, A., Zalucki, M.P., 2012. Estimating temperature-dependent developmental rates of

potato tuberworm, *Phthorimaea operculella* (Lepidoptera: Gelechiidae). Insect Sci. 19,

609-620.

Gonzàlez, L., Carvajal, J., 2003. Life cycle of *Caligus rogercresseyi* (Copepoda: Caligidae)

parasite of Chilean reared salmonids. Aquaculture 220, 101-117.

Greenberg, S.M., Sétamou, M., Sappington, T.W., Liu, T.-X., Coleman, R.J., Armstrong, J.S.,

2005. Temperature-dependent development and reproduction of the boll weevil

(Coleoptera: Curculionidae). Insect Sci. 12, 449-459.

Guerrero, F., Blanco, J.M., Rodríguez, V., 1994. Temperature-dependent development in marine

copepods: a comparative analysis of models. J. Plankton Res. 16, 95-103.

Haghani, M., Fathipour, Y., Talebi, A.A., Baniameri, V., 2007. Temperature-dependent

development of *Diglyphus isaea* (Hymenoptera: Eulophidae) on *Liriomyza sativae*

(Diptera: Agromyzidae) on cucumber. J. Pest Sci. 80, 71-77.

Hamasaki, K., Sugizaki, M., Dan, S., Kitada, S., 2009. Effect of temperature on survival and

developmental period of coconut crab (*Birgus latro*) reared in the laboratory. Aquaculture

292, 259-263.

Harcourt, D.C., Yee, J.M., 1982. Polynomial algorithm for predicting the duration of insect life

stages. Environ. Entomol. 11, 581-584.

Hartnoll, R.G., 1982. Growth, in: The Biology of the Crustacea, Vol. 2: Embryology,

Morphology, and Genetics. Abele, L.G. (Ed.). Academic Press, New York, pp. 111-196.

Heip, C., 1974. A comparison between models describing the influence of temperature on the

development rate of copepods. Biologisch Jaarboek Dodonaea 42, 121-125.

Herzig, A., 1983. The ecological significance of the relationship between temperature and

duration of embryonic development in planktonic freshwater copepods. Hydrobiologia

100, 65-91.

Hilbert, D.W., Logan, J.A., 1983. Empirical model of nymphal development for the migratory

grasshopper, *Melanoplus sanguinipes* (Orthoptera: Acrididae). Environ. Entomol. 12, 1-5.

Holling, C.S., 1965. The functional response of predators to prey density and its role in mimicry

and population regulation. Mem. Entomol. Soc. Can. 45, 1-60.

Huang, Z., Ren, S., Musa, P.D., 2008. Effects of temperature on development, survival,

longevity, and fecundity of the *Bemisia tabaci* Gennadius (Homoptera: Aleyrodidae)

predator, *Axinoscymnus cardilobus* (Coleoptera: Coccinellidae). Biol. Control 46, 209-

215.

822 Huey, R.b., Svenson. R.D., 1979. Integrating thermal physiology and ecology of ectotherms: a

823   discussion of approaches. Am. Zool. 19, 357-366.

824 Huntley, M.E., López, M.D.G., 1992. Temperature-dependent production of marine copepods: a

825   global synthesis. Am. Nat. 140, 201-242.

826 Ikemoto, T., Takai, K., 2000. A new linearized formula for the law of total effective temperature

827   and the evaluation of line-fitting methods with both variables subject to error. Environ.

828   Entomol. 29, 671-682.

829 Ismael, D., Anger, K., Moreira, G.S., 1997. Influence of temperature on larval survival,

830   development, and respiration in *Chasmagnathus granulata* (Crustacea, Decapoda).

831   Helgoländer Meeresun. 52, 463-475.

832 Jacas, J.A., Peña, J.E., Duncan, R.E., Ulmer, B.J., 2008. Thermal requirements of *Fidiobia*

833   *dominica* (Hymenoptera: Platygastridae) and *Haeckeliania sperata* (Hymenoptera:

834   Trichogrammatidae), two exotic egg parasitoids of *Diaprepes abbreviatus* (Coleoptera:

835   Curculionidae). BioControl 53, 451-460.

836 Jaffari, S., Fathipour, Y., Faraji, F., 2012. Temperature-dependent development of *Neoseiulus*

837   *barkeri* (Acari: Phytoseiidae) on *Tetranychus urticae* (Acari: Tetranychidae) at seven

838   constant temperatures. Insect Sci. 19, 220-228.

839 Janisch, E., 1932. The influence of temperature on the life-history of insects. Trans. Entomol.

840   Soc. Lond. 80, 137-168.

841 Jenkins, E. J., Kutz, S.J., Hoberg, E.P., Polley, L., 2006. Bionomics of larvae of

842   *Parelaphostronglyus odocoilei* (Nematoda: Protostrongylidae) in experimentally infected

843   gastropod intermediate hosts. J. Parasitol. 92, 298-305.

Kamps, D.M., 1978. The effect of temperature on the development time and brood size of

Diaptomus pallidus Herrick. Hydrobiologia 61, 75-80.

Kang, K.H., Zhang, Z.F., Bao, Z.M., Zhou, B., Han, C.B., 2009. Influence of water temperature

on spawning induction and larval development of the sea squirt *Halocynthia ritteri* (Oka,

1906). Aquaculture Res. 40, 513-518.

Karimi-Malati, A., Fathipour, Y., Talebi, A.A., 2014. Development response of *Spodoptera*

exigua to eight constant temperatures: Linear and nonlinear modeling. J. Asia-Pac.

Entomol. 17, 349-354.

Khan, M., Gregg, P., Mensah, R., 2009. Effect of temperature on the biology of *Creontiades*

dilutus (Heteroptera: Miridae). Austral. J. Entomol. 48, 210-216.

Kim, D.-S., Riedl, H., 2005. Effect of temperature on development and fecundity of the

predaceous plant bug *Deraeocoris brevis* reared on *Ephestia kuehniella* eggs. BioControl

50, 881-897.

Kim, S.C., Song, J.-H., Kim, D.-S., 2008. Effect of temperature on the development and

fecundity of the Cryptic Mealybug, *Pseudococcus cryptus*, in the laboratory. J. Asia-Pac.

Entomol. 11, 149-153.

Klein Breteler, W.C.M., Gonzalez, S.R., Schogt, N., 1995. Development of *Pseudocalanus*

elongatus (Copepoda, Calanoida) cultured at different temperatures and food conditions.

Mar. Ecol. Prog. Ser. 119, 99-110.

Koda, K., Nakamura, H., 2010. Effects of temperature on the development and survival of an

endangered butterfly, *Shijimiaeoides divinus barine* (Leech) (Lepidoptera: Lycaenidae).

Entomol. Sci. 13, 29-34.

Kontodimas, D.C., Eliopoulos, P.A., Stathas, G.J., Economou, L.P., 2004. Comparative

temperature-dependent development of *Nephus includens* (Kirsch) and *Nephus bisignatus*

(Boheman) (Coleoptera: Coccinellidae), preying on *Planococcus citri* (Risso)

(Homoptera: Pseudococcidae): evaluation of a linear and various non-linear models using

specific criteria. Environ. Entomol. 33, 1-11.

Krüger, R.F., Kirst, F.D., de Souza, A.S.B., 2010. Rate of development of forensically-important

Diptera in southern Brazil. Revista Brasileira de Entomologia 54, 624-629.

Lactin, D.J., Holliday, N.J., Johnson, D.L., Craigen, R., 1995. Improved rate model of

temperature-dependent development by arthropod. Environ. Entomol. 24, 68-75.

Lamb, R.J., Gerber, G.H., Atkinson, G.F., 1984. Comparison of developmental rate curves

applied to egg hatching data of Entomoscelis Americana Brown (Col: Chrysomelidae).

Environ. Entomol. 13, 868-872.

Lavagnini, T.C., de Freitas, S., Bezerra, A.L., 2009. Aspectos biológicos de *Chrysoperla*

raimundoi Freitas & Penny (Neuroptera, Chrysopidae). Revista Brasileira de

Entomologia 53, 629-634.

Lefebvre, F., Pasquerault, T., 2004. Temperature-dependent development of *Ophyra aenescens*

(Wiedemann, 1830) and *Ophyra capensis* (Wiedemann, 1818) (Diptera, Muscidae).

Forensic Sci. Int. 139, 75-79.

Li, D., 2002. The combined effects of temperature and diet on development and survival of a

crab spider, *Misumenops tricuspidatus* (Fabricius) (Araneae: Thomisidae). J. Therm. Biol.

27, 83-93.

Lind, M.I., Johansson, F., 2007. The degree of adaptive phenotypic plasticity is correlated with

the spatial environmental heterogeneity experienced by island populations of *Rana*

temporaria. J. Evo. Biol. 20, 1288-1297.

Lindley, J.A., 1990. Regressions for estimating development times of the pelagic larvae of

Paguridae and Porcellanidae. J. Plankton Res. 12, 673-678.

Liu, S.-S., Meng, X.-D., 1999. Modelling development time of *Myzus persicae* (Hemiptera:

Aphididae) at constant and natural temperatures. Bull. Entomol. Res. 89, 53-63.

Logan, D.P., Kettle, C.G., 2007. Temperature-dependent development and distribution in the soil

profile of pupae of greyback canegrub *Dermolepida albohirtum* (Waterhouse)

(Coleoptera: Scarabaeidae) in Queensland sugarcane. Austral. J. Entomol. 46, 17-22.

Logan, J.A., 1988. Toward an expert system for development of pest simulation models. Environ.

Entomol. 17, 359-376.

Logan, J.A., Wollkind, D.J., Hoyt, S.C., Tanigoshi, K., 1976. An analytical model for description

of temperature dependent rate phenomena in arthropods. Environ. Entomol. 5, 1133-1140.

Luypaert, G., Witters, J., Huylenbroeck, J.V., Maes, M., De Riek, J., De Clercq, P., 2014.

Temperature-dependent development of the broad mite *Polyphagotarsonemus latus*

(Acari: Tarsonemidae) on *Rhododendron simsii*. Exp. Appl. Acarol. 63, 389-400.

MacKenzie, B.R., 1988. Assessment of temperature effects on interrelationships between stage

durations, mortality, and growth in laboratory-reared *Homarus americanus* Milne

Edwards larvae. J. Exp. Mar. Biol. Ecol. 116, 87-98.

Marchioro, C.A., Forester, L.A., 2011. Development and survival of the diamondback moth,

Plutella xylostella (L.) (Lepidoptera: Yponomeutidae) as a function of temperature:

909   effects of number of generations in tropical and subtropical regions. Neotrop. Entomol.

910   40, 533-541.

911   Matsuda, H., Yamakawa, T., 1997. Effects of temperature on growth of the Japanese spiny

912   lobster, *Panulirus japonicus* (V. Siebold) phyllosomas under laboratory conditions. Mar.

913   Freshwater Res. 48, 791-796.

914   Mazzei, K.C., Newman, R.M., Loos, A., Ragsdale, D.W., 1999. Development rates of the native

915   milfoil weevil, *Eurhychiopsis lecontei*, and damage to Eurasian watermilfoil at constant

916   temperatures. Biol. Control 16, 139-143.

917   McLaren, I.A., 1963. Effects of temperature on growth of zooplankton and the adaptive value of

918   vertical migration. J. Fish. Res. Board Can. 20, 685-727.

919   McLaren, I.A., 1995. Temperature-dependent development in marine copepods: comments on

920   choices of models. J. Plankton Res. 17, 1385-1390.

921   McLaren, I.A., Cooley J.M., 1972. Temperature adaptation of embryonic development rate

922   among frogs. Physiol. Zool. 45, 223-228.

923   McLaren, I.A., Corkett, C.J., Zillioux, E.J., 1969. Temperature adaptations of copepod eggs from

924   the Arctic to the Tropics. Biol. Bull. 137, 486-493.

925   McQuaid, C.D., Branch, G.M., 1985. Trophic structure of rocky intertidal communities: response

926   to wave action and implications for energy flow. Mar. Ecol. Prog. Ser. 22, 153-161.

927   Mehrnejad, M.R., 2012. Biological parameters of *Elasmus nudus* (Hymenoptera, Eulophidae), a

928   parasitoid of the pistachio fruit hull borer moth, *Arimania komaroffi* (Lepidoptera,

929   Pyralidae). Biocontrol Sci. Technol. 22, 659-670.

930   Menge, B.A., 1978. Predation intensity in a rocky intertidal community: effect of an algal

931   canopy, wave action and desiccation on predator feeding rates. Oecologia 34, 17-35.

Miller, C.B., Lynch, D.R., Carlotti, F., Gentleman, W., Lewis, C.V.W., 1998. Coupling of an

individual-based population dynamic model of *Calanus finmarchicus* to a cricualtion

model for the Georges Bank region. Fish. Oceanogr. 7, 219-234.

Miller, D.C.M., Moloney, C.L., van der Lingen, C.D., Lett, C., Mullon, C., Field, J.G., 2006.

Modelling the effects of physical–biological interactions and spatial variability in

spawning and nursery areas on transport and retention of sardine *Sardinops sagax* eggs

and larvae in the southern Benguela ecosystem. J. Mar. Syst. 61, 212-229.

Milne Edwards, H., 1837. Histoire naturelle des Crustacés, comprenant l'anatomie, la

physiologie et la classification de ces animaux. Volume 2. Librairie encyclopédique de

Roret, Paris.

Mironidis, G.K., Savopoulou-Soultani, M., 2009. Development, survival and growth rate of the

Hyposoter didymator-Helicoverpa armigera parasitoid-host system: effect of host instar

at parasitism. Biol. Control 49, 58-67.

Mobarakian, M., Zamani, A.A., Karmizadeh, J., Naghadeh, N.M., Emami, M.S., 2014.

Modelling development of *Callosobruchus maculatus* and *Anisopteromalus calandrae* at

various constant temperatures using linear and non-linear models. Biocontrol Sci. Tech.

24, 1308-1320.

Mochizuki, S., Kayaba, Y., Tanida, K., 2006. Larval growth and development in the caddisfly

Cheumatopsyche brevilineata under natural thermal regimes. Entomol. Sci. 9, 129-136.

Mols, P.J.M., van de Ende, E., Blommers, L.H.M., 1998. Embryonic and larval development of

Orthosia (Lep., Noctuidae) species used for optimizing timing of monitoring and control

in apple orchards. J. Appl. Entomol. 122, 431-439.

Morimoto, S., Imamura, T., Visarathanonth, P., Miyanoshita, A., 2007. Effects of temperature on

the development and reproduction of the predatory bug *Joppeicus paradoxus* Puton

(Hemiptera: Joppeicidae) reared on *Tribolium confusum* eggs. Biol. Control 40, 136-141.

Naish, P.J., Hartwell, S., 1988. Exponentially modified Gaussian functions – a good model for

chromatographic peaks in isocratic HPLC? Chromatographia 26, 285-296.

Newman, B.K., Papadopoulos, I., Vorsatz, J., Woolridge, T.H., 2006. Influence of temperature

on the larval development of *Upogebia africana* and *U. capensis* (Decapoda:

Thalassinidae: Upogebiidae) in the laboratory. Mar. Ecol. Prog. Ser. 325, 165-180.

Nietschke, B.S., Magarey, R.D., Borchert, D.M., Calvin, D.D., Jones, E., 2007. A developmental

database to support insect phenology models. Crop Protect. 26, 1444-1448.

Nondillo, A., Redaelli, L.R., Botton, M., Pinent, S.M.J., Gitz, R., 2008. Exigências Térmicas e

Estimativa do Número de Gerações Anuais de *Frankliniella occidentalis* (Pergande)

(Thysanoptera: Thripidae) em Morangueiro [Thermal Requirements and Estimate of the

Annual Number of Generations of *Frankliniella occidentalis* (Pergande) (Thysanoptera:

Thripidae) on Strawberry Crop]. Neotrop. Entomol. 37, 646-650.

Pakyari, H., Fathipour, Y., Enkegaard, A., 2011. Estimating development and temperature

thresholds of *Scholothrips longicornis* (Thysanoptera: Thripidae) on eggs of two-spotted

spider mite using linear and nonlinear models. J. Pest Sci. 84, 153-163.

Palmer, M.A., Coull, B.C., 1980. The prediction of development rate and the effect of

temperature for the meiobenthic copepod, *Microarthridion littorale* (Poppe). J. Exp. Mar.

Biol. Ecol. 48, 73-83.

Papanikolaou, N.E., Milonas, P.G., Kontodimas, D.C., Demiris, N., Matsinos, Y.G., 2013.

Temperature-dependent development, survival, longevity and fecundity of the

aphidophagous ladybird beetle, *Propylea quatuordecimpunctata* (Coleoptera:

Coccinellidae). Ann. Entomol. Soc. Am. 106, 228-234.

Park, H.-H., Ahn, J.-J., Park, C.-G., 2014. Temperature-dependent development of

Cnaphalocrocis medianalis Guenée (Lepidoptera: Pyralidae) and their validation in semi-

field conditions. J. Asia-Pac. Entomol. 17, 83-91.

Paul, A.J., Paul, J.M., 1999. Development of larvae of the golden king crab *Lithodes aequispinus*

(Anomura: Lithodidae) reared at different temperatures. J. Crustac. Biol. 19, 42-45.

Picard, R., Cook, D., 1984. Cross-validation of regression models. J. Am. Stat. Assoc.79, 575-

583.

Prasad, Y.G., Prabhakar, M., Sreedevi, G., Rao, G.R., Venkateswarlu, B., 2012. Effect of

temperature on development, survival and reproduction of mealybug, *Phenacoccus*

solenopsis Tinsley (Hemiptera: Pseudococcidae) on cotton. Crop Protect. 39, 81-88.

Quinn, B.K., 2014. Assessing potential influence of larval development time and drift on large-

scale spatial connectivity of American lobster (*Homarus americanus*). M.Sc. Thesis,

University of New Brunswick, Saint John, NB.

Quinn, B.K., 2016 (in press). Threshold temperatures for performance and survival of American

lobster larvae: a review of current knowledge and implications to modeling impacts of

climate change. Fish. Res. http://dx.doi.org/10.1016/j.fishres.2016.09.022.

Quinn, B.K., Rochette, R., 2015. Potential effect of variation of water temperature on

development time of American lobster larvae. ICES J. Mar. Sci. 72 (Supplement 1), i79-

i90.

Quinn, B.K., Rochette, R., Ouellet, P., Sainte-Marie, B., 2013. Effect of temperature on

development rate of larvae from cold-water American lobster (*Homarus americanus*). J.

Crustac. Biol. 33, 527-536.

Ramalho, F.S., Wanderley, P.A., Malaquias, J.B., Rodrigues, K.C.V., Souza, J.V.S., Zanuncio,

J.C., 2009. Temperature-dependent development rates of *Bracon vulgaris*, a parasitoid of

boll weevil. Phytoparasitica 37, 17-25.

Ratkowsky, D.A., Lowry, R.K., McMeekin, T.A., Stokes, A.N., Chandler, R.E., 1983. Model for

bacterial culture growth rate throughout the entire biokinetic temperature range. J.

Bacteriol. 154, 1222-1226.

Reitzel, A.M., Miner, B.G., McEdward, L.R., 2004. Relationships between spawning date and

larval development time for benthic marine invertebrates: a modeling approach. Mar.

Ecol. Prog. Ser. 280, 13-23.

Roberts, S.D., Dixon, C.D., Andreacchio, L., 2012. Temperature dependent larval duration and

survival of western king prawn, *Penaeus (Melicertus) latisulcatus* Kishinouye, from

Spencer Gulf, South Australia. J. Exp. Mar. Biol. Ecol. 411, 14-22.

Rothlisberg, P.C., 1979. Combined effects of temperature and salinity on the survival and growth

of the larvae of *Pandalus jordani* (Decapoda: Pandalidae). Mar. Biol. 54, 125-134.

Roy, M., Brodeur, J., Cloutier, C., 2002. Relationship between temperature and development rate

of *Stethorus punctillum* (Coleoptera: Coccinellidae) and its prey *Tetranychus mcdaniali*

(Acarina: Tetranychidae). Environ. Entomol. 31, 177-187.

Sànchez-Ramos, I., Àlvarez-Alfageme, F., Castañera, P., 2007. Development and survival of the

cheese mites, *Acarus farris* and *Tyrophagus neiswanderi* (Acari: Acaridae), at constant

temperatures and 90% relative humidity. J. Stored Products Res. 43, 64-72.

Schoolfield, R.M., Sharpe, P.J.H., Magnuson, C.E., 1981. Non-linear regression of biological

1022    temperature-depednent rate models based on absolute reaction-rate theory. J. Theor. Biol.

1023    88, 719-731.

1024 Schultze, K., Anger, K., 1997. Larval growth patterns in the Aesop shrimp *Pandalus montagui*. J.

1025    Crustac. Biol. 17, 471-479.

1026 Sharpe, P.J.H., DeMichele, D.W., 1977. Reaction kinetics of poikilotherms development. J.

1027    Theor. Biol. 64, 649-670.

1028 Shi, P., Ge, F., 2010. A comparison of different thermal performance functions describing

1029    temperature-dependent development rates. J. Therm. Biol. 35, 225-231.

1030 Singh, M., Sharma, S.B., 1994. Temperature effects on development and reproduction of

1031    *Heterodera cajani* on Pigeonpea. J. Nematol. 26, 241-248.

1032 Smits, N., Brière, J.-F., Fargues, J., 2003. Comparison of non-linear temperature-depednent

1033    development rate models applied to *in vitro* growth of entomopathogenic fungi. Mycol.

1034    Res. 107, 1476-1484.

1035 Somero, G.N., 2004. Adaptation of enzymes to temperature: searching for basic "strategies".

1036    Comp. Biochem. Phys. B 139, 321-333.

1037 Son, Y., Lewis, E.E., 2005. Modelling temperature-dependent development and survival of

1038    *Otiorhynchus sulcatus* (Coleoptera: Curculinonidae). Agr. Forest Entomol. 7, 201-209.

1039 Stavrinides, M.E., Lara, J.R., Mills, N.J., 2010. Comparative influence of temperature on

1040    development and biological control of two common vineyard pests (Acari:

1041    Tetranychidae). Biol. Control 55, 126-131.

1042 Stinner, R.E., Guitierez, A.P., Butler, Jr., G.D., 1974. An algorithm for temperature-dependent

1043    growth rate simulation. Can. Entomol. 106, 519-524.

Tauti, M., 1925. J. Imp. Fish Inst. Tokyo, 21, (cited in Winberg, 1971 and Guerrero et al., 1994).

Taveras, R., Hilje, L., Carballo, M., 2004. Development of *Hypsipyla grandella* (Zeller)

(Lepidoptera: Pyralidae) in response to constant temperatures [Desenvolvimento de

Hypsipyla grandella (Zeller) (Lepidoptera: Pyralidae) em diferentes temperaturas

constants]. Neotrop. Entomol. 33, 1-6.

Taylor, F., 1981. Ecology and evolution of physiological time in insects. Am. Nat. 117, 1-23.

Thessalou-Legaki, M., 1990. Advanced larval development of *Callianassa tyrrhena* (Decapoda:

Thalassinidea) and the effect of environmental factors. J. Crustac. Biol. 10, 659-666.

Thompson, B.M., 1982. Growth and development of *Pseudocalanus elongatus* and *Calanus* sp.

In the laboratory. J. Mar. Biol. Assoc. U.K. 62, 359-372.

Tofangsazi, N., Buss, E.A., Meagher, R., Mascarin, G.M., Arthurs, S.P., 2012. Thermal

requirements and development of *Herpetogramma phaeopteralis* (Lepidoptera:

Crambidae: Spilomelinae). J. Econ. Entomol. 105, 1573-1580.

Tun-Lin, W., Burkot, T.R., Kay, B.H., 2000. Effects of temperature and larval diet on

development rates and survival of the dengue vector *Aedes aegypti* in north Queensland,

Australia. Medical and Vet. Entomol. 14, 31-37.

Ulmer, B.J., Jacas, J.A., Peña, J.E., Duncan, R.E., Castillo, J., 2006. Effect of temperature on life

history of *Aprostocetus vaquitarum* (Hymenoptera: Eulophidae), an egg parasitoid of

Diaprepes abbreviatus (Coleoptera: Curculionidae). Biol. Control 39, 19-25.

Voss, S.C., Cook, D.F., Hung, W.-F., Dadour, I.R., 2012. Survival and development of the

forensically important blow fly, *Calliphora varifrons* (Diptera: Calliphoridae) at constant

temperatures. Forensic Sci. Med. Pathol. 10, 314-321.

Wang, R.-S., Lan, Z.-X., Ding, Y.-Q., 1982. Studies on mathematical models of the relationship

between insect development and temperature. Acta. Ecol. Sin. 2, 47-57. [In Chinese]

Winberg, G.G., 1971. Methods for the Estimation of Production of Aquatic Animals. Academic

Press, London.

Yao, S., Huang, Z., Ren, S., Mandour, N., Ali, S., 2011. Effects of temperature on development,

survival, longevity, and fecundity of *Serangium japonicum* (Coleoptera: Coccinellidae), a

predator of *Bemisia tabaci* Gennadius (Homoptera: Aleyrodidae). Biocontrol Sci.

Technol. 21, 23-34.

Zhao, F., Zhang, W., Hoffman, A.A., Ma, C.-S., 2014. Night warming on hot days produces

novel impacts on development, survival and reproduction in a small arthropod. J. Anim.

Ecol. 83, 769-778.

1077

1078

1079

1080

1081

1082

1083 **Electronic Supplementary Material:**

1084

1085 **Table S1.** Development function parameter estimates, R^2, and AIC_C for American lobster data

1086 plotted in Figure 1.

1087

Table S2. List of studies and species obtained for the literature review and meta-analyses,

including taxonomic grouping, function type(s) used, best function, whether the best and used

function were the same, temperature range tested, performance (R^2, AIC_C rank, and Δ_i) of each

of the 33 functions tested ('na' = study not possible to retested in meta-analysis), and

consequences of using original studies' rather than the actual best function to fit datasets in terms

of prediction error, R^2 decrease, and information loss (Δ_i).

1094

1095

1096

1097

1098

1099

1100

1101

1102

1103

1104

1105

1106   **Tables and Figures:**

1107

Figure 1. Examples of the curves formed by different development functions, when these are

used to calculate development time (y-axes, days) at different temperatures (x-axes, °C). Types

of functions plotted here are those presented in Table 1, separated into those with k-values

(number of parameters + 1) of (A) 3, (B) 4, (C) 5, or (D) 6-8. Actual functions (see Table S1)

were derived from and fitted to data for total development times (combined time to complete

larval stages I-III) of American lobster (*Homarus americanus* (H. Milne Edwards, 1837)) larvae,

as reported by Quinn et al. (2013); observed development times are plotted (squares) along with

predictions of development functions (lines). Coefficients, R^2, AIC_C, and Δ_i values for these

functions are presented in Table S1. '*' = function used by Quinn et al. (2013) to fit the data, '!'

= actual "best" function for these data, and '+' = worst function for these data.

1118

1119

1120 **Figure 2.** Usage in each development study reviewed of one or more different types of

1121 development function. The percentage (%) of species datasets obtained in initial literature review

1122 (n = 99 total, see Table S2) that were tested with each number of functions is plotted on the y-

1123 axes and broken down by taxa (gray bars = arachnids, black = crustaceans, and white = insects).

1124 For names and details of development functions (#1-33) the reader is referred to Table 1.

1125

1126

Figure 3. Usage in each development study reviewed of specific development function types.

The percentage (%) of the 99 species datasets obtained in initial literature review (see Table S2)

that were (A) tested with each type of function (i.e., all used functions) and (B) concluded to be

best represented by each type of function (i.e., best used function) is plotted on the y-axes.

Results are also broken down by taxonomic group, as in Fig. 2. For names and details of

development functions (#1-33) the reader is referred to Table 1.

1133

Figure 4. Percentage (%) of species datasets re-analyzed for meta-analysis (n = 79) for which the development function found to be best for the data (lowest AIC_C value, ranking based on AIC_C = 1) in this study (i.e., actual best function) (A) agreed or not with the best function as used in its original published study or (B) was among the set of all function(s) used within its original published study. Results are also broken down by taxonomic group, as in Fig. 2. For names and details of development functions (#1-33) the reader is referred to Table 1.

Figure 5. Percentage (%) of species datasets re-analyzed for meta-analysis (n = 79) for which each development function type was found to be best for the data (lowest AIC_C value, ranking based on AIC_C = 1) in this study. Results are broken down by taxonomic group, as in Fig. 2. For names and details of development functions (#1-33) the reader is referred to Table 1.

Figure 6. Overall performance of different development functions (x-axes) assessed based on (A-C) R^2 values, (D-F) ranking based on AIC_C values, and (G-I) Δ_i values (y-values). Function performance was assessed separately for each taxon: (A, D, G) arachnids (n = 10 species for each function), (B, E, H) crustaceans (n = 28), and (C, F, I) insects (n = 41). Possible rankings ranged from 1 ("best" model) to 33 ("worst"). Values plotted are mean values per function and taxonomic group taken across all species datasets within that group ± 95 % C.I.s. For names and details of development functions (#1-33) the reader is referred to Table 1. Results of post-hoc comparisons among functions with Tukey's HSD test for each taxonomic group are presented in Table 2.

1159

Figure 7. Pearson's correlation coefficient values (R) calculated between the ranges of

temperatures (°C) tested in all original studies (n = 99) and rank (out of 33) of development

functions based on AIC_C for each species dataset. Statistical significance of correlation

coefficients is indicated by labels above each plotted point as follows: * $p \leq 0.05$, ** $p \leq 0.01$,

*** $p < 0.001$; non-significant ($p > 0.05$) results are not labelled. For names and details of

development functions (#1-33) the reader is referred to Table 1.

1166

1167

1168

Table 1. Names and forms of the 33 different development functions examined in this study. Functions are arranged and numbered

first in order of their value of k (number of parameters + 1, used in AIC_C analyses), and then alphabetically by their common name. In

all equations D is development time (in days), T is temperature (°C), A, B, C, F, a, b, c, d, f, and g are fitted constants. Additional

fitted constants in some equations with biological meaning and constrained values are as follows: T_{min} = biological minimum

temperature; T_{max} = biological maximum temperature; T_{opt} = temperature at which development rate is maximized; K = thermal

constant, or the number of degree days required to complete development (Kontodimas et al., 2004); ΔT = range of temperatures over

which development occurs; D_{min} = minimum development time. Functions that are fitted to development rate (1/D) are here

represented in their inverted form, from which D can be directly calculated. For clarity lowercase letters are used for functions fitted to

development rate, and uppercase letters are used for functions fitted directly to time (D).

1178

Function	k	Equation	Reference(s)
(1) Arrhenius	3	$D = 1/(a*EXP(b/(T)))$	Guerrero et al., 1994
(2) Heip or power	3	$D = A*T^B$	Anger, 2001; Heip, 1974; Guerrero et al., 1994
(3) Hyperbola	3	$D = A/(T-T_{min})$	Hamasaki et al., 2009; Heip, 1974
(4) Ikemoto and Takai	3	$D*T = K+T_{min}*D$ $D = 1/((T-T_{min})/K)$	Ikemoto and Takai, 2000; Papanikolaou et al., 2013

(5) Linear rate, or thermal summation	3	$D = 1/(a+b*t)$ $D = K/(T-T_{min})$ $T_{min} = -a/b$ $K = 1/b$	Campbell et al., 1974; Guerrero et al., 1994; Kontodimas et al., 2004
(6) Linear time, or sum of temperatures	3	$D = A+B*T$ $T_{max} = -A/B$	Guerrero et al., 1994; Winberg, 1971
(7) Tauti or exponential	3	$D = A*EXP(B*T)$ $D = 1/(a*EXP(b*t))$ $A = 1/a, B = -b$	Anger, 2001; Guerrero et al., 1994; Tauti, 1925
(8) Bĕlehrádek	4	$D = A*(T-T_{min})^B$	Anger, 2001; Bĕlehrádek, 1935; Guerrero et al., 1994; Heip, 1974; McLaren, 1963
(9) Brière-1	4	$D = 1/(a*T*(T-T_{min})*SQRT(T_{max}-T))$	Brière et al., 1999; Shi and Ge, 2010
(10) Gaussian	4	$D = 1/(a*EXP(-0.5*((T-b)/c)^2))$	Shi and Ge, 2010; Taylor, 1981
(11) Kontodimas-16	4	$D = 1/(a*(T-T_{min})^2*(T_{max}-T))$	Kontodimas et al., 2004

(12) Lactin-1	4	$D = 1/(EXP(a*T)-EXP(a*T_{max}*((T_{max}-T)/ \Delta T)))$	Lactin et al., 1995; Zhao et al., 2014
(13) Modified Arrhenius	4	$D = 1/(a*EXP(b/(T-T_{min})))$	Guerrero et al., 1994
(14) Pradham-Taylor, or Taylor	4	$D = 1/(a*EXP(-0.5*((T-T_{opt})/b)^2))$	Golizadeh and Zalucki, 2012; Mobarakian et al., 2014; Roy et al., 2002
(15) Quadratic rate	4	$D = 1/(a*T^2+b*T+c)$	Mehrnejad, 2012
(16) Quadratic time	4	$D = A*T^2+B*T+C$	de Oliveira et al., 2009; Quinn et al., 2013
(17) Sigmoid, logistic, or Davidson	4	$D = 1/(c/(1+EXP(a-b*T)))$	Davidson, 1942; Kontodimas et al., 2004
(18) 3rd order polynomial rate, or Harcourt	5	$D = 1/(a*T^3+b*T^2+c*T+d)$	Harcourt and Yee, 1982; Kontodimas et al., 2004
(19) 3rd order	5	$D = A*T^3+B*T^2+C*T+F$	Bayoh and Lindsay, 2003; Harcourt and Yee, 1982

polynomial time

(20) Brière-2	5	$D = 1/(a*T*(T-T_{min})*(T_{max}-T)^{(1/b)})$	Brière et al., 1999; Shi and Ge, 2010
(21) Holling Type III, or Hilbert and Logan	5	$D = 1/(a*((T^2/(T^{\wedge 2}+b^2))*EXP((T_{max}-T)/\Delta T)))$	Hilbert and Logan, 1983; Holling, 1965; Kontodimas et al., 2004
(22) Janisch	5	$D = (D_{min}/2)*(EXP(K*(T-T_{max}))+EXP(-A*(T-T_{max})))$	Analytis, 1981; Janisch, 1932; Kontodimas et al., 2004
(23) Lactin-2	5	$D = 1/(EXP(a*T)-EXP(a*T_{max}*((T_{max}-T)/\Delta T))+b)$	Kontodimas et al., 2004; Lactin et al., 1995
(24) Lamb, or Taylor non-symmetric Gauss	5	$D = 1/(a*EXP(-0.5*((T-T_{opt})/T_{min})^2))$, if $T \leq T_{opt}$ $D = 1/(a*EXP(-0.5*((T-T_{opt})/T_{max})^2))$, if $T > T_{opt}$	Kontodimas et al., 2004; Lamb et al., 1984; Taylor, 1981
(25) Logan-6 (or Logan-1)	5	$D = 1/(a*(EXP(b*T)-EXP(b*T_{max}-((T_{max}-T)/\Delta T))))$	Logan et al., 1976; Shi and Ge, 2010; Kontodimas et al., 2004
(26) Modified Gaussian	5	$D = 1/(a*EXP(-0.5*(ABS(T-b)/c)^d))$	Angilletta Jr., 2006; Naish and Hartwell, 1988; Shi and Ge, 2010
(27) Ratkowsky	5	$D = 1/((a*(T-T_{min})*(1-EXP(b*(T-T_{max}))))^2)$	Ratkowsky et al., 1983; Shi and Ge, 2010

(28) Stinner	5	$D = 1/(c/(1+EXP(a+b*T)))$, if $T \leq T_{opt}$ $D = 1/(c/(1+EXP(a+b*(2*T_{opt}-T))))$, if $T > T_{opt}$	Kontodimas et al., 2004; Stinner et al., 1974
(29) Analytis	6	$D = 1/(a*(T-T_{min})\wedge b*(T_{max}-T)^c)$	Analytis 1977, 1981; Kontodimas et al., 2004
(30) Logan-10 (or Logan-2)	6	$D = 1/(a*((1/(1+K*EXP(-b*T)))-EXP(-(T_{max}-T)/\Delta T)))$	Kontodimas et al., 2004; Logan, 1988; Logan et al., 1976; Shi and Ge, 2010
(31) Performance	6	$D = 1/(m*(T-T_{min})*(1-EXP(K*(T-T_{max}))))$	Huey and Stevenson, 1979; Shi and Ge, 2010
(32) Sharpe and DeMichele	7	$D = 1/(T*(EXP(a-(b/T)))/(1+EXP(c-(d/T))+EXP(f-(g/T))))$	Kontodimas et al., 2004; Schoolfield et al., 1981; Sharpe and DeMichele, 1977
(33) Wang-Lan-Ding (W-L-D)	8	$D = 1/((K*(1-EXP(-a*(T-T_{min})))*(1-EXP(b*(T-T_{max}))))/(1+EXP(-c*(T-d))))$	Shi and Ge, 2010; Wang et al., 1982

1179

1180

Table 2. Results of post-hoc comparisons of fit (R^2) and rank (based on comparisons of AIC_C

values) among different development functions within each arthropod subphylum using Tukey's

HSD test. Functions with different letters in a particular column had significantly different values

of R^2 or rank; comparisons were not made among subphyla (i.e., among columns). Functions

labelled with the letter 'A' had the overall best performance (highest mean R^2, rank closest to 1)

while letters from B to M indicate progressively poorer function performance. Means of the R^2

and rank data compared in this table are plotted in Fig. 6. Post-hoc comparisons were not made

of Δ_i values because these did not differ significantly overall among functions. For names and

details of development functions (#1-33) the reader is referred to Table 1.

1190

Function	R^2			Rank		
	Arachnida	**Crustacea**	**Insecta**	**Arachnida**	**Crustacea**	**Insecta**
1	AB	ABCD	ABCD	BCDEF	CDEFGH	EFGHIJ
2	A	AB	AB	ABC	A	ABCD
3	A	AB	AB	A	A	AB
4	AB	ABCD	ABCD	BCDEF	CDEFGH	FGHIJKL
5	AB	ABC	ABCD	ABCDEF	CDEFG	FGHIJKL
6	B	ABC	ABCD	DEF	DEFGH	LM
7	AB	ABC	ABC	ABCDE	ABC	DEFGH
8	A	AB	A	A	A	A
9	A	D	ABCD	ABCDEF	EFGHI	CDEF
10	AB	ABC	ABCD	ABCD	ABCD	BCDE
11	A	ABCD	ABCD	BCDEF	CDEFGH	DEFGHI

12	AB	ABCD	ABCD	ABCDEF	DEFGH	GHIJKLM
13	AB	ABC	ABCD	CDEF	CDEFGH	EFGHIJ
14	A	ABC	ABCD	ABCD	ABCD	BCDE
15	AB	ABC	ABCD	CDEF	ABCDE	EFGHIJKL
16	A	A	AB	AB	AB	ABC
17	A	ABC	ABCD	ABCDE	ABCD	BCDE
18	A	ABC	ABCD	ABCDE	BCDEF	EFGHIJ
19	A	ABC	ABCD	ABC	ABCD	BCDE
20	AB	CD	ABCD	ABCD	EFGHI	BCDE
21	A	ABCD	ABCD	EF	HI	JKLM
22	AB	ABC	ABC	ABCDEF	CDEFGH	EFGHIJKL
23	AB	ABC	ABCD	BCDEF	EFGHI	GHIJKLM
24	AB	ABCD	ABCD	ABCDEF	FGHI	EFGHIJKL
25	AB	ABCD	CD	BCDEF	EFGHI	KLM
26	A	AB	ABCD	ABC	ABCDE	BCDE
27	AB	BCD	ABCD	ABCDE	GHI	EFGHIJ
28	A	ABC	ABCD	ABCDEF	CDEFGH	EFGHIJK
29	AB	ABC	ABC	ABCDEF	CDEFGH	DEFG
30	B	ABC	BCD	F	HI	M
31	AB	ABC	ABCD	BCDEF	GHI	HIJKLM
32	AB	ABC	ABC	ABCDE	EFGHI	DEFG
33	AB	ABCD	D	EF	I	IJKLM

1191

Table 3. Consequences of fitting development data with functions used or found to be best in

original published studies rather than the best function for each of 79 reanalyzed species datasets,

as determined in this study. Consequences were assessed by calculating differences between

used and actual best functions in terms of (1) increased mean error (absolute, in days) between

observed and predicted development times, (2) increased percent (%) prediction error, (3)

usually poorer fit (R^2), (4) poorer percent (%) fit (R^2), and (5) information loss (Δ_i of used

models; $\Delta_i = 0$ is the best model). resulting. Mean values $\pm$ 95 % confidence intervals (95 % C.I.)

for each difference measure taken across all retested datasets are shown (with the range of values

in parentheses), as well as the results of one-sample t-tests comparing these differences to zero;

statistically-significant p-values ($p \leq 0.05$) are presented in bold text.

1202

Measure of difference between used and actual best functions	Mean $\pm$ 95 % C.I. (Minimum – Maximum)	Significance of difference from zero
Mean error (days)	4.042 ± 3.696 (-0.024 – 132.342)	$t_{78} = 2.144$, **p = 0.035**
Mean error (%)	85.913 ± 53.666 (-1.690 – 1725.529)	$t_{78} = 3.138$, **p = 0.002**
R^2	-0.091 ± 0.039 (-0.873 – 0.002)	$t_{78} = -4.620$**, p < 0.001**
R^2 (%)	-10.736 ± 4.595 (-100 – 0.172)	$t_{78} = -4.579$**, p < 0.001**
Δ_i of used function	225.011 ± 118.316 (0 – 3958.691)	$t_{78} = 3.727$, **p < 0.001**

1203