A critical review of the use and performance of different function types for modeling

temperature-dependent development of arthropod larvae

Brady K. Quinn*

Department of Biological Sciences, University of New Brunswick, 100 Tucker Park Road, Saint

John, NB, Canada E2L 4L5

*Corresponding author: bk.quinn@unb.ca, 1-506-343-7676

Highlights:

• Temperature-dependent development functions of arthropod larvae were reviewed

• Data from published studies were re-tested and fit with eight different function types

• 86.5 % of published studies did not fit their data with the best function of those tested

• Performance differed among functions and was related to taxon and temperature range tested

• Function type impacted predicted development times, so using the best function matters

ABSTRACT

Temperature-dependent development influences production rates of economically- and

ecologically-important arthropod species, including crustaceans important to fisheries and

agricultural pests. Numerous candidate equation types (development functions) exist to describe

the effect of temperature on development time, yet most studies use only a single type of

equation and there is no consensus as to which, if any model predicts development rates better

than the others, nor what the consequences of selecting a potentially incorrect model equation are

on predicted development times. In this study, a literature search was performed of studies fitting

development functions to development of arthropod larvae (87 species). The published data of

some (52) of these species were then fit with eight commonly-used development functions.

Overall performance of each function type and consequences of using a function other than the

best one to model data were assessed. Performance was also related to taxonomy and the range

of temperatures examined. The majority (86.5 %) of studies were found to not use the best

function out of those tested. Using the incorrect model lead to significantly less accurate (e.g.,

mean difference ± SE 7.9 ± 1.7 %, range: 1-18 %) predictions of development times than the best

function. Overall, Heip Power, Quadratic, Bělehrádek, and Modified Arrhenius functions

performed well, Linear Rate and Tauti Exponential were intermediate, and the Linear Sum and

Brière-2 functions performed poorly. More complex functions only performed well when wide

temperature ranges were tested, which tended to be confined to studies of insects or arachnids

compared with those of crustaceans. Results indicate the biological significance of choosing the

best-fitting model to describe temperature-dependent development time data.

Key Words: Temperature; arthropod; larval development; development functions; molting rate

1. Introduction:

Temperature affects biota at all levels, ranging from effects at the fundamental biochemical and physiological levels (Bělehrádek, 1935; Coutant and Talmage 1976; Somero, 2004) to effects on individual organisms (Brière et al., 1999; MacKenzie, 1988), populations (Aiken and Waddy, 1986; Cooper et al., 2012; McLaren et al., 1969), communities, and ecosystems (Menge, 1978; McQuaid and Branch, 1985). Through its effects on the physical and chemical properties of biologically active molecules, such as enzymes, temperature affects the rate at which numerous life processes occur, including metabolism, oxygen consumption, photosynthesis, movement, survival, growth, and embryonic development (Bělehrádek, 1935; Brière et al. 1999; Corkett, 1972; Coutant and Talmage 1976; Du et al., 2007; Geffen and Nash, 2012; Herzig, 1983; McLaren et al., 1969). Temperature also significantly affects larval development rate of poikilothermic animals, including some vertebrate larvae (Lind and Johansson, 2007; Kang et al., 2009; Miller et al., 2006) and those of invertebrates (e.g., de Severyn et al., 2000; Jenkins et al. 2006; Singh and Sharma, 1994). Temperature has particularly strong impacts on moulting and development of arthropods (Anger, 1984; Corkett and McLaren, 1970; Easterbrook et al., 2003; Hamasaki et al., 2009; Koda and Nakamura, 2010; MacKenzie, 1988; Marchioro and Forester, 2011; McLaren, 1963).

Within certain tolerance limits (Bělehrádek, 1935; Brière et al., 1999; Campbell et al., 1974; Shi and Ge, 2010) rates of biological processes of poikilotherms, including larval development, are positively correlated with temperature; thus, higher temperatures generally result in more rapid development than lower temperatures. This has important ecological implications, as environmental temperatures can influence generation times, production cycles, and population dynamics of such organisms. Higher or lower temperatures could, for example,

lead to changes in the amount and/or timing of peak secondary marine production of copepods

(Huntley and López, 1992; McLaren, 1963), outbreaks of agricultural pests (Easterbrook et al.,

2003) or vector-borne diseases (Bayoh and Lindsay, 2003), or introduction and establishment of

invasive species into new areas (de Rivera et al., 2007). Water temperatures could also influence

patterns of recruitment to populations of marine invertebrates, including crustaceans such as

lobsters and crabs, on which human fisheries depend (Aiken and Waddy, 1986; Anger, 1984;

Caddy, 1986; MacKenzie, 1988; Rothlisberg, 1979).

When modeling ecology and population dynamics of arthropods, equations are used to

represent the functional relationship between environmental temperature and development rate or

time of larvae. These equations, hereafter referred to as development functions, are derived by

rearing larvae at different controlled temperatures in a lab or hatchery setting, observing

development times of multiple larvae at each temperature, and then using regression analyses to

fit an equation relating temperature to development time (or its inverse, rate) to the data

obtained. There are countless potential forms of equation that can be used to fit such data,

including various linear, simple curvilinear, and complex non-linear functions (e.g., see reviews

by Anger, 2001; Angilleta Jr., 2006; Blanco et al. 1995; Guerrero et al., 1994; Heip, 1974;

McLaren, 1995; Shi and Ge, 2010; Smits et al. 2003). These functions differ in form,

assumptions, procedures used to derive their parameters, and most importantly in terms of the

development times predicted. For example, development times of American lobster, *Homarus*

americanus (H. Milne Edwards, 1837), larvae predicted with these eight different development

function types can differ from each other and the data used to derive them by ≥ 40 days at the

same temperatures (see Fig. 1; Table S1). However, this is no clear standard rule or consensus as

to what is the "best" type of development function to apply to these kinds of data. Researchers

are generally left to choose the type of development function to use on their own, and will often

select one or a very few forms that have the best apparent match to their characters of their data

or has been used by other studies on the same or related species (e.g., Edgar and Andrew, 1990;

McLaren et al., 1969). Given the potential for different functions to make very different

predictions of development times (e.g., Fig. 1), however, development function choice should be

given more consideration in studies on these species.

It is possible that certain function types may in general be better representations of the

relationship between temperature and development of arthropod larvae, or specific sub-groups

within the Arthropoda (e.g., arachnids vs. crustaceans vs. insects), for example because they

come closer to capturing thermal performance relations of enzymes and other biomolecules

mediating moulting and development cycles in these taxa (Brière et al., 1999; Huey and

Stevenson, 1979; Somero, 2004). As a result, such functions might also achieve better fit to

development data, be able to more-closely match real observed development times, and make

better predictions of development in nature. Differences in methodologies used in studies on

different taxa, for example the fact that the range of temperatures tested is generally wider for

insects and arachnids than crustaceans; (reviewed by Hartnoll, 1982; Quinn and Rochette, 2015),

might also lead to apparent taxonomic differences in function performance and should be

investigated. Several previous studies have compared the characteristics of different

development function types in general (Anger, 2001; Blanco et al., 1995; Guerrero et al., 1994;

McLaren, 1995), while others have attempted to fit multiple function types to data for one or two

specific species under study in order to select the best function for their data (e.g., Angilletta Jr.,

2006; Heip, 1974; Shi and Ge, 2010; Smits et al., 2003). Many other studies seem to choose one

or very few function(s) semi-arbitrarily, without discussing alternatives (e.g., de Oliveria et al.,

2009; Thompson 1982; see also Results). However, no previous study has attempted to assess the

degree to which one versus multiple types of development functions are used in published

studies, compared performance of different types of development functions across multiple

species, or assessed the overall impact of function choice to predictions made with such

functions. Such a large-scale analysis is needed, though, because it could potentially allow

functions that tend to better represent development data in general to be identified, which can

then allow for more informed decisions by future studies on arthropod larval development.

In this study, a critical literature review was conducted to assess whether and to what

extent studies of temperature-dependent development of arthropod larvae attempt to represent

their data with more than one development function type, and also which specific types of

functions tend to be used. Then, data from previously published studies were extracted and

retested to derive multiple different development functions for the same datasets. The best model

type for each dataset was determined, and whether or not published studies actually used the best

function type for their data was recorded. Overall performance of different function types were

assessed by comparing overall function rankings, proportion of variance explained, and

information loss across datasets. Any taxonomic patterns (e.g., whether particular function types

performed better for arachnids than for crustaceans and/or insects) and whether performance was

related to the range of temperatures tested were also noted. The consequences of using different

function types were then tested by comparing the difference of predicted versus observed

development times, fit, and information loss of the function type used in original studies versus

that of the best function for the data.

2. Materials and methods:

2.1. Literature review:

A literature search was conducted through Web of Science (Thompson Reuters, 2015) for

the terms "temperature" AND "development". An initial search was carried out on 12 September

2012, through which the majority of the data in the present study were obtained; this search

yielded 1,052 results. A second search was carried out on 19 November 2014, which returned 35

additional results not available or published online at the time of the initial search. These 1,087

total search results were them further examined, and several criteria were used to remove non-

relevant results. Accessible peer-reviewed studies that reported larval development rates or times

of arthropods at different temperatures and derived a regression equation(s) (i.e., development

function) from their data were sought out. Studies that looked exclusively at at growth (size

increase), which is a distinct process from development (Forster et al., 2011), were excluded.

After applying these criteria, 73 studies of 87 different arthropod species were obtained

for subsequent examination and analyses (Table S2). Several specific types of development

function were frequently utilized in these studies (Table S2); a sub-sample of these functions are

presented in Table 1 and discussed in the next section (2.2.1, below). Studies were published

between 1970 and 2014, and conducted in several different countries on marine, freshwater, and

terrestrial species (Table S2) of various taxa within the Arachnida (Phylum Arthropoda:

Subphylum Chelicerata), Crustacea, and Insecta (Table S2). To assess whether published studies

tested multiple development functions on their data, each study was carefully read and the

number and types of development functions used to fit the data for each study species were

noted. Even if results from multiple functions were not reported, if alternative functions than

reported were at least mentioned in the Methods sections of studies they were counted as having

considered > 1 function. Also if any study tested multiple development functions on their data

and concluded one of these to be the "best" function for their data this was also noted. The

percent (%) of the 87 species datasets from the literature search on which one, two, or more

functions were tested, and the % of datasets on which different types of functions were tested,

were then calculated.

2.2. Meta-analysis:

2.2.1. Development functions considered in this study:

In the present study, eight development function types were examined (Table 1; Figure

1). These included two functions derived by simple linear fitting to development data (the Linear

Rate and Linear Sum functions), three simplistic curvilinear (non-linear) functions (Heip Power,

Tauti Exponential, and Quadratic functions), two more-complex curvilinear functions including a

nonzero minimum temperature (T_{min}) for development (Bĕlehrádek and Modified Arrhenius

functions) and one very complex, asymmetrical non-linear function including both a minimum

(T_{min}) and maximum (T_{max}) temperature for development (Brière-2) (Table 1). These functions

were found in the literature review in the present study (see section 2.1 and Results) to be used

quite frequently in studies of arthropod larvae. Additional development functions also exist,

including several functions resembling the Brière-2 one (i.e., including T_{min} and T_{max} parameters)

but with very complex structures requiring specialized fitting procedures not readily applicable

in many statistical software packages (e.g., Huey and Stevenson, 1979; Ikemoto and Takai, 2000;

Lactin et al., 1995; Logan, 1988; Schoolfield et al., 1981; Wang et al., 1982; see review by Shi

and Ge, 2010); as such, these additional functions were not considered further in this study.

The Linear Rate function is fit to development rate (d) data and becomes a hyperbola

after transformation back to development time (D) (d= 1/D; Campbell et al., 1974; Table 1). The

Linear Sum function is a true linear model that is simply fit to D (Winberg, 1971; Table 1). Both

of these linear functions can be used to derive starting estimates for the values of T_{min} and T_{max}

(Campbell et al., 1974; Table 1) to be used in deriving Bĕlehrádek, Modified Arrhenius, and

Brière-2 functions (see below). The Heip function is a simple Power fit of D versus temperature

with an implicit minimum threshold for development of 0°C (Heip, 1974). The Tauti function is

likewise a simple exponential curve fit to time data, which is equivalent to the Exponential

function often fit to rates (Guerrero et al., 1994; Tauti, 1925). The Quadratic function is a

second-order symmetrical polynomial function with positive curvature, in which above a

particular temperature D begins to increase with temperature beyond a minimum value (e.g., de

Oliveira et al., 2009; Quinn and Rochette, 2015). The Bĕlehrádek (Bĕlehrádek, 1936; McLaren,

1963) and Modified Arrhenius (Guerrero et al., 1994) functions are based on power and modified

exponential curves, respectively, but with an added parameter (a nonzero minimum temperature,

T_{min}) included at which development rate approaches zero (Modified Arrhenius) while time

approaches infinity (Bĕlehrádek) at a vertical asymptote; if biologically-realistic values of T_{min}

cannot be obtained, then this parameter can be set to 0°C, making these functions equivalent in

this case to the Heip Power and regular Arrhenius functions (Guerrero et al., 1994). It should be

noted that both the regular and Modified Arrhenius functions relate development rate to the

inverse of temperature, rather than directly to temperature itself (see Table 1), and are thus

distinct from typical exponential curves. Lastly, the Brière-2 function is normally fit to rate and

has two vertical asymptotes at T_{min} and T_{max}, at which development time approaches infinity

(rate approaches 0) (Brière et al., 1999). The Brière-2 development curve is left-skewed, with

T$_{max}$ occurring at temperatures very close to that at which minimum development times occur

(Brière et al., 1999). There is also a Brière-1 function, in which the value of the b-parameter (see

Table 1) is constrained to a value of 2 (Brière et al., 1999), but because during preliminary

analyses far-better fits were obtained when allowing this parameter to vary than when

constraining it (results not shown), the more-limited Brière-1 version of this function was not

considered further in the present study.

2.2.2. Re-analysis of species datasets from published studies:

215   To assess whether the function(s) used in published studies were actually the "best"

functions for these published datasets, data were extracted for re-analysis from as many of the

studies obtained through the literature search described above as possible. Studies or datasets

from the literature search that did not examine at least one of the eight development function

types described in the Introduction (Table 1; Figure 1) and previous section (2.2.1) or did not

present their data in a way that allowed it to be extracted for retesting (e.g., only mean

development times resented, without any measure of error), had to be excluded. Therefore, only

52 species datasets out of the 87 studies from the literature search could be retested (Table S2);

these included 7 arachnids (all mites, in the Subclass Acari), 20 Crustaceans (9 copepods and 11

decapods), and 25 insects in 6 orders (5 Coleoptera, 3 Diptera, 5 Hemiptera, 4 Hymenoptera, 6

Lepidoptera, and 2 Thysanoptera) (Table S2). Raw data or means ± error (standard deviation

(SD), standard error (SEM), etc.) and sample sizes were extracted from tables or figures in

published papers for each of these 52 species and used to generate datasets for reanalysis.

228   Each study dataset was analyzed with linear and nonlinear regressions (Table 1) between

temperature and development time or rate, as appropriate. These regression were carried out

using IBM SPSS Statistics 22 (SPSS Inc., 2014). To simplify analyses, only total development

times or rates (i.e., summed across multiple larval stages) were examined and data for individual

stages were not. For development functions including thermal limits (T_{min} or T_{max}; see Table 1

and section 2.2.1), unconstrained regressions were initially carried out, with starting values for

these parameters set to values estimated from linear functions (see Table 1 and section 2.2.1,

above). However, results were not accepted if this yielded biologically unrealistic estimates, such

as $T_{min} < 0°C$ in a species not known to survive and develop at sub-zero temperatures,

unreasonably low T_{min} (e.g., -100°C) or high T_{max} (e.g., 100°C), or T_{min} or T_{max} within the

temperature range for which successful development was reported in the original study. In this

case, constrained regressions were carried out (e.g., $T_{min} \geq 0°C$, $T_{min} <$ minimum T with

successful development, $T_{max} >$ maximum T with successful development, etc.) until satisfactory

values were obtained.

Once a regression equation corresponding to each development function was obtained for

each dataset, AIC_C values (Akaike's Information Criterion (AIC) corrected for finite sample size;

Akaike, 1973; Anderson, 2008) were calculated using the residual sum of squares (RSS) between

observed development times and those predicted by each function for each study (Anderson,

2008). Development functions were then ranked for each species dataset based on AIC_C-values.

The "best" possible rank was 1, corresponding to the lowest AIC_C value, and the "worst"

possible rank – if there were no ties – was 8, which corresponded to the highest AIC_C value

among the functions tested. For each species dataset, whether or not the function used in its

original published study was the same as that determined to be the "best" function for the data in

this study was noted. RSS values were also used to calculate R^2-values for each function, on each

dataset. These values were used as a measure of function performance (see next section, 2.2.3),

indicating the proportion of the variation in observed development times that was explained by

each temperature-dependent development function.

2.2.3. Assessing and comparing the overall performance of different functions:

To determine whether any particular type(s) of function tended to do "better" than others,

the overall performance of each development function across species datasets was assessed using

three measures: average ranking (determined using AIC_C), R^2, and Δ_i values. These measures of

performance were also compared across different taxonomic groups (arachnids, crustaceans, and

insects) as it was possible that certain functions might better represent development of animals in

a certain group(s) better than animals in others; this could be due to real biological differences

among taxa or to different experimental methodologies used for their rearing.

Calculation of function rankings and R^2-values for each function on each dataset was

described above (previous section). Δ_i values were calculated once the best function for a given

dataset was determined to assess the information potentially lost by using a function other than

the best one (Anderson, 2008); a lower Δ_i value is better, and indicates less information loss. The

Δ_i value for a given function, "i", is calculated by subtracting the AIC_C value of the best function

of those tested from its AIC_C value (Anderson 2008); therefore, the best function will have $\Delta_i =$

0 . A function with a lower Δ_i, higher R^2, or lower (better) ranking value on average than other

functions was considered to have performed better overall than other functions.

Separate two-way ANOVAs were carried out in IBM SPSS Statistics 22 (SPSS Inc.,

2014) to compare each of these three measures of performance (rankings, R^2, and Δ_i) across

different development functions (factor with 8 levels; Table 1), as well as among different

taxonomic groups (factor with 3 levels: Arachnida, Crustacea, and Insecta). If a statistically-

significant ($p \leq 0.05$) interaction between function and taxon was found, the data for that

measure were split by taxon and then separate one-way ANOVAs comparing different functions

were carried out for each taxon. If significant differences among functions were found, Tukey's

Honestly Significant Difference (HSD) test was used to perform post-hoc comparisons among

specific function types.

2.2.4. Assessing the consequences of development function choice:

The consequences of choosing one function versus another to predict larval development

at different temperatures were assessed by comparing whether and how much the best function

identified in this study predicted observed development times relative to the function originally

used in the study from which each species dataset was obtained (hereafter the "original"

function). Three measures, described in detail below, were calculated for each of the 52 species

datasets to assess consequences of function choice. These were mean "improvement", R^2

increase, and original study function Δ_i.

To calculate the first of these, "improvement", the absolute deviance (in days) between

predicted (using a development function) and mean observed development time was calculated

for both the best and original study functions, at each temperature tested in original studies. The

absolute deviance for the original function was then subtracted from that for the best function at

each temperature, to determine how much predictions were improved (i.e., became closer to

observed values) by using the best versus original function. These differences were then

averaged across all temperatures to calculate a mean absolute "improvement" (in days) per each

dataset that was due to using the best versus original functions. Mean improvement per dataset

was also expressed as a percent improvement by performing the aforementioned calculations, but

before averaging differences between deviance of best and original functions across temperatures

these differences were divided by the original function's deviance and multiplied by 100 %; this

translated the improvement from a "raw" measure (in days) to a percentage (%).

302   R^2 increase was simply calculated for each dataset by subtracting the R^2-value of that

303   dataset's original function from that of its best function. Percent R^2 increase was also calculated

304   for each dataset by dividing the R^2 by the original function's R^2, and then multiplying by 100 %.

305   A large R^2 increase implied that a greater proportion of the variation in development time was

306   explained by the best function than the original one.

307   The Δ_i of the original function for each dataset was also examined to assess the extent of

potential information lost by using these, rather than the best functions, to fit the data. As

described above, a lower Δ_i (closer to $0 = \Delta_i$ of best function) is better, and implies the function

retains more useful information than a function with a higher Δ_i. Generally a function with $\Delta_i < 2$

contains some useful information, even if it is not the "best" function, whereas a function with Δ_i

> 14 is highly unlikely to be informative (Anderson 2008). A high original function Δ_i value

would imply that the original function was considerably less informative than the best function.

314   If the best function for a given dataset was the same as the original function, all of the

measures described above would have a value of zero; if any measure were not significantly

different from zero, then, it would imply that using the best versus original function did not result

in meaningfully different predicted development times. Therefore, the distributions of mean

"improvement" and R^2 increase when best versus original functions were used to predict

development times, as well as original function Δ_i values, across all species datasets were

compared to zero (null hypothesis of no differences) using analysis of confidence intervals

(Cummings et al., 2007). Mean values $\pm$ 95 % confidence intervals (95 % C.I.) of each

measurement were calculated across all datasets. If the 95% C.I.s did not overlap with zero the

null hypothesis could be rejected and a significant ($p < 0.05$) impact of function choice

concluded.

2.3. Potential relationship between thermal range and function performance:

One interesting pattern noted during the literature review in this study was that studies of

temperature-dependent development differed considerably in their methodology, particularly

regarding the range of temperatures tested. Thermal ranges varied considerably among studies in

general, from as narrow as 6°C to as wide as 35°C (Table S2). Differences also appeared to exist

between studies of different taxonomic groups, particularly between studies of crustaceans

versus those of arachnids and insects (Hartnoll, 1982; Quinn and Rochette, 2015; Table S2). To

confirm whether such taxonomic differences were significant, the thermal range for each species

dataset subjected to detailed re-analysis (n =52 total) was calculated as the maximum

temperature tested in its original study minus the minimum temperature tested; this included any

temperatures at which successful development was not observed (i.e., survival = 0 %), as this

implies a developmental threshold (like T_{min} and T_{max} in the Brière-2 function, for example).

One-way ANOVA was then used to compare thermal ranges among taxonomic groups.

More complex functions have lower power to model smaller datasets (Shi and Ge, 2010).

This is especially true for complex development functions containing threshold temperatures

(T_{min} and T_{max}) if these are fit to data recorded over narrow thermal ranges not approaching a

species' real thermal limits (Shi and Ge, 2010). In such cases, thermal thresholds have to be

extrapolated too far beyond actual observations, resulting in excessively extreme estimates for

these parameters. One could thus expect that complex functions like Brière-2 may perform better

when wider thermal ranges are tested, potentially approaching or encompassing real thermal

limits. To test whether or not more complex functions would be selected as better functions when

wider thermal ranges were tested, Pearson's correlation coefficients (R) were calculated between

the thermal ranges calculated for each dataset and the ranking of each function per dataset. Thus,

eight separate correlation analyses were carried out (one for each type of development function),

each consisting of 52 thermal range-ranking pairs (one pair per dataset). Whether correlations

were significant ($p \leq 0.05$), and if significant R-coefficients were positive or negative was

examined. As lower ranking values implied better function performance (see above), a positive

correlation for a given function implied that the function did worse when a wider range of

temperatures was tested, whereas a negative correlation meant that the function did better when a

wider thermal range was tested.

3. Results:

<u>3.1. Results of literature review – usage of different functions:</u>

Out of 87 different species datasets, over half (56.5 %, or 49 datasets) were reported to

have been fit with only one development function and the majority (89.2 %, or 78 datasets) were

fitted with four or fewer functions (Fig. 2A). In most cases studies that used 2-4 functions

examined insects or arachnids (Fig. 2A) and used 1-3 complex functions (like the Brière-2

function) plus the Linear Rate function to derive starting values for T_{min} and T_{max} parameters in

these complex functions (Fig. 2B; Table S2). The only taxonomic group for which > 4 different

functions were tested was Insecta, for which 10.6 % of all species datasets (9 in total,

representing 18.1 % of insects) were tested with 5-17 different development functions (Fig. 2A).

The most frequently-used development functions overall, and particularly among studies

of the Insecta and Arachnida, were the Linear Rate (34.9 % of all datasets, 48.7 % of insect

datasets, 29.5 % of arachnids) and Brière-2 functions (7.6 % of all datasets, 5.0 % of insects,

35.3 % of Arachnida), in addition to various other complex functions (30.2 % of all datasets,

37.5 % of Insecta, 35.3 % of arachnids) not examined in detail in this study (Fig. 2B; Table S2;

see Methods, section 2.2.1.). No other function types were used for arachnids, and the only other

functions used for insects were the Heip Power and Quadratic functions, and these were only

used rarely (5.0 and 3.8 % of insect datasets, respectively; Fig. 2B). All eight development

functions examined in this study – in addition to a few other types - were used in previous

studies of Crustacea (Fig. 2B; Table S2), with the most common being the Heip Power and

Bělehrádek functions in 6.8 and 8.3 % of all datasets, respectively (Fig. 2B), representing 25.7

and 31.3 % of Crustacean datasets.

3.2. Results of meta-analysis – did previous studies use the "best" function for their data?

The development function(s) used to fit species datasets and/or concluded to have been

the "best" function for these data in their original studies were found, in the vast majority of

cases, not to be the best function for the data out of the eight functions examined (Fig. 3). When

AIC_C was used to rank functions, the best function for 86.5 % of 52 study datasets was

concluded to be a different one than that selected in previous studies (Fig. 3). For all three

taxonomic groups, a considerable majority of datasets (75.0 % for Crustacea, 92.1 % for Insecta,

and 100 % for Arachnida) were found to be best fit using a different function than that used in

original studies of these species (Fig. 3).

3.3. Overall performance of different functions:

There was a significant interaction ($F_{14, 392} = 4.180$, $p < 0.001$) between the effects of

taxonomic group and development function type on overall performance of the eight different

development functions as assessed by ranking functions with AIC_C. Therefore differences in

rankings among functions were compared for each taxonomic group separately. In all three

taxonomic groups, ranking did significantly differ among different development functions

(Arachnida: $F_{7, 48} = 22.061$, $p < 0.001$; Crustacea: $F_{7, 48} = 11.103$, $p < 0.001$; Insecta: $F_{7, 48} =$

32.833, $p < 0.001$). Differences in rankings among functions were actually very similar across

the different taxonomic groups (Fig. 4). There were some differences among taxa in terms of the

placement of functions of more intermediate mean rankings; for instance, the Tauti Exponential

function was ranked well among Crustacea (Fig. 4C) but not for the other taxa (for Arachnida it

had one of the poorest ranks (Fig. 4A) and it was intermediate for Insecta (Fig. 4B)) and the

Linear Rate and Brière-2 functions were ranked poorly for the Arachnida and Crustacea (Fig.

4A, C) but intermediately for Insecta (Fig. 4B). However, the functions with the "best" (lowest

values, closest to 1) and "worst" (highest values, furthest from 1) mean rankings were mostly

consistent among groups. The best-ranked functions included the Heip Power, Quadratic, and

Bělehrádek functions for all three taxonomic groups, as well as the Modified Arrhenius function

in both Arachnida and Insecta (Fig. 4). The worst functions included the Linear Sum function in

all three taxa and the Brière-2 function in both Arachnida and Crustacea (Fig. 4).

3.4. Consequences of function choice:

R^2-values differed among the eight development functions ($F_{7, 392} = 7.973$, $p < 0.001$;

Fig. 5A) but not among taxonomic groups ($F_{2, 392} = 1.635$, $p = 0.196$); there was no significant

interaction between function and taxonomy ($F_{14, 392} = 1.138$, $p = 0.323$). Mean R^2-values were

fairly high for all development functions and ranged between 0.7 and 0.9 (Fig. 5A), meaning

that, on average, all temperature-dependent development functions tested were able to explain

the majority of variation in observed development times. Significant differences were found

among functions, however. The Heip Power, Quadratic, Bĕlehrádek, and Modified Arrhenius

functions had the highest overall R^2-values (approximately 0.9) and explained significantly more

variation in development time than the Tauti Exponential and Linear Rate functions, which had

intermediate R^2-values (approximately 0.8) (Fig. 5A). The Linear Sum and Brière-2 functions

had the lowest R^2-values, of 0.7 on average and explained significantly less variation in

development time than all other functions except the Linear Rate function (Tukey's HSD test, p

< 0.001; Fig. 5A). The explanatory power of development functions therefore differed by as

much as 10-20% on average depending on which was used (Fig. 5A).

Δ_i values also differed among functions ($F_{7, 392} = 6.178$, $p < 0.001$) but not among taxa (F

$_{2, 392} = 1.766$, $p = 0.172$; non-significant interaction: $F_{14, 392} = 0.812$, $p = 0.656$; Fig. 5B). All

mean Δ_i values were high (mean $\Delta_i \geq 44.5$) because each function was not selected as the best

function by AIC_C at least once, and in many cases the difference between the AIC_C values of the

best function and the 2nd best function (Δ_i) were quite large (as evidenced by the variance in Fig.

5B). The lowest average Δ_i values of 44.5 and 47.5 were calculated for the Bĕlehrádek and

Modified Arrhenius functions, and the highest value (mean $\Delta_i = 451.2$) for the Linear Sum

function (Fig. 5B). There were therefore large differences in the amount of information attained

depending on which function was used.

Mean improvement ± 95 % C.I. in how well predicted development times matched

observed time when the best versus the originally-used function were utilized was 1.9 ± 0.4 days,

and mean improvement per dataset ranged from < 1 to > 5 days (Fig. 6A). These improvements

could be fairly large in terms of percent improvement over predictions using original function,

which ranged from 1 to 18 % or greater, with an average value ± 95 % C.I. s of 7.9 ± 1.7 % (Fig.

6B). When the best function was found in this study to not be the same as the function used in

the original studies from which datasets were obtained (n = 42/52 datasets), the R^2 increased by

an average ± 95 % C.I.s of 0.105 ± 0.045 (Fig. 6C), which is significantly greater than zero (p <

0.05) meaning that ~10 % of the variation in the data was better explained using the best

function. The difference in R^2-values between originally-used and best functions could be much

greater in many cases, though, as percent increase in R^2 by using the best function ranged from

as low as 1% to as high as 80% for certain study datasets (Fig. 6C); mean percent increase in R^2

± 95 % C.I. was 5.935 ± 4.340 % (Fig. 6C). Additionally the mean Δ_i of the originally-used

functions was 261.9 ± 99.0 (Fig. 6D), indicating a substantial loss of information relative to the

actual best function, at least as determined in the present study. All measures of improvement

resulting from using the best rather than original functions on species datasets were significantly

different from zero (analysis of confidence intervals, p ≤ 0.05; Fig. 6).

3.5. Range of temperature tested in studies versus function performance:

Thermal ranges tested in original studies differed significantly among studies of different

taxonomic groups ($F_{2, 49}$ = 12.607, p < 0.001; see also Table S2). Interestingly, studies of

crustaceans tended to be carried out over significantly smaller thermal ranges (mean ± 95% C.I.

= 10.7 ± 1.4 °C) than those of arachnids (17.5 ± 3.0 °C' Tukey's HSD test, p =0.003) or insects

(16.82 ± 2.0 °C; Tukey's HSD test, p < 0.001) implying notable differences in the way thermal

effects are investigated in these taxa; thermal ranges studied for Insects and Arachnids did not

differ, however (Tukey's HSD test, p = 0.937).

The range of temperatures examined in previous studies was significantly correlated with

performance (ranking) of several of the development functions tested in this study (Table 2).

Rankings of the Linear Sum, Heip Power, and Tauti Exponential functions by AIC_C were

weakly, but significantly and positively correlated with temperature range (Table 2), meaning

that these functions tended to do more poorly (higher value = poorer rank, further from 1) when

larger thermal ranges were tested. Conversely, rankings of the Modified Arrhenius, Bělehrádek,

and Brière-2 functions ranked by AIC_C were negatively correlated with temperature range (Table

2), meaning that these functions performed better (lower value = better rank, closer to 1) when

studies tested a wider range of temperatures.

4. Discussion:

<u>4.1. Use and performance of different development functions in previous studies:</u>

Modeling functional relationships between temperature and life history characters is an

essential component of studying the biology of poikilothermic organisms (Angilletta Jr., 2006;

Bělehrádek, 1935; Shi and Ge, 2010). Predictions of generation times (Huntley and Lopez,1992),

timing of seasonal events (Bayoh and Lindsay, 2003), dispersal potential (de Rivera et al., 2007),

and recruitment to adult populations of Arthropoda (Aiken and Waddy, 1986; Caddy, 1986)

produced by such modeling efforts are thus sensitive to the types of temperature-dependent

larval development functions incorporated in these. Development times of arthropod larvae

predicted for the same species and temperatures by different function types can differ

substantially, which has important impacts on predictions made. Using the best possible function

to represent a given species' and/or study's dataset should thus be a crucial component of the

study of temperature-dependent arthropod larval development, which should precede reporting

and use of the results of such studies in models. However, in the present study this important step

was found to be largely bypassed by the majority of studies. Particular function types tended to

be used more often than others for particular taxonomic groups with little or no clear justification

for the choice made, while consideration of alternative function types was rarely reported in

published papers. In most cases the function used in original published studies was not actually

the best one for the datasets presented. Further, fitting these data with the best model resulted in

better fit, less disagreement between predicted and observed development times, decreased

information loss, and presumably also better predictive ability. These results demonstrate that

development function choice is an important but often-ignored step in research on arthropod

larval development, which should be given greater consideration in future studies.

495   Choosing one particular development function might have some justification if any

function(s) could be said to be better overall than others. In the present study, no single function

type was found to be the best or worst, but the Heip Power, Quadratic, Bĕlehrádek, and Modified

Arrhenius functions tended to perform quite well overall, while the Linear Sum and Brière-2

functions tended to perform quite poorly. Functions that performed well overall might be

recommended as good starting points for fitting development data, and those that did poorly

overall could conversely be used with caution. However, even these functions were not

universally found to be the best or worst for all datasets. For example, the Linear Sum function

was actually found to be the best for American lobster (*Homarus americanus* (H. Milne

Edwards, 1837)) data from Quinn et al. (2013), while the Heip Power, Bĕlehrádek, and other

overall "good" functions performed more poorly on these data (Table S1). Also, the Brière-2

function, which performed poorly overall, did somewhat better on insect data than for other taxa

and actually was among the best models for some insect species datasets (Table S2). Therefore,

it is difficult to make general statements about which function is always best to use; this must

rather be assessed on a case-by-case basis, for each species and study. Results in this study

showed that not using the best function for a given dataset can result in very different predicted

development times, which could lead to very different (and potentially erroneous) inferences and

predictions of species biology by modeling studies (Miller et al., 1998; Miller et al., 2006;

Quinn, 2014; Reitzel et al., 2004). The practice among many fields of study has been to fit data

with a development function type that has been used in previous studies on similar species; for

example, the frequent use of the Bĕlehrádek function on copepod crustaceans (Anger, 2001;

Corkett and McLaren, 1970; Hamasaki et al. 2009) or Linear Rate + complex function(s) on

insects and arachnids (Golizadeh and Zalucki, 2012; Shi and Ge, 2010; Smits et al., 2003; Table

S2). However, based on results of the present study this practice should be discontinued.

4.2. Importance of temperature range tested to function performance:

An interesting finding in the present study was that overall performance of several

function types was correlated with the range of temperatures tested in original published studies.

Specifically, as the range of temperatures tested increased performance (i.e., likelihood to be

ranked as the best model) of the simplest functions examined (Linear Sum, Heip Power, and

Tauti Exponential) decreased while that of the most complex functions (Bĕlehrádek, Modified

Arrhenius, and Brière-2) increased. This result does make sense, however, if one considers the

"real" nature of temperature-biological rate relationships. Because the actual performance of the

enzymes mediating larval development most certainly have upper and lower functional threshold

temperatures, beyond which development cannot progress (Brière et al., 1999; Quinn and

Rochette, 2015; Somero, 2004), one can assume that for most species the "true" relationship

between temperature and development time resembles the Brière-2 function, or similar complex

asymmetrical curves (e.g., Huey and Stevenson, 1979; Shi and Ge, 2010). A study carried out

over a very wide range of temperatures should be able to approach or exceed thermal thresholds

and therefore identify these limiting temperatures, and thus be best explained by a complex,

Brière-2-type function. However, if one carries out their study over a more narrow thermal

range, they will only be able to "observe" a certain section of the development curve, which

could be located relatively far from one or both threshold temperatures. This could result in the

observed temperature-development data having a distinctly Linear, Quadratic, or Power

function-like shape, such that one of these alternative, simpler functions would be identified as

the "best" function for the data over this specific range. Indeed, this seems to have been the case

in several of the datasets examined, in which thermal ranges and/or sample sizes were relatively

small (e.g., Quinn et al., 2013; Corkett and McLaren, 1970; Hamasaki et al., 2009; Carlotti et al.,

2007; Table S1, S2) and the best function was determined to be one of the simpler functions,

such as the Linear Sum or Heip function, even though these should be the least-realistic

(Bělehrádek, 1935; Brière et al., 1999; Somero, 2004). Importantly, when this occurs the best

function will be the one that provides the most informative description of development times

over a very specific range of temperatures, but its performance is likely to degrade if

extrapolation beyond this range is attempted. Ultimately the "best" function should be of a

complex form resembling a Brière-2-type function, but most studies, especially of Crustacea, are

not conducted over sufficiently wide temperature ranges to be allow good estimates of such

functions' parameters to be derived.

The vast majority of insects and arachnids have terrestrial and/or freshwater aquatic

habitats, in which temporal variability in air and water temperatures can be very large (Pakyari et

al., 2011; Sanchez-Ramos et al., 2007; Stavrinides et al. 2010). As a result, the likelihood of

these organisms and their larvae being exposed to extreme temperatures exceeding thresholds for

moulting, development, and/or survival can be high. Conversely, many crustaceans (and all of

those examined in the present review; Table S2) inhabit the marine realm as larvae and/or adults

(Paul and Paul, 1999; Roberts et al., 2012; Thompson, 1982). While it is not impossible that

marine crustacean larvae could encounter temperatures too high for development or survival

(e.g., such warm extremes could occur in shallow coastal areas, highly-stratified water columns,

intertidal zones at low tide, or more generally due to future climate change; Caffara et al., 2012;

Quinn and Rochette, 2015), they are thought to be far more likely to encounter lower limiting

temperatures, especially in the deeper ocean or temperate regions (Hartnoll, 1982; MacKenzie,

1988). This perceived difference in limiting temperatures appears to have lead studies on

temperature-dependent development in these groups along different paths. Studies of insects and

arachnids often use very wide temperature ranges with the intent of capturing lower and upper

limiting temperatures for development in their species because these physiological limits are

known to be essential to modeling these species in their natural environments (Shi and Ge, 2010;

Smits et al., 2003). Studies of crustaceans, to the contrary, tend to be limited to more narrow

thermal ranges deemed "ecologically-relevant" (i.e., likely to be encountered by the species in

nature); occasionally these include lower limits, but in general physiological limits, especially

upper ones, are rarely sought (Hartnoll, 1982).

While there is certainly logic behind the use of narrower, more-relevant thermal ranges in

studies of Crustacea, this approach still has potential to result in errors for two main, related

reasons. First, the type of development function used changes predicted development times both

at and between (interpolation) observed temperatures and especially outside of these

(extrapolation) (Angilletta Jr., 2006; Campbell et al., 1974; Quinn and Rochette, 2015). Second,

thermal development limits actually change the shape of the "real" and estimated (i.e., fitted by

regression) development curve, for example by decreasing its curvature when lower and upper

limits are further apart and increasing curvature when these are closer together (Bělehrádek,

1935; Brière et al., 1999; Shi and Ge, 2010; personal observations by author). All else being

equal, these difference in curvature can result in very different development times at the same

temperatures. As a result, it is important to know the physiological limits of a given species when

modeling its development. Even if a species rarely encounters temperatures close to these limits,

development times calculated at intermediate temperatures will be impacted by the values of

these limits; if one ignores these limits and uses a different function type, or attempts to estimate

limits by extrapolation from a narrow thermal range, there is great potential for erroneous

predictions of development times to be made.

4.3. Discussion of potential limitations and next steps:

In the present review, published studies on arthropod larvae were obtained through

literature searches through Web of Science (Thompson Reuters, 2015). These searches were by

no means comprehensive – many other studies of temperature-dependent development of

arthropod larvae exist that were not indexed in this search engine – but it was extensive and did

provide a good sample of such studies encompassing many different years, regions, and

arthropod taxa (Table S2). This sample of the relevant literature was thus appropriate and useful

for the purposes of the present review of development function usage and performance. An

expanded search using additional search tools in a future study could obtained data for other

taxonomic groups within the Arthropoda (e.g, myriapods, more non-Acari arachnids (spiders),

other orders of Crustacea and Insecta, etc.), or even outside of this phylum (e.g., molluscs: de

Severyn et al., 2000; nematodes: Jenkins et al., 2006; Singh and Sharma, 1994; urochorates:

Kang et al., 2009; vertebrates: McLaren and Cooley, 1972; Miller et al. 2006), which may reveal

additional patterns in study design, taxonomy, and function usage of interest. However, overall

patterns and conclusions of the present study would likely hold true. To simplify analyses, this

study conducted analyses on total larval development time data rather than on individual larval

stages. However, in most species development time of each larval stages has a distinct response

to temperature, requiring different developmental equations to be derived for each stage (Corkett

and McLaren, 1970; Hartnoll, 1982). Often survival to and through later larval stages is very

low, so power to fit more complex functions to later-stage development data can be limited. As a

result, the best function can potentially differ among larval stages of the same species, in the

same study; indeed, this was noted for American lobster data in the present study (Table S1). A

future study should investigate stage-specific changes in the "best" development function(s), to

confirm whether such patterns could impact the overall performance of different function types,

prediction error, and so on. However, it would make mathematical sense for similar patterns to

be found through such a detailed review to those noted in the present study, given that similar

factors (e.g., sample sizes and thermal ranges tested) would impact function performance.

As noted in the Introduction and Methods (2.2.1), numerous development function types

exist in addition to the eight examined in detail in this study. Many of these are extremely

complex (e.g., Angilletta Jr., 2006; Huey and Stevenson, 1979; Logan, 1988; Schoolfied et al.,

1981; Shi and Ge, 2010) and used very frequently in studies of arthropod (mainly non-Crustacea)

larvae (Golizadeh and Zalucki, 2012; Huang et al., 2008; Mobarakian et al., 2014; Table S2). It

is possible that one or more of the functions not examined herein could actually be the closest to

"real" development relationships and/or perform better overall than all others. For example, Shi

and Ge (2010) concluded that the Performance function (Huey and Stevenson, 1979) was best

out of the 12 functions they tested on development data for diamondback moth, *Plutella*

xylostella (L.). This and other additional functions should thus be investigated in a future,

expanded review. However, findings of this review that one or few functions are used by most

studies, the best function was not the one used in most original published studies, and that using

the non-best function results in poorer predictions would not change through consideration of

such additional functions. In fact, more complex functions are unlikely to achieve much higher

performance on many of the datasets obtained in the present study, particularly those for

Crustacea reared over narrow, non-extreme thermal ranges; if anything, functions with greater

complexity than the Brière-2 one are most likely to do worse on many of these datasets due

smaller sample sizes (Shi and Ge, 2010).

In this study function performance was assessed mainly in terms of fit (R^2 and observed

versus predicted values) and information loss (AIC_C and Δ_i). These gave good indications of how

appropriate each function and its parameter estimates were for particular datasets (e.g., how well

sample sizes supported estimation of more complex functions). However, future studies could

take more thorough approaches to assessing predictive ability of different functions. One

approach to be used in the future to assess function performance could be cross-validation

(Picard and Cook, 1984; Anderson, 2008). Even better would be actually testing development

functions on new data, for example by predicting development times for a particular species at

different temperatures using different functions derived in a prior study, and then measuring new

development times and comparing these to predictions. If future studies attempted this, very

thorough tests and new evidence in favour of one function or another might be obtained.

4.4. Implications of development time predictions based on different functions:

Differences in development times predicted for the same species and temperature among

development functions have potential to impact various types of predictions relevant to arthropod

biology and ecology. Larval survival is usually inversely related to larval duration, such that

slower development results in fewer potential recruits to adult populations (Reitzel et al., 2004;

Roberts et al., 2012). Most life history and bio-physical models account for this by reducing

larval numbers in simulated cohorts by a certain percentage at each model time step, resulting in

substantial, exponential losses per each additional step spend in larval development (e.g., Miller

et al., 1998; Quinn, 2014). In many crustaceans, the larval phase of the life cycle is the main

dispersive phase (Anger, 1984; de Rivera et al., 2007; MacKenzie, 1988). Lengthening the larval

developmental period of such larvae can dramatically alter how far and to where larvae drift in

simulations with ocean currents; for example, simulations by Quinn (2014) showed that slowing

larval development (and thus lengthening drift time) of American lobster larvae by 60 % could

result in increased drift distances of larvae by up to ca. 500 km. If drift were overestimated in

such models, for instance due to use of inappropriate development functions, then the degree of

population mixing would be overestimated and an incorrect estimate of population structure

obtained; underestimation by the same means could also lead to considerable errors. Likewise if

dispersal ability of larvae of an invasive species, such as the green crab *Carcinus maenas* (L.) (de

Rivera et al., 2007), were underestimated in this way potential invasions to new regions that

could be predicted may be missed. Using development functions to estimate the timing of

seasonal peaks in abundance of disease vectors (Bayoh and Lindsay, 2003), agricultural pests

(Campbell et al., 1994; Easterbrook et al., 2003; Stavrinides et al., 2010), or species that serve as

important food sources to others (e.g., copepod secondary productivity in the ocean: Carlotti et

al., 2007) also depends on being able to make good estimates of larval development. For many

species very small differences in development time similar to the difference in "errors" of best

and original studies' functions are enough to dramatically alter the nature and implications of

modeled predictions (e.g., ≤ 1-5 days; Gadino and Walton, 2012). The type of development

function used can thus have large impacts on predictions, so it is important that studies attempt to

find the best model for their data. Importantly, much research is now being initiated to assess

how future climate change will impact many species, including arthropods and their larvae

(Caffara et al., 2012; Quinn and Rochette, 2015). Use of non-best development functions within

such research clearly could result in erroneous predictions as well and so should be avoided.

4.5. Recommendations and conclusions:

Based on results of this study, it is recommended that future studies examining effects of

temperature on development of arthropod larvae consider and attempt to fit multiple alternative

development function types to their data to determine the best way to model their results and

report that this was attempted. No one function type is better or worse overall, but the range of

temperatures to be considered and potential use of results (e.g., for extrapolation to temperatures

not observed) can be used as a guide when deciding which functions are most likely to provide

good representations of data. In general simpler functions (Heip Power, Tauti Exponential) could

provide better descriptions of development observed over relatively narrow thermal ranges (but

provide poor extrapolation ability). Conversely, wider ranges encompassing lower (Bĕlehrádek,

Modified Arrhenius) and/or upper (Brière-2 and similar functions) limiting temperatures for

development can support more complex functions, which potentially resemble more closely true

enzymatic and biological thermal performance curves (Brière et al., 1999; Somero, 2004) and

may allow modeling over all temperatures potentially encountered. Studies on crustaceans in

particular should be conducted over wider thermal ranges in the future so that limiting

temperatures of these species can be identified and more complex, presumably realistic

development functions reliably fit to data for these organisms. Considering different potential

function types to find the best for each dataset should lead to better predictions of larval

development times in support of subsequent research on these important species.

Acknowledgements:

Thanks are due to Rémy Rochette, Joël Chassé, Jeff Houlahan, and Heather Hunt for advice and

guidance during this project, and the University of New Brunswick, Saint John Campus, for

providing resources that made the review and analyses possible.

References:

Aiken, D.E., Waddy, S.L., 1986. Environmental influence on recruitment of the American

lobster, *Homarus americanus*: a perspective. Can. J. Fish. Aquat. Sci. 43, 2258-2270.

Akaike, H., 1973. Information theory as an extension of the maximum likelihood principle, in:

Petrov, B.N., Csaki, F. (Eds.), Second International Symposium on Information Theory.

Akademiai Kiado, Budapest, pp. 267-281.

Al-Saffar, Z.Y., Grainger, J.N.R., Aldrich, J., 1996. Temperature and humidity affecting

development, survival and weight loss of the pupal stage of *Drosophila melanogaster*,

and the influence of alternating temperature on the larvae. J. Therm. Biol. 21, 389-396.

Anderson, D.R., 2008. Model Based Inference in the Life Sciences: A Primer on Evidence.

Springer Science+Business Media, LLC, New York.

Anger, K., 1984. Development and growth in larval and juvenile *Hyas coarctatus* (Decapoda,

Majidae) reared in the laboratory. Mar. Ecol. Prog. Ser. 19, 115-123.

Anger, K., 2001. Crustacean Issues 12: The Biology of Decapod Crustacean Larvae. A.A.

Balkema, Rotterdam.

Angilletta, Jr., M.J., 2006. Estimating and comparing thermal performance curves. J. Therm.

Biol. 31, 541-545.

Bayoh, M.N., Lindsay, S.W., 2003. Effect of temperature on the development of the aquatic

stages of *Anopheles gambiae* sensu stricto (Diptera: Culicidae). Bull. Entomol. Res. 93,

375-381.

Bĕlehrádek, J., 1935. Temperature and living matter. Protoplasma Monographia, No. 8.

Borntraeger, Berlin.

Blanckenhorn, W.U., 1997. Effects of temperature on growth, development and diapause in the

yellow dung fly – against all the rules? Oecologia 111, 318-324.

Blanco, J. M., Guerrero, F., Rodríguez, V., 1995. The fate of comparisons of models in

temperature-dependent growth of copepods: a reply to the comments by McLaren. J.

Plankton Res. 17, 1391-1394.

Brière, J.F., Pracross, P., Rioux, A.Y., Pierre, J.S., 1999. A novel rate model of temperature-

dependent development for arthropods. Environ. Entomol. 28, 22-29.

Caddy, J.F., 1986. Modelling stock-recruitment processes in Crustacea: some practical and

theoretical perspectives. Can. J. Fish. Aquat. Sci. 43, 2330-2344.

Caffara, A., Rinaldi, M., Eccel, E., Rossi, V., Pertot, I., 2012. Modelling the impact of climate

change on the interaction between grapevine and its pests and pathogens: European

grapevine moth and powdery mildew. Agr. Ecosyst. Environ. 148, 89-101.

Campbell, A., Frazer, B.D., Gilbert, N., Gutierrez, A.P., Mackauer, M., 1974. Temperature

requirements of some aphids and their parasites. J. Appl. Ecol. 11, 431-438.

Carlotti, F., Bonnet, D., Halsband-Lenk, C., 2007. Development and growth rates of

Centropages typicus. Progress in Oceanography 72, 164-194.

Cave, R.D., Schiacchetano, C., Diaz, R., 2009. Temperature-dependent development of the

cycad Aulacapsis scale, *Aulacapsis yasumatsui* (Hemiptera: Diaspididae). Florida

Entomol. 92, 578-581.

Choy, S., 1991. Embryonic and larval biology of *Liocarcinus holsatus* and *Necora puber*

(Crustacea: Brachyura: Portunidae). J. Exp. Mar. Biol. Ecol. 148, 77-92.

Compolo, O., Malacrinò, A., Laudani, F., Maione, V., Zappalà, L., Palmeri, V., 2014. Population

dynamics and temperature-dependent development of *Chrysomphalus aonidum* (L.) to

aid sustainable pest management decisions. Neotrop. Entomol. 43, 453-464.

Cooper, B.S., Tharp II, J.M., Jernberg, I.I., Angilletta Jr., M.J., 2012. Developmental plasticity of

thermal tolerances in temperate and subtropical populations of *Drosophila melanogaster*.

J. Therm. Biol. 37, 211-216.

Corkett, C.J., 1972. Development rate of copepod eggs of the genus *Calanus*. J. Exp. Mar. Biol.

Ecol. 10, 171-175.

Corkett, C.J., McLaren, I.A., 1970. Relationships between development rate of eggs and older

stages of copepods. J. Mar. Biol. Assoc. U.K. 50, 161-168.

Coutant, C.C., Talmage, S.S., 1976. Thermal effects. J. Wat. Poll. Contr. Fed. 48, 1486-1544.

Cummings, G., Fiddler, F., Vaux, D.L., 2007. Error bars in experimental biology. J. Cell Biol.

177, 7–11.

de Oliveira, S.A., Souza, B., Auad, A.M., da Silva, D.M., Souza, L.S., Carvalho, C.A., 2009.

Desenvolvimento e reprodução de *Sipha flava* (Forbes) (Hemiptera: Aphididae) em

diferentes temperaturas. Neotrop. Entomol. 38, 311-316.

de Rivera, C.E., Hitchcock, N.G., Teck, S.J., Steves, B.P., Hines, A.H., Ruiz, G.M., 2007. Larval

development rate predicts range expansion of an introduced crab. Mar. Biol. 150, 1275-

1288.

de Severeyn, Y. G., Severeyn, H., Grant, W., Reverol, Y., 2000. Effect of water temperature on

larval development of the bivalve mollusk *Tivela mactroides*: evaluation in the laboratory

and via simulation. Ecol. Model. 129, 143-151.

Du, W.-G., Hu, L.-J., Lu, J.-L., Zhu, L.-J., 2007. Effects of incubation temperature on embryonic

development rate, sex ratio and post-hatching growth in the Chinese three-keeled pond

turtle, *Chinemys reevesii*. Aquaculture 272, 747-753.

Easterbrook, M.A., Fitzgerald, J.D., Pinch, C., Tooley, J., Xu, X.-M., 2003. Development time

and fecundity of three important arthropod pests of strawberry in the United Kingdom.

Ann. Appl. Biol. 143, 325-331.

Edgar, A.J., Andrew, T.E., 1990. A simple method for fitting Bělehrádek's equation to

embryonic development data of zooplankton. Hydrobiologia 194, 177-181.

Eliopoulos, P.A., Kontodimas, D.C., Stathas, G.J., 2010. Temperature-dependent development of

Chilocorus bipustulatus (Coleoptera: Coccinellidae). Mol. Ecol. Evol. 39, 1352-1358.

Emmert, C.J., Mitzell III, R.F., Andersen, P.C., Frank, J.H., Stimac, J.L., 2008. Effects of

contrasting diets and temperatures on reproduction and prey consumption by

Proprioseiopsis asetus (Acari: Phytoseiidae). Exp. Appl. Acarol. 44, 11-26.

Forster, J., Hirst, A.G., 2012. The temperature-size rule emerges from ontogenetic difference

between growth and development rates. Functional Ecol. 26, 483-492.

Forster, J., Hirst, A.G., Woodward, G., 2011. Growth and development rates have different

thermal responses. Am. Nat. 178, 668-678.

Gadino, A.N., Walton, V.M., 2012. Temperature-related development and population parameters

for *Typhlodromus pyri* (Acari: Phytoseiidae) found in Oregon vineyards. Exp. Appl.

Acarol. 58, 1-10.

Gao, G.-Z., Perkins, L.E., Zalucki, M.P., Lu, Z.-Z., Ma, J.-H., 2013. Effect of temperature on the

biology of *Acyrthosiphon gossypii* Mordvilko (Homoptera: Aphididae) on cotton. J. Pest.

Sci. 86, 167-172.

Geffen, A.J., Nash, R.D.M., 2012. Egg development rates for use in egg production methods

(EPMs) and beyond. Fish. Res. 117-118, 48-62.

Golizadeh, A., Zalucki, M.P., 2012. Estimating temperature-dependent developmental rates of

potato tuberworm, *Phthorimaea operculella* (Lepidoptera: Gelechiidae). Insect Sci. 19,

609-620.

Gonzàlez, L., Carvajal, J., 2003. Life cycle of *Caligus rogercresseyi* (Copepoda: Caligidae)

parasite of Chilean reared salmonids. Aquaculture 220, 101-117.

Greenberg, S.M., Sétamou, M., Sappington, T.W., Liu, T.-X., Coleman, R.J., Armstrong, J.S.,

2005. Temperature-dependent development and reproduction of the boll weevil

(Coleoptera: Curculionidae). Insect Sci. 12, 449-459.

Guerrero, F., Blanco, J.M., Rodríguez, V., 1994. Temperature-dependent development in marine

copepods: a comparative analysis of models. J. Plankton Res. 16, 95-103.

Haghani, M., Fathipour, Y., Talebi, A.A., Baniameri, V., 2007. Temperature-dependent

development of *Diglyphus isaea* (Hymenoptera: Eulophidae) on *Liriomyza sativae*

(Diptera: Agromyzidae) on cucumber. J. Pest Sci. 80, 71-77.

Hamasaki, K., Sugizaki, M., Dan, S., Kitada, S., 2009. Effect of temperature on survival and

developmental period of coconut crab (*Birgus latro*) reared in the laboratory. Aquaculture

292, 259-263.

Hartnoll, R.G., 1982. Growth, in: The Biology of the Crustacea, Vol. 2: Embryology,

Morphology, and Genetics. Abele, L.G. (Ed.). Academic Press, New York, pp. 111-196.

Heip, C., 1974. A comparison between models describing the influence of temperature on the

development rate of copepods. Biologisch Jaarboek Dodonaea 42, 121-125.

Herzig, A., 1983. The ecological significance of the relationship between temperature and

duration of embryonic development in planktonic freshwater copepods. Hydrobiologia

100, 65-91.

Huang, Z., Ren, S., Musa, P.D., 2008. Effects of temperature on development, survival, longevity, and fecundity of the *Bemisia tabaci* Gennadius (Homoptera: Aleyrodidae) predator, *Axinoscymnus cardilobus* (Coleoptera: Coccinellidae). Biol. Control 46, 209-215.

Huey, R.b., Svenson. R.D., 1979. Integrating thermal physiology and ecology of ectotherms: a discussion of approaches. Am. Zool. 19, 357-366.

Huntley, M.E., López, M.D.G., 1992. Temperature-dependent production of marine copepods: a global synthesis. Am. Nat. 140, 201-242.

Ikemoto, T., Takai, K., 2000. A new linearized formula for the law of total effective temperature and the evaluation of line-fitting methods with both variables subject to error. Environ. Entomol. 29, 671-682.

Jacas, J.A., Peña, J.E., Duncan, R.E., Ulmer, B.J., 2008. Thermal requirements of *Fidiobia dominica* (Hymenoptera: Platygastridae) and *Haeckeliania sperata* (Hymenoptera: Trichogrammatidae), two exotic egg parasitoids of *Diaprepes abbreviatus* (Coleoptera: Curculionidae). BioControl 53, 451-460.

Jaffari, S., Fathipour, Y., Faraji, F., 2012. Temperature-dependent development of *Neoseiulus barkeri* (Acari: Phytoseiidae) on *Tetranychus urticae* (Acari: Tetranychidae) at seven constant temperatures. Insect Sci. 19, 220-228.

Jenkins, E. J., Kutz, S.J., Hoberg, E.P., Polley, L., 2006. Bionomics of larvae of *Parelaphostronglyus odocoilei* (Nematoda: Protostrongylidae) in experimentally infected gastropod intermediate hosts. J. Parasitol. 92, 298-305.

Kamps, D.M., 1978. The effect of temperature on the development time and brood size of *Diaptomus pallidus* Herrick. Hydrobiologia 61, 75-80.

Kang, K.H., Zhang, Z.F., Bao, Z.M., Zhou, B., Han, C.B., 2009. Influence of water temperature

on spawning induction and larval development of the sea squirt *Halocynthia ritteri* (Oka,

1906). Aquaculture Res. 40, 513-518.

Karimi-Malati, A., Fathipour, Y., Talebi, A.A., 2014. Development response of *Spodoptera*

exigua to eight constant temperatures: Linear and nonlinear modeling. J. Asia-Pac.

Entomol. 17, 349-354.

Khan, M., Gregg, P., Mensah, R., 2009. Effect of temperature on the biology of *Creontiades*

dilutus (Heteroptera: Miridae). Austral. J. Entomol. 48, 210-216.

Kim, D.-S., Riedhl, H., 2005. Effect of temperature on development and fecundity of the

predaceous plant bug *Deraeocoris brevis* reared on *Ephestia kuehniella* eggs. BioControl

50, 881-897.

Kim, S.C., Song, J.-H., Kim, D.-S., 2008. Effect of temperature on the development and

fecundity of the Cryptic Mealybug, *Pseudococcus cryptus*, in the laboratory. J. Asia-Pac.

Entomol. 11, 149-153.

Klein Breteler, W.C.M., Gonzalez, S.R., Schogt, N., 1995. Development of *Pseudocalanus*

elongatus (Copepoda, Calanoida) cultured at different temperatures and food conditions.

Mar. Ecol. Prog. Ser. 119, 99-110.

Koda, K., Nakamura, H., 2010. Effects of temperature on the development and survival of an

endangered butterfly, *Shijimiaeoides divinus barine* (Leech) (Lepidoptera: Lycaenidae).

Entomol. Sci. 13, 29-34.

Krüger, R.F., Kirst, F.D., de Souza, A.S.B., 2010. Rate of development of forensically-important

Diptera in southern Brazil. Revista Brasileira de Entomologia 54, 624-629.

Lactin, D.J., Holliday, N.J., Johnson, D.L., Craigen, R., 1995. Improved rate model of

temperature-dependent development by arthropod. Environ. Entomol. 24, 68-75.

Lavagnini, T.C., de Freitas, S., Bezerra, A.L., 2009. Aspectos biológicos de *Chrysoperla*

raimundoi Freitas & Penny (Neuroptera, Chrysopidae). Revista Brasileira de

Entomologia 53, 629-634.

Lefebvre, F., Pasquerault, T., 2004. Temperature-dependent development of *Ophyra aenescens*

(Wiedemann, 1830) and *Ophyra capensis* (Wiedemann, 1818) (Diptera, Muscidae).

Forensic Sci. Int. 139, 75-79.

Li, D., 2002. The combined effects of temperature and diet on development and survival of a

crab spider, *Misumenops tricuspidatus* (Fabricius) (Araneae: Thomisidae). J. Therm.

Biol. 27, 83-93.

Lind, M.I., Johansson, F., 2007. The degree of adaptive phenotypic plasticity is correlated with

the spatial environmental heterogeneity experienced by island populations of *Rana*

temporaria. J. Evo. Biol. 20, 1288-1297.

Liu, S.-S., Meng, X.-D., 1999. Modelling development time of *Myzus persicae* (Hemiptera:

Aphididae) at constant and natural temperatures. Bull. Entomol. Res. 89, 53-63.

Logan, D.P., Kettle, C.G., 2007. Temperature-dependent development and distribution in the soil

profile of pupae of greyback canegrub *Dermolepida albohirtum* (Waterhouse)

(Coleoptera: Scarabaeidae) in Queensland sugarcane. Austral. J. Entomol. 46, 17-22.

Logan, J.A., 1988. Toward an expert system for development of pest simulation models.

Environ. Entomol. 17, 359-376.

Luypaert, G., Witters, J., Huylenbroeck, J.V., Maes, M., De Riek, J., De Clercq, P., 2014.

Temperature-dependent development of the broad mite *Polyphagotarsonemus latus*

(Acari: Tarsonemidae) on *Rhododendron simsii*. Exp. Appl. Acarol. 63, 389-400.

MacKenzie, B.R., 1988. Assessment of temperature effects on interrelationships between stage

durations, mortality, and growth in laboratory-reared *Homarus americanus* Milne

Edwards larvae. J. Exp. Mar. Biol. Ecol. 116, 87-98.

Marchioro, C.A., Forester, L.A., 2011. Development and survival of the diamondback moth,

Plutella xylostella (L.) (Lepidoptera: Yponomeutidae) as a function of temperature:

effects of number of generations in tropical and subtropical regions. Neotrop. Entomol.

40, 533-541.

Mazzei, K.C., Newman, R.M., Loos, A., Ragsdale, D.W., 1999. Development rates of the native

milfoil weevil, *Eurhychiopsis lecontei*, and damage to Eurasian watermilfoil at constant

temperatures. Biol. Control 16, 139-143.

McLaren, I.A., 1963. Effects of temperature on growth of zooplankton and the adaptive value of

vertical migration. J. Fish. Res. Board Can. 20, 685-727.

McLaren, I.A., 1995. Temperature-dependent development in marine copepods: comments on

choices of models. J. Plankton Res. 17, 1385-1390.

McLaren, I.A., Cooley J.M., 1972. Temperature adaptation of embryonic development rate

among frogs. Physiol. Zool. 45, 223-228.

McLaren, I.A., Corkett, C.J., Zillioux, E.J., 1969. Temperature adaptations of copepod eggs from

the Arctic to the Tropics. Biol. Bull. 137, 486-493.

McQuaid, C.D., Branch, G.M., 1985. Trophic structure of rocky intertidal communities: response

to wave action and implications for energy flow. Mar. Ecol. Prog. Ser. 22, 153-161.

Mehrnejad, M.R., 2012. Biological parameters of *Elasmus nudus* (Hymenoptera, Eulophidae), a

parasitoid of the pistachio fruit hull borer moth, *Arimania komaroffi* (Lepidoptera,

Pyralidae). Biocontrol Sci. Technol. 22, 659-670.

Menge, B.A., 1978. Predation intensity in a rocky intertidal community: effect of an algal

canopy, wave action and desiccation on predator feeding rates. Oecologia 34, 17-35.

Miller, C.B., Lynch, D.R., Carlotti, F., Gentleman, W., Lewis, C.V.W., 1998. Coupling of an

individual-based population dynamic model of *Calanus finmarchicus* to a cricualtion

model for the Georges Bank region. Fish. Oceanogr. 7, 219-234.

Miller, D.C.M., Moloney, C.L., van der Lingen, C.D., Lett, C., Mullon, C., Field, J.G., 2006.

Modelling the effects of physical–biological interactions and spatial variability in

spawning and nursery areas on transport and retention of sardine *Sardinops sagax* eggs

and larvae in the southern Benguela ecosystem. J. Mar. Syst. 61, 212-229.

Milne Edwards, H., 1837. Histoire naturelle des Crustacés, comprenant l'anatomie, la

physiologie et la classification de ces animaux. Volume 2. Librairie encyclopédique de

Roret, Paris.

Mironidis, G.K., Savopoulou-Soultani, M., 2009. Development, survival and growth rate of the

Hyposoter didymator-Helicoverpa armigera parasitoid-host system: effect of host instar

at parasitism. Biol. Control 49, 58-67.

Mobarakian, M., Zamani, A.A., Karmizadeh, J., Naghadeh, N.M., Emami, M.S., 2014.

Modelling development of *Callosobruchus maculatus* and *Anisopteromalus calandrae* at

various constant temperatures using linear and non-linear models. Biocontrol Sci. Tech.

24, 1308-1320.

Mochizuki, S., Kayaba, Y., Tanida, K., 2006. Larval growth and development in the caddisfly

Cheumatopsyche brevilineata under natural thermal regimes. Entomol. Sci. 9, 129-136.

Mols, P.J.M., van de Ende, E., Blommers, L.H.M., 1998. Embryonic and larval development of

Orthosia (Lep., Noctuidae) species used for optimizing timing of monitoring and control

in apple orchards. J. Appl. Entomol. 122, 431-439.

Morimoto, S., Imamura, T., Visarathanonth, P., Miyanoshita, A., 2007. Effects of temperature on

the development and reproduction of the predatory bug *Joppeicus paradoxus* Puton

(Hemiptera: Joppeicidae) reared on *Tribolium confusum* eggs. Biol. Control 40, 136-141.

Newman, B.K., Papadopoulos, I., Vorsatz, J., Woolridge, T.H., 2006. Influence of temperature

on the larval development of *Upogebia africana* and *U. capensis* (Decapoda:

Thalassinidae: Upogebiidae) in the laboratory. Mar. Ecol. Prog. Ser. 325, 165-180.

Nondillo, A., Redaelli, L.R., Botton, M., Pinent, S.M.J., Gitz, R., 2008. Exigências Térmicas e

Estimativa do Número de Gerações Anuais de *Frankliniella occidentalis* (Pergande)

(Thysanoptera: Thripidae) em Morangueiro [Thermal Requirements and Estimate of the

Annual Number of Generations of *Frankliniella occidentalis* (Pergande) (Thysanoptera:

Thripidae) on Strawberry Crop]. Neotrop. Entomol. 37, 646-650.

Pakyari, H., Fathipour, Y., Enkegaard, A., 2011. Estimating development and temperature

thresholds of *Scholothrips longicornis* (Thysanoptera: Thripidae) on eggs of two-spotted

spider mite using linear and nonlinear models. J. Pest Sci. 84, 153-163.

Palmer, M.A., Coull, B.C., 1980. The prediction of development rate and the effect of

temperature for the meiobenthic copepod, *Microarthridion littorale* (Poppe). J. Exp. Mar.

Biol. Ecol. 48, 73-83.

Park, H.-H., Ahn, J.-J., Park, C.-G., 2014. Temperature-dependent development of

Cnaphalocrocis medianalis Guenée (Lepidoptera: Pyralidae) and their validation in semi-

field conditions. J. Asia-Pac. Entomol. 17, 83-91.

Paul, A.J., Paul, J.M., 1999. Development of larvae of the golden king crab *Lithodes aequispinus*

(Anomura: Lithodidae) reared at different temperatures. J. Crustac. Biol. 19, 42-45.

Picard, R., Cook, D., 1984. Cross-validation of regression models. J. Am. Stat. Assoc.79, 575-

583.

Prasad, Y.G., Prabhakar, M., Sreedevi, G., Rao, G.R., Venkateswarlu, B., 2012. Effect of

temperature on development, survival and reproduction of mealybug, *Phenacoccus*

solenopsis Tinsley (Hemiptera: Pseudococcidae) on cotton. Crop Protect. 39, 81-88.

Quinn, B.K., 2014. Assessing potential influence of larval development time and drift on large-

scale spatial connectivity of American lobster (*Homarus americanus*). M.Sc. Thesis,

University of New Brunswick, Saint John, NB.

Quinn, B.K., Rochette, R., 2015. Potential effect of variation of water temperature on

development time of American lobster larvae. ICES J. Mar. Sci. 72 (Supplement 1), i79-

i90.

Quinn, B.K., Rochette, R., Ouellet, P., Sainte-Marie, B., 2013. Effect of temperature on

development rate of larvae from cold-water American lobster (*Homarus americanus*). J.

Crustac. Biol. 33, 527-536.

Reitzel, A.M., Miner, B.G., McEdward, L.R., 2004. Relationships between spawning date and

larval development time for benthic marine invertebrates: a modeling approach. Mar.

Ecol. Prog. Ser. 280, 13-23.

Roberts, S.D., Dixon, C.D., Andreacchio, L., 2012. Temperature dependent larval duration and

survival of western king prawn, *Penaeus (Melicertus) latisulcatus* Kishinouye, from

Spencer Gulf, South Australia. J. Exp. Mar. Biol. Ecol. 411, 14-22.

Romalho, F.S., Wanderley, P.A., Malaquias, J.B., Rodrigues, K.C.V., Souza, J.V.S., Zanuncio,

J.C., 2009. Temperature-dependent development rates of *Bracon vulgaris*, a parasitoid of

boll weevil. Phytoparasitica 37, 17-25.

Rothlisberg, P.C., 1979. Combined effects of temperature and salinity on the survival and growth

of the larvae of *Pandalus jordani* (Decapoda: Pandalidae). Mar. Biol. 54, 125-134.

Sànchez-Ramos, I., Àlvarez-Alfageme, F., Castañera, P., 2007. Development and survival of the

cheese mites, *Acarus farris* and *Tyrophagus neiswanderi* (Acari: Acaridae), at constant

temperatures and 90% relative humidity. J. Stored Products Res. 43, 64-72.

Schoolfield, R.M., Sharpe, P.J.H., Magnuson, C.E., 1981. Non-linear regression of biological

temperature-depednent rate models based on absolute reaction-rate theory. J. Theor. Biol.

88, 719-731.

Shi, P., Ge, F., 2010. A comparison of different thermal performance functions describing

temperature-dependent development rates. J. Therm. Biol. 35, 225-231.

Singh, M., Sharma, S.B., 1994. Temperature effects on development and reproduction of

Heterodera cajani on Pigeonpea. J. Nematol. 26, 241-248.

Smits, N., Brière, J.-F., Fargues, J., 2003. Comparison of non-linear temperature-depednent

development rate models applied to *in vitro* growth of entomopathogenic fungi. Mycol.

Res. 107, 1476-1484.

Somero, G.N., 2004. Adaptation of enzymes to temperature: searching for basic "strategies".

Comp. Biochem. Phys. B 139, 321-333.

Son, Y., Lewis, E.E., 2005. Modelling temperature-dependent development and survival of

998   *Otiorhynchus sulcatus* (Coleoptera: Curculinonidae). Agr. Forest Entomol. 7, 201-209.

999   Stavrinides, M.E., Lara, J.R., Mills, N.J., 2010. Comparative influence of temperature on

1000   development and biological control of two common vineyard pests (Acari:

1001   Tetranychidae). Biol. Control 55, 126-131.

1002   Tauti, M., 1925. J. Imp. Fish Inst. Tokyo, 21, (cited in Winberg, 1971 and Guerrero et al., 1994).

1003   Taveras, R., Hilje, L., Carballo, M., 2004. Development of *Hypsipyla grandella* (Zeller)

1004   (Lepidoptera: Pyralidae) in response to constant temperatures [Desenvolvimento de

1005   *Hypsipyla grandella* (Zeller) (Lepidoptera: Pyralidae) em diferentes temperaturas

1006   constants]. Neotrop. Entomol. 33, 1-6.

1007   Thessalou-Legaki, M., 1990. Advanced larval development of *Callianassa tyrrhena* (Decapoda:

1008   Thalassinidea) and the effect of environmental factors. J. Crustac. Biol. 10, 659-666.

1009   Thompson, B.M., 1982. Growth and development of *Pseudocalanus elongatus* and *Calanus* sp.

1010   In the laboratory. J. Mar. Biol. Assoc. U.K. 62, 359-372.

1011   Tofangsazi, N., Buss, E.A., Meagher, R., Mascarin, G.M., Arthurs, S.P., 2012. Thermal

1012   requirements and development of *Herpetogramma phaeopteralis* (Lepidoptera:

1013   Crambidae: Spilomelinae). J. Econ. Entomol. 105, 1573-1580.

1014   Tun-Lin, W., Burkot, T.R., Kay, B.H., 2000. Effects of temperature and larval diet on

1015   development rates and survival of the dengue vector *Aedes aegypti* in north Queensland,

1016   Australia. Medical and Vet. Entomol. 14, 31-37.

1017   Ulmer, B.J., Jacas, J.A., Peña, J.E., Duncan, R.E., Castillo, J., 2006. Effect of temperature on life

1018   history of *Aprostocetus vaquitarum* (Hymenoptera: Eulophidae), an egg parasitoid of

1019   *Diaprepes abbreviatus* (Coleoptera: Curculionidae). Biol. Control 39, 19-25.

Voss, S.C., Cook, D.F., Hung, W.-F., Dadour, I.R., 2012. Survival and development of the

forensically important blow fly, *Calliphora varifrons* (Diptera: Calliphoridae) at constant

temperatures. Forensic Sci. Med. Pathol. 10, 314-321.

Wang, R.-S., Lan, Z.-X., Ding, Y.-Q., 1982. Studies on mathematical models of the relationship

between insect development and temperature. Acta. Ecol. Sin. 2, 47-57. [In Chinese]

Winberg, G.G., 1971. Methods for the Estimation of Production of Aquatic Animals. Academic

Press, London.

Yao, S., Huang, Z., Ren, S., Mandour, N., Ali, S., 2011. Effects of temperature on development,

survival, longevity, and fecundity of *Serangium japonicum* (Coleoptera: Coccinellidae), a

predator of *Bemisia tabaci* Gennadius (Homoptera: Aleyrodidae). Biocontrol Sci.

Technol. 21, 23-34.

Zhao, F., Zhang, W., Hoffman, A.A., Ma, C.-S., 2014. Night warming on hot days produces

novel impacts on development, survival and reproduction in a small arthropod. J. Anim.

Ecol. 83, 769-778.

1034

1035 **Electronic Supplementary Material:**

1036

1037 **Table S1.** Development function parameter estimates, R^2, and AIC_C for American lobster data

1038 plotted in Figure 1.

1039

1040 **Table S2.** List of studies and species obtained for the literature review and meta-analyses,

1041 including taxonomic grouping, function type(s) used, best function, whether the best and used

1042 function were the same, and temperature range tested.

 Tables and Figures:

Table 1. Names and forms of the 8 different development functions examined in this study.

Functions ranged in complexity from linear [2] to simple curvilinear [1, 3, 4, 5], to complex

curvilinear (including a "biological minimum temperature" parameter, T_{min}) [6, 7], to complex

asymmetrical (including both T_{min} and a "biological maximum temperature", T_{max}) [8]. If a T_{min}

("biological minimum temperature") or T_{max} ("biological maximum temperature") value is

implied within or estimable from a given function, this is indicated in the "Notes" column. In all

equations D is development time (in days), T is temperature (°C), A, B, C, a, and b are fitted

constants, and e is the Euler number. See section 2.2.1 for other details of these functions.

1052

Function [a]	Equation [b, c]	Notes
[1] Linear Rate[1]	$D = 1/(a + bT)$	$T_{min\ est} = -a/b$
[2] Linear Sum[2]	$D = A + BT$	$T_{max\ est} = -A/B$
[3] Heip Power[3]	$D = AT^B$	$T_{min} = 0°C$
[4] Tauti [d] **Exponential**[4]	$D = Ae^{BT}$	No T_{min}
[5] Quadratic[5]	$D = AT^2 + BT + C$	No T_{min}, $D_{min} = C$
[6] Bělehrádek[6, 7]	$D = A(T-T_{min})^B$	
[7] Modified Arrhenius[8]	$D = 1/[ae^{b(1/(T-Tmin))}]$	
[8] Brière-2[9]	$D = 1/[aT(T-T_{min})(T_{max}-T)^{(1/b)}]$	

[a] References for development functions are as follows: [1]Campbell et al. 1974; [2]Winberg (1971);
[3]Heip (1974); [4]Tauti (1925); [5]de Oliveira et al. (2009); [6]Bělehrádek (1935); [7]McLaren (1963);
[8]Guerrero et al. (1994); [9]Brière et al. (1999); [b] Functions that are usually fitted to development
rate (d = 1/D) are here represented in their inverted form, from which D can be directly
calculated; [c] For clarity lowercase letters (a, b) are used for functions fitted to d, and uppercase
letters (A, B, C) are used for functions fitted directly to D;[d] The Exponential function as
presented here [4] is mathematically equivalent to the Tauti function, $D = 1/[ae^{bT}]$, such that a =
1/A and b = -B, so the two are treated interchangeably.

Table 2. Correlations between the ranges of temperatures (°C) tested in original studies and rank

(out of 8) of development functions based on AIC_C for each species dataset. Pearson's

correlation coefficient values (R) and their associated p-values (evaluated for n = 52 pairs) are

presented; statistically-significant p-values (p ≤ 0.05) are indicated in bold text.

Function	R	p
Linear Rate	+0.067	0.637
Linear Sum	+0.387	**0.005**
Heip Power	+0.425	**0.002**
Tauti Exponential	+0.427	**< 0.001**
Quadratic	+0.056	0.696
Bělehrádek	-0.356	**0.010**
Modified Arrhenius	-0.510	**< 0.001**
Brière-2	-0.324	**0.019**

1065

1066

1067

Figure 1. (*Preceding page*) Examples of the curves formed by different development functions,

when these are used to calculate development time (y-axes, days) at different temperatures (x-

axes, °C). Types of functions plotted here are those presented in Table 1. Actual functions (see

Table S1) were derived from and fitted to data for total development times (combined time to

complete larval stages I-III) of American lobster (*Homarus americanus* (H. Milne Edwards,

1837)) larvae, as reported by Quinn et al. (2013); mean observed development times $\pm$ 95 %

C.I.s are plotted (squares) along with predictions of development functions (lines). Coefficients,

R^2, AIC_C, and Δ_i values for these functions are presented in Table S1. '*' = function used by

Quinn et al. (2013) to fit the data, '!' = actual "best" function for these data, and '+' = worst

function for these data.

1079

1080

Figure 2. Usage in each development study reviewed of (A) one or more different types of

development function and (B) specific development function types. The percentage (%) of

species datasets obtained in initial literature review (n = 87 total, see Table S2) that were tested

with each number and type of functions is plotted on the y-axes and broken down by taxa (gray

bars = arachnids, black = insects, and white = crustaceans). Tauti Expo = Tauti Exponential

function, and "Mod Arrhenius" = Modified Arrhenius function. *The "Other" category in (B)

includes all function types not otherwise listed; for clarity, if more than one "Other" function was

tested by original studies per dataset (see Table S2) it is not counted toward the total shown here.

1089

Figure 3. Percentage (%) of species datasets re-analyzed for meta-analysis (n = 52) for which the

development function found to be best for the data (lowest AIC_C value, ranking based on AIC_C =

1) in this study agreed with the best function as used in its original published study (Best

function used = Yes) or not (Best function used = No). Results are also broken down by

taxonomic group, as in Fig. 2.

1095

Figure 4. (*Preceding page*) Overall performance (assessed as ranking based on AIC_C values, y-

axes) of each development function (x-axes). Function performance was assessed separately for

each taxon: (A) arachnids (n = 7 species for each function), (B) insects (n = 25), and (C)

crustaceans (n = 20). Possible rankings ranged from 1 ("best" model) to 8 ("worst"). Values

plotted are mean rankings per function and taxonomic group taken across all species datasets

within that group ± 95 % C.I.s. Different letters above each mean indicate functions that ranked

significantly differently for each taxon (Tukey's HSD test, $p \leq 0.05$).

1103

Figure 5. Overall performance of different development functions (x-axes) assessed based on (A) R^2 values and (B) Δ_i values. Values are means $\pm$ 95% C.I.s calculated across all species datasets (n = 52), in all taxa for each function. Different letters above each mean indicate functions whose performance differed significantly (Tukey's HSD test, $p \leq 0.05$).

1110

Figure 6. Improvement in terms of (A) reduced mean error (absolute, in days) between observed

and predicted development times, (B) reduced percent (%) prediction error, or (C) improved fit

(R^2) resulting from using the best function for each species dataset (as determined in this study)

as opposed to the function used or found to be best in original published studies. The percentage

(%) of species datasets reanalyzed (n = 52, see Table S2) that showed different levels of

improvement is plotted on the y-axes. Also shown are the consequences of using the non-best

function in terms of (D) information loss (Δ_i of used models); note that $\Delta_i = 0$ is the best model.