

Adipocyte JAK2 mediates hepatic insulin sensitivity and the diabetogenic action of Growth Hormone

Kevin C. Corbit¹, João Paulo G. Camporez², Jennifer L. Tran¹, Camella G.
Wilson¹, Rachel J. Perry², Gerald I. Shulman^{2,3,4}, *Michael J. Jurczak^{5,6}, *Ethan J.
Weiss^{1,6}

*Co-senior authors

¹Cardiovascular Research Institute, University of California, San Francisco, San
Francisco, CA 94143

Department of ²Internal Medicine, ³Cellular and Molecular Physiology and the
⁴Howard Hughes Medical Institute, Yale University School of Medicine, New
Haven, CT, 06520

⁵Department of Medicine, Division of Endocrinology and Metabolism, University
of Pittsburgh, Pittsburgh, PA, 15261

⁶To whom correspondence should be addressed

ABSTRACT

For nearly 100 years, Growth Hormone (GH) has been known to regulate insulin sensitivity and risk of diabetes. However, the tissue governing the effects of GH signaling on insulin and glucose homeostasis remains unknown. Excess GH reduces fat mass and insulin sensitivity. Conversely, GH insensitivity (GHI) is associated with increased adiposity, augmented insulin sensitivity, and protection from diabetes. Here we induce adipocyte-specific GHI through conditional deletion of *Jak2* (JAK2A), an obligate transducer of GH signaling. Similar to whole-body GHI, JAK2A mice had increased adiposity and extreme insulin sensitivity. Loss of adipocyte *Jak2* augmented hepatic insulin sensitivity and conferred resistance to diet-induced metabolic stress without overt changes in circulating fatty acids. While GH injections induced hepatic insulin resistance in control mice, the diabetogenic action was absent in JAK2A mice. Collectively, our results show that adipose tissue governs the effects of GH on insulin and glucose homeostasis. Further, we show that JAK2 mediates liver insulin sensitivity via an extra-hepatic, adipose tissue-dependent mechanism.

Argentinian Physician Scientist, Bernardo Houssay demonstrated that injection of anterior pituitary extract worsens glycemic control in dogs ^{1,2}. In contrast, loss of anterior pituitary function leads to hypoglycemia and increased sensitivity to insulin ³. Similar results were observed in humans where hypophysectomy ameliorates not only insulin resistance ⁴⁻⁶ but diabetic complications as well ^{7,8}. More recently, it has been demonstrated that GH is responsible for much of the pituitary-derived diabetogenic activity ⁹. Both loss- and gain-of-function studies in humans and rodents support a role for GH in the biology of insulin responsiveness. Specifically, loss of GH receptor (GHR) function in humans and mice is associated with insulin sensitivity and protection against age-related diabetes ^{10,11}. Conversely, acromegalic patients with excessive GH secretion and transgenic *Gh* over-expressing mice have increased mortality and insulin resistance ¹²⁻¹⁴. Collectively, robust physiologic and genetic data support a prominent role for GH signaling in insulin/glucose homeostasis and the etiology of diabetes.

One of the major physiological functions of GH is controlling adipose tissue lipolysis ^{15,16}. Recent studies have demonstrated a critical role for insulin-mediated suppression of adipocyte lipolysis in the acute inhibition of hepatic gluconeogenesis through reductions in hepatic acetyl-CoA, leading to decreased pyruvate carboxylase activity and flux. Further, macrophage-induced lipolysis was shown to promote increased rates of hepatic gluconeogenesis and fasting hyperglycemia by promoting increased hepatic acetyl CoA content and pyruvate

carboxylase activity/flux, as well as increased conversion of glycerol to glucose (Perry et al., 2015). Although the ability for GH to promote lipolysis directly is ambiguous, data supporting a causal role for lipolysis in GH-mediated insulin resistance are robust ^{17,18}. The role, if any, of lipolytic activity in Growth Hormone Deficiency or GHI-associated augmentation of insulin sensitivity is entirely unknown.

The tissue(s) mediating the effects of GH signaling on insulin and glucose homeostasis has been elusive. Loss of GH signaling in liver through conditional deletion of the GH receptor *Ghr* ¹⁹, *Stat5* ²⁰, or *Jak2* ²¹ confers lean body mass, fatty liver, and insulin resistance. In stark contrast, mice ²² and humans ¹⁰ with global disruption of GHR have increased adiposity and insulin sensitivity.

There are conflicting results regarding skeletal muscle GHI on whole-body insulin sensitivity ^{23,24}, none of which phenocopy global GHI. Mice with beta-cell specific disruption of GHR show little effect on fasting insulin levels or insulin content in the pancreas on normal chow ²⁵. Recently, mice with fat-specific disruption of GHR were described to have increased fat mass but no change in insulin or glucose homeostasis ²⁶. However, these mice were generated using the *Fabp4:Cre* which is known to have activity outside of adipose tissue ²⁷.

In an effort to determine the specific role of adipose tissue in the metabolic activity of GH, we deleted *Jak2* from adipocytes using *Adiponectin:Cre*. Similar to

whole-body GHI, and as described earlier ²⁸, the resulting JAK2A mice had increased adiposity. Yet they also had improved whole-body insulin sensitivity. Interestingly, while chronic systemic GH exposure promoted hepatic insulin resistance and lipolysis in control mice, JAK2A animals were refractory to the diabetogenic action of GH. Prominent mechanisms regulating hepatic insulin sensitivity, including reductions in free fatty acids and liver lipid content, failed to account for the insulin sensitizing effects observed in JAK2A mice. Collectively, our work demonstrates that adipose tissue regulates the diabetogenic activity of GH and that a JAK2-dependent, adipose-derived factor mediates whole-body insulin sensitivity.

RESULTS

Adipocyte-specific deletion of *Jak2* augments adiposity and insulin responsiveness. We disrupted *Jak2* specifically in adipocytes using *Adiponectin:Cre*²⁹ on an inbred C57BL/6 background. On normal chow, body weight between control and JAK2A cohorts did not differ (Figure 1a). However, JAK2A mice had increases in both absolute and percent body fat (Figures 1b and 1c). Both visceral epididymal and subcutaneous inguinal fat pad mass were higher in JAK2A animals (Figures 1d and 1e). Despite this, fasting JAK2A mice were hypoglycemic and trended toward hypoinsulinemia (Figures 1f and 1g), implying increased whole-body insulin sensitivity. Consistent with this, the JAK2A cohort was hyper-responsive to insulin during the insulin tolerance test (ITT, Figure 1h). Thus, despite increased adiposity JAK2A mice have augmented whole-body insulin responsiveness.

Increased hepatic insulin sensitivity and suppression of endogenous glucose production in JAK2A mice. To definitively determine tissue-specific insulin sensitivity, hyperinsulinemic-euglycemic clamps were performed. There was no difference in clamped plasma insulin levels between groups during the clamp (Supplemental Figure 1b) and plasma glucose levels were matched at approximately 120 mg/dl (Figure 2a). As compared to the control cohort, JAK2A animals required a higher glucose infusion rate (GIR) to maintain euglycemia, confirming augmented whole-body insulin sensitivity (Figures 2b and 2c). While basal endogenous glucose production (EGP) was unchanged between groups

(Figure 2d), clamped JAK2A mice suppressed EGP following insulin infusion nearly 100% and to a much greater extent than control animals (Figures 2e and 2f), demonstrating improved hepatic insulin sensitivity. Whole-body glucose disposal (Figure 2g) and skeletal muscle- (Figure 2h) and adipose tissue-specific (Figure 2i) glucose uptake did not differ between the cohorts. Therefore, disruption of adipocyte *Jak2* confers improved whole-body insulin sensitivity almost entirely via augmented hepatic insulin sensitivity and suppression of hepatic glucose production.

Enhanced insulin-mediated suppression of lipolysis in JAK2A mice. It was recently shown that reductions in adipocyte lipolysis and fatty acids (FA) are major regulators of insulin-mediated suppression of hepatic glucose output^{30,31}. Since GH is a major *in vivo* regulator of lipolysis and loss of adipocyte *Jak2* greatly augmented insulin-induced reductions in EGP, we examined FA levels in the clamped mice. Plasma FA did not statistically differ between the euglycemic control and JAK2A cohorts (Figure 2j). Following insulin infusion the JAK2A cohort had lower absolute plasma FA (Figure 2k). However, no difference was appreciated between control and JAK2A mice when expressed as a percent suppression of basal FA (Figure 2l). Thus, while reduced lipolysis may contribute to augmented hepatic insulin sensitivity in JAK2A mice, the magnitude of difference between insulin-mediated suppression of EGP and FA (compare Figures 2e and 2k) presents the possibility of potential alternative mechanisms.

JAK2A mice are resistant to diet-induced metabolic derangement. To determine the susceptibility to metabolic stress, we next challenged mice with a high-fat diet (HFD). After ten weeks of HFD, body weight (Figure 3a), total (Figure 3b) and percent (Figure 3c) fat mass, and epididymal fat mass (Figure 3d) were unchanged between the control and JAK2A cohorts. The inguinal subcutaneous fat pads weighed more in JAK2A mice (Figure 3e). Both fasting blood glucose and insulin were reduced in JAK2A animals (Figures 3f and 3g). During ITT the JAK2A cohort maintained remarkable insulin responsiveness, suggesting whole-body insulin sensitivity was preserved despite HFD (Fig 3h).

JAK2A mice maintain exquisite hepatic insulin sensitivity following HFD without changes in FA levels. Hyperinsulinemic-euglycemic clamp experiments showed conclusively that JAK2A mice retained exquisite insulin sensitive following HFD. Plasma insulin (Supplemental Figure 1d) and glucose levels (Fig. 4a) were matched in response to a fixed and variable rate infusion, respectively, during the clamp. At steady-state, GIR was nearly doubled in JAK2A animals (Figure 4c). Similar to mice on chow, basal EGP was unchanged between the groups (Figure 4d). However, while clamped control mice demonstrated hepatic insulin resistance, EGP was completely suppressed following insulin infusion in JAK2A animals, and furthermore appeared negative (Figures 4e and 4f). Negative EGP as measured by tracer dilution during a hyperinsulinemic euglycemic clamp is a well-documented phenomenon that typically occurs in extremely insulin sensitive models with high rates of glucose turnover, likely due

to variable mixing of the exogenous 3-³H-glucose tracer with the endogenous glucose pool ³². Variable mixing of the 3-³H-glucose tracer also impacts measures of whole-body glucose disposal, which appeared unchanged between the two cohorts (Figure 4g). However, plasma clearance and tissue accumulation of the non-metabolizable glucose tracer ¹⁴C-2-deoxyglucose (2-DG) is not impacted by variable mixing and is therefore a more direct and reliable measure of glucose disposal. Tissue-specific glucose transport determined by 2-DG clearance was significantly increased in both skeletal muscle (Figure 4h) and adipose tissue (Figure 4i) in JAK2A mice. These data demonstrate that, despite the technical limitations of the 3-³H-glucose tracer under the conditions used, both hepatic and peripheral insulin sensitivity were improved in HFD JAK2A mice. In contrast, basal (Figure 4j) and insulin-mediated suppression of plasma FA (Figures 4k and 4l) did not differ between control and JAK2A animals. Therefore, adipocyte *Jak2* mediates whole-body insulin sensitivity even when challenged with HFD. Further, JAK2A mice augmented hepatic insulin sensitivity independent of statistically significant changes in FA levels. Collectively, JAK2A mice are resistant to the metabolic derangements of HFD.

Reduced hepatic triacylglycerol levels do not account for increased insulin sensitivity in chow fed JAK2A mice. Increased tissue lipid deposition strongly correlates with insulin resistance. Therefore, we reasoned that livers of JAK2A mice would have reduced lipid burden. Surprisingly, total levels of hepatic triacylglycerol (TAG) were unaltered between control and JAK2A chow fed

cohorts (Supplemental Figure 1e). Therefore hepatic insulin sensitivity in chow-fed JAK2A mice is not associated with reduced total liver lipid content. Following HFD hepatic TAG content was increased in both control and JAK2A animals (Supplemental Figure 1e). Hepatic TAG levels in HFD-fed JAK2A mice were lower than the control cohort and hence may contribute to preserved insulin sensitivity in JAK2A animals following dietary challenge.

Acute GH treatment induces hepatic insulin resistance dependent on adipocyte *Jak2*. Having established that insulin sensitivity is augmented in the setting of loss-of-function JAK2A mutants, we next determined the role of adipose tissue JAK2 in gain-of-function GH signaling. To this end, control and JAK2A mice were treated with vehicle or supra-physiological doses of recombinant mouse GH daily for five days. GH treatment induced hyperinsulinemia in control and, to a lesser extent, in JAK2A mice (Supplemental Figure 1f). Following the hyperinsulinemic clamp plasma insulin levels were equivalent in all groups (Supplemental Figure 1g). Similar to our previous chow-fed clamp study (Figure 2), vehicle-injected JAK2A mice required a higher GIR compared to the control cohort (Figures 5b and 5c). Systemic GH treatment reduced GIR by ~35% in control animals and by ~19% in JAK2A mice (Figure 5c). Basal EGP did not differ between groups (Figure 5d). Consistent with our previous study, JAK2A vehicle-treated mice demonstrated augmented EGP suppression following insulin infusion (Figure 5f). Conversely, GH treatment significantly reduced insulin-mediated EGP suppression in control animals,

confirming the diabetogenic activity of GH (Figure 5f). Remarkably, GH injections had no effect on hepatic insulin sensitivity in the JAK2A cohort (Figures 5e and 5f). GH treatment diminished whole-body glucose disposal (Figure 5g) and skeletal muscle glucose uptake (Figure 5h) in both control and JAK2A mice. No differences in adipose tissue glucose uptake were observed (Figure 5i). Thus, only five days of systemic GH exposure induces hepatic insulin resistance in an adipocyte *Jak2*-dependent manner.

GH promotes adipocyte lipolysis indirectly via JAK2-dependent inhibition of insulin action. In hyperinsulinemic-euglycemic clamped mice, circulating basal FA levels were unchanged in GH-injected animals (Figure 5j). Following insulin infusion, GH-injected control animals failed to suppress plasma FA to the levels of vehicle-treated mice, revealing that GH promoted lipolysis indirectly via inhibitory effects on insulin activity (Figure 5k). The ability of GH to antagonize reductions in circulating FA was absent in JAK2A mice (Figures 5k and 5l), demonstrating that adipocyte *Jak2* transduces the GH signal to interfere with insulin-mediated suppression of lipolysis. This correlated with retention of hepatic insulin sensitivity (Figure 5e). Collectively, we conclude that adipose tissue mediates GH-dependent antagonism of insulin's activity in both hepatocytes and adipocytes.

Similar to our earlier clamp experiment (Figure 2), neither basal (Figure 5j) nor percent suppression of circulating FFA (Figure 5l) differed between vehicle

treated control and JAK2A animals. Again, the lack of a demonstrable effect on lipolysis occurred concurrent with augmented hepatic insulin sensitivity (Figure 5e). A nearly identical pattern was observed for palmitate turnover (Supplemental Figures 1h-1j). Therefore, hepatic insulin resistance in GH-treated control animals directly correlates with adipocyte *Jak2*-dependent lipolysis. Further, the augmented hepatic insulin sensitivity of JAK2A mice may be mediated by a mechanism(s) other than or in addition to inhibition of lipolysis.

Adipocyte *Jak2* governs hepatic insulin signaling. Given that neither liver lipid content nor lipolysis strongly correlated with hepatic insulin sensitivity in mice lacking adipocyte *Jak2*, we measured hepatic insulin signaling in control and JAK2A cohorts. To this end, we continually exposed mice to recombinant GH for 28 days via mini-osmotic pumps. Subsequently, hepatic insulin responsiveness was assessed via levels of phosphorylated AKT (pAKT) following injections into the inferior vena cava (i.v.c). Vehicle infused control animals responded robustly to i.v.c. insulin (Figure 6). Conversely, chronic GH exposure strongly abrogated insulin-induced hepatic pAKT, consistent with the diabetogenic action of GH observed during the clamp studies and previous work³³. However, GH treatment did not impair insulin-induced pAKT in JAK2A livers, demonstrating that adipocyte JAK2 mediates hepatic insulin responsiveness to GH.

DISCUSSION

Houssay and colleagues first described their landmark studies on extreme insulin sensitivity in hypophysectomized dogs nearly 100 years ago ². In 1987, Mayer Davidson lamented that “the mechanism of GH effects on carbohydrate and lipid metabolism is far from clear.” Further, “Several studies suggest that the insulin antagonistic effects of GH may be secondary to the increased lipolysis via the glucose-fatty acid cycle. It is disappointing to note that this was the conclusion of the last general review of this area more than 20 years ago, and we have made little progress in substantiating or refuting this hypothesis.” ³⁴ Now, almost 30 years after this review, there is still no clear understanding of the cellular or tissue-level basis underlying the effects of GH on insulin and glucose homeostasis.

Global GHI, in both the physiologic and genetic context, confers insulin sensitivity throughout life. Conversely, acromegalic patients with excessive GH secretion are prone to insulin resistance, risk of diabetes and increased mortality. In fact, it has been proposed that GH hypersecretion may be the cause as much as the consequence of poor diabetic control ³⁵. These effects have been well documented in dogs, mice, and teleosts ³⁶, demonstrating robust evolutionary conservation. However, the vast array of loss-of-function studies describing inactivation of GHR, JAK2, or STAT5 in metabolic tissues including liver, skeletal muscle, pancreatic beta cell, and even adipose tissue has yielded complex and difficult to interpret results and failed to answer the key question of where GH

signaling (at the cell or tissue level) mediates its effects on whole-body insulin and glucose metabolism. Here, we show conclusively that abrogation of GH signaling in adipocytes augments whole-body and hepatic insulin sensitivity, protects from HFD-induced whole-body and hepatic insulin resistance, and prevents GH-mediated hepatic insulin resistance.

Our work relies on JAK2 as a signaling intermediate to infer GH action. While there is clear evidence that GH signaling is entirely dependent on JAK2 in fat and other tissues, it is possible or even likely that JAK2 mediates signals downstream of other cytokine signaling receptors in fat. Conditional deletion of *Ghr* with AP2:Cre results in obesity with no improvements in glucose homeostasis ²⁶. Removal of *Jak2* from adipose using the same *Cre* transgene was reported to also induce adiposity with age-related insulin resistance ³⁷. This is in contrast to our JAK2A mice that, while obese, are exquisitely insulin sensitive, even when challenged with HFD or supra-physiologic amounts of GH. Our work here and previously ²⁸ utilized *Adiponectin:Cre* which has been shown to be considerably more adipocyte-specific ³⁸. Thus, in the absence of an *Adiponectin:Cre*-driven *Ghr* deletion phenotype, it is not yet possible to conclude if the insulin sensitivity conferred by lack of adipose JAK2 is entirely due to interruption of GH signaling. However, we do definitively show that adipocyte JAK2 is required for GH to induce adipose tissue lipolysis and hepatic insulin resistance.

Insulin resistance strongly correlates with hepatic steatosis. Mechanistically, it has been shown that the subcellular localization of diacylglycerols (DAG) and not absolute levels of liver lipid mediate hepatic insulin resistance. Specifically, DAG and PKC ϵ are membrane-associated in insulin resistant rats but lipid droplet bound in the insulin sensitive state, preventing PKC ϵ translocation and activation at the plasma membrane where PKC ϵ inhibits Insulin Receptor kinase activity through phosphorylation of threonine 1150 ³⁹. These findings are germane, as in some cases liver lipid and hepatic insulin resistance do not correlate, such as CGI-58 ASO treatment ³⁹. For instance, we recently published work demonstrating that resolution of hepatic steatosis in hepatocyte-specific *Jak2* knockout mice (JAK2L) via concomitant deletion of *Cd36* is not sufficient to reverse the insulin resistance observed in JAK2L animals ⁴⁰. Others have shown discordance between insulin resistance and tissue lipid deposition, such as in Chanarin-Dorfman Syndrome and genetic forms of hepatic steatosis ^{41,42}.

Further, insulin resistance and adiposity fail to correlate in cases of Cowden and Laron Syndromes ^{43,44}. In fact, one of the natural consequences of augmented adipose tissue insulin sensitivity is reduced lipolysis, and hence increased adiposity, as is experienced by patients taking insulin-sensitizing agents such as thiazolidinedione ⁴⁵. Here we report that chow-fed JAK2A have increased adiposity, but fail to demonstrate statistical differences in the rate of either basal or insulin-attenuated lipolysis.

In 1963 Philip Randle put forth the notion that FAs may themselves induce insulin resistance ⁴⁶. Indeed, increasing levels of circulating FAs highly correlate with insulin resistance ⁴⁷. Further, clinical trials testing agents that reduce FAs have shown increased insulin sensitivity outcomes ⁴⁸⁻⁵¹. One of the many actions of insulin is to inhibit adipose tissue lipolysis and hence FA release into the circulation. GH has also been proposed to be a major regulator of adipose tissue lipolysis ¹⁶. Indeed, acromegalic patients have enhanced lipolysis and increased adipose expression of genes that regulate lipolysis ⁵². Two independent trials reported that administration of the anti-lipolytic drug acipimox increases insulin sensitivity in GH-treated patients ^{17,18}. Collectively, these data support a predominant role for lipolysis in GH-mediated insulin resistance. Our work here supports these findings. We showed that GH-mediated lipolysis correlates with hepatic insulin resistance and that GH promoted adipose tissue lipolysis through JAK2, but indirectly via perturbation of insulin action. Therefore, the predominant diabetogenic mechanism of GH is likely due to increased adipose tissue lipolysis promoting hepatic insulin resistance.

However, it is unknown if abrogated adipose tissue lipolysis mediates the enhancement of insulin sensitivity observed under loss-of-function conditions, such as in Laron Syndrome and other forms of GHI. As opposed to systemic GH treatment, our results do not support a role for modulation of lipolysis in JAK2A-associated insulin sensitivity. Specifically, we did not observe differences in circulating FA under clamped conditions between high fat diet fed control and

JAK2A mice, despite augmented hepatic insulin sensitivity in the latter. This is consistent with clinical results reporting that inhibition of GH signaling in Type 1 diabetics improves hepatic insulin sensitivity without a clear effect on lipolysis ⁵³. While the results from gain- and loss-of-function GH conditions appear disparate, the effects of GH on lipolysis may be determined by the nutritional state of the organism ⁵⁴. Here, we present the possibility that the acromegalic and GHI states may mediate insulin sensitivity by different mechanisms. Alternatively, adipocyte JAK2 may antagonize insulin sensitivity by additional mechanisms independent of GH signaling.

Recently two groups reported that insulin-induced reductions in hepatic glucose output can occur independently of liver insulin signaling ^{55,56}. This surprising finding supports older literature showing that insulin inhibits hepatic glucose output indirectly via effects on adipose tissue ⁵⁷⁻⁶⁰ and suggests that extra-hepatic tissue(s) receive insulin signals to mediate hepatic insulin sensitivity. In support of this, selective enhancement of adipocyte insulin sensitivity imparts whole-body glucose homeostasis ⁶¹. Further, transplanting insulin sensitive adipose tissue into insulin resistance animals restores insulin responsiveness ⁶²⁻⁶⁴. Finally, overt lack of adipose tissue function is sufficient to induce insulin resistance ⁶⁵. Therefore adipose tissue may be the dominant regulator of whole-body insulin sensitivity.

Exactly how adipose tissue mediates hepatic insulin sensitivity is unknown, but recent mechanisms have been proposed that include lipolysis ^{30,31}. We did not observe consistent differences in adipose tissue lipolysis or liver lipid content in chow-fed, HFD-fed or GH-treated insulin sensitized JAK2A mice. Further, we have not measured consistent or significant changes in classic insulin sensitizing factors such as Adiponectin and Leptin ²⁸. Therefore, adipocyte JAK2 may mediate insulin sensitivity by a novel mechanism.

Here, we report that loss of adipocyte *Jak2* is sufficient to impart whole-body insulin sensitivity independent of adiposity and liver lipid content, suggesting the existence of a paracrine regulator of GH diabetogenic activity and highlighting cross talk between adipose and liver. Our results show that acroegaly-associated hepatic insulin resistance is a result of attenuated insulin-mediated adipose tissue lipolysis. Finally, we provide evidence to support alternative mechanisms regulating gain- and loss-of-function GH signaling-mediated effects on insulin responsiveness. Isolation of the JAK2-dependent, adipocyte-derived factor(s) mediating whole-body insulin sensitivity may someday lead to new anti-diabetic therapeutics.

REFERENCES

- 1 Houssay, B. A. The Hypophysis and Metabolism. *New England Journal of Medicine* **214**, 961-971, doi:doi:10.1056/NEJM193605142142001 (1936).
- 2 Houssay, B. A. & Biasotti, A. The Hypophysis, Carbohydrate Metabolism And Diabetes. *Endocrinology* **15**, 511-523, doi:doi:10.1210/endo-15-6-511 (1931).
- 3 Houssay, B. A. Carbohydrate Metabolism. *New England Journal of Medicine* **214**, 971-986, doi:doi:10.1056/NEJM193605142142002 (1936).
- 4 Cushing, H. Further Concerning a Parasympathetic Center in the Interbrain: VII. The Effect of Intraventricularly-Injected Histamine. *Proc Natl Acad Sci U S A* **18**, 500-510 (1932).
- 5 Chabanier, H. & Copeman, W. S. A New System of Treatment of Diabetes Mellitus. *Br Med J* **1**, 897-898 (1926).
- 6 Luft, R. & Olivecrona, H. Experiences with hypophysectomy in man. *J Neurosurg* **10**, 301-316, doi:10.3171/jns.1953.10.3.0301 (1953).
- 7 Luft, R., Olivecrona, H., Ikkos, D., Kornerup, T. & Ljunggren, H. Hypophysectomy in man; further experiences in severe diabetes mellitus. *Br Med J* **2**, 752-756 (1955).
- 8 Javid, M., Gordon, E. S. & Erickson, T. C. Hypophysectomy in severe diabetes. I. Neurosurgical aspects. *J Neurosurg* **15**, 504-511, doi:10.3171/jns.1958.15.5.0504 (1958).
- 9 Greenberg, E. Growth Hormone and Diabetes Mellitus. *Diabetes* **14**, 43-45 (1965).

- 10 Guevara-Aguirre, J. *et al.* Growth hormone receptor deficiency is associated with a major reduction in pro-aging signaling, cancer, and diabetes in humans. *Sci Transl Med* **3**, 70ra13, doi:10.1126/scitranslmed.3001845 (2011).
- 11 List, E. O. *et al.* Endocrine parameters and phenotypes of the growth hormone receptor gene disrupted (GHR^{-/-}) mouse. *Endocr Rev* **32**, 356-386, doi:10.1210/er.2010-0009 (2011).
- 12 Capatina, C. & Wass, J. A. 60 YEARS OF NEUROENDOCRINOLOGY: Acromegaly. *J Endocrinol* **226**, T141-160, doi:10.1530/JOE-15-0109 (2015).
- 13 Valera, A. *et al.* Glucose metabolism in transgenic mice containing a chimeric P-enolpyruvate carboxykinase/bovine growth hormone gene. *FASEB J* **7**, 791-800 (1993).
- 14 Steger, R. W., Bartke, A. & Cecim, M. Premature ageing in transgenic mice expressing different growth hormone genes. *J Reprod Fertil Suppl* **46**, 61-75 (1993).
- 15 Moller, N. & Jorgensen, J. O. Effects of growth hormone on glucose, lipid, and protein metabolism in human subjects. *Endocr Rev* **30**, 152-177, doi:10.1210/er.2008-0027 (2009).
- 16 Chaves, V. E., Junior, F. M. & Bertolini, G. L. The metabolic effects of growth hormone in adipose tissue. *Endocrine* **44**, 293-302, doi:10.1007/s12020-013-9904-3 (2013).

- 17 Nielsen, S., Moller, N., Christiansen, J. S. & Jorgensen, J. O. Pharmacological antilipolysis restores insulin sensitivity during growth hormone exposure. *Diabetes* **50**, 2301-2308 (2001).
- 18 Segerlantz, M., Bramnert, M., Manhem, P., Laurila, E. & Groop, L. C. Inhibition of the rise in FFA by Acipimox partially prevents GH-induced insulin resistance in GH-deficient adults. *J Clin Endocrinol Metab* **86**, 5813-5818, doi:10.1210/jcem.86.12.8096 (2001).
- 19 Fan, Y. *et al.* Liver-specific deletion of the growth hormone receptor reveals essential role of growth hormone signaling in hepatic lipid metabolism. *J Biol Chem* **284**, 19937-19944, doi:10.1074/jbc.M109.014308 (2009).
- 20 Barclay, J. L. *et al.* GH-dependent STAT5 signaling plays an important role in hepatic lipid metabolism. *Endocrinology* **152**, 181-192, doi:10.1210/en.2010-0537 (2011).
- 21 Sos, B. C. *et al.* Abrogation of growth hormone secretion rescues fatty liver in mice with hepatocyte-specific deletion of JAK2. *J Clin Invest* **121**, 1412-1423, doi:10.1172/JCI42894 (2011).
- 22 Bartke, A. & Westbrook, R. Metabolic characteristics of long-lived mice. *Front Genet* **3**, 288, doi:10.3389/fgene.2012.00288 (2012).
- 23 Mavalli, M. D. *et al.* Distinct growth hormone receptor signaling modes regulate skeletal muscle development and insulin sensitivity in mice. *The Journal of Clinical Investigation* **120**, 4007-4020 (2010).

- 24 Vijayakumar, A. *et al.* Targeted Loss of GHR Signaling in Mouse Skeletal Muscle Protects Against High-Fat Diet-Induced Metabolic Deterioration. *Diabetes* **61**, 94-103, doi:10.2337/db11-0814 (2012).
- 25 Wu, Y. *et al.* Growth hormone receptor regulates β cell hyperplasia and glucose-stimulated insulin secretion in obese mice. *The Journal of Clinical Investigation* **121**, 2422-2426 (2011).
- 26 List, E. O. *et al.* The Role of GH in Adipose Tissue: Lessons from Adipose-Specific GH Receptor Gene-Disrupted Mice. *Molecular Endocrinology* **27**, 524-535, doi:10.1210/me.2012-1330 (2013).
- 27 Mullican, S. E. *et al.* A Novel Adipose-Specific Gene Deletion Model Demonstrates Potential Pitfalls of Existing Methods. *Molecular Endocrinology* **27**, 127-134, doi:doi:10.1210/me.2012-1267 (2013).
- 28 Nordstrom, S. M., Tran, J. L., Sos, B. C., Wagner, K.-U. & Weiss, E. J. Disruption of JAK2 in adipocytes impairs lipolysis and improves fatty liver in mice with elevated GH. *Molecular Endocrinology*, doi:10.1210/me.2013-1110 (2013).
- 29 Eguchi, J. *et al.* Transcriptional control of adipose lipid handling by IRF4. *Cell Metab* **13**, 249-259, doi:10.1016/j.cmet.2011.02.005 (2011).
- 30 Titchenell, P. M. *et al.* Direct Hepatocyte Insulin Signaling Is Required for Lipogenesis but Is Dispensable for the Suppression of Glucose Production. *Cell Metab* **23**, 1154-1166, doi:10.1016/j.cmet.2016.04.022 (2016).

- 31 Perry, Rachel J. *et al.* Hepatic Acetyl CoA Links Adipose Tissue Inflammation to Hepatic Insulin Resistance and Type 2 Diabetes. *Cell* **160**, 745-758, doi:<http://dx.doi.org/10.1016/j.cell.2015.01.012> (2015).
- 32 Finegood, D. T., Bergman, R. N. & Vranic, M. Estimation of endogenous glucose production during hyperinsulinemic-euglycemic glucose clamps. Comparison of unlabeled and labeled exogenous glucose infusates. *Diabetes* **36**, 914-924 (1987).
- 33 Gao, Y. *et al.* The role of PTEN in chronic growth hormone-induced hepatic insulin resistance. *PLoS One* **8**, e68105, doi:10.1371/journal.pone.0068105 (2013).
- 34 Davidson, M. B. Effect of Growth Hormone on Carbohydrate and Lipid Metabolism. *Endocr Rev* **8**, 115-131, doi:10.1210/edrv-8-2-115 (1987).
- 35 Press, M., Tamborlane, W. V. & Sherwin, R. S. Importance of raised growth hormone levels in mediating the metabolic derangements of diabetes. *N Engl J Med* **310**, 810-815, doi:10.1056/NEJM198403293101302 (1984).
- 36 McMenamin, S. K., Minchin, J. E., Gordon, T. N., Rawls, J. F. & Parichy, D. M. Dwarfism and increased adiposity in the gh1 mutant zebrafish vizzini. *Endocrinology* **154**, 1476-1487, doi:10.1210/en.2012-1734 (2013).
- 37 Shi, S. Y. *et al.* Adipocyte-specific deficiency of Janus kinase (JAK) 2 in mice impairs lipolysis and increases body weight, and leads to insulin resistance with ageing. *Diabetologia* **57**, 1016-1026, doi:10.1007/s00125-014-3185-0 (2014).

- 38 Lee, K. Y. *et al.* Lessons on conditional gene targeting in mouse adipose tissue. *Diabetes* **62**, 864-874, doi:10.2337/db12-1089 (2013).
- 39 Cantley, J. L. *et al.* CGI-58 knockdown sequesters diacylglycerols in lipid droplets/ER-preventing diacylglycerol-mediated hepatic insulin resistance. *Proc Natl Acad Sci U S A* **110**, 1869-1874, doi:10.1073/pnas.1219456110 (2013).
- 40 Wilson, C. G. *et al.* Hepatocyte-Specific Disruption of CD36 Attenuates Fatty Liver and Improves Insulin Sensitivity in HFD-Fed Mice. *Endocrinology* **157**, 570-585, doi:10.1210/en.2015-1866 (2016).
- 41 Nur, B. G., Gencpinar, P., Yuzbasioglu, A., Emre, S. D. & Mihci, E. Chanarin-Dorfman syndrome: Genotype-Phenotype Correlation. *Eur J Med Genet* **58**, 238-242, doi:10.1016/j.ejmg.2015.01.011 (2015).
- 42 Lallukka, S. & Yki-Jarvinen, H. Non-alcoholic fatty liver disease and risk of type 2 diabetes. *Best Pract Res Clin Endocrinol Metab* **30**, 385-395, doi:10.1016/j.beem.2016.06.006 (2016).
- 43 Pal, A. *et al.* PTEN mutations as a cause of constitutive insulin sensitivity and obesity. *N Engl J Med* **367**, 1002-1011, doi:10.1056/NEJMoa1113966 (2012).
- 44 Guevara-Aguirre, J. *et al.* GH Receptor Deficiency in Ecuadorian Adults Is Associated With Obesity and Enhanced Insulin Sensitivity. *J Clin Endocrinol Metab* **100**, 2589-2596, doi:10.1210/jc.2015-1678 (2015).

- 45 Carey, D. G. *et al.* Effect of rosiglitazone on insulin sensitivity and body composition in type 2 diabetic patients [corrected]. *Obes Res* **10**, 1008-1015, doi:10.1038/oby.2002.137 (2002).
- 46 Randle, P. J., Garland, P. B., Hales, C. N. & Newsholme, E. A. The glucose fatty-acid cycle. Its role in insulin sensitivity and the metabolic disturbances of diabetes mellitus. *Lancet* **1**, 785-789 (1963).
- 47 Boden, G. Gluconeogenesis and glycogenolysis in health and diabetes. *J Investig Med* **52**, 375-378 (2004).
- 48 Fulcher, G. R., Walker, M., Catalano, C., Agius, L. & Alberti, K. G. Metabolic effects of suppression of nonesterified fatty acid levels with acipimox in obese NIDDM subjects. *Diabetes* **41**, 1400-1408 (1992).
- 49 Fulcher, G. R. *et al.* A double blind study of the effect of acipimox on serum lipids, blood glucose control and insulin action in non-obese patients with type 2 diabetes mellitus. *Diabet Med* **9**, 908-914 (1992).
- 50 Hadigan, C., Liebaw, J., Torriani, M., Andersen, R. & Grinspoon, S. Improved triglycerides and insulin sensitivity with 3 months of acipimox in human immunodeficiency virus-infected patients with hypertriglyceridemia. *J Clin Endocrinol Metab* **91**, 4438-4444, doi:10.1210/jc.2006-1174 (2006).
- 51 Daniele, G. *et al.* Chronic reduction of plasma free fatty acid improves mitochondrial function and whole-body insulin sensitivity in obese and type 2 diabetic individuals. *Diabetes* **63**, 2812-2820, doi:10.2337/db13-1130 (2014).

- 52 Hochberg, I. *et al.* Gene Expression Signature in Adipose Tissue of Acromegaly Patients. *PLoS One* **10**, e0129359, doi:10.1371/journal.pone.0129359 (2015).
- 53 Thankamony, A. *et al.* Short-term administration of pegvisomant improves hepatic insulin sensitivity and reduces soleus muscle intramyocellular lipid content in young adults with type 1 diabetes. *J Clin Endocrinol Metab* **99**, 639-647, doi:10.1210/jc.2013-3264 (2014).
- 54 Bergan, H. E., Kittilson, J. D. & Sheridan, M. A. Nutritional state modulates growth hormone-stimulated lipolysis. *Gen Comp Endocrinol* **217-218**, 1-9, doi:10.1016/j.ygcen.2015.04.017 (2015).
- 55 Titchenell, P. M., Chu, Q., Monks, B. R. & Birnbaum, M. J. Hepatic insulin signalling is dispensable for suppression of glucose output by insulin in vivo. *Nat Commun* **6**, 7078, doi:10.1038/ncomms8078 (2015).
- 56 Sullivan, O. *et al.* FoxO1 integrates direct and indirect effects of insulin on hepatic glucose production and glucose utilization. *Nature communications* **6**, 7079, doi:10.1038/ncomms8079 (2015).
- 57 Lewis, G. F., Zinman, B., Groenewoud, Y., Vranic, M. & Giacca, A. Hepatic Glucose Production Is Regulated Both by Direct Hepatic and Extrahepatic Effects of Insulin in Humans. *Diabetes* **45**, 454-462, doi:10.2337/diab.45.4.454 (1996).
- 58 Girard, J. The Inhibitory Effects of Insulin on Hepatic Glucose Production Are Both Direct and Indirect. *Diabetes* **55**, S65-S69, doi:10.2337/db06-S009 (2006).

- 59 Lewis, G. F., Vranic, M., Harley, P. & Giacca, A. Fatty Acids Mediate the Acute Extrahepatic Effects of Insulin on Hepatic Glucose Production in Humans. *Diabetes* **46**, 1111-1119, doi:10.2337/diab.46.7.1111 (1997).
- 60 Levine, R. & Fritz, I. B. The Relation of Insulin to Liver Metabolism. *Diabetes* **5**, 209-222, doi:10.2337/diab.5.3.209 (1956).
- 61 Morley, T. S., Xia, J. Y. & Scherer, P. E. Selective enhancement of insulin sensitivity in the mature adipocyte is sufficient for systemic metabolic improvements. *Nat Commun* **6**, 7906, doi:10.1038/ncomms8906 (2015).
- 62 Foster, M. T. *et al.* Subcutaneous Adipose Tissue Transplantation in Diet-Induced Obese Mice Attenuates Metabolic Dysregulation While Removal Exacerbates It. *Physiol Rep* **1**, doi:10.1002/phy2.15 (2013).
- 63 Gavrilova, O. *et al.* Surgical implantation of adipose tissue reverses diabetes in lipoatrophic mice. *J Clin Invest* **105**, 271-278, doi:10.1172/JCI7901 (2000).
- 64 Klebanov, S., Astle, C. M., DeSimone, O., Ablamunits, V. & Harrison, D. E. Adipose tissue transplantation protects ob/ob mice from obesity, normalizes insulin sensitivity and restores fertility. *J Endocrinol* **186**, 203-211, doi:10.1677/joe.1.06150 (2005).
- 65 Bindlish, S., Presswala, L. S. & Schwartz, F. Lipodystrophy: Syndrome of severe insulin resistance. *Postgrad Med* **127**, 511-516, doi: 10.1080/00325481.2015.1015927 (2015).

- 66 Ayala, J. E. *et al.* Standard operating procedures for describing and performing metabolic tests of glucose homeostasis in mice. *Dis Model Mech* **3**, 525-534, doi:10.1242/dmm.006239 (2010).
- 67 Jurczak, M. J. *et al.* Dissociation of inositol-requiring enzyme (IRE1alpha)-mediated c-Jun N-terminal kinase activation from hepatic insulin resistance in conditional X-box-binding protein-1 (XBP1) knock-out mice. *J Biol Chem* **287**, 2558-2567, doi:10.1074/jbc.M111.316760 (2012).

METHODS

Animals and diets. Derivation of *Adiponectin:CRE;Jak2^{lox/lox}* (JAK2A) mice were previously described ²⁸. For these studies *Jak2^{lox/lox}* were used as controls. However, for these studies, the *Jak2^{lox/lox}* mice were backcrossed onto the C57BL/6 background for at least nine generations. For chow studies, mice were fed PicoLab Mouse Diet 20 (Lab Diet #5058; Percent calories provided by protein 23%, fat 22%, carbohydrate 55%) and for high fat diet studies mice were fed Research Diets D12492 (percent calories provided by protein 20%, fat 60%, carbohydrate 20%).

Dietary studies. Six week-old control and JAK2A mice were fed high fat diet or maintained on chow for ten weeks. Total fat mass was determined by Dual-energy X-ray absorptiometry. Following a 16-hour overnight fast blood glucose

and serum insulin levels were determined by hand-held glucometer (X) and ELISA (X), respectively. For insulin tolerance testing, mice were fasted for four hours (09:00-13:00) followed by intraperitoneal injection of 2 U/g insulin (Novolin® Novo Nordisk, Bagsvaerd Denmark). Blood glucose levels were determined by tail prick using a hand-held glucometer at the times indicated.

Hyperinsulinemic-euglycemic clamp studies. Clamp studies were performed according to recent recommendations of the NIH-funded Mouse Metabolic Phenotyping Consortium and as previously described ^{66,67}. Briefly, mice recovered one week after receiving surgery to implant an indwelling jugular vein catheter prior to clamp studies. Mice were fasted overnight and received a basal infusion of 3-³H-glucose and U-¹³C-palmitate conjugated to BSA to determine fasting glucose and palmitate turnover. A primed/continuous infusion of insulin, 3-³H-glucose and U-¹³C-palmitate was administered alongside a variable infusion of 20% dextrose to maintain euglycemia during the hyperinsulinemic portion of the study. A 10 µCi bolus injection of ¹⁴C-2-deoxyglucose was given at 90 min to determine tissue-specific glucose uptake, which was calculated from the area under the curve of ¹⁴C-2-deoxyglucose detected in plasma and the tissue content of ¹⁴C-2-deoxyglucose-6-phosphate. Blood was collected by tail massage at set intervals and glucose levels measured by a glucose oxidase method. Insulin infusion rates were as follows; chow 2.5 mU·kg⁻¹·min⁻¹; HFD 4.0 mU·kg⁻¹·min⁻¹; GH 2.5 mU·kg⁻¹·min⁻¹. Glucose and palmitate turnover were determined as the ratios of the 3-³H-glucose and U-¹³C-palmitate infusion rates to the specific

activity or plasma enrichment corrected to the contribution of the infusion rate for plasma glucose and palmitate, respectively, during the last 40 min of the hyperinsulinemic infusion. Calculation of endogenous glucose production and tissue specific glucose uptake were described previously ⁶⁷. For chow studies mice were approximately 16 weeks of age and HFD study mice were age-matched to chow studies and fed HFD for 4 weeks prior to clamps. For Growth Hormone (GH) treatment mice received daily vehicle (0.03M NaHCO₃, 0.15 NaCl, pH 9.5) or 5 mg/kg recombinant mouse GH (Dr. A. F. Parlow, National Hormone and Peptide Program, UCLA, Torrance, CA) by subcutaneous injection for 5 days prior to clamp.

Mini-osmotic pump studies. Control and JAK2A mice were implanted with 28-day Alzet mini-osmotic pumps (Durect Corporation, Cupertino CA) to deliver recombinant mouse GH at a dose of 5 mg/kg/day. GH was resuspended in GH buffer (0.03M NaCO₃, 0.15M NaCl, pH 9.5). At the end of 28 days, mice were fasted overnight (16 hrs) followed by injection with 0.01U insulin via the inferior vena cava. After 5 minutes, the liver was removed and immediately snap frozen in liquid nitrogen. Liver tissue was prepared for gel electrophoresis as previously described ⁴⁰. For detection of pAkt and total Akt, liver was homogenized in RIPA buffer (50 mM Tris, 150 mM NaCl, 1% Triton-X100, 1% deoxycholate, 0.1% SDS, 1mM EDTA, pH 7.4) supplemented with protease and phosphatase inhibitors (Thermo Fisher Scientific, Waltham MA). After incubating on ice for 20 minutes, the homogenate was centrifuged for 20 minutes at 15000 RCF at 4C. The protein

concentration of the supernatant was determined using BCA assay and 20 g of protein used for gel electrophoresis. P-Akt serine-473 and Akt antibodies (Cell Signaling) were used at a dilution of 1:2000 followed by anti-Rabbit secondary at a dilution of 1:10000. Images were collected using Chemidoc Imager (Bio-Rad, Hercules, CA).

Statistics and graphics. All statistical tests and figures were done using GraphPad Prism v6.0

Acknowledgements

This study was supported by National Institutes of Health (NIH) Grants 1R01DK091276 (to E.J.W.), DK076169 (M.J.J. & E.J.W.), DK099402 (M.J.J.), DK059635, DK40936, DK45735 (G.I.S.). We also gratefully acknowledge the support of the James Peter Read Foundation, the University of California, San Francisco (UCSF) Cardiovascular Research Institute, the UCSF Diabetes Center (P30 DK063720), and the UCSF Liver Center (P30 DK026743).

Author contributions

J. L. T. performed experiments for Figures 1 and 3. J. P. G. C. and R. J. P. executed experiments for Figures 2, 4, and 5. C. G. W. carried out experiments for Figure 6. G. I. S. provided critical insight and reviewed the manuscript. K. C. C. wrote the manuscript, performed statistical analyses, and made the figures. M. J. J. wrote the manuscript and designed and supervised the clamp

experiments. E. J. W. conceived of the study, wrote the manuscript, performed statistical analyses, made the figures.

Contributing financial interests

The authors declare no competing financial interests.




FIGURE 1. Loss of adipocyte *Jak2* augments insulin responsiveness in chow fed mice.

a.) Total body weight and b.) fat mass in control (CON) and JAK2A mice. c.) Percent body fat as a fraction of total body weight. d.) Total epididymal visceral and e.) inguinal subcutaneous fat. f.) Fasting plasma glucose and g.) insulin. h.) Insulin tolerance testing expressed as a percent of basal (fasting) glucose in control (open circles) and JAK2A (black triangles) mice. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001 by T test (b-f) and OneWay ANOVA (h). (g)=grams. N=7-11 for both cohorts. Data are +/- S.E.M.




FIGURE 2. Adipocyte *Jak2* mediates hepatic insulin sensitivity in chow fed mice.

a.) Blood glucose levels and b.) glucose infusion rate (GIR) during glucose infusion to achieve euglycemia in control (CON, open circles) and JAK2A (black triangles) mice. c.) GIR at euglycemia in control and JAK2A mice. d.) Basal and e.) clamped endogenous glucose production (EGP) in control and JAK2A mice. f.) Percent suppression of EGP following insulin infusion in control and JAK2A mice. g.) Whole-body glucose uptake in control and JAK2A mice. h.) 2-deoxyglucose (2-DG) uptake in gastrocnemius and i.) epididymal visceral fat in control and JAK2A mice. j.) Basal and k.) clamped plasma free fatty acid (FFA) levels in control and JAK2A mice. l.) Percent suppression of plasma FFA following insulin infusion in control and JAK2A mice. *p<0.05, **p<0.01, ***p<0.001 by OneWay ANOVA (a and b) and T test (c, e, f, and k). N=9 for both cohorts. Data are +/- S.E.M.




FIGURE 3. JAK2A mice maintain exquisite insulin responsiveness following dietary challenge.

a.) Total body weight and b.) fat mass in control (CON) and JAK2A mice. c.) Percent body fat as a fraction of total body weight. d.) Total epididymal visceral and e.) inguinal subcutaneous fat. f.) Fasting plasma glucose and g.) insulin. h.) Insulin tolerance testing expressed as a percent of basal (fasting) glucose in control (open circles) and JAK2A (black triangles) mice. ** $p < 0.01$, **** $p < 0.0001$ by T test (e, f, and g) and OneWay ANOVA (h). (g)=grams. N=8-16 for both cohorts. Data are \pm S.E.M.




FIGURE 4. Loss of adipocyte *Jak2* imparts hepatic and whole-body insulin sensitivity in high fat diet fed mice.

a.) Plasma glucose levels and b.) glucose infusion rate (GIR) during glucose infusion to achieve euglycemia in control (CON, open circles) and JAK2A (black triangles) mice. c.) GIR at euglycemia in control and JAK2A mice. d.) Basal and e.) clamped endogenous glucose production (EGP) in control and JAK2A mice. f.) Percent suppression of EGP following insulin infusion in control and JAK2A mice. g.) Whole-body glucose uptake in control and JAK2A mice. h.) 2-deoxyglucose (2-DG) uptake in gastrocnemius and i.) epididymal visceral fat in control and JAK2A mice. j.) Basal and k.) clamped plasma fatty acid (FA) levels in control and JAK2A mice. l.) Percent suppression of plasma FA following insulin infusion in control and JAK2A mice. *p<0.05, **p<0.01, ***p<0.001 by OneWay ANOVA (a and b) and T test (c, e, f, h, and i). N=8-11 for both cohorts. Data are +/- S.E.M.




FIGURE 5. Adipocyte *Jak2* is required for Growth Hormone-induced hepatic insulin resistance and lipolysis.

a.) Plasma glucose levels and b.) glucose infusion rate (GIR) during glucose infusion to achieve euglycemia in vehicle (Veh)-injected control (CON, open black circles) and JAK2A (closed black triangles) mice and Growth Hormone (GH)-injected control (open green circles) and JAK2A (closed green triangles) cohorts. c.) GIR at euglycemia in vehicle and GH-injected control and JAK2A mice. d.) Basal and e.) clamped endogenous glucose production (EGP) in vehicle- and GH-injected control and JAK2A mice. f.) Percent suppression of EGP following insulin infusion in vehicle- and GH-injected control and JAK2A mice. g.) Whole-body glucose uptake in vehicle- and GH-injected control and JAK2A mice. h.) 2-deoxyglucose (2-DG) uptake in gastrocnemius and i.) epididymal visceral fat in vehicle- and GH-injected control and JAK2A mice. j.) Basal and k.) clamped plasma free fatty acid (FFA) levels in control and JAK2A mice. l.) Percent suppression of plasma FFA following insulin infusion in control and JAK2A mice. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$ by OneWay ANOVA (a and b). N=6-8 for both cohorts. Data are \pm S.E.M.




FIGURE 6. Adipocyte *Jak2* mediates insulin-induced hepatic AKT activation.

a.) Western blot of insulin-stimulated liver lysates with antibodies against phosphorylated (pAKT) and total AKT (AKT) in control and JAK2A mice following continuous delivery of vehicle (-) or recombinant Growth Hormone (GH, +) for 28 days. d.) Densitometric ratio of pAKT/AKT in control and JAK2A mice treated (+GH) or not (-GH) with GH. *p<0.05, **p<0.01, ****p<0.0001 by OneWay ANOVA. Data are +/- S.E.M.