1

The RNA-binding protein RBP10 controls a regulatory cascade that defines

bloodstream-form trypanosome identity.

Elisha Mugo and Christine Clayton

DKFZ-ZMBH Alliance, Zentrum für Molekulare Biologie der Universität Heidelberg,

University of Heidelberg. Im Neuenheimer Feld 282, D69120 Heidelberg, Germany.

Abstract

Gene expression control in the pathogen *Trypanosoma brucei* relies almost

exclusively on post-transcriptional mechanisms, so RNA binding proteins must

assume the burden that is usually borne by transcription factors. *T. brucei* multiply in

the blood of mammals as bloodstream forms, and in the midgut of Tsetse flies as

procyclic forms. We show here that a single RNA-binding protein, RBP10, defines the

bloodstream-form trypanosome differentiation state. Depletion of RBP10 from

bloodstream-form trypanosomes gives cells that can grow only as procyclic forms;

conversely, expression of RBP10 in procyclic forms converts them to bloodstream

forms. RBP10 binds to procyclic-specific mRNAs containing an UAUUUUUU motif,

targeting them for translation repression and destruction. Products of RBP10 target

mRNAs include not only the major procyclic surface protein and enzymes of energy

metabolism, but also protein kinases and stage-specific RNA-binding proteins:

consequently, alterations in RBP10 trigger a regulatory cascade.

Introduction

In multicellular organisms, cell differentiation is usually unidirectional, with

stabilisation of differentiated states by epigenetic mechanisms, and reversal only in

when healthy gene regulation is disrupted. In contrast, differentiation in unicellular

organisms can be reversible, or involve cycles in which the direction of each

transition is fixed. In each case differentiation is initiated by environmental stimuli,

which activate signal transduction cascades; these in turn result in post-translational

modifications that modify transcription factor activity. The consequent changes in

mRNA synthesis, and hence protein synthesis, trigger a cascade of downstream

regulatory events, affecting additional signalling molecules, transcription factors, and

post-transcriptional modulators. Differentiation states are then stabilised by

modification of DNA and chromatin.

Kinetoplastids are unicellular flagellates, and include many pathogens of mammals

and plants, including the leishmanias and trypanosomes that cause human disease.

Their life cycles involve a series of unidirectional transitions, which are driven by

environmental changes as they move between mammalian and invertebrate hosts.

The kinetoplastid studied in this paper, *Trypanosoma brucei*, causes human sleeping

sickness in Africa and diseases of ruminants throughout the tropics. *T. brucei* multiply

as bloodstream form trypomastigotes in the blood and tissue fluids of mammals,

escaping immunity by antigenic variation of Variant Surface Glycoprotein (VSG) [1].

High cell density triggers growth arrest and differentiation to stumpy forms [2], which,

upon uptake by a Tsetse fly, convert to proliferative procyclic forms in the midgut.

Procyclic forms express EP and GPEET procyclin surface proteins [3], and have

numerous metabolic adaptations including increased mitochondrial metabolism. In

the following 2 weeks, trypanosomes migrate towards the salivary glands [4].

Epimastigotes, expressing the surface protein BARP [5], are found attached to the

proventriculus and salivary glands; here, a subset of parasites undergoes meiosis

and mating [6,7]. Finally, growth-arrested, VSG-expressing metacyclic forms are

seen swimming freely in the salivary glands; these are now ready for infection of a

new mammal. Trypanosome developmental transitions are marked by many changes

in mRNA [8-12] and protein [13-16] abundance.

Despite extensive changes in gene expression through the *T. brucei* life cycle, and

the existence of stable differentiation states, the normal paradigm for regulation does

not apply. This is because - as in other Kinetoplastid protists - transcription by *T.*

brucei RNA polymerase II is polycistronic, with no regulation at the level of individual

mRNAs. Monocistronic mRNAs are excised by 5' *trans* splicing of a 39nt capped

leader sequence, and by 3' polyadenylation [17,18]. Regulation is thus exclusively at

the levels of mRNA processing, degradation, and translation [10-12,19-22]. The

sequences required for regulation of translation and mRNA decay often lie in the 3'-

untranslated regions (3'-UTRs) of the mRNAs. Many groups of trypanosome mRNAs

are coordinately regulated [23] and in a few cases, co-regulated genes have been

shown to share regulatory elements that are bound by specific RNA-binding proteins

[24,25]. RNA-binding proteins thus assume the burden that is carried by transcription

factors in other eukaryotes, by regulating distinct mRNA subsets [19,26]. Although an

RNAi machinery exists, it is not essential for cell proliferation or differentiation [27],

and microRNAs have not been found [28].

The very abundant *VSG* and procyclin mRNAs are made, exceptionally, by RNA

polymerase I [29], and their transcription is developmentally regulated by epigenetic

mechanisms [1]. Procyclin is not subject to antigenic variation, and the presence of

even a small amount on the bloodstream-form surface would enable the

development of host immunity. Suppression of procyclin expression is therefore

essential for trypanosome pathogenicity. However, procyclin gene transcription is

only 10-fold less active in bloodstream forms than in procyclic forms [30]. This means

that additional suppressive mechanisms are required: the mRNAs are both extremely

rapidly degraded and very poorly translated [31-35], and trafficking of procyclin

proteins to the surface is suppressed [36]. A 26nt motif within the 3'-UTR of procyclin

mRNAs is required for their degradation and translational suppression [33,34].

However, until now, the protein(s) responsible for preventing procyclin expression

were unknown.

RBP10 is a cytosolic protein with a single RNA recognition motif. Proteome analyses

[16] and Western blot results [37] indicate that RBP10 is expressed in proliferating

bloodstream forms, but not in stumpy forms or procyclics; *RBP10* mRNA also has

low abundance in procyclic forms and in salivary gland trypanosomes [38]. Depletion

of RBP10 by RNA interference (RNAi) in the bloodstream form is lethal, and forced

RBP10 expression in procyclics inhibits growth [37]. Microarray analyses on cells

with RBP10 RNAi or forced expression suggested a function for RBP10 either in

stabilizing bloodstream-form specific mRNAs, or destabilizing procyclic-specific

mRNAs, but it was unclear which effects were due to growth inhibition and the direct

RNA targets of RBP10 were unknown.

In this paper we set out to identify the direct mRNA targets of RBP10 and to

determine its role in mRNA control. We discovered that the mRNAs that are bound to

RBP10 in bloodstream forms are preferentially targeted for translation inhibition and

destruction, and that cells without RBP10 can survive only by differentiating to the

procyclic form. Conversely, cells expressing RBP10 can survive only as bloodstream

forms. The presence of RBP10 thus defines bloodstream-form trypanosome identity.

Results

Attachment of the RBP10 C-terminal domain to a reporter mRNA causes
translation inhibition and mRNA degradation

Inducible expression of lambdaN-peptide-tagged RBP10 in cells containing a *CAT-*

boxB reporter will result in attachment of the RBP10 to the reporter RNA via the

lambdaN-boxB interaction. We previously showed that after 24h lambdaN-RBP10

induction, there was a decrease in boxB-tagged chloramphenicol acetyltransferase

(*CAT*) reporter expression [37,39,40]. New deletion analyses (Fig 1) revealed that the

C-terminal 88 amino acids of RBP10 was sufficient for the repressive effect (Fig 1 A,

fragment F3). To investigate the mechanism in more detail we examined whether the

CAT-boxB mRNA comigrated with polysomes in sucrose gradients. In the absence of

tethered protein, the *CAT-boxB* reporter migrated in the dense sucrose fraction with

large polysomes (Fig 1 B). With lambdaN-RBP10-F3 expression, *CAT-boxB* mRNA

was reduced in abundance and the remainder migrated towards the top of the

gradient. The control mRNA (encoding alpha-tubulin) was unaffected (Fig 1 B). The

RBP10 C-terminus thus prevents translation initiation and promotes mRNA

destruction. Interestingly, although a homologue of RBP10 is present in all

sequenced Kinetoplastids, only the RRM domain is conserved: this C-terminal

sequence is specific to *Trypanosoma* (S1 Fig).

Fig 1

Tethering of the C-terminal portion of RBP10 to a reporter mRNA inhibits its translation and causes its destruction.

A. Various fragments of RBP10 were inducibly expressed with an N-terminal lambdaN peptide and a C-terminal myc tag, in a cell line expressing a *CAT* reporter mRNA with 5 boxB sequences in the 3'-UTR. Cartoons showing the fragments are on the left. The numbers above the full-length protein are amino acid residues. The effects on the amounts *CAT* mRNA and CAT protein were measured by Northern blot and enzyme assay, respectively. Levels are expressed as mean ± standard deviation for at least 3 replicates, relative to a line with no lambdaN peptide protein (top bars)

B. Bloodstream-form trypanosomes expressing the reporter in (A), and with tetracycline-inducible expression of lamdaN-RBP10-F3, were used. The reporter with or without tethered protein is shown at the top. Lysates from cells grown without (left) or with (right) tetracycline were subjected to sucrose gradient centrifugation. The panel below the cartoons show the absorption profiles at 254 nm as the fractions were collected. The migration of *CAT* and beta-tubulin (*TUB*) mRNAs on the gradient was detected by Northern blotting; representative blots are shown. The graphs show Northern signals, expressed as the percentage of the total signal, with arithmetic mean and standard deviation for three independent biological replicates.

RBP10 can interact with an eIF4E interacting protein

We investigated the mechanism of RBP10-mediated repression by looking for interaction partners. We made a bloodstream-form cell line containing a single copy of RBP10 bearing a cleavable N-terminal affinity purification tag (S2 Fig A). After tandem affinity purification the most consistent co-purifying proteins were components of the proteasome (S1 Table). While this could be a mechanism for repression, it is also possible that some of the affinity-tagged RBP10 was targeted to the proteasome because it was incorrectly folded. The only interaction partner linked to mRNA degradation was CAF40, a component of the NOT deadenylase complex, but since no other components of the complex were present the significance of the interaction was unclear. Furthermore, V5-CAF40 pull-down failed to identify any RBP10 peptides (data not shown). To find out whether other proteins - including those of unknown function - might be important we investigated direct interactions of RBP10 by 2-hybrid screening. Nearly 80 potential partners were seen (S2 Table), of

which 40 had not been seen in other screens. This suggested that RBP10 is prone to

non-specific interactions. Potential partners included RBP10 itself, and four other

RNA binding proteins. Interestingly, 8 of the 33 specifically interacting proteins had

previously been show to suppress expression in the tethering assay [39,40]. There

was however no overlap between the two-hybrid results and those from affinity

purification.

A single fragment of a trypanosome eIF4E-interacting protein (4EIP) was scored as a

possible RBP10 partner in the two-hybrid screen. Leishmania 4EIP interacts with

eIF4E1 [41], which is one of the six trypanosomatid homologues of the cap-binding

translation initiation factor eIF4E. In *T. brucei*, both 4EIP and eIF4E1 are extremely

strong suppressors of reporter expression in the tethering assay [39,40]. We

confirmed the two-hybrid interaction between full length RBP10 and 4EIP (Fig 2 A);

and found that the interaction was mediated by the C-terminal 88 amino acids (F3)

that are active in the tethering assay. This would be consistent with RBP10 acting via

4EIP recruitment. To test the interaction *in vivo*, we used trypanosomes expressing

RBP10 that was N-terminally V5-tagged at the endogenous locus, then expressed C-

terminally myc-tagged 4EIP from a strong promoter. Immunoprecipitation of myc-

tagged 4EIP indeed pulled down a very small proportion of V5-RBP10 (Fig 2 B). It is

therefore possible that RBP10 acts by recruiting 4EIP. However, depletion of 4EIP

does not reproducibly inhibit bloodstream-form trypanosome growth ([42] and our

preliminary results), which argues against this hypothesis.

Fig 2

Interaction of RBP10 with 4E-IP

A. Two-hybrid assay showing yeast growth. Upper panels show selection against

non-interacting pairs.

B. Immunoprecipitation of 4E-IP-Myc pulls down V5-tagged RBP10. The number of

cell equivalents loaded for the eluate was 33x that of input and unbound fractions.

The inverse precipitation did not confirm the interaction, but this could be because of

an imbalance of protein abundances in the cells. (The abundance of 4E-IP is not

known).

RBP10 represses procyclic-form specific mRNAs

To analyse the role of RBP10 in gene expression in bloodstream forms, we depleted RBP10 by RNAi for 15h (S2 Fig B), before parasite proliferation and translation (S3 Fig) were affected. High-throughput cDNA sequencing (RNA-Seq) revealed nearly 200 mRNAs that changed significantly in total and/or polysomal RNA fractions (S3 Table, sheets 3-7). 36% of the mRNAs that increased after RNAi were preferentially expressed in procyclic forms, while 44% of those that decreased were preferentially expressed in bloodstream forms (Table 1 and Fig 3 A). Increases in polysomal RNA after RNAi were very slightly stronger than for total RNA. *RBP10* was the only mRNA with strongly decreased polysomal loading after RNAi, whereas about 60 mRNAs moved into polysomes (S3 Table sheet 3). 19 of these 60 mRNAs have a higher ribosome density in procyclic forms than in bloodstream forms [11], suggesting that their translation is inhibited in bloodstream forms. An example is the *COXVI* mRNA (S3 Table sheet 3, S4 Fig A,B), which is known to be developmentally regulated mainly at the level of translation [43]. Only 4 mRNAs had lower polysome occupancy after RNAi (S1 Table sheet 3). Our results almost certainly under-estimate the extent of translation changes, since we measured only whether an mRNA was in the polysomal fraction or not. For example, if an mRNA had a few ribosomes before RNAi, and more ribosomes after RNAi, this would not have been detected.

Fig 3

mRNAs bound to and/or regulated by RBP10.

Results for mRNA levels are displayed as proportional Venn diagrams made using http://jura.wi.mit.edu/bioc/tools/venn.php and http://www.eulerdiagrams.org/eulerAPE/

BS = Bloodstream form, PC = procyclic form. Categories were: Bound: Supplementary Table S3 sheet 4. PC>BS and BS>PC: at least 2x significantly regulated [10]. BS rbp10 down or up: affected by RNAi, Supplementary Table S1 sheets 6 & 7; PC+RBP10 down or up: affected by RBP10 expression in procyclics, Supplementary Table S2 sheets 6 & 7. Fischer exact test values, calculated for the dataset in the centre, are in blue italics.

A) Relationship between developmental regulation and the effects of RBP10 depletion in bloodstream forms

B) Relationship between developmental regulation and the effects of RBP10
expression in procyclic forms

C) Expression of RBP10 in procyclic forms related to the effects *rbp10* RNAi in
bloodstream forms.

D) Developmental regulation of the mRNAs bound by RBP10.

E) Effect of bloodstream-form *rbp10* RNAi on the mRNAs bound by RBP10.

D) Effect of procyclic-form RBP10 expression on the mRNAs bound by RBP10.

G) Overlaps between bound mRNAs, procyclic-specific mRNAs, and the mRNAs that
increased after *rbp10* RNAi in bloodstream forms

H) Overlaps between bound mRNAs, the mRNAs that increased after *rbp10* RNAi in
bloodstream forms, and the mRNAs that decreased after RBP10 expression in
procyclic forms

I) Abundances of RBP10-bound mRNAs compared to all other mRNAs.

Next we analysed the effect of expressing C-terminally myc-tagged RBP10 (RBP10-myc) in procyclic forms. After only 6h induction, (S2 Fig C, S4 Fig) we saw decreases in procyclic-form-specific mRNAs, and increases in bloodstream-form specific mRNAs (Fig 3 B and S4 Table). A subset of mRNAs showed opposite reactions in the bloodstream RNAi and procyclic over-expression experiments, consistent with being immediate RBP10 targets (Fig 3 C).

We had previously attempted to identify RBP10-bound mRNAs, but the methods that we used had insufficient sensitivity [37]. Since a subsequent study showed that RBP10 is indeed bound to mRNA [40], we made another attempt. This time, we used our bloodstream-form cell line in which all RBP10 bears a cleavable tag (S2 Fig A), and compared bound and unbound mRNAs by RNASeq. 400 mRNAs were identified (S5 Fig A and S5 Table, sheet 3). 12% of them are more abundant in procyclic forms

than in bloodstream forms (Fig 3 D). Consistent with the tethering results, 15% of the

RBP10 targets increased after *rbp10* RNAi in bloodstream forms, but only 1%

decreased (Fig 3 E). Only 16 target mRNAs decreased after RBP10 expression in

procyclic forms (Fig 3 F); this low number might be due to the very short duration of

RBP10-myc expression. The RBP10-bound mRNAs that are more abundant in

procyclic forms and increased after *rbp10* RNAi (Fig 3 G) are listed in Table 1: six of

them are also decreased after RBP10 expression in procyclics (Fig 3 H). Scatter

plots showing the effects of RNAi on bound RNA abundance and polysomal loading

are shown in S5 Fig C and D.

Table 1

21 mRNAs that are bound to RBP10, are more abundant (RNA PC>BS) or better

translated in procyclic forms than in bloodstream forms according to ribosome

profiling (Riboprof PC>BS), and are increased after *rbp10* RNAi in bloodstream

forms. For details see S5 Table sheet 8. The latter includes many additional mRNAs

that fulfilled fewer conditions.

id	Annotation	RNA PC>BS	RiboProf PC>BS	de-creased by RBP10 in PC
Metabolism				
9.5900	glutamate dehydrogenase (GDH)	yes	yes	no
10.2350	pyruvate dehydrogenase complex	yes	yes	no
10.4280	Complex III subunit	yes	yes	no
1.4100	cytochrome oxidase subunit IV (COXIV)	yes	yes	no
11.15550	NADH-cytochrome b5 reductase,	yes	yes	no
4.4990	ubiquinol-cytochrome c reductase	yes	yes	no
7.210	proline dehydrogenase	yes	yes	no
9.4310	tricarboxylate carrier	yes	yes	no
9.7470	purine nucleoside transporter NT10	yes	yes	yes
11.6280	pyruvate phosphate dikinase (PPDK)	yes	yes	no
11.8990	cation transporter, putative	yes	yes	no
Surface				
10.10250	Procyclins #	yes	yes	yes
7.6850	trans-sialidase (TbTS)	yes	yes	yes
8.7340	trans-sialidase	no	yes	yes
4.3500	Amastin-like protein	yes	yes	no
RNA binding				
7.2660	ZC3H20	yes	no	no
7.2680	ZC3H22	yes	yes	yes
Unknown function				
9.13200	hypothetical protein	yes	yes	yes
11.16300	hypothetical protein	yes	yes	no
7.5550	hypothetical protein	yes	yes	no
10.3300	DUF4206 (cysteine-rich)	yes	yes	no
10.2360	hypothetical protein	yes	no	no

Overall, the RBP10-bound mRNAs are significantly less abundant than unbound

mRNAs (Fig 3 I). At the proteome level [16], the products of RBP10-bound mRNAs

are enriched in proteins that increase within 2-24h of the initiation of stumpy form

differentiation, or early during conversion to procyclic forms. There is no enrichment

for proteins that appear later in differentiation, or only in mature procyclics, and

unregulated proteins or those that are decreased in procyclic forms are under-

represented (S5 Table, sheet 6).

RBP10-bound mRNAs share a UAUUUUUU motif

A comparison of the 3'-untranslated region (3'-UTR) sequences of the most reliably

RBP10-bound mRNAs with those of unbound mRNAs (S5 Table, sheet 5 and S1-S3

texts) revealed that the motif $UA(U)_6$ is highly enriched in the 3'-UTRs of RBP10

targets (Fig 4 A). It is present 1-9 times in 225 out of the 255 most strongly bound 3'-

UTRs (Fig 4 B, S1 text). Transcriptome-wide, the number of $UA(U)_6$ motifs was

strongly associated with RBP10 binding (Fig 4 C). Nevertheless, many mRNAs that

contain $UA(U)_6$ were not pulled down with RBP10. Although there are some

exceptions [44,45], RRM domains usually bind to single-stranded RNA [46-48]; the

unbound motifs may have inappropriate conformations, or they might instead be

bound by competing proteins.

Fig 4

RBP10 targets a UAUUUUUU motif

A) The 3'-UTRs of 188 mRNAs that were at least 3x enriched in both RBP10-bound

samples (Supplementary Table S5 sheet 5) were compared with those of mRNAs

that did not bind (<0.7x enrichment in both experiments) using DREME. Only 3'-

UTRs annotated in tritrypDB were used (S1 text, S2 text, S3 text). The best-scoring

motif found is shown.

B) Numbers of $UA(U)_6$ motifs in 255 manually annotated bound 3'-UTRs (S1 text, S5

Table sheet 5). One mRNA each had 7, 8 and 9 motifs. The numbers of motifs

shown here are sometimes higher than with automatically annotated 3'-UTRs from

the genome database, because some of the automatic 3'-UTRs are truncated.

C) Effect of the $UA(U)_6$ motif on RBP10 binding: median ± 25th percentiles, with 95%

confidence limits and outliers. Database 3'-UTRs were analyzed. Since these are

often truncated, and are missing for 30% of genes, the numbers of motifs are under-

estimated. Significant differences (Student t-test) are shown, For the mRNAs that

were enriched according to DeSeq (S5 Table, sheet 3), 284/353 annotated UTRs

contained UA(U)$_6$.

D) Reporters used in (E) and (F) [34]. The red bars are UAUUUUUU elements, which

are highlighted in the 26mer sequence.

E) RBP10 binding to the *EP* 3'-UTR requires the 26mer. Cells were UV-irradiated,

and RBP10 was pulled down with anti-RBP10 (upper panel, Western blot). RNAs

were detected by reverse transcription and PCR using gene -specific primers (lower

panel).

F) *rbp10* RNAi specifically affects expression of a reporter containing the 26mer. CAT

activity was measured 24h after RNAi induction.

UA(U)$_6$ was previously identified in several mRNAs encoding procyclic-specific

proteins [43]. The 26mer 3'-UTR element that is required for *EP1* procyclin mRNA

instability [33,34] contains two UA(U)$_6$ motifs (Fig 4 B) and these are single-stranded

in vivo [49], as is required for RRM domain binding. Deletion of the region containing

UA(U)$_6$ from the COXV 3'-UTR increased protein expression without much effect on

the mRNA level [43], consistent with translation control. The mRNAs encoding

phosphoglycerate kinase (PGK) constitute a third example. In bloodstream-form

trypanosomes, most PGK is encoded by the gene *PGKC* and is targeted to

microbodies called glycosomes (S6 Fig A); in procyclic forms, a different gene,

PGKB, is expressed and the enzyme is in the cytosol [50,51] (S6 Fig C). The *PGKB*

and PGKC coding regions are identical except for a C-terminal targeting sequence in

PGKC, so they were not distinguished in our initial RNASeq analysis. However, the

3'-UTRs are completely different, and determine developmental regulation [52]. The

UA(U)$_6$ motif is present twice within a 3-UTR segment that is required for suppression

of PGKB expression in bloodstream forms [53] (S1 text). Using our RNASeq pull-

down data, we compared the abundances of several 15nt sequences specific to the

PGKB and *PGKC* mRNAs and found that that *PGKB* mRNA is almost exclusively in

the bound fraction, whereas most *PGKC* mRNA is unbound. A similar analysis gave

RBP10 binding ratios of 2.8:1 for *EP1* and 1.3:1 for *GPEET* mRNAs. The *GPEET*

mRNA 3'-UTR has UA(U)$_5$ repeats, but no UA(U)$_6$, and it is not yet clear whether it is

bound by RBP10.

To test whether the 26mer was required for binding of the *EP1* 3'-UTR to RBP10, we

used cell lines that expressed chloramphenicol acetyltransferase (CAT) reporter

mRNAs with either the intact wild-type *EP1* 3'-UTR, or a version without the 26mer

(*ep1Δ26*) (Fig 4 D) [34]. Co-immunoprecipitation of the *CAT* mRNA with RBP10

depended on the presence of the 26mer (Fig 4 E). Depletion of RBP10 by RNAi also

increased expression from the *CAT-EP* reporter but not from *CAT-ep1Δ26* (Fig 4 F).

These results demonstrated both that RBP10 binds to the *EP* 3'-UTR via the 26mer

sequence, and that the 26mer is necessary for RBP10-mediated regulation of an

mRNA containing the *EP1* 3'-UTR.

Expression of RBP10 determines bloodstream form identity

To test the role of RBP10 in differentiation, we first examined conversion from the

bloodstream form to the procyclic form. Our monomorphic bloodstream forms with

rbp10 RNAi, or treated with the differentiation-inducer *cis* aconitate, are unable to

multiply under procyclic culture conditions. We therefore used differentiation-

competent trypanosomes (EATRO 1125, Antat1.1 strain). As expected [3], growth of

these cells at high density in methyl cellulose for three days resulted in expression of

the a -form marker PAD1 [54] and stumpy morphology; RBP10 was also reduced

(Fig 5 A, lanes 2 &3), consistent with its absence in the stumpy-form proteome [54].

The cells were now treated with 6mM cis-aconitate in bloodstream-form medium at

27°C for 17h. Upon transfer to procyclic medium without cis-aconitate, the cells

started to grow as procyclics within 24h (Fig 5B).

Fig 5

RBP10 depletion primes bloodstream forms for differentiation to procyclic forms

A) Expression of PAD1 and RBP10 in trypanosomes incubated at maximal density in

methyl cellulose-containing medium for 3 days (left panel), or after *rbp10* RNAi (right

panel). The RBP10 signal is weak, probably because the samples were not

denatured in order to allow PAD1 detection.

B) Growth of trypanosomes incubated at maximal density in methyl cellulose-

containing medium for 3 days, followed by cis-aconitate treatment for 17h at 27°C.

Time 0 is the time of transfer to procyclic conditions.

C) Cumulative growth curve of bloodstream-form (BS) trypanosomes with and

without RNAi. WT = wild-type (*tet* repressor only); -RNAi: RNAi cell line with no

tetracyline: +RNAi: RNAi cell line with tetracycline.

D) Expression of RBP10, PIP39, and EP procyclin were measured by Western

blotting up to 3d after tetracycline induction of RNAi.

E) 17h after RNAi induction at 37°C in bloodstream form medium (+RNAi), or

incubation at high density in procyclic medium with 6mM cis-aconitate at 27°C (+CA),

trypanosomes were placed in procyclic medium (MEM-pros) at 27°C with neither

tetracycline nor cis-aconitate. The cell densities of the cultures were monitored.

Negative controls were either wild-type or uninduced RNAi lines, without cis-

aconitate. Cell densities are shown; they started to increase at 72h.

F) Protein expression after transfer to procyclic medium at 27°C after 17h pre-

treatment as in (E).

G) Immunofluorescence: Cell morphology and expression of GPEET procyclin after

3d cultivation in procyclic medium. DNA is stained with DAPI.

H) The cartoon shows the way in which the position of the kinetoplast, relative to the

nucleus (P-K/K-N), was measured. The box plot shows the median with 25th

percentiles and 95% confidence limits. Some outliers have been deleted. The full

results, together with the numbers of trypanosomes measured, are in S7 Fig A.

BS - normal bloodstream forms; PC- normal procyclic forms; CA - 17h cis-aconitate

pretreatment; RNAi or Ri; 17h *rbp10* RNAi induction; +RBP10 - induced expression

of RBP10 for 2 days. There were no significant differences between the 3-day and 6-

day differentiating populations. All other pairs were significantly different (Student t-

test, $p < 0.05$).

Induction of *rbp10* RNAi in differentiation-competent cells in log phase resulted in

growth inhibition (Fig 5 C), weak PAD1 expression (Fig 5 A, lanes 6-8) and, within

17h, expression of the differentiation-regulating phosphatase PIP39 [55] (Fig 5 D).

After 17h RNAi, transfer of the cells to procyclic-form medium at 27°C (without cis-

aconitate) allowed them to survive and eventually start to grow (Fig 5 E), with

expression of EP procyclin (Fig 5 F, lanes 1-7) and GPEET procyclin (Fig 5 G)

proteins. As expected, cells without RNAi induction died (Fig 5 E). The RNAi

experiment had to be done without high density pre-incubation in methyl cellulose

since RNAi can only be induced in growing trypanosomes. As a control, we therefore

grew differentiation-competent cells to maximum density in liquid medium, and

treated them with cis-aconitate for 17h at 27°C [56] before transfer to procyclic

medium without cis-aconitate, also at 27°C. In these cells, weak RBP10 expression

persisted for 2 days (Fig 5 F, lanes 8-13) but the growth and procyclin expression

kinetics were similar to those seen after *rbp10* RNAi (Fig 5 E, F, G).

The rather slow resumption of growth after both *rbp10* RNAi and 17h cis-aconitate

treatment, and the persistence of RBP10 in the cis-aconitate experiment, suggested

that the population might be a mixture of differentiation-competent and incompetent

cells. Indeed, both differentiating cultures were mixtures of normal-looking,

proliferating, and dying or abnormal forms, and even after 3 days in procyclic culture,

about 9% of the cells in both cultures retained bloodstream-form morphology and

PGK localization (S6 Fig D). In bloodstream forms, the kinetoplast is at the posterior

end of the cell (Fig 5 G), which means that the distance between the posterior and

the kinetoplast (P-K) is much less than the distance from the kinetoplast to the

nucleus (K-N) (Fig 5 H and S7 Fig A, B; "BS"). Procyclic forms are longer and the

kinetoplast is nearer to the nucleus (Fig 5 G), giving a much higher P-K/K-N ratio (Fig

5 H and S7 Fig A, B; "PC"). The differentiating cultures showed very heterogeneous

P-K/K-N ratios, with medians that were intermediate between those of bloodstream

and procyclic forms (Fig 5 H, S7 Fig A-C). As expected, cells with low ratios typical of

bloodstream forms were preferentially GPEET negative (S7 Fig D,E).

We next inducibly expressed RBP10-myc in differentiation-competent procyclic

forms. For the results shown, we made bloodstream forms with inducible RBP10-

myc, differentiated them into procyclic forms, grew these for more than 3 months,

and then induced RBP10 expression. However similar results were obtained by

differentiating bloodstream forms into procyclic forms, and then making three new

cell lines with the inducible RBP10-myc plasmid. After RBP10-myc induction, cell

growth was inhibited, as expected [37] (Fig 6 A). After 48h, expression of procyclin

was strongly reduced, alternative oxidase increased, and untagged RBP10 was

detected (Fig 6 B, C). At this point the cell population was very heterogeneous. Some

cells had normal procyclic morphology and expressed EP and GPEET procyclin;

some were clearly abnormal; while another subset had clear bloodstream-form

morphology (Fig 6D). This was reflected in the P-K/K-N ratios, which ranged from

procyclic to bloodstream-form patterns (Fig 5 G, S7 Fig F). Flow cytometry analysis

revealed that about 50% of cells had decreased, or absent, procyclin expression (Fig

6 E, F). Cells with bloodstream-form morphology were usually (but not always)

procyclin negative (S7 Fig G). Notably, *VSG* mRNA was present (Fig 6 C). We do not

know which VSGs are expressed in the bloodstream-form-morphology cells, but a

subset of them stained with a polyclonal antibody against VSG117 (Fig 6 D, G, H).

PGK staining in the bloodstream-form-like cells seemed reduced and somewhat

more punctate than in procyclic forms, but had not yet assumed a purely glycosomal

pattern (S6 Fig E). Remarkably, the RBP10-induced cells could now survive only if

they were transferred to bloodstream-form growth conditions (Figure 6A). After 6

days in these conditions, all surviving cells had bloodstream-form morphology and

60% of them cross-reacted with the anti-VSG117 antibody.

 Fig 6

RBP10 expression in procyclic forms causes differentiation to bloodstream forms

A) Cell counts in a typical experiment. Expression of RBP10-myc was induced using

tetracycline at 27°C in procyclic-form medium. The left-hand panel is a cumulative

growth curve. After 48h cells were transferred to bloodstream-form medium at 37°C;

in the right-hand panel cell densities are shown.

B) Expression of EP procyclin and RBP10 after induction of RBP10-myc expression.

Cells were cultured in procyclic medium at 27°C. Expression of EP Procyclin protein

decreased after 48h of RBP10-myc induction, while untagged RBP10 becomes

detectable.

C) As in (B) except that culture continued for 4 days. Proteins that are expressed

more in bloodstream forms than in procyclic forms were detected by Western

blotting. TAO is trypanosome alternative oxidase; HNRNPH is an RNA-binding

protein. *VSG* mRNA was detected by RT-PCR using a spliced leader primer and a

primer that hybridizes to a conserved region within the *VSG* 3'-UTR.

D) Trypanosomes were taken after 48h RBP10-myc induction. They were stained for

EP procyclin and phosphorylated GPEET procyclin (green), VSG117 (magenta) and

DNA (cyan). The stain is overlaid with differential interference contrast (grey). The

panel shows a typical green procyclic form, a bloodstream form with two kinetoplasts,

and a third cell which resembles a bloodstream form (terminal kinetoplast) but was

not stained by anti-VSG117. This cell may express a VSG that does not react with

the anti-VSG117 antibodies.

E) Quantitation of surface EP procyclin by flow cytometry, comparing cells with (day

2) and without (day 0) induced RBP10-myc expression.

F) Quantitation of surface phospho-GPEET procyclin by flow cytometry, comparing

cells with (day 2) and without (day 0) induced RBP10-myc expression.

G) Quantitation of staining with anti-VSG117 by flow cytometry, comparing cells with

(day 2) and without (day 0) induced RBP10-myc expression.

H) Measurement of surface protein expression by FACSscan. Values for the different

windows are mean and standard deviation from 4 replicates. Similar results were

obtained by manual counting of stained smears.

The conversion of procyclic forms to bloodstream forms was highly reproducible. It

was not due to persistence of a few bloodstream forms in the procyclic cultures,

since survival always depended on induced RBP10 expression (Fig 6 A). RBP10

induction for two days was required; this was also the time needed to detect native

RBP10 expression by Western blotting. Moreover, our preliminary results indicate

that transient expression of RBP10 from an episome is sufficient to generate

bloodstream forms. In contrast, attempts to obtain bloodstream forms by RBP10

expression in Lister 427 cells that have been in procyclic form culture for more than

two decades were not successful.

Interestingly, epimastigote-like cells were never seen after induction of RBP10-myc

expression in our differentiation-competent cells (24h or 48h), and the epimastigote-

specific surface protein BARP was not detected by Western blotting or

immunofluorescence. Although we may have missed transient formation of

epimastigotes, this is unlikely since the behaviour of the trypanosomes was clearly

not synchronous. Moreover, the presence of dividing cells with bloodstream-form

morphology, suggested that RBP10 induction might be causing a direct conversion of

procyclic forms to bloodstream forms, "jumping" past the non-dividing metacyclic

form. We therefore compared the mRNA changes after 6h RBP10-myc induction with

those found in salivary gland parasites (a mixture of epimastigotes and metacyclics)

(S4 Table, sheets 5 and 6). The transcriptome comparisons and morphological

analysis suggested that after 2 days of RBP10 expression the cell population might

include both metacyclic and bloodstream forms. Interestingly, however, the mRNA

encoding the meiosis protein MND1, which is specific to a pre-metacyclic

intermediate form, is bound by RBP10 and *decreases* upon RBP10 induction.

Evidence so far thus suggests that RBP10 induction causes a developmental switch

that by-passes the epimastigote stage.

Discussion

Our results indicate that the presence of RBP10 defines the identity of a

trypanosome as a replication-competent bloodstream form.

RBP10 binds to procyclic-specific mRNAs with a UA(U)$_6$ motif in the 3'-UTR,

suppressing translation and causing mRNA destruction. However, many RNAs that

are bound by RBP10 did not change in either abundance, or in the percentage in

polysomes, after RBP10 RNAi or forced expression. As noted above, our assay did

not measure ribosome density, so it is likely that many changes in translation went

undetected. However, our results are completely consistent with work on other RNA-
binding proteins: usually, only some of the bound mRNAs change after the protein is
removed (e.g. [57]). For most mRNAs, many different proteins can bind along the
same 3'-UTR - a sequence of 300nt is likely to bind at least twenty proteins - and it is
their combinatorial effects that determine mRNA behaviour.

RBP10 target mRNAs include several that have been studied in considerable detail,
including those encoding cytochrome complexes and the procyclins. However, the
direct effects of RBP10 are by no means sufficient to explain its far-reaching
influence, since many mRNAs that change in translation or abundance after RBP10
depletion are not bound by RBP10. This suggests that RBP10 depletion may initiate
a cascade of events. Our results show how this could occur. Notably, the direct
RBP10 targets include the mRNAs encoding three potential RNA-binding proteins,
ZC3H20, ZC3H21, and ZC3H22, These proteins are normally expressed only in
procyclic forms [15,16,58]. ZC3H20 binds to, and stabilises, at least two procyclic-
specific mRNAs [58], of which one, *MCP12*, is not bound by RBP10 but was
increased after *rbp10* RNAi. ZC3H22 suppresses expression in the tethering assay
[39], and is required for proliferation of procyclic forms [15,42]. Suppression of
ZC3H21 and ZC3H22 expression is clearly very important to bloodstream-form
trypanosome survival: their mRNA 3-UTRs contain, respectively, no fewer than 5 and
7 RBP10 binding motifs.

In addition to controlling ZC3H20-22, RBP10 binds to and represses mRNAs
encoding two procyclic-specific protein kinases (Tb927.8.6490, Tb927.11.15010) and
one procyclic specific protein phosphatase (Tb927.10.8050). Also bound by RBP10,
but with no detected effect of RNAi, are the transcripts encoding protein kinases
RDK1 and NRKA. NRKA [15] is up-regulated after the onset of stumpy form
differentiation; it is possible that RBP10 indeed regulates its translation but we did not
detect this because of the insensitivity of our assay. RDK1 is a differentiation
repressor [59] and is more abundant in bloodstream forms, making regulation by
RBP10 unlikely. Similar to RBP10, depletion of *RDK1* primes bloodstream forms to
differentiate to procyclic forms. More than 20 mRNAs show similar changes after
RNAi targeting either *RDK1* or *RBP10*, and 16 of these are both bound and regulated
by RBP10. Surprisingly, the *RBP10* transcript was not significantly affected after
RDK1 RNAi; however, the protein level and phosphorylation status of RBP10 were
not determined.

The changes in RNA regulators, protein kinases and phosphatases, and perhaps
also the changes in metabolism that are directly induced by RBP10 loss (e.g. [60])

will trigger downstream effects on signaling pathways. Such secondary effects of

rbp10 RNAi must include the increase in PIP39 (Tb927.9.6090), a key regulator of

differentiation [55], and decreases in mRNAs encoding the RNA-binding proteins

PUF11, RBP9, ZC3H31, ZC3H46, ZC3H48, DRBD5 and HNRNPH (S1 Table, sheet

7). These in turn will trigger further changes in gene expression.

So far, the only condition known to allow differentiation of *T. brucei* procyclic forms to

bloodstream forms *in vitro* is induced expression of the putative RNA-binding protein

RBP6. This mimics the natural differentiation pathway, with formation of

epimastigotes within 24h and appearance of VSG-expressing metacyclic forms 4-5

days later [61]. *RBP6* mRNA has low abundance in both procyclic and bloodstream

forms, and is maximal in salivary gland parasites [38]. RBP6 action is therefore

transient, although it might possibly define the epimastigote form; its mechanism of

action is unknown. Depletion of the RNA-binding protein DRBD18 from procyclic

forms caused a partial switch to a bloodstream-form or epimastigote expression

pattern; the mRNAs that increased included those encoding RBP6 and RBP10, but

the ability to differentiate further was not tested [62]. Heat shock can also cause

small increases in epimastigote-specific transcripts [63]. The rapid conversion of

procyclic forms to bloodstream forms by RBP10 expression, in contrast, might be

regarded as a form of *trans*-differentiation since the intermediate epimastigote stage

was not detected. In animal cells, *trans*-differentiation can be achieved through

changes in expression of transcription factors, either by direct manipulation or by

changing expression of microRNAs that control transcription factor expression [64-

66]. However, these changes have so far always been unidirectional. *Trans*

differentiation that works in both directions according to the presence or absence of a

single protein has, to our knowledge, not previously been observed.

It has long been known that stumpy forms cannot revert to long-slender bloodstream

forms, and recent results indicate that the differentiation of stumpy forms to

procyclics is an irreversible bi-stable switch [15]. We suggest that expression of

RBP10 governs a bi-stable switch during the transition from bloodstream forms to

stumpy forms, and from procyclic forms to bloodstream forms. The bi-stable

character can be explained by the fact that RBP10 directly suppresses procyclic-

specific post-transcriptional regulators. While RBP10 is present, the bloodstream-

form state is self-reinforcing; loss of RBP10 enables procyclic regulators to be

expressed, while expression of RBP10 in procyclic forms tips the balance towards a

bloodstream-form expression pattern. In conclusion, our results demonstrate how

differentiation states can be maintained by a post-transcriptional cascade.

Materials and methods

Parasite culture and plasmid constructs

T. brucei Lister 427 [67] and pleomorphic AnTat 1.1 (EATRO 1125) [56] cells expressing the Tet-repressor were used. Lister 427 were used for all experiments except those shown in Figs 5 and 6. Bloodstream form parasites were cultured in HMIi-9 medium supplemented with 10% fetal bovine serum at 37^{o}C with 5% CO2. The procyclic forms were grown at 27^{o}C in MEM-Pros medium supplemented with 10% heat inactivated fetal bovine serum. Stable cell lines were generated and maintained as described in [56,67] except that selection of the pleomorphic cells was done with 8μg/ml hygromycin and 2μg/ml blasticidin. The tetracycline inducible constructs for *RBP10* RNAi and over expression were described in [37]. All new constructs and oligonucleotides are in S6 Table. A bloodstream-form cell line expressing all RBP10 with a tandem affinity purification tag at the N-terminus was generated by inserting the tag at the 5' end of one endogenous *RBP10* open reading frame (pHD2506) and deletion of the second copy (pHD2061). Plasmids for the tethering assays are in S6 Table. The tethering constructs were separately transfected in a cell line constitutively expressing the *CAT* reporter with 5 copies of *boxB* preceding the *ACT* 3' UTR. Expression of the fusion protein was induced for 24h using tetracycline (100ng/ml) and the CAT assays carried out as described in [25].

Cross-linking and RNA immunoprecipitation

3×10^{9} bloodstream form cells expressing TAP-RBP10 were irradiated using UV (400 mj/cm2), washed in cold PBS and the cell pellet snap frozen in liquid nitrogen. The RNA immunoprecipitation was done as described in [24]. Briefly, the extracts were incubated with the beads, and the unbound fraction was collected. After washing, bound RBP10 was eluted using TEV protease. After digestion with 50μg proteinase K at 42°C for 15 minutes to reduce cross-linked protein, RNA was isolated from both the bound and unbound fractions using Trifast reagent (Peqlab, GMBH). To assess the quality of the purified RNA, an aliquot of the sample was analysed by Northern blotting and the blot hybridized with a splice leader probe. Total RNA from the unbound fraction was depleted of ribosomal RNA (rRNA) using RNAse H as described in [68] except that a cocktail of 50-base DNA oligos complementary to trypanosome rRNAs was used [63]. The recovered RNA from both bound and unbound samples was then analysed by RNA-Seq.

Tandem affinity purification (TAP) and genome-wide yeast two-hybrid screen

Approximately 2×10^{10} bloodstream form cells expressing in-situ TAP-RBP10 were harvested and used for TAP as previously described [69]. Three biological replicates were done, with or without RNAse A treatment; triplicate results from a cell line with inducible GFP-TAP served as the background control. The eluate was separated on a 10% SDS-polyacrylamide gel for only 2 cm and the proteins visualized by colloidal Coomassie staining. The gel area was excised and analysed by mass spectroscopy. For the yeast two-hybrid screen, the RBP10 open reading frame was cloned into the pGBKT7 plasmid (pHD2361). The construct was used as a bait to screen a library made up of *T. brucei* random genomic DNA fragments. The screening, high throughput sequencing and analysis were done as previously described [70].

Polysome fractionation

Approximately $3-5 \times 10^8$ cells were collected by centrifugation (850 x g, 10 minutes, $21^{o}C$), and then treated in serum free media with 100µg/ml cycloheximide for 7 minutes at room temperature. The pellet was washed in 1 ml ice cold PBS, lysed in 350µl lysis buffer (20mM Tris pH7.5, 20mM KCl, 2mM MgCl2, 1mM DTT, 1000u RNasin (Promega), 10µg/ml leupeptin, 100µg/ml cycloheximide, 1× complete protease Inhibitor without EDTA (Roche), 0.2% (vol/vol) IGEPAL) by passing 15-30 times through a 21G needle, followed by centrifugation (15000 x g, 10 minutes, $4^{o}C$) to clear the lysate. KCl was adjusted to 120 mM and the clarified lysate loaded on top of a 4 ml continuous linear 15-50% sucrose (w/v) gradient in polysome buffer (20mM Tris pH7.5, 120mM KCl, 2mM MgCl2, 1mM DTT, 10µg/ml leupeptin, 100µg/ml cycloheximide). After 2 hours of ultracentrifugation (40000 rpm, $4^{o}C$; Beckman SW60 rotor), 400µl fractions were collected using Teledyne Isco Foxy Jr. gradient fractionator system and RNA was isolated using Trifast reagent (Peqlab, GMBH). In the case of RBP10 tethering, lambdaN-RBP10 was induced (24h) and the distribution of the *CAT* reporter and *alpha-tubulin* mRNAs in the collected fractions detected by Northern blotting as described in [71].

For RNASeq, bloodstream form cells plus or minus *RBP10* RNAi for 15h and, the procyclic form cells with or without RBP10-myc over-expression for 6h were used. In this case 250µl fractions were collected from each gradient and RNA was isolated after pooling the fractions into two groups; i) lighter fractions including monosomes, ii) denser fractions with at least two ribosomes. The amount of mRNA in the pooled samples was assessed by Northern blotting using splice leader RNA as probe. Also, total RNA was prepared from the input samples (~10% of total cell lysate) to quantify

645 changes in the steady state mRNAs levels. All samples were depleted of rRNAs (as
646 above) prior to analysis by RNA-seq.

High throughput RNA sequencing and bioinformatic analysis

RNA-seq was done at the CellNetworks Deep Sequencing Core Facility at the
University of Heidelberg. For library preparation, NEBNext Ultra RNA Library Prep Kit
for Illumina (New England BioLabs Inc.) was used. The libraries were multiplexed (6
samples per lane) and sequenced with a HiSeq 2000 system, generating 50 bp
single-end sequencing reads.
The quality of the raw sequencing data was checked using FastQC
(http://www.bioinformatics.babraham.ac.uk/projects/fastqc), and the sequencing
primers removed using Cutadapt [72]. The data was aligned to the *T. brucei* TREU
927 reference genome using Bowtie [73], then sorted and indexed using SAMtools
[74]. Reads aligning to open reading frames of the TREU 927 genome were counted
using custom python scripts. Analysis for differentially expressed genes was done in
R using the DESeq package [75] with an adjusted p-value cut-off of 0.05.
Comparative analysis was limited to a list of unique genes modified from [9]. Gene
annotations are manually updated versions of those in TritrypDB
(http://tritrypdb.org/tritrypdb/), and categories were assigned manually. Statistical
analysis was done in R. The 3' UTR motif enrichment search was done using
DREME [76]; annotated 3' UTRs sequences were downloaded from tritrypDB and we
considered only the mRNAs with 3' UTR >20 nt. Manual 3'-UTR annotation was done
using the RNASeq reads and poly(A) site data [9,11,77] in tritrypDB.

Analysis of 3'-UTR reporters

Bloodstream form cells expressing the *CAT* reporter with either full length *EP1* 3'
UTR (pHD1610) or a mutant version (pHD1611) lacking the 26mer (*EP1Δ26*)
instability element [34] were used for RNA immunoprecipitation . For the pull down,
4×10^8 cells (without the stem-loop) were irradiated using UV (400 mj/cm^2), washed in
cold PBS and the cell pellet lysed in 350µl lysis buffer (10mM Tris pH 7.5, 10mM
NaCl, 1000u RNasin (Promega), 1× complete protease Inhibitor without EDTA
(Roche), 0.1% IGEPAL) by passing 15 times through a 21G needle. The lysate was
cleared by centrifugation at 15000g for 10 minutes at 4^{o}C, NaCl was adjusted to
150mM followed by incubation with 50µl anti-RBP10 [37] coupled agarose beads for
2 hours at 4^{o}C. After washing the beads 5 times with IPP150 buffer (10mM Tris pH
7.5, 150mM NaCl, 0.1% IGEPAL), the beads were treated with 20µg proteinase K at
42^{o}C for 15 minutes and RNA was isolated from both bound and unbound fractions

using Trifast reagent (peqlab, GMBH). Equal amounts of the recovered RNA (eluate

and flow through fractions) were reverse-transcribed using RevertAid First Strand

cDNA Synthesis Kit (Thermal Scientific) according to the manufacturer's instructions.

2µl of the cDNA was used as template in a 50µl PCR reaction to detect the *CAT* and

alpha-tubulin genes. PCR was done using Q5 DNA polymerase and buffer (NEB)

with 0.5µm of the following primers, for *CAT* (CZ5725; CZ689) and *alpha-tubulin*

(CZ5725; CZ6168); the forward primer is the same for both since it anneals to the

splice leader. Aliquots (10µl) were removed after 27, 31 and 36 cycles and analysed

by agarose gel electrophoresis.

To determine if the regulation of the *EP* mRNA by RBP10 depends on the 26mer

instability element, a stem-loop construct (pHD1984) targeting *RBP10* was

transfected into the two cell lines [37]. *RBP10* RNAi was induced using 100ng/ml

tetracycline for 17h or 24h and CAT activity was measured.

RBP10 and trypanosome differentiation

For bloodstream-procyclic form conversion, pleomorphic AnTat 1.1 bloodstream form

cells with stem-loop RNAI (pHD1984) targeting *RBP10* were used. 17h after RNAi

induction, the cells were pelleted, resuspended (~8x10^5 cells/ml) in procyclic form

(MEM-pros) medium and incubated at 27^{o}C. As positive controls, wild type or

uninduced cells (2x10^6 cells/ml) were treated with 6mM cis-aconitate (Sigma) at

27^{o}C; after 17h the cells were transferred into procyclic form media (~8x10^5 cells/ml)

and maintained at 27^{o}C. Samples were taken at different times for Western blotting

(~5x10^6) and for morphological analysis.

To convert procyclic to bloodstream forms, AniTat 1.1 bloodstream form cells with an

inducible RBP10-myc construct (pHD2098) were differentiated to procyclic forms

using cis-aconitate as described above. The cells were cultured in presence of

hygromycin (8µg/ml) and phleomycin (0.2µg/ml) for more than 3 months to generate

well-established procyclic forms. RBP10-myc was induced using 100ng/ml

tetracycline. Marker proteins were detected by Western blot. VSG transcripts were

detected by semi-quantitative RT-PCR as previously described [61] using primers

CZ6308/CZ6309. To obtain growing bloodstream forms, RBP10-myc was induced for

48 hours, the cells were pelleted, resuspended (2x10^5 cells/ml) in HMI-9 medium and

then incubated at 37^{o}C with 5% CO_2. The cell density was monitored for 6 days or

more; wild type or uninduced cells served as control.

Morphological analysis

To stain surface proteins, dried smears were fixed with 100% methanol at -20°C for 15 minutes, rehydrated in 1x PBS for 15 minutes, blocked for 20 minutes with 20% FCS, then labeled with mouse anti-EP (1:500; Cedarlane, Canada), mouse anti-phospho-GPEET (1:500; Cedarlane, Canada) and rabbit anti-VSG-117 (1:500, from either G. Cross or P. Overath). Anti-BARP staining was done as described in [5], using a procyclic cell line with the *BARP* open reading frame in the procyclin locus as a positive control. For PGK staining, $2x10^6$ cells were collected by centrifugation, re-suspended, fixed in 4% paraformadehyde (in 1x PBS) for 18 minutes, sedimented again for 2 minutes, re-suspended in PBS and allowed to settle on poly-L-lysine coated slides for 30 minutes. Before staining, slides were blocked with 20% fetal calf serum (1x PBS) for 20 minutes. Cells were permeabilised with 0.2% (v/v) Triton-X 100 (in 1xPBS) for 15 minutes at room temperature, washed twice then incubated for one hour with rabbit anti PGK antibody (1:1500, [78]). The cells were washed 3 times before being stained with fluor-conjugated secondary antibody (1:500; mouse-Cy3 or rabbit-Alexa-488; Molecular probes, Eugene). Cellular DNA was stained with 100ng/ml DAPI in 1x PBS for 15 minutes. Images were taken using Olympus Cell-R microscope. Random fields were photographed (by E.M.) and analysed (by C.C.) using Fiji [79]. We attempted blind analysis but this was not possible because the differences between samples were too obvious.

Flow cytometry

Approximately $5x10^6$ cells were fixed with 2% formaldehyde/0.05% glutaraldehyde at 4°C for at least 1 hour. The cells were pelleted, washed twice with PBS then incubated with 200µl (2% BSA in PBS) mouse anti-EP (Cedarlane, Canada; 1:500), anti phosph-GPEET (Cedarlane, Canada; 1:500), or rabbit anti-VSG-117 (1:500) for one hour on ice. After washing twice, the cells were stained with the secondary antibody (1:500; mouse-Cy5 or rabbit-Alexa-488; Molecular probes, Eugene) for one hour. Cells stained only with the secondary antibody and the unstained cells served as the negative controls. Flow cytometry was performed with BD FACSCanto II flow cytometer and the data analysed using FlowJo software (TreeStar Inc.).

Western blotting

Western blotting was done as previously described [71]. Antibodies were: anti-RBP10 (rat, 1:2000, [37]), anti-HNRNPH (rabbit,1:5000, [20]), anti-TAO (rabbit,1:100, [78]), anti-EP (mouse,1:2000, Cedarlane, Canada), anti-phospho-GPEET (mouse,1:2000, Cedarlane, Canada), anti-BARP (rabbit,1:2500, [5]), and anti-PIP39

748 (rabbit,1:1000, [55]) and anti-PAD1 (rabbit,1:1000, [54] with electrophoresis under

749 non-denaturing conditions) .

Data availability.

751 All sequence datasets are available at Array Express with the accession number E-
752 MTAB-4564.

Ethical approval

754 The research did not involve animals or humans and did not require ethical approval.

Acknowledgements

For antibodies we thank Keith Matthews (Edinburgh University, Scotland; PIP39 and

PAD1), Isabel Roditi (University of Bern, Switzerland, BARP); Shula Michaeli (Bar-

Ilan University, Israel, anti-HNRNPH), Paul MIchels (anti-PGK, Edinburgh University,

Scotland) and Minu Chauduri (Meharry Medical College, Nashville, Tennessee, USA,

AOX). The anti-VSG117 antibody was either from Peter Overath (Max-Planck Institut,

Tübingen, Germany) or George Cross (Rockefeller University, USA). We also thank

Keith, Isabel and Shula for useful discussions. All RNA-Seq libraries were prepared

by David Ibberson at Bioquant, Heidelberg, and sequenced at EMBL. We thank

Claudia Helbig and Ute Leibfried for technical assistance, Dorothea Droll for the 4EIP

yeast 2-hybrid constructs, and Kevin Leiss for counting the numbers of elements in

all 3'-UTRs.

References

1. Horn D (2014) Antigenic variation in African trypanosomes. Mol Biochem Parasitol
195: 123-129.
2. Vassella E, Reuner B, Yutzy B, Boshart M (1997) Differentiation of African
trypanosomes is controlled by a density sensing mechanism which signals
cell cycle arrest via the cAMP pathway. J Cell Sci 110: 2661-2671.
3. Vassella E, Acosta-Serrano A, Studer E, Lee SH, Englund PT, et al. (2001)
Multiple procyclin isoforms are expressed differentially during the
development of insect forms of Trypanosoma brucei. J Mol Biol 312: 597-607.
4. Rotureau B, Van Den Abbeele J (2013) Through the dark continent: African
trypanosome development in the tsetse fly. Front Cell Infect Microbiol 3: 53.
5. Urwyler S, Studer E, Renggli CK, Roditi I (2007) A family of stage-specific alanine-
rich proteins on the surface of epimastigote forms of Trypanosoma brucei.
Mol Microbiol 63: 218-228.
6. Peacock L, Bailey M, Carrington M, Gibson W (2013) Meiosis and Haploid
Gametes in the Pathogen Trypanosoma brucei. Curr Biol.

7. Peacock L, Ferris V, Sharma R, Sunter J, Bailey M, et al. (2011) Identification of
the meiotic life cycle stage of *Trypanosoma brucei* in the tsetse fly. Proc Natl
Acad Sci U S A 108: 3671-3676.
8. Nilsson D, Gunasekera K, Mani J, Osteras M, Farinelli L, et al. (2010) Spliced
leader trapping reveals widespread alternative splicing patterns in the highly
dynamic transcriptome of *Trypanosoma brucei*. PLoS Pathog 6: e1001037.
9. Siegel T, Hekstra D, Wang X, Dewell S, Cross G (2010) Genome-wide analysis of
mRNA abundance in two life-cycle stages of *Trypanosoma brucei* and
identification of splicing and polyadenylation sites. Nucleic Acids Res 38:
4946-4957.
10. Fadda A, Ryten M, Droll D, Rojas F, Färber V, et al. (2014) Transcriptome-wide
analysis of mRNA decay reveals complex degradation kinetics and suggests
a role for co-transcriptional degradation in determining mRNA levels. Mol
Microbiol 94: 307-326.
11. Jensen BC, Ramasamy G, Vasconcelos EJ, Ingolia NT, Myler PJ, et al. (2014)
Extensive stage-regulation of translation revealed by ribosome profiling of
Trypanosoma brucei. BMC Genomics 15: 911.
12. Vasquez JJ, Hon CC, Vanselow JT, Schlosser A, Siegel TN (2014) Comparative
ribosome profiling reveals extensive translational complexity in different
Trypanosoma brucei life cycle stages. Nucleic Acids Res 42: 3623-3637.
13. Gunasekera K, Wuthrich D, Braga-Lagache S, Heller M, Ochsenreiter T (2012)
Proteome remodelling during development from blood to insect-form
Trypanosoma brucei quantified by SILAC and mass spectrometry. BMC
Genomics 13: 556.
14. Urbaniak MD, Guther ML, Ferguson MA (2012) Comparative SILAC proteomic
analysis of *Trypanosoma brucei* bloodstream and procyclic lifecycle stages.
PLoS One 7: e36619.
15. Domingo-Sananes MR, Szoor B, Ferguson MA, Urbaniak MD, Matthews KR
(2015) Molecular control of irreversible bistability during trypanosome
developmental commitment. J Cell Biol 211: 455-468.
16. Dejung M, Subota I, Bucerius F, Dindar G, Freiwald A, et al. (2016) Quantitative
proteomics uncovers novel factors involved in developmental dfferentiation of
Trypanosoma brucei. PLoS Pathog 12: e1005439.
17. Michaeli S (2011) Trans-splicing in trypanosomes: machinery and its impact on
the parasite transcriptome. Future Microbiol 6: 459-474.
18. Clayton C, Michaeli S (2011) 3' processing in protists. Wiley Interdiscip Rev RNA
2: 247-255.
19. Clayton C (2013) The regulation of trypanosome gene expression by RNA-
binding proteins. PLoS Pathog 9: e1003680.
20. Gupta SK, Kosti I, Plaut G, Pivko A, Tkacz ID, et al. (2013) The hnRNP F/H
homologue of *Trypanosoma brucei* is differentially expressed in the two life
cycle stages of the parasite and regulates splicing and mRNA stability.
Nucleic Acids Res 41: 6577-6594.
21. Gupta SK, Chikne V, Eliaz D, Tkacz ID, Naboishchikov I, et al. (2014) Two
splicing factors carrying serine-arginine motifs, TSR1 and TSR1IP, regulate
splicing, mRNA stability, and rRNA processing in Trypanosoma brucei. RNA
Biol 11.
22. Antwi E, Haanstra J, Ramasamy G, Jensen B, Droll D, et al. (2016) Integrative
analysis of the *Trypanosoma brucei* gene expression cascade predicts
differential regulation of mRNA processing and unusual control of ribosomal
protein expression. BMC Genomics 17: 306.
23. Queiroz R, Benz C, Fellenberg K, V S, Hoheisel J, et al. (2009) Transcriptome
analysis of differentiating trypanosomes reveals the existence of multiple
post-transcriptional regulons. BMC Genomics 10: 495.

24. Archer SK, van Luu D, de Queiroz R, Brems S, Clayton CE (2009) *Trypanosoma*
brucei PUF9 regulates mRNAs for proteins involved in replicative processes
over the cell cycle. PLoS Pathog 5: e1000565.
25. Droll D, Minia I, Fadda A, Singh A, Stewart M, et al. (2013) Post-transcriptional
regulation of the trypanosome heat shock response by a zinc finger protein.
PLoS Pathog 9: e1003286.
26. Kolev NG, Ullu E, Tschudi C (2014) The emerging role of RNA-binding proteins in
the life cycle of *Trypanosoma brucei*. Cell Microbiol 16: 482-489.
27. Janzen CJ, van Deursen, F., Shi H, Cross GAM, Matthews KR, et al. (2006)
Expression site silencing and life-cycle progression appear normal in
Argonaute1-deficient *Trypanosoma brucei* Mol Biochem Parasitol 14: 102-
107.
28. Tschudi C, Shi H, Franklin J, Ullu E (2012) Small interfering RNA-producing loci
in the ancient parasitic eukaryote *Trypanosoma brucei*. BMC Genomics 13:
427.
29. Gunzl A, Bruderer T, Laufer G, Schimanski B, Tu LC, et al. (2003) RNA
polymerase I transcribes procyclin genes and variant surface glycoprotein
gene expression sites in *Trypanosoma brucei.* Eukaryot Cell 2: 542-551.
30. Biebinger S, Rettenmaier S, Flaspohler J, Hartmann C, Peña-Diaz J, et al. (1996)
The PARP promoter of *Trypanosoma brucei* is developmentally regulated in a
chromosomal context. Nucleic Acids Res 24: 1202-1211.
31. Furger A, Schürch N, Kurath U, Roditi I (1997) Elements in the 3' untranslated
region of procyclin mRNA regulate expression in insect forms of
Trypanosoma brucei by modulating RNA stability and translation. Mol Cell
Biol 17: 4372-4380.
32. Hehl A, Vassella E, Braun R, Roditi I (1994) A conserved stem-loop structure in
the 3' untranslated region of procyclin mRNAs regulates expression in
Trypanosoma brucei. Proc Natl Acad Sci USA 91: 370-374.
33. Schürch N, Furger A, Kurath U, Roditi I (1997) Contribution of the procyclin 3'
untranslated region and coding region to the regulation of expression in
bloodstream forms of *Trypanosoma brucei*. Mol Biochem Parasit 89: 109-121.
34. Hotz H-R, Hartmann C, Huober K, Hug M, Clayton CE (1997) Mechanisms of
developmental regulation in *Trypanosoma brucei*: A polypyrimidine tract in the
3'-untranslated region of a trypanosome surface protein mRNA affects RNA
abundance and translation. Nucleic Acids Res 25: 3017-3025.
35. Hotz H-R, Biebinger S, Flaspohler J, Clayton CE (1998) *PARP* gene expression:
regulation at many levels. Mol Biochem Parasitol 91: 131-143.
36. Engstler M, Boshart M (2004) Cold shock and regulation of surface protein
trafficking convey sensitization to inducers of stage differentiation in
Trypanosoma brucei. Genes Dev 18: 2798-2811.
37. Wurst M, Selinger B, Jha B, Klein C, Queiroz R, et al. (2012) Expression of the
RNA Recognition Motif protein RBP10 promotes a bloodstream-form
transcript pattern in *Trypanosoma brucei*. Mol Microbiol 83: 1048-1063.
38. Telleria EL, Benoit JB, Zhao X, Savage AF, Regmi S, et al. (2014) Insights into
the trypanosome-host interactions revealed through transcriptomic analysis of
parasitized tsetse fly salivary glands. PLoS Negl Trop Dis 8: e2649.
39. Erben E, Fadda A, Lueong S, Hoheisel J, Clayton C (2014) Genome-wide
discovery of post-transcriptional regulators in *Trypanosoma brucei*. PLoS
Pathogens 10: e1004178.
40. Lueong S, Merce C, Fischer B, Hoheisel J, Erben E (2016) Gene expression
regulatory networks in *Trypanosoma brucei*: insights into the role of the
mRNA-binding proteome. Mol MIcrobiol 100: 457-471.
41. Zinoviev A, Leger M, Wagner G, Shapira M (2011) A novel 4E-interacting protein
in *Leishmania* is involved in stage-specific translation pathways. Nucleic
Acids Res 39: 8404-8415.

42. Alsford S, Turner D, Obado S, Sanchez-Flores A, Glover L, et al. (2011) High
throughput phenotyping using parallel sequencing of RNA interference targets
in the African trypanosome. Genome Res 21: 915-924.
43. Mayho M, Fenn K, Craddy P, Crosthwaite S, Matthews K (2006) Post-
transcriptional control of nuclear-encoded cytochrome oxidase subunits in
Trypanosoma brucei: evidence for genome-wide conservation of life-cycle
stage-specific regulatory elements. Nucleic Acids Res 34: 5312-5324.
44. Arieti F, Gabus C, Tambalo M, Huet T, Round A, et al. (2014) The crystal
structure of the Split End protein SHARP adds a new layer of complexity to
proteins containing RNA recognition motifs. Nucleic Acids Res 42: 6742-
6752.
45. Morgan CE, Meagher JL, Levengood JD, Delproposto J, Rollins C, et al. (2015)
The First Crystal Structure of the UP1 Domain of hnRNP A1 Bound to RNA
Reveals a New Look for an Old RNA Binding Protein. J Mol Biol 427: 3241-
3257.
46. Wang X, Tanaka Hall T (2001) Structural basis for recognition of AU-rich element
RNA by the HuD protein. Nat Struct Biol 8: 141-145.
47. Deo R, Bonanno J, Sonenberg N, Burley S (1999) Recognition of polyadenylate
RNA by the poly(A)-binding protein. Cell 98: 835-845.
48. Handa N, Nureki O, Kurimoto K, Kim I, Sakamoto H, et al. (1999) Structural basis
for recognition of the tra mRNA precursor by the Sex-lethal protein. Nature
398: 579-585.
49. Drozdz M, Clayton CE (1999) Structure of a regulatory 3'-untranslated region
from *Trypanosoma brucei*. RNA 5: 1632-1644.
50. Misset O, Opperdoes FR (1987) The phosphoglycerate kinases from
Trypanosoma brucei. Eur J Biochem 162: 493-500.
51. Gibson WC, Swinkels BW, Borst P (1988) Post-transcriptional control of the
differential expression of phosphoglycerate kinase genes in *Trypanosoma
brucei*. J Mol Biol 201: 315-325.
52. Blattner J, Clayton CE (1995) The 3'-untranslated regions from the *Trypanosoma
brucei* phosphoglycerate kinase genes mediate developmental regulation.
Gene 162: 153-156.
53. Quijada L, Hartmann C, Guerra-Giraldez C, Drozdz M, Irmer H, et al. (2002)
Expression of the human RNA-binding protein HuR in *Trypanosoma brucei*
induces differentiation-related changes in the abundance of developmentally-
regulated mRNAs. Nucleic Acids Res 30: 1-11.
54. Dean S, Marchetti R, Kirk K, Matthews K (2009) A surface transporter family
conveys the trypanosome differentiation signal. Nature 459: 213-217.
55. Szoor B, Ruberto I, Burchmore R, Matthews KR (2010) A novel phosphatase
cascade regulates differentiation in *Trypanosoma brucei* via a glycosomal
signaling pathway. Genes Dev 24: 1306-1316.
56. Benz C, Mulindwa J, Ouna B, Clayton C (2011) The *Trypanosoma brucei* zinc
finger protein ZC3H18 is involved in differentiation. Mol Biochem Parasitol
177: 148-151.
57. Kershaw CJ, Costello JL, Talavera D, Rowe W, Castelli LM, et al. (2015)
Integrated multi-omics analyses reveal the pleiotropic nature of the control of
gene expression by Puf3p. Sci Rep 5: 15518.
58. Ling A, Trotter J, Hendriks E (2011) A zinc finger protein, TbZC3H20, stabilises
two developmentally regulated mRNAs in trypanosomes. J Biol Chem 286:
20152-20162.
59. Jones NG, Thomas EB, Brown E, Dickens NJ, Hammarton TC, et al. (2014)
Regulators of Trypanosoma brucei cell cycle progression and differentiation
identified using a kinome-wide RNAi screen. PLoS Pathog 10: e1003886.

60. Vassella E, Probst M, Schneider A, Studer E, Renggli C, et al. (2004) Expression
of a major surface protein of *Trypanosoma brucei* insect forms is controlled by
the activity of mitochondrial enzymes. Mol Biol Cell 15: 3986-3993.
61. Kolev NG, Ramey-Butler K, Cross GA, Ullu E, Tschudi C (2012) Developmental
progression to infectivity in *Trypanosoma brucei* triggered by an RNA-binding
protein. Science 338: 1352-1353.
62. Lott K, Mukhopadhyay S, Li J, Wang J, Yao J, et al. (2015) Arginine methylation
of DRBD18 differentially impacts its opposing effects on the trypanosome
transcriptome. Nucleic Acids Res 43: 5501-5523.
63. Minia I, Merce C, Terrao M, Clayton C (2016) Many mRNAs that increase after
heat shock of procyclic-form *Trypanosoma brucei* are also induced during
differentiation to mammalian-infective forms. submitted.
64. Wang H, Li X, Gao S, Sun X, Fang H (2015) Transdifferentiation via transcription
factors or microRNAs: Current status and perspective. Differentiation 90: 69-
76.
65. van der Meulen T, Huising MO (2015) Role of transcription factors in the
transdifferentiation of pancreatic islet cells. J Mol Endocrinol 54: R103-117.
66. Gao T, McKenna B, Li C, Reichert M, Nguyen J, et al. (2014) Pdx1 maintains
beta cell identity and function by repressing an alpha cell program. Cell Metab
19: 259-271.
67. Alibu VP, Storm L, Haile S, Clayton C, Horn D (2004) A doubly inducible system
for RNA interference and rapid RNAi plasmid construction in *Trypanosoma*
brucei. Mol Biochem Parasitol 139: 75-82.
68. Adiconis X, Borges-Rivera D, Satija R, DeLuca DS, Busby MA, et al. (2013)
Comparative analysis of RNA sequencing methods for degraded or low-input
samples. Nat Methods 10: 623-629.
69. Estévez A, Kempf T, Clayton CE (2001) The exosome of *Trypanosoma brucei*.
EMBO J 20: 3831-3839.
70. Singh A, Minia I, Droll D, Fadda A, Clayton C, et al. (2014) Trypanosome MKT1
and the RNA-binding protein ZC3H11: interactions and potential roles in post-
transcriptional regulatory networks. Nucleic Acids Res 42: 4652-4668.
71. Minia I, Clayton C (2016) Regulating a post-transcriptional regulator: protein
phosphorylation, degradation and translational blockage in control of the
trypanosome stress-response RNA-binding protein ZC3H11. PLoS
Pathogens 12: e1005514.
72. Martin M (2011) Cutadapt removes adapter sequences from high-throughput
sequencing reads. EMBnetjournal 17: 10-12.
73. Langmead B, Trapnell C, Pop M, Salzberg S (2009) Ultrafast and memory-
efficient alignment of short DNA sequences to the human genome. Genome
Biol 10: R25.
74. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, et al. (2009) The Sequence
Alignment/Map format and SAMtools. Bioinformatics 25: 2078-2079.
75. Anders S, Huber W (2010) Differential expression analysis for sequence count
data. Genome Biol 11: R106.
76. Bailey T (2011) DREME: Motif discovery in transcription factor ChIP-seq data.
Bioinformatics 27: 1653-1659.
77. Kolev N, Franklin J, Carmi S, Shi H, Michaeli S, et al. (2010) The transcriptome of
the human pathogen *Trypanosoma brucei* at single-nucleotide resolution.
PLoS Pathog 6: e1001090.
78. Zomer AW, Allert S, Chevalier N, Callens M, Opperdoes FR, et al. (1998)
Purification and characterisation of the phosphoglycerate kinase isoenzymes
of *Trypanosoma brucei* expressed in *Escherichia coli*
. Biochim Biophys Acta 1386: 179-188.

79. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, et al. (2012) Fiji:
an open-source platform for biological-image analysis. Nat Methods 9: 676-
682.
80. Fritz M, Vanselow J, Sauer N, Lamer S, Goos C, et al. (2015) Novel insights into
RNP granules by employing the trypanosome's microtubule skeleton as a
molecular sieve. Nucl Acids Res 43: 8013-8032.
81. Mony BM, Macgregor P, Ivens A, Rojas F, Cowton A, et al. (2013) Genome-wide
dissection of the quorum sensing signalling pathway in *Trypanosoma brucei*.
Nature 505: 681-685.

1007

Supporting information

S1 Text

3'-untranslated regions of mRNAs that were at least 3x enriched in both RBP10 pull-

downs.

S2 Text

3'-untranslated regions of mRNAs that were less than 0.7x enriched in both RBP10

pull-downs.

S3 Text

3'-untranslated regions of mRNAs that were at least 3x enriched on average in

RBP10 pull-downs, according to DESeq (Padj <8 E-5), but one replicate was less

than 3x enriched.

S1 Table

Proteins that co-purified with TAP-tagged RBP10.

Endogenously TAP-tagged RBP10 was purified three times in the presence or

absence of RNase. The controls were three independent purifications of GFP-TAP

Sheet 1: Proteins enriched in the purifications. Peptide counts and ratios are shown.

The highest peptide count from the GFP-TAP replicates was used as a baseline. The

proteins shown in this table were present as at least 2 peptides, present in at least

two similar purifications, and were on average at least 2-fold enriched relative to

GFP-TAP. Functions of interest are highlighted in different colours. "Average PGKB

=tet/-tet" are tethering screen results [39]; a value above 1.5 suggests that the

protein can suppress expression. "in starvation granules?" is 1 if positive and 0 if

negative [80]. The number of peptides found by purification of MKT1-TAP are also

indicated [70].

Sheet 2: All raw data.

S2 Table

Results of a yeast 20-hybrid screen with a random fragment library.

The library [70] was screened with pGBKT7-RBP10 and results evaluated by
sequencing.

Sheet 1: Raw data for pGBKT7-RBP10. All reads that started from position -36
onwards (relative to the ATG) are included.

Sheet 2: Raw data for pGBKT7 (unselected background, filtered for at least 4 reads)

Sheet 3: Results for pGBKT7-RBP10, filtered to exclude loci with less than 10 reads,
and open reading frames with less than 2 independent fragments. Numerous
cysteine protease fragments have been removed.

Sheet 4: Possible interaction partners. The ratio to the maximum background count is
at least 4. Proteins that also scored as interacting with MKT1 or CFB1 [70] (indicating
less specificity) are listed separately. The "regulator" column shows results from the
tethering screen [39].

S3 Table

Effect of RBP10 depletion on gene expression in bloodstream forms.

RNAi was induced for 15h and mRNA was collected for RNASeq from total cells, the
free and monosomal fractions, and the polysomal fractions. Data are for two
independent experiments which are labelled A and B.

Throughout the Tables, the colour code for ratios is as follows: Red is >2, orange is
1.5-2.0, green is 0.667 - 0.5 and blue is <0.5. Shading of columns does not follow a
particular colour code but is designed to make the tables easier to navigate. Many
numbers have been rounded to 2 or 3 points beyond the decimal in order to
decrease the sizes of the files. Ribosome profiling results (Riboprof) are from [11],
and developmental regulation data for total RNA are from [10].

Sheet 1: Raw read counts for all unique genes, chosen according to [9] with some
later additions.

Sheet 2: Columns E - P Results were converted to reads per million reads (RPM).
For each experiment, we then corrected the RPM for the monosomes and polysomes
according to the fraction of the total mRNA that was in each fraction (sheet 7 and S3
Fig, numbers not included in the Table). We then calculated the percentage of each

mRNA that was on polysomes (columns Q-T). Column U contains the average of ratio between RNAi and control, calculated for the total input RNA - (K+L) / (E+F). Column V is the same ratio for the fraction of RNA not on polysomes, and column W is the ratio for the polysomal fractions (S+T) / (Q+R). To find the lowest possible estimate of the ratio in W, we took the lowest value for the proportion in polysomes after *rbp10* RNAi, and divided by the highest value from wild-type. From this column (X) we could find mRNAs that reliably move *into* polysomes after RNAi. Column Y is similar but shows the *highest* estimate for the ratio; this can be used to find mRNAs that reliably move *away* from polysomes after RNAi. Since there were only 3 genes in this category (column Y) they were not considered further.

Sheet 3: mRNAs that move onto polysomes after *rbp10* RNAi. Same as sheet 2, except that only genes for which column W is more than 1.5, and column X is more than 1.2, are included. (This means that the average increase is at least 1.5x and the minimum is 1.2x.) Columns Y and Z are ribosome profiling results from [11], giving the ribosome density ratios for bloodstream forms (BS) divided by procyclic forms (PC). Slender bloodstream forms from blood are slBS and cultured bloodstream forms are cBS. Column AB is results from S5 Table, sheet 2 (binding of the RNA to RBP10, ratio of bound to unbound) and column AC is the adjusted P-values for the results in column AB.

Sheet 4: DeSeq analysis of the data in Sheet 1.

Sheet 5: Genes showing at least a 1.5-fold increase either in the amount of total RNA, or the amount of RNA in polysomes, or both, according to DeSeq (adjusted P-value threshold 0.05). Column AA is the RNA-binding ratio from S5 Table, sheet 2,

Sheet 6: As sheet 5, except that these are genes showing at least a 1.5-fold decrease either in the amount of total RNA, or the amount of RNA in polysomes, or both, according to DeSeq (adjusted P-value threshold 0.05).

Sheet 7: Quantitation of spliced leader signals for the free/monosome and polysome fractions used for RNA-Seq

Sheet 8: Statistical analyses to compare the proteomes of differentiating trypanosomes [16] with the effects of RBP10 RNAi on RNA levels.

S4 Table

Effect of RBP10 expression on gene expression in procyclic forms.

Expression of RBP10 was induced with tetracycline addition for 6 hours, and RNA
samples prepared from duplicates as in Supplementary Table S3. Controls are the
same cells without tetracycline.

Sheet 1: Columns D-O are raw reads. Columns P-S are corresponding RPM values
for input samples.

Sheet 2: Proportions in free/monosome and polysome fractions were calculated as
described for Supplementary Table S3, sheet 2. Columns Q-X: Effects on both
polysomes and free/monosome fractions were also calculated in a similar way.

Sheet 3: As sheet 2, but only genes showing regulated translation according to the
following criteria: (a) movement to the free/monosome fraction from the polysomes.
This is the top section (rows 2-320). The smallest value for the proportion in
free/monosomes with RBP10 expression was divided by the largest value calculated
value for the proportion in free/monosomes without RBP10 induction. If this number
was greater than 1.5, the gene was included. In addition, we divided the largest
number for the proportion in polysomes plus RBP10 expression by the smallest value
for the proportion in polysomes without RBP10 induction. If this number was less
than 0.667 the gene was also included. Genes that satisfied both criteria are in bold.
Additional columns are as in Table S3. Rows 322-326 are the four RNAs showing
movement onto polysomes.

Sheet 4: DeSeq analysis of the results in Sheet 1.

Sheet 5: Genes showing at least a 1.5-fold increase either in the amount of total
RNA, or the amount of RNA in polysomes, or both, according to DeSeq (adjusted P-
value threshold 0.05).

Sheet 6: As sheet 5, except that these are genes showing at least a 1.5-fold
decrease either in the amount of total RNA, or the amount of RNA in polysomes, or
both, according to DeSeq (adjusted P-value threshold 0.05). Columns V-Z are results
for salivary gland trypanosomes, calculated from the raw data in [38]; statistical
analysis was not possible due to the absence of replicas.

Sheet 7: Quantitation of spliced leader signals for the free/monosome and polysome
fractions used for RNASeq.

S5 Table

RNAs bound to tagged RBP10.

Results for two independent experiments, A and B, are shown.

Sheet 1: Raw read counts for all samples, calculation of reads per million reads

(RPM), and ratios of bound/unbound for each experiment.

Sheet 2: DeSeq analysis of the raw data in sheet 1

Sheet 3: Set of mRNAs that were at least 3-fold bound according to DeSeq. This is a

slightly more relaxed stringency than that used in sheet 5. The largest P-value was

7E-5. H-J: Results for developmental regulation [10]; K - calculated from [38]; L-T:

results from Supplementary Tables S3 and S4; U, W, Z: [13]; V: [14]; X, Y, AA: [15];

AB: [16]; AC: [81]; AD: [42].

Sheet 4: Statistics used to create the Venn diagrams in Figure 2.

Sheet 5: Set of bound RNAs used for the motif search. Filters were (a) at least 10

reads in all samples and (b) a bound/unbound ratio of >3 in both experiments. The 3'-

UTR lengths are those annotated in TritrypDB, except where noted. Only annotated

3'-UTRs were used for the initial motif search.

S6 Table

Plasmids and oligonucleotides

S1 Fig

Alignment of RBP10 from different kinetoplastids

1146

A. Bloodstream form TAP-RBP10 cell line

B. Bloodstream form RNAi

Anti RBP10

C. Procyclic form inducible expression of RBP10-myc

1147

S2 Fig

Cell line characterisation and RBP10 expression

A) Characterisation of bloodstream-form trypanosomes with a sequence encoding an

N-terminal tandem affinity purification tag integrated in frame with the *RBP10* coding

region. The left hand panel is a western blot showing the presence or absence of

TAP-RBP10 and native RBP10, with genotypes above the lanes. The right-hand

panel is a cumulative growth plot for the cell line used in the interactome

experiments.

B) RNAi cell line: Western blots showing the time course of RBP10 decrease after

tetracycline addition

C) Western blots showing the time course of expression of RBP10-myc in Lister 427

procyclic forms after tetracycline addition.

1160

1161

S3 Fig

RNAi targeting *RBP10* in bloodstream forms: samples used for RNASeq

A. Typical Sucrose gradient profiles of samples

B. Total mRNA in the sucrose gradient fractions, detected using the spliced leader
probe

C. Average quantitation of the spliced leader signal

1168

1169

S4 Fig

Expression of RBP10 in procyclic forms: samples used for RNASeq

A. Typical Sucrose gradient profiles of samples

B. Total mRNA in the sucrose gradient fractions, detected using the spliced leader

probe,

C. Average quantitation of the spliced leader signal

1176

1177

1178

S5 Fig

Effects of RNAi and analysis of RBP10-bound RNAs.

A. Analysis of binding of individual mRNAs to RBP10. The classes are ratios of
bound/unbound; the number of different open reading frames in each class is on the
y axis.

B. Scatter plot comparing the effect of *rbp10* RNAi on total RNA (x axis) with the
effect on polysomal RNA (y axis) for mRNAs that were at least 3x enriched according
to DeSeq; all P-values were less than 8E-5. The black line is the regression line and
the red and green lines show the 95% confidence limits for the data. The blue
shadow encloses total mRNAs that were less than 1.5x affected, and the pink
shadow encloses polysomal RNAs that were less than 1.5x affected. The cyan line is
perfect correlation. The box beneath the graph lists relevant TritrypDB accession
numbers. The gene numbers on the plot are also accession numbers, with "Tb927."
removed.

C. As (B), but here the y axis shows the effect of RNAi on the percentage of the
mRNA in polysomes.

1195

S6 Fig

Relocation of phosphoglycerate kinase during differentiation.

Cells were fixed with fomaldehyde, pernmeabilized with triton x-100, and stained
using a polyclonal antibody to phosphglycerate kinase (PGK). The grey-scale panels
show PGK alone and the differential interference contrast panels show DNA in cyan
and PGK in magenta.

A. Bloodstream forms

B. Bloodstream forms after incubation with cis aconitate for 17h at 27°C.

C. Procyclic forms.

D. Bloodstream forms with 17h *rbp10* RNAi followed by culture for 3 days under
procyclic-form culture conditions. The selection shows on procyclic-like trypanosome
(left) and one which still has bloodstream-form morphology (right).

E. Procyclic forms after 2 days induction of expression of RBP10-myc. Cells with
terminal kinetoplasts are labelled "b" and cells with more procyclic morphology are
labelled "p". Cells that appear to be dividing are indicated with asterisks. A very
abnormal cell is indicated with "?". Cells if unclear status are not labelled.

1212
1213

1214

S7 Fig

Morphological analysis of differentiation.

Cells were treated as indicated, fixed, and stained for DNA (cyan), EP-procyclin,

phopho-GPEET procyclin, or VSG. (magenta). The posterior-kinetoplast and

kinetoplast-nucleus distances were measured in Image J and the ratios calculated.

A. Summary of all results. This is the same as Fig 5H, but all of the outliers are
included, and the numbers of cells analysed are below each lane. BS - normal
bloodstream forms; PC- normal procyclic forms; CA - 17h cis-aconitate pretreatment;
RNAi or Ri; 17h *rbp10* induction; +RBP10 - induced expression or RBP10 for 2 days.

B. Cell distributions after cis-aconitate-induced differentiation. The x axis shows the
P-K/K-N ration, and y-axis shows the percentage of cells with that ratio. Controls are
bloodstream forms (magenta) and established procyclic forms (green). The other
lines are for 3 and 6 days after transfer to procyclic medium (MEM).

C. As B, but with transfer after 17h *rbp10* RNAi.

D. Phospho-GPEET expression 3 days after cis-aconitate-stimulated differentiation.
The cells indicated with the cyan line (BS_CA_3d MEM) were subdivided into
GPEET positive and GPEET negative.

E. Phospho-GPEET expression 3 days after RNAi-stimulated differentiation; other
details as in D.

F. Cell distributions in procyclic forms after induced expression of RBP10.

G. As (F), but also showing staining with anti-phospho-GPEET and anti-VSG117.
Cells wit no GPEET have bloodstream-form kinetoplast positions, and a subset of
them cross-reacts with anti-VSG117 antibodies. The remainder may express VSGs
that do not cross react with the anti-117 antibody.
1239