Cellobiose consumption uncouples extracellular glucose

sensing and glucose metabolism in *Saccharomyces*

cerevisiae

Kulika Chomvong[1], Daniel I. Benjamin[2], Daniel K. Nomura[2] and Jamie H.D. Cate[3,4,5,#]

[1]Department of Plant and Microbial Biology, University of California, Berkeley, CA

94720

[2]Department of Nutritional Sciences and Toxicology, University of California, Berkeley,

CA 94720

[3]Department of Molecular and Cell Biology, University of California, Berkeley, CA 94720

[4]Department of Chemistry, University of California, Berkeley, CA 94720

[5]Physical Biosciences Division, Lawrence Berkeley National Laboratory, Berkeley, CA

94720

[#]Corresponding author (email: jcate@lbl.gov)

KC: kchomv@gmail.com

DIB: dbenjami@stanford.edu

DKN: dnomura@berkeley.edu

JHDC: jcate@lbl.gov

Running title: ATP and optimal glucose consumption

Final character count: 46,908

Keywords: cellobiose/glucose sensors/metabolomics/PMA1

Abstract

Glycolysis is central to energy metabolism in most organisms, and is highly

regulated to enable optimal growth. In the yeast *Saccharomyces cerevisiae*, feedback

mechanisms that control flux through glycolysis span transcriptional control to

metabolite levels in the cell. Using a cellobiose consumption pathway, we decoupled

glucose sensing from carbon utilization, revealing new modular layers of control that

induce ATP consumption to drive rapid carbon fermentation. Alterations of the beta

subunit of phosphofructokinase (*PFK2),* H$^+$-plasma membrane ATPase (*PMA1),* and

glucose sensors *(SNF3, RGT2)* revealed the importance of coupling extracellular

glucose sensing to manage ATP levels in the cell.   Controlling the upper bound of

cellular ATP levels may be a general mechanism used to regulate energy levels in cells,

via a regulatory network that can be uncoupled from ATP concentrations under

perceived starvation conditions.

Importance

Living cells are fine-tuned through evolution to thrive in their native environments.

Genome alterations to create organisms for specific biotechnological applications may

result in unexpected and undesired phenotypes. We used a minimal synthetic biological

system in the yeast *Saccharomyces cerevisiae* as a platform to reveal novel

connections between carbon sensing, starvation conditions and energy homeostasis.

Introduction

Most microorganisms favor glucose as their primary carbon source, as reflected

in their genetic programs hard-wired for this preference. Central to carbon metabolism is

glycolysis, which is finely tuned to the dynamic state of the cell due to the fact that

glycolysis first consumes ATP before generating additional ATP equivalents. To avoid

catastrophic depletion of ATP, the yeast *Saccharomyces cerevisiae* has evolved a

transient ATP hydrolysis futile cycle coupled to gluconeogenesis (1). Glycolysis in yeast

is also tightly coupled to glucose transport into the cell, entailing three extracellular

glucose sensing mechanisms and at least one intracellular glucose signaling pathway

(2).

Synthetic biology and metabolic engineering of yeast holds promise to convert

this microorganism into a "cell factory" to produce a wide range of chemicals derived

from renewable resources or those unattainable through traditional chemical routes.

However, many applications require tapping into metabolites involved in central carbon

metabolism, a daunting challenge as living cells have numerous layers of feedback

regulation that fine-tune growth to changing environments. Cellular regulation evolved

intricate networks to maintain and ensure cell survival. For example, *S. cerevisiae* has

evolved to rapidly consume high concentrations of glucose through fermentation, while

repressing the expression of other carbon consumption pathways, an effect termed

glucose repression. When perturbed genetically, regulatory networks such as those in

glucose repression often generate undesirable or unexpected phenotypes.

For yeast to be useful in producing large volumes of renewable chemicals or

biofuels, it will be important to expand its carbon utilization to include multiple sugars in

the plant cell wall. One promising approach that helps overcome glucose repression

and allows simultaneous utilization of different sugars is cellobiose consumption (3).

Cellobiose is a disaccharide with two units of glucose linked by a ß-1,4 glycosidic bond.

Cellobiose consumption using a minimal additional pathway in yeast—a cellodextrin

transporter (CDT-1) and intracellular ß-glucosidase (4)–avoids glucose repression by

importing carbon in the form of cellobiose instead of glucose. The cellodextrin

transporter allows cellobiose to enter the cell where it is hydrolyzed to glucose and

consumed via the native glycolytic consumption pathway. By moving glucose production

into the cell, the *Neurospora crassa*-derived cellobiose consumption pathway is nearly

the minimal synthetic biological module imaginable in *S. cerevisiae*, comprised of just

two genes. Nevertheless, in *S. cerevisiae* the cellobiose consumption pathway is inferior

to consumption of extracellular glucose in terms of rate and results in a prolonged lag

phase (5). Previous efforts to understand the impact of cellobiose consumption on the

physiology of *S. cerevisiae* using transcriptional profiling revealed that cellobiose

improperly regulates mitochondrial activity and amino acid biosynthesis, both of which

are tightly coupled to the transition from respiration to fermentation (5).

Since glycolytic enzymes are regulated mostly at the post-transcriptional level (6),

we probed cellobiose consumption in *S. cerevisiae* at the metabolite level. We found

that key metabolites in glycolysis are highly imbalanced, leading to low flux through

glycolysis and slow fermentation. We also found that excess ATP levels drive the

imbalance, and identified a new potential regulatory role of glucose sensors in cellular

ATP homeostasis.

Results

Metabolite profiling of cellobiose utilizing *S. cerevisiae*

S. cerevisiae cells engineered with the cellobiose consumption pathway exhibits

a prolonged lag phase, with decreased growth and carbon consumption rates in

comparison to when glucose is provided (Figure S1A) (5). We hypothesized that

cellobiose consumption results in an ATP deficit in glycolysis relative to glucose, due to

the fact that the cellodextrin transporter (CDT-1) in the cellobiose utilizing pathway is a

proton symporter, requiring ATP hydrolysis for cellobiose uptake (7). Moreover, under

anaerobic conditions, ATP production is limited to substrate-level phosphorylation,

further restricting ATP availability. We measured the steady-state concentrations of ATP

and other metabolites in central carbon metabolism in yeast fermenting cellobiose

compared to glucose. Of the 48 compounds analyzed, the abundance of 25 compounds

was significantly different between cellobiose and glucose-fed samples (Figure S1B).

Surprisingly, ATP levels increased by 6-fold in the cellobiose grown cells (Figure 1A).

The relative abundance of compounds in glycolysis—fructose 6-phosphate (F6P),

glucose 6-phosphate (G6P), glucose and pyruvate—increased by 444-, 81-, 7- and 3-

fold, respectively, while that of phosphoenolpyruvate (PEP) decreased by 2-fold (Figure

1A, B). These results suggest that the yeast cells underwent drastic physiological

changes, reflected in metabolite levels, when cellobiose was provided in place of

glucose.

Phosphofructokinase-1 inhibition by excess ATP

Given the dramatic buildup of glucose, G6P and F6P intermediates prior to the

phosphofructokinase (*PFK1*) reaction in glycolysis (Figure 1B, 2A), we reasoned that

Pfk1 might be a major bottleneck in cellobiose consumption. Pfk1 catalyzes the

phosphorylation of F6P, using one ATP and yielding fructose 1,6-bisphosphate

(F1,6BP) as a product. As the second committed step in glycolysis, Pfk1 is heavily

regulated—with ATP acting as an allosteric inhibitor and AMP and fructose 2,6-

bisphosphate serving as activators (8-10).

To test whether allosteric inhibition of Pfk1 by ATP limited cellobiose

fermentation, a mutation in Pfk1 that makes the enzyme ATP-insensitive (P722L in the

Pfk1 beta subunit, note the original publication referred to this mutation as P728L (11))

was introduced into the chromosomally encoded *PFK1* gene (mutation here termed

pfk1m) in the cellobiose utilizing strain. This mutation was previously shown to reduce

ATP inhibition but also AMP and fructose 2,6-bisphosphate activation of Pfk1 in *S.*

cerevisiae (11). We chose this mutation over an ATP-insensitive, AMP/F2,6BP-sensitive

mutant phosphofructokinase (10) because the latter's phenotype has not been

evaluated in *S. cerevisiae*. High initial cell densities were used hereafter, as the focus of

this study is sugar consumption rather than growth.

Consistent with allosteric inhibition of Pfk1 by ATP, the cellobiose consumption

efficiency (E_c) of the *pfk1m* strain increased by 33% in comparison to the strain with

wild-type Pfk1 (Figure 2B). In these high cell densities, negligible changes in growth rate

were observed (Figure 2C). The relative abundance of G6P and F1,6BP decreased by

47% and 34%, respectively, while that of ATP remained relatively unchanged (Figure

2D). The unchanged ATP level was expected as the ATP requirement for the cellobiose

consumption pathway was likely offset by the ATP generated as part of carbon

metabolism. These results indicate that the 6-fold increase in cellular ATP

concentrations allosterically inhibited Pfk1*,* resulting in accumulation of glucose, G6P

and F6P, which eventually slowed down cellobiose consumption.

Limited activity of plasma membrane ATPase

Although the *pfk1m* strain partially increased the rate of cellobiose fermentation,

cellular ATP remained elevated relative to glucose fermentation. It is unlikely that ATP

production was the cause of the difference, as fermentation is limited to substrate-level

phosphorylation under anaerobic conditions regardless of carbon source. We therefore

tested whether the activity of one of the major ATP sinks in yeast, the plasma

membrane ATPase (Pma1) was responsible for the ATP buildup. Pma1 hydrolyzes 25-

40% of cellular ATP in yeast (12) and is heavily regulated by glucose (13).

A constitutively active mutant form of *PMA1* (*pma1-Δ916*, here abbreviated *pt*)

(14) was introduced into the endogenous *PMA1* locus in the cellobiose utilizing strain.

This mutation results in high Pma1 ATPase activity even in carbon starvation conditions

(14). The E_c of the *pt* strain was 4 times that of the control (Figure 2B), whereas the

growth rate of *pt* strain was only slightly faster than the Pma1 WT strain (Figure 2C). As

expected, we observed a 66% decrease in cellular ATP levels in the *pt* strain in

comparison to the wild-type control (WT, i.e. cellobiose pathway only) (Figure 2D). In

addition, the concentrations of G6P and F1,6BP decreased by 58% and 93%,

respectively, relative to strains expressing wild-type *PMA1*. Notably, these

concentrations dropped more than when the ATP-insensitive *PFK1* mutant was

introduced (Figure 2D). These results suggest that increased Pma1 ATPase activity

improved cellobiose fermentation. We hypothesize that the drastic decrease in F1,6BP

level and the fast growth rate were the result of rapid glycolytic flux, as the cells

experience low cellular ATP levels in the *pt* strain.

Next, we observed the phenotypes of *pfk1m-pt* double mutant strain. The

cellobiose consumption profile of a *pfk1m-pt* double mutant was identical to that of the

pt strain (Figure 2A). However, the growth rate and relative abundance of G6P, F1,6BP

and ATP of the *pfk1m-pt* differed from the *pt* strain (Figure 2C,D). In fact, their levels

were similar to those in the *pfk1m* strain. These results imply that while the ATP might

be hydrolyzed rapidly due to the *pt* effect, the removal of ATP inhibition on *pfk1* allowed

enough ATP to be regenerated downstream that no growth burst was observed. The

underlying explanation of the mixed phenotypes will require future experiments to

dissect how Pfk1 exerts allosteric control on glycolysis and ATP levels.

Carbon starvation-like state of the plasma membrane ATPase

Although cellobiose theoretically provides the same energy and carbon

availability to cells as glucose, its releases glucose only after intracellular hydrolysis by

β-glucosidase. Thus the cellobiose consumption system used here does not generate

extracellular glucose, which acts as a crucial signaling molecule for yeast carbon

metabolism. Taken together with the observation that increased ATPase activity in the

pt strain increased cellobiose consumption efficiency, we wondered whether the limited

Pma1 activity in cellobiose-fed cells is due to the absence of extracellular glucose in the

media. Transcriptionally, the presence of glucose increases *PMA1* mRNA levels by 2-4

times via the regulation of Rap1, Gcr1 and Sir2 (15-17). Consistent with the requirement

for extracellular glucose sensing, previous RNA sequencing experiments revealed a

40% decrease in *PMA1* transcript levels when cellobiose was provided in place of

glucose (5). However, although transcriptional regulation of *PMA1* is important, it is

slower than post-transcriptional regulation and results in smaller changes (13, 18).

In the presence of glucose, phosphorylation of Ser-899 decreases Pma1's K_m

and Ser-911/Thr-912 increases Pma1's V_{max} for ATP, respectively (13, 19, 20). Given

the 6-fold excess amount of ATP observed in cellobiose utilizing conditions (Figure 1A),

the effective velocity of the Pma1 reaction is likely approaching V_{max} regardless of the

phosphorylation status at Ser-899 (21). The K_m of ATP hydrolysis by Pma1 has been

reported to increase approximately 3-fold from 1.2 mM in glucose-fermenting cells to 4.0

mM in glucose-starved cells, while a 10-fold decrease in V_{max} was reported in the same

study (22). We reasoned that the 6-fold increase of ATP in cellobiose-fed cells should

result in ATP concentrations in excess of the K_m for Pma1, resulting in Pma1 activity

being limited by its V_{max}. Thus, in this study, we did not investigate the phosphorylation

of Ser-899 and chose to investigate whether V_{max}-determining phosphorylation states of

Ser-911 and Thr-912 might play a major role in establishing the efficiency of cellobiose

fermentation (Figure 3A).

Combinatorial mutations of Ser-911/Thr-912 to alanine and aspartic acid were

introduced into the endogenous *PMA1* gene to prevent or mimic phosphorylation,

respectively. We were unable to obtain strains with *pma1*-S911A/T912A and *pma1*-

S911A/T912D, potentially because the combinations were lethal. All mutant strains

whose Pma1 S911 position was mutated to aspartic acid consumed cellobiose more

efficiently in comparison to when the S911 position remained unchanged (Figure 3B).

By contrast, mutating Pma1 T912 to aspartic acid did not show a correlation with the

cellobiose consumption phenotype. These results suggest that phosphorylation of Pma1

at S911 was lacking when cellobiose was provided as a sole carbon source.

Positive effects of extracellular glucose sensor deletions

According to the above mutational analysis, the Pma1 phosphorylation state of

cellobiose-fed cells was similar to that in carbon starvation conditions (Figure 3A) (20).

In the previously published experiments (20), carbon-starved cells were prepared by

incubating mid-exponential phase cells in media without glucose. In such conditions,

neither extracellular nor intracellular glucose is present. For the cellobiose-fed cells,

based on the relatively high level of intracellular glucose we detected (Figure 1A), it is

unlikely that the intracellular glucose induced Pma1 carbon starvation. Additionally,

since intracellular glucose metabolism is expected in cellobiose-fed cells, its effect on

Pma1 carbon starvation was also ruled out (23). We therefore tested the role of

extracellular glucose in regulating Pma1 activity. In cellobiose-fed cells, glucose is not

provided as part of the media, and thus the extracellular glucose is absent. We

hypothesize that the extracellular glucose is likely essential for full activation of Pma1

through S911 phosphorylation.

Snf3*,* Rgt2 and Gpr1 are the three known sensors of extracellular glucose in *S.*

cerevisiae. Snf3 and Rgt2 mainly regulate glucose transport while Gpr1 controls cell

physiology via an interaction with Gpa2 to activate protein kinase A and cAMP synthesis

(23). To mimic the presence of extracellular glucose, constitutively active mutations

(*snf3* R229K*, rgt2* R231K and *gpa2* R273A) were introduced into the endogenous loci to

probe the role of each glucose-sensing pathway (24, 25). Surprisingly, the cellobiose

consumption efficiency of all the three mutant strains decreased by ~25% (Figure S2A).

We then inverted the genetic modifications by deleting *SNF3, RGT2* and/or *GPA2*.

Notably, the triple glucose-sensing deletion strain (*snf3Δrgt2Δgpa2Δ, or srgΔ*) showed a

275% increase in E_c (Figure 4A).

Combinatorial deletions revealed that the Gpr1 pathway did not contribute to

improved cellobiose fermentation, but combining the *SNF3* and *RGT2* deletions (*srΔ*)

was necessary and sufficient to replicate the E_c of the triple deletion strain (Figure 4A,

Figure S2B, C). Consistent with the observed E_c values, the intracellular ATP levels of

srgΔ and *srΔ* decreased by 41% and 18% respectively, while those in the individual-

deletion strains *snf3Δ* and *rgt2Δ* remained unchanged (Figure 4A). These results reveal

a negative correlation between E_c and cellular ATP levels (Figure 4B) and that Snf3 and

Rgt2 acted together to regulate cellular ATP levels, in addition to regulating glucose

transport.

Although the additional deletion of *GPA2* (*gpa2Δ*) in the *srΔ* strain did not further

improve E_c (Fig 4A), it reduced the relative abundance of ATP by 28%, implying that the

Gpr1 pathway has a separate mechanism to control cellular ATP levels that does not

directly affect carbon metabolism. Consistently, *gpa2Δ* had a negative or neutral impact

on cellobiose consumption (Figure S2). The decrease in ATP level may be a result of

altered cellular activities, controlled by Gpr1 via the Tor and cAMP-PKA-Ras pathways

(23). The relationship between Gpr1-regulated ATP levels and carbon metabolism

remains to be discovered. Since *gpa2Δ* did not have a direct effect on E_c, it was not

investigated further in this study.

Snf3/Rgt2 regulation of cellular ATP levels

To examine whether Snf3/Rgt2 regulated the cellular ATP level in cellobiose

fermentations via Pma1, an *snf3Δ rgt2Δ pma1-Δ916* strain (*srΔ-pt*) was constructed.

Notably, the E_c of the *srΔ-pt* strain increased more than 4 times in comparison to the

wild-type control (Figure 4A, Figure 5A). The improvement was additive, within the

range of the ΔE_c summation of *srΔ* and *pma1-Δ916* strains relative to wild-type (Figure

5A). Although ATP levels decreased in a nearly linear fashion as a function of E_c (Figure

4B), it is not presently possible to ascertain whether the *srΔ* and *pma1-Δ916* mutations

act entirely independently due to limitations in measurement accuracy (Figure 5A).

To further determine the relationship between Snf3/Rgt2 and Pma1, the

vanadate-specific ATPase activity of Pma1 (26) from different strains consuming

cellobiose was analyzed (Figure 5B). Consistent with the constitutively active nature of

the *pt* mutation, the activities of Pma1 in the *pt* and *srΔ-pt* strains were higher than

those observed in the WT or *srΔ* strains, respectively. Addition of *srΔ* decreased the

Pma1 ATPase activities by 25% and 32% in WT and *pt* strains, respectively. In other

words, the absence of Snf3 and Rgt2 led to a partial decrease in Pma1 ATPase activity,

which implied that Snf3/Rgt2 partially activated Pma1 ATPase activity in the absence of

glucose.

Discussion

277   To identify the effects of a minimal alteration to carbon metabolism in yeast, we

278   chose a cellobiose-consumption pathway composed of two genes and analyzed its

279   cellular metabolite profiles in comparison to cells provided with glucose, yeast's

280   preferred carbon source (Figure 6). Here we focus on the cellobiose consumption

281   efficiency (E_c), as E_c linearly correlated with ethanol production rate, while ethanol yield

282   remained mostly unchanged (Figure S3).

283   More than half of the metabolites significantly changed in abundance when

cellobiose was provided in place of glucose. The buildup of G6P, F6P and ATP in *S.*

cerevisiae fermenting cellobiose suggested that Pfk1 was one of the bottlenecks in the

process. Pfk1 is subjected to complex allosteric regulation, including inhibition by ATP

and activation by AMP and fructose 2,6-bisphosphate (F2,6BP) (8-10). The Pfk1

bottleneck was partially relieved in cells expressing an ATP/AMP/F2,6BP-insensitive

PFK1 allele, while the ATP level remained elevated. These results contrast with

previous studies that identified ATP depletion and the buildup of fructose-1,6-

bisphosphate–the metabolite immediately downstream of Pfk1–as a weak link in

glycolysis (1). Although we did not investigate the effect of AMP and F2,6BP activation

since their changes between glucose and cellobiose conditions were less than 2 fold

and they did not meet the significance threshold of a p-value less than 0.01, it is

possible that they could influence cellobiose consumption efficiency.

ATP is central to a cell's energy currency, but too much ATP is not necessary

beneficial (27, 28). In fact, we observed a negative correlation between cellular ATP

levels and cellobiose consumption efficiency (Figure 4B). A similar correlation has been

reported for glucose as a carbon source, suggesting metabolic uncoupling of energy

homeostasis in yeast cells (29). We propose that intracellular glucose concentrations–

generated by cellobiose hydrolysis in our experiments–and glucose metabolism (23) are

insufficient to trigger glucose activation of key metabolic pathways and enzyme activity.

For example, we found that the ATP-dependent proton pump Pma1 existed in a carbon-

starvation like state during cellobiose fermentation, and was partially responsible for the

aberrant accumulation of ATP. These results suggest that neither intracellular glucose

nor glucose metabolism are sufficient to fully activate Pma1. A previous study showed

the lack of phosphorylation of S899 and S911/T912 in Pma1, in a

hexokinase/glucokinase deletion strain (*hxk1Δ hxk2Δ glk1Δ*) provided with glucose,

suggesting that phosphorylation of these residues requires glucose metabolism (30).

Together with our results, we propose that the activation of Pma1 through S911

phosphorylation requires both extracellular glucose and glucose metabolism. Our

results reveal that the cellobiose utilization system allows uncoupling of glucose

metabolism and intracellular glucose from extracellular glucose signaling. Future

experiments will be required to reveal why ATP was not consumed by other cellular

processes triggered under starvation (31).

Cytosolic pH is also a key regulator of carbon utilization (32), and is likely to be

impacted by the use of the proton symporter CDT-1 for cellobiose import and the

resulting low activity of Pma1. High cytosolic pH is necessary and sufficient to activate

Tor-Ras-PKA activities, which are downstream of Gpr1 glucose sensing pathway (32).

By contrast, the proton symporter CDT-1 and low activity of Pma1 may result in a low

cytosolic pH. However, cytosolic pH alone is unlikely to determine the cellobiose

consumption efficiency (E_c) as the strain with an ATP/AMP/F2,6BP-insensitive *PFK1*

allele showed improved E_c but unaltered cellular ATP levels. Furthermore, the

inactivation of the Gpr1 pathway resulted in decreased cellular ATP but unaltered E_c.

Glucose storage, i.e. in the form of trehalose, may be interconnected through the Gpr1

pathway because Ras-cAMP activates trehalase required to break down trehalose—a

phenomenon observed when gluconeogenesis is switched to glycolysis (33). Trehalose

cycling has been shown to lead to an imbalanced state of glycolysis (1). The

relationship between carbon storage and E_c will require future studies to examine this

relationship.

We also found that the well-studied extracellular glucose sensors Snf3/Rgt2

exhibited a novel role in cellular ATP homeostasis partially through the major plasma

membrane ATPase Pma1. Deletion of extracellular glucose sensors (Snf3/Rgt2)

increased cellobiose consumption efficiency and partially restored ATP levels.

Interestingly, the absence of the Snf3/Rgt2 decreased Pma1 ATPase activities, an

effect that should have led to an increase in ATP level. The restored low ATP level

observed in the *sr*Δ strain implied that Snf3/Rgt2 regulated cellular ATP level with

additional mechanism(s) other than through Pma1. It is known that deletion of *SNF3*

and *RGT2* slows down glucose consumption (34), due to the inability of these strains to

degrade Mth1/Std1, which block the promoter regions of hexose transporters required

for optimal glucose import (35-38). Unlike glucose, cellobiose does not signal Mth1

degradation even with intact Snf3/Rgt2 (38). Thus, genes downstream of Mth1

regulation, including hexose transporters, are not expected to be responsible for the

improved E_c and decreased cellular ATP levels observed in *sr*Δ. Consistent with this

model, no growth defect is observed in a *mth1*Δ strain growing on glucose, suggesting

that Snf3/Rgt2 has additional regulatory nodes other than Mth1 (34). Future

transcriptional profiling and ribosome profiling experiments will be required to reveal the

additional Snf3/Rgt2 roles in cellular ATP homeostasis.

The present systems-level study of a minimal synthetic biology pathway for

cellobiose consumption revealed the dramatic impact of decoupling extracellular and

intracellular glucose sensing, resulting in an overabundance of ATP in cells. The

inability of *S. cerevisiae* to catabolize ATP for cellular processes in the presence of

intracellular glucose and glucose metabolism but in the absence of extracellular glucose

resulted in slow fermentation. Thus, ATP levels must be kept in a relatively narrow

range for optimal fermentation and to allow robust startup of glycolysis, yet yeast seems

to lack a direct mechanism to monitor ATP concentrations. For example, a dynamic

model showed that a small concentration difference of inorganic phosphate, a product

of ATP hydrolysis, could alter cell fate from a stable glycolytic steady state to an

imbalanced dead-end state (1). Here, we found that the extracellular glucose sensing by

Snf3/Rgt2 required for optimal glucose fermentation (2) can be uncoupled from the role

of these receptors in regulating ATP homeostasis under carbon starvation conditions. It

will be important in the future to map the fully regulatory pathways of ATP homeostasis

leading from Snf3/Rgt2 and, independently, terminating in Pma1.

Materials & Methods

Yeast strains, media and anaerobic fermentation

The *S. cerevisiae* background strain used in this study was S288C *ura3::P$_{PGK1}$-*

cdt-1 N209S/F262Y-*T$_{ADH1}$ lyp1::P$_{TDH3}$-gh1-1 (codon optimized)-T$_{CYC1}$* derived by

chromosomal DNA library selection (39). The strain was subjected to further

modifications using the CRISPRm system (39). The list of strains constructed,

CRISPRm guides and primers used are included in Table S1.

Seed cultures for cellobiose fermentation experiments were grown in 20 g/L

glucose modified optimal minimal media (oMM) (5) and harvested at mid-log phase. All

cellobiose fermentation experiments were conducted under strict anaerobic conditions,

in 80 g/L cellobiose oMM media at an initial OD$_{600}$ of 20, using 10 mL serum flasks

containing 5 mL fermentation in 2-5 biological replicates. The flasks were incubated at

30°C, 220 rpm. The cellobiose consumption efficiency (E$_c$) was defined as the inverse

of the area under the curve of extracellular cellobiose concentration over time.

Analytical analysis of yeast metabolites

382   Extracellular cellobiose concentrations were determined by high performance

383   liquid chromatography on a Prominence HPLC (Shimadzu, Kyoto, Japan) equipped with

384 Rezex RFQ-FastAcid H 10 x 7.8 mm column. The column was eluted with 0.01 N of

385 H_2SO_4 at a flow rate of 1 mL/min, 55°C.

386  For the metabolite profiling comparison between glucose and cellobiose

387 (modified from (40)) , equal amounts of yeast cells at mid-exponential phase of

388 anaerobic sugar consumption (10 g/L cellobiose or glucose) were harvested (final pellet

389 OD_{600} equivalent to 5). The samples were quenched in 180 µL of 40:40:20

390 acetonitrile:methanol:water. Following the addition of 10 nmols of d3 serine (as an

391 internal standard), the mixtures were vortexed and centrifuged at 13,000 rpm for 10

392 minutes. The supernatants were injected onto an Agilent 6460 QQQ LC-MS/MS and the

393 chromatography was achieved by normal phase separation with a Luna NH_2 column

394 (Phenomenex) starting with 100% acetonitrile with a gradient to 100% 95:5 water

395 acetonitrile. 0.1% formic acid or 0.2% ammonium hydroxide with 50 mM ammonium

396 acetate was added to assist with ionization in positive and negative ionization mode,

397 respectively. Five biological replicates were used for each sample analyzed.

398  For targeted intracellular metabolite comparisons, yeast cells equivalent to 20

399 OD_{600} units were harvested and filtered through a 0.8 µm nylon membrane, prewashed

400 with 3 mL water, followed by another 3 mL water wash after cell filtration. The

401 membranes were placed in 1.5 mL extraction solution (0.1 M formic acid, 15.3 M

402 acetonitrile) flash-frozen in liquid nitrogen, and stored at -80°C. Before analysis, the

403 extracts were vortexed for 15 minutes and centrifuged to collect the supernatants at 4°C.

404 Glucose 6-phosphate and fructose 1,6-bisphosphate were separated and identified

405 using a 1200 Series liquid chromatography instrument (Agilent Technologies, Santa

406 Clara, CA). 1 µL of each sample was injected onto an Agilent Eclipse XDB-C18 (2.1 mm

i.d., 150 mm length, 3.5 µm particle size) column with a Zorbax SB-C8 (2.1 mm i.d.,

12.5 mm length, 5 µm particle size) guard column and eluted at 25 °C and a flow rate of

0.2 mL/min with the following gradient (modification from (41)): 15 min isocratic 100%

buffer A (10 mM tributylamine/15 mM acetic acid), then in 15 min with a linear gradient

to 60% buffer B (methanol), 2 min isocratic 60% B, then 10 min equilibration with 100%

buffer A. The eluent from the column was introduced into a mass spectrometer for 25

minutes after the first 10 minutes. Mass spectrometry (MS) was performed on an LTQ

XL ion trap instrument (Thermo Fisher Scientific, San Jose, CA) with an ESI source

operated in negative ion mode. The MS settings were capillary temperature 350 °C, ion

spray voltage 4.5 kV, sheath gas flow: 60 (arbitrary units), auxiliary gas flow 10

(arbitrary units), sweep gas flow 5 (arbitrary units). For the MS/MS product ion scan, the

scan range was m/z 80 to m/z 300. The compounds G6P at m/z 259.1 and F16BP at

m/z 339.1 were isolated with an m/z 2 isolation width and fragmented with a normalized

collision-induced dissociation energy setting of 35% and with an activation time of 30

ms and an activation Q of 0.250.

The significance threshold between cells provided with cellobiose and cells

provided with glucose was set at a p-value of 0.01. Five biological replicates were used

in each sample group. Only the metabolites with higher than a 2 fold-change between

each sample group were included in the analysis.

Plasma membrane isolation

Strains were subjected to cellobiose fermentation under anaerobic conditions.

Yeast cells with an OD_{600} equivalent to 40 were harvested at mid-log phase and flash

frozen in liquid nitrogen. Membrane fractions were extracted based on the protocol

published in (42).

Pma1 ATPase activity assay

The ATPase assay described in (26) was modified as follows. 30 µg of the

isolated membrane fraction was incubated in assay buffer (50 mM MES pH 5.7, 10 mM

$MgSO_4$, 50 mM KCl, 5 mM NaN_3, 50 mM KNO_3) with and without 3 mM orthovanadate

for 25 minutes at 30 °C. 1.8 mM ATP was added to start the 100 uL reactions. The

reactions were incubated at 30 °C for 15 minutes, then the membranes were isolated

from the reactions by centrifugation at 13,000x g for 10 minutes at 4°C. The released

inorganic phosphate was measured in the supernatant using the ATPase/GTPase

Activity Assay Kit (Sigma-Aldrich) based on the manufacturer's protocol. The specific

Pma1 ATPase activities were calculated by subtracting the concentration of released

inorganic phosphate in reactions provided with orthovanadate from those without.

Yeast cell-based cellobiose uptake assay

The cell-based cellobiose uptake assay was modified from (43). Yeast strains

were grown to mid-exponential phase in 2% oMM glucose, washed with assay buffer (5

mM MES, 100 mM NaCl, pH 6.0) three times and resuspended to a final OD_{600} of 10.

Equal volumes of the cell suspension and 200 µM cellobiose were mixed to start the

reactions, which were incubated at 30°C with continuous shaking for 15 minutes. The

reactions were stopped by adding 150 µL of supernatants to 150 µL 0.1 M NaOH. The

concentrations of the remaining cellobiose were measured using an ICS-3000 Ion

Chromatography System (Dionex, Sunnyvale, CA, USA) equipped with a CarboPac®

PA200 carbohydrate column. The column was eluted with a NaOAc gradient in 100 mM

NaOH at a flow rate of 0.4 mL/min, 30°C.

Acknowledgements

The authors thank Raissa Estrela, Dr. Xin Li, Dr. Ligia Acosta-Sampson, Dr.

Yuping Lin and Dr. Matt Shurtleff for helpful discussions. This work was supported by

funding from Energy Biosciences Institute to JHDC and from National Institutes of

Health (R01CA172667) to DKN.

Conflict of Interest

The authors declare that they have no conflict of interest.

Author contributions

All authors contributed to the design of the experiments. KC and DIB carried out the

experiments. DKN and JHDC aided in interpretation of the systems-level experiments.

KC wrote the manuscript, with editing from DIB, DKN, and JDHC.

References:

1.   **van Heerden JH**, **Wortel MT**, **Bruggeman FJ**, **Heijnen JJ**, **Bollen YJM**,
Planque R, **Hulshof J**, **O'Toole TG**, **Wahl SA**, **Teusink B**. 2014. Lost in
Transition: Start-Up of Glycolysis Yields Subpopulations of Nongrowing Cells.
Science **343**:1245114–1245114.

2.   **Youk H**, **van Oudenaarden A**. 2009. Growth landscape formed by perception
and import of glucose in yeast. Nature **462**:875–879.

3.   **Ha S-J**, **Galazka JM**, **Kim SR**, **Choi J-H**, **Yang X**, **Seo J-H**, **Glass NL**, **Cate JHD**,
Jin Y-S. 2011. Engineered Saccharomyces cerevisiae capable of simultaneous
cellobiose and xylose fermentation. Proc Natl Acad Sci USA **108**:504–509.

4.   **Galazka JM**, **Tian C**, **Beeson WT**, **Martinez B**, **Glass NL**, **Cate JHD**. 2010.
Cellodextrin transport in yeast for improved biofuel production. Science **330**:84–
86.

5.   **Lin Y**, **Chomvong K**, **Acosta-Sampson L**, **Estrela R**, **Galazka JM**, **Kim SR**, **Jin**
Y-S, **Cate JH**. 2014. Leveraging transcription factors to speed cellobiose
fermentation by Saccharomyces cerevisiae. Biotechnology for Biofuels **7**:126.

6.   **Daran-Lapujade P**, **Rossell S**, **van Gulik WM**, **Luttik MAH**, **de Groot MJL**,
Slijper M, **Heck AJR**, **Daran J-M**, **de Winde JH**, **Westerhoff HV**, **Pronk JT**,
Bakker BM. 2007. The fluxes through glycolytic enzymes in Saccharomyces
cerevisiae are predominantly regulated at posttranscriptional levels. Proceedings
of the National Academy of Sciences **104**:15753–15758.

7.   **Kim H**, **Lee W-H**, **Galazka JM**, **Cate JHD**, **Jin Y-S**. 2014. Analysis of cellodextrin
transporters from Neurospora crassa in Saccharomyces cerevisiae for cellobiose
fermentation. Appl Microbiol Biotechnol **98**:1087–1094.

8.   **Bañuelos M**, **Gancedo C**, **Gancedo JM**. 1977. Activation by phosphate of yeast
phosphofructokinase. J Biol Chem **252**:6394–6398.

9.   **Avigad G**. 1981. Stimulation of yeast phosphofructokinase activity by fructose
2,6-bisphosphate. Biochem Biophys Res Commun **102**:985–991.

10. **Nissler K**, **Otto A**, **Schellenberger W**, **Hofmann E**. 1983. Similarity of activation
of yeast phosphofructokinase by AMP and fructose-2,6-bisphosphate. Biochem
Biophys Res Commun **111**:294–300.

11. **Rodicio R**, **Strauss A**, **Heinisch JJ**. 2000. Single point mutations in either gene
encoding the subunits of the heterooctameric yeast phosphofructokinase abolish
allosteric inhibition by ATP. J Biol Chem **275**:40952–40960.

12. **Gradmann D**, **Hansen UP**, **Long WS**, **Slayman CL**, **Warncke J**. 1978. Current-
voltage relationships for the plasma membrane and its principal electrogenic
pump in Neurospora crassa: I. Steady-state conditions. J Membr Biol **39**:333–367.

13. **Serrano R**. 1983. In vivo glucose activation of the yeast plasma membrane
ATPase. FEBS Lett **156**:11–14.

14. **Mason AB**, **Allen KE**, **Slayman CW**. 2014. C-terminal truncations of the
Saccharomyces cerevisiae PMA1 H+-ATPase have major impacts on protein
conformation, trafficking, quality control, and function. Eukaryotic Cell **13**:43–52.

15. **Rao R**, **Drummond-Barbosa D**, **Slayman CW**. 1993. Transcriptional regulation

by glucose of the yeast PMA1 gene encoding the plasma membrane H(+)-
ATPase. Yeast **9**:1075–1084.

16.   **García-Arranz M**, **Maldonado AM**, **Mazón MJ**, **Portillo F**. 1994. Transcriptional
control of yeast plasma membrane H(+)-ATPase by glucose. Cloning and
characterization of a new gene involved in this regulation. J Biol Chem
269:18076–18082.

17.   **Kang WK**, **Kim YH**, **Kang HA**, **Kwon K-S**, **Kim J-Y**. 2015. Sir2 phosphorylation
through cAMP-PKA and CK2 signaling inhibits the lifespan extension activity of
Sir2 in yeast. Elife **4**.

18.   **Eraso P**, **Mazón MJ**, **Portillo F**. 2006. Yeast protein kinase Ptk2 localizes at the
plasma membrane and phosphorylates in vitro the C-terminal peptide of the H+-
ATPase. Biochimica et Biophysica Acta (BBA) - Biomembranes **1758**:164–170.

19.   **Portillo F**, **Eraso P**, **Serrano R**. 1991. Analysis of the regulatory domain of yeast
plasma membrane H+-ATPase by directed mutagenesis and intragenic
suppression. FEBS Lett **287**:71–74.

20.   **Lecchi S**, **Nelson CJ**, **Allen KE**, **Swaney DL**, **Thompson KL**, **Coon JJ**,
Sussman MR, **Slayman CW**. 2007. Tandem phosphorylation of Ser-911 and Thr-
912 at the C terminus of yeast plasma membrane H+-ATPase leads to glucose-
dependent activation. J Biol Chem **282**:35471–35481.

21.   **Eraso P**, **Portillo F**. 1994. Molecular mechanism of regulation of yeast plasma
membrane H(+)-ATPase by glucose. Interaction between domains and
identification of new regulatory sites. J Biol Chem **269**:10393–10399.

22.   **Goossens A**, **La Fuente de N**, **Forment J**, **Serrano R**, **Portillo F**. 2000.
Regulation of yeast H(+)-ATPase by protein kinases belonging to a family
dedicated to activation of plasma membrane transporters. Mol Cell Biol **20**:7654–
7661.

23.   **Rolland F**, **Winderickx J**, **Thevelein JM**. 2002. Glucose-sensing and -signalling
mechanisms in yeast. FEMS Yeast Res **2**:183–201.

24.   **Ozcan S**, **Dover J**, **Rosenwald AG**, **Wolfl S**, **Johnston M**. 1996. Two glucose
transporters in Saccharomyces cerevisiae are glucose. Proceedings of the
National Academy of Sciences **93**:12428–12432.

25.   **Xue Y**, **Batlle M**, **Hirsch JP**. 1998. GPR1 encodes a putative G protein-coupled
receptor that associates with the Gpa2p Galpha subunit and functions in a Ras-
independent pathway. EMBO J **17**:1996–2007.

26.   **Viegas CA**, **Sá-Correia I**. 1991. Activation of plasma membrane ATPase of
Saccharomyces cerevisiae by octanoic acid. J Gen Microbiol **137**:645–651.

27. **Pontes MH**, **Sevostyanova A**, **Groisman EA**. 2015. When Too Much ATP Is
Bad for Protein Synthesis. J Mol Biol **427**:2586–2594.

28. **Browne SE**. 2013. When too much ATP is a bad thing: a pivotal role for P2X7
receptors in motor neuron degeneration. J Neurochem **126**:301–304.

29. **Larsson C**, **Nilsson A**, **Blomberg A**, **Gustafsson L**. 1997. Glycolytic Flux Is
Conditionally Correlated with ATP Concentration in. J Bacteriol **179**:7243–7250.

30. **Mazón MJ**, **Eraso P**, **Portillo F**. 2015. Specific phospho-antibodies reveal two
phosphorylation sites in yeast Pma1 in response to glucose. FEMS Yeast Res.

31. **Thomsson E**, **Larsson C**, **Albers E**, **Nilsson A**, **Franzén CJ**, **Gustafsson L**.
2003. Carbon starvation can induce energy deprivation and loss of fermentative
capacity in Saccharomyces cerevisiae. Appl Environ Microbiol **69**:3251–3257.

32. **Dechant R**, **Peter M**. 2014. Cytosolic pH: A conserved regulator of cell growth?
Mol Cell Oncol **1**:e969643–4.

33. **Thevelein JM**, **Hohmann S**. 1995. Trehalose synthase: guard to the gate of
glycolysis in yeast? Trends Biochem Sci **20**:3–10.

34. **Choi K-M**, **Kwon Y-Y**, **Lee C-K**. 2015. Disruption of Snf3/Rgt2 glucose sensors
decreases lifespan and caloric restriction effectiveness through Mth1/Std1 by
adjusting mitochondrial efficiency in yeast. FEBS Lett **589**:349–357.

35. **Moriya H**, **Johnston M**. 2004. Glucose sensing and signaling in Saccharomyces
cerevisiae through the Rgt2 glucose sensor and casein kinase I. Proceedings of
the National Academy of Sciences **101**:1572–1577.

36. **Flick KM**, **Spielewoy N**, **Kalashnikova TI**, **Guaderrama M**, **Zhu Q**, **Chang H-C**,
Wittenberg C. 2003. Grr1-dependent inactivation of Mth1 mediates glucose-
induced dissociation of Rgt1 from HXT gene promoters. Molecular Biology of the
Cell **14**:3230–3241.

37. **Lakshmanan J**, **Mosley AL**, **Ozcan S**. 2003. Repression of transcription by Rgt1
in the absence of glucose requires Std1 and Mth1. Curr Genet **44**:19–25.

38. **Jouandot D**, **Roy A**, **Kim J-H**. 2011. Functional dissection of the glucose
signaling pathways that regulate the yeast glucose transporter gene (HXT)
repressor Rgt1. J Cell Biochem **112**:3268–3275.

39. **Ryan OW**, **Skerker JM**, **Maurer MJ**, **Li X**, **Tsai JC**, **Poddar S**, **Lee ME**,
DeLoache W, **Dueber JE**, **Arkin AP**, **Cate JHD**. 2014. Selection of chromosomal
DNA libraries using a multiplex CRISPR system. Elife **3**.

40. **Benjamin DI**, **Louie SM**, **Mulvihill MM**, **Kohnz RA**, **Li DS**, **Chan LG**, **Sorrentino**
A, **Bandyopadhyay S**, **Cozzo A**, **Ohiri A**, **Goga A**, **Ng S-W**, **Nomura DK**. 2014.

Inositol phosphate recycling regulates glycolytic and lipid metabolism that drives
cancer aggressiveness. ACS Chem Biol **9**:1340–1350.

41.   **Luo B**, **Groenke K**, **Takors R**, **Wandrey C**, **Oldiges M**. 2007. Simultaneous
determination of multiple intracellular metabolites in glycolysis, pentose
phosphate pathway and tricarboxylic acid cycle by liquid chromatography-mass
spectrometry. J Chromatogr A **1147**:153–164.

42.   **Kaiser CA**, **Chen EJ**, **Losko S**. 2002. Subcellular fractionation of secretory
organelles. Meth Enzymol **351**:325–338.

43.   **Li X**, **Yu VY**, **Lin Y**, **Chomvong K**, **Estrela R**, **Park A**, **Liang JM**, **Znameroski
EA**, **Feehan J**, **Kim SR**, **Jin Y-S**, **Glass NL**, **Cate JHD**. 2015. Expanding xylose
metabolism in yeast for plant cell wall conversion to biofuels. Elife **4**.

Figure Legends

Figure 1. Metabolite profile of cells provided with glucose or cellobiose. (A) Significant changes of intracellular metabolite levels in cells provided with cellobiose compared to cells provided glucose as a sole carbon source. (B) Schematic representation of metabolite changes. The green and red dots represented higher and lower relative metabolite levels in cells provided with cellobiose compared to cells provided with glucose. Statistical analyses are described in the Materials and Methods.

Figure 2. Manipulation of phosphofructokinase (*PFK1*) and plasma membrane ATPase *(PMA1)*. (A) Schematic representation of cellobiose consumption route in the upper glycolytic pathway. Glc, G6P, F6P and ATP are highlighted, and were found in higher abundance when cellobiose was provided in comparison to glucose. (B) Cellobiose consumption profile and (C) cell density profile of the strains with ATP-insensitive Pfk1 (*pfk1m*), constitutively active Pma1 (*pt*) and the combination of both mutations (*pfk1m-pt*) in comparison to the cellobiose pathway-only strain, here used as wild-type (WT). (D) Relative abundance of G6P, F1,6BP and ATP levels of the WT, *pfk1m, pt* and *pfk1m-pt* strains, relative to the WT strain fermenting cellobiose. The experiments were carried out in 5 biological replicates, with standard errors from the mean shown.

Figure 3. Carbon starvation-like state of the plasma membrane ATPase (*PMA1*) in cellobiose-fermenting cells. (A) Phosphorylation states of Pma1 residues S911 and T912 under carbon starvation, glucose metabolizing and cellobiose metabolizing conditions. (B) Cellobiose consumption efficiency (E_c) of cells expressing Pma1 with phosho-mimic or phosphorylation-preventing mutations at positions serine 911 (S911)

and threonine 912 (T912). Shown are the mean and standard deviation for 3 biological

replicates. The experiments were carried out in 2 biological replicates, with standard

errors from the mean shown.

Figure 4. Effect of glucose sensor deletions. (A) Cellobiose consumption efficiency

(E_c) and cellular ATP levels of strains with different glucose sensor deletions and

different combinations of constitutively-active Pma1 mutations. Shown are the averages

and standard deviations of three biological replicates. (B) Correlation of E_c and cellular

ATP. Standard deviations for three biological replicates are shown for each point.

Figure 5. Glucose sensor deletions and cellular ATP levels. (A) Additive effect of the

increase in E_c and the decrease in cellular ATP of *sr*∆ and *pt* in *sr*∆-*pt* strain. The

experiments were carried out in 5 biological replicates, with standard errors form the

mean shown. (B) Specific Pma1 ATPase of WT*, pt, sr*∆ and *sr*∆-*pt* strains measured

from normalized membrane fractions of cells harvested at mid-log phase. The

experiments were carried out in 3 biological replicates, with standard errors from the

mean shown.

Figure 6. Schematic representation of ATP homeostasis and cellular regulation of

cellobiose fermentation. Excess ATP inhibited phosphofructokinase (Pfk1), resulting

in an upper glycolytic metabolite buildup and slow cellobiose consumption. The buildup

of ATP may be caused by low activity of Pma1, with no phosphorylation at position 911.

Additionally, glucose sensors Snf3 and Rgt2 together influenced cellular ATP levels via

Pma1 and other mechanisms to be identified.

Supplementary Figure legends

Figure S1. Metabolite profile of cells provided with glucose or cellobiose. (A) cell

density and sugar consumption profiles of the strains used for metabolite profiling

experiments. (B) Heatmap representation of steady state intracellular metabolite

abundance of cells provided with glucose or cellobiose under anaerobic conditions.

Shown are the results of 5 biological replicates. Asterisks (*) mark identified compounds

that were significantly different between glucose and cellobiose conditions at a

significance threshold of a p-value < 0.01.

Figure S2. Cellobiose consumption profiles and efficiency of Snf3, Rgt2 and Gpa2

mutants. (A) Cellobiose consumption profiles of strains with consitutively active Snf3

(R229K), Rgt2 (R231K) and Gpa2 (R273A). (B) Cellobiose consumption profiles of

SNF3, *RGT2*, and *GPA2* double deletion combinations. (C) Cellobiose consumption

profiles of *SNF3* and *RGT2* single deletion in comparison to the *sr∆, srg∆ and sr∆-pt*

strains. (D) Cellobiose consumption profiles of *GPA2* deletion (*g∆)* in comparison to the

WT strain. In panels A, B, C, and D, the results of 3 biological replicates are shown.

Figure S3. Relationship of ethanol productivity parameters and cellobiose

consumption efficiency (E_c). Ethanol production rate and the final concentration of

ethanol are plotted against E_c.

660
661   **Table S1:** List of strains constructed, CRISPRm guides and primers.

A

Carbon Starvation   Glucose   Cellobiose

B

Δ916		
D	D	
D	T	
A	T	
D	A	
S	T	← Pma1-WT
S	A	
S	D	

S911   T912

E_c

0   1   2   3   4