System-wide automatic extraction of functional signatures in *Pseudomonas aeruginosa* with eADAGE

Jie Tan[1], Georgia Doing[2], Kimberley A. Lewis[2], Courtney E. Price[2], Kathleen M. Chen[3], Kyle C. Cady[4,5], Barret Perchuk[4,5], Michael T. Laub[4,5], Deborah A. Hogan[2], Casey S. Greene[3,6,7,*]

1. Department of Molecular and Systems Biology, Geisel School of Medicine at Dartmouth, Hanover, NH, USA 03755
2. Department of Microbiology and Immunology, Geisel School of Medicine at Dartmouth, Hanover, NH, USA 03755
3. Department of Systems Pharmacology and Translational Therapeutics, University of Pennsylvania, Philadelphia, PA, USA 19104
4. Department of Biology, Massachusetts Institute of Technology, Cambridge, MA, 02139, USA.
5. Howard Hughes Medical Institute, Cambridge, MA, 02139, USA.
6. Institute for Translational Medicine and Therapeutics, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA 19104
7. Institute for Biomedical Informatics, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA 19104

*To whom correspondence should be addressed: csgreene@mail.med.upenn.edu

Running title: Extracting signatures from public data

Abstract

Abundant public expression data capture gene expression across diverse conditions. These steady state mRNA measurements could reveal the transcriptional consequences of cells' genetic backgrounds or their responses to the environment. However, public data remain relatively untapped, in part because extracting biological signal as opposed to technical noise remains challenging. Here we introduce a procedure, termed eADAGE, that performs unsupervised integration of public expression data using an ensemble of neural networks as well as heuristics that, given a dataset, help users identify an appropriate level of model complexity. This ensemble modeling approach captures biological pathways more clearly than existing methods, enabling analyses that span entire public gene expression compendia such as that for the bacterium *Pseudomonas aeruginosa*. These analyses reveal a previously undiscovered feature of the phosphate starvation response apparent in public data: a sensor kinase, KinB, that is required for full activation of the response to phosphate at intermediate concentrations. Our molecular validation experiments confirm this role of KinB and our screen of a histidine kinase knock out collection confirmed the prediction's specificity. Public data are captured from a broad range of conditions in diverse organism backgrounds and may provide a unique opportunity to identify these subtle and context-specific regulatory interactions. Algorithms that extract biological signal from these data, such as eADAGE, can highlight opportunities to discover mechanisms that are apparent from but unrealized in public data.

Keywords

denoising autoencoders/ensemble modeling/gene expression/*Pseudomonas aeruginosa*/PhoB crosstalk

Introduction

Available gene expression data are outstripping our knowledge about the organisms that we're measuring. Ideally each organism's data reveals the principles underlying gene regulation and consequent pathway activity changes in every condition in which gene expression is measured. Extracting this information requires new algorithms, but many commonly used algorithms are supervised. These algorithms require curated pathway knowledge to work effectively, and in many species such resources are biased in various ways (Schnoes *et al*, 2013; Gillis & Pavlidis, 2013; Greene & Troyanskaya, 2012). Annotation transfer can help, but such function assignments remain challenging for many biological processes (Jiang *et al*, 2016). An unsupervised method that doesn't rely on annotation transfer would bypass the challenges of both annotation transfer and biased knowledge.

Along with our wealth of data, abundant computational resources can now power deep unsupervised applications of neural networks (Vincent *et al*, 2008). Denoising autoencoders (DAs), an unsupervised neural network method, are well suited to gene expression data (Tan *et al*, 2015). DAs are more robust than linear approaches such as ICA or PCA in the context of public data, which employ heterogeneous experimental designs, lack shared controls and provide limited metadata (Tan *et al*, 2016b). Investigators can use models trained on public

43 data to reexamine their own data to identify the processes or pathways that are particularly
44 perturbed in their experiment.
45
As a result of a training process that uses stochastic gradient descent, each DA model
represents a local minimum. Because the number of parameters is large, there are many
distinct models that can explain the data. One way to address this challenge is by finding
patterns that are stable across multiple models, which Yu highlights for its importance in
statistical inference (Yu, 2013). Even if run-to-run variability in models obscures some biological
features, these stable cross-model patterns may clearly resolve biological pathways. To directly
target stability, we introduce an unsupervised ensemble modeling procedure that improves
model robustness by constructing an ensemble neural network model. Our approach is
inspired by consensus clustering (Monti *et al*, 2003), which now has become a standard part of
clustering applications for biological datasets.
56
While this new generation of unsupervised data integration methods provides a powerful lens
through which to study complex biological systems, it also raises questions: how many
biological signatures are in the data; and how many samples are needed to find those features?
Heuristics should include data-driven analyses that can be applied in any setting, as well as
knowledge-based heuristics that are suitable in cases where pathway annotations are available.
We introduce both data- and knowledge-based heuristics to suggest an appropriate model size
for a given data compendium.
64
We previously constructed a 50-node ADAGE (Analysis using Denoising Autoencoders for Gene
Expression) model of *P. aeruginosa* because this balanced the need for breadth of the model
with the limits of manual annotation required for the initial validation of the method (Tan *et al*,
2016b). Our heuristics indicate that the existing *P. aeruginosa* compendium can support 300
node models, which we use in this work. We compare ADAGE models with models generated
through an ensemble form of ADAGE (eADAGE) and find that eADAGE models have greater
stability. The eADAGE models are not only more robust, but also more successfully capture
pathway features in the data. While ADAGE allows comparison of pathway activities within an
experiment, eADAGE models also enable analyses that cut across an organism's gene
expression compendium. We analyze *P. aeruginosa*'s responses to multiple types of media with
eADAGE. This cross-compendium analysis reveals a gene expression signature for the response
to low-phosphate in certain media, which our molecular validation confirms. Our analyses of
this signature across the compendium revealed a previously undiscovered link between KinB
and PhoB activity. PhoB is known to mediate the low phosphate response and is canonically
regulated by a histidine kinase, PhoR. However, our molecular validation shows PhoA activity,
which indicates PhoB activity, independent of PhoR in some conditions and dependent on KinB
in some conditions. Our experiments confirm that KinB regulates PhoB activity at physiologically
relevant phosphate concentrations and that this activity is specific to KinB among kinases.
83
84 In summary, eADAGE more precisely and robustly represents biological processes and pathways.
85 It allows biologists to identify not only differentially active signatures within one experiment,

86 but also cross-compendium patterns that reveal undiscovered regulatory mechanisms captured
87 within existing public data.
88

Results

Analysis of the effects of model size on pathway characterization of ADAGE models from *P. aeruginosa* gene expression.

While ADAGE models are constructed without the use of any curated information, we can use experimentally-derived knowledge on gene functions to provide heuristic information about the number and types of pathways captured by a model and determine how this varies with model size. We define a functional signature learned by an ADAGE model as a set of genes that contribute the highest positive or highest negative weights to a specific node (see methods for detail). Therefore, one node results in two gene signatures, one on each high weight side. These high-weight (HW) genes are often involved in a common biological process as demonstrated by the fact that there is often a statistically significant enrichment in specific KEGG pathways within each signature. For models of different sizes (10-1000 nodes), we determined the number of KEGG pathways significantly associated with at least one gene signature in a model, referred to as KEGG pathway coverage for that model, and found that KEGG pathway coverage increased as model size increased until a model size of approximately 300 (Figure 1A). The number of pathways per node (including pathways associated with both the positive and negative signatures in a node) for all nodes with at least one associated KEGG pathway decreased as model size increased (Figure EV1), suggesting that multiple pathways were grouped in small models and were separated into more discrete features in large models with more nodes.

Methods that use pathway coverage to identify the appropriate model size for a collection of samples require curated pathways, but a parallel analysis can be performed without this requirement. During ADAGE training, neural networks are trained to reconstruct the input from data with noise added, and calculating the reconstruction error does not require any curated information. The reconstruction error can also be used to estimate model sizes that can be supported by the available data. The reconstruction error quickly decreases as model size increases and reaches a plateau at model size of approximately 300 (Figure 1B). Further increasing model size does not improve reconstruction, suggesting that the available data are insufficient to support larger models. Therefore, considering the knowledge-driven and data-driven heuristics together, we identified a 300-node model as most appropriate for the existing *P. aeruginosa* gene expression compendium.

Analysis of the effects of sample number in the training set on ADAGE models

The expression compendium contains 1051 samples from 125 experiments. We next performed a subsampling analysis in which we trained ADAGE models on randomly selected sets of 100, 200, 500, and 800 expression profiles. We examined the number of KEGG pathways associated with at least one gene signature (pathway coverage) as a function of the size of the training set (Figure 1C). In the 50-node models, the size used in (Tan *et al*, 2016b), the average KEGG pathway coverage at each training size increased significantly up to 500 samples (Tukey's HSD adjusted p-values < 0.05 between models trained with 100, 200, and 500 samples), but

differences beyond 500 training samples were not significant (Tukey's HSD adjusted p values >
0.05 between models trained with 500, 800, and 1051 samples). For 300-node models, pathway
coverage showed significant increases (Figure 1C) between the models constructed with 100,
200, 500, and 800 samples (Tukey's HSD adjusted p-values < 0.05) but not between 800 and
1051 (Tukey's HSD adjusted p-value > 0.05). The slower increase in pathway coverage when
sample size is relatively large suggests redundancy in the compendium, potentially due to
biological replicates or experiments probing similar processes. This highlights the importance of
data that capture diverse processes.
138
Using the subsampling strategy, we also evaluated the reconstruction error of each model on
its training set and a randomly chosen held out test set of 200 samples. As sample size
increased, training reconstruction errors increased slightly while testing reconstruction errors
dropped dramatically (Figure 1D). We fitted exponential models between sample size and the
differences of training and testing errors (R^2= 0.78 for 50-node models and R^2= 0.83 for 300-
node models). We extrapolated from these models to predict that testing errors would
approximately match training errors when sample size was 782 for 50-node models and 1076
for 300-node models. These results suggested that smaller models were less sensitive to sample
size, likely because they have fewer parameters to fit and also that our 1051 sample
compendium was sufficient to train a 300-node model.
149
eADAGE: ensemble modeling improves the model breadth, depth, and robustness
Individual ADAGE models capture a local minimum, with models of the same size capturing
different pathways. This occurs because each ADAGE model is initialized with random weights,
and the training processes are sensitive to these initial conditions. eADAGE, in which we built
an ensemble version of individual ADAGE models, took advantage of this variation to enhance
model robustness. Each eADAGE model integrated nodes from 100 individual ADAGE models
(Figure 2A). To unite nodes from different models, we applied consensus clustering on nodes'
weight vectors. Our previous ADAGE analyses showed that HW genes characterized each node's
biological significance, so we designed a weighted Pearson correlation to incorporate gene
weights in building eADAGE models (see methods). We compared eADAGE to two primary
baseline methods: individual ADAGE models and corADAGE, which combined nodes with an
unweighted Pearson correlation. For direct comparison, the model sizes of ADAGE, eADAGE,
and corADAGE were all fixed to 300 nodes, which we found to be appropriate for the current
P.a expression compendium.
164
eADAGE models exhibited greater KEGG pathway coverage than those generated by other
methods. We evaluated ADAGE, corADAGE, and eADAGE for the number of covered KEGG
pathways (Figure 2B). Both corADAGE and eADAGE covered significantly more KEGG pathways
than ADAGE (t-test p-value of 1.04e-6 between corADAGE (n=10) and ADAGE (n=1000) and t-
test p-value of 1.41e-6 between eADAGE (n=10) and ADAGE (n=1000)). Moreover, eADAGE
models covered, on average, 10 more pathways than corADAGE (t-test p-value of 1.99e-3, n=10
for both groups), confirming the critical roles of an ADAGE node's HW gene signatures in
defining biological pathways. Genes that participate in multiple pathways can influence
pathway enrichment analysis, a factor termed pathway crosstalk (Donato *et al*, 2013). If

eADAGE signatures tended to include genes that participated in many pathways, this could also
drive the increase in number of observed pathways. To control for this, we performed crosstalk
correction (Donato *et al*, 2013). After correction, the total number of covered pathways
dropped approximately by half (Figure EV2), but eADAGE still covered significantly more
pathways than corADAGE (t-test p-value of 1.60e-3) and ADAGE (t-test p-value of 6.16e-07).
These results suggested that eADAGE effectively integrates multiple models to more broadly
capture pathway signals embedded in diverse gene expression compendia.
181
We next evaluated how specifically and completely signatures learned by the models capture
known KEGG pathways. We use each gene signature's FDR corrected p-value for enrichment of
a KEGG pathway as a combined measure, as this captures both the sensitivity and specificity. If
a pathway was significantly associated with multiple gene signatures in a model, we only
considered its most significant association. We found that 71% of pathways were more
significantly enriched (had lower median p-values) in corADAGE models (n=10) when compared
to individual ADAGE models (n=100) (Figure EV3). This increased to 87% for eADAGE (n=10). We
also directly compared eADAGE and corADAGE by this measure and observed that 74% of
pathways were more significantly enriched in eADAGE. Our earlier evaluation of pathway-based
heuristics showed that different pathways were best captured at different model sizes (Figure
EV3). We next compared the 300-node eADAGE model to individual models of each size.
Although the 300-node eADAGE models were constructed only from 300-node ADAGE models,
we found that 69% pathways were more significantly enriched (i.e. lower median p-values) in
eADAGE models than ADAGE models of any size, including those with more nodes than the
eADAGE models (Figure EV3). Three example pathways that are best captured either when
model size is small, large, or in the middle are all well captured in the 300-node eADAGE model
(Figure 2C). These results demonstrate that eADAGE's ensemble modeling procedure captures
signals across model sizes more effectively than individual ADAGE and corADAGE models. Thus
eADAGE more completely and precisely captures the gene expression signatures of biological
pathways.
202
We designed eADAGE to provide a more robust analysis framework than individual ADAGE
models. To assess this, we examined the percentage of models that covered each pathway
(coverage rate) between ADAGE and eADAGE (Figure EV4). The pathways covered by each
individual ADAGE model were highly variable. Most KEGG pathways were covered by less than
half of individual models but more than half of eADAGE models (Figure EV5), suggesting that
eADAGE models were more robust than individual ADAGE models. We excluded all pathways
always covered by both individual ADAGE and eADAGE models and observed that 72% of the
remaining pathways were covered more frequently by eADAGE than ADAGE. This suggests that
their associations are stabilized through the ensemble construction procedures. In summary,
these comparisons of eADAGE and ADAGE reveal that not only are more pathways captured
more specifically, but also those that are captured are captured more consistently.
214
Principle component analysis (PCA) and independent component analysis (ICA) have been
previously used to extract biological features and build functional gene sets (Engreitz *et al*, 2010;
Raychaudhuri *et al*, 2000; Gong *et al*, 2007; Alter *et al*, 2000; Raychaudhuri *et al*, 2000; Lutter *et*

al, 2009; Frigyesi *et al*, 2006; Chen *et al*, 2008; Roden *et al*, 2006; Ma & Kosorok, 2009). We
performed PCA and generated multiple ICA models from the same *P. aeruginosa* expression
compendium and evaluated their KEGG pathway coverage following the same procedures used
for eADAGE. eADAGE substantially and significantly outperforms PCA in terms of pathway
coverage (Figure 2D). We observed that low-order PCs tend to be associated with more
pathways than high-order PCs, which is consistent with the higher variance explained by low-
order PCs. ICA and eADAGE covered a similar number of pathways at the significance cutoff of
FDR 0.05. However, we observed that eADAGE represented KEGG pathways more precisely
than ICA. Specifically, among pathways significantly enriched in either approach, 68% pathways
exhibited more significant enrichment in eADAGE. Increasing the significance threshold for
pathway coverage demonstrates the advantage of eADAGE (Figure 2D).
229
Taken together, the eADAGE method outperformed PCA, ICA, ADAGE, and corADAGE in
capturing KEGG pathways of *P. aeruginosa*. eADAGE had higher pathway coverage, covered
pathways more specifically, and more robustly than existing methods. Though the primary goal
of eADAGE is not function prediction, it captures pathway signals in transcriptional data
significantly more effectively than existing unsupervised data integration methods. These
results clearly indicate that eADAGE extracts biological patterns from gene expression
compendia.
237
Elucidating functional signatures that are indicative of growth medium
Gene expression analysis experiments are generally designed to include only a small number of
variables. Thus, it is not surprising that in the majority of publically available *P. aeruginosa* gene
expression experiments, the growth medium was held constant while there was variation in
other parameters such as strain, environmental parameters (i.e. pH, temperature, growth
phase, or gaseous atmosphere), or culture amendments (antibiotics, etc.). Though it was rare
for the base medium to vary within a single experiment, our manual annotation showed that 78
different base media were used across the gene expression compendium (Table EV1). These
media exhibit great diversity in the number of different medium components, the abundance of
different nutrients, and the form in which nutrients were provided. Some medium groups
contained numerous samples from multiple experiments other media types were unique to a
single experiment.
250
For biological evaluation, we built a single new eADAGE model with 300 nodes. In this model
we identified cross-media signals of biological importance. The model's weight matrix (Table
EV2), positive and negative gene signatures for each node (Table EV3) and signature activities
for each sample in the compendium (Table EV4) are provided. For each gene signature, we
used its mean activity for samples prepared from cells grown in a given medium compared to
the mean activity of the signature across all the remaining samples in the compendium
normalized to the total range to provide an activation score. We then further reduced this set
by filtering to all signature-medium relationships with an activation score greater than or equal
to 0.4 (Table EV5), which included only the top 2.4% of potential pairs. To capture pan-media
patterns we identified signatures that had multiple signature-media activation scores above this
threshold and averaged their activation scores. Table EV6 lists the media groups on which gene

signatures were most differentially active (a complete list of signature-media associations is in
Table EV7).
264
The signature Node164pos had the highest pan-media activation score. To evaluate the basis
for the high activation score, we examined the signature's underlying activity across media. This
revealed that Node164pos was highly active in King's A medium, peptone medium, and
NGM+<0.1mM phosphate (NGMlowP), but not in NGM+25mM phosphate (NGMhighP) (Figure
3A). The difference in Node164pos activity between NGMlowP and NGMhighP suggested that
the genes in this signature respond to phosphate concentrations. Consistent with this proposal,
a KEGG pathway enrichment analysis of Node164pos genes suggested a strong enrichment in
genes involved in phosphate acquisition (Table EV6). The other two media in which *P.*
aeruginosa gene expression leads to consistently high Node164pos activity (Peptone and King's
A) also had low phosphate concentrations (0.4 mM) relative to other media in the compendium.
For example, commonly used LB has a phosphate concentration of approximately 4.5 mM
(Bertani, 2004) and many others have concentrations above 20 mM.
277
Many of the phosphate responsive genes in Node164pos are known to be under the control of
the transcription factor PhoB in *P. aeruginosa* (Santos-Beneit, 2015; Blus-Kadosh *et al*, 2013;
Bielecki *et al*, 2015) and *phoB* itself is in this signature. Analysis of genes in Node164pos finds
that this signature has the largest overlap with the PhoB regulon (see methods) in comparison
to all other gene signatures learned by this eADAGE model (FDR q-value of 8.1e-29 in
hypergeometric test). The transcript levels of genes in Node164pos are higher in peptone,
King's A, and NGM+<0.1 phosphate medium relative to the other samples in the compendium
including NGM+ 25 mM phosphate (Figure 3B).
286
Among the highest weight genes in Node164pos is a gene that encodes alkaline phosphatase
(PhoA), an enzyme with an activity that can be easily measured using a colorimetric assay. As
expected, PhoA activity (blue color) was high when *P. aeruginosa* was grown on NGMlowP and
not when grown on NGMhighP (Figure 4A). The same trend was observed in another medium,
MOPS, with the same high and low phosphate concentrations. Also consistent, PhoA was not
active on the phosphate replete medium LB. Furthermore, PhoA activity was dependent on
PhoB and the PhoB-activating histidine kinase PhoR, which is consistent with previous
publications (Bielecki *et al*, 2015). These results provide striking evidence that low phosphate
media induced PhoB activity as predicted by the eADAGE analysis and previous
characterizations of the *P. aeruginosa* phosphate response.
297
Interestingly, Node164pos activity exhibited a wide spread in PIA medium (0.8 mM phosphate),
with six samples having high activities and the other six samples having low activities (Figure
3A). Each set of six samples came from a different study. All of the strains in the first sample set,
published in 2012, in which Node164pos was low used a PAO1 *kinB*::Gm[R] mutant background
(Damron *et al*, 2012). The second, published in 2013, used a PAO1 strain with *kinB* intact and
showed high Node164pos activity on PIA (Damron *et al*, 2013). The fact that *kinB* mutant
samples in PIA show significantly lower Node164pos activity suggests that KinB may be a
regulator of PhoB on PIA.

306
Expectedly, PhoA activity was high in peptone and King's A media, as in NGMlowP. Further, as
predicted by eADAGE Node164pos activity, PhoA activity was evident and was KinB dependent
on PIA medium (Figure 4B). Notably, PhoA activity was still dependent on PhoB and PhoR as it
was on peptone and King's A. Over time the ΔphoR mutants developed PhoA activity on all
three media, but the ΔkinB on PIA did not (Figure 4C). Recovery of PhoA activity in ΔphoR
mutants suggests that there are PhoR-independent paths for PhoB activation. The co-
dependence on KinB and PhoR suggest that these kinases do not perform redundant functions
but rather regulate PhoB in conjunction with each other. To determine if the deletion of kinases
non-specifically altered PhoB activation, we screened 63 in-frame deletion mutants each lacking
a histidine kinase (Table EV8) to identify mutants altered in PhoA activation on PIA. Other than
ΔphoR, ΔkinB was the only strain lacking PhoA activity on NGMlowP. This suggests that PhoA
activity is not altered by changes in generic kinase activity but rather the interaction with KinB is
specific.
320
Notably, although PIA, peptone and King's A contain low phosphate levels compared to rich
media such as LB, PIA has a higher phosphate concentration (0.8mM) than do peptone and
King's A (0.4mM). To test whether the moderately low level of phosphate in PIA provokes KinB
regulation of PhoA, we conducted a titration experiment in MOPS minimal medium. PhoA
activity was consistent with that on NGM and other media. Further, while the PAO1 wild type
(WT) showed PhoA activity at 0.5 mM, PAO1 ΔkinB did not (Figure 4D). This shows KinB
regulation of PhoB in moderately low phosphate and not in lower phosphate. Noting that PIA
has a phosphate concentration of 0.8 mM and our titration experiment in MOPS showed kinB-
sensitivity at 0.5 mM, it is likely that the precise concentration at which KinB regulates PhoA
depends on the background medium. Node164pos activity across the compendium suggested,
and experimental evidence confirmed, that KinB regulates PhoA at moderately low phosphate
levels. To our knowledge, KinB has not been previously implicated in the activation of PhoB,
although cross-compendium eADAGE analysis of two PIA experiments was capable of revealing
this relationship.
335
336 In summary, eADAGE effectively extracted biologically meaningful features, accurately
337 indicated their activity in multiple media spanning numerous independent experiments, and
338 revealed a novel regulatory mechanism. By summarizing gene-based expression information
339 into biologically relevant features, eADAGE greatly simplifies analyses that cut across large gene
340 expression compendia.
341

Discussion
For unsupervised approaches, it has been very challenging to determine the appropriate model
complexity. In supervised learning problems, predictors can be assessed through cross-
validation accuracies. To our knowledge, there is not yet a similar well-established approach to
estimate appropriate model size for unsupervised feature construction of gene expression data.
Here we develop heuristics that target two aspects of the problem: the model needs to be well
supported by the amount of available data and the extracted features should well resemble
known biological processes. Our data-driven heuristics can be applied to organisms for which

gene-process annotations are lacking. We expect that additional data will support larger models,
especially data that measure experimental conditions that are not tested in the existing
compendium.
353
We also contribute a novel eADAGE algorithm. This algorithm combines multiple ADAGE
models into one ensemble model to address model variability due to stochasticity and local
minima. The algorithm is inspired by consensus clustering, which reconciles the differences in
cluster assignments in multiple runs. Comparable approaches have also been applied for ICA,
where researchers have used the centrotypes of multiple ICA models as the final model
(Frigyesi *et al*, 2006). The ICA centrotype approach for ADAGE corresponds to corADAGE, and
our comparison of eADAGE and corADAGE shows that eADAGE not only covers more biological
pathways, but also results in cleaner representations of biological pathways. The results of our
direct comparison suggest that placing particular emphasis on the genes most associated with a
particular signature may be a useful property for other unsupervised feature construction
algorithms in biology.
365
Our eADAGE model revealed patterns that were obvious in the analysis of the large data
compendium containing experiments performed in 78 different media, but that were not
necessarily evident in individual experiments. For example, our cross-compendium analysis of
expression patterns specific to certain media determined those media in which *P. aeruginosa*
was in a phosphate limited, PhoB-ON state. PhoB is a global regulator, and thus an awareness
of the different states of the PhoB regulon in different media will likely provide important
insight into medium-specific phenotypes of relevance. Using eADAGE, we uncovered a subtle
aspect of the phosphate starvation response that depends on a histidine kinase not previously
associated with the PhoB-dependent phosphate response pathway. Bacteria have evolved
many mechanisms to insulate response pathways from each other (Podgornaia & Laub, 2013),
but cross-talk, wherein a sensor kinase from one pathway phosphorylates a response regulator
from another, has been suggested to explain the complexity of signaling networks, including
that of PhoB (Ninfa *et al*, 1988; Fisher *et al*, 1995). Thus, it is possible that both PhoR and KinB
directly activate PhoB. While there are many examples of one kinase partially compensating
when the cognate kinase is absent, it is more challenging to find conditions where two kinases
are needed for full response regulator activation (Verhamme *et al*, 2002). Alternatively, KinB
may influence PhoB activity indirectly by regulating activities that affect PhoB levels, PhoB
dephosphorylation, or PhoB localization, or protein-protein interactions. Future work will
further address the mechanistic aspects of this model. We propose that moderate levels of
phosphate, like those in PIA, provide a niche for crosstalk: the activity of PhoR is low enough
that the interaction with KinB is needed for full PhoB activity on this medium. Together, PhoR
and KinB may enable a more sensitive and effective response to phosphate limitation. Without
the eADAGE analysis across multiple media, we would not have found this nuanced mechanism.
389
In the 300-node eADAGE model used for medium analysis, 81% genes in *P.a.* PAO1 genome are
included in at least one gene signature. Among all signatures, 59% have a KEGG pathways
associated with them. Among the remaining 41% of signatures, 17 of them had ten or fewer
genes and may not have associated with KEGG pathways for this reason. Some others were

associated with known pathways that have not yet been annotated in KEGG, such as
Node174pos which encodes genes encoded by the P2 phage or Node150neg which contains
genes involved in the formation of surface associated communities referred to as biofilms. Thus
eADAGE can group functionally linked genes in ways that may facilitate the identification or
annotation of pathways.
399
There are now abundant public gene expression data. Cross-compendium analyses provide the
opportunity to efficiently use existing data to identify regulatory patterns that are evident
across multiple experiments, datasets, and labs. To tap this potential, we will require algorithms
that robustly integrate these diverse datasets in a manner that is not tied to only aspects of
biology that are well understood. We expect that robust unsupervised data integration
methods, like eADAGE, will play a key role in this process.
406

Materials and Methods
Data processing
We followed the same procedures for data collection, processing, and normalization from (Tan
et al, 2016b) and updated the *P. aeruginosa* gene expression compendium to include newly
uploaded datasets on GPL84 platform from the ArrayExpress database (Rustici *et al*, 2013) on
31 July 2015. The updated *P. aeruginosa* compendium contains 125 datasets with 1051
individual genome-wide assays. We provide the *P. aeruginosa* expression compendium (Dataset
EV1) along with all the code used in this paper (Tan *et al*, 2016a). The eADAGE repository is also
tracked under version control at https://bitbucket.org/greenelab/eadage.
416

Construction of ADAGE models
We constructed ADAGE models as described in (Tan *et al*, 2016b). To summarize the process
and outputs, we constructed a denoising autoencoder for the gene expression compendium.
Denoising autoencoders model the data in a lower dimension than the input space, and the
models are trained with random gene expression measurements set to zero. Thus an ADAGE
model must learn gene-gene dependencies to fill in this missing information. Once the ADAGE
model is trained, each node in the hidden layer contains a weight vector. These positive and
negative weights represent the strength of each gene's connection to that node.
425

Gene signatures as sign-specific high-weight gene sets
In previous work (Tan *et al*, 2016b) we defined high-weight (HW) genes as those in the
extremes of the weight distribution on the positive or negative side of a node. Here, we use a
more granular definition that accounts for sign specificity. Each node's gene weights are
approximately normal and centered at zero in ADAGE models (Tan *et al*, 2015, 2016b). We
defined positive HW genes as those that were more than 2.5 standard deviations from the
mean on the positive side, and negative HW genes as those that were more than 2.5 standard
deviations from the mean on the negative side. After this split, a model with n nodes provides
2n gene signatures. Because a node is simply named by the order that it occurs in a model, we
named two gene signatures derived from one node as "NodeXXpos" and "NodeXXneg".
436

KEGG pathway enrichment analysis

To evaluate the biological relevance of gene signatures extracted by an ADAGE model, we
tested how they relate to known KEGG pathways (Kanehisa, 2000). We tested a signature's
association with each KEGG pathway using hypergeometric test and corrected the p-value by
the number of KEGG pathways we tested following the Benjamini–Hochberg procedure. We
used a false discovery rate of 0.05 as the significance cutoff.
443
Genes can be annotated to multiple pathways. To control for this effect in our analysis, we also
performed a parallel analysis after applying crosstalk correction as described in (Donato *et al*,
2013). This approach uses expectation maximization to map each gene to the pathway in which
it has the greatest predicted impact. A gene-to-pathway membership matrix, defined using
KEGG pathway annotations, initially makes the assumption that each gene's role in all of its
assigned pathways remains constant independent of context. We then applied pathway
crosstalk correction using genes' weights for each node in the ADAGE model. We used the
expectation maximization algorithm to maximize the log-likelihood of observing the
membership matrix given each node's weight vector. This process inferred an underlying gene-
to-pathway impact matrix and iteratively estimated the probability that a particular gene g
contributed the greatest fraction of its impact to some pathway P. Upon convergence, we
assigned each gene to the pathway in which it had the maximum impact. The resulting pathway
definitions do not share genes. We then used these corrected definitions for an analysis parallel
to the KEGG process described above.
458
Reconstruction error calculation
The training objective of ADAGE is to take a sample with added noise and return the originally
measured expression values. The error between the reconstructed data and the initial data is
the 'reconstruction error.' To summarize the difference over all genes we used cross-entropy
between the original sample and the reconstruction, which has been widely used with these
methods and in this domain (Vincent *et al*, 2008; Tan *et al*, 2016b). This matches the statistic
used during training of the model. To calculate reconstruction error for a model, we use the
mean reconstruction error across samples.
467
Model size and sample size heuristics
One important parameter of a denoising autoencoder model is the number of nodes in the
hidden layer, which we refer to as the model size. To evaluate the impact of model size and
choose the most appropriate size, we built 100 ADAGE models at each model size of 10, 50, 100,
200, 300, 500, 750, and 1000, using different random seeds. The random seed determines the
initialization statuses of the weight matrix and bias vectors in ADAGE construction and thus
different random seeds will result in training stopped at different local minimums. Other
training parameters were kept the same and set to the values identified as suitable for a gene
expression compendium (Tan *et al*, 2015). In total, 800 ADAGE models with 100 at each model
size were generated in the model size evaluation experiment.
478
To evaluate the impact of sample size on the performance of ADAGE models, we randomly
generated subsets of the *P. aeruginosa* expression compendium with sample size of 100, 200,
500, and 800. We then trained 100 ADAGE models at each sample size, each with a different

combination of 10 different random subsets and 10 different random training initializations. To
evaluate each model, we randomly selected 200 samples not used during training as its testing
set. We performed this subsampling analysis at model size 50 and 300. In total, 800 ADAGE
models were built in the sample size evaluation experiment.
486
Construction of eADAGE models
We constructed ensemble ADAGE (eADAGE) models by combining many individual ADAGE
models in to a single model. For each eADAGE model we combined 100 individual ADAGE
models. The 100 models were trained with identical parameters but distinct random seeds. For
an eADAGE model of size 300, we trained 100 individual models with 300 nodes each, which
provided 30000 total nodes. Each node has a weight vector. We have previously observed that
high-weight genes provided the most information to each node (Tan *et al*, 2016b), so we
calculated a weighted Pearson correlation between each node's weight vectors. Our weighted
Pearson correlation used (|node1 weight|+|node2 weight|)/2 as the weight function for each
gene. We compared this to an unweighted Pearson correlation (corADAGE) as well a baseline
ADAGE model.
498
After calculating correlation (weighted for eADAGE and unweighted for corADAGE), we
converted the correlation to distance by calculating (1- correlation)/2. This provided a
30000*30000 distance matrix storing distances between every two nodes. We clustered this
distance matrix using the Partition Around Medoids (PAM) clustering algorithm (Park & Jun,
2009).We implemented clustering in R using the ConsensusClusterPlus package (Wilkerson &
Hayes, 2010) from Bioconductor with the ppam function from Sprint package to perform
parallel PAM (Piotrowski *et al*, 2011). We set the number of clusters to match the individual
ADAGE model (e.g. 300) allowing for direct comparison between the eADAGE and ADAGE
methods.
508
Clustering assigned each node to a cluster ranging from 1 to 300. We combined nodes assigned
to the same cluster by calculating the average of their weight vectors. These 300 averaged
vectors formed the weight matrix of the eADAGE model. Because the ensemble model is built
from the weight matrices of individual models, it does not have the parameters that form the
bias vectors. We built 10 eADAGE and 10 corADAGE models from 1000 ADAGE models with
each ensemble model built upon 100 different individual models. The individual eADAGE model
used for biological analysis in this work was constructed with random seed 123, which was
arbitrarily chosen before model construction and evaluation.
517
PCA and ICA model construction
We constructed PCA and ICA models and defined each model's weight matrix following the
same procedures in (Tan *et al*, 2016b). To compare with the 300-node eADAGE, we generated
models of matching size (300 components). For ICA, we evaluated 10 replicates. PCA provides a
single model. PCA and ICA models were evaluated through the KEGG pathway enrichment
analysis described above.
524
Media annotation of the *P. aeruginosa* compendium

A team of *P. aeruginosa* biologists annotated the media for all samples in the compendium by
referring to information associated with each sample in the ArrayExpress (Rustici *et al*, 2013)
and/or GEO (Edgar, 2002) databases and along with the original publication, if reported. Each
sample was annotated by two curators separately. Conflicting annotations, if they occurred,
were resolved by a third curator. The media annotation for all samples in the compendium
were provided in Table EV1.
532

Activity calculation for a gene signature

534   We calculated a signature's activity for a specific sample as $A = W \cdot E / N$, in which W is the
535   weight vector of genes in that signature, E is a vector of genes' expression values after zero-one
536   normalization in that sample, and N is the number of genes. It can be viewed as an averaged
537   weighted sum of genes' expression levels. We normalized a signature's activity by the number
538   of genes (N) in that signature, because different signatures have different number of genes.
539

Identification of signatures activated across media

We calculated an activation score to identify gene signatures with dramatically elevated or
reduced activity in a specific medium. We grouped samples by their medium annotation. For
each gene signature and medium combination, we calculated the difference between the mean
activity of the signature for samples in that medium as well as the mean activity across the
remainder of samples in the compendium. We divided this absolute difference in the means by
the range of activity for all samples across the compendium. This score captures the proportion
by which the mean activity in a medium differs relative to the total difference across the
compendium. We termed this ratio the activation score.
549

To identify the most specifically active signatures for each medium, we constructed a table for
all pairs with an activation score greater than or equal to 0.4 (Table EV5). This was highly
stringent: it captured only the top 2.4% of the potential signature-medium pairs. To identify
pan-media signatures, we limited signatures to those that were active in multiple media
(greater or equal to 0.4) and averaged their activation scores. These signatures exhibit parallel
patterns for multiple media across multiple distinct experiments.
556

Definition of the PhoB regulon

558   A PhoB regulon for the PAO1 genome was adapted from the PhoB regulon of PA14 in (Bielecki
559   *et al*, 2015) in order to be comparable to ADAGE models. Of the 187 genes in the PA14 regulon,
560   160 were in the PAO1 reference genome (www.pseudomonas.com).
561

Strains and Media

Strains used were WT, Δ*phoB* (DH2633, O'Toole lab collection), Δ*phoR* (DH2516) and Δ*kinB*
(DH2517), all in the PA14 background. All strains were maintained on LB with 1.5% agar and
grown at 37 °C. For cross-media and phosphate concentration comparisons, BCIP assays (see
methods below) were conducted on different base media with 1.5% agar (Fisher): King's A
(pancreatic digest of gelatin (Difco) 20g/L; $MgCl_2$ 1.4g/L; K_2SO_4 10g/L; glycerol 10ml/L) (King *et al*,
1954), LB (Tryptone (Fisher) 10g/L; Yeast Extract (Fisher) 5g/L; NaCl 5g/L) (Bertani, 2004), MOPS
(morpholinepropanesulfonic acid 40mM; Glucose 20 ml/L; K_2SO_4 2.67mM; K_2HPO_2 0mM, 25mM

or 0.1 – 1 mM) (Neidhardt *et al*, 1974), NGM (Pancreatic digest of gelatin 2.5g/L; cholesterol
5mg/L; NaCl 3g/L; MgSO$_4$ 1mM; CaCl$_2$ 1mM; KCl 25mM; Potassium Phosphate buffer pH6 0 or
25 mM) (Zaborin *et al*, 2009), Peptone (Pancreatic digest of gelatin 10g/L; MgSO$_4$ 1.5g/L; K$_2$SO$_4$
10g/L) (Lundgren *et al*, 2013), Pseudomonas Isolation Agar (PIA, prepared as per instructions,
BioWorld).

BCIP assay
Various media were supplemented with 5-bromo-4-chloro-3-indolyl phosphate (BCIP) DMF
solution to a final concentration of 60 µg/mL. BCIP assay plates were inoculated with 5 µl of
overnight *P.a* culture in LB broth. Colonies were grown for 16 hours at 37 °C then matured at
room temperature until imaging. Images were collected 16 and 32 hours post inoculation.

Screen of a histidine kinase knock out collection
Molecular techniques to construct the histidine kinase (HK) knock out collection were carried
out as previously described (Ha *et al*, 2014). For each strain in the HK collection, a BCIP assay
was performed on PIA. Plates were struck with an overnight *P.a.* culture concentrated two-fold
by centrifugation. Plates were incubated at 37 °C 12-16 hours and matured at room
temperature for an additional 12-16 hours alkaline phosphatase activity was determined
qualitatively, based on blue color.

Acknowledgements
This work was supported in part by a grant from the Gordon and Betty Moore Foundation
(GBMF 4552) to CSG. This work was supported by National Institutes of Health (NIH) grant RO1-
AI091702 to DAH. MTL is an investigator of the Howard Hughes Medical Institute. This work
was supported by a pilot grant from the Cystic Fibrosis Foundation (STANTO15R0) to CSG and
DAH. The authors acknowledge Gregory Way and René Zelaya for helpful code review.

Author contributions
JT, DAH and CSG conceived and designed the research. JT, GD and KMC performed
computational analyses. GD, KAL and CEP performed molecular experiments. KC, BP and MTL
constructed and contributed the histidine kinase knock out collection. JT, GD, KMC, DAH and
CSG wrote the manuscript, and KAL, CEP, KMC, KD, BP and MTL provided critical feedback.

Conflict of interest
The authors have no conflicts of interest to report.

Reference
Alter O, Brown PO & Botstein D (2000) Singular value decomposition for genome-wide
expression data processing and modeling. *Proc. Natl. Acad. Sci. U. S. A.* **97:** 10101–6
Bertani G (2004) Lysogeny at mid-twentieth century: P1, P2, and other experimental systems. *J.*
Bacteriol. **186:** 595–600
Bielecki P, Jensen V, Schulze W, Gödeke J, Strehmel J, Eckweiler D, Nicolai T, Bielecka A, Wille T,
Gerlach RG & Häussler S (2015) Cross talk between the response regulators PhoB and TctD

allows for the integration of diverse environmental signals in *Pseudomonas aeruginosa*.
Nucleic Acids Res. **43:** 6413–25

Blus-Kadosh I, Zilka A, Yerushalmi G & Banin E (2013) The effect of *pstS* and *phoB* on quorum
sensing and swarming motility in *Pseudomonas aeruginosa*. *PLoS One* **8:** e74444

Chen L, Xuan J, Wang C, Shih I-M, Wang Y, Zhang Z, Hoffman E, Clarke R, Devore J, Peck R,
Tusher V, Tibshirani R, Chu G, Storey J, Xiao W, Leek J, Tompkins R, Davis R, Conesa A,
Nueda M, et al (2008) Knowledge-guided multi-scale independent component analysis for
biomarker identification. *BMC Bioinformatics* **9:** 416

Damron FH, Barbier M, McKenney ES, Schurr MJ & Goldberg JB (2013) Genes required for and
effects of alginate overproduction induced by growth of *Pseudomonas aeruginosa* on
Pseudomonas isolation agar supplemented with ammonium metavanadate. *J. Bacteriol.*
195: 4020–36

Damron FH, Owings JP, Okkotsu Y, Varga JJ, Schurr JR, Goldberg JB, Schurr MJ & Yu HD (2012)
Analysis of the *Pseudomonas aeruginosa* regulon controlled by the sensor kinase KinB and
sigma factor RpoN. *J. Bacteriol.* **194:** 1317–30

Donato M, Xu Z, Tomoiaga A, Granneman JG, Mackenzie RG, Bao R, Than NG, Westfall PH,
Romero R & Draghici S (2013) Analysis and correction of crosstalk effects in pathway
analysis. *Genome Res.* **23:** 1885–93

Edgar R (2002) Gene Expression Omnibus: NCBI gene expression and hybridization array data
repository. *Nucleic Acids Res.* **30:** 207–210

Engreitz JM, Daigle BJ, Marshall JJ & Altman RB (2010) Independent component analysis: mining
microarray data for fundamental human gene expression modules. *J. Biomed. Inform.* **43:**
932–44

Fisher SL, Jiang W, Wanner BL & Walsh CT (1995) Cross-talk between the histidine protein
kinase VanS and the response regulator PhoB. Characterization and identification of a VanS
domain that inhibits activation of PhoB. *J. Biol. Chem.* **270:** 23143–9

Frigyesi A, Veerla S, Lindgren D, Höglund M, Quackenbush J, Jutten C, Herault J, Chiappetta P,
Roubaud M, Torrésani B, Hyvärinen A, Oja E, Liebermeister W, Lee S, Batzoglou S,
Martoglio A, Miskin J, Smith S, MacKay D, Saidi S, et al (2006) Independent component
analysis reveals new and biologically significant structures in microarray data. *BMC*
Bioinformatics **7:** 290

Gillis J & Pavlidis P (2013) Assessing identity, redundancy and confounds in Gene Ontology
annotations over time. *Bioinformatics* **29:** 476–82

Gong T, Xuan J, Wang C, Li H, Hoffman E, Clarke R & Wang Y (2007) Gene module identification
from microarray data using nonnegative independent component analysis. *Gene Regul.*
Syst. Bio. **1:** 349–63

Greene CS & Troyanskaya OG (2012) Accurate evaluation and analysis of functional genomics
data and methods. *Ann. N. Y. Acad. Sci.* **1260:** 95–100

Ha D-G, Richman ME & O'Toole GA (2014) Deletion mutant library for investigation of
functional outputs of cyclic diguanylate metabolism in *Pseudomonas aeruginosa* PA14.
Appl. Environ. Microbiol. **80:** 3384–93

Jiang Y, Oron TR, Clark WT, Bankapur AR, D'Andrea D, Lepore R, Funk CS, Kahanda I, Verspoor
KM, Ben-Hur A, Koo DCE, Penfold-Brown D, Shasha D, Youngs N, Bonneau R, Lin A,
Sahraeian SME, Martelli PL, Profiti G, Casadio R, et al (2016) An expanded evaluation of

protein function prediction methods shows an improvement in accuracy. *Genome Biol.* **17:** 184

Kanehisa M (2000) KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids Res.* **28:** 27–30

KING EO, WARD MK & RANEY DE (1954) Two simple media for the demonstration of pyocyanin and fluorescin. *J. Lab. Clin. Med.* **44:** 301–7

Lundgren BR, Thornton W, Dornan MH, Villegas-Peñaranda LR, Boddy CN & Nomura CT (2013) Gene PA2449 is essential for glycine metabolism and pyocyanin biosynthesis in *Pseudomonas aeruginosa* PAO1. *J. Bacteriol.* **195:** 2087–100

Lutter D, Langmann T, Ugocsai P, Moehle C, Seibold E, Splettstoesser WD, Gruber P, Lang EW & Schmitz G (2009) Analyzing time-dependent microarray data using independent component analysis derived expression modes from human macrophages infected with *F. tularensis holartica*. *J. Biomed. Inform.* **42:** 605–611

Ma S & Kosorok MR (2009) Identification of differential gene pathways with principal component analysis. *Bioinformatics* **25:** 882–9

Monti S, Tamayo P, Mesirov J & Golub T (2003) Consensus clustering: a resampling-based method for class discovery and visualization of gene expression microarray data. *Mach. Learn.* **52:** 91–118

Neidhardt FC, Bloch PL & Smith DF (1974) Culture medium for enterobacteria. *J. Bacteriol.* **119:** 736–47

Ninfa AJ, Ninfa EG, Lupas AN, Stock A, Magasanik B & Stock J (1988) Crosstalk between bacterial chemotaxis signal transduction proteins and regulators of transcription of the Ntr regulon: evidence that nitrogen assimilation and chemotaxis are controlled by a common phosphotransfer mechanism. *Proc. Natl. Acad. Sci. U. S. A.* **85:** 5492–6

Park H-S & Jun C-H (2009) A simple and fast algorithm for K-medoids clustering. *Expert Syst. Appl.* **36:** 3336–3341

Piotrowski M, Forster T, Dobrezelecki B, Sloan TM, Mitchell L, Ghazal P, Mewsissen M, Petrou S, Trew A & Hill J (2011) Optimisation and parallelisation of the partitioning around medoids function in R. In *2011 International Conference on High Performance Computing & Simulation* pp 707–713. IEEE

Podgornaia AI & Laub MT (2013) Determinants of specificity in two-component signal transduction. *Curr. Opin. Microbiol.* **16:** 156–62

Raychaudhuri S, Stuart JM & Altman RB (2000) Principal components analysis to summarize microarray experiments: application to sporulation time series. *Pac. Symp. Biocomput.***:** 455–66

Roden JC, King BW, Trout D, Mortazavi A, Wold BJ, Hart CE, Tavazoie S, Hughes J, Campbell M, Cho R, Church G, Tamayo P, Slonim D, Mesirov J, Zhu Q, Kitareewan S, Dmitrovsky E, Lander E, Golub T, Eisen M, et al (2006) Mining gene expression data by interpreting principal components. *BMC Bioinformatics* **7:** 194

Rustici G, Kolesnikov N, Brandizi M, Burdett T, Dylag M, Emam I, Farne A, Hastings E, Ison J, Keays M, Kurbatova N, Malone J, Mani R, Mupo A, Pedro Pereira R, Pilicheva E, Rung J, Sharma A, Tang YA, Ternent T, et al (2013) ArrayExpress update--trends in database growth and links to data analysis tools. *Nucleic Acids Res.* **41:** D987-90

Santos-Beneit F (2015) The Pho regulon: a huge regulatory network in bacteria. *Front. Microbiol.*

6: 402

Schnoes AM, Ream DC, Thorman AW, Babbitt PC & Friedberg I (2013) Biases in the
experimental annotations of protein function and their effect on our understanding of
protein function space. *PLoS Comput. Biol.* **9:** e1003063

Tan J, Doing G, Lewis KA, Price CE, Chen KM, Cady KC, Perchuk B, Laub MT, Hogan DA & Greene
CS (2016a) eADAGE-1.0.0rc1. *Zenodo*

Tan J, Hammond JH, Hogan DA & Greene CS (2016b) ADAGE-Based Integration of Publicly
Available *Pseudomonas aeruginosa* Gene Expression Data with Denoising Autoencoders
Illuminates Microbe-Host Interactions. *mSystems* **1:** e00025-15

Tan J, Ung M, Cheng C & Greene CS (2015) Unsupervised feature construction and knowledge
extraction from genome-wide assays of breast cancer with denoising autoencoders. *Pac.*
Symp. Biocomput. **20:** 132–43

Verhamme DT, Arents JC, Postma PW, Crielaard W & Hellingwerf KJ (2002) Investigation of in
vivo cross-talk between key two-component systems of *Escherichia coli*. *Microbiology* **148:**
69–78

Vincent P, Larochelle H, Bengio Y & Manzagol P-A (2008) Extracting and composing robust
features with denoising autoencoders. In *Proceedings of the 25th International Conference*
on Machine learning - ICML '08 pp 1096–1103. New York, New York, USA: ACM Press

Wilkerson MD & Hayes DN (2010) ConsensusClusterPlus: a class discovery tool with confidence
assessments and item tracking. *Bioinformatics* **26:** 1572–3

Yu B (2013) Stability. *Bernoulli* **19:** 1484–1500

Zaborin A, Romanowski K, Gerdes S, Holbrook C, Lepine F, Long J, Poroyko V, Diggle SP, Wilke A,
Righetti K, Morozova I, Babrowski T, Liu DC, Zaborina O & Alverdy JC (2009) Red death in
Caenorhabditis elegans caused by *Pseudomonas aeruginosa* PAO1. *Proc. Natl. Acad. Sci. U.*
S. A. **106:** 6327–32

Figure Legends

Figure1: Knowledge- and data-driven heuristics for ADAGE

A   Knowledge-driven model size heuristics on pathway coverage. As model size increases,
pathway coverage also increases at first and then levels out at size 300. The red line goes
through the median value at each model size.

B   Data-driven model size heuristics on reconstruction error. As model size increases, the
reconstruction error drops quickly at the beginning and levels out at size 300. The red line goes
through the median reconstruction errors at each model size.

C   Knowledge-driven sample size heuristics on pathway coverage. For 50-node models,
pathway coverage increases with sample size and peaks at 500 samples. 300-node models
cover more pathways than 50-node modes in general and maintain a slow growing trend of
pathway coverage at the maximum sample size.

D   Data-driven sample size heuristics on reconstruction error. In both 50- and 300-node
models, the reconstruction errors on the test set get closer to the reconstruction errors on the
train set as sample size increases.

Figure 2: The construction and performance of eADAGE.

A　　eADAGE construction workflow. 100 individual ADAGE models were built using the same
input dataset (step 1). Nodes from all models were extracted (step 2) and clustered based on
the similarities in their associated weight vectors (step 3). Nodes derived from different models
were rearranged by their clustering assignments (step 4). Weight vectors from nodes in the
same cluster were averaged and thus becoming the final weight vector of a newly constructed
node in an eADAGE model (step5).
B　　Pathway coverage comparison between individual ADAGE and ensemble ADAGE.
eADAGE models (n=10) covers significantly more pathways than both corADAGE (n=10) and
ADAGE (n=1000).
C　　The enrichment significance of three example pathways in different models. The three
pathways show different trends as model size increases in individual ADAGE, however, their
median significance levels in eADAGE are comparable or better than all individual models with
different sizes. The grey dotted line indicates FDR q-value of 0.05 in pathway enrichment.
D　　Comparison among PCA, ICA, and eADAGE in pathway coverage at different significance
levels. eADAGE outperforms PCA at all significance levels. eADAGE and ICA show similar
pathway coverage at the cutoff q-value = 0.05. However, ICA covers less pathways than
eADAGE as the significance cutoff becomes more stringent.
761
Figure 3: Node164pos is active in a NGM+<0.1phosphate, peptone, King's A, and PIA media
A　　Activity of Node164pos in each medium type. NGM+<0.1phosphate, peptone, and King's
A media have evident elevation in Node164pos's activity. PIA medium show a wide range in
Node164pos's activity. All other media have very low activities.
B　　Gene expression heatmaps of genes in Node164pos across samples in
NGM+<0.1phosphate, peptone, King's A, and PIA media. Heatmap color range is determined
by the Z-scored gene expression of all samples in the compendium (Figure EV6). These genes
are highly expressed in all samples grown on NGM + <0.1mM phosphate, peptone, King's A, and
half of samples on PIA, but not expressed in samples grown on NGM + 25mM phosphate.
771
Figure 4: PhoA activity, as seen by the colorimetric BCIP assay in various media
A　　PhoA activity, as seen by the blue-colored product of BCIP cleavage, is dependent on
low phosphate concentrations, *phoB, phoR* and, in NGM, *kinB*.
B　　PhoA is active in King's A, Peptone and PIA and is dependent on *phoB* and *phoR* on
King's A and peptone but dependent on *kinB* as well on PIA at 16 hours.
C　　PhoA is active in King's A, Peptone and PIA and is dependent on *phoB,* but no longer
phoR, while still dependent on *kinB* on PIA after 32 hours.
D　　PhoA activity is dependent on phosphate concentrations < 0.6 mM, *phoB, phoR* and *kinB*
as well at 0.5 mM phosphate in MOPS. Concentration 0.2 mM (not shown) mimics 0.1mM and
concentrations 0.7mM – 0.9mM (not shown) mimic 1.0 mM.
782

Expanded View Figure Legends

784　Figure EV1: The relationship between model size and the number of KEGG pathways a node
785　significantly associated with. Pathways associated with positive and negative signatures of a
786　node were added together. When model is small, one node needs to account for multiple KEGG

787   pathways. As model size grows, more nodes become available and pathways also tend to
788   spread into different nodes.
789
790   Figure EV2: Pathway coverage comparison between individual ADAGE and ensemble ADAGE
791   after correcting pathway crosstalk effects. eADAGE models (n=10) covers significantly more
792   pathways than both corADAGE (n=10) and ADAGE (n=1000).
793
794   Figure EV3: The association significance of each KEGG pathway in the 300-node eADAGE models
795   (n = 10), 300-node corADAGE models (n=10) and ADAGE models with different number of
796   nodes (n = 100 for each model size).
797
798   Figure EV4: The coverage rate of each KEGG pathway in 300-node ADAGE models (n=1000) and
799   300-node eADAGE (n=10) models.
800
801   Figure EV5: The distribution of KEGG pathway coverage rates in 300-node ADAGE models
802   (n=1000) and 300-node eADAGE models (n=10). eADAGE shows a higher density in distribution
803   on the high coverage end.
804
805   Figure EV6: Z-scored gene expression heatmap of genes in signature Node164pos across all
806   samples in the compendium.
807

808   **Expanded View Datasets and Tables**
809   Dataset EV1: *Pseudomonas aeruginosa* gene expression compendium. Gene expression values
810   in each sample have been background corrected and quantile normalized.
811
812   Table EV1: Medium annotation for each sample in the compendium. Some medium names are
813   abbreviated and their actual ingredients are provided in a separate sheet in the excel file.
814
815   Table EV2: Weight matrix of the eADAGE model highlighted in this paper. The weight matrix
816   defines how much each gene contributes to each node.
817
818   Table EV3: Genes in each signature in the eADAGE model. Filtering the weight matrix provides
819   these gene sets. Specifically, genes in a signature are those that were more than 2.5 standard
820   deviations from the mean on the positive side, or those that were more than 2.5 standard
821   deviations from the mean on the negative side.
822
823   Table EV4: The activity of each signature for each sample in the compendium. A signature has
824   high activity when its gene weights are required to reconstruct the gene expression.
825
826   Table EV5: A complete list of signatures activated in one medium with activation scores higher
827   than or equal to 0.4.
828

829   Table EV6: Top 5 eADAGE signatures that were active in a group of media. Each signature was
830   annotated by its percentage of uncharacterized genes, its associated KEGG pathways, and a
831   manual inspection of genes in the signature.
832
833   Table EV7: A complete list of signatures activated in a group of media.
834
835   Table EV8: Library of histidine kinase deletion mutants in PA14 used for a BCIP screen in PIA. All
836   strains in the collection of PA14 histidine kinase mutants described by the number from the
837   Hogan lab collection (DH number), the locus of the deleted gene, gene name (if available) and
838   position in the storage plate.

839 Figure 1

Figure 2

843 Figure 3

Figure 4