Genome-Wide Association Study of Ionomic Traits on Diverse Soybean Populations from Germplasm Collections

Greg Ziegler[1,2], Randall Nelson[3], Stephanie Granada[2], Hari B. Krishnan[4,5], Jason D. Gillman[4,5] and Ivan Baxter[1,2]

[1]USDA-ARS Plant Genetics Research Unit, St. Louis, Missouri
[2]Donald Danforth Plant Science Center, St. Louis, Missouri
[3] USDA-ARS, Soybean/Maize Germplasm, Pathology and Genetics Research Unit and Department of Crop Sciences, University of Illinois, Urbana, Illinois
[4]USDA-ARS Plant Genetics Research Unit, Columbia, Missouri
[5]Plant Science Division, University of Missouri, Columbia, Missouri

Abstract

The elemental content of a soybean seed is a determined by both genetic and environmental factors and is an important component of its nutritional value. The elemental content is chemically stable, making the samples stored in germplasm repositories an intriguing source of experimental material. To test the efficacy of using samples from germplasm banks for gene discovery, we analyzed the elemental profile of seeds from 1653 lines in the USDA Soybean Germplasm Collection. We observed large differences in the elemental profiles based on where the lines were grown, which lead us to break up the genetic analysis into multiple small experiments. Despite these challenges, we were able to identify candidate SNPs controlling elemental accumulation as well as lines with extreme elemental accumulation phenotypes. Our results suggest that elemental analysis of germplasm samples can identify SNPs in linkage disequilibrium to genes, which can be leveraged to assist in crop improvement efforts.

Introduction

One of the biggest challenges facing agricultural research today is finding ways to improve crop yield and nutrition while farming in increasingly erratic climates and on more marginal lands. Throughout modern agriculture, crops have been bred for maximal yield under optimal environmental conditions. Farming marginal soils with insufficient fertilization or irrigation leads to dramatic decreases in crop yield. In addition, plants grown on marginal soils may exhibit a reduced nutritional profile, which is an important consideration for staple crops. To properly address these issues, we need to develop a more complete understanding of the genetic mechanisms underlying a plant's response to various environmental stresses (Baxter and Dilkes 2012).

An important aspect underlying a plant's response to environmental stresses is its ability to regulate mineral nutrients. Apart from carbon and oxygen, a plant relies entirely on the bioavailable nutrients in the soil in which it is growing for survival. Soil nutrient bioavailability can

45 vary drastically, not just as a result of soil composition, but also as a side effect of drought and
46 flood conditions, changes in soil pH, and changes in the soil microbiome (FAO 1996).
47 Understanding the uptake, regulation, transport, and storage of mineral nutrients under a variety
48 of environmental conditions is essential to deciphering the complex relationship between a plant
49 and its environment.
50
Single-seed ionomic profiles have proven both highly heritable and susceptible to environmental
perturbations in maize (Baxter *et al.* 2014). This makes the study of the seed ionome a powerful
tool for matching a plant's genetic characteristics with its response to environmental
perturbations. Both environmental and genetic properties can effect multiple elements in
combination, resulting in genetic loci that might control different elements in different
environments (Baxter 2015; Asaro *et al.* 2016). Additionally, once collected, apart from the
possibility of external contamination, the elemental content of a seed sample is fixed. Tissue for
ionomic analysis doesn't need to be specially stored or quickly analyzed after collection.
Conveniently, this allows for the ionomic analysis of excess tissue collected for other purposes,
without the necessity of a separate field experiment. Here we demonstrate the utility of
leveraging existing germplasm by performing a genome-wide association study on ionomic traits
in seed tissue measured from diverse soybean lines selected from the USDA Soybean
Germplasm Collection.
64
65 **Results**
66
67 **Experimental Design**
68
The mission of the USDA-ARS National Plant Germplasm System (NPGS) is "to acquire,
evaluate, preserve and provide a national collection of genetic resources to secure the
biological diversity that underpins a sustainable U.S. agricultural economy." Some of these
collections are the target for high-density genotyping projects making them ideal populations for
genome-wide association studies. However, the prohibitive cost of controlled field trials to
measure novel phenotypes can limit their utility for genetics research. In this experiment, we
used existing germplasm to find novel genotype-phenotype associations without the expensive
overhead of independent field trials. Although this experiment is limited by the inability to grow
plants in a common environment, the high heritability of ionomics traits (Baxter *et al.* 2014), as
well as the stability of the ionome in stored tissue (Baxter *et al.* 2014), makes ionomic
phenotyping an ideal test case for mining germplasm resources. To test the power of ionomics
to find genetic factors underpinning elemental accumulation, we analyzed seeds from 1653
soybean [Glycine max (L.) Merr.] lines representing the diversity found in the USDA Soybean
Germplasm Collection stored at Urbana, IL.
83
A core collection of 1685 accessions of the USDA Soybean Germplasm Collection represents a
substantial amount of the genetic diversity in the entire collection. The core collection contains
approximately 10% of the total number of introduced soybean accessions. The 1653 soybean
lines used in this study comprised all of the 1685 accessions available when the research was
started. For accessions in maturity groups 000 through VIII for which field evaluation data were

available the core was selected using origin, qualitative and quantitative data. Accessions were
divided in groups based on origin and then further subdivided based on maturity group, which
classifies soybean accessions based on photoperiod and temperature response. A total of 81
strata were established. A multivariate proportional sampling strategy within each stratum was
determined to be the optimal procedure for identifying a sample of accessions that best
represents the diversity of the total collection. Field evaluation data were not available for
accessions in maturity groups IX and X, but because these accessions are adapted to sub-
tropical and tropical conditions and are likely to have unique genetic diversity, a sample of 10%
of these accessions was added to the core collection based on multivariate analysis of the
qualitative data. A full explanation of the development of the core collection can be found in
Oliveira et al. (2010). The seeds available in the NPGS for this core collection come from grow-
outs that span 12 years at three locations (Urbana, IL, Stoneville, MS, and Upala, Costa Rica)
(Table 1). The selection of which lines to grow for line maintenances in a given year is
independent of the strata used to select the core collection, making each growout year an
independent experiment to look for loci controlling elemental accumulation. Additonally, analysis
of the first two principal components from the SNP dataset shows no apparent bias between
genetic architecture and growout (Supplemental Figure 1).
106
107   **Table 1. Number of lines and markers in each GWAS dataset. There is no overlap between lines in the**
108   **datasets. Markers are the number of segregating SNPs in each dataset, filtered for minor allele frequency >**
109   **0.05.**

Location	Growout Year	Lines	GWAS Markers
Stoneville	1999	104	33962
Stoneville	2004	121	34571
Stoneville	2006	59	35192
Urbana	2000	109	36432
Urbana	2001	69	36032
Urbana	2002	94	36151
Urbana	2003	147	35783
Urbana	2004	89	35490
Urbana	2005	87	35559
Urbana	2006	143	36065
Urbana	2007	98	36091
Urbana	2008	58	35432
Urbana	2009	102	36489
Costa Rica	9 years combined	111	31479

110
111
Phenotypes
Using the elemental analysis pipeline described in Ziegler et al. (2013, see methods), we
analyzed ~6 seeds from each line, measuring the levels of 20 elements in each seed
(Supplemental Table 1). While 1653 lines were analyzed in total, 262 of these lines were from
grow-outs containing fewer than 50 lines in the dataset. We excluded these lines from further

analysis and all following analysis is based on the remaining 1391 lines (elemental profiles for
excluded lines are included in the Supplemental Table 1). We performed an ANOVA
significance test to assess whether there are significant environmental effects on the phenotypic
data gathered from lines grown in separate locations and in separate years at the same
location. Although a distinct set of lines were grown in each grow-out, lines were assigned to a
grow-out without regard to population structure. As a result, we would expect, in the absence of
environmental effects, phenotypic measurements to be similar. The ANOVA test indicates a
significant location effect, and for Stoneville and Urbana, significant effects for growth year, for
most elements measured (p<0.01 with Bonferroni correction, Table 2). This effect can also be
seen in the phenotypic distribution (before transformation) for many of the traits (Figure 1 and
Supplemental Figure 2). These results clearly demonstrate that most of the year growouts were
unique environments, supporting their analysis as individual experiments. The lack of significant
differences by year for many elements in Costa Rica (13 out of 21) may be indicative of a lack of
statistical power due to the small number of lines grown per year. Because there were not
enough lines in any one grow-out from Costa Rica for a GWAS analysis, the only way we were
able to analyze the Costa Rica data was by combining data across all 10 years.

133
134 **Table 2. Analysis of grow out location and year effect on elemental accumulation. The *p*-value for each**
135 **element from an ANOVA of a linear model with Location or Location x Year interaction. The significance**
136 **cutoff was set at *p* < 0.01 with Bonferroni correction. NS=Not Significant**

Element	Location	Costa Rica x Year	Stoneville x Year	Urbana x Year
Seed Weight	NS	NS	6.87E-07	0.0001776
B	0.0001174	NS	1.24E-07	NS
Na	3.06E-307	NS	NS	NS
Mg	0.0003425	5.24E-08	7.19E-09	2.19E-29
Al	9.17E-31	8.70E-13	2.62E-11	3.56E-36
P	5.72E-27	1.26E-05	NS	3.29E-16
S	6.49E-34	NS	3.58E-10	6.23E-35
K	2.37E-24	1.16E-05	1.46E-07	2.12E-06
Ca	1.63E-19	NS	6.78E-13	1.17E-26
Mn	9.80E-45	0.0003116	3.03E-15	1.53E-17
Fe	7.12E-29	NS	8.44E-09	2.36E-34
Co	3.42E-148	NS	1.10E-19	3.65E-12
Ni	3.04E-173	5.90E-13	5.75E-06	2.37E-33
Cu	1.33E-243	NS	1.05E-14	1.40E-29
Zn	1.34E-145	NS	6.38E-08	9.29E-30
As	1.66E-57	NS	5.50E-12	NS
Se	0	0.0001141	1.13E-16	2.23E-14
Rb	0	4.39E-08	6.75E-44	2.17E-15
Sr	0	NS	7.59E-06	3.34E-18
Mo	0	NS	3.68E-40	6.66E-44
Cd	3.25E-45	NS	5.48E-26	3.79E-07

139
140 **Figure 1. Molybdenum accumulation in single soybean seeds (mg/kg) across experimental grow-outs.**

Comparison of elemental concentrations of replicate seeds from the same line in each grow-out does indicate the presence of a genotypic effect on elemental concentrations. Concentrations in seeds from the same line were usually more similar to each other than they were to the population as a whole (Figure 2 and Supplemental Figure 3).

147
148 **Figure 2. Distribution of Cadmium phenotype (linear model residuals, see Methods) in lines from a single**
149 **growout: Stoneville, MS, 1999. Lines are ordered by median of between 2 and 8 seed replicates.**

151
The Box-Cox procedure (Box and Cox 1964) was used to estimate appropriate transformation
functions for the phenotype data to meet the assumptions of GWAS for normally distributed
dependent variables. The Box-Cox algorithm suggested that 138 of the 294 traits (14
environments x 21 phenotypes) needed no transformation and an additional 151 needed only
minor transformations to control for the long-tail distributions often seen in concentration data
(inverse, inverse square root, log, or square root) (Supplemental Table 2). Because most traits
appear to only need minor transformations, for uniformity and ease of interpretation, all of the
traits in which a transformation was recommended were transformed using a log transformation.
160
161 **Population Structure**
162
The first two principal components obtained using the 36,340 polymorphic SNPs from the entire
1391 lines in the dataset explained 15% of the total SNP variance and the first 10 principal
components explained 28% of the total variance. Variance explained by each PC drops rapidly
after the first 10 PCs with 50% variance not reached until PC76. The first two principal
components separate the population into groups roughly corresponding to each lines country of
origin, with South Korean and Japanese accessions forming distinct clades while Chinese,
Russian and other accessions form a much less cohesive block (Figure 3).

170
171 **Figure 3. Principal component analysis of the genotypes of 1391 soybean lines. Colored by country of origin:**
172 **China (532), Japan (267), South Korea (200), Russia (61), Other or unknown country of origin (331).**

MLMM GWAS

Using the SoySNP50k chip data (Song *et al.* 2013), we performed a GWAS study using a multi-locus mixed model (MLMM) to identify associated loci for each of 21 phenotypes (20 elements, seed weight) in 13 distinct grow-outs of diverse soybean lines and the Costa Rica dataset of grow-outs pooled across years (Table 1). The MLMM procedure starts with an EMMAX scan of all markers and then iteratively adds the markers with the highest association to the model and rescans. The MLMM procedure returns a list of cofactors that together describe the total estimated narrow-sense heritability of a given trait (which we will refer to as the all cofactor model). By definition, MLMM will create a model containing at least one cofactor for each trait. Of the models generated, 84 models met the stopping criteria after only one SNP was added to the model. The average model contained 11 SNPs, with no traits reaching the maximum 40 SNP model (e.g. not converging on a model describing all of the phenotypic variance). The largest model contained 29 SNPs, for iron in the 2009 Urbana grow-out. The 294 GWAS tests returned 1756 unique SNPs. While these most complex models likely contain factors that account for phenotypic variance merely by chance (e.g., false positives), many of these cofactors are likely real.

A simpler model, which includes only a subset of the total cofactors, can be selected using a model selection parameter (Segura *et al.* 2012). Segura et al. proposed two model selection criteria: the extended Bayesian information criterion (EBIC) and the multiple-Bonferroni criterion (mBonf) (Segura *et al.* 2012). Although both criteria produced generally similar results, we found the EBIC criteria to be less stringent than mBonf. Due to the relatively small sample size in many of our grow-outs, we have chosen the more inclusive EBIC criteria in an attempt to include more moderate effect loci in our model at the cost of a higher false positive rate. QQ-plots for both the null model, containing no cofactors, and the optimal EBIC model were generated to assess whether there were uncontrolled confounding effects in our model arising from cryptic relatedness and population structure. While there was some inflation of p-values in the null model, the MLMM procedure of iteratively including large-effect loci into the model successfully controls for this p-value inflation and the distribution of p-values in the EBIC models closely follows the expected null distribution except for the significantly associated loci (Figure 4 and Supplemental Figure 4).

207
Figure 4. Quantile-quantile plot of the observed *p*-values against expected *p*-values from the GWAS analysis
for sulfur accumulation. The MLMM algorithm includes cofactors that reduce inflation of *p*-values (green
line). The model without cofactors indicates presence of *p*-value inflation (blue line). The expected
distribution of *p*-values under the null hypothesis (red line).

The EBIC model selection method returned the MLMM model containing no cofactors for about
half of the GWAS tests (164/294). The remaining 130 tests returned a total of 573 unique SNPs.
When looking at the combined set of SNPs returned across all grow-outs, of the 21 phenotypes
tested, at least one SNP was returned for each trait, with seed weight returning the most (96)
and boron returning the least (6). Table 3 contains information about the number of cofactors
returned in each model (EBIC and all) for each trait and Supplemental Table 3 contains the
complete list of SNPs returned.
220
Overall, despite a large number of tests for association (294), a relatively small number of SNPs
were identified. Given the ability of the multi cofactor model to reduce the levels of spurious
false positives, a large number of even the full model SNPS are likely to be real. However, given
the large number of independent growouts and the partial independence of the elemental traits,
we are able to apply more stringent criteria confidence in associations. Below, we list several
sets of SNPs associated with elemental traits, ordered from 'most confident' to 'lower
confidence'.   Since the likelihood of the same false associations being found more than once
for the same trait in separate grow-outs with independent sets of lines is small, we looked for
SNPs returned in multiple scans, which are likely to be real. Across these 130 experiments, 10
SNPs were returned more than once. Of these 10 SNPs, the exact same SNP was found for the
same element in a different grow-out two times (ss715604985 and ss715605104, both for
cadmium), different elements in the same grow-out once (ss715608340 for Ca and Sr), and
different elements in different growouts 7 times (Table 4). The same element/multiple location
and multiple element/same location SNPs constitute our highest confidence set for SNPs
affecting the ionome, but likely greatly underestimate the useful information in the dataset.
236

Table 3. Number of SNP cofactors returned by each GWAS experiment. Each cell contains the number of cofactors in the EBIC selected model and the all cofactor model, respectively. See methods for criteria for inclusion of a SNP in the EBIC or all cofactor model.

Growout/ Element	Al	As	B	Ca	Cd	Co	Cu	Fe	K	Mg	Mn	Mo	Na	Ni	P	Rb	S	Seed Weight	Se	Sr	Zn	Total
00U	1/1	0/1	3/7	4/10	12/13	0/10	0/3	0/14	0/3	18/19	8/10	1/4	0/1	0/13	0/12	0/3	0/13	2/16	0/10	2/10	4/20	55/193
01U	8/8	1/1	1/1	1/8	1/1	2/4	0/2	0/7	2/6	1/1	3/5	0/8	1/1	0/1	0/1	0/1	0/1	7/8	17/18	1/4	0/1	46/88
02U	0/2	0/11	0/1	1/4	10/13	0/14	0/4	0/3	0/7	0/1	1/2	0/8	0/1	2/11	5/10	2/3	0/9	14/16	1/3	0/14	0/9	36/146
03U	2/3	0/2	0/2	0/1	3/19	2/7	0/4	0/8	0/11	0/12	1/3	1/11	0/2	0/6	3/7	0/1	0/8	26/26	3/6	0/11	0/7	41/157
04S	1/9	0/1	0/4	2/6	3/3	0/3	0/1	0/6	3/5	0/14	0/1	0/1	0/4	0/3	1/11	1/1	0/4	1/24	0/11	4/12	0/8	16/132
04U	0/1	0/1	0/3	5/5	1/1	0/2	0/1	1/7	0/3	0/1	1/1	0/1	0/2	1/2	0/1	0/1	2/6	0/15	1/2	0/7	0/1	12/64
05U	0/10	0/1	1/1	2/4	3/6	3/6	0/2	0/23	0/4	0/5	2/5	0/1	0/1	0/1	1/1	0/1	2/13	17/18	14/16	1/1	0/2	46/122
06S	0/4	8/8	0/5	0/1	0/1	0/2	0/1	0/5	2/10	1/1	0/1	0/3	0/1	1/5	16/17	0/8	0/2	3/4	15/15	0/5	5/6	51/105
06U	0/1	0/2	0/1	1/7	1/15	0/1	1/10	5/13	3/10	0/9	0/6	0/3	0/1	1/11	0/1	0/1	0/10	3/12	1/14	0/11	0/1	16/140
07U	0/1	0/1	1/2	1/1	2/5	1/2	1/1	0/1	3/3	0/9	1/3	1/2	0/2	2/3	0/3	1/4	0/1	1/10	1/4	0/3	0/3	16/64
08U	1/2	2/3	0/1	14/15	1/4	20/20	8/8	9/10	0/1	12/12	0/1	0/1	0/1	0/1	9/11	2/3	0/1	5/7	1/2	3/4	0/1	87/109
09U	1/1	0/1	0/1	19/20	0/10	0/14	0/14	29/29	1/1	0/2	1/2	22/22	18/18	1/1	1/1	0/21	19/19	17/18	0/1	0/10	0/1	129/207
99S	2/2	0/5	0/1	1/11	1/12	1/13	0/10	0/2	0/1	1/6	0/15	1/1	0/4	0/7	0/1	1/11	0/4	0/15	0/17	0/1	0/20	8/159
CR	0/11	0/1	0/3	0/8	4/7	0/11	0/1	2/3	7/8	3/11	1/7	7/9	0/3	0/4	0/9	0/8	0/9	0/12	0/1	2/13	0/12	26/151
Total	16/56	11/39	6/33	51/101	42/110	29/109	10/62	46/131	21/73	36/103	19/62	33/75	19/42	8/69	36/86	7/67	23/100	96/201	54/120	13/106	9/92	585/1837

Because each grow-out contains an independent set of lines, the set of SNPs tested differs
between grow-outs depending upon the SNP minor allele frequency in each dataset.
Additionally, common SNPs between growouts will still differ in allele frequency, which could
result in neighboring SNPs, still in LD with the causal variant, being returned for different GWAS
experiments. Therefore, looking for only exact overlaps between datasets may be overly
restrictive. Soybean has been estimated to have a linkage disequilibrium (LD) decay distance of
between 360Kbp in euchromatic regions and 9.6Mbp in heterochromatic regions (Hwang et al.,
2014). To better search for overlaps between our datasets while also taking into account the
large variability in LD range across the soybean genome, we grouped all of the SNPs returned
across experiments by whether they are in LD with one another. Although many factors affect
the ability to detect an association between a QTL and the actual causative loci, the minimal r^2
for detection between the loci is generally estimated to be between 0.2 and 0.33 (Ardlie *et al.*
2002; Qanbari *et al.* 2010; Wallace *et al.* 2014) with a value of 0.2 previously being used to
define LD range in the soybean genome (Hwang *et al.* 2014). Therefore, we defined an overlap
between SNPs as whether a pair of SNPs has an $r^2 > 0.2$. When this approach was applied to
the all cofactors model, the same locus was returned for the same phenotype in different grow-
outs 18 times, a different phenotype in the same grow-out 44 times and different phenotypes in
different growouts 237 times (Supplemental Table 4). Often a SNP returned as significant in the
EBIC model for one growout, will have a corresponding SNP in the all cofactor model of another
growout, indicating that the signal is there in other populations, but at too weak a level to meet
strict significance thresholds.
262
Another line of evidence that the SNPs identified are real is the co-location with candidate
genes. Due to the large regions of linkage disequilibrium in the soybean genome, each of the
30,000 SNPs in our experiment is linked to dozens to hundreds of genes. Many plant
processes, including root structure/function, water relations, and inter, intra and extra-cellular
structures, can alter the elemental accumulation (Baxter *et al.* 2009; Tian *et al.* 2010; Chao *et al.*
2011, 2013; Barberon 2017). Each SNP is therefore likely to be associated with several
plausible candidate genes. We looked under the SNPs of our overlap sets for strong
candidates- those with orthologs associated directly with elemental phenotypes. Table 5
contains a list of SNPs found on or near candidate or already characterized genes. Many of the
candidates are under SNPs associated with individual elements to which they or their orthologs
were previously linked, or with chemically related elements (i.e Mn, Co, Cd with Fe or Se with
S). The presence of these strong candidates under the detected SNPS supports the evidence
from overlap that they are real associations.
276
277 **Table 4. SNPs returned in the EBIC selected model in two or more grow-outs.**

Chromosome	Base Pair	Environment	Trait	logP	Model	Overlap Type
9	4612586	99S	Cd	10.06	EBIC	Same Element, Different Location
9	4612586	04U	Cd	5.39	EBIC	Same Element, Different Location
9	4991159	00U	Cd	18.68	EBIC	Same Element, Different Location
9	4991159	02U	Cd	18.95	EBIC	Same Element, Different Location
9	4991159	03U	Cd	11.88	EBIC	Same Element, Different Location

9	4991159	06U	Cd	6.77	EBIC	Same Element, Different Location
10	5863544	04S	Ca	6.20	EBIC	Different Element, Same Location
10	5863544	04S	Sr	7.68	EBIC	Different Element, Same Location
2	46468030	03U	Seed Weight	11.73	EBIC	Different Element, Different Location
2	46468030	05U	Se	29.18	EBIC	Different Element, Different Location
5	41315343	06S	Mg	4.82	EBIC	Different Element, Different Location
5	41315343	09U	Mo	4.58	EBIC	Different Element, Different Location
10	5179735	05U	S	5.73	EBIC	Different Element, Different Location
10	5179735	06S	Ni	7.36	EBIC	Different Element, Different Location
13	19554349	07U	Ni	6.66	EBIC	Different Element, Different Location
13	19554349	09U	Ca	18.06	EBIC	Different Element, Different Location
13	22047323	02U	Cd	14.82	EBIC	Different Element, Different Location
13	22047323	06S	K	5.59	EBIC	Different Element, Different Location
13	26504428	00U	Cd	6.30	EBIC	Different Element, Different Location
13	26504428	03U	Seed Weight	10.48	EBIC	Different Element, Different Location
19	84371	08U	Cu	16.51	EBIC	Different Element, Different Location
19	84371	09U	Fe	51.76	EBIC	Different Element, Different Location

278 **Table 5. Returned SNPs overlapping candidate or already characterized genes. Bold font indicates lines returned in the more conservative EBIC model**
279 **for at least one growout. SNP basepairs are mapped to soybean reference genome build Glyma1.1.**

280 Chromosome	Base Pair (of most significant SNP)	Environment(s)	Trait(s)	-logP (Of most significant SNP)	Candidate Gene
9	**4991159**	00U; 02U; 03U; 06U	Cd	18.95	HMA13; Glyma.09g055600 (Benitez et al., 2012); (Fang et al., 2016)
2	43023030	99S;CR	Cd	20.67	Glyma.02g215700 is similar to At2-MMP which is induced during cadmium stress to leaves (Golldack et al., 2002)
3	**40883820**	02U; 99S	Se	21.15	NRAMP metal transporter (Glyma.03g181400); Aluminum Sensitive 3 (ALS3; Glyma.03g175800)
5	33737561	CR; 09U	Ca	36.24	Multidrug resistance-associated protein 3 (MRP3, Glyma.05g145000); AtMRP5 implicated in Calcium homeostasis in Arabidopsis (Gaillard et al., 2008)
14	**47003645**	06S; 03U	Co	17.91	ZIP metal ion transporter (Glyma.14g196200); Overlaps with a Zn and Rubidium (in all cofactor)
15	**410656**	04S; 07U	Mn	7.11	CAX2 (Glyma.15g001600), implicated in Mn transport (Shigaki et al., 2002); NRAMP6 (Glyma.15g003500), Mn transport; MGT2 (Glyma.15g002700) and MGT4 (Glyma.15g005200), magnesium transport
2	**5555909**	07U	Fe; Zn; P; Cu	6.91	ATOX1 (Glyma.02g068700), Copper transport
1	**54551283**	01U; CR; 00U; 04U	Al; Rb; Mo; Co; K	7.64	ALMT (Glyma.01g223300), Aluminum activated malate transport, malate is a chelator for aluminum and critical in detoxification
2	44460357	09U; 02U	Co; Ca	10.96	Heavy metal transport/detoxification (Glyma.02g222600, Glyma.02g222700);Potassium transporter 1 (Glyma.02g228500); Phosphate transporter 4;3 (Glyma.02g224200)
3	**5165511**	09U; 06U	Fe; Mn	36.05	YSL6 (Glyma.03g040200); FPN1 ferroportin (Glyma.03g042500)
7	**5480577**	06S; 06U	As; Ni	22.46	Heavy metal transport/detoxification (Glyma.07g065800); NRAMP2 (Glyma.07g058900)
11	**17367460**	04U; 06U	Fe; Se	21.13	ABC Transporter (Glyma.11g194700, Glyma.11g196100)
19	**84371**	08U; 09U	Cu; Fe	51.76	ATOX1 (Glyma.19g001000), Copper transport
3	**5455217**	00U; 04U	Mg; Co	7.45	iron regulated 1 (Glyma.03g042500); iron regulated 2 (Glyma.03g042400); YSL6 (Glyma.03g040200)
15	**1222084**	05U	Se	29.64	Sulphate Transporter (Glyma.15g014000) (El Kassis et al., 2007; Cabannes et al., 2011); Sulfite Transporter (Glyma.15g015600)
9	**4799335**	06S	K	4.31	Potassium Transporter (Glyma.09g052700)
7	**5900018**	06U	Fe	5.07	Overlap with IDC for FRO2 (Mamidi et al. 2014); Glyma.07g067700; Also Glyma.07g065800 a heavy metal detox
9	**4518093**	09U	Mo	17.96	Molybdenum Cofactor sulfurase (Glyma.09g050100)
9	**3807440**	09U	S	31.98	Glyma.09g045200 Heavy Metal Transport; Close to all cofactor selenium
5	8074553	00U; 06S	Fe	7.06	Stabilizer of iron transporter (AGO10, PNH, ZLL; Glyma.05g011300), in IDC datset (Mamidi et al. 2014)
3	45338714	03U	Fe	8.30	NAS3; Glyma.03g231200; Overlaps IDC (Mamidi et al. 2014)

Verification of High and Low Sulfur and Phosphorus accumulating lines

To test whether the elemental accumulation of ionomic traits in the lines in our panel are intrinsic to the genetics of the lines or an artifact of the environmental and field conditions, we performed two experiments in which we selected the highest and lowest accumulating lines for sulfur and phosphorus and regrew the seeds in controlled field and greenhouse conditions. Eight lines, four with a high phosphorus phenotype and four with a low phosphorus phenotype were selected for regrowth in a field in Columbia, MO. Three of the four high phosphorus lines exhibited a high phosphorus phenotype in the regrow experiment, while the low phosphorus lines had phenotypes closer to the control line level (Figure 5 and Table 6). Broad-sense heritability for phosphorus between the GRIN growout concentrations and this experiment was 0.65 (Supplemental Table 5).

Table 6. Accessions chosen for validation of phosphorus accumulation. High and low phosphorus accumulating lines were chosen to regrow to test the reproducibility of ionomic traits. Values listed in the table are mg Phosphorus/kg tissue.

Accession	Regrow Phosphorus (mg/kg)	Regrow Phosphorus Standard Error	Regrow Number of Seeds Tested	Collection Phosphorus	Collection Phosphorus Standard Error	Collection Number of seeds tested	Phosphorus Level
PI081042-1	5464.77	127.08	12	4149.66	109.15	5	Low
PI424159B	5965.40	160.35	12	4305.02	168.68	5	Low
PI475822B	5830.14	179.63	11	5819.22	335.34	6	Low
PI567691	6121.47	186.62	11	6001.76	372.65	6	Low
PI086081	6665.44	123.66	12	8280.90	123.01	6	High
PI423813	7100.48	198.13	14	8421.17	481.09	6	High
PI089772	6432.51	130.76	12	8785.44	300.08	6	High
PI567721	5622.10	193.65	12	9602.50	504.11	5	High

In a separate experiment, 10 lines total, four low sulfur accumulating lines and six high sulfur accumulating lines were selected and regrown in both a field and greenhouse trial. In both the field and greenhouse experiment, all of the six high sulfur lines had a higher sulfur accumulation than the four low accumulating lines. Interestingly, the field grown varieties had a larger difference in sulfur accumulation between the high and low varieties (Figure 5 and Table 7). Although not selected for accumulation of other elements, there was also a correlation between measured values in the germplasm collection and the regrow set for many other elemental phenotypes tested (Supplemental Figures 5 and 6). Broad-sense heritability for sulfur between the GRIN growout concentrations, the greenhouse, and the field growouts was 0.64 (Supplemental Table 5).

310 **Table 7**. Accessions chosen for validation of sulfur accumulation. **High and low sulfur accumulating lines were chosen to regrow to test the reproducibility**
311 **of ionomic traits. Values listed in the table are mg sulfur/kg tissue.**

Accession	Regrow Field Sulfur (mg/kg)	Regrow Field Standard Error	Regrow Field Number of Seeds Tested	Regrow Greenhouse Sulfur (mg/kg)	Regrow Greenhouse Standard Error	Regrow Greenhouse Number of Seeds Tested	Collection Sulfur (mg/kg)	Collection Sulfur Standard Error	Collection Number of seeds tested	Sulfur Level
PI096322	3674.77	82.01	6	3303.99	86.76	6	2694.52	75.46	7	Low
PI229327	3183.07	69.30	6	NA	NA	NA	2764.57	62.35	7	Low
PI507411	3190.73	26.38	4	3126.35	84.73	6	2797.00	67.14	8	Low
PI603599A	3584.44	48.23	6	3075.94	114.71	8	2874.06	64.85	8	Low
PI603162	4336.25	45.05	6	3703.22	70.82	6	3771.84	71.02	8	High
PI339734	4856.20	158.22	6	4875.50	68.81	4	3774.48	21.99	2	High
PI437377	4728.93	112.23	6	3413.30	82.30	6	3847.54	82.38	7	High
PI603910B	4301.96	64.81	5	4074.24	80.70	5	3925.33	71.42	8	High
PI082278	4703.29	51.39	5	4265.62	99.98	6	3929.56	117.16	7	High
PI424078	NA	NA	NA	4791.33	187.03	5	4245.06	78.57	5	High

(a)

(b)

Figure 5. Confirmation grow out of high and low sulfur and phosphorus accumulating lines. A, Regrow versus original concentration of 8 lines selected for high and low phosphorus accumulation. Correlation between GRIN concentration and regrow was 0.24. B, Regrow versus original concentration of 10 lines selected for high and low sulfur accumulation, regrown in both greenhouse and field environments. Error bars indicate the standard error of the replicate seeds. Correlation (r^2) between GRIN seed concentrations and the regrown high and low varieties grown in the greenhouse and in the fields were 0.61 and 0.84, respectively.

Discussion

Analysis of ionomic traits has led to a deeper understanding of the complex regulatory system organisms use to maintain homeostasis of essential elements (Baxter *et al.* 2008; Baxter 2010; Atwell *et al.* 2010; Yu *et al.* 2012). To broaden our understanding of how genetic and environmental components affect the ionome, we have developed a high-throughput ionomic phenotyping system that can rapidly measure 20 ionomic traits and seed weight in agronomically important crops, such as soybean, maize, sorghum and cotton. To assess the utility of our phenotyping system for genome wide association studies in soybean, we measured the ionome of a diverse set of more than 1300 soybean lines, divided into 14 independent populations grown in three locations over the course of a decade. Coupled with a high-resolution genetic map (Song *et al.* 2013), we performed a genome wide association study using a multi-locus mixed model procedure (Segura *et al.* 2012). We were also able to show that lines selected from these experiments for extreme phenotypes of elemental accumulation were likely to display similar phenotypes in follow up experiments.

In spite of the limited number of lines in each grow-out, one of the strengths of this study is the number of distinct field replications. Although there was no overlap between lines for any of the 14 grow-outs, we found many genetic interactions that were robust across environments and genotypes. We report several different sets of SNPs corresponding to different levels of stringency in the individual experiments and the way we compared results between the experiments. These range from the 1756 SNPs from the full models, which likely contain several false positive associations, to the two SNPs that were returned in multiple experiments for the same element. Hundreds of SNPs in the total dataset are likely to be real due to their inclusion in a more conservative model or due to being found in several locations once LD is taken into account. Several of these mapped directly to what could be considered *a priori* candidate genes that have either already been characterized in soybean or are close orthologs of metal homeostasis proteins in *A. thaliana* and other species (Table 5). The discovery of orthologs of known *Arabidopsis* genes in soybean experiments highlights the value of studies in model organisms, where the genetics and growth habits are more amenable to large scale studies. Many more overlaps between different phenotypes found in different locations suggests genetic by environmental effect on which phenotype is affected by a causal locus. Many of the SNPs which overlap across environments are novel associations with no obvious gene candidates and are strong candidates for follow-up studies to determine their relationship to plant nutrient homeostasis.

The strongest element-loci association in our study was for the cadmium phenotype which is associated with a gene that codes for HMA13, a P_{1B}-ATPase (HMA13; Glyma.09g055600) previously implicated in seed cadmium concentration in soybean (Benitez *et al.* 2012). A previous GWAS study on iron deficiency chlorosis found seven loci strongly associated with the disease phenotype (Mamidi *et al.* 2014). Our analysis returned 3 of the seven loci found in that study, all associated with seed Fe, including the two strongest associations from the IDC panel: a locus associated with nicotianamine synthase 3 (NAS3; Glyma.03g231200) and a locus associated with a stabilizer of iron transporter (AGO10; Glyma.05g011300). If gene discovery of

358 small to medium effect loci is the goal of a study, using samples from germplasm banks may not
359 be appropriate, but even with all the caveats about statistical power and gene by environment
360 interactions, we found loci that had strong candidates for some elements. These results could
361 be used to prioritize genes and lines for further characterization experiments.
362
363

Conclusion
Using state-of-the-art association mapping techniques we were able to use the data we
collected using our high-throughput ionomic phenotyping pipeline to identify both lines with
extreme phenotypes and loci associated with elemental traits. Many of these associations were
strong enough to occur across a diverse set of environmental conditions, while others were
found in only one of the environments tested. While there are likely many more associations in
our GWAS dataset that we haven't yet explored, this experiment serves as a proof of concept of
using stored seed to perform GWAS on ionomic traits. While our efforts were focused on the
identification of markers associated with elemental traits, the SNPs identified were associated
with many *a priori* candidate genes. The use of seeds as the phenotyped tissue allows for the
direct association of the consequences of allelic difference of SNPs and associated candidate
genes with traits that affect the tissue with the most agronomic importance in soybeans. While
planned experiments with more replication and higher numbers of lines will always have more
power to identify genetic and environmental factors driving elemental accumulation in the seed,
this study demonstrates the utility of leveraging available samples to screen germplasm.
379

380 **Materials and Methods**
381

382 **Germplasm**
383

A diverse panel of 1653 soybean accessions was selected from the core soybean collection of
the USDA Soybean Germplasm Collection, as described in the results. Because the mission of
NPGS is to maintain a viable collection of plant germplasm, the collections are periodically
regrown to maintain viable seed. The size of the soybean germplasm collection necessitates
that only a subset of the complete germplasm collection is grown-out each year. Furthermore,
the diverse panel of accessions belongs to a variety of maturity groups and was grown-out in
three separate locations: Stoneville, MS, Urbana, IL, and Upala, Costa Rica. The 1653 lines in
the panel are, thus, broken into 13 distinct year and location sets, with no overlap of lines
between years or locations (Table 1). The Costa Rica dataset had no individual years with
enough lines (>50) to perform a successful association analysis. However, by creating three
additional datasets by combining data from each location, regardless of year, we were able to
analyze data from the Costa Rica grow-outs.
396

397 **Confirmation Growouts**
398

Small plots of four low sulfur accumulating lines and six high sulfur accumulating lines were
grown in Mexico silt loam soil at Bradford Research and Extension Center, Columbia, Missouri.
Cultural practices were typical of those utilized for soybean production in the Midwest US. The

same set of plants were also grown in environmentally controlled greenhouse in 6 liter pots
containing PRO-MIX (Premier Horticulture, Quebec, Canada) medium amended with Osmocote
Classic controlled release fertilizer (Scotts, OH). Greenhouse settings were 16 h day length with
30/18°C day/night temperatures.
406
Small plots of differential phosphorus lines were grown out in 2012 at South Farm Agricultural
Research Center (Columbia, MO, Latitude 38.908189, Longitude -92.278693, Mexico silt loam
soil) as single plots of 5 feet long with a 3 foot gap between rows and 30 inches between rows.
Field conditions were typical of soybean production in the Midwest US, with NPK Fertilizer
applied at rates appropriate according to soil analyses (10.6/50/75) and two pre-emergent
herbicides were applied before planting: Authority First (Authority First Corp, Philadelphia, PA)
applied at 6.45 oz/acre; and Stealth applied at 1 qt/ac (Loveland Products, Loveland, CO, USA).
Post-emergent herbicides were also used: Ultra Blazer (UPI, King of Prussia, PA, USA) applied
at 1.5pt/acre; Basagran (Arysta LifeScience North America, LLC, Cary, NC, USA) applied at
1.5pt/acre and Select Max (Valent Biosciences Corp., Libertyville, IL, USA) applied at 24
oz/acre. At maturity, plots were bulk harvested and threshed and a subsample was used for
ICP-MS analysis.
419
420 **Ionomic Phenotyping by ICP-MS**
421
422 Samples were phenotyped on two separate occasions for the elemental concentrations for B,
423 Na, Mg, Al, P, S, K, Ca, Mn, Fe, Co, Ni, Cu, Zn, As, Se, Rb, Mo, and Cd following the analytical
424 methods described in Ziegler et al. (2013). Seed weight is also recorded for each sample
425 analyzed, so it was also included as a phenotype in our study.
426
A simple weight normalization procedure to correct measured sample concentrations for seed
size was found to introduce artifacts, especially for elements whose concentration is at or near
the method detection limit. This could either be due to a systematic over or under reporting of
elemental concentrations by the ICP-MS procedure or a violation of the assumption that all
elemental concentrations scale linearly with weight. We used an alternative method to normalize
for seed weight following the method recently reported in Shakoor et al. (2016). A linear model
was developed modeling unnormalized seed concentrations against seed weight and the
analytical experiment the seed was run in. The residuals from this linear model were then
extracted and used as the elemental phenotype. For each element, the phenotypic
measurement was taken as the median of the elemental concentrations from the 2 or 8 seeds
measured from each line (after outlier removal of measurements with a median absolute
deviation of >10 where we had enough samples). To meet the normality assumptions required
for GWAS, an analysis using the Box-Cox algorithm was used to determine an appropriate
transformation for each trait (Box and Cox 1964). Since each grow-out has a distinct set of lines,
which may result in different phenotypic distributions, transformations were performed
separately for each element in each dataset listed in Table 1. Transformations were selected
based upon the 95% confidence interval returned by the Box-Cox function implemented in the R
package MASS (Box and Cox 1964; Venables *et al.* 2002).
445

446 **GWAS**
447
All of the lines included in this analysis (and all of the annual accessions in the Soybean
Germplasm Collection in 2010) have been genotyped using the SoySNP50K beadchip and are
available at soybase.org (Song *et al.* 2013). Separate genotype files were generated for each
grow-out that contain only the lines present in that grow-out. The genotype files were each
filtered to remove SNPs with a minor allele frequency less than 0.05 and missing SNPs were
imputed as the average allele for that SNP. The number of SNPs for each grow-out varied
between 31,479 and 36,340. The final number of SNPs used for association mapping of each
grow-out are listed in Table 1. SNPs were called using the Glyma1.1 reference genome. All
SNP base pair locations reported are from a map to Glyma1.1.
457
Both kinship and structural components were included in the mixed model and were calculated
using the filtered genotype matrix containing all 1391 lines found across all 13 grow-outs. The
kinship matrix was calculated using the VanRaden method as implemented in GAPIT
(VanRaden 2008; Lipka *et al.* 2012). To correct for population stratification a principal
component analysis was performed. The first ten principal components were used as fixed
effects in the mixed model.
464
Association mapping was performed using a multilocus mixed model (MLMM) approach that
performs a stepwise mixed-model regression with forward inclusion and backward elimination of
genotypic markers included as fixed effects (Segura *et al.* 2012). In this model forward steps are
performed until the heritable variance estimate reaches 0 (indicating the current model includes
covariates that explain all of the heritable phenotypic variance) or a maximum number of
forward-inclusion steps have been performed, which we set at 40.
471
MLMM implements two model selection methods to determine the optimal mixed model from the
set of step-wise models calculated: the extended Bayesian information criterion (EBIC, Chen
and Chen 2008) and the multiple-Bonferroni criterion (mbonf, Segura *et al.* 2012). The EBIC
model uses the Bayesian information criteria to select a model taking into account both number
of SNPs in the analysis as well as number of cofactors in the model. In our analysis, the EBIC
was usually less conservative (eg. selected larger models). A larger model likely increases the
number of type 1 errors, but it is less likely to miss true associations. Because we are
performing a further selection step comparing results across independent experiments, we used
the EBIC models for further analysis. Additionally, we also analyzed the cofactors returned by
the final forward inclusion model (maximum model), which includes either the maximum 40
cofactors or the total number of cofactors needed to explain the estimated heritability.
483
484 SNPs included as cofactors in either the EBIC model or the maximum model were compared
485 across GWAS experiments. SNPs were determined to overlap with a neighboring SNP if it had
486 an r^2 LD of >0.2.
487
488 **Calculation of Linkage Disequilibrium**
489

Linkage disequilibrium, expressed as a correlation coefficient between markers (r^2), was
calculated using the filtered SNP data set containing all 1391 lines from the experiment and the
LD function of the 'genetics' R package (Warnes *et al.* 2013).

Germplasm and Data Availability

Lines used can be found at the USDA Soybean Germplasm Center. All scripts and data used
can be found at www.ionomicshub.org and https://github.com/baxterlab/SoyIonomicsGWAS.

Figure/Table Legends

Supplemental Figure 1. Principal component analysis of the genotypes of 1391 soybean
lines. Colored by GRIN growout.

Supplemental Figure 2. Elemental accumulation in soybean seeds across experimental
grow-outs.

Supplemental Figure 3. Distribution of all elemental phenotypes in all grow-outs. Lines
are ordered by the median of between 2 and 8 seed replicates.

Supplemental Figure 4. QQ-plots for all GWAS experiments performed.

Supplemental Figure 5. Regrow versus original concentration for all phenotypes in the
phosphorus selection experiment.

Supplemental Figure 6. Regrow versus original concentration for all phenotypes in the
sulfur selection experiment.

Supplemental Table 1. Raw ionomics data and phenotypes after transformation for
GWAS for all lines in the experiment.

Supplemental Table 2. Box-Cox suggested transformations for ionomics phenotypes.

Supplemental Table 3. All SNPs returned in either 'All Cofactor', 'EBIC', or 'Multiple
Bonferroni' models for all GWAS experiments.

Supplemental Table 4. SNPs returned in two or more grow-outs based on Linkage
Disequilibrium calculation.

Supplemental Table 5. Broad-sense heritabilities calculated for ionomic traits in the
sulfur and phosphorus confirmation experiments.

Literature Cited

Ardlie, K. G., L. Kruglyak, and M. Seielstad, 2002 Patterns of linkage disequilibrium in the
human genome. Nat. Rev. Genet. 3: 299–309.

Asaro, A., G. Ziegler, C. Ziyomo, O. Hoekenga, B. Dilkes *et al.*, 2016 The Interaction of
Genotype and Environment Determines Variation in the Maize Kernel Ionome. G3
Genes Genomes Genet. g3.116.034827.

Atwell, S., Y. S. Huang, B. J. Vilhjalmsson, G. Willems, M. Horton *et al.*, 2010 Genome-wide
association study of 107 phenotypes in a common set of Arabidopsis thaliana inbred
lines. Nature 465: 627–631.

Barberon, M., 2017 The endodermis as a checkpoint for nutrients. New Phytol. 213: 1604–1610.

Baxter, I., 2010 Ionomics: The functional genomics of elements. Brief. Funct. Genomics 9: 149–
156.

Baxter, I., 2015 Should we treat the ionome as a combination of individual elements, or should
we be deriving novel combined traits? J. Exp. Bot. 66: 2127–2131.

Baxter, I., and B. P. Dilkes, 2012 Elemental Profiles Reflect Plant Adaptations to the
Environment. Science 336: 1661–1663.

Baxter, I., P. S. Hosmani, A. Rus, B. Lahner, J. O. Borevitz *et al.*, 2009 Root Suberin Forms an
Extracellular Barrier That Affects Water Relations and Mineral Nutrition in Arabidopsis.
PLOS Genet. 5: e1000492.

Baxter, I. R., O. Vitek, B. Lahner, B. Muthukumar, M. Borghi *et al.*, 2008 The leaf ionome as a
multivariable system to detect a plant's physiological status. Proc. Natl. Acad. Sci. 105:
12081–12086.

Baxter, I. R., G. Ziegler, B. Lahner, M. V. Mickelbart, R. Foley *et al.*, 2014 Single-Kernel Ionomic
Profiles Are Highly Heritable Indicators of Genetic and Environmental Influences on
Elemental Accumulation in Maize Grain (Zea mays) (X. Zhang, Ed.). PLoS ONE 9:
e87628.

Benitez, E. R., M. Hajika, and R. Takahashi, 2012 Single-Base Substitution in P1B-ATPase
Gene Is Associated with a Major QTL for Seed Cadmium Concentration in Soybean. J.
Hered. 103: 278–286.

Box, G. E. P., and D. R. Cox, 1964 An Analysis of Transformations. J. R. Stat. Soc. Ser. B
Methodol. 26: 211–252.

Cabannes, E., P. Buchner, M. R. Broadley, and M. J. Hawkesford, 2011 A comparison of sulfate
and selenium accumulation in relation to the expression of sulfate transporter genes in
Astragalus species. Plant Physiol. 157: 2227–2239.

Chao, D.-Y., B. Dilkes, H. Luo, A. Douglas, E. Yakubova *et al.*, 2013 Polyploids exhibit higher
potassium uptake and salinity tolerance in Arabidopsis. Science 341: 658–659.

Chao, D.-Y., K. Gable, M. Chen, I. Baxter, C. R. Dietrich *et al.*, 2011 Sphingolipids in the root
play an important role in regulating the leaf ionome in Arabidopsis thaliana. Plant Cell
23: 1061–1081.

Chen, J., and Z. Chen, 2008 Extended Bayesian information criteria for model selection with
large model spaces. Biometrika 95: 759–771.

El Kassis, E., N. Cathala, H. Rouached, P. Fourcroy, P. Berthomieu *et al.*, 2007
Characterization of a Selenate-Resistant Arabidopsis Mutant. Root Growth as a
Potential Target for Selenate Toxicity. Plant Physiol. 143: 1231–1241.

FAO, 1996 *Global climate change and agricultural production: direct and indirect effects of*
changing hydrological, pedological, and plant physiological processes (F. A. Bazzaz &
W. G. Sombroek, Eds.). Food and Agriculture Organization of the United Nations ; Wiley,
[Rome] : Chichester ; New York.

Gaillard, S., H. Jacquet, A. Vavasseur, N. Leonhardt, and C. Forestier, 2008 AtMRP6/AtABCC6,
an ATP-Binding Cassette transporter gene expressed during early steps of seedling
development and up-regulated by cadmium in Arabidopsis thaliana. BMC Plant Biol. 8:
22.

Golldack, D., O. V. Popova, and K.-J. Dietz, 2002 Mutation of the Matrix Metalloproteinase At2-
MMP Inhibits Growth and Causes Late Flowering and Early Senescence in Arabidopsis.
J. Biol. Chem. 277: 5541–5547.

Hwang, E.-Y., Q. Song, G. Jia, J. E. Specht, D. L. Hyten *et al.*, 2014 A genome-wide association
study of seed protein and oil content in soybean. BMC Genomics 15: 1.

Lipka, A. E., F. Tian, Q. Wang, J. Peiffer, M. Li *et al.*, 2012 GAPIT: genome association and
prediction integrated tool. Bioinformatics 28: 2397–2399.

Mamidi, S., R. K. Lee, J. R. Goos, and P. E. McClean, 2014 Genome-Wide Association Studies
Identifies Seven Major Regions Responsible for Iron Deficiency Chlorosis in Soybean
(Glycine max). PLoS ONE 9: e107469.

Oliveira, M. F., R. L. Nelson, I. O. Geraldi, C. D. Cruz, and J. F. F. de Toledo, 2010 Establishing
a soybean germplasm core collection. Field Crops Res. 119: 277–289.

Qanbari, S., E. C. G. Pimentel, J. Tetens, G. Thaller, P. Lichtner *et al.*, 2010 The pattern of
linkage disequilibrium in German Holstein cattle. Anim. Genet. 41: 346–356.

Segura, V., B. J. Vilhjálmsson, A. Platt, A. Korte, Ü. Seren *et al.*, 2012 An efficient multi-locus
mixed-model approach for genome-wide association studies in structured populations.
Nat. Genet. 44: 825–830.

Shakoor, N., G. Ziegler, B. P. Dilkes, Z. Brenton, R. Boyles *et al.*, 2016 Integration of
experiments across diverse environments identifies the genetic determinants of variation
in Sorghum bicolor seed element composition. Plant Physiol. pp.01971.2015.

Shigaki, T., J. K. Pittman, and K. D. Hirschi, 2003 Manganese Specificity Determinants in the
ArabidopsisMetal/H+ Antiporter CAX2. J. Biol. Chem. 278: 6610–6617.

Song, Q., D. L. Hyten, G. Jia, C. V. Quigley, E. W. Fickus *et al.*, 2013 Development and
Evaluation of SoySNP50K, a High-Density Genotyping Array for Soybean. PLoS ONE 8:
e54985.

Tian, H., I. R. Baxter, B. Lahner, A. Reinders, D. E. Salt *et al.*, 2010 Arabidopsis NPCC6/NaKR1
Is a Phloem Mobile Metal Binding Protein Necessary for Phloem Function and Root
Meristem Maintenance. Plant Cell 22: 3963–3979.

VanRaden, P. M., 2008 Efficient methods to compute genomic predictions. J. Dairy Sci. 91:
4414–4423.

Venables, W. N., B. D. Ripley, and Venables, W. N, 2002 *Modern applied statistics with S*.
Springer, New York.

Wallace, J. G., P. J. Bradbury, N. Zhang, Y. Gibon, M. Stitt *et al.*, 2014 Association Mapping
across Numerous Traits Reveals Patterns of Functional Variation in Maize. PLoS Genet
10: e1004845.

Warnes, G., G. Gorjanc, F. Leisch, and M. Man, 2013 *genetics: Population Genetics*.

Yu, D., J. M. C. Danku, I. Baxter, S. Kim, O. K. Vatamaniuk *et al.*, 2012 High-resolution
genome-wide scan of genes, gene-networks and cellular systems impacting the yeast
ionome. BMC Genomics 13: 623.

Ziegler, G., A. Terauchi, A. Becker, P. Armstrong, K. Hudson *et al.*, 2013 Ionomic Screening of
Field-Grown Soybean Identifies Mutants with Altered Seed Elemental Composition.
Plant Genome 6: 0.