Frequency of mosaicism points towards mutation-prone early cleavage cell

divisions in cattle.

Chad Harland[1#], Carole Charlier[1#], Latifa Karim[2], Nadine Cambisano[2], Manon

Deckers[2], Myriam Mni[1], Erik Mullaart[3], Wouter Coppieters[2], Michel Georges[1].

[1] Unit of Animal Genomics, GIGA-R & Faculty of Veterinary Medicine, University

of Liège, Belgium. [2] GIGA Genomics Platform, University of Liège, Belgium. [3] CRV,

Arnhem, The Netherlands.

 # Contributed equally to this work

Correspondence: michel.georges@ulg.ac.be

It has recently become possible to directly estimate the germ-line de novo

mutation (*dnm*) rate by sequencing the whole genome of father-mother-

offspring trios, and this has been conducted in human[1-5], chimpanzee[6],

mice[7], birds[8] and fish[9]. In these studies *dnm*'s are typically defined as

variants that are heterozygous in the offspring while being absent in both

parents. They are assumed to have occurred in the germ-line of one of the

parents and to have been transmitted to the offspring via the sperm cell or

oocyte. This definition assumes that detectable mosaïcism in the parent in

which the mutation occurred is negligible. However, instances of

detectable mosaïcism or premeiotic clusters are well documented in

humans and other organisms, including ruminants[10-12]. We herein take

advantage of cattle pedigrees to show that as much as ~30% to ~50% of

***dnm*'s present in a gamete may occur during the early cleavage cell**

divisions in males and females, respectively, resulting in frequent

detectable mosaïcism and a high rate of sharing of multiple *dnm*'s between

siblings. This should be taken into account to accurately estimate the

mutation rate in cattle and other species.

To study the process of *dnm*'s in the cattle germ-line, we sequenced the whole

genome of 54 animals from four pedigrees. Grand-parents, parents and offspring

(referred to as probands) were sequenced at average 26-fold depth (min = 21),

and grand-offspring at average 21-fold depth (min = 10). The source of DNA was

venous blood for females and sperm for males. The genome of one male proband

(Pr 1) was sequenced both from semen (26-fold depth) and blood DNA (37-fold

depth) (Figure 1A).

Using the standard definition, we identified 190 candidate *dnm*'s as variants that

were (i) detected in a proband, (ii) absent in both parents (and grand-parents

when available), (iii) transmitted to at least one grand-offspring, and (iv) not

previously reported in unrelated individuals from the 1,000 Bulls project[13]

(Suppl. Figure 1&2 and Suppl. Table 1). For confirmation, we developed

amplicons spanning 113 candidate *dnm*'s and sequenced them at average depth

of ~2,187 in the 54 animals plus 55 relatives (Figure 1A). This confirmed the

genuine nature of 110/113 variants, demonstrating the excellent specificity of

our bioinformatics pipeline. The three remaining ones were also detected in one

of the parents (although not in the grand-parents) in the confirmation, and

momentarily ignored.

We first examined what proportion of *dnm*'s detected in a proband might

actually have occurred during its development rather than being inherited via

the sperm or oocyte. An unambiguous distinction between the two types of

dnm's is their degree of linkage with either the paternal or maternal haplotype

upon transmission to the next generation (i.e. the grand-offspring in Figure 1A).

Dnm's that have occurred in the germ-line of the sire will show *perfect* linkage

with the proband's paternal haplotype in the grand-offspring (i.e. always

transmitted with the paternal haplotype, never transmitted with the maternal

haplotype), while *dnm*'s that have occurred in the germ-line of the dam will show

perfect linkage with the proband's maternal haplotype in the grand-offspring. On

the contrary, *dnm*'s that have occurred during the development of the proband

will be in *complete* (but imperfect) linkage with either the paternal or maternal

haplotype (i.e. sometimes transmitted with the maternal haplotype, never

transmitted with the paternal haplotype, or sometimes transmitted with the

paternal haplotype, never transmitted with the maternal haplotype)(Suppl.

Figure 1). Across the four pedigrees, 124 variants were in perfect linkage with

the paternal haplotype, 32 in perfect linkage with the maternal haplotype, 10 in

complete (but imperfect) linkage with the paternal haplotype and 21 in complete

(but imperfect) linkage with the maternal haplotype (Figure 1B). If the 10+21

dnm's indeed occurred during the development of the proband rather then being

inherited from the sire or dam, the *dnm* dosage (defined as the proportion of

reads spanning the *dnm* site that carry the mutant allele) is expected to be < 50%

in the proband but equal to 50% in the grand-offspring inheriting the *dnm*. The

mean dosage was 0.26 in the proband, and 0.52 in the grand-offspring, and this

difference was highly significant (p < 10^{-6}). The corresponding means were 0.48

and 0.49 (p = 0.40) for the 124+32 mutations showing perfect linkage with

either the paternal or maternal haplotype (Figure 1C).

We conclude that in cattle ~17% of *dnm*'s detected in an animal using standard

procedures are not inherited from the sire or dam but correspond to premeiotic

clusters generated during the development of the individual.   This is a lower

bound, as *dnm* will only be detected and recognized as having occurred in the

proband if (i) the *dnm* dosage is sufficiently high for the proband to be called

heterozygote, (ii) the *dnm* is transmitted to at least one grand-offspring, and (iii)

complete (but imperfect) linkage is demonstrated in the grand-offspring (Suppl.

Figure 3). We will refer to this type of *dnm*'s as Proband-Mosaic (PM), while the

others will be referred to as Sire-Non-Mosaic (SNM) (meaning that the sire is not

mosaic for a mutation transmitted via his sperm), or as Dam-Non-Mosaic (DNM)

(meaning that the dam is not mosaic for a mutation transmitted via her oocyte).

The proportion of PM (but not SNM and DNM) mutations differed significantly

between probands (p = 0.004); neither differed significantly between sexes (p >

0.30).   Of particular interest, the three PM mutations of proband 1 were

detected in both sperm and blood DNA (Suppl. Table 1), indicating that they

occurred early in development (see hereafter).

If detectable mosaicism for *dnm*'s is common in the individual in whom they

occurred, requiring their absence in the DNA of the parents (as typically done)

will cause genuine *dnm*'s to be eliminated. We took advantage of the grand-

parents available in three pedigrees to recover such events as variants that were

(i) absent in the grand-parents, (ii) detected in either sire or dam with a dosage

significantly < 50% (Suppl. Table 1), (iii) transmitted to the proband with a

dosage of ~50%, (iv) transmitted to at least one grand-offspring with a dosage of

~50%, and (v) not previously reported in unrelated individuals[13].   We will refer

to these types of mutations as Sire-Mosaic (SM) and Dam-Mosaic (DM),

respectively (meaning that the sire/dam is detectably mosaic for a *dnm*

transmitted via the sperm/oocyte)(Suppl. Figure 1).   We detected 61 such

candidate events, including the 3/113 variants mentioned above (Suppl. Table 1

and Suppl. Figure 2).   We developed amplicons for 34, and sequenced (average

1,498-fold depth) all 54 individuals plus 55 relatives (including ≥ 5 half-sibs of

the probands) (Figure 1A).   We took advantage of whole genome sequence

information that became available for 27 half-sibs, to trace the inheritance of the

remaining 24 candidate variants.   The ensuing data indicated that 11/61

candidates were genuine *dnm*'s but occurred in the germ-line of one of the

grand-parents rather than one of the parents (dosage ~50% in the sire or dam,

and perfect linkage in the half-sibs).   The SM/DM status was unambiguously

demonstrated for 40 (dosage < 50% in the sire or dam in the confirmation,

transmission to half-sibs, and complete (but imperfect) linkage) and strongly

supported for the remaining 10 (dosage < 50% in the sire/dam in the

confirmation or complete (but imperfect) linkage yet without transmission)

(Suppl. Figure 2 and Figure 1B).   Further supporting the genuine nature of the

SM/DM mutations, the dosage was 0.12 on average in the corresponding parent,

while being 0.51 in descendants ($p < 10^{-6}$) (Figure 1C).

For *dnm*'s that were detectably mosaic in sperm (SM and PM in male probands),

allelic dosage was significantly correlated with rate of transmission (p = 0.025)

and strength of linkage (p = 0.0002). These correlations were not significant for

dnm's that were detectably mosaic in blood (DM and PM in female probands).

This suggests that the degree of mosaicism in the soma is a poor indicator of the

degree of mosaicism in the germ line (Figure 1D).   Accordingly, the rate of

transmission of the three PM mutations of proband 1 to its 53 offspring was

better predicted by their dosage in sperm than in blood (Suppl. Figure 4).

Considering SNM/SM and DNM/DM mutations jointly, we conclude that on

average a sire is detectably mosaic (in sperm) for 29% of *dnm*'s present in a

sperm cell, while a dam is detectably mosaic (in blood) for 51% of *dnm*'s present

in an oocyte.   These are lower bounds as we only considered *dnm*'s for which the

dosage was significantly < 0.5 in the parent (condition (ii) above).   These figures

are possibly consistent with recent reports in the mouse (~25%)[7], but

considerably larger than current estimates in human (~5%)[14].   They are

certainly larger than expected assuming that the mutation rate per cell division

is uniform throughout development, and it suggests that the mutation rate is

higher for early cell divisions (Suppl. Figure 5).   Moreover, when analyzing the

transmission patterns of SM and DM mutations to the half-sibs of the proband (in

whom the *dnm*'s were detected), we were struck by the fact that (i) >60% of half-

sibs share at least one *dnm* with the proband, while <50% are expected (p =

0.05), and (ii) half-sibs sharing multiple *dnm*'s with the proband appeared

surprisingly common (Suppl. Table 2). These findings also indicate that a

substantial proportion of *dnm*'s must occur early in development and be present

in the precursor cells common to the soma and germ line (Suppl. Figure 5).

In mammals, after fertilization, cleavage, and segregation of (i) the inner cell

mass from the trophoblast, (ii) the epiblast from the hypoblast, (iii) the

embryonic epiblast from the amniotic ectoderm, a small number of epiblast-

derived cells located in the wall of the yolk sac in the vicinity of the allantois are

induced to become primordial germ cells (PGCs). These migrate to the primitive

gonad where they expand and produce >1 million gametogonia. Oogonia initiate

meiosis prior to birth in females. Spermatogonia will resume mitotic divisions at

puberty allowing (i) the maintenance of a pool of stem cell like spermatogonia,

and (ii) sustained spermiogenesis involving ~3 additional mitotic divisions

followed by meiosis (Suppl. Figure 6).   We simulated the process of de novo

mutagenesis in the male and female germ cell lineages assuming (i) uniform pre-

and post-natal mutation rates per cell division, and (ii) 40 PGCs sampled at

random from the embryonic epiblast-derived cells[15].   Pre- and post-natal

mutation rates were adjusted to match the observed number of mutations per

gamete (34 in sperm, 14 in oocytes).   Under these conditions, we virtually never

observed the level of mosaicism, nor the sharing between sibs characterizing the

real data (Figure 2). We (i) increased the relative mutation rate during the early

cell divisions (keeping the mutation rate per gamete constant)(10 and 20-fold

increase during the first 4, 7, 11, 15 and 18 cell divisions; Suppl. Figure 6), (ii)

reduced the number of induced PGCs (4, 10, or 40), and (iii) varied the

relatedness between PGCs (i.e. sampled randomly amongst all embryonic

epiblast-derived cells or from a sub-sector)(Suppl. Figure 6).   Increasing the

mutation rate during the very first cell divisions matched the real data much

better (Figure 2).   To quantitatively evaluate model fitting we used (i) the

proportion of PM, SM and DM mutations with corresponding rate of mosaicism

in sperm and soma, and (ii) the proportion of sibs sharing 0, 1, 2, ... *dnm*'s with a

proband, to compute the likelihood of the data under different scenarios (see

M&M). A 20-fold increased mutation rate during the first four cell divisions,

combined with 4 related PGCs fitted the data best (Table 1 and Suppl. Table 2).

The data were $\geq 10^{16}$ times less likely under models assuming a uniform mutation

rate throughout development, and $\geq 10^5$ times less likely assuming an increased

mutation rate passed the 7^{th} cell division (after segregation of inner cell mass

and throphoblast)(Table 1 and Suppl. Table 2).

When accounting for genome coverage, the estimated number of *dnm*'s per

gamete (SNM+SM, DNM+DM) averaged 46.6 for sperm cells and 18.1 for oocytes

(male/female ratio of 2.6), corresponding to an average mutation rate of

$\sim 1.2 \times 10^{-8}$ per base pair per gamete. Including an estimate (from the simulations)

of the number of missed SM (~ 3.3) / DM (~ 1.1) and misclassified PM mutations

(~ 2.9 to ~ 8 depending on the proband), yields an average mutation rate of

$\sim 1.17 \times 10^{-8}$ per base pair per gamete and a male/female ratio of 2.4.   The

standard approach of ascertaining *dnm*'s (i.e. erroneously considering PM

mutations, ignoring SM and DM mutations) would have yielded a mutation rate

of 0.9×10^{-8} per bp per gamete, with a 2.5-fold higher mutation rate in bulls than

in cows.

Two hundred twenty of the 237 identified *dnm*'s were nucleotide substitutions,

17 small insertion-deletions.   The non-mosaic classes of mutations (SNM and

DNM) were ~ 30-fold enriched in CpG>TpG transitions as expected.   This

signature was also present but less pronounced for mosaic mutations (PM, SM

and DM).   Mosaic mutations were ~ 2.6-fold enriched in C>A and/or G>T

transversions, largely due to GpCpA>GpApA and TpCpT>TpApT substitutions

(Figure 3).   This was unlikely to be an artifact for reasons spelled out in Suppl.

Figure 7.   It is noteworthy that this is exactly the same mutational signature as

the one recently reported for human embryonic somatic mutations[16].   There

was no obvious difference between the profile of *dnm*'s in the male and female

germ line (data not shown). In general, *dnm*'s appeared uniformly scattered

across the genome (Suppl. Figure 8).

The enrichment of C>A/G>T transversions in the mosaic mutations caused the

overall Ti/Tv ratio to be 1.33, well below expectations. This was likely due to

sampling variation (meaning that Ti/Tv ratios might differ between families and

that we by chance sampled families at the low end), as the Ti/Tv ratio was 1.99

for 2,530 candidate *dnm*'s detected with the same bioinformatics pipeline in a

follow-up study of 113 probands (excluding the ones analyzed in this work), i.e.

closer to expectations and the 2.2 Ti/Tv ratio of SNPs segregating in the

Holstein-Friesian dairy cattle population (MAF ≤ 0.01; rare allele considered to

be the derived allele). However, the spectrum of the 2,530 *dnm*'s remained

significantly different from the SNP spectrum, with an excess of C>A/G>T

transversions in the mosaic class of mutations, an excess of C>T/G>A transitions

in both mosaic and non-mosaic mutations, and a paucity of T>C/A>G transitions

in both mosaic and non-mosaic mutations (Suppl. Figure 7). This could point

towards recent alterations of the mutational profile in domestic cattle. It is

worth noting in this regard that most analyzed animals were bred using artificial

insemination and/or in vitro embryo production. It seems unlikely that artificial

insemination with frozen semen could explain the observed familial clustering of

specific *dnm*'s. However, it is conceivable that in vitro maturation, fertilization

and culture of oocytes and embryos affect the *dnm* rate, possibly by perturbing

DNA replication. It is important to determine whether this is the case, especially

as the same methods are increasingly used in human reproduction.

Dnm's occurring during the development of an individual, should a priori affect

the maternal and paternal chromosome with equal probability.   When

considering the PM, SM and DM jointly, 48 mosaic mutations occurred on the

maternal chromosome versus 31 on the paternal chromosome (p = 0.11).   This

trend suggests that the maternal and paternal chromosomes might be

epigenetically distinct during early development and that this may affect their

mutability.

Our work points towards the fact that direct estimates of mutation rates from

sequencing families may have to be revisited, taken PM, SM and DM status into

account, to obtain more accurate estimates of the mutation rate per gamete and

per generation. This may affect both the overall mutation rate as well as its

male/female ratio.   However, our analyses suggest that the effect is likely to be

modest and would, for instance, be insufficient to explain the present 2-fold

discrepancy between direct and indirect estimates in human studies[17,18].   We

confirmed by simulation that the rate of mosaicism does not significantly affect

the rate of nucleotide substitution per generation or average fixation time[19]

(Suppl. Figure 9).

Our work calls for a careful reevaluation of the importance of mosaicism for

dnm's in humans.   If more common than presently appreciated, the recurrence

risk of *dnm*-dependent disorders in sibs may be higher than generally assumed

[11,17].   Moreover, a non-negligible proportion of true *dnm*'s may have been ignored

(because they were detected at low dosage in the parents) in *dnm*-dependent

searches for genes underlying inherited disorders hence reducing the potential

power of such studies.

Acknowledgements

This work was funded by the DAMONA Advanced ERC project to Michel Georges.

Carole Charlier is Senior Research Associate from the Fonds de la Recherche

Scientifique - FNRS (F.R.S.-FNRS). Chad Harland has been funded in part by

Livestock Improvement Corporation (New Zealand). We are grateful to Erik

Mullaart and CRV (Arnhem the Netherlands) for providing us with the sperm and

blood samples. We used the supercomputing facilities of the Consortium des

Equipements de Calcul Intensif en Fédération Wallonie Bruxelles (CECI) funded

by the F.R.S-FNRS.

Authors contributions

254   MG, CH, CC: designed the experiments. EM: provided samples. LK, NC, MD, WC:

255   performed the sequencing. CH, MG, CC: analyzed data. MG, CH, CC: wrote the

256   paper.

Data availability

259   All sequence data will be made freely available in public databases.

References

1. Roach JC et al. Analysis of Genetic Inheritance in a Family Quartet by Whole-

Genome Sequencing. *Science* **328**: 636–639 (2010).

2. Conrad DF et al. Variation in genome-wide mutation rates within and between

human families. *Nat Genet* **43**: 712–714 (2011.).

3. Kong A et al. Rate of de novo mutations and the importance of father/'s age to

disease risk. *Nature* **488**: 471–475 (2012).

4. Campbell CD et al. Estimating the human mutation rate using autozygosity in a

founder population. *Nat Genet* **44**: 1277–1281 (2012).

5. Michaelson JJ et al. Whole-Genome Sequencing in Autism Identifies Hot Spots

for De Novo Germline Mutation. *Cell* **151**: 1431–1442 (2012).

6. Venn O et al. Strong male bias drives germline mutation in chimpanzees.

Science **344**:1272-1275 (2014).

7. Lindsay SJ et al. Striking differences in patterns of germline mutation between

mice and humans. doi: https://doi.org/10.1101/082297

8. Smeds L, Qvarnström A, Ellegren H Direct estimate of the rate of germline

mutation in a bird. *Genome Res.* **26**: 1211-1218 (2016).

9. Feng C et al. Moderate nucleotide diversity in the Atlantic herring is associated

with a low mutation rate. *eLife*, in press (2017).

10. Woodruff RC & Thompson JN. Have premeiotic clusters of mutation been

overlooked in evolutionary theory? *J. Evol. Biol.* **5:**457-464 (1992).

11. Campbell IM et al. Somatic mosaicism: implications for disease and

transmission genetics. *Trends Genet.* **31**:382-392 (2015).

12. Smit M et al. Mosaicism of Solid Gold supports the causality of a noncoding A-

to-G transition in the determinism of the callipyge phenotype. *Genetics* **163**:453-

456 (2003).

13. Daetwyler HD et al. Whole-genome sequencing of 234 bulls facilitates

mapping of monogenic and complex traits in cattle. *Nat Genet* **46**:858-865

(2014).

14. Rahbari R et al. Timing, rates and spectra of human germline mutation.

Nature Genetics **48:** 126-133 (2016).

15. Ohinata et al. Blimp1 is a critical determinant of the germ cell lineage in mice.

Nature **437**:207-213 (2005).

16. Ju YS et al. Somatic mutations reveal asymetric cellular dynamics in the early

human embryo. *Nature* **543**: 714-718 (2017).

17. Scally A. Mutation rates and the evolution of germline structure. *Philos Trans*

R Soc Lond B Biol Sci **371:** 20150137 (2016).

18. Segurel L, Wyman MJ, Przeworski M. Determinants of mutation rate variation

in the human germline. *Annu Rev Genomics Hum Genet* **15**:47-70 (2014).

19. Woodruff RC & Thomson JN. The fundamental theorem of neutral evolution:

rates of substitution and mutations should factor in premeiotic clusters. *Genetics*

125: 333-339 (2005).

Figure 1: (A) Four pedigrees (I, II, III, IV) used for the detection of *dnm*'s. GP: grand-parents (PGS: paternal grand-sires, PGD,
paternal grand-dams, MGS: maternal grand-sires, MGD: maternal grand-dams), S: sires, D: dams, Pr: probands, HS: half-sibs (of the
proband), GO: grand-offspring. The five probands are labeled in red. Animals in blue were genome-sequenced at average depth of 23
and used for the detection of *dnm*'s. Animals in gray were used for confirmation by whole genome (average sequence depth of 20) or
targeted sequencing (see Supplemental Methods). DNA was extracted from venous blood for females, and semen from males, except for
Proband 1 for which both semen and blood DNA were analyzed. **(B) Numbers and types of *dnm*'s detected in the five probands (Pr1,**
Pr2, Pr3, Pr4), Pr5). Green: PM = proband mosaic, Light blue: SNM = sire non mosaic, Dark blue: SM = sire mosaic, Light red: DNM =
dam non mosaic, Dark red: DM = dam mosaic. For each bar, the lower light section corresponds to the actual number of detected *dnm*'s,
the upper darker section to an extrapolation to the whole genome based on the estimated coverage. ND = not done (because the
corresponding grand-parents were not sequenced). **(C) Cumulative frequency distribution of *dnm* dosage estimated as the**
proportion of reads carrying the mutation. Green circles: PM mutations in the probands. Light blue circles: SNM mutations in the
probands. Light red circles: DNM mutations in the probands. Dark blue circles: SM mutations in the sires. Dark red circles: DM
mutations in the dams. Grey circles: corresponding PM, SNM, DNM, SM and DM mutations in the grand-offspring. The three SM and one
DM variant with dosage of 0, were shared between the proband and at least one half-sib yet not detectable in the semen or blood of the
corresponding parent. **(D)** Relationship between the *dnm* dosage (fraction of mutant reads) and the rate of transmission to offspring
(left) and strength of linkage (right) for mutations that are detectably mosaic in the sperm of a male parent (upper), or in the blood of a
female parent (lower). Green circles: PM mutations, blue circles: SM mutations, red circles: DM mutations. The corresponding
correlations were significant in males (p = 0.015 and 1.4×10^{-5}) but not in females (p = 0.60 and 0.16).

Figure 2: (A) *Dnm*'s with detectable mosaicism in sperm DNA of male probands (PM – sperm) or sires (SM – sperm), or in blood DNA of
female probands (PM – blood) or dams (DM – blood), ranked by observed rate of mosaicism. Colored lines: real data. Pr1-5: proband 1-
5. Dark gray shaded area: 95% confidence interval obtained from simulations assuming uniform mutation rate per cell division and 40
unrelated PGCs. Light gray shaded area: 95% confidence interval obtained from simulations assuming 20-fold higher mutation rate
during the first 4 cell divisions, and 4 related PGCs (Table 1). **(B)** Distribution of the proportion of half-sibs (HS) of the probands that
share 0, or at least 1 (1+) of the *dnm*'s detected in the corresponding proband. Red bars: real observations for *dnm's* transmitted by the
dam (DM+DNM). Blue bars: real observations for *dnm's* transmitted by the sire (SM+SNM). Dark grey bars: expectation under the null
hypothesis of uniform prenatal mutation rate per cell division and 40 unrelated PGCs. Light grey bars: expectation under the best
alternative model assuming a 20x increased mutation rate during the first 4 cell division and 4 related PGCs (Table 1).

Figure 3: (A) SNM and DNM (i.e. *dnm*'s assumed to have occurred in the later stages of gametogenesis): fold excess or deficiency over expected for specific nucleotide substitutions accounting for trinucleotide context. Inset: Proportion of *dnm*'s corresponding to the six possible types of nucleotide substitutions. **(D)** Idem for PM, SM and DM (i.e. *dnm*'s assumed to have occurred in the early stages of gametogenesis). ***: $p < 0.001$, *: $p < 0.05$ (accounting for multiple testing by Sidak correction).

399 **Table 1: Relative likelihood of the observations under different models of**
400 **gametogenesis**

Fold increase mutation rate	During first x cell divisions	Number of PGCs	Related PGCs or not	Log(LR)
20x	**4**	**4**	**T**	0.00
10x	4	4	T	-0.35
20x	4	4	**F**	-0.40
10x	**7**	4	F	-1.57
20x	4	**10**	T	-1.74
20x	4	**40**	T	-3.45
10x	**11**	4	T	-5.09
20x	**15**	4	T	-7.03
10x	**18**	4	T	-9.46
1x	7	4	T	-16.17

The first four columns correspond to the parameters that were tested in the model: (i)
the fold increase of the mutation rate (1x, 10x, 20x), (ii) during the x first cell divisions
(4, 7, 11, 15, 18), (iii) the number of PGCs (4, 10, 40), and (iv) the ontogenetic
relatedness of the PGCs (F(alse) or T(rue)). Log(LR) corresponds to the logarithm
(base 10) of the likelihood of the data relative to the best model (first line).
Parameters are in bold when the corresponding model is the best given that
parameter value. We only show results for models that are the best given at least one
parameter value. Likelihoods of all models are given in Suppl. Table 3.

Methods

Whole genome sequencing. DNA was extracted from sperm (for males) or

whole blood (females and one male) for the four families and their relatives

using standard procedures.   Familial relationships were confirmed by

genotyping all DNAs with the 10K Illumina SNP chip. We constructed 550bp

insert size whole genome Illumina Nextera PCR free libraries following the

protocols recommended by the manufacturer. All samples where then sequenced

on Illumina HighSeq 2000 instruments, using the 2x100bp paired end protocol

by the GIGA Genomics platform (University of Liege). Data was mapped using

BWA mem (version 0.7.9a-r786)[20] to the BosTau6 reference genome. Alignments

were processed according to the GATK[21] best practices version 2 with PCR

duplicates marked, Indel realignment and Base Quality Score Recalibration using

known sites. GATK HaplotypeCaller (version 3.4) was used according to the N+1

workflow to generate variants from the alignments. Common variants were then

compared to a 10K Illumina SNP chip for each individual to confirm the identity

of the library.

Detection of de novo mutations.   We developed a suite of scripts to identify

dnm's from a vcf file produced by GATK and containing sequence information

about members of four-generation pedigrees such as the ones described in Fig. 1.

The first ("4_phaser_4_gen.pl") generates the linkage phase for the parents (sire

and dam), the proband, and the grand-offspring. Phasing is done based on high

quality variant positions and genotypes (f.i. QUAL score ≥ 50,000; PL scores ≥

100; sequence depth ≤ 2.5x the average sequence depth).   The outcome is

knowledge   of   the   grand-parental   origin   of   the   paternal   and   maternal

chromosomes of the proband including the identification of cross-over events, as

well as the grand-parental origin of the chromosomes transmitted by the

proband to the grand-offspring including the identification of cross-over events.

The second module ("5_de_novo_detector_4_gen.pl") identifies the candidate

dnm's per se. It first identifies diallelic variant positions for which all grand-

parents, sire, dam and proband have a genotype and sequence coverage between

set limits (f.i. 10 and 60). The proportion of variants sites satisfying these depth

limits was used to estimate the proportion of the genome (of the total of the

2,670,422,299 base pairs in the bovine Bostau6 build) that was explored.

Candidate *dnm*'s were then identified as sites for which (i) the QUAL score was ≥

100, (ii) the proband had genotype 0/1 with corresponding PL-scores ≥ 40, (iii)

none of the grand-parents had reads with the alternate allele (AD = x,0), (iv)

either the sire or the dam had no reads with the alternate allele, and (v) reads

with the alternate allele were found in the grand-offspring. The same script also

determines the genotype frequencies (0/0, 0/1 and 1/1) at the corresponding

position in sequenced individuals outside of the pedigree that are not

descendants of the sire or the dam. The third module

("6_germline_assigner_4_gen.pl") combines the output of the first and second

module to determine in which individual the *dnm* is most likely to have occurred

(one of the four grand-parents, sire or dam, or proband) and on which grand-

parental chromosome it occurred. All candidate *dnm*'s were manually curated

using the Integrated Genome Viewer (IGV)[22]. *Dnm*'s with mutant reads in either

sire or dam (even if called 0/0 by GATK) were relabeled as SM or DM, provided

that the *dnm* segregated in complete (but imperfect) linkage with the paternal or

maternal haplotype (respectively), in the half-sibs of the proband.

To test the corresponding pipeline, we identified 11,255 variants (of which

10,093 SNPs) for which Pr2 was heterozygous and which were not present in

unrelated individuals including from the 1,000 Bulls project.   The corresponding

"genotype fields" of the parents and grand-offspring were modified in the vcf file

such that the genotype (GT) was set at 0/0 and the unfiltered allele depth (AD) of

the derived allele set at 0.   The Ti/Tv ratio for the corresponding SNPs was

1.9922.   The proportion of the genome explored was estimated at 76% as

described above.   The pipeline detected 9,325 variants (83% sensitivity) of

which 8,409 SNPS (81% sensitivity).   The Ti/Tv ratio amongst detected *dnm*'s

was 1.9989, indicating that the pipeline did not introduce a Ti/Tv bias.

Confirmation of candidate dnm's. PCR primers were then designed for each

candidate passing the quality check in IGV using BatchPrimer[23] targeting a

product size of 200-1000bp with at least one primer being present in unique

(non-repeat) sequence (as identified by repeatmasker). The resulting amplicons

were sequenced on an Illumina MiSeq instrument using the 2x250bp paired end

protocol. The sequenced amplicons were aligned to the BosTau6 reference

genome using BWA mem and candidate de novo mutations were checked in IGV

and variants were called using freebayes (v1.0.2-15-g357f175)[24].

Modeling gametogenesis.   (i) Data types: To compare the adequacy of the

different gametogenesis models we computed the likelihood of three types of

data.   The first is the degree of mosaicism in the parent across *dnm*'s detected in

a given gamete.   Thus we may have detected n SM and m SNM *dnm*'s in a given

sperm cell.   The n SM *dnm*'s have dosages in the paternal sperm DNA of x_1, x_2,

x_3, ... $x_n > 0$ while the m SNM *dnm*'s have a dosage of 0. We have three such lists

for Pr3, Pr4 and Pr5. Likewise, we may have detected n OM and m ONM *dnm*'s in

a given oocyte. The n OM *dnm*'s have dosages in the maternal blood DNA of x_1, x_2,

x_3, ... $x_n > 0$ while the m ONM *dnm*'s have a dosage of 0. We have four such lists

for Pr2, Pr3, Pr4 and Pr5.

The second data set consists in lists of PM *dnm*'s and their dosage (x_1, x_2, x_3, ... x_n

> 0) in sperm (Pr1 and Pr2) or blood DNA (Pr3, Pr4, Pr5).

The third data type consists in the number of *dnm*'s detected in a gamete

transmitted to a proband, and the numbers of those shared by the studied half-

sibs of the proband. Thus, we may have detected n SM and m SNM *dnm*'s in a

sperm cell (or oocyte) transmitted to a given proband, of which half-sib 1 will

share x_1, half-sib 2 x_2, ..., where $x_i \le n$.

(ii) Computing probabilities under various models of gametogenesis: For data

type 1, we simulated the process of de novo mutation in the female and male cell

lineages described in Suppl. Figure 6.   For the null hypothesis, the mutation rate

per cell division before birth was set at an average of 0.77 (Poisson distributed),

such that the number of *dnm*'s per oocyte averaged 14 (as observed).   The

mutation rate per cell division after birth was set at an average of 0.3 (Poisson

distributed), such that the number of *dnm*'s per sperm cell averaged 34 (as

observed).   For alternative hypotheses, the mutation rate for the early cleavage

cell divisions (4, 7, 11, 15 and 18 first cell divisions, corresponding to the

different development stages in Suppl. Figure 6) was increased 10- or 20-fold

when compared to the remaining prenatal cell divisions, for which the mutation

rate was concomitantly reduced such that the overall number of *dnm*'s per

oocyte remained unaffected (average of 14).   We further tested 4, 10 and 40

induced PGCs, and unrelated or related induced PGCs as described in Suppl. Fig.

6. For all 90 possible scenarios, we determined by simulation what proportion of

dnm's found in a sperm cell (respectively oocyte) were characterized by a dosage

in paternal sperm (respectively maternal soma) of 0-0.05, 0.05-0.10, ...   These

proportions were then used as probabilities in computing the likelihood of the

data (i.e. a series *dnm*'s with corresponding rate of mosaicism in the parental

tissue) under the corresponding model.   Given our experimental design, SM and

DM are only recognized as such, if (i) their dosage in the sire (SM) or dam (DM)

is significantly < 0.5, and (ii) they show complete (but imperfect) linkage in the

available half-sibs.   We considered a fixed number of eight half-sibs in the

simulations. These conditions were included in the simulations. Thus a mutation

was only considered if it satisfied these two criteria.   The dosage that was

considered was not the true dosage for that mutation, but the "realized" dosage

assuming a sequence depth of 24.

To compute the likelihood of the second type of data, we simulated

gametogenesis in exactly the same way as for data type 1. We then randomly

sampled *n* gametes, where *n* corresponds to the number of GO (hence 5 for Pr1,

Pr3-5, and 11 for Pr2).   For all *dnm*'s in these *n* gametes, we then determined the

dosage in the germ-line (Pr1, Pr2) or soma (Pr3-5).   For all 90 possible scenarios,

we determined by simulation what proportion of PM *dnm*'s detected in sperm of

blood DNA were characterized by a dosage of 0-0.05, 0.05-0.10, ...   These

proportions were then used as probabilities in computing the likelihood of the

data (i.e. a series PM *dnm*'s with corresponding dosage in sperm or blood) under

the corresponding model. With the real data, PM mutations are only recognized

as such (i) if they are transmitted to at least one of the *n* offspring, (ii) if the

proband is called heterozygous for the corresponding *dnm* by GATK, and (iii) if

we demonstrate complete (but imperfect) linkage in the GO.   Condition (i) is

achieved in the simulation by sampling *n* gametes at random. Condition (ii) and

(iii) were modeled in the simulations. We considered five and eleven GO to

match the real data. Thus a mutation was only considered if it satisfied these

two criteria. The dosage that was considered was not the true dosage for that

mutation, but the "realized" dosage assuming a sequence depth of 24.

For data type 3, we modified the simulations in order to exactly generate a

predetermined number *n* of *dnm*'s in a given "reference" gamete. Thus if in the

real data an oocyte was characterized by 11 *dnm*'s (f.i. Pr3 and Pr4), we would in

the simulations distribute 11 *dnm*'s across the (7+4+4+3+18) cell divisions

leading to a simulated reference oocyte and track their segregation (according to

their point of occurrence) across the entire germ line lineage. Under the null

hypothesis of uniform prenatal mutation rate, all 36 cell divisions would have

equal chance to be hit by anyone of the 11 mutations. Under the alternative

hypotheses, early cleavage cell divisions would have a 10- or 20-fold higher

chance than the remaining ones. Under the hypothesis of related PGCs the

segregation pattern of early mutations in the germ line lineage would be

concomitantly affected (see Suppl. Fig. 6). We would then samples gametes at

random from the same germ line tree and count the number of mutations shared

with the "reference" gamete. This would generate a frequency distribution of

gametes sharing 0, 1, 2, ... , *n* *dnm*'s with the reference gamete.   The

corresponding frequencies were then used as probabilities in computing the

likelihood of the data (i.e. a series half-sibs sharing 0, 1, … , *n dnm*'s with the

reference gamete transmitted to the proband) under the corresponding model.

Likelihoods of the data under the 90 tested models were then simply computed

as the product of the probabilities of all *dnm*'s (data type 1 and 2) and half-sibs

(data type 3) extracted from the simulations performed under the corresponding

model.

(iii) Estimating the number of missed SM and DM and misclassified PM

mutations: The simulations for dataset 1 allowed us to estimate the number of

SM and DM mutations missed either because the "realized" dosage was to high in

the parent, or because we could not demonstrate complete (but imperfect)

linkage in the half-sibs. Likewise the simulations for dataset 2 allowed us to

estimate the number of PM mutations that, although detected (realized dosage

sufficient to be called heterozygous by GATK), were misclassified as SNM or DNM

mutations because showing perfect linkage in the available GO.   Under the best

biological model (20x increased mutation rate during the first 4 cell divisions, 4

related PGCs, see Table 1), these numbers were: (i) average loss of 3.3 SM

mutations, (ii) average loss of 1.1 lost DM mutations, (iii) average gain of 1.45

SNM and 1.45 DNM mutations for a male proband with 11 GO, (iv) average gain of

4 SNM and 4 DNM mutations for a male proband with 5 GO, and (v) average gain

of 1.8 SNM and 1.8 DNM mutations for a female proband with 5 GO.

20. Li H & Durbin R. Fast and accurate short read alignment with Burrows-

Wheeler transform. *Bioinformatics* **25:** 1754-1760 (2009).

21. McKenna A *et al.* The Genome Analysis Toolkit: a MapReduce framework for

analyzing next-generation DNA sequencing data. *Genome Res.* **20:** 1297-1303

(2010).

22.Robinson JT et al. Integrative Genomics Viewer. *Nature Biotechnology* **29**:24–

26 (2011).

23. You FM *et al.* BatchPrimer3: a high throughput web application for PCR and

sequencing primer design. *BMC Bioinformatics* **9:**253 (2008).

24. Garrison E, Marth G. Haplotype-based variant detection from short-read

sequencing. arXiv preprint arXiv:1207.3907 [q-bio.GN]. (2012) .

588