Structure, culture, and predicted function of the gut microbiome of the Mormon cricket

Anabrus simplex **(Orthoptera: Tettigoniidae)**

Chad C. Smith[1], Robert B. Srygley[2], Frank Healy[3] Karthikeyan Swaminath[1], Ulrich G. Mueller[1]

[1]Department of Integrative Biology, University of Texas at Austin, 1 University Station C0990,

Austin, TX 78724

[2]USDA-Agricultural Research Service, Northern Plains Area Research Laboratory, 1500 Central

Avenue, Sidney MT 59270

[3]Department of Biology, Trinity University, 1 Trinity Pl., San Antonio, TX 78212

Corresponding author: Chad C. Smith

Email: chad.smith123@gmail.com

Header: Structure of the Mormon cricket gut microbiome

SUMMARY

1. The gut microbiome of insects plays an important role in their ecology and evolution, participating in nutrient acquisition, immunity, and behavior. Microbial community structure within the gut is heavily influenced by differences among gut regions in morphology and physiology, which determine the niches available for microbes to colonize.

2. We present a high-resolution analysis of the structure of the gut microbiome in the Mormon cricket *Anabrus simplex,* an insect known for its periodic outbreaks in the western United States and nutrition-dependent mating system. The Mormon cricket microbiome was dominated by eleven taxa from the Lactobacillaceae, Enterobacteriaceae, and Streptococcaeae. While most of these were represented in all gut regions, there were marked differences in their relative abundance, with lactic-acid bacteria (Lactobacillaceae) more common in the foregut and midgut and enteric (Enterobacteriaceae) bacteria more common in the hindgut.

3. Differences in community structure were driven by variation in the relative prevalence of three groups: a *Lactobacillus* in the foregut, *Pediococcus* lactic-acid bacteria in the midgut, and *Pantoea agglomerans*, an enteric bacterium, in the hindgut. These taxa have been shown to have beneficial effects on their hosts in insects and other animals by improving nutrition, increasing resistance to pathogens, and modulating social behavior.

4. Using PICRUSt to predict gene content from our 16S rRNA sequences, we found enzymes that participate in carbohydrate metabolism and pathogen defense in other orthopterans. These were predominately represented in the hindgut and midgut, the most important sites for nutrition and pathogen defense.

5. Phylogenetic analysis of 16S rRNA sequences from cultured isolates indicated low levels of divergence from sequences derived from plants and other insects, suggesting that these bacteria are likely to be exchanged between Mormon crickets and the environment.

6. Our study shows strong spatial variation in microbiome community structure, which influences predicted gene content and thus the potential of the microbiome to influence host function.

Keywords: symbiosis, alimentary tract, immunity, nutrition, PICRUSt, carbohydrate metabolism, lactic acid bacteria, Enterobacteriaceae, katydid,bacteria, metagenomic

INTRODUCTION

Insects are the most speciose and abundant taxa in the animal kingdom, playing a key ecological role in many of the world's ecosystems. Symbioses between insects and their microbial associates has undoubtedly contributed to their success, providing the capability to degrade recalcitrant food, to supplement nutrient-deficient diets, to protect them from their natural enemies, and to modulate the expression of social behavior (Engel & Moran 2013; Douglas 2015). Among the niches available to occupy the host, the gut houses the largest and most diverse microbiome in insects (Engel & Moran 2013; Douglas 2015) and other animals (Ley *et al.* 2008; Cho & Blaser 2012). Gut morphology and physiology vary markedly along the alimentary tract, resulting in an environmental gradient that influences, and is influenced by, the microbial communities that populate it (Dillon & Dillon 2004; Engel & Moran 2013).

The insect gut consists of three regions that are analogous to that in mammals, the foregut, the midgut and the hindgut, each of which contributes to a different aspect of gut function (Douglas 2013). The foregut serves as the entry point for food, where it is stored in the crop before passing through the proventriculus, a valve that can also be modified to mechanically filter food (Woodring & Lorenz 2007; Douglas 2013) and even microbes (Lanan *et al.* 2016). Digestion and absorption of nutrients begins at the midgut, which, in some species, contains specialized crypts that house microbes that aid in insect nutrition (Kikuchi, Meng & Fukatsu 2005; Bistolas *et al.* 2014). Host immune factors have been shown to play an important role in regulation of commensal microbes in the midgut (Ryu, Ha & Lee 2010; Buchon, Broderick & Lemaitre 2013), some of which protect the host from pathogens (Forsgren *et al.* 2010). Following the midgut is the hindgut, which is comprised of the ileum, colon, and rectum. Malphigian tubules permeate the anterior hindgut, excreting nitrogenous waste and other solutes

from the hemocoel that can provide nutrients for dense populations of microbes (Bignell 1984).

In some species, dense bristle-like structures in the ileum (Woodring & Lorenz 2007) and rectal

papillae (Hunt & Charnley 1981) provide attachment sites for bacteria, some of which fix

nitrogen (Tai *et al.* 2016), degrade recalcitrant plant polymers (Kaufman & Klug 1991; Engel &

Moran 2013), and prevent infection (Dillon & Charnley 2002).

The Mormon crickets *Anabrus simplex* (Orthoptera: Tettigoniidae) is an economically

important shield-backed katydid distributed throughout the Western United States. Mormon

crickets can form dense aggregations of millions of individuals spread over 10 kilometers long

and several kilometers wide, feeding on forbes, grasses, and agricultural crops as they march in

migratory bands across the landscape (MacVean 1987; Simpson *et al.* 2006). Mormon crickets

are also emerging as a model for the study of how social interactions and diet influence

immunity (Srygley *et al.* 2009; Srygley & Lorch 2011) and the microbiome (Smith *et al.* 2016).

Differences in population density are linked to reproductive behavior, as in high density

populations, protein-limited females compete for access to males to gain access to a

proteinaceous "nuptial gift" males produce for females during copulation (Gwynne 1984). While

consumption of male nuptial gifts by females does not influence the composition of the

microbiome, sexually inactive females experience a dramatic decline in *Pediococcus* lactic-acid

gut bacteria compared to sexually active females (Smith *et al.* 2016). Lactic-acid bacteria are

common associates of the alimentary tract and regarded for their beneficial effects on immune

function and nutrition in animals, including insects (Forsgren *et al.* 2010; Storelli *et al.* 2011;

Erkosar *et al.* 2015).

We characterize the structure of the gut microbiome of Mormon crickets and infer their

evolutionary relationships using a combination of culture-dependent and culture-independent

approaches. Our aims are to determine whether gut microbial communities vary along the

alimentary tract in the Mormon cricket and to infer their potential to influence host function

based on their known taxonomic associations with other insects and by employing bioinformatic

tools that predict metabolic capabilities from 16S rRNA sequences. We also establish methods

for isolating Mormon cricket gut microbiota in culture to permit future experimental

manipulations of the gut microbiome and build genomic resources to infer their evolution and

function.

<u>MATERIALS AND METHODS</u>

Animal collection and tissue processing

Mormon crickets were obtained from field (n=5) and laboratory-raised (n=8) collections. Wild

females were caught in EK Mountain (43°47'58"N, 106°50'31"W, 1752 m) near Kaycee,

Wyoming in the summer of 2014, immediately preserved in 100% ethanol, and stored at -80°C

until dissection. Laboratory-raised Mormon crickets were derived from eggs collected from

individuals caught on Paint Rock Road (44°27'52"N, 107°27'37"W, 2654 m) in the Bighorn

Mountains and fed a mixture of wheat bran, wheat germ, sunflower, mixed bird seeds, tropical

fish flakes, fresh Romaine lettuce (added daily), and water *ad libitum*.

Mormon crickets were dissected using flame-sterilized tools after rinsing in 1% bleach

for 3 min followed by two rinses in autoclaved distilled water to remove bacteria on the

exoskeleton. DNA from the foregut (crop and proventriculus), midgut (ventriculus), ileum, and

rectum (Fig. 1) of laboratory-raised crickets was extracted with MoBio Powersoil$^{©}$ as in Smith et

al. (2016). Foregut (crop and proventriculus), midgut (ventriculus), and hindgut tissue (ileum and

rectum combined) was extracted from field-collected animals using a bead-beating/phenol-

chloroform extraction protocol (see supplementary material). DNA extraction methods can

influence the representation of taxa in 16S rRNA metagenomic studies (Yuan *et al.* 2012),

however our aim here is not to make inferences about differences between field and laboratory-

raised animals but differences among tissue types. We include the source of the animal (field or

laboratory) and as a covariate in our statistical analyses to account for variation due to

source/DNA extraction method (see Statistics).

Sequencing and Bioinformatics

The variable V4 region of 16S rRNA gene was amplified with universal primers (Hyb515F: 5'-

GTGYCAGCMGCCGCGGTA -3', Hyb806R: 5'-GGACTACHVGGGTWTCTAAT-3') and

sequenced on the Illumina Miseq V3 platform. DADA2 1.1.5 (Callahan *et al.* 2016) was used to

process the raw sequencing data and taxonomy was assigned with the Greengenes 13.8 database

at 97% identity (see supplementary material). Sequence variants that comprised an average of

less than 1% of the reads recovered within a given Mormon cricket were removed prior to

analysis using phyloseq 1.16.2 (McMurdie & Holmes 2013).

Metagenomic predictions

We used PICRUSt (Phylogenetic Investigation of Communities by Reconstruction of

Unobserved States) v1.1.0 (Langille *et al.* 2013) to estimate the functional gene content of our

samples. PICRUSt generates metagenomic predictions from 16S rRNA data using annotations of

sequenced genomes in the IMG database. Nearest sequenced taxon index (NSTI) values were

small (mean $\pm$ sd: 0.03 ± 0.01, range: 0.006-0.040), indicating the taxa in our samples were

closely related to the genomes in the IMG database (see supplementary material). Greengenes

IDs used by PICRUSt to construct the phylogenetic tree were assigned to sequence variants

using Qiime 1.9 (Caporaso *et al.* 2010), and the Kyoto Encyclopedia of Genes and Genomes

(KEGG) database was used for functional classification.

Bacterial Abundance

The abundance of bacteria was estimated using qPCR following Powell et. al (2014) from

laboratory-raised (n=8) and field-caught (n=8) Mormon crickets (see supplementary

information). Universal 16S rRNA gene primers 27F (5'-AGAGTTTGATCCTGGCTCAG-3')

and 355R (5'-CTGCTGCCTCCCGTAGGAGT-3') were used to amplify all copies of the 16S

rRNA gene in tissue samples from laboratory (n=8) and field caught individuals (n=8) and copy

number quantified with standard curves from the cloned target sequence (Powell *et al.* 2014)

Culturing and phylogenetic analysis

Five lab-reared female Mormon crickets were surface sterilized in 1% bleach for three min,

rinsed twice in sterile water and dissected using flame-sterilized tools. Gut tissue was

homogenized for 10 s with a bead beater using autoclaved 3.2mm stainless steel beads in sterile

PBS. Homogenates were plated onto trypsin soy agar, brain heart infusion agar, nutrient agar, or

Man–Rogosa–Sharpe agar (BD), cultured in anaerobic or Campy (low O_2) Gaspak pouches

(Becton, Dickinson and Company, Franklin Lakes, NJ) at 37°C for 24-48 hours, and individual

colonies passaged three times to obtain pure isolates. DNA was then extracted with chelex and

the 16S gene amplified for Sanger sequencing using 27F (5'-AGAGTTTGATCCTGGCTCAG-

3') and 1492R (5'-GGTTACCTTGTTACGACTT-3') primers (see supplementary material).

Phenotypic assays

Fresh overnight cultures of all isolates were used for microscopic analysis. Lactobacillaceae

isolates were cultured in Man–Rogosa–Sharpe medium and Enterobacteriaceae were cultured in

nutrient broth or LB medium. Biochemical tests were done following Bridson (1998). Motility

was determined using SIM medium and microscopic examination of culture wet mounts. Man–

Rogosa–Sharpe or nutrient broth containing 1 g/L potassium nitrate was used for nitrate

reduction tests. Fermentation tests were done anaerobically in Man–Rogosa–Sharpe and nutrient

broth media with the addition of indicated sugars to 1% w/v final concentration.

Statistics

Analyses were performed in R 3.3.1 (R Core Development Team 2013). Sequence tables were

rarified at 1300 reads using phyloseq (McMurdie & Holmes 2013), resulting in the exclusion of

hindgut samples from two field-caught females that had a low number of reads. Alpha diversity

was compared among tissue types and between origin of subject (field vs. lab) with a linear

mixed model (Bates *et al.* 2013), entering the individual ID as a random effect to account for

within-subject correlations in diversity. Species richness, Chao1, and the Shannon-Weiner

diversity index were calculated. Post-hoc comparisons among gut regions were performed using

a Tukey test (Hothorn, Bretz & Westfall 2008).

Beta diversity among gut tissue types and between animal source (field vs. lab) was

assessed with a distance-based redundancy analysis (db-RDA) in vegan 2.3 (Oksanen *et al.*

2015), specifying a principal components ordination of Bray-Curtis distances. Statistical

significance of the terms in the db-RDA model was determined by 999 permutations of the

distance matrix in vegan, restricting the permutations to within each individual to retain the

nested structure of the data. The same procedure was also used to examine variation among

tissue types in the abundance of KEGG pathways, except nonmetric multidimensional scaling

was used for ordination.

We assessed the difference in taxon abundance among tissue types in univariate analyses

by fitting the data to a negative binomial generalized linear mixed model (Bates *et al.* 2013),

specifying the individual ID as the random effect and the tissue type and animal source (field vs.

lab) as fixed effects. A similar procedure was used to assess differences in 16S rRNA gene copy

number between tissue types and animal source, except a normal distribution was specified.

Likelihood ratio tests were used to determine the statistical significance of each factor (Venables

& Ripley 2002). Goodness-of-fit was assessed by with a Chi-square test (Faraway 2006) and

homoscedasticity was assessed by examination of residual plots. Nonparametric methods were

used in univariate analyses of the metagenomic predictions because no distribution provided a

reasonable fit to the data. P-values were adjusted for multiple tests using the false discovery rate

(Benjamini & Hochberg 1995).

RESULTS

Spatial structure of the gut microbiome

We recovered 11 dominant sequence variants from field and lab-raised individuals (Fig. 2), with

the remaining 749 sequence variants comprising <1% of the sequences from a given Mormon

cricket. Field and laboratory-raised individuals shared 7 of the 11 sequence variants, including

the most abundant *Pediococcus acidilactici* OTU that varied with mating status in a previous

study (*P. acidilactici* 102222; Smith *et al.* 2016). The remaining five shared sequence variants

were two Lactobacilliaceae (*Lactobacillus sp* and *P. acidilactici* 2), two Enterobacteriaceae

(*Pantoea agglomerans* and a *Klebsiella sp*) and one Streptococcaceae (*Lactococcus garvieae*).

Field-caught Mormon crickets had three taxa that were not shared with laboratory-raised

individuals, while lab-raised individuals had two taxa that were not shared with field individuals

(Fig. 2). Guts from two laboratory individuals were almost completely comprised of the enteric

bacterium *Pantoea agglomerans* (99.3% and 80.8% of reads respectively), so we conducted our

analysis with and without these individuals.

Species richness and diversity differed among gut regions and were higher in field

compared to lab-raised animals (Table 1, Fig. 3). There was no significant interaction between

collection source and tissue type (Table 1), indicating that differences in alpha diversity among

tissue types were shared between lab and field-caught animals. We found that the midgut was the

most diverse part of the gut with two of the three measures of alpha diversity (species richness

and the Chao1 diversity estimator), while the hindgut and foregut had similar levels of richness

and diversity. The third metric (Shannon-Weiner) also found the foregut to be the least diverse

region, but differed in that the midgut and hindgut had similar levels of species diversity (Table

1, Fig 3).

The db-RDA analysis revealed that the structure of the gut microbiome also varied

among gut regions and between field and laboratory animals (Table 2, Fig. 4a). The non-

significant interaction in this analysis, however, indicates that the differences in community

structure among tissue types were consistent between field and laboratory-raised individuals

(Table 2). To determine which members of the gut microbiome varied among gut regions, we

plotted the taxa scores from db-RDA analyses of field and laboratory Mormon crickets (Figure

S1a). Three groups of bacteria appeared to separate along the gut axis: a *Lactobacillus sp.* lactic-

acid bacterium associated with the foregut, *Pediococcus* lactic-acid bacteria were associated with

the midgut, and *Pantoea agglomerans*, an enteric bacterium, was found in association with the

hindgut. Inspection of the plots from laboratory animals, where the ileum and rectum of the

hindgut were dissected separately, indicate that *P. agglomerans* is more abundant in the rectum,

while the composition of the ileum, which is separated from the rectum by the colon, closely

resembled that of the midgut (Figure S1b).

Univariate analyses of these three groups largely confirmed the pattern in the ordination

(Table 3, Fig. 2, Fig. S2). The interaction between tissue type and source was not significant in

any of the analyses and dropped to estimate the differences in abundance between tissue types.

Lactobacillus sp. was three times more common in the foregut than in the midgut (β=1.4 $\pm$ 0.50,

p=0.02) and seven times more abundant in the foregut than in the hindgut (β=2.0 $\pm$ 0.51,

p<0.001). *Pediococcus* were similar in abundance in the midgut and hindgut but 4.7 times more

common in these areas than the foregut (β=1.1 $\pm$ 0.36, p=0.006). *P. agglomerans* was 209 times

more abundant in the hindgut than in the foregut (β=3.8 $\pm$ 0.87, p<0.001) and twelve times more

abundant in the hindgut than in the midgut (β=2.5 $\pm$ 0.82, p=0.007).

Abundance of bacteria from 16S rRNA qPCR

The number of copies of bacterial 16S rRNA genes was significantly different among tissue

types, as indicated by the significant interaction between tissue type and the source of the

Mormon crickets (Analysis of deviance: Source, $F_{1,14}$=25.9, p<0.001; tissue type, $F_{3,161}$=7.8,

p<0.001; Interaction, $F_{3,161}$=2.8, p=0.04, Fig. 7). We decomposed the interaction to determine

how the total number of 16S rRNA copies differed among tissue types within field and

laboratory-raised animals. The major difference between the two sources was that in wild

Mormon crickets, the midgut had the lowest abundance of all gut regions, while in laboratory-

raised individuals, both the midgut and the ileum had the lowest abundance of bacterial 16S

rRNA genes (Table S1, Fig. 5).

PICRUSt metagenomic predictions

PICRUSt analysis of 16S rRNA sequence variants recovered 5,891 KEGG orthologs associated

with 328 metabolic pathways. The representation of the predicted KEGG pathways differed

significantly among gut regions in both the full and reduced datasets, while the source of the

animals had a significant influence in the full dataset but not the reduced dataset (Table 2, Fig.

4b). Neither analysis, however, showed an interaction between tissue type and whether an animal

was wild or lab-reared, indicating that metagenomic predictions differed among tissue types in

similar ways (Table 2). Univariate analyses found significant differences among tissue types in

most KEGG pathways (Table S2), including those that could affect host-microbe interactions via

their role in nutrition, immunity, degradation of xenobiotics, and production of secondary

metabolites (Fig. 6). In these functional groups, the hindgut exhibited the most abundant

representation of each KEGG category, followed by the midgut and then the foregut (Fig. 6).

<u>Nutrition</u>

We searched our metagenomic predictions for specific bacterial genes known to

contribute to host nutrition in orthopterans. We queried our database for enzymes capable of

metabolizing the complex plant carbohydrates xylan, pectin, raffinose, and galactomannan,

which are metabolized by gut bacteria in the house cricket *Achetus domesticus* (Kaufman & Klug

1991), and cellulose, an important component of the plant cell wall. We found KEGG orthologs

involved in the metabolism of all these complex plant polymers, except galactomannan. The

pectin metabolic pathway was also incomplete. Only pectinesterase, the first of three enzymes

involved in pectin metabolism, was found among the eleven dominant taxa in Mormon crickets,

although the remaining two enzymes were represented in the minority members (i.e. <1% of 16S

rRNA sequences, see *Sequencing and Bionformatics*) of the gut microbiome.

The abundance of KEGG orthologs for carbohydrate metabolism in our samples were

most pronounced in the hindgut (Table S3, Fig. 7a) and dominated by the enteric bacteria,

particularly *Klebsiella sp*. and Enterobacteriaceae 1 (Fig 7b). Lactic-acid bacteria, however, were

also represented in predictions for raffinose metabolism and enzymes capable of participating in

the degradation of cellubiose to glucose via cellubiose glucohydrolase, but not in degrading

cellulose to cellubiose (Fig 7b).

Gut bacteria might also play a role in the production of the essential amino acid

phenylalanine via the shikimate pathway, which is found in microbes and plants but not in

animals (Herrmann & Weaver 1999). Phenylalanine is required for stabilization of the cuticle

following molting (Bernays & Woodhead 1984) and is converted to tyrosine, the precursor of

melanin, a key component of the insect immune response (González-Santoyo & Córdoba-

Aguilar 2012). All enzymes in the shikimate pathway were represented in our metagenomic

predictions (Fig. S3), although prephenate hydrogenase, which is required for phenylalanine

synthesis, was only represented in *Lactococcus garviae. L. garviae* abundance thus might

influence the availability of phenylalanine for Mormon crickets, unless they are able to acquire it

in sufficient quantities directly from their diet.

<u>Immunity</u>

In the locust *Schistocera gregaria* (Orthoptera)*,* four phenols have been shown to

increase resistance to microbial pathogens (Dillon & Charnley 1988, 1995): hydroquinone, 3,4-

dihydroxybenzoic acid, p-hydroxybenzoic acid, and 4,5-dihydroxybenzoic acid. We found

enzymes associated with the production of all these compounds except for 4,5-dihydroxybenzoic

acid, which was not annotated in the KEGG database. Hydroquinone production was represented

by the enzyme arbutin 6-phosphate glucohydrolase, which metabolizes arbutin, a phenolic

glycoside present in leaf and fruit tissue of many plants (Xu *et al.* 2015).

Two enzymes were found capable of producing 3,4-dihydroxybenzoic acid. The first,

vanillate monooxygenase, demethylates vanillic acid, a compound derived from lignin (Bugg *et*

al. 2011). This is also the pathway proposed for 3,4-dihydroxybenzoic acid production in locusts

based on the abundance of vanillic acid in their feces (Dillon & Charnley 1988, 1995). The

second, p-hydroxybenzoate 3-monooxygenase, oxidizes p-hydroxybenzoic acid, one of the other

antimicrobial phenols in locusts (Dillon & Charnley 1995). The most likely source of p-

hydroxybenzoic acid in the diet of Mormon crickets is benzoic acid, which is a precursor to

salicyclic acid in plants (Raskin 1992). The enzyme responsible for catalyzing the conversion of

benzoic acid to p-hydroxybenzoic acid (benzoate 4-monooxygenase), however, was not found

among the 11 dominant taxa in our samples, although it was present in the minority members of

the Mormon cricket gut microbiome. Production of p-hydroxybenzoic acid in appreciable

concentrations is thus less likely than for hydroquinone or 3,4-dihydroxybenzoic acid.

Like carbohydrate metabolism, the hindgut (Fig. 7c) and enteric bacteria (Fig. 7d)

dominated the abundance of KEGG orthologs implicated in the production of antimicrobial

phenols in our samples, with the exception of hydroquinone, which was represented to varying

degrees among the lactic-acid bacteria. Notably, *P. agglomerans,* which has been reported to

participate in the production of 3,4-dihydroxybenzoic acid in locusts (Dillon & Charnley 1995),

was not among taxa responsible for the occurrence of vanillate monooxygenase in our samples

(Fig 7d).

Finally, we searched for three other known contributors to pathogen defense: bacterocins,

antibiotics, and lactate dehydrogenase, which provides protection from pathogens in the gut by

reducing pH (Servin 2004). We found lactate dehydrogenase to be equally represented among

gut regions (Fig 7c), and lactic-acid bacteria were the main contributors to our samples (Fig. 7d).

We found three bacteriocins in the KEGG database: nisin, mutacin, and *blp*-derived bacterocins.

None of these were found in our metagenomics predictions, perhaps not surprising considering

their association with *Streptococcus,* which was not among the top 11 taxa in our samples (Fig.

2). The bacteriocins we would expect to find based on taxonomy (e.g. pediocin for *Pediococcus*)

were not annotated in the KEGG database.

Turning to the antibiotics, we found enzymes involved in the production of streptomycin,

penicillin, and novobiocin, but not all enzymes required for their synthesis were present (data not

shown). We did find β-lactamase, which confers resistance to β-lactam antibiotics (e.g.

penicillins, cephalosporins, monobactams, and carbapenems; Drawz & Bonomo 2010),

represented among all the Enterobacteriaceae and *Pediococcus* taxa in our samples, but not

among *Lactobacillis sp.* other Lactobacillaceae (data not shown). This suggests that lactate

production and antibiotic resistance could play a role in microbe-microbe interactions in the

Mormon cricket gut microbiome.

Phylogenetic analysis of cultured isolates

Thirteen strains were cultured from the Mormon cricket gut based on 99% sequence similarity of

their near full-length 16S rRNA genes (mean $\pm$ sd: 1406 $\pm$ 30bp). Six were lactic-acid bacteria

(Lactobacillaceae) and seven were enteric bacteria (Enterobacteriaceae).

The lactic-acid bacteria fell into two clades in our phylogenetic analysis (Fig. 8). The first

clade was comprised of *Pediococcus acidilactici* isolates derived from environmental sources,

such as plants and various human foodstuffs, as well as strains from the human gut. Similarity to

sequences from the BLAST search was high (>99.5%) and branch lengths were short, indicating

that *Pediococcus* from the Mormon cricket gut are not highly derived from their relatives, as has

been found for *Lactobacillus* species isolated from bees (Fig. 8; McFrederick *et al.* 2013).

Our search for *Pediococcus* sequences from insect guts in Genbank recovered sequences

from the termites *Macrotermes bellicosus* and *M. subhyalinus*, which formed their own well-

supported clade (Fig 8). Cultured *Pediococcus acidilactici* shared 100% sequence identity in the

V4 region with the *P. acidilactici 1* phylotype sequenced using the Illumina platform in this

study and with the *P. acidilactici* (102222) phylotype associated with variation in mating status

in Mormon crickets (Smith *et al.* 2016). Morphologically, *Pediococcus acidilactici* were

nonmotile and spherical (0.8 – 1.0 μm), often dividing to form pairs as described for other

Pediococcus. As other members of the genus, the *P. acidilactici* were gram-positive, non-motile,

facultatively anaerobic, grow at low pH, and produce lactate from lactose (Table S2).

The second clade of lactic-acid bacteria was comprised primarily of plant-associated

Lactobacillus. Unlike *P. acidilactici*, these *Lactobacillus* formed a distinct clade with good

branch support (Fig 8), indicating it is genetically distinct enough at the 16S rRNA locus to

distinguish itself from other clades in the phylogeny. Similar to *P. acidilacitici*, these

Lactobacillus had high sequence similarity (>99.5%) to other members of the clade and a short

branch length, indicating that while it is distinct enough to form its own clade, it is not highly

derived from its relatives at the 16S rRNA locus.

Our Genbank search for *Lactobacillus* isolated from insect guts found sequences from

ants, bees, and termites, and fruit flies, all of which fell into a different clade than *Lactobacillus*

isolated from Mormon crickets. *Lactobacillus* from these taxa thus appear to have a different

evolutionary history. *Lactobacillus* isolates shared 100% sequence identity in the V4 region with

the Lactobacillaceae 2 phylotype sequenced using the Illumina platform in this study.

Morphologically, these *Lactobacillus* appear as non-motile straight rods, approximately 1.3-2

μm in length and 0.8-1.0 μm wide and are gram-positive, non-motile, facultatively anaerobic,

grow at low pH, and produce lactate from lactose (Table S2).

The seven Enterobacteriaceae strains were most similar to *Enterobacter* strains in our

BLAST search, which recovered sequences from a variety of plant and animal sources (sequence

similarity=98.7-99.8%). Our survey of Genbank found *Enterobacter* from alimentary tracts of a

diverse group of insects, including termites, cockroaches, flies, beetles, stink bugs, bees, ants,

and moths. Like other studies (Brenner *et al.* 2005), however, the 16S rRNA gene did not have

enough signal to resolve relationships among *Enterobacter* and its relatives (data not shown) so

we present a simpler phylogeny with the Mormon cricket isolates and type strains from the

family (Fig. 9).

We found that our Mormon cricket isolates were interspersed with *Enterobacter*,

Klebsiella, and *Escherichia* type strains. A multilocus sequencing approach is thus needed to

improve the inference (Brenner *et al.* 2005). All seven strains isolated from Mormon crickets had

100% identity at the V4 region with the *Klebsiella* phylotype sequenced on the Illumina

platform, however the phylogenetic (Fig. 9) and phenotypic data (Table S2) suggest that

Klebsiella is unlikely to be a correct taxonomic assignment. Unlike most *Klebsiella*, cultured

strains were motile, which is more typical of *Enterobacter* and other Enterobacteriaceae

(Brenner *et al.* 2005). Morphologically, all isolates were straight rods, approximately 0.8-1.0 µm

in length and 0.6-0.8 µm wide. Strains were gram-negative and facultatively anaerobic (Table

S2).

<u>DISCUSSION</u>

We found striking differences in the diversity and structure of the gut microbiome in the

Mormon cricket *Anabrus simplex*. While most taxa were represented in the foregut, midgut and

hindgut, there were dramatic differences in relative abundance within the Lactobacillaceae and

between the Lactobacillaceae and Enterobacteriaceae, the main families recovered in our culture

and culture-independent studies. Predictions of their metabolic capabilities using PICRUSt

suggest the potential for these gut bacteria to participate in the metabolism of complex

carbohydrates and defense against microbial pathogens, particularly among the enteric bacteria

in the midgut and hindgut, and to a lesser extent, the lactic-acid bacteria. Finally, our

phylogenetic analysis of cultured isolates found that Mormon cricket gut bacteria are not highly

derived from related bacteria associated with plants or the guts of other animals, suggesting that

gut bacteria are either acquired from the environment in each generation or have not been

restricted to Mormon crickets over appreciable periods of evolutionary time. Our findings have

important implications for our understanding of the ecological and evolutionary processes that

influence the assembly and function of gut microbial communities, as it suggests that host-

microbe and microbe-microbe interactions shape the abundance and distribution of the gut

microbiome.

Our finding that bacterial abundance is lower in the midgut is in agreement with reports

from other orthopterans (Ulrich, Buthala & Klug 1981; Hunt & Charnley 1981) and insects

(Köhler *et al.* 2012), and has been attributed to characteristics that make the midgut less

hospitable to bacteria than other regions of the alimentary tract (Douglas 2015). The midgut in

insects secretes a host of digestive enzymes, is immunologically active, and lined by the

peritrophic membrane, which acts as a protective barrier that restricts microbes to the lumen and

protects the epithelium (Douglas 2015). In the two orthopterans that have been studied in detail,

bacteria are found in the midgut lumen but not in association with the epithelium (Hunt &

Charnley 1981; Mead, Khachatourians & Jones 1988). As a consequence, midgut bacteria might

need to be continually replenished from ingested food (Blum *et al.* 2013) because the peritrophic

membrane is continually shed into the hindgut. In some insects, specialized midgut crypts

provide niches that microbes colonize (Kikuchi *et al.* 2005; Bistolas *et al.* 2014), however we did

not observe analogous structures in Mormon crickets (Fig. 1).

The midgut is particularly vulnerable to pathogens because the lack of an endocuticle

leaves the epithelium exposed once the peritrophic membrane is penetrated (Lehane &

Billingsley 1996). The Mormon cricket midgut was populated by lactic-acid bacteria, with

Pediococcus specifically exhibiting greater abundance in the midgut (and hindgut) than in the

foregut. Lactic-acid bacteria are known for their beneficial effects in insects, increasing

resistance to parasites in bees (Forsgren *et al.* 2010) and promoting development in fruit flies by

enhancing proteolytic activity (Erkosar *et al.* 2015) and upregulating host ecdysone and insulin-

like peptides (Storelli *et al.* 2011). Lactic-acid bacteria are also known to suppress pathogenic

bacteria by reducing pH through the production of lactate and by producing a number of

antimicrobial compounds, such as hydrogen peroxide and bacteriocins (Cintas *et al.* 2001).

A previous study found that sexual interactions in Mormon crickets influences the

abundance of three *Pediococcus* phylotypes (Smith *et al.* 2016), however spatial information on

where in the gut *Pediococcus* is located has been unavailable until now. *Pediococcus* in the

midgut could provide immunological or nutritional benefits to Mormon crickets, as has been

shown for *P. acidilactici* in other animals (Castex *et al.* 2008, 2009). We found that the capacity

for lactate production in our samples was dominated by *Pediococcus* and other lactic-acid

bacteria, although the abundance of the enzyme mediating lactate production was not higher in

the midgut relative to other regions based on our metagenomics predictions. The cultured isolates

of *P. acidilactici* obtained from Mormon crickets in this study will enable future experimental

and comparative genomic approaches to evaluate these hypotheses.

Lactic-acid bacteria were also common in the foregut, which was dominated by a

Lactobacillus that averaged 73.9% of the sequences recovered from this region. Bignell (1984)

noted that the foregut of insects tends to be the most acidic compartment, however studies that

measure the physiochemical environment and characterize microbiome composition of the

foregut are rare (but see Köhler *et al.* 2012). This is because the endocuticle, lack of

differentiated cells for absorption of nutrients, and frequent purging of consumed material into

the midgut provides little opportunity for foregut microbes to contribute to host nutrition. The

large differences in community structure between the foregut and the rest of the alimentary tract

in our study does illustrate the dramatic transition in microbial communities between what is

ingested and what can colonize the more distal regions of the gut. Our metagenomics predictions

also suggest that the foregut is not the site of extensive carbohydrate metabolism or pathogen

defense for most of the pathways we examined.

In contrast to the foregut and midgut, the hindgut was characterized by a dramatic

increase in enteric bacteria (Enterobacteriaceae). Ordination of the laboratory Mormon cricket

samples indicated that the rectum, not the ileum, was primarily responsible for the difference in

community structure in the hindgut. Enterobacteriaceae comprised 83.5% of the sequences from

the rectum compared to 57.5% from the ileum in laboratory-raised animals, which was more

similar to the midgut in community structure (Fig. 4a). This distinction is potentially important

because higher digestive efficiency in conventional compared to germ-free crickets has been

attributed to microbial colonization of the ileum in the orthopteran *A domesticus* (Kaufman &

Klug 1991).

Metabolism of the specific complex carbohydrates attributed to bacteria in this study

were also identified in our metagenomic predictions and localized to the hindgut, as well as

enzymes involved in the production of the essential amino acid phenylalanine via the shikimate

pathway. Phenylalanine is a precursor for tyrosine, which is required to stabilize the cuticle

during molting (Bernays & Woodhead 1984) and in phenoloxidase synthesis, an important

component of the insect immune system (González-Santoyo & Córdoba-Aguilar 2012).

Tradeoffs between allocation of tyrosine to immune function and cuticle formation during

development in Mormon crickets (Srygley 2012) might be impacted by microbial contributions

to amino acid production if sufficient quantities of these amino acids are not obtained directly

from their diet.

Of the three enteric bacteria represented in this study, *P agglomerans* was common to

both field and lab individuals and increased in abundance in the hindgut. *Pantoea* are known

plant pathogens and have been associated with a variety of medical conditions in humans

(Walterson & Stavrinides 2015). In insects, however, *Pantoea* have been shown to have

mutualistic associations with their host. They are required for the completion of development in

stinkbugs (Hosokawa *et al.* 2016; but see Dillon & Charnley 2002), produce compounds that

attract insects to their host plants in flies (Robacker, Lauzon & He 2004; Maccollom *et al.* 2009),

and in the orthopteran *Schistocerca gregaria*, produce a key component of the locust aggregation

pheromone (Dillon, Vennard & Charnley 2000, 2002) and reduce susceptibility to microbial

pathogens through the production of phenols (Dillon & Charnley 1986, 1995).

Our metagenomics predictions suggest that enteric bacteria in Mormon crickets might be

capable of producing at least two of the antimicrobial phenols identified in *S. gregaria*, although

P. agglomerans was not identified as an important contributor in our study. This illustrates a

limitation of PICRUSt, as genomes in the IMG database used to make inferences about gene

content may miss important among-strain variation in metabolic capabilities. *P. agglomerans*

derived from *S. gregaria* are likely to have acquired this capability independently, unless the

metabolic pathway is different from the one analyzed here or the taxonomic designation reported

by Dillon and Charnley (1986, 1995) is incorrect.

Metagenomic analyses are also dependent upon annotation of the relevant pathways in

the KEGG database. We were unable, for example, to assess the potential for the Mormon

crickets microbiome to produce bacteriocins or the aggregation pheromone guaiacol, a bacterial

metabolite produced by *P. agglomerans* in *S. gregaria* (Dillon *et al.* 2000, 2002), because the

predicted pathways are not annotated in the KEGG database. The role of the gut microbiome in

protecting Mormon crickets from their own pathogens (MacVean & Capinera 1991) and its

influence on aggregation behavior (MacVean 1987; Simpson *et al.* 2006) is thus an important

direction for future research.

Conclusion

Variation in morphology and physiology is thought to differentiate niches within the gut that

influence the organization of the microbiome. Our study describes at high resolution how

bacterial communities vary among gut regions, and suggests that host-microbe and/or microbe-

microbe interactions have a role in how microbial communities are assembled and maintained.

While the metagenomics predictions gleaned from our study suggests that some of these bacteria

might benefit Mormon cricket nutrition, immunity, and perhaps even modulate social behavior,

experiments are needed to evaluate this possibility. Our establishment of methods for culturing

Mormon cricket gut bacteria will enable experimental and comparative genomic approaches in

the future to infer the ecological and evolutionary consequences of host-microbe symbiosis.

<u>Data Accessibility</u>

Sequences are archived under NCBI BioProject PRJNA362233.

Acknowledgements

504   We thank Laura Senior, USDA-ARS, for help with field collections and rearing laboratory

505   insects. We also thank Bruce Shambaugh and the Cheyenne office of USDA-APHIS, PPQ for

506   assisting with field collections, and Nancy Moran, UT-Austin, for providing the standard for the

507   qPCR experiment. Work was funded by National Science Foundation award DEB-1354666 and

508   the W.M. Wheeler Lost Pines Endowment from the University of Texas at Austin to UGM.

References

Bates, D., Maechler, M., Bolker, B. & Walker, S. (2013) *lme4: Linear Mixed-Effects Models Using S4 Classes*.

Benjamini, Y. & Hochberg, Y. (1995) Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society. Series B (Methodological)*, **57**, 289–300.

Bernays, E. & Woodhead, S. (1984) The need for high levels of phenylalanine in the diet of *Schistocerca gregaria* nymphs. *Journal of Insect Physiology*, **30**, 489–493.

Bignell, D.E. (1984) The arthropod gut as an environment for microorganisms. *Invertebrate-Microbial Interactions* (eds J.M. Anderson, A.D.M. Rayner & D.W.H. Walton), pp. 205–227. Cambridge University Press, Cambridge.

Bistolas, K.S.I., Sakamoto, R.I., Fernandes, J.A.M. & Goffredi, S.K. (2014) Symbiont polyphyly, co-evolution, and necessity in pentatomid stinkbugs from Costa Rica. *Microbial Symbioses*, **5**, 349.

Blum, J.E., Fischer, C.N., Miles, J. & Handelsman, J. (2013) Frequent replenishment sustains the beneficial microbiome of *Drosophila melanogaster*. *mBio*, **4**, e00860-13.

Brenner, D.J., Krieg, N.R., Staley, J.T., Garrity, G.M., Boone, D.R., De Vos, P., Goodfellow, M., Rainey, F.A. & Schleifer, K.-H. (eds). (2005) *Bergey's Manual® of Systematic Bacteriology*. Springer US, Boston, MA.

Bridson, E.Y. (ed). (1998) *The Oxoid Manual*, 8th ed. Oxoid LTD, Basingstoke, United Kingdom.

Buchon, N., Broderick, N.A. & Lemaitre, B. (2013) Gut homeostasis in a microbial world: insights from *Drosophila melanogaster*. *Nature Reviews Microbiology*, **11**, 615–626.

Bugg, T.D., Ahmad, M., Hardiman, E.M. & Singh, R. (2011) The emerging role for bacteria in lignin degradation and bio-product formation. *Current Opinion in Biotechnology*, **22**, 394–400.

Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A. & Holmes, S.P. (2016) DADA2: High-resolution sample inference from Illumina amplicon data. *Nature Methods*, **13**, 581–583.

Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., Costello, E.K., Fierer, N., Peña, A.G., Goodrich, J.K., Gordon, J.I., Huttley, G.A., Kelley, S.T., Knights, D., Koenig, J.E., Ley, R.E., Lozupone, C.A., McDonald, D., Muegge, B.D., Pirrung, M., Reeder, J., Sevinsky, J.R., Turnbaugh, P.J., Walters, W.A., Widmann, J., Yatsunenko, T., Zaneveld, J. & Knight, R. (2010) QIIME allows analysis of high-throughput community sequencing data. *Nature Methods*, **7**, 335–336.

Castex, M., Chim, L., Pham, D., Lemaire, P., Wabete, N., Nicolas, J.-L., Schmidely, P. &
Mariojouls, C. (2008) Probiotic *P. acidilactici* application in shrimp *Litopenaeus*
stylirostris culture subject to vibriosis in New Caledonia. *Aquaculture*, **275**, 182–193.

Castex, M., Lemaire, P., Wabete, N. & Chim, L. (2009) Effect of dietary probiotic *Pediococcus*
acidilactici on antioxidant defences and oxidative stress status of shrimp *Litopenaeus*
stylirostris. *Aquaculture*, **294**, 306–313.

Cho, I. & Blaser, M.J. (2012) The human microbiome: at the interface of health and disease.
Nature Reviews Genetics, **13**, 260–270.

Cintas, L.M., Casaus, M.P., Herranz, C., Nes, I.F. & Hernández, P.E. (2001) Review:
Bacteriocins of Lactic Acid Bacteria. *Food Science and Technology International*, **7**,
281–305.

Dillon, R. & Charnley, A. (1986) Inhibition of *Metarhizium anisopliae* by the gut bacterial flora
of the desert locust, *Schistocerca gregaria*: Evidence for an antifungal toxin. *Journal of*
invertebrate pathology, **47**, 350–360.

Dillon, R.J. & Charnley, A.K. (1988) Inhibition of *Metarhizium anisopliae* by the gut bacterial
flora of the desert locust: characterisation of antifungal toxins. *Canadian Journal of*
Microbiology, **34**, 1075–1082.

Dillon, R.J. & Charnley, A.K. (1995) Chemical Barriers to Gut Infection in the Desert Locust: In
Vivo Production of Antimicrobial Phenols Associated with the Bacterium Pantoea
agglomerans. *Journal of Invertebrate Pathology*, **66**, 72–75.

Dillon, R. & Charnley, K. (2002) Mutualism between the desert locust *Schistocerca gregaria*
and its gut microbiota. *Research in Microbiology*, **153**, 503–509.

Dillon, R.J. & Dillon, V.M. (2004) The gut bacteria of insects: nonpathogenic interactions.
Annual review of entomology, **49**, 71–92.

Dillon, R.J., Vennard, C.T. & Charnley, A.K. (2000) Pheromones: Exploitation of gut bacteria in
the locust. *Nature*, **403**, 851–851.

Dillon, R. j., Vennard, C. t. & Charnley, A. k. (2002) A Note: Gut bacteria produce components
of a locust cohesion pheromone. *Journal of Applied Microbiology*, **92**, 759–763.

Douglas, A.E. (2013) Alimentary canal, digestion and absorption. *The Insects: Structure and*
Function, 5th ed (eds S.. J. Simpson, A.E. Douglas & R.F. Chapman), pp. 46–80.
Cambridge University Press, Cambridge.

Douglas, A.E. (2015) Multiorganismal Insects: Diversity and Function of Resident
Microorganisms. *Annual Review of Entomology*, **60**, 17–34.

Drawz, S.M. & Bonomo, R.A. (2010) Three decades of β-Lactamase inhibitors. *Clinical*
Microbiology Reviews, **23**, 160–201.

Engel, P. & Moran, N.A. (2013) The gut microbiota of insects – diversity in structure and
function. *FEMS Microbiology Reviews*, **37**, 699–735.

Erkosar, B., Storelli, G., Mitchell, M., Bozonnet, L., Bozonnet, N. & Leulier, F. (2015) Pathogen
virulence impedes mutualist-mediated enhancement of host juvenile growth via inhibition
of protein digestion. *Cell Host & Microbe*, **18**, 445–455.

Faraway, J.J. (2006) *Extending the Linear Model with R: Generalized Linear, Mixed Effects and*
Nonparametric Regression Models. Chapman & Hall/CRC, New York.

Forsgren, E., Olofsson, T.C., Vásquez, A. & Fries, I. (2010) Novel lactic acid bacteria inhibiting
Paenibacillus larvae in honey bee larvae. *Apidologie*, **41**, 99–108.

González-Santoyo, I. & Córdoba-Aguilar, A. (2012) Phenoloxidase: a key component of the
insect immune system. *Entomologia Experimentalis et Applicata*, **142**, 1–16.

Gwynne, D.T. (1984) Sexual selection and sexual differences in Mormon crickets (Orthoptera:
Tettigoniidae, *Anabrus simplex*). *Evolution*, **38**, 1011–1022.

Herrmann, K.M. & Weaver, L.M. (1999) The shikimate pathway. *Annual review of plant*
biology, **50**, 473–503.

Hosokawa, T., Ishii, Y., Nikoh, N., Fujie, M., Satoh, N. & Fukatsu, T. (2016) Obligate bacterial
mutualists evolving from environmental bacteria in natural insect populations. *Nature*
Microbiology, **1**, 15011.

Hothorn, T., Bretz, F. & Westfall, P. (2008) Simultaneous inference in general parametric
models. *Biometrical Journal*, **50**, 346–363.

Hunt, J. & Charnley, A.K. (1981) Abundance and distribution of the gut flora of the desert
locust, Schistocerca gregaria. *Journal of Invertebrate Pathology*, **38**, 378–385.

Kaufman, M.G. & Klug, M.J. (1991) The contribution of hindgut bacteria to dietary
carbohydrate utilization by crickets (Orthoptera: Gryllidae). *Comparative Biochemistry*
and Physiology Part A: Physiology, **98**, 117–123.

Kikuchi, Y., Meng, X.-Y. & Fukatsu, T. (2005) Gut symbiotic bacteria of the genus burkholderia
in the broad-headed bugs *Riptortus clavatus* and *Leptocorisa chinensis* (Heteroptera:
Alydidae). *Applied and Environmental Microbiology*, **71**, 4035–4043.

Köhler, T., Dietrich, C., Scheffrahn, R.H. & Brune, A. (2012) High-resolution analysis of gut
environment and bacterial microbiota reveals functional compartmentation of the gut in
wood-feeding higher termites (*Nasutitermes* spp.). *Applied and Environmental*
Microbiology, **78**, 4691–4701.

Lanan, M.C., Rodrigues, P.A.P., Agellon, A., Jansma, P. & Wheeler, D.E. (2016) A bacterial
filter protects and structures the gut microbiome of an insect. *The ISME Journal*.

Langille, M.G.I., Zaneveld, J., Caporaso, J.G., McDonald, D., Knights, D., Reyes, J.A.,
615   Clemente, J.C., Burkepile, D.E., Vega Thurber, R.L., Knight, R., Beiko, R.G. &
616   Huttenhower, C. (2013) Predictive functional profiling of microbial communities using
617   16S rRNA marker gene sequences. *Nature Biotechnology*, **31**, 814–821.

618   Lehane, M.J. & Billingsley, P.F. (eds). (1996) *Biology of the Insect Midgut*. Springer
619   Netherlands, Dordrecht.

620   Ley, R.E., Lozupone, C.A., Hamady, M., Knight, R. & Gordon, J.I. (2008) Worlds within
621   worlds: evolution of the vertebrate gut microbiota. *Nature Reviews Microbiology*, **6**, 776–
622   788.

623   Maccollom, G.B., Lauzon, C.R., Sjogren, R.E., Meyer, W.L. & Olday, F. (2009) Association and
624   attraction of blueberry maggot fly curran (Diptera: Tephritidae) to *Pantoea*
625   *(enterobacter) agglomerans*. *Environmental Entomology*, **38**, 116–120.

626   MacVean, C.. (1987) Ecology and management of the Mormon cricket, *Anabrus simplex*.
627   *Integrated Pest Management on Rangeland: A Shortgrass Prairie Perspective*, pp. 116–
628   136. Westview Press, Boulder, CO.

629   MacVean, C.M. & Capinera, J.L. (1991) Pathogenicity and transmission potential of Nosema
630   locustae and Vairimorpha n. sp. (Protozoa: Microsporida) in Mormon crickets (Anabrus
631   simplex; Orthoptera: Tettigoniidae): A laboratory evaluation. *Journal of Invertebrate*
632   *Pathology*, **57**, 23–36.

633   McFrederick, Q.S., Cannone, J.J., Gutell, R.R., Kellner, K., Plowes, R.M. & Mueller, U.G.
634   (2013) Specificity between *Lactobacilli* and hymenopteran hosts is the exception rather
635   than the rule. *Applied and Environmental Microbiology*, **79**, 1803–1812.

636   McMurdie, P.J. & Holmes, S. (2013) Phyloseq: an R package for reproducible interactive
637   analysis and graphics of microbiome census data. *PLoS ONE*, **8**, e61217.

638   Mead, L.J., Khachatourians, G.G. & Jones, G.A. (1988) Microbial ecology of the gut in
639   laboratory stocks of the migratory grasshopper, *Melanoplus sanguinipes* (fab.)
640   (Orthoptera: Acrididae). *Applied and Environmental Microbiology*, **54**, 1174–1181.

641   Oksanen, J., Blanchet, F.G., Kindt, R., Legendre, P., Minchin, P.R., O'Hara, R.B., Simpson,
642   G.L., Solymos, P., Stevens, M.H.H. & Wagner, H. (2015) *Vegan: Community Ecology*
643   *Package*.

644   Powell, J.E., Martinson, V.G., Urban-Mead, K. & Moran, N.A. (2014) Routes of acquisition of
645   the gut microbiota of the honey bee *Apis mellifera*. *Applied and Environmental*
646   *Microbiology*, **80**, 7378–7387.

647   R Core Development Team. (2013) *R: A Language and Environment for Statistical Computing*.
648   R Foundation for Statistical Computing, Vienna, Austria.

Raskin, I. (1992) Role of salicylic acid in plants. *Annual review of plant biology*, **43**, 439–463.

Robacker, D.C., Lauzon, C.R. & He, X. (2004) Volatiles production and attractiveness to the
Mexican fruit fly of *Enterobacter agglomerans* isolated from apple maggot and Mexican
fruit flies. *Journal of Chemical Ecology*, **30**, 1329–1347.

Ryu, J.-H., Ha, E.-M. & Lee, W.-J. (2010) Innate immunity and gut–microbe mutualism in
Drosophila. Developmental & Comparative Immunology, **34**, 369–376.

Servin, A.L. (2004) Antagonistic activities of lactobacilli and bifidobacteria against microbial
pathogens. *FEMS Microbiology Reviews*, **28**, 405–440.

Simpson, S.J., Sword, G.A., Lorch, P.D. & Couzin, I.D. (2006) Cannibal crickets on a forced
march for protein and salt. *Proceedings of the National Academy of Sciences of the
United States of America*, **103**, 4152–4156.

Smith, C.C., Srygley, R.B., Dietrich, E.I. & Mueller, U.G. (2016) Partitioning the effect of
mating and nuptial feeding on the microbiome in gift-giving insects. *Environmental
Microbiology Reports*.

Srygley, R.B. (2012) Ontogenetic changes in immunity and susceptibility to fungal infection in
Mormon crickets *Anabrus simplex. Journal of Insect Physiology*, **58**, 342–347.

Srygley, R.B. & Lorch, P.D. (2011) Weakness in the band: nutrient-mediated trade-offs between
migration and immunity of Mormon crickets, *Anabrus simplex. Animal Behaviour*, **81**,
395–400.

Srygley, R.B., Lorch, P.D., Simpson, S.J. & Sword, G.A. (2009) Immediate protein dietary
effects on movement and the generalised immunocompetence of migrating Mormon
crickets *Anabrus simplex* (Orthoptera: Tettigoniidae). *Ecological Entomology*, **34**, 663–
668.

Storelli, G., Defaye, A., Erkosar, B., Hols, P., Royet, J. & Leulier, F. (2011) *Lactobacillus
plantarum* promotes *Drosophila* systemic growth by modulating hormonal signals
through TOR-dependent nutrient sensing. *Cell Metabolism*, **14**, 403–414.

Tai, V., Carpenter, K.J., Weber, P.K., Nalepa, C.A., Perlman, S.J. & Keeling, P.J. (2016)
Genome evolution and nitrogen fixation in bacterial ectosymbionts of a protist inhabiting
wood-feeding cockroaches. *Applied and Environmental Microbiology*, **82**, 4682–4695.

Ulrich, R.G., Buthala, D.A. & Klug, M.J. (1981) Microbiota associated with the gastrointestinal
tract of the common house cricket, *Acheta domestica. Applied and Environmental
Microbiology*, **41**, 246–254.

Venables, W.N. & Ripley, B.D. (2002) *Modern Applied Statistics with S*, Fourth. Springer, New
York.

Walterson, A.M. & Stavrinides, J. (2015) Pantoea: insights into a highly versatile and diverse
genus within the Enterobacteriaceae. *FEMS Microbiology Reviews*, fuv027.

Woodring, J. & Lorenz, M.W. (2007) Feeding, nutrient flow, and functional gut morphology in
the cricket *Gryllus bimaculatus*. *Journal of Morphology*, **268**, 815–825.

Xu, W.-H., Liang, Q., Zhang, Y.-J. & Zhao, P. (2015) Naturally occurring arbutin derivatives
and their bioactivities. *Chemistry & Biodiversity*, **12**, 54–81.

Yuan, S., Cohen, D.B., Ravel, J., Abdo, Z. & Forney, L.J. (2012) Evaluation of methods for the
extraction and purification of dna from the human microbiome. *PLoS ONE*, **7**, e33865.

Figures and Tables

Table 1. Analysis of deviance comparing alpha diversity between source populations (wild or laboratory) and among tissue types (foregut, midgut, and hindgut). Values represent the F-statistic (p-value) for each term. Statistically significant terms (p<0.05) are indicated in bold. Degrees of freedom were estimated using the Kenward-Rogers approximation. In the reduced dataset, two individuals from the laboratory-reared population were removed (see Methods).

	Reduced Dataset			Full Dataset		
	Species Richness	Chao1	Shannon-Weiner	Species Richness	Chao1	Shannon-Weiner
Source	**13.9 (0.003)**	**11.0 (0.007)**	**9.17(0.01)**	**14.0(0.002)**	**10.5 (0.006)**	**8.22 (0.013)**
Tissue type	**5.85 (0.010)**	**4.79 (0.02)**	**7.07 (0.005)**	**6.77 (0.004)**	**5.68 (0.008)**	**8.44 (0.001)**
Interaction	0.51 (0.61)	1.02 (0.38)	0.98 (0.39)	0.66 (0.53)	1.15 (0.33)	1.28 (0.29)

Table 2. Permutation tests of distance-based redundancy analyses of DADA2 16S rRNA sequence variants and PICRUSt metagenomic predictions. Source population (wild or laboratory) and tissue type (foregut, midgut and hindgut) were entered as factors.

	Reduced Dataset		Full Dataset	
	F	*P*	*F*	*P*
DADA2				
Source	**8.99**	**<0.001**	**7.75**	**<0.001**
Tissue type	**9.85**	**<0.001**	**5.49**	**<0.001**
Interaction	0.26	0.47	0.52	0.84
PICRUSt				
Source	0.23	0.76	**4.31**	**0.04**
Tissue type	**1.61**	**0.003**	**8.09**	**0.002**
Interaction	0.16	0.35	0.92	0.43

Table 3. Likelihood ratio tests from GLMMs fitting the abundance of sequence variants to source population (wild or laboratory) and tissue type (foregut, midgut or hindgut). Values are Chi-square (p-value). LAC1=*Lactobacillus sp.*, PED=*Pediococcus*, PAG=*Pantoea agglomerans*.

	Reduced Dataset			Full Dataset		
	LAC1	PED	PAG	LAC1	PED	PAG
Source	1.32 (0.25)	0.80 (0.37)	2.36 (0.12)	0.18 (0.66)	0.51 (0.48)	0.01 (0.96)
Tissue type	**16.3 (<0.001)**	**9.25 (0.01)**	**20.7 (<0.001)**	**12.7 (0.002)**	**16.1 (<0.001)**	**41.9 (<0.001)**
Interaction	0.36 (0.84)	0.84 (0.66)	3.86 (0.15)	0.3 (0.86)	0.95 (0.62)	2.67 (0.26)

Table 4. Permutation test of distance-based redundancy analysis of PICRUSt metagenomics predictions between source populations (wild or laboratory) and among tissue types (foregut, midgut and hindgut).

	Reduced Dataset		Full Dataset	
	F	*P*	*F*	*P*
Illumina 16S rRNA				
Source	**8.99**	**<0.001**	**7.75**	**<0.001**
Tissue type	**9.85**	**<0.001**	**5.49**	**<0.001**
Interaction	0.26	0.47	0.52	0.84
KEGG Pathways				
Source	0.23	0.76	**4.31**	**0.04**
Tissue type	**1.61**	**0.003**	**8.09**	**0.002**
Interaction	0.16	0.35	0.92	0.43

Figure 1. External morphology of the (a) foregut, (b) midgut, and (c) hindgut in the Mormon cricket. E=esophagus, Cr=crop, PV=proventriculus, Ce=cecum, V=ventriculus, MT=Malphigian tubules, Ile=ileum, C=colon, R=rectum. Malphigian tubules have been trimmed to illustrate their entry point into the hindgut.

Figure 2. Mean relative abundance of the eleven dominant sequence variants from (a) field-caught and (b) laboratory-raised Mormon crickets from 16S rRNA Illumina sequencing.

Figure 3. Alpha diversity in field-caught and laboratory-raised Mormon crickets.

Figure 4. Ordination of sample scores from the db-RDA of the reduced dataset for (a) Illumina 16S sequencing and (b) PICRUSt metagenomic predictions.

Figure 5. Abundance of bacterial 16S rRNA genes in the Mormon crickets gut. Bars indicate significant differences between regions (* p<0.05).

Figure 6. Relative abundance of KEGG pathways related to nutrition, immunity, xenobiotic degradation, and secondary metabolite production among tissue types.

Figure 7. Abundance of KEGG orthologs related to (a) carbohydrate metabolism and (b) antimicrobial compound production among tissue types, and the contributions of taxonomic groups (c,d) to ortholog abundance. Key for colors representing taxonomic groups are in Figure 2.

Fig 8. Maximum likelihood estimation of phylogenetic relationships among lactic-acid bacteria 16S rRNA sequences from Mormon cricket gut isolates and their relatives. Branches with bootstrap support <75% are collapsed.

Figure 9. Maximum likelihood estimation of phylogenetic relationships among enteric bacteria 16S rRNA sequences from Mormon cricket gut isolates and type strains. Branches with bootstrap support <50% are collapsed.