1

Title

Maintaining tiger connectivity and minimizing extinction into the next century: Insights from

landscape genetics and spatially-explicit simulations

Authors

Prachi Thatte[1*], Aditya Joshi[2], SrinivasVaidyanathan[3], Erin Landguth[4]and Uma Ramakrishnan[1*]

Affiliations

1- National Center for Biological Sciences, Tata Institute of Fundamental Research

2- Wildlife Conservation Trust

3- Foundation for Ecological Research, Advocacy and Learning

4- School of Public and Community Health Sciences, University of Montana, 32 Campus Drive,

Missoula, MT 59812 USA.

Corresponding authors: prachi.thatte@gmail.com; uramakri@ncbs.res.in

National Centre for Biological Sciences, Tata Institute of Fundamental Research, Bellary Road,

Bangalore 560065. Phone number: +91 80 23666030/31

Abstract

Habitat loss is the greatest threat to large carnivores around the world. Maintenance of

functional connectivity in fragmented landscapes will be important for long-term species

persistence. Here, we merge landscape genetics analyses and spatially-explicit simulations to

understand future persistence and extinction of tigers (*Panthera tigris*) in Central India. Tigers

in this landscape are restricted to Protected Areas (PAs) and forest fragments embedded within

a mosaic of agricultural fields and human settlements. We examined current population

connectivity of tigers across nine reserves (using 116 non-invasively sampled individuals and 12

microsatellites). Genetic data was used to infer resistance-to-movement. Our results suggest

that dense human settlements and roads with high traffic are detrimental to tiger movement.

We used landscape genetic simulations to model 86 different scenarios that incorporated

impacts of future land-use change on inferred population connectivity and extinction. Our

results confirm that genetic variability (heterozygosity) will decrease in the future and small

and/or isolated PAs will have a high risk of local extinction. The average extinction risk of small

PAs reduced by 23-70% on adding a 5 km buffer around exiting boundaries. Unplanned

development results in 35% lower heterozygosity and 56% higher average extinction probability

for tigers even within protected areas. Increasing tiger numbers in such a scenario decreases

extinction probability just by 12 % (from 56% to 44%). Scenarios where habitat connectivity was

enhanced and maintained, stepping-stone populations were introduced/maintained, and tiger

numbers were increased, led to low overall extinction probability (between 3-21%). Our

simulations provide a means to quantitatively evaluate the effects of different land-use change

scenarios on connectivity and extinction, linking basic science to land-use change policy and

planned infrastructure development.

Introduction

The current rates and magnitude of species decline and extinction are higher than ever

before (Barnosky *et al.* 2011; Dirzo *et al.* 2014). Most mammals retain less than half of their

historical range, resulting in substantial population decline and habitat fragmentation

(Morrison *et al.* 2007; Dirzo *et al.* 2014). Due to their large area requirements, slower life

histories and low densities, large carnivores are especially vulnerable to habitat fragmentation

and isolation. A majority (77%) of large carnivores continue to undergo worldwide decline, with

populations at risk of local extirpation due to habitat loss (Ripple *et al.* 2014).

Conservation efforts, including population monitoring, legal protection, creation of

protected areas, reintroductions and translocations, have ensured recovery of species, such as

the grey wolf (*Canis lupus*) in North America and brown bears (*Ursus arctus*) in northern

Europe, among others (Ripple & Beschta 2012; Hagen *et al.* 2015). Long-term persistence of

such threatened populations requires identifying and maintaining connectivity among habitat

patches (Jackson *et al.* 2016). Among large carnivores, significant attention and resources are

invested in recovery and conservation of the tiger, an iconic species with less than 4000

individuals left in the wild.

Tigers have lost four subspecies and 93% of their historical range, and what remains of

their existing range is highly fragmented. With nearly 65% of the world's wild tigers (Jhala, Y.

V.;Qureshi, Q.;Gopal 2015) and substantial genetic variation (Mondol *et al.* 2009a), India is a

stronghold for tiger survival. Recent reports suggest that conservation and management efforts

in India over the last three decades have led to a 30% increase in tiger numbers (Jhala, Y.

V.;Qureshi, Q.;Gopal 2015). Despite what appears to be a demographic recovery, the median

number of tigers within individual protected areas (PA) in India is low (median: 19, range: 2-

215; (Wikramanayake *et al.* 2010; Jhala, Y. V.;Qureshi, Q.;Gopal 2015)). Most populations by

themselves may not be viable, and the continued survival of tigers could be contingent on

maintaining connectivity between PAs.

Several independent genetic studies in the high priority tiger conservation landscape of

Central India confirm that PAs exchange dispersing individuals and are fairly well connected

(Joshi *et al.* 2013; Sharma *et al.* 2013a; Yumnam *et al.* 2014; Reddy *et al.* 2017). About 35% of

India's tigers are estimated to live outside PAs (Jhala, Y. V.;Qureshi, Q.;Gopal 2015) and may

play a critical role in maintaining connectivity. However, India is a country with over a billion

people, an economy growing at 7% annually, poised for rapid urbanization and the ensuing

increase in associated infrastructure. Among the planned infrastructure, existing highways in

the landscape are being widened to meet the demands of increasing traffic (e.g., National

Highway 7 which bisects a critical corridor is being widened after a prolonged legal battle owing

to the conflict between tiger conservation and development activities (Srivastava & Tyagi

2016)), further fracturing an already highly fragmented landscape. Landscapes outside PAs

(~95% of India's area) are about to change dramatically, and tiger management and

conservation is currently only focused on protected areas.

Earlier studies correlating genetic connectivity with landscape elements have revealed

that tiger movement is negatively impacted by human settlements within the Central Indian

landscape (CIL) (Joshi *et al.* 2013). Urban populations in India are projected to double from 410

million in 2014 to 814 million in 2050 (United Nations *et al.* 2014). Additionally, built-up areas

have been increasing almost 3 times faster than population in nearly all large Indian cities

(Sudhira 2011). Along with the urbanization, demands for better road and railway connectivity

between cities is also projected to increase (National Transport Policy Development Committee

2013). Such landscape transitions will negatively impact tiger connectivity. Small, isolated

populations have known genetic consequences, including low variation (Frankham 1996;

Boersen *et al.* 2003), increased inbreeding and increased disease susceptibility (Spielman *et al.*

2004; Trinkel *et al.* 2011), and heightened extinction risk (Saccheri *et al.* 1998). However, these

are general predictions, and we do not know how future land-use change will specifically

impact connectivity and persistence of tigers in the landscape.

Few studies (Tian *et al.* 2011, 2014) have attempted to understand how future climate

and landscape change might affect persistence of tigers. Tian et al. (2014) used population

viability analysis (PVA) to simulate the effect of climate change and habitat fragmentation on

future persistence of Amur tigers by incorporating factors affecting species distribution, but not

dispersal. Persistence of tigers in complex and changing landscapes such as CIL requires

modeling population persistence based on empirical genetic data, factors affecting dispersal,

predicted landscape/climate change and interactions of these factors with demography.

Globally, very few such intensive modeling exercises have been conduct to predict future

persistence of endangered species (Landguth *et al.* 2014; Benson *et al.* 2016; Brown *et al.*

2016).

In this paper, we examine genetic connectivity among tiger populations in the CIL,

including individuals within and outside PAs. We use this data on gene flow to infer the effect of

different landscape features on dispersal and connectivity. We then carry out forward- time,

spatially-explicit, individual-based simulations to understand how genetic diversity, connectivity

and extinction probability will change under nine different development scenarios. We examine

these scenarios, while accounting for tigers inside and outside PAs. We also test the effect of

increasing tiger numbers and the effects of assumptions about dispersal, modeling a total of 86

scenarios.

Materials and Methods

In this study, we collected genetic samples from wild populations (*Study area and*

sampling), generated genetic data and conducted population genetic analyses (*Genotyping and*

population genetic analysis). Landscape genetic analyses allowed us to infer landscape

elements impacting connectivity (*Landscape genetic analyses*). Future scenarios were simulated

assuming various criteria for landscape change, including specific management relevant

scenarios, and tiger demography (*Landscape genetic simulations*). All simulations inferred

genetic variability, inbreeding, connectivity and extinction probability in 2100. A flow chart of

the methods is presented in Figure S1 and more detail is described as follows.

Study area and sampling

The CIL is a global priority tiger conservation landscape. With ~34% forest cover and an

estimated 688 (596- 780) tigers (Jhala, Y. V.;Qureshi, Q.;Gopal 2015), it is a stronghold for tiger

conservation. The PAs in the landscape are embedded in a heterogeneous matrix of multiple

land-use types.

Non-invasive (scat) samples (n= 580) were collected between October 2012 and April

2014 from potential areas (PAs and forested areas outside PAs which are a part of territorial

forest divisions and forest development corporations) in the state of Maharashtra, Madhya

Pradesh and Chhattisgarh. We sampled eleven PAs: (1) Kanha Tiger Reserve (KTR), (2) Pench

Tiger reserve (PTR), (3) Bandhavgarh Tiger Reserve (BTR), (4) Achanakmar Tiger Reserve (ATR),

(5) Nagzira (NGZ) and (6) Nawegaon (NAW) (which together comprise a Tiger Reserve) (7)

Satpura Tiger Reserve (STR), (8) Tadoba-Andhari Tiger Reserve (TATR), (9) Bor Tiger Reserve

(BOR), (10) Umred-Karhandla Wildlife Sanctuary (UK) and (11) Tipeshwar Wildlife Sanctuary

(TIP). We also sampled in three territorial forest divisions outside PAs: (1) Balaghat Forest

Division (BAL), (2) Central-Chanda Forest Division (CHA) and (3) Bramhapuri Forest Division

(BPR) (Figure 1). Sampling was also carried out in Sitanadi-Udanti Tiger Reserve (S-U) and Panna

Tiger Reserve (PAN). We did not find any tiger scat samples in S-U and samples from PAN were

not used to optimize the resistance layers since tigers in PAN have been reintroduced from KTR

and PTR after they went extinct locally in 2006. However, we included PAN in the forward-time

simulations. See supplementary material S1 for more details.

Genotyping and population genetic analysis

In order to quantify genetic connectivity between different PAs, we first extracted DNA

using standardized methods and identified individuals(Mukherjee *et al.* 2007; Mondol *et al.*

2009b). We then calculated heterozygosity based differentiation statistics (PopGenReport

(Adamack & Gruber 2014), MMOD (Winter 2012), and HIERFSTAT (Goudet 2005) in R (Ihaka &

Gentleman 2012)). For further details about genetic analysis, refer to the supplementary

material S2.

Landscape genetics analysis

We calculated inter-PA genetic distances based on proportion of shared alleles (D_{PS};

(Bowcock *et al.* 1994)) using Microsatellite Analyzer (MSA, version 4.05; (Dieringer &

Schlutterer 2003)). The relationship between the observed genetic structure and the landscape

variables likely to affect tiger dispersal was systematically evaluated using a multi-model

inference and optimization approach (Shirk *et al.* 2010) described next.

We selected landscape variables known to affect tiger dispersal based on Joshi et al.

(2013) (i.e., land cover, human settlement layer, roads and railway lines along with density of

linear features) to build resistance models. We did not use topography related variables as used

in Reddy et al. (2017). Tigers are habitat generalists (ranging from 0- 3000 meters above sea

level), their occurrence is associated with forests in the CIL and they have been reported to

avoid desserts and short grasslands (Kitchener and Dugmore 2000, Yumnam et al. 2014).

Athreya et al. (2014) revealed that tigers disperse through and use less rugged areas in Central

India. Additionally, over the last 300 years most forest cover loss has been in the low elevation

and less rugged areas. As a result, the current forest cover is concentrated in highly rugged and

relatively high-slope areas ((Sharma *et al.* 2013b), supplementary figure S3).We used MODIS

land-cover data (MCD12Q1) and reclassified it into five broad land-cover types which we ranked

in order of increasing resistance: (i) forest, (ii) degraded and scrub forest, (iii) agriculture

(including fallow/ wasteland) and (iv) built-up (developed with buildings and non-building

structures) areas. We developed a layer of human settlements by merging urban and peri-

urban areas derived from nightlight data (available at National Geophysical Data Centre) and

rural areas from population density data (available at

http://www.worldpop.org.uk/data/get_data/). Since all villages in the study landscape are not

electrified, using nightlights alone would have underrepresented human settlements within the

landscape. A vector layer of national highways, state highways and major public roads was

reclassified into 5 categories based on the intensity of traffic on the road (based on Passenger

Car Unit (PCU) data for 2006 from the Ministry of Road Transport and Highways). PCU

represents the number of vehicles in terms of passenger cars and accounts for all types of

vehicles. See supplementary material S3 for more details. Railway lines and roads were used to

generate a layer representing the density of linear features.

Landscape resistance values were inferred from the genetic data using multi-model

optimization approach described in Shirk et al (2010). Each landscape variable was related to

landscape resistance using a mathematical model (supplementary material S4). Using genetic

data as the response variable and systematically varying the model parameters (maximum

resistance and a shape parameter to account for the relationship of the variable with

resistance), we identified the best fitting model parameters for each variable. The best fitting

model was identified as the one with the most significant correlation with genetic data, after

controlling for the effect of geographic distance using partial Mantel tests (Smouse *et al.* 1986).

The landscape variables that explained significant variation after controlling for geographic

distance in univariate models were retained for further analysis. The retained variables were

combined (additively) and optimized again in a multivariate context to account for interactions

between different landscape variables(Shirk *et al.* 2010). We then used the estimated

landscape resistance data to understand how future landscape change may alter the resistance

surface (and therefore, connectivity) as described below.

Landscape genetic simulations

We used CDPOP (Landguth & Cushman 2010) to simulate 20 non-overlapping

generations (~100 years for tigers) of mating and dispersal among individuals for each scenario

(scenarios described in next section). CDPOP is a spatially-explicit, individual-based simulator of

population genetic processes. It simulates mating and dispersal in a finite population assigned

to fixed locations, recording alleles of all individuals every generation. We initialized individuals

in the simulations as being characterized by 12 loci with at most 11 alleles per locus. For the

PAs with genetic data, individuals were assigned alleles based on the allele frequency

distribution for that PA. For PAs without genetic data, we assigned allele frequencies based on

STRUCTURE results (Figure S5) or using other information (see S5(a) for details). Individuals

were initially seeded on the landscape based on the current tiger estimates (Table S4). In each

generation, females gave birth to offspring (number based on a normal distribution with mean

3±2) with an equal sex ratio at birth. Adults died, and the vacant locations were occupied by

dispersing individuals. Probability governing mating and dispersal related movement was based

on proximity of individual locations specified by the pairwise cost-distance matrices between

individuals (calculated using 'gdistance' in R), which were scenario-specific (see next section). A

negative exponential function (Sutherland *et al.* 2000) with median dispersal distance of 85 km

(Bowman *et al.* 2002) and two different maximum dispersal distances of 300km (Patil *et al.*

2011) and 500km (Bowman *et al.* 2002; Natesh *et al.* 2017) was used to model natal dispersal. If

all locations were occupied, any remaining offspring not assigned to a location were eliminated

(Balloux 2001).The probabilistic functions governing birth rate, mating and dispersal introduced

demographic stochasticity in the model. See Table S1 for more details on simulation

parameters. At the end of each simulation, we calculated genetic diversity indices

(heterozygosity, inbreeding and allelic richness) and differentiation indices (Global and pairwise

Fst, G'st, Jost D) (Jombart 2008; Adamack & Gruber 2014). Probability of tiger extinction for

each PA was the number of times the tiger population within the PA went extinct out of 100

replicate simulation runs. We calculated sex ratio skew and fluctuations in population size

across generations.

Simulation scenarios

In order to explore how future landscape change may affect dispersal and connectivity

of tiger populations, we carried out spatially-explicit genetic simulations under different

landscape change scenarios. All scenarios we present are plausible, and derived from covariates

which are known to influence land-use land-cover change. The objective of these scenarios was

to identify the kind of landscape-wide, land-use changes that will facilitate or impede tiger

connectivity, and not to identify where land-use change will occur in the future. Scenarios of

landscape change were developed using Land Change Modeler (LCM) in IDRISI (SELVA,

http://www.clarklabs.org) based on the change in land-use land-cover (LULC) from a

combination of land cover (MODIS) and human settlements data and road expansion from 2001

to 2012. Future LULC maps were generated for 2020, 2040, 2060, 2080. See supplementary S3a

for details.

Simulations to assess change in genetic connectivity in the future were carried out for

nine landscape change scenarios (labelled F1-F9, see Table 1 for the scenario description and

rationale). For each of these scenarios, the landscape was parameterized using optimized

resistance values based on the landscape genetic analysis described in the previous section.

Cost-distance matrices were calculated for all the future maps that were generated. During

simulations, the distance matrices changed after every 4 generations (~20 years) to account for

the changing landscape except for scenario F1 (no landscape change) and F9 (PAs were fenced

at the 1st time step of 20 years and remained fenced after that). Along with scenarios F1 to F9,

we carried out additional simulations for scenarios F1 and F4 where we added a 5km buffer

around the smaller PAs in the landscape (< 400km^2). We evaluated the effect of increased PA

size on connectivity and extinction estimates. Simulations for the first eight scenarios were

carried out under 4 sub-scenarios:(a) with tigers restricted to PAs (current numbers constant),

(b) with tigers inside and outside PAs (outside individuals distributed randomly within existing

forest patches), (c) with tigers inside and outside PAs (outside individuals clustered in space to

form 'stepping-stone' populations between PAs), and (d) with tigers restricted to PAs (numbers

increase). See supplementary material S5b for details.

The simulations included eleven additional unsampled PAs: PAN, S-U, Indravati Tiger

Reserve (IND), Melghat Tiger Reserve (MEL), Ratapani Wildlife Sanctuary (RAT), Kawal Tiger

Reserve (KAW), Tamorpingla Wildlife Sanctuary (TAM), Sanjay Tiger Reserve (SAN), Noradehi

Wildlife Sanctuary (NOR), Barnawapara Wildlife Sanctuary (BAR)and Chaprala Wildlife

Sanctuary (CHH). These areas were included in future scenario simulations as some have tigers,

while others have high probability of tiger occupancy, or are protected reserves.

Results

Population Genetic Analysis

Out of 289 samples identified as tigers, data for at least 8 out of 12 microsatellite loci could be generated from 127 samples, 116 of which were identified as unique individuals. The P(ID) (the probability of two different individuals having the same genotype) was 1.48×10^{-11} and the more conservative measure Sib P(ID) (PID when all individuals in the population are assumed to be siblings) was 5.1×10^{-5}, indicating that even related individuals would have a very low probability of having identical genotypes. The estimated allelic dropout across loci was 0.062 and the frequency of false alleles was 0.013 (comparable to other studies in the landscape; (Caragiulo *et al.* 2015)). Six out of the 12 loci showed significant deviation from Hardy- Weinberg equilibrium, suggesting the presence of genetic structure. The mean number of alleles per locus was 8.7 and expected heterozygosity was 0.723 (Supplementary Table S2). Global F_{ST} was estimated to be 0.169, with highest pairwise differentiation between BTR and BPR (Tables S2 and S3). Table 2 presents D_{PS}, a measure of contemporary genetic differentiation.

Functional connectivity:

Human settlements were the most important (highest magnitude of correlation) landscape variable explaining genetic distance between PAs. Land-use and traffic intensity on roads also explained significant variation, even after accounting for geographic distance (Table 3). These three variables (traffic intensity on roads, human settlements and land-use) were retained for multivariate optimization. The parameter estimates for each landscape variable are presented in Table 3.

Shape parameter (the parameter that determines the shape of the relationship between

the landscape variable and resistance) and maximum resistance of the optimum models of all

the three landscape variables changed on combining, suggesting interaction between these

variables. The non-linear transformations (x >1) indicate that low resistance is offered by

smaller and middle values assigned to the variable and the resistance increases steeply with

very high values. For example, roads with low and moderate traffic offer negligible resistance to

movement, however, the resistance increases steeply with very high traffic. Correlation

between the pairwise cost distance among populations (estimated from the combined

resistance surface) and genetic distance was high (0.7857 after controlling for the geographic

distance, 0.8166 without controlling for geographic distance). The correlation value for isolation

by distance model (geographical distance alone) was 0.624

Future connectivity

Overall, genetic diversity reduced over time in all the simulation scenarios (Figure S6).

However, restoring and protecting corridors between PAs lead to the least decline in genetic

variation (~20%). Heterozygosity decreased faster and reduced to a lower final value at the end

of 100 years in the scenarios with lower dispersal threshold (300km). Within both the dispersal

categories, the loss of genetic diversity was greater in the scenarios where the forest cover loss

was higher (Figure S6). Our results suggest that inbreeding did not increase appreciably over

100 years (F<0.25 in all scenarios). Figure 2 summarizes the implications of three different

management decisions on structural connectivity, allelic richness, inbreeding and extinction in a

subset of the simulated scenarios.

Irrespective of land-use change scenario, dispersal threshold and tiger demographic

trajectory, small isolated PAs (TIP and BOR) had the highest risk of extinction. Reducing the

dispersal threshold from 500km to 300km doubled the average extinction probability of some

populations (BTR, PTR, MEL and STR, see Figure 3a and 3b). Currently well connected, but small

PAs (UK, CHH, NGZ and NAW) had high extinction probability only in the scenarios where forest

cover around them was lost. Some large PAs that currently have a very low number of tigers (<

10 tigers, KAW, IND, RAT and S-U) also had high extinction probability except in the sub-

scenarios where tiger numbers increased. Small population size was associated with high

variance and highly skewed sex ratios (Figures S7 and S8). The extinction probability of small

PAs was also governed by their isolation and associated re-colonization probability (Figure 3).

Adding a buffer around the small PAs reduced their overall extinction probability. Among the

isolated small PAs (TIP and BOR), addition of buffer reduced the extinction probability by ~23%.

Adding a buffer zone around the small, currently connected populations reduced the extinction

probability to a large extent (~70%), but only in the simulations where the landscape around

the buffers changed (F4). The benefit of having a buffer around the small PAs was the highest in

the sub-scenario with stepping stone populations (sub-scenario c) between PAs.

Change in connectivity: specific infrastructure projects

Increase in mined area and associated increase in built-up areas lead to ~15 times

higher extinction probability in small and medium sized PAs of BTR, SAN, UK and KAW due to

their proximity to coal fields. Presence of NH7 as a barrier without any mitigation structures

(scenario F6) increased the F_{ST} between KTR and PTR ~4 times compared to scenario F1 and

scenario F7 (Figure S9). NH7 bisects the corridor between these two PAs. NH6 bisects the

corridor between NGZ and NAW. The scenario with NH6 as a barrier (scenario F6) leads to ~19

times and ~65 times higher probability of extinction for dispersal threshold of 500 km and 300

km respectively. Higher increase in the genetic differentiation in the latter case (between NGZ-

NAW, as compared to KTR-PTR) could be due to the smaller current population sizes of NGZ and

NAW. Across land-use change scenarios, increasing tiger number (sub-scenario d) within and

outside PAs (sub- scenarios c) lead to ~37% lower extinction probabilities overall and ~11%

lower reduction in heterozygosity.

Discussion

The St. Petersburg declaration of 2010 envisaged doubling tiger numbers across all tiger nations by 2022 (Global Tiger Recovery Program 2011). Our results from CIL demonstrate that maintaining or establishing connectivity and ensuring protection will be critical to meet such tiger conservation targets. Along with corridor conservation, designing, notifying and maintaining stepping-stone populations between PAs will be beneficial for maintaining future connectivity. To facilitate regional planning, spatially explicit analysis could help identify locations where infrastructure could be developed without affecting tiger connectivity. The current results are general prescriptions, but findings from this study provide evidence that can be translated into spatially explicit conservation planning exercises. Additionally, our framework can be applied globally to other landscapes and species (where genetic data are available) to improve connectivity in the context of development.

What impacts current tiger connectivity?

Human footprint on the landscape had the strongest impact on connectivity. Dense human settlements and roads with high traffic offered highest resistance to movement. Degraded forests and plantations offered negligible resistance and agriculture-village matrix offered low resistance to tiger movement. However, long contiguous stretches of agriculture-village matrix could lead to accumulation of cost over space and impede movement. Our results are supported by empirical data on tiger movement. Recent data from radio-collared tigers reveals that long distance dispersing tigers do not avoid agriculture-village matrix and cross low traffic roads (Athreya *et al.* 2014; Krishnamurthy *et al.* 2016). Both, traffic intensity and dense human settlements have a non-linear relationship with resistance, suggesting that only very

349   high intensity traffic and high-density human settlements offered high resistance to movement

350   (Table 4).

351   Roads are known to negatively impact genetic diversity and differentiation in animal

species, especially for mammals and amphibians (Shirk *et al.* 2010; Holderegger & Di Giulio

2010). However, most of these studies have been carried out in developed/industrialized

countries where road density is high and many of the roads are fenced (Holderegger & Di Giulio

2010). Although India has the second largest road network in the world, 46% of the roads are

currently not surfaced (National Transport Policy Development Committee 2013)and very few

segments are fenced. As a result, even the existing national highways connecting major centers

may be permeable to tiger movement on stretches with low traffic volumes and when other

landscape features promote dispersal. However, this is bound to change in the future as roads

will be widened to accommodate increasing traffic volumes.

Future change in diversity

362   Genetic diversity reduced over time in all simulated scenarios. Results suggest that even

establishing corridors would not suffice to maintain current level of heterozygosity. Our results

are supported by Bay et al. (2013), where simulations of mitochondrial diversity revealed that

even with geneflow, the number of tigers essential to maintain current heterozygosity would

exceed the carrying capacity of PAs in peninsular India (Bay *et al.* 2013). Our simulations

revealed that both increasing tiger numbers and maintaining connectivity were essential for

preventing drastic reduction of heterozygosity.

Stepping-stone corridors preserve connectivity

Our simulations revealed that loss of forest cover due to diversion of land for agriculture

or infrastructure lead to high genetic differentiation. However, increasing number of tigers

within existing PAs and allowing clusters of individuals to survive outside PAs decreased the

observed genetic differentiation and inbreeding. Presence of breeding clusters of tigers outside

PAs also reduced the probability of extinction. These intervening clusters aid in dispersal

between the larger, more robust populations, thus forming 'stepping-stone corridors'.

Scenarios that incorporated stepping-stone corridors resulted in 10% higher genetic variation

and 6-86% lower extinction probability compared to those without.

Habitat restoration and protection is critical

Dinerstein et al. (2006) recommended restoring habitat to increase population

connectivity between tiger conservation landscapes. Our results demonstrated that such

habitat restoration to establish corridors reduced extinction probability by 68% (Scenario F8,

Figure 3a and 3b) and was critical for population persistence in the future. Such landscape

restoration may require careful selection of areas so as to benefit both people and wildlife

(Defries *et al.* 2007). This may be difficult to achieve between all pairs of PAs, especially those

that have negligible structural connectivity between them. Our results revealed that increasing

tiger numbers allowed large but currently degraded PAs (with few tigers) to achieve low

extinction probability, underscoring the importance of better PA management and protection.

Increased tiger numbers buffered against demographic stochasticity in small PAs and decreased

the overall extinction probability. This was best demonstrated in the scenario where we added

a buffer around the small PAs and extinction probability dropped by 23-70%. The government

of India is taking steps to notify 'Eco-sensitive Zones' around PAs to regulate developmental

activities (Mathur 2012). However, our results suggest that unless connectivity is restored and

stepping-stone populations protected (also suggested by (Chundawat *et al.* 2016)), even such

rescue effects would fail for small PAs when the landscape around them became unsuitable.

Such extinction debt poses a significant challenge for conservation while these tiger

populations still persist.

Low levels of inbreeding in Central Indian tigers of the future

Kenney et al. (2014) investigated the effect of inbreeding depression on population

viability in tigers and found that even populations as large as (with 63- 80 > 1 year old tigers)

the big populations in our study have a high future risk of extinction due to inbreeding

depression if connectivity was not maintained. Our results suggest that inbreeding did not

increase appreciably over the next 100 years ($F<0.25$ in all scenarios). Levels of inbreeding were

lower than those known to impact fitness in mammals based on studies in the wild and

captivity (Ralls & Ballou 1982; Keller 2002). Hence, we did not simulate the effect of inbreeding

depression on survival and extinction. However, further increase in the inbreeding co-efficient

over time may lead to inbreeding depression and increase extinction risk of even large

populations. Our simulations suggest that inbreeding coefficient is likely to increase faster if

tiger numbers do not increase in the future (Figure S12).

Model assumptions and caveats

Overall, our framework provides the first snapshot of the potential to combine

landscape genetics with spatially explicit simulations for tiger conservation. Future studies

should investigate how different land-use change data and dispersal strategies would affect

tiger connectivity in CIL. For example, future landscape change projection in our simulations

was based on landscape change over one decade (2001- 2012). Would longer decadal datasets

help us better understand LULC change in the CIL? Additionally, our model does not allow for

future land-use policy changes. For example, accelerated development could result in loss of

connectivity in less than 100 years. However, we do incorporate scenarios which demonstrate

the impacts of specific infrastructure projects currently being undertaken or envisaged in the

future (Scenarios F5-F7).

While modeling tiger dispersal, we did not consider the variation in behavioral response

of individuals to land use change and the probability of locating a corridor. Modeling how

individuals incorporate information about surrounding habitat while dispersing would improve

the simulation results (Colbert *et al.* 2009). However, such data is currently not available.

Future research should evaluate the influence of different dispersal strategies on tiger

connectivity. Finally, our simulations investigating tiger connectivity assume a maximum

dispersal distance, which is difficult to estimate in the wild. The two maximum dispersal

distances of 300 and 500km that we used were based on available data (Bowman *et al.* 2002;

Patil *et al.* 2011; Natesh *et al.* 2017). A possibility of longer than 500 km dispersal cannot be

ruled out, and hence we carried out simulations with a longer maximum dispersal distance (as

suggested in (Joshi *et al.* 2013), results in supplementary material). This longer threshold led to

lower genetic differentiation (in scenario F1) than currently observed, suggesting that this

particular dispersal scenario may be overestimating the probability of long distance dispersals.

However, even with such enhanced dispersal distances, the key populations that needing

management attention, the impact of infrastructure projects, and scenarios that lead to better

outcomes did not change (Supplementary Figures S6, S9-S12).

 Implications for Conservation Planning

Our results reveal that multiple actions like delineating and maintaining corridors, with

stepping-stone populations between PAs, along with increasing tiger numbers are necessary to

maintain long-term viable tiger populations at a landscape scale. Our results highlight the

immediate need for regional land-use management and planning exercises aimed at managing

tiger populations as a network of PAs connected with corridors.

Land-use change

Currently, infrastructure development does not incorporate conservation goals while

developing project plans, which could seriously undermine ongoing tiger conservation efforts

within the landscape. Nearly 50% of India's population is projected to live in cities by 2030

(World Bank Group 2015), and research has shown that built- up area increases faster than

population increase in urban areas (Sudhira *et al.* 2004). Along with urbanization, coal

requirement for electricity generation is projected to increase ~2.5 times by 2031-32

(Fernandes 2012). To provide better connectivity between cities and to accommodate

increasing road traffic (estimated to grow at about 13% per year over the next 20 years),

massive infrastructure development projects are being undertaken (National Transport Policy

Development Committee 2013). Our simulations reveal the impacts of upcoming development

on tiger connectivity. Ideally, expansion of the current road network should include realignment

of new roads to avoid critical tiger habitat (see (Raman 2011)). We show that widening national

highways without installing crossing structures (NH6 and NH7, Scenario F6) in an area critical to

tiger connectivity will increase genetic differentiation between populations on either side 19 to

65 fold. When alternative routes are not available, we strongly recommend planning and

installing mitigation structures (under and over-passes) for wildlife passage before roads are

built or widened. This is more economical than retrofitting existing roads and should be

considered during the environment impact assessment of new infrastructure projects (Glista *et*

al. 2009). Our results (Figure S9) highlight the importance of installing crossing structures for

maintaining connectivity (F_{ST} increases by 80% from an average 0.025 to 0.13 in the absence of

such structures). Currently, such regional level planning which aligns conservation goals with

developmental plans is in its infancy in India. Our results provide impetus to such efforts by

highlighting landscape variables that need to be considered while developing infrastructure

plans.

Forest diversion and mining

Diversion of forest for mining is one of the major causes for loss of forest cover and structural

connectivity within the CIL. Over the last 3 decades, 40% of the total forest land diverted has

been for mining (Centre for Science and Environment 2012). Coal mining alone accounted for

65% of the total land diverted for mining between 2007 and 2011 (Centre for Science and

Environment 2012). If, along with other minerals being mined, all the coal blocks in the study

landscape were opened for mining, the mined area and associated increase in built- up area

would lead to ~22 % higher extinction probability even for large PAs in proximity to these

blocks. We emphasize the urgent need to protect the delineated corridors (as in Qureshi *et al.*

2014) for preserving connectivity to prevent extinctions.

Conclusion

Several species globally face threats due to anthropogenic impacts, similar to the tigers

in our study area. Landscapes with high conservation value, especially in the tropics, have been

changing rapidly due to increasing human population and exploitation of natural resource

leading to conversion and degradation of habitat (Crooks *et al.* 2011; Elmhagen *et al.* 2015;

Newbold *et al.* 2015). Our results highlight the urgent need for informed development plans

that consider biodiversity and connected wildlife populations in addition to human

development goals. With habitats of most large mammals getting increasingly fragmented, our

approach of combining landscape genetics with forward-time simulations to estimate extinction

probability and loss of connectivity provides a valuable tool for conservation management.

Such an approach can help identify populations vulnerable to landscape change, improving

conservation and management of populations, species and landscapes to ensure long-term

persistence.

Acknowledgements

We are thankful to the Maharashtra, Madhya Pradesh and Chhattisgarh Forest

Departments for providing the necessary permissions and logistical support for this research.

We sincerely thank the forest department staff, field assistants and volunteers for their help

and support during the fieldwork. We thank the Indian Resource Center for Clark Labs at

Ashoka Trust for Research in Ecology and the Environment for letting us use their facilities. We

thank V. V. Robin, Meghana Natesh, Y. V. Jhala and an anonymous reviewer for helpful

comments on the manuscript. We are grateful to the National Tiger Conservation Authority

(NTCA) of India for partially funding this study. We thank the National Centre for Biological

Sciences (NCBS) for institutional support. PT was supported by Council for Industrial and

Scientific Research fellowship and NCBS. UR was supported by the DAE outstanding Scientist

Award.

References

Adamack AT, Gruber B (2014) PopGenReport⬚: simplifying basic population genetic analyses in

R (S Dray, Ed,). *Methods in Ecology and Evolution*, **5**, 384–387.

Areendran G, Rao P, Raj K, Mazumdar S, Puri K (2013) Land use / land cover change dynamics

analysis in mining areas of Singrauli district in Madhya Pradesh, India. *Tropical Ecology*, **54**,

239–250.

Athreya V, Navya R, Punjabi G a *et al.* (2014) Movement and activity pattern of a collared

tigress in a human-dominated landscape in central India. *Tropical Conservation Science*, **7**,

75–86.

Balloux F (2001) EASYPOP (Version 1.7): A Computer Program for Population Genetics

Simulations. *Journal of Heredity*, **92**, 301–302.

Barnosky AD, Matzke N, Tomiya S *et al.* (2011) Has the Earth's sixth mass extinction already

arrived? *Nature*, **471**, 51–7.

Bay RA, Ramakrishnan U, Hadly EA (2013) A call for tiger management using "reserves" of

genetic diversity. *The Journal of heredity*, **105**, 295–302.

Benson JF, Mahoney PJ, Sikich JA *et al.* (2016) Interactions between demography, genetics, and

landscape connectivity increase extinction probability for a small population of large

carnivores in a major metropolitan area. *Proceedings of the Royal Society of London B:*

Biological Sciences, **283**.

Boersen MR, Clark JD, King TL (2003) Estimating black bear population density and genetic

diversity at Tensas River, Louisiana using markers microsatellite. *Wildlife Society Bulletin*,

31, 197–207.

Bowcock AM, Ruiz-Linares A, Tomfohrde J *et al.* (1994) High resolution of human evolutionary

trees with polymorphic microsatellites. *Nature*, **368**, 455–7.

Bowman J, Jaeger J a. G, Fahrig L (2002) Dispersal Distance of Mammals Is Proportional To

Home Range Size. *Ecology*, **83**, 2049–2055.

Brown JL, Weber JJ, Alvarado-Serrano DF *et al.* (2016) Predicting the genetic consequences of

future climate change: The power of coupling spatial demography, the coalescent, and

historical landscape changes. *American journal of botany*, **103**, 153–63.

Caragiulo A, Pickles RSA, Smith JA *et al.* (2015) Tiger (Panthera tigris) scent DNA: a valuable

conservation tool for individual identification and population monitoring. *Conservation

Genetics Resources*, **7**, 681–683.

Centre for Science and Environment (2012) *Forest and Environment Clearances*. Centre for

Science and Environment, New Delhi, India.

Chundawat RS, Sharma K, Gogate N, Malik PK, Tamim A (2016) Size matters⍰: Scale mismatch

between space use patterns of tigers and protected area size in a Tropical Dry Forest.

Biological Conservation, **197**, 146–153.

Colbert J, Le Galliard JF, Cote J, Meylan S, Massot M (2009) Informed dispersal, heterogeneity in

animal dispersal syndromes and the dynamics of spatially structured populations. *Ecology

Letters, **12**, 197–209.

Crooks KR, Burdett CL, Theobald DM, Rondinini C, Boitani L (2011) Global patterns of

fragmentation and connectivity of mammalian carnivore habitat. *Philosophical*

transactions of the Royal Society of London. Series B, Biological sciences, **366**, 2642–51.

Defries RS, Hansen A, Turner BL, Reid R, Liu J (2007) Land use change around protected areas:

management to balance human needs and ecological function. *Ecological Applications*, **17**,

1031–1038.

Dieringer D, Schlotterer C (2003) microsatellite analyser (MSA): a platform independent analysis

tool for large microsatellite data sets. *Molecular Ecology Notes*, **3**, 167–169.

Dinerstein E, Loucks C, Heydlauff A *et al.* (2006) *Setting Priorities for the Conservation and*

Recovery of Wild Tigers: 2005–2015. A User's Guide. WWF, WCS, Smithsonian, and NFWF-

STF, Washington, D.C. – New York.

Dirzo R, Young HS, Galetti M *et al.* (2014) Defaunation in the Anthropocene. *Science*, **345**, 401–

406.

Dooley MP, Folkerts-landau D, Garber PM (2014) *The revived Bretton Woods system's first*

decade. National Bureau of Economic Research, Cambridge.

Dutta T, Sharma S, Mcrae BH, Sarathi P, Defries R (2015) Connecting the dots : mapping habitat

connectivity for tigers in central India. *Regional Environmental Change*.

Elmhagen B, Eriksson O, Lindborg R (2015) Implications of climate and land-use change for

landscape processes, biodiversity, ecosystem services, and governance. *Ambio*, **44**, S1–S5.

van Etten J (2012) R package gdistance: distances and routes on geographical grids (version 1.1-

1).

Fernandes A (2012) *How coal mining is thrashing tigerland*. Greenpeace, India.

Forest Survey of India (2015) *India State of Forest Report 2015*. Forest Survey of India,

Dehradun, India.

Frankham R (1996) Relationship of Genetic Variation to Population Size in Wildlife.

Conservation Biology, **10**, 1500–1508.

Glista DJ, DeVault TL, DeWoody JA (2009) A review of mitigation measures for reducing wildlife

mortality on roadways. *Landscape and Urban Planning*, **91**, 1–7.

Global Tiger Recovery Program (2011) *Global Tiger Recovery Program 2010- 2022*. The World

Bank, Washington, DC.

Goudet J (2005) hierfstat, a package for r to compute and test hierarchical F-statistics.

Molecular Ecology Notes, **5**, 184–186.

Hagen SB, Kopatz A, Aspi J, Kojola I, Eiken HG (2015) Evidence of rapid change in genetic

structure and diversity during range expansion in a recovering large terrestrial carnivore.

Proceedings. Biological sciences / The Royal Society, **282**, 20150092.

Holderegger R, Di Giulio M (2010) The genetic effects of roads: A review of empirical evidence.

Basic and Applied Ecology, **11**, 522–531.

Ihaka R, Gentleman R (2012) R: A Language for Data Analysis and Graphics. *Journal of*

Computational and Graphical Statistics.

Jackson CR, Marnewick K, Lindsey PA, Røskaft E, Robertson MP (2016) Evaluating habitat

connectivity methodologies: a case study with endangered African wild dogs in South

Africa. *Landscape Ecology*, **31**, 1433–1447.

Jhala, Y. V.;Qureshi, Q.;Gopal R. (2015) *Status of tigers in India, 2014*. National Tiger

Conservation Authority, New Delhi & The Wildlife Institute of India, Dehradun.

Jombart T (2008) adegenet: a R package for the multivariate analysis of genetic markers.

Bioinformatics (Oxford, England), **24**, 1403–5.

Joshi A, Vaidyanathan S, Mondol S, Edgaonkar A, Ramakrishnan U (2013) Connectivity of Tiger

(Panthera tigris) Populations in the Human-Influenced Forest Mosaic of Central India. *PLoS*

ONE, **8**, e77980.

Keller L (2002) Inbreeding effects in wild populations. *Trends in Ecology & Evolution*, **17**, 230–

241.

Krishnamurthy R, Cushman SA, Sarkar MS *et al.* (2016) Multi-scale prediction of landscape

resistance for tiger dispersal in central India. *Landscape Ecology*, **31**, 1355–1368.

Landguth EL, Muhlfeld CC, Waples RS *et al.* (2014) Combining demographic and genetic factors

to assess population vulnerability in stream species. *Ecological Applications*, **24**, 1505–

1524.

Laurance WF, Balmford a. (2013) A global map for road building. *Nature*, **495**, 308–309.

Lele S, Rosencranz A (2008) Supreme Court and India's Forests. *Economic and Political Weekly*,

11–14.

Mathur VB (2012) *India's action plan for implementing the convention on biological diversity's*

programme of work on protected areas. Wildlife Institute of India, Dehradun; Ministry of

Environment and Forests, New Delhi, India.

Ministry of Environment and Forests (2008) *National Action Plan on Climate Change*. Ministry of

Enivronment and Forests, New Delhi, India.

Ministry of Mines (2011) *Unlocking the Potential of the Indian Minerals Sector*. Strategy Paper

for Ministry of Mines, New Delhi, India.

Mondol S, Karanth KU, Ramakrishnan U (2009a) Why the Indian subcontinent holds the key to

global tiger recovery. *PLoS genetics*, **5**, e1000585.

Mondol S, Ullas Karanth K, Samba Kumar N *et al.* (2009b) Evaluation of non-invasive genetic

sampling methods for estimating tiger population size. *Biological Conservation*, **142**, 2350–

2360.

Morrison JC, Sechrest W, Dinerstein E, Wilcove DS, Lamoreux JF (2007) Persistence of Large

Mammal Faunas as Indicators of Global Human Impacts. *Journal of Mammalogy*, **88**, 1363–

1380.

Mukherjee N, Mondol S, Andheria A, Ramakrishnan U (2007) Rapid multiplex PCR based species

identification of wild tigers using non-invasive samples. *Conservation Genetics*, **8**, 1465–

1470.

Natesh M, Alta G, Nigam P *et al.* (2017) Conservation priorities for endangered Indian tigers

through a genomic lens. *Scientific Reports*, **7**.

National Transport Policy Development Committee (2013) *Trends in Growth and development

of transport*. Planning Commission, India.

Newbold T, Hudson LN, Hill SLL *et al.* (2015) Global effects of land use on local terrestrial

biodiversity. *Nature*, **520**, 45–50.

Patil N, Samba Kumar N, Gopalaswamy AM, Ullas Karanth K (2011) Dispersing tiger makes a

point. *Oryx*, **45**, 472–475.

Prakash A, Gupta RP (2016) Land-use mapping and change detection in a coal mining area - a

case study in the Jharia coalfield , India. *International Journal of Remote Sensing*, **19**, 391–

410.

Qureshi Q, Saini S, Basu P *et al.* (2014) *Connecting tiger populations for long-term conservation.*

National Consevation Authority of India & Wildlife Institute of India, Dehradun. TR2014-02.

Ralls K, Ballou J (1982) Effect og inbreeding on juvenile mortality in some small mammal

species. *Laboratory Animals*, **16**, 159–166.

Raman TRS (2011) *Framing ecologically sound policy on linear intrusions affecting wildlife*

habitats: Background paper for the national board of wildlife. Nature Conservation

Foundation, Mysore.

Reddy PA, Cushman SA, Srivastava A, Sarkar MS, Shivaji S (2017) Tiger abundance and gene flow

in Central India are driven by disparate combinations of topography and land cover.

Diversity and Distributions, 1–12.

Ripple WJ, Beschta RL (2012) Trophic cascades in Yellowstone: The first 15years after wolf

reintroduction. *Biological Conservation*, **145**, 205–213.

Ripple WJ, Estes JA, Beschta RL *et al.* (2014) Status and ecological effects of the world's largest

carnivores. *Science*, **343**, 1241484.

Saccheri I, Kuussaari M, Kankare M, Vikman P, Hanski I (1998) Inbreeding and extinction in a

646  butterfly metapopulation. *Nature*, **392**, 491–494.

647 Sharma S, Dutta T, Maldonado JE *et al.* (2013a) Selection of microsatellite loci for genetic

648  monitoring of sloth bears. *Ursus*, **24**, 164–169.

649 Sharma S, Dutta T, Maldonado JE *et al.* (2013b) Forest corridors maintain historical gene flow in

650  a tiger metapopulation in the highlands of central India. *Proceedings of the Royal Society B:*

651  *Biological Sciences*, **280**, 20131506.

652 Shirk AJ, Wallin DO, Cushman SA, Rice CG, Warheit KI (2010) Inferring landscape effects on gene

653  flow: a new model selection framework. *Molecular Ecology*, **19**, 3603–3619.

654 Smouse PE, Long JC, Sokal RR (1986) Multiple Regression and Correlation Extensions of the

655  Mantel Test of Matrix Correspondence. *Systematic Zoology*, **35**, 627.

656 Spielman D, Brook BW, Briscoe DA, Frankham R (2004) Does Inbreeding and Loss of Genetic

657  Diversity Decrease Disease Resistance? *Conservation Genetics*, **5**, 439–448.

658 Srivastava R, Tyagi R (2016) Wildlife corridors in india. *Environmental Law Review*, **18**, 205–223.

659 Sudhira HS (2011) *Urban landcover and landcover change dataset of Indian cities*. Indian

660  Institute of Human Settlements Working Paper, Mimeo.

661 Sudhira HSS, Ramachandra TV V., Jagadish KSS (2004) Urban sprawl: metrics, dynamics and

662  modelling using GIS. *International Journal of Applied Earth Observation and*

663  *Geoinformation*, **5**, 29–39.

664 Sutherland GD, Harestad AS, Price K, Lertzman KP (2000) Scaling of Natal Dispersal Distances in

665  Terrestrial Birds and Mammals. *Conservation Ecology*, **4**, 16.

Tian Y, Wu J, Smith AT *et al.* (2011) Population viability of the Siberian Tiger in a changing

landscape: Going, going and gone? *Ecological Modelling*, **222**, 3166–3180.

Tian Y, Wu J, Wang T, Ge J (2014) Climate change and landscape fragmentation jeopardize the

population viability of the Siberian tiger (Panthera tigris altaica). *Landscape Ecology*, **29**,

621–637.

Trinkel M, Cooper D, Packer C, Slotow R (2011) Inbreeding depression increases susceptibility to

bovine tuberculosis in lions: an experimental test using an inbred-outbred contrast

through translocation. *Journal of wildlife diseases*, **47**, 494–500.

United Nations, Department of Economic and Social Affairs, Population Division (2014) *World*

Urbanization Prospects: The 2014 Revision, Highlights (ST/ESA/SER.A/352).

Wikramanayake E, Dinerstein E, Loucks C *et al.* (2010) *Roads to Recovery or Catastrophic Loss*:

How Will the Next Decade End for Wild Tigers? (R Tilson, P Nyhus, Eds,). Elsevier Inc.

Wildlife institute of India (2016) *Eco-friendly measures to mitigate impacts of linear*

infrastructure on wildlife. Wildlife Insitute of India, Dehradun, India.

Winter DJ (2012) MMOD: an R library for the calculation of population differentiation statistics.

Molecular ecology resources, **12**, 1158–60.

World Bank Group (2015) *Population Estimates and Projections*. The World Bank, Washington,

DC.

Yadav PK, Kapoor M, Sarma K (2012) Land Use Land Cover Mapping , Change Detection and

Conflict Analysis of Nagzira-Navegaon Corridor , Central India Using Geospatial

Technology. *International Journal of Remote Sensing and GIS*, **1**, 90–98.

Yumnam B, Jhala Y V, Qureshi Q *et al.* (2014) Prioritizing tiger conservation through landscape

genetics and habitat linkages. *PloS one*, **9**, e111207.

Figures and Tables

Figure 1. Study area: The figure contains a map of India with the study landscape highlighted

and enlarged. In the enlarged study area, protected areas are marked by black outline and

genetically identified individual tigers as red dots.

Figure 2. Structural connectivity, allelic richness, inbreeding and extinction after 100 years

under selected management scenarios: The 3 management scenarios in this figure have

corresponding panels with two sub-scenarios (a- tiger number does not increase and d-

increase in tiger number) and 4 plots each representing management outcomes for (L to R): 1-

structural connectivity, 2-allelic richness, 3-inbreeding estimate, and 4-extinction probability.

The structural connectivity is the same for sub-scenarios a and d. Hence there is a single

structural connectivity plot for each of the management outcomes. Completely isolated PAs

(scenario F9) are represented in black. The outcomes for different scenarios can be compared

within a column. The coloured areas in each plot are the protected areas and their colour

represents the scenario-specific outcome. The gradient from blue to red represents low to high

(structural connectivity, allelic richness, inbreeding coefficient and extinction probability). Blank

PAs in the plots within columns 2 and 3 are the PAs which go extinct by the year 2100. Blank

PAs in column 4 (extinction probability plots) are the ones which do not show extinction in any

of the 100 replicate simulation runs.

Figure 3. Extinction Probability

(a) Matrix representing extinction probability for each population for each of the scenarios after

100 years- dispersal threshold 300km (b) Matrix representing extinction probability for each

population for each of the scenarios after 100 years- dispersal threshold 500km. x-axis

represents the PAs and y axis represents the scenarios and sub-scenarios (c) Scatterplot of

population size (average of current size and size in sub-scenario d) vs. isolation index calculated

as the average cost distance between populations. Each point represents a population and the

colour represents it's average extinction probabilityacross scenarios. NOR, TAM, BAR do not

have any tigers currently and were only included in the simulation sub-scenario d where tiger

numbers increase. Hence they do not have colour in panels (a) and (b) for scenarios a, b and c.

Scenarios: F1-No change in landscape, F2- Forest cover constant, F3- Area under agriculture

constant, F4- Unconstrained landscape change, F5- Mines, F6- NH6 and NH7 as barriers, F7-

NH6 and NH7 as barriers with gaps, F8- Corridors, F9- PAs fenced. Sub- scenarios: a- Tigers only

inside protected areas, b- Tigers inside and outside (random), c- Tigers inside and outside

(clustered), d- Tiger numbers increase.

Table 1. Landscape change scenarios for forward time simulations.

Scenario	Description	Rationale
F1- No landscape change*	Status quo	Null scenario
F2- Forest cover constant	Landscape change modeled while keeping the forest cover constant	The Green India mission under the National Action Plan on Climate Change (Ministry of Environment and Forests 2008) advocates achieving a forest cover of 33%. Current forest cover is 21% (Forest Survey of India 2015). In 1996, the supreme court of India redefined the scope of Forest Conservation Act 1980 and banned tree felling inside forests across India (Lele & Rosencranz 2008).
F3- Agriculture area constant	Landscape change modeled while keeping the area under agriculture constant	Global food demand is projected to double by 2050. Even if use of technology to intensify agriculture increases yield, area under agriculture is not expected to reduce in the future(Laurance & Balmford 2013)
F4- Unrestricted change*	Landscape change modeled based on change from 2000 to 2012	India's Gross Domestic Product (GDP) growth rate was higher than ever before in the decade from 2001- 2011. Although this rate reduced after 2011, the recent government's development driven policies are likely to increase the economic growth rate (Dooley *et al.* 2014). The rate of granting forest clearances has also been highest from 2002- 2011 within the last three decades. 387952 hectare of forest land was

		diverted during this decade for defense, mining, irrigation, power projects, industries and infrastructure projects (Centre for Science and Environment 2012)
F5- Effect of mines and associated landuse change	In order to evaluate the effect of mines, we let the rest of the landscape remain constant (like in F1) except for the increase in area of mines and associated built-up area over the next 100 years. The mining area (mine+ built-up) increased 3.6 times every 20 years based on a study in central India (Areendran *et al.* 2013)	The central Indian region is rich in mineral deposits. The mining sector currently contributes ~2% to India's GDP and the Ministry of Mines, Government of India has targeted to increase this share to 5% of GDP (Ministry of Mines 2011). The Government of India amended the Mines and Minerals (Development and Regulation) Act in 2015 in order to expedite environmental clearances and issuance of licenses. This amendment also provides for the creation of District Mineral Foundations (DMF) to work towards developing mining affected areas. Research in Central India has shown that mining leads to landuse change and an increase in built-up areas around mines (Areendran *et al.* 2013; Prakash & Gupta 2016) and the setting up of DMF will only increase the rate of this conversion. Data on mines were obtained from the Ministry of Mines database (http://mcas.nic.in/Mining_plan_web_Query.asp) and Greenpeace India (Fernandes 2012). The data from Ministry of Mines were converted into a spatial layer based on their GPS locations and area of the mines. Data on coal mines described in (Fernandes 2012) was obtained as shapefiles through personal communication.
F6- Highways as barriers	Landscape does not change except two national highways (NH6 and NH7) which cut across the landscape are converted into barriers to movement	Road traffic is estimated to grow at about 13 % per annum over the next 20 years (National Transport Policy Development Committee 2013). NH7, which runs north to south, bisects a critical corridor in the landscape and has recently been cleared to be widened from two to four lane capacity. NH6, which runs east to west and bisects another critical corridor, is also being considered for widening. Yadav et al. (2012) have observed agriculture and built-up area encroachment along NH6 in the forested area which connects two PAs (NGZ and NAW) (Yadav *et al.* 2012), thus potentially increasing the resistance to movement of tigers. This scenario is

		a case study to specifically look at the effects of these highways, if they were to become barriers in the future, on the corridors they bisect.
F7- Highways as barriers with wildlife crossings	Landscape does not change except two national highways (NH6 and NH7) which cut across the landscape are converted into barriers with provision for wildlife crossing at points where they bisect critical corridors. There was one gap in each highway. The gaps were 1 pixel wide and were placed near the locations where the NH6 and NH7 are currently narrow (2lanes wide, without a divider in between) and there is forest cover on the either side.	Guidelines to mitigate impacts of linear infrastructure on wildlife recommend a 100 m mitigation structure per 1 km length of infrastructure in critical tiger corridors (Wildlife institute of India 2016). Although we cannot test the effectiveness of different kinds of structures which can provide connectivity across roads in this simulation, we investigate the effect of having a gap in the barrier which can potentially maintain connectivity.
F8- Habitat restoration to establish corridors between all PAs	The corridors were designated based on the least cost paths (generated using the gdistance package(van Etten 2012) in R (Ihaka & Gentleman 2012)) between PAs and the proposed corridor between Kanha and Pench.	Restoration of habitat and establishing corridors between PAs has been recommended to maintain and even increase the connectivity in the landscape (Dinerstein *et al.* 2006; Dutta *et al.* 2015). We generated least cost paths between PAs and we assigned a value of 1 to all the pixels in these paths (equivalent to forest). All non-forest pixels within the least cost paths were converted to forest to represent restoration of habitat. These paths overlap with the corridors delineated in previous studies (Qureshi *et al.* 2014; Dutta *et al.* 2015). We test how beneficial establishing these corridors would be.
F9- Fenced PAs	All the protected areas have fence around them in the future preventing dispersal	Extreme scenario to investigate the effect of fencing on genetic variation and extinctions in the future.

731   * We carried out additional simulations for scenarios F1 and F4 where we added a 5 km buffer

732   around the smaller PAs in the landscape (< 400 km^2). We evaluated the effect of increased PA

733   size on connectivity and extinction estimates.

734   Table 2. Genetic differentiation- DPS

	BTR (n= 20)	BPR (n= 18)	BOR (n= 4)	KTR (n= 22)	CHA (n=8)	PTR (n= 15)	STR (n= 4)	TATR (n= 26)	UK (n= 5)
BTR	0								
BPR	0.629	0							
BOR	0.669	0.456	0						
KTR	0.548	0.597	0.641	0					
CHA	0.621	0.286	0.359	0.592	0				
PTR	0.501	0.627	0.612	0.351	0.637	0			
STR	0.541	0.612	0.598	0.3541	0.592	0.416	0		
TATR	0.639	0.213	0.413	0.571	0.267	0.612	0.591	0	
UK	0.700	0.358	0.373	0.624	0.474	0.635	0.624	0.322	0

736   Table 3. Univariate optimization results

Univariate Optimization					
Landscape Variable	Maximum Resistance	Shape Parameter	Mantel's r	Partial Mantel's r	Significance (partial)

	(R_{max})	(x)			
Human Settlement	10	0.001	0.859	0.778	0.001
Land-use	10000	10	0.806	0.678	0.001
Linear density	10	0.01	0.683	0.383	0.027
Roads (traffic)	10000	10	0.776	0.603	0.001
Multivariate Optimization					
Human Settlement	1000	5	0.8166	0.7857	0.001
Land-use	100	50			
Roads (traffic	1000	10			

737 Non-linear exponential transformations were the best models for all these variables