Phylogenetic Characterization of the Energy-taxis Receptor Aer in *Pseudomonas* and Phenotypic

Characterization in *P. pseudoalcaligenes* KF707

Sean C. Booth[a]#* and Raymond J. Turner[a]#

Department of Biological Sciences, University of Calgary, Calgary, AB, Canada[a]

#Address correspondence to Raymond J. Turner, turnerr@ucalgary.ca and Sean C. Booth,

scbooth@ucalgary.ca

* Current address: Department of Zoology, Department of Biochemistry, University of Oxford,

UK

+1-403-220-3581

Running Title: *Pseudomonas* Energy-Taxis Receptors

Keywords: chemotaxis, *Pseudomonas*, aerotaxis, energy-taxis, phylogenetics, motility

Abstract

Chemotaxis allows bacteria to sense gradients in their environment and respond by directing their swimming. Aer is a receptor that, instead of responding to a specific chemoattractant, allows bacteria to sense cellular energy levels and move towards favourable environments. In *Pseudomonas,* the number of apparent Aer homologs differs between the only two species it had been characterized in, *P. aeruginosa* and *P. putida.* Here we combined bioinformatic approaches with deletional mutagenesis in *P. pseudoalcaligenes* KF707 to further characterize Aer. It was determined that the number of Aer homologs varies between 0-4 throughout the *Pseudomonas* genus, and they were phylogenetically classified into 5 subgroups. We also used sequence analysis to show that these homologous receptors differ in their HAMP signal transduction domains. Genetic analysis also indicated that some Aer homologs have likely been subject to horizontal transfer. *P. pseudoalcaligenes* KF707 was unique among species for having three Aer homologs as well as the receptors CttP and McpB. Phenotypic characterization in this species showed the most prevalent homolog of Aer was key, but not essential for energy-taxis. This study demonstrates that energy-taxis in *Pseudomonas* varies between species and provides a new naming convention and associated phylogenetic details for Aer chemoreceptors.

Introduction

Chemotaxis is the ability to sense and swim along chemical gradients, and is a widespread, important behaviour in bacteria [1]. Canonically, it functions by extracellular compounds interacting with membrane-bound chemoreceptors, generally called methyl-accepting chemotaxis proteins (MCPs), causing a phosphorylation signal cascade through CheA and CheY to alter the direction of flagellar rotation [1], which in turn allows the cell to direct swimming through concentration gradients. The first chemoreceptors were characterized in *E. coli*, but now many more have been described, particularly in *Pseudomonas* [2]. Many receptors for specific ligands are being identified through a high-throughput approach [3], but some specialized receptors do not recognize extracellular signals.

Energy-taxis is a behavior in bacteria that enables swimming towards optimal environments for producing energy by sensing intracellular signals [4]. This is closely related to aerotaxis (swimming towards oxygen) as in *E. coli*, a receptor was discovered which enabled taxis towards air bubbles [5]. This inner membrane receptor was named 'Aer' though it does not bind oxygen, but instead enables taxis towards metabolizable carbon sources [6] by detecting the cellular energy state via a cytosolic PAS (Per-Arnt-Sim) domain with a flavin adenine dinucleotide (FAD) co-factor [7]. A similar receptor was also discovered in *P. aeruginosa*, along with a second distinct protein, named Aer-2 as both contributed to aerotaxis [8]. Aer-2 is a cytosolic protein that binds oxygen through a heme group [9]. In *P. putida* three *aer*-like genes were found, but only one was important for energy-taxis [10]. This was also named 'Aer2' so we will refer to the cytosolic receptor Aer-2 as McpB hereafter. It has been shown to bind oxygen, but does not affect flagellar activity as it is key to the assembly of, and signals through the Che2 chemosensory system (whose function is unknown) [11–13].

Aer has been characterized by gene inactivation in *P. putida* PRS2000 [14], *P. aeruginosa*
PA01 [8], *P. putida* KT2440 [10] and *P. putida* F1 [15]. The *P. putida* species all have 3 *aer-*
like genes but not *mcpB*, whereas *P. aeruginosa* only has one, indicating there is variation within
the genus for both loci. The species our group studies, *P. pseudoalcaligenes* KF707, has 3 *aer-*
like genes despite being more closely related to *P. aeruginosa* than *P. putida* [16, 17] and three
chemosensory clusters, corresponding to the *P. aeruginosa* F6, F7 and pil-chp systems [18],
making it possible to investigate the function of multiple *aer*-like genes and *mcpB* in the same
organism. In *P. aeruginosa*, McpB was originally implicated with Aer as an aerotaxis receptor
[8], but its function remains uncertain as despite its ability to directly bind oxygen, its signal is
not transduced to the flagellum [9]. The *mcpB* gene is part of the *che2* gene cluster, which is
preceded by another chemoreceptor, *mcpA* [13]. This gene was demonstrated to mediate positive
chemotaxis towards tetrachloroethylene and renamed *cttP* [19]. As CttP has no obvious ligand-
binding region and in *E. coli,* its MCPs can mediate attractant and repellent responses to phenol
that do not involve their ligand binding regions [20]; the observed taxis to TCE may also have
been fortuitous, indicating that CttP has some other function. Additionally, CttP co-localizes
with the che (F7) chemosensory system while McpB is necessary for the formation of the che2
(F6) chemosensory complex [13] which led us to question whether CttP and McpB may
somehow influence energy-taxis.

In this study we present a combination of a bioinformatic characterization of Aer throughout
the *Pseudomonas* genus with a genetic knockout characterization of three Aer homologs, McpB
and CttP in *P. pseudoalcaligenes* KF707. A phylogeny of Aer was built using sequences
obtained from 65 *Pseudomonas* species providing insight into the distribution throughout the
genus and enabling the definition of five 'Aer' groups. Only *P. pseudoalcaligenes* KF707 had

the unique feature of possessing three Aer homologs, McpB and CttP making it the ideal

candidate to investigate if these receptors had related functions. Using single and combinatory

deletion mutants, we found that the most common *Pseudomonas* Aer homolog had the most

influence on energy-taxis. Together these results provide a definition of the Aer energy-taxis

receptor as a family with varied distribution in *Pseudomonas* and implicate the most common

Aer homolog as the primary energy-taxis receptor in *P. pseudoalcaligenes* KF707.

<u>Results</u>

To investigate how the number of Aer-like receptors, differences in their amino acid

sequences and how they are related phylogenetically, Aer sequences from 65 *Pseudomonas*

species were analyzed. Species with completely sequenced genomes were selected, but for those

with many strains only a few representatives that have been highly studied were included (e.g. *P.*

aeruginosa). Species with incomplete (draft) sequences were also included in an attempt to

ensure representation from all major *Pseudomonas* clades [16, 17]. 144 protein sequences were

obtained from the NCBI database using the *P. aeruginosa* PA01 Aer sequence (NP_250252.1) as

a BLAST [21] query sequence. All hits with >95% sequence coverage were included. These

sequences were aligned using COBALT [22] as this alignment algorithm ensures that conserved

domains are aligned despite a lack of similarity elsewhere in the sequence.

From this alignment a maximum likelihood phylogeny was constructed using PHYML

(Supplementary Figure 1) [23, 24]. Two additional phylogenies were generated, one including

Aer from *E. coli* K12, and one including seventeen Aer-like sequences from the eight closest

related species to *Pseudomonas* [25]. The topology of these two unrooted phylogenies

(Supplementary Figures 2 and 3) were identical to the *Pseudomonas*-only tree, which was rooted

based on the similar placement of the outgroups in both trees (Supplementary Figure 4). The

rooted phylogeny allowed clear delineation of Aer sequences into subgroups.

Phylogenetic Grouping of Aer

The generated phylogeny implied that there are several sub-families of Aer homologs

(Figure 1B). These groups were sub-aligned by themselves, then analyzed to determine if the

group assignments were accurate. Sequence Harmony/Multi Relief [26] was used to determine,

for each position, whether that AA is conserved within each group, and whether it is divergent

between the two groups (Supplementary Figure 5). Each pair of groups was compared in this

fashion and most intergroup comparisons indicated that 25-35% of AAs were above the

distinction cut-off, indicating the group distinctions were correct. Conversely, two proposed

subgroupings of Aer.g1 were demonstrated to differ from the rest of Aer.g1 by much less than

any other between-group comparisons as only 10% of AAs were above the cut-off. These two

groups were thus included as part of Aer.g1 (named Aer.g1A and Aer.g1B), reducing the total

number of groups to five. Throughout this process, Aer.g5 was consistently excluded as it only

contains 2 sequences, which are highly divergent from all others.

Distribution of Groups within *Pseudomonas*

To determine the prevalence of each Aer homolog group within the *Pseudomonas* genus,

the number of homologs from each group that each species possessed was counted and a

hierarchically clustered heatmap was generated (Supplementary Figure 6). This analysis showed

that all included species, except *P. denitrificans* ATCC13867, had an Aer.g1 homolog.

Conversely, only species related to *P. aeruginosa* had McpB and CttP, which were always found

together. There were very few species that had McpB/CttP and multiple Aer homologs, only

Pseudomonas sp 21, *P. nitroreducens* HBP1, *P. resinovorans* NBRC 106553 and *P.*

pseudoalcaligenes KF707 fit this category. Aer.g1 was duplicated only in *P. fluorescens* and

related subgroups 1-8 [17], and these duplications matched the prior subdivision of Aer.g1 into

Aer.g1A and Aer.g1B. Duplications of Aer.g2 occurred only (and always) in *P. stuzeri* and the

closely related *P. balearica*. Unlike the other duplications, these were tandem duplications.

Duplications of Aer.g3 only occurred in *P. protegens* CHA0 and Pf-5.

Relationship between *Pseudomonas* and Aer Phylogenies

To understand how the phylogeny of Aer related to the phylogeny of *Pseudomonas*, a

species phylogeny was generated (Figure 1A) using concatenated alignments of *gyrB*, *rpoB*, and

rpoD. This phylogeny matched well with previously published *Pseudomonas* phylogenies [17,

27]. Examination of how the various Aer subgroups, as well as McpB/CttP, mapped onto the

genus phylogeny revealed their likely evolutionary trajectories (Figure 1A). The presence of

aer.g1 in all species, except *P. denitrificans* indicates that it is a basal *Pseudomonas* gene. Both

subdivisions of Aer.g1 occurred only in the more derived *P. fluorescens* subgroup. Conversely,

aer.g2 is distributed throughout the genus with no clear pattern. It is present in the more ancestral

P. stutzeri group as well as the more derived *P. fluorescens* subgroup but is not present in many

species in between. *aer*.g3 is present, except for *P. protegens,* exclusively in the *P. putida* group,

though not all members of the clade have it. *aer*.g4 is present mostly in the related *P. straminea,*

P. oleovorans, P. putida, P. lutea and *P. syringae* subgroups. Notably, it is not present in the *P.*

chlororaphis or *P. fluorescens* subgroups, despite these being more derived than the

aforementioned groups. *aer.g4* was also found in *P. resinovorans* and *P. pseudoalcaligenes*

KF707, but not the rest of their *P. aeruginosa* subgroup. *aer.g5* is only present in two species,

both in the *P. oleovorans* subgroup. *mcpB* and *cttP*, which were always found together, were

exclusively present in the *P. aeruginosa* subgroup.

Figure 1: Phylogeny of select *Pseudomonas* species (A) and their Aer protein sequences (B). Species phylogeny is based on nucleotide

sequences of concatenated *gyrB*, *rpoB* and *rpoD*. Numbers along branches indicate bootstrap

support values from 100 replicates and scale bar indicates average number of nucleotide changes

per position. Colours indicate the presence of particular Aer groups, matching the grouping

defined by the cladogram (B), which was confirmed in subsequent analysis. The root for the Aer

protein tree was estimated to be between Aer.g2 (dotted outline) and the other groups based on

inspection of unrooted phylogenies including Aer from *E. coli* or similar sequences from closely

related non-*Pseudomonas* species. See Supplementary Figures 1-3 for details. For both trees,

nodes with support values below 50 were collapsed. Previously characterized Aer sequences

from *P. aeruginosa* (triangle) and *P. putida* (squares) are indicated, as well as those from *P.*

pseudoalcaligenes KF707 (black stars).

<u>Genetic Features of *aer* Homologs</u>

The grouping of Aer into 5 subgroups based on protein sequence information was

confirmed and expanded on by examining features of the underlying *aer* genes. The upstream

and downstream regions of the *aer* genes were inspected to determine whether the genomic

context was consistent across species (Figure 2). Additionally, the first two upstream and

downstream genes of each *aer* gene were identified (Supplementary Table 1). The frequency of

each of the associated genes and the general genomic context for each Aer homolog were thus

examined to identify the most commonly associated genes (summarized in Figure 2). Each

homolog was part of an apparently unique gene cluster: Aer.g1 with an aconitate hydratase (in

69% of species) and a CAAX amino protease (33%), Aer.g2 with a PAS-containing diguanylate

cyclase/phosphodiesterase (88%), Aer.g3 with a different PAS-containing diguanylate

cyclase/phosphodiesterase (88%) and Aer.g4 with a LysR-type transcriptional regulator (64%)

and a DTW-domain containing protein (74%). Beyond these, the second upstream and

downstream genes tended to vary more, though some upstream genes were clearly conserved:

Aer.g1, 23S rRNA methyl transferase (42%); Aer.g3, C4 dicarboxylate transporter (63%);

Aer.g4, agmatinase (56%).

Figure 2: Frequency of occurrence and orientation of genes upstream and downstream from *aer*

homologs of select *Pseudomonas* species. Genes with the same orientation and high frequency of

occurrence with *aer* are outlined. Genes with noteworthy functions are coloured, and are

discussed in the text, such as the 3 different diguanylate cyclases/phosphodiesterases (A, dark

red; B, red; C, light red). Numbers beside gene functions indicate the frequency that they were

found within each group or sub group. Gene lengths are approximate, *aer* is about 1.5kb long.

Aer.g1* indicates Aer.g1 sequences not including Aer.g1A and Aer.g1B.

Evidence for Horizontal Transfer of *aer* Homologs

While examining the genomic contexts of *aer* homologs, indicators of mobile genetic elements were found nearby in many cases (Supplementary Table 2). Along with the varied distribution of Aer homologs within the *Pseudomonas* genus, and the findings from the genus phylogeny that distantly related species had similar complements of *aer* genes, this implied that they may have been subject to horizontal transfer. Initial inspection of the Aer phylogeny identified 3 sequences that diverged from the expected species phylogeny. These 3 sequences (WP_015271024.1, *P. putida* HB3267; WP_014754514.1, *P. putida* ND6; and WP_013791017.1, *P. fulva* 12-X) were each found clustered with sequences from unrelated species whereas other Aer sequences were consistently clustered with sequences from closely related species. Of these 3 genes only the *aer.g3* gene from *P. putida* ND6 was associated with mobile elements. Mobile elements were found within 5 Kbp of *aer* homolog genes (Supplementary Table 2). When mobile elements were found, they were often located immediately up and/or downstream of the *aer* homolog and its associated gene(s) described in the above section. Tanglegrams were used to identify instances where the *gyrB*/*rpoB*/*rpoD* based species phylogeny did not match with the *aer* subgroup phylogenies (Supplementary Figures 7-10). Three instances for Aer.g2 and one for Aer.g3 were identified where the *aer* gene had likely been transferred horizontally. Probable instances of horizontal transfer were also identified from mapping the *aer* subgroups onto the genus phylogeny (Figure 1A). *P. resinovorans* and *P. pseudoalcaligenes* KF707 have Aer.g4 despite being distantly related to the other strains that possess this homolog. Also, both *P. protegens* strains have Aer.g3 despite being more distantly related to the *P. putida* subgroup strains that only have this homolog.

<u>Amino Acid Sequence Comparison of Groups</u>

As genomic analysis of the Aer subgroups confirmed the AA sequence-based groupings

from the SHMR analysis, further sequence analysis was pursued to uncover sequence features

that distinguished the various Aer groups. First, the domain architecture from all sequences were

compared (Supplementary Figure 11) by submitting all sequences to SMART [28]. All

sequences were similar to the expected domain architecture of Aer from *P. aeruginosa* PA01,

consisting of an N-terminal Per/Arnt/Sim (PAS) domain, transmembrane helices and then the

cytoplasmic kinase control (called MA for methyl-accepting in SMART) region which includes

the CheW/CheA interface. HAMP domains between the transmembrane region and MA domain

were identified almost exclusively in Aer.g2 sequences, likely due to the difficulty in identifying

this domain [29]. To further identify regions of the protein that are specific to the subgroups,

results from the SHMR analysis for each inter-group comparison were plotted against the entire

length of the Aer protein (Figure 3A) and compared to the overall conservation at each AA

position (Supplementary Figure 12). As expected, the CheW interface region of the MCP-signal

domain and the PAS ligand binding domain were the most conserved regions, and did not differ

between groups. Inspection of the SHMR scores for each intergroup comparison showed that the

regions determining the specificity of each group were the same for each Aer group (Figure 3A),

except for Aer.g2, which had a group-specific region immediately N-terminal to the PAS domain.

Sequences from this group also had variable length C-terminal extensions. The most notable

group-specific region was in between the transmembrane helices and the beginning of the kinase

control (MA) domain. As this is the location of the HAMP domain in *E. coli* Aer, this region was

examined more closely.

Figure 3: Multi-relief comparison of Aer groups showing group-specificity (A) and group-specific Weblogos of HAMP domains (B). Black points indicate positions with a multi-relief score above 0.6. This cutoff indicates positions that are conserved within groups but divergent between. Domains are PAS (Pern/Art/Sin) ligand binding, TMH (transmembrane helices), HAMP (Histidine kinases, adenylate cyclases, methyl-accepting chemotaxis proteins and phosphatases), and CheW/CheA Interface also called Kinase Control (subdivided into A, signaling; B, flexible bundle; C, methylation). The HAMP domain weblogos in B were defined as the 54 AA anchored around Asp33. Characteristic HAMP domain residues were obtained from the SMART database, helices are marked in grey as defined in [1], and for each subgroup in yellow as determined by PSIPRED [2, 3]. Amino acids are coloured for hydrophobicity.

<u>Group-Specific Domain Differences</u>

First, the CheW/CheA interface region was examined more closely as its subdomains

have been clearly defined [30]. In bacterial chemoreceptors this domain consists of pairs of

heptads, separated into three subdomains. Central to the domain is a glutamate residue which

denotes the 'zero' position from which paired heptads are counted outwards until the C-terminus

on one side and the end of the domain on the other. As there were 20 heptads to the C-terminus

in all sequences, except Aer.g2 which had variable length extensions, all Aer from *Pseudomonas*

fall in the 40H class [30]. The three sub-domains (from the center outwards), signaling, flexible

bundle, and methylation were marked and used to compare the conservation (Supplementary

Figure 13) and group-specificity (Supplementary Figure 14) of the various Aer groups.

Conservation showed that corresponding heptads of the more central methylation subdomain

were better conserved than the flexible bundle subdomain. This corresponded with indications of

group-specific features in the flexible bundle subdomain and in the outer heptads (furthest from

the signaling subdomain) of the methylation subdomain.

The PAS and HAMP domains were also compared between all groups. The PAS domains

from all Aer groups were not notably different (Supplementary Figure 14). Conversely, the

region where a HAMP domain was expected, but only identified by SMART in Aer.g2, differed

between each Aer group (Figure 3B). The predicted 54 AA HAMP domains could be anchored

by a conserved aspartate and glycine [31], though the 51st and 54th residue of this domain marked

the first AA of the kinase control domain of Aer.g2 and the other groups, respectively. Aer.g2

matched the SMART database HAMP domain residues the closest. Though all groups differed

substantially from one another in this region, secondary structure prediction [32, 33] identified

helices in each group which matched the expected HAMP domain architecture [31]. In Aer.g1,

Aer.g3 and Aer.g4, the final seven AAs of the domain were well conserved, forming a heptad

with the motif SZEARL(K/Q). Only the Z was present in Aer.g2. With regards to the subdivision

of Aer.g1, both the PAS and HAMP domains supported keeping all sequences as a single group

(Supplementary Figures 15 and 16). Aer.g2 also differed from the other three groups at its C-

terminus as the final heptad of the kinase control domain differed and its members feature an up

to 9 AA extension (Supplementary Figure 17).

<u>Phenotypic Results</u>

To investigate the functions of Aer homologs, CttP and McpB, and to see whether they

were linked, deletion mutants were constructed in *P. pseudoalcaligenes* KF707. Single and

combination deletions were constructed using 2-step homologous recombination, the suicide

vector pG19II and SacB sucrose counter-selection [34]. To test energy-taxis, soft agar swim

plates were used [35]. In this assay cells consume carbon source from a central inoculation point

then expand outwards via a combination of division, chemotaxis and energy-taxis. Non-

chemotactic strains cannot expand beyond the point of inoculation but non-energy tactic cells

will produce a ring of expanding cells that is smaller than the wild-type (Supplementary Figure

18). To ensure that energy-taxis was being compared between the different mutants, the results

were compared to a taxis-negative CheA::KmR insertional inactivation mutant [36], and the

chemotactic ability of all strains was also confirmed in classical chemotaxis swim plates

(Supplementary Figure 19). Comparison of these metabolism independent results to the

metabolism dependent results of the energy-taxis swim plates allowed for differences in energy-

taxis capabilities to be observed between strains. All 24 mutants generated in this study, along

with the wild-type and CheA control were tested in triplicate in media containing pyruvate

(Figure 4) or succinate (Supplementary Figure 20). Ring diameters were measured from the

furthest point reached from the inoculation centre (Supplementary Figure 21), normalized to the

wild-type, then tested for differences using Tukey's Honest Significant Differences test

(Supplementary Table 3).

Figure 4: Mean normalized energy-taxis diameters in 50mM pyruvate of strains of *P.*

pseudoalcaligenes KF707 with deletions of *mcpB, cttP* and *aer* homologs. Boxes filled in dark-

grey indicate significant differences from the wild-type based on Tukey's Honest Significant

Differences test with a confidence value of 0.95. Values were normalized to the wild-type within

each experiment at 24 and 48 h, allowing datapoints from both times to be combined (n ≥ 5).

Wild-type strains were normalized to the mean of technical replicates within each experiment.

The *cheA::KmR* mutant was not grown with antibiotic present.

The diameter of the chemotactic negative CheA mutant was about 10% of the wild-type, whereas energy-taxis negative mutants with diameters about 60% smaller were significantly different from the wild-type (Figure 4). All the singular deletions of Aer.g1, Aer.g2, Aer.g4, McpB and CttP had no effect on energy-taxis; only strains that had Aer.g1 deleted in combination with at least one other receptor showed changes in their energy-taxis phenotype. Conversely, deletion of any combination of these receptors, including the quadruple McpB/CttP/Aer.g2/Aer.g4 mutant did not adversely affect energy-taxis. The McpB/Aer.g2/Aer.g4 triple mutant had a diminished diameter, about 80% of the wild-type, but this was not significant. Results from plates containing succinate (Supplementary Figure 20) were similar to those obtained using pyruvate, but were more variable. The observed differences in swim diameters were not due to differences in growth rate or swimming speed as the changes in swim diameters between the 24 and 48 h measurements were similar between all strains (Supplementary Figure 22, Supplementary Table 4). Growth rates in liquid media also did not differ (Supplementary Figure 23). Additionally, all strains were tested for chemotaxis towards pyruvate, glucose (Supplementary Figure 19) and succinate (data not shown) using traditional swim plate assays. All strains except the CheA::KmR mutant were able to swim towards pyruvate, succinate and glucose. As Aer.g1 played a pivotal role in energy-taxis, the wild-type and quintuple McpB/CttP/Aer.g1/Aer.g3/Aer.g4 mutant were complemented with Aer.g1 on a plasmid which showed the expected increase and recovery, respectively, of energy-taxis in succinate plates (Supplementary Figure 24, Supplementary Table 5). Together these experiments showed that it was only the combined deletion of Aer.g1 with any of the other tested receptors that caused a reduction of energy-taxis.

Discussion

Protein sequence and genetics classify Aer homologs into 5 groups

Aer has previously been described in *P. putida* and *P. aeruginosa* [8, 37–39], though the two species differed in the number of apparent homologs indicating there is interspecies variation. Here we constructed a protein sequence-based phylogeny which separated Aer into 5 different groups. All groups varied in the same regions of the protein, except Aer.g2 which had a unique region at the C terminus, which does not resemble the C-terminal CheR interface pentapeptide of McpB [40]. The most conserved regions included the PAS ligand binding domain and CheW/CheA interface region which confer the necessary functions of FAD cofactor binding/signal reception [41] and signal transduction, respectively. The conserved regions in the signaling region were better conserved than the predicted PAS domain, though these domains are known to differ at the sequence level [41]. Interestingly, it was the cytoplasmic region between the transmembrane helices and kinase control domain where a HAMP domain was expected that differed most between groups (Figure 3). In Aer from *E. coli* the HAMP domain in this region [42] interacts with the PAS domain and is essential for the function of the protein as the HAMP domain transfers the signal through the protein [11]. The motif (5)RP[…]DEXG(36) is representative of 'canonical' HAMP domains where the signal is transduced through the membrane [31], but was only found in Aer.g2. As the signal in Aer can be passed directly from the cytoplasmic PAS domain to the HAMP domain as in *E. coli* [43], the HAMP domains of the other Aer groups were likely able to evolve without disrupting this function, as predicted [31]. This implies that Aer.g1, Aer.g3, Aer.g4 and Aer.g5 have diverged from their common ancestor with Aer.g2, which matches the phylogenetic results.

 <u>Aer.g1 is the most prevalent homolog</u>

Aer from *E. coli* has been well-studied, so was used to estimate the root of the

Pseudomonas Aer phylogeny. After adding its sequence to the *Pseudomonas* alignment and

removing any gaps created (*E. coli* Aer is of the 36H class), the subsequent phylogeny placed it

in between Aer.g1 and Aer.g2 (Supplementary Figure 2). To confirm that this was an appropriate

branch to root the tree, an additional phylogeny was generated adding closely related non-

Pseudomonas aer-like genes (Supplementary Figure 3). Again, most sequences placed between

Aer.g1 and Aer.g2 indicating that the Aer.g1/3/4/5 and Aer.g2 groups diverged from a common

ancestor.

Aer.g1 was the most prevalent as all the *Pseudomonas* species included in this study had

at least one homolog except *P. denitrificans* ATCC13867. As this species did not have any Aer

homologs, this means that Aer is not part of the *Pseudomonas* core genome, though it is likely

ancestral and was lost in *P. denitrificans*. Unlike Aer.g1, Aer.g2 was not present in all

Pseudomonas species, but it was present in species from most subgroups. This suggests that both

Aer.g1 and Aer.g2 were present in the last common ancestor of all *Pseudomonads*.

Almost all *aer.g1* genes were adjacent to an aconitase gene (Figure 3), indicating that

they may be co-regulated or co-transcribed. In *Helicobacter pylori,* TlpD, which controls tactic

behaviour in low-energy conditions, has been found to interact with aconitase [44], possibly

indicating some common connection between aconitase and energy-taxis receptors. A gene

encoding a predicted CAAX amino protease was found adjacent to one third of Aer.g1 homologs.

This type of membrane-bound protease aids in prenylating proteins to ease their membrane

localization [45]. The Aer.g1 proteins whose genes were located beside these predicted proteases

did not have the expected CAAX motif and there were no obvious features at the sequence level

that distinguished these sequences from other Aer homologs indicating that the genomically co-

localized CAAX protease may not be involved in their functioning.

<u>Other Aer groups may have alternate functions</u>

As the 5 previously characterized Aer receptors are Aer.g1 homologs with the same

energy-taxis function, it is possible that the other homolog groups may have a related, but

different and/or more specific function. The three Aer homologs of *P. putida* KT2440 were

found to be differentially expressed [10], supporting this hypothesis. In *E. coli* Aer is also a

thermosensor [46] indicating another possible role, or they could aid in tuning the energy-taxis

response similarly to how Tsr and Tar from *E. coli* enable taxis towards the ideal pH from both

higher and lower initial pHs [47]. The function of the additional Aer homologs could also be

related to biofilm formation as Aer.g2 and Aer.g3 were each associated with their own PAS-

domain containing diguanylate cyclase/phosphodiesterase. Though Aer is expected to only act as

a Che1 (F6) chemosensor [48], the recent revelation that not all chemosensors mediate

chemotaxis [49], and are instead involved in so-called 'alternative cellular functions' such as

cyclic di-GMP signaling, makes this frequent association of phosphodiesterase/diguanylate

cyclases   with   Aer.g2   and   Aer.g3   noteworthy.   Deletion   of   the   diguanylate

cyclase/phosphodiesterase associated with Aer.g3 in *P. putida* KT2440 caused a general defect

in motility [10] and was later shown to be a bi-functional cyclic di-GMP phosphodiesterase and

diguanylate cyclase [50]. These two homolog groups may thus be involved in energy-sensing

behaviour in biofilms as c-di-GMP is an important regulator in the transition from planktonic to

sessile growth modes. In *P. aeruginosa*, increased levels of cyclic di-GMP decreases the

frequency of flagellar motor switching through MapZ as it inhibits CheR1 from methylating

some MCPs [51]. As this shows a direct connection between chemotaxis and cyclic di-GMP,

understanding the function of the Aer homolog-associated cyclase/phosphodiesterases will be of

interest.

<u>Possibility of horizontal transfer of *aer* homologs</u>

Chemoreceptors have been posited to be subject to horizontal gene transfer [52], as has

been demonstrated in plasmids of *E. coli/Shigella* [53]. Here we presented multiple lines of

evidence that suggest genes from the *aer* group have been horizontally transferred in the

chromosome of *Pseudomonas* species. In the Aer phylogeny, three homologs from *P. putida*

HB3267, ND6 and *P. fulva* 12-X each clustered beside Aer sequences from unrelated species

indicating they may have been obtained from those species. Incongruencies between the Aer

phylogeny and genus phylogeny were also found using tanglegrams. These figures highlight

other instances where *aer* homolog sequences were more closely related than the species hosting

these genes, which could indicate non-vertical inheritance. Additionally, we noted several cases

where strains of the same species differed in which Aer homologs they possessed.

The observed variation in the distribution of Aer homologs is unsurprising as even within

the *P. aeruginosa* pan-genome from 7 strains there are 2,000 accessory genes compared to 5,000

core genes [54]. The fact that *aer.g1* almost qualifies as part of the genus core genome (*P.*

dentrificans being the only outlier) indicates that is a very useful gene in the varied environments

inhabited by *Pseudomonas*. The existence of the various homologs speaks to their utility as they

each may confer a more specific function. As many *Pseudomonas* species live in the soil and

rhizosphere, which promotes horizontal gene transfer [55], there would be ample opportunity for

these *aer* homolog genes to spread. Interestingly they appear to be restricted to *Pseudomonas* as

reverse BLAST searches consistently returned *Pseudomonas* sequences as the closest hits.

Genes influencing energy-taxis in *P. pseudoalcaligenes* KF707

Genes influencing energy-taxis in *P. pseudoalcaligenes* KF707

P. pseudoalcaligenes KF707 was the only species that had three Aer homologs as well as

McpB and CttP. We have previously examined its chemotactic ability towards biphenyl [56], and

how its multiple *cheA* genes affect biofilm formation [36] and swimming motility [18]. Aer

from *P. aeruginosa* PA01 [8], *P. putida* F1 [15] and *P. putida* KT2440 [10] have been previously

characterized but are all part of the Aer.g1 homolog group. In the present study, all three *aer*

homologs in *P. pseudoalcaligenes* KF707, representing *aer.g1*, *aer.g2* and *aer.g4*, as well as

mcpB and *cttP* were deleted individually and in combination. The energy-taxis phenotypes of

these mutants were compared based on their diameters in soft agar energy-taxis plates. As the

growth rate of all the strains in these plates, and in liquid culture did not differ, as well as their

chemotactic ability in swim plates, the observed differences were taken as indication that the

perturbed genes influenced energy-taxis in some way.

No single deletions affected the energy-tactic ability of *P. pseudoalcaligenes* KF707.

This differs from *P. putida* KT2440 which became energy-taxis negative when only its Aer.g1

homolog was inactivated [38]. Both species have Aer.g4, but KF707 has Aer.g2 and KT2440

Aer.g3. In KF707, co-deletion of either *aer.g4* or *aer.g2* in the Δ*aer.g1* background caused a

reduction of energy-taxis, but deletion of *aer.g2* and *aer.g4* together had no effect. The three Aer

homologs in KT2440 are differentially expressed and thus were not truly redundant [38], but in

KF707 its Aer homologs appear to be at least partially redundant. In *Ralstonia solanacearum*

both of its "Aer1" and "Aer2" receptors were necessary for aerotaxis and restored energy-taxis in

a deficient *E. coli* mutant [57]. As these receptors are similar to Aer from *Pseudomonas* and *E.*

coli, it was likely energy-taxis enabling the swimming of *R. solanacearum* towards oxygen.

Interestingly these two receptors appear not to be redundant as inactivation of either one caused a

defect. In *Vibrio cholerae*, the number of Aer homologs also differs between species. The El tor

biotype has three, compared to two in strain O395N1 [58]. The additional Aer in the El tor

biotype is most similar to Aer.g2, though all the sequences from *Vibrio* and *Ralstonia* appear to

be quite different (identity below 50% on <90% coverage) than *Pseudomonas* sequences making

it difficult to categorize these homologs according to the naming scheme developed here for

Pseudomonas. Only one homolog in *V. cholera* was found to affect energy-taxis, leaving the

exact function of the other homologs unknown, similar to *Pseudomonas.*

Aer and McpB were first characterized as aerotaxis receptors in *P. aeruginosa* PA01 [8].

McpB has been shown to bind oxygen [9], and its overexpression in *P. aeruginosa* and *E. coli*

abolished chemotaxis [59] but its function as an aerotaxis receptor remains inconclusive. Here

we showed that single deletion of *mcpB* did not affect energy-taxis, but co-deletion with *aer.g1*

resulted in a similar reduction in energy-taxis as co-deletion with either of the other *aer*

homologs. The McpB protein has a C-terminal extension that specifically allows CheR2, and

only CheR2, to methylate it [40], and is required for the complexation of other Che2 proteins

[13], making it appear that the entire Che2 pathway exists to transduce the signal sensed by

McpB. This was recently confirmed using sequence analysis of all *P. aeruginosa* PAO1

chemoreceptors [48]. Our results were thus surprising as no connection was expected between

McpB and the che1 system, which transduces the signal of Aer, though the functions of Aer and

McpB in aerotaxis were linked in their initial characterization [8]. As their phenotypic outputs

may overlap, *P. denitrificans* may be an ideal organism to study McpB, as it has no Aer

homologs.

The *che2* gene cluster is preceded by *cttP* so we investigated the possibility of its

involvement in energy-taxis as well. In *P. aeruginosa* PA01, CttP was characterized as a receptor

for positive chemotaxis towards tetra/tri-chloroethylene (PCE/TCE) [19]. As this organism does

not perform reductive dehalogenation, intentional taxis towards these compounds is puzzling. In

E. coli, the receptors Tar and Tsr can mediate attractant and repellent responses to phenol

through its direct interaction with the transmembrane helices or cytoplasmic HAMP signaling

domains [20]. We thus hypothesized that a similar interaction may have mediated the observed

PCE/TCE chemotaxis and that CttP actually has a different function. CttP is only found in

organisms with McpB, and the gene is located immediately upstream of the *che2* gene cluster.

Here we showed that co-deletion of *cttP* and a*er.g1* in *P. pseudoalcaligenes* KF707 reduced

energy-taxis comparable to co-deletion of *aer.g2* or *aer.g4*. CttP has an unusual domain

architecture compared to other MCPs as it has no predicted ligand-binding domain and a single

N-terminal predicted transmembrane domain [60]. It does have the expected CheW/CheA

interface domain, but also a long (>100 AA) C-terminal extension not seen in most MCPs. It

could be possible that CttP does not directly sense any signal, but interacts with MCP receptor

complexes and/or influences their interaction with the Che complex. Overexpression of *cttP*

abolished all chemotaxis in both *P. aeruginosa* and *E. coli* [59], possibly indicating that its

function is in modulating signaling from the che1 system.

Despite being the first gene in the *che2* gene cluster, *cttP* is involved only with *che1*

genes as the *che2* system is dedicated to processing the signal sensed by *mcpB* [48]. It is thus

puzzling that *cttP* was always found in species that have *mcpB*, and that deletion of *mcpB* in

combination with *aer.g1* diminished energy-taxis in *P. pseudoalcaligenes* KF707. This could be

due to cross-talk between the systems, which is supported by a recent study that showed that the

che2 system is an ancestral F7 chemosensory system that evolved and co-opted parts of the *che*

(F6) system to become the well characterized system present in *E. coli* [61]. Here it was

suggested that since the F7 system appears to function in stress response, but that CttP co-

evolved with it and interacts with the F6 system[13] it may bridge the input/output of the two

systems.

<u>Conclusions</u>

In *Pseudomonas,* Aer is not a single receptor for energy-taxis, but a family of receptors

which differ mostly in the HAMP domain. The number of Aer homologs varies widely between

species and they have likely been horizontally transferred. The most prevalent homolog in the

Pseudomonas genus, here named Aer.g1, mediates energy-taxis in *P. aeruginosa* and *P. putida*,

but in *P. pseudoalcaligenes* KF707 Aer.g2 and Aer.g4, can also influence energy-taxis. McpB

and CttP also appear to have some effect. These findings indicate that Aer and energy-taxis in

Pseudomonas are more variable and complicated than the single receptor found in the archetypal

species *P. aeruginosa.*

 Materials and Methods

Protein Sequences

All sequences were obtained using NCBI databases and tools [62]. BLAST searches on the

NCBI website [21] (pBLAST for draft and completed genomes, tBLASTn for whole-genome

shotgun genomes) were performed using *P. aeruginosa* PAO1 Aer (NP_0250252.1) as a query

sequence. From each species, all hits with >95% sequence coverage, no matter how low the

sequence identity, were selected for inclusion. BLAST did not return any results with coverage

values between 67% and 95%, indicating that all included sequences were likely to truly be Aer

sequences. Expect values were always below 1×10^{-100}. Four sequences were removed as they

were redundant entries resulting from incorrect start site annotations resulting in two proteins

with the same C-terminus but slightly different N-termini. Sequence accession numbers were

then obtained from the international nucleotide sequence database collaboration (INSDC) [63].

Amino Acid Alignment

Full length sequences were aligned using COBALT (on the NCBI website) [22] with default

settings. Gaps were manually removed by adjusting the alignment in Jalview 2.10 [64] then re-

aligning using MUSCLE with default settings [65]. Names were cleaned up using a custom

script in R to reduce the names to just the species, strain and accession number in a presentable

fashion [66]. Aer from *E. coli* (AGX35053.1), and seventeen sequences obtained by BLAST

query using PAO1 Aer (NP_250252.1) against *Chromohalobacter salexigens, Marinobacter*

hydrocarbonoclasticus VT8, Marinobacter sp. ELB17, Hahella chejuensis KCTC 2396,

Marinomonas sp. MWYL1, Marinomonas sp. MED121, Neptuniibacter caesariensis

[Oceanospirillum sp. MED92], Bermanella marisrubri [Oceanobacter sp. RED65], Reinekea

blandensis (sp. MED297) were added to the *Pseudomonas*-only alignment which was then re-

aligned and all gaps trimmed so phylogenies were built only based on conserved regions.

Maximum likelihood phylogenies were built using PHYML [23], as implemented by the South

of France Bioinformatics platform [24]. PHYML options: Amino-Acids, 100 bootstrap replicates,

JTT amino-acid substitution model, BEST tree topology search operation, tree topology, branch

length and rate parameters were optimized. Consense was then used to generate a consensus tree

from the bootstrap replicates using the majority rule (extended).

Sequence Harmony and Multi-Relief Analysis to Compare Groups

The multi-Harmony server was used to apply sequence harmony and multi-relief

(SHMR) to validate groupings made based on the ML tree grouping and alignment. Groups were

manually decided based on the tree topology and corresponding alignment, initially making for 7

groups. These seven groups were compared in pair-wise fashion using SHMR [26]. Unique

regions of each group of Aer homologs were identified by comparing the SHMR scores for each

pair-wise comparison with the overall conservation score of each AA and the domain

architecture of the Aer protein. Conservation scores were obtained from Jalview [64], and along

with the SHMR scores were smoothed and plotted using ggplot2 [67] in R. Smoothing was

performed by calculating the average of the 3 proceeding and following AAs for each position.

The domain architecture of Aer was obtained from the conserved domain database using Aer

from *P. aeruginosa* as a query (NP_250252.1) (CDD) [68]. WebLogos were generated using the

Weblogo generator tool [69].

Distribution of Groups

Based on the ML tree grouping, the number of times a strain appeared in each group was

counted. A hierarchically clustered heatmap using Bray-Curtis distance and average clustering

was made in R. Presence of Aer-2 (NP_248866.1) and CttP (WP_003106690.1) were determined

using BLAST searches specifically against the strains.

Detection of Evidence of Horizontal Gene Transfer

Graphical representations of the Aer homologs nucleotide sequences were manually

inspected on NCBI. The first two genes upstream and downstream of the *aer* homolog were

noted, along with any mobile elements (transposase, integrase and inverted repeats) within 5kb.

For each different upstream and downstream gene their frequency of occurrence was calculated

for each homolog group. The frequency of occurrence of each type of mobile element was also

calculated, as well as for the complete set of homologs.

Generation of DNA Sequence Phylogeny

Gene sequences were obtained by BLAST [21] using sequences of *gyrB, rpoB* and *rpoD*

from *P. aeruginosa* PAO1. Sequences were aligned separately by MAFFT [70] and positions

with gaps in any strain were removed. Sequences from all three genes were then concatenated

and a ML phylogeny was generated using DIVEIN [71]. DIVEIN parameters: substitution model,

GTR; equilibrium frequencies, optimized; proportion of invariable sites and gamma distribution

parameter, estimated; number of substitution rate categories, 4; tree searching, NNI+SPR; tree

optimized for topology and branch lengths; 100 bootstrap replicates. Tanglegrams were

generated in Dendroscope [72] by pruning branches from the genus DNA tree so that only strains

that matched the particular Aer homolog group were included.

Culture Growth

For molecular biology, cultures were routinely cultured in lysogeny broth (LB, 5 g/L

yeast extract, 10 g/L Tryptone, 10 g/L NaCl). For energy-taxis experiments *P. pseudoalcaligenes*

KF707 strains were grown overnight (16 h) in minimal salts media containing 10 mM pyruvate

or succinate. Minimal salts media contained (in g/L) K_2HPO_4, 3; NaH_2PO_4, 1.15; NH_4Cl, 1; KCl,

0.15; $MgSO_4$, 0.15; $CaCl_2$, 0.01; $FeSO_4$, 0.0025. The latter four were sterile filtered and added

after autoclaving.

<u>Generation of Deletion Constructs and Mutants</u>

Deletions were generated using two-step homologous recombination [34] with

pG19II[73]. Nucleotide sequences for Aer.g1, Aer.g2, Aer.g4, McpB and CttP were obtained

from the draft genome sequence of *P. pseudoalcaligenes* KF707 [74]. Genomic DNA was

isolated by the phenol/chloroform method [75]. Primers were designed using Primer BLAST

[76] and purchased from IDT (IDT, USA). Deletion constructs with BamHI or HinDIII

restriction sites and complementary overlaps were ligated into pG19II using T4 ligase

(Invitrogen, USA) and transformed into *E. coli* Top10F' using standard methods [75].

Sequencing (Eurofins, USA) confirmed colonies were puddle mated with *P. pseudoalcaligenes*

KF707 and *E. coli* HB101 carrying the helper plasmid pRK2013 [77]. Transconjugants were

selected on AB glucose Gm20 plates. Second recombination events were selected for by pre-

growth on LB no salt then selection on LB 10% sucrose. Colony PCR and sequencing (Eurofins,

USA) were used to confirm the deletion. *P. pseudoalcaligenes* KF707 *cheA::Tn5* from Tremaroli

et al 2011 [36] is called the 'cheA' mutant throughout the text as its cheA1 (equivalent to *P.*

aeruginosa PA01) has been disrupted.

<u>Energy-taxis Swim Plates</u>

Swim plates were made by making minimal salts media with 0.3% agar and 50mM

succinate or pyruvate. Strains were grown overnight in MSM containing 10mM of the

appropriate carbon source. Plates were inoculated by dipping a sterilized needle into the culture

then stabbing into the plate. The diameter of growth for each strain was measured at the

maximum distance away from the inoculation centre that bacteria were visible, at 24 and 48 h

either manually using a ruler or digitally using a photograph and ImageJ [78]. Experiments were

repeated at least 3 times for all strains in each media, always including at least one wild-type to

be used as a normalizing control. Collected data were processed in R to normalize the size of the

growth diameter to the corresponding wild-type size at 24 and 48 h. Tukey's Honest Significant

Differences test was used to determine if the differences between strains were significant for

each carbon source. As the TukeyHSD() function in R [66] tests ALL pair-wise comparisons,

only those comparing each strain to the wild-type are presented here. However, this makes these

results more robust as the false positive correction for a confidence level of 0.95 was applied to

all 325 comparisons which were tested.

Chemotaxis Swim Plates

Strains were grown up overnight as before, then 1 mL was pelleted, washed once with 1

mL minimal salts media (no carbon source) then resuspended in 100 µL minimal salts media (no

carbon source). 20 µL was spotted at the edge of a minimal salts media plate containing no

carbon source and 0.3% agar. Either an agar plug containing 50 mM carbon source or small

amount of crystals was placed in the centre and plates were incubated overnight at 30°C. Plates

were photographed and positive chemotaxis was interpreted as an arc of cells nearer to the centre

of the plate than the cells had been spotted.

Growth Curves

Overnight cultures of all strains were normalized to OD 0.1 then diluted 1/100 into 200

µL minimal salts media with 10mM pyruvate in a 96 well microtiter plate. The plate was

incubated at 30°C, shaking at 150RPM and the OD600 was checked every 6h.

Complementation

Aer.g1 was complemented into the wild-type and the quintuple mutant (Aer-

2/CttP/Aer.g1/Aer.g2/Aer.g4). Aer.g1 was cloned into pSEVA_342 [79], and was conjugated

into *P. pseudoalcaligenes* KF707 the same as for the deletion constructs, only selection was

performed on AB glucose plates with 30µg/mL chloramphenicol. For the energy-taxis assays,

overnight cultures were grown with 30µg/mL chloramphenicol. Instead of directly comparing to

the wild-type, these strains were compared based on their change in energy-taxis diameters over

24h. All plasmids used in this study are summarized in Supplementary Table 6.

Acknowledgements

The authors would like to thank Stephana Cherak for her technical assistance, and Dr. Iain

George and Dr. Gordon Chua for use of their imaging setup for photographing the swim plates.

We thank Dr. Martina Cappelletti for assistance with the colony PCR design and her insights into

KF707 physiology and genomic interpretation. This project was funded by Natural Sciences and

Engineering Research Council (NSERC) of Canada Discovery Grant RGPIN/219895-2010 to

RJT.

References

1.   **Wadhams GH, Armitage JP**. Making sense of it all: Bacterial chemotaxis. *Nat Rev*

Microbiol 2004;5:1024–1037.

2.   **Sampedro I, Parales RE, Krell T, Hill JE**. Pseudomonas chemotaxis. *FEMS Microbiol*

Rev 2015;39:17–46.

3.   **Krell T**. Tackling the bottleneck in bacterial signal transduction research: high-throughput

identification of signal molecules. *Mol Microbiol* 2015;96:685–8.

4.   **Alexandre G, Greer-Phillips S, Zhulin IB**. Ecological role of energy taxis in

microorganisms. *FEMS Microbiol Rev* 2004;28:113–26.

5.   **Bibikov SI, Biran R, Rudd KE, Parkinson JS**. A signal transducer for aerotaxis in

Escherichia coli. *J Bacteriol* 1997;179:4075–4079.

6.   **Rebbapragada A, Johnson MS, Harding GP, Zuccarelli AJ, Fletcher HM,** *et al.* The

Aer protein and the serine chemoreceptor Tsr independently sense intracellular energy

levels and transduce oxygen, redox, and energy signals for Escherichia coli behavior. *Proc*

Natl Acad Sci U S A 1997;94:10541–6.

7.   **Samanta D, Widom J, Borbat PP, Freed JH, Crane BR**. Bacterial Energy Sensor Aer

Modulates the Activity of the Chemotaxis Kinase CheA Based on the Redox State of the

Flavin Cofactor. *J Biol Chem* 2016;291:25809–25814.

8.   **Hong CS, Shitashiro M, Kuroda A, Ikeda T, Takiguchi N,** *et al.* Chemotaxis proteins

and transducers for aerotaxis in Pseudomonas aeruginosa. *FEMS Microbiol Lett*

2004;231:247–252.

9.   **Watts KJ, Taylor BL, Johnson MS**. PAS/poly-HAMP signalling in Aer-2, a soluble

haem-based sensor. *Mol Microbiol* 2011;79:686–99.

10.   **Sarand I, Österberg S, Holmqvist S, Holmfeldt P, Skärfstad E, *et al.*** Metabolism-

dependent taxis towards (methyl)phenols is coupled through the most abundant of three

polar localized Aer-like proteins of Pseudomonas putida. *Environ Microbiol*

2008;10:1320–1334.

11.   **Garcia D, Watts KJ, Johnson MS, Taylor BL**. Delineating PAS-HAMP interaction

surfaces and signalling-associated changes in the aerotaxis receptor Aer. *Mol Microbiol*

2016;100:156–172.

12.   **Garcia D, Orillard E, Johnson MS, Watts KJ**. Gas Sensing and Signaling in the PAS-

Heme Domain of the Pseudomonas aeruginosa Aer2 Receptor. *J Bacteriol*

2017;199:e00003-17.

13.   **Güvener ZT, Tifrea DF, Harwood CS**. Two different Pseudomonas aeruginosa

chemosensory signal transduction complexes localize to cell poles and form and remould

in stationary phase. *Mol Microbiol* 2006;61:106–18.

14.   **Nichols NN, Harwood CS**. An aerotaxis transducer gene from Pseudomonas putida.

FEMS Microbiol Lett 2000;182:177–183.

15.   **Luu R a., Schneider BJ, Ho CC, Nesteryuk V, Ngwesse SE, *et al.*** Taxis of

Pseudomonas putida F1 toward phenylacetic acid is mediated by the energy taxis receptor

AER2. *Appl Environ Microbiol* 2013;79:2416–2423.

16.   **Bodilis J, Nsigue Meilo S, Cornelis P, De Vos P, Barray S**. A long-branch attraction

artifact reveals an adaptive radiation in pseudomonas. *Mol Biol Evol* 2011;28:2723–6.

17.   **Gomila M, Peña A, Mulet M, Lalucat J, García-Valdés E**. Phylogenomics and

systematics in Pseudomonas. *Front Microbiol* 2015;6:214.

18.   **Fedi S, Triscari-Barberi T, Nappi MR, Sandri F, Booth SC,** *et al.* The Role of cheA

Genes in Swarming and Swimming Motility ofPseudomonas pseudoalcaligenes KF707.

Microbes Environ 2016;31:169–172.

19.   **Shitashiro M, Tanaka H, Hong CS, Kuroda A, Takiguchi N,** *et al.* Identification of

chemosensory proteins for trichloroethylene in Pseudomonas aeruginosa. *J Biosci Bioeng*

2005;99:396–402.

20.   **Pham HT, Parkinson JS**. Phenol sensing by Escherichia coli chemoreceptors: a

nonclassical mechanism. *J Bacteriol* 2011;193:6597–604.

21.   **Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ**. Basic local alignment search

tool. *J Mol Biol* 1990;215:403–10.

22.   **Papadopoulos JS, Agarwala R**. COBALT: Constraint-based alignment tool for multiple

protein sequences. *Bioinformatics* 2007;23:1073–1079.

23.   **Guindon S, Gascuel O**. A Simple, Fast, and Accurate Algorithm to Estimate Large

Phylogenies by Maximum Likelihood. *Syst Biol* 2003;52:696–704.

24.   **Guindon S, Dufayard J, Lefort V**. New Algorithms and Methods to Estimate Maximum-

Likelihood Phylogenies : Assessing the Performance of PhyML 3 . 0. *Syst Biol*

2010;59:307–21.

25. **Williams KP, Gillespie JJ, Sobral BWS, Nordberg EK, Snyder EE,** *et al.* Phylogeny

of gammaproteobacteria. *J Bacteriol* 2010;192:2305–14.

26. **Brandt BW, Feenstra KA, Heringa J**. Multi-Harmony: detecting functional specificity

from sequence alignment. *Nucleic Acids Res* 2010;38:W35-40.

27. **Özen AI, Ussery DW**. Defining the Pseudomonas genus: where do we draw the line with

Azotobacter? *Microb Ecol* 2012;63:239–48.

28. **Schultz J, Copley RR, Doerks T, Ponting CP, Bork P**. SMART: a web-based tool for

the study of genetically mobile domains. *Nucleic Acids Res* 2000;28:231–4.

29. **Airola M V., Watts KJ, Crane BR**. Identifying divergent HAMP domains and poly-

HAMP chains. *Journal of Biological Chemistry*. Epub ahead of print 2010. DOI:

10.1074/jbc.L109.075721.

30. **Alexander RP, Zhulin IB**. Evolutionary genomics reveals conserved structural

determinants of signaling and adaptation in microbial chemoreceptors. *Proc Natl Acad Sci*

U S A 2007;104:2885–90.

31. **Dunin-Horkawicz S, Lupas AN**. Comprehensive analysis of HAMP domains:

Implications for transmembrane signal transduction. *J Mol Biol* 2010;397:1156–1174.

32. **Zimmermann L, Stephens A, Nam SZ, Rau D, Kübler J,** *et al.* A Completely

Reimplemented MPI Bioinformatics Toolkit with a New HHpred Server at its Core. *J Mol*

Biol. Epub ahead of print 2018. DOI: 10.1016/j.jmb.2017.12.007.

33. **Jones DT**. Protein secondary structure prediction based on position-specific scoring

matrices. *J Mol Biol*. Epub ahead of print 1999. DOI: 10.1006/jmbi.1999.3091.

34.   **Hmelo LR, Borlee BR, Almblad H, Love ME, Randall TE,** *et al.* Precision-engineering

711   the Pseudomonas aeruginosa genome with two-step allelic exchange. *Nat Protoc*

712   2015;10:1820–1841.

713   35.   **Harwood CS, Nichols NN, Kim M, Diitty JL, Parales RE**. Identification of the pcaRKF

714   Gene Cluster from Pseudomonas. *Biotechniques* 1994;176:6479–6488.

715   36.   **Tremaroli V, Fedi S, Tamburini S, Viti C, Tatti E,** *et al.* A histidine-kinase cheA gene

716   of Pseudomonas pseudoalcaligens KF707 not only has a key role in chemotaxis but also

717   affects biofilm formation and cell metabolism. *Biofouling* 2011;27:33–46.

718   37.   **Nichols NN, Harwood CS**. An aerotaxis transducer gene from Pseudomonas putida.

719   *FEMS Microbiol Lett* 2000;182:177–183.

720   38.   **Sarand I, Österberg S, Holmqvist S, Holmfeldt P, Skärfstad E,** *et al.* Metabolism-

721   dependent taxis towards (methyl)phenols is coupled through the most abundant of three

722   polar localized Aer-like proteins of Pseudomonas putida. *Environ Microbiol*

723   2008;10:1320–1334.

724   39.   **Luu R a., Schneider BJ, Ho CC, Nesteryuk V, Ngwesse SE,** *et al.* Taxis of

725   Pseudomonas putida F1 toward phenylacetic acid is mediated by the energy taxis receptor

726   AER2. *Appl Environ Microbiol* 2013;79:2416–2423.

727   40.   **García-Fontana C, Corral Lugo A, Krell T**. Specificity of the CheR2 methyltransferase

728   in Pseudomonas aeruginosa is directed by a C-terminal pentapeptide in the McpB

729   chemoreceptor. *Sci Signal* 2014;7:ra34.

41.   **Henry JT, Crosson S**. Ligand-Binding PAS Domains in a Genomic, Cellular, and

Structural Context. *Annu Rev Microbiol* 2011;65:261–286.

42. **Taylor BL**. Aer on the inside looking out: paradigm for a PAS?HAMP role in sensing

oxygen, redox and energy. *Mol Microbiol* 2007;65:1415–1424.

43. **Watts KJ, Ma Q, Johnson MS, Taylor BL**. Interactions between the PAS and HAMP

Domains of the Escherichia coli Aerotaxis Receptor Aer. *J Bacteriol* 2004;186:7440–7449.

44. **Behrens W, Schweinitzer T, McMurry JL, Loewen PC, Buettner FFR,** *et al.*

Localisation and protein-protein interactions of the Helicobacter pylori taxis sensor TlpD

and their connection to metabolic functions. *Sci Rep* 2016;6:23582.

45. **Pei J, Mitchell DA, Dixon JE, Grishin N V.** *Expansion of Type II CAAX Proteases*

Reveals Evolutionary Origin of γ-Secretase Subunit APH-1. 2011. Epub ahead of print

2011. DOI: 10.1016/j.jmb.2011.04.066.

46. **Nishiyama S, Ohno S, Ohta N, Inoue Y, Fukuoka H,** *et al.* Thermosensing function of

the Escherichia coli redox sensor Aer. *J Bacteriol* 2010;192:1740–3.

47. **Yang Y, Sourjik V**. Opposite responses by different chemoreceptors set a tunable

preference point in *Escherichia coli* pH taxis. *Mol Microbiol* 2012;86:1482–1489.

48. **Ortega DR, Fleetwood AD, Krell T, Harwood CS, Jensen GJ,** *et al.* Assigning

chemoreceptors to chemosensory pathways in Pseudomonas aeruginosa. *Proc Natl Acad*

Sci 2017;201708842.

49. **Bardy SL, Briegel A, Rainville S, Krell T**. Recent advances and future prospects in

bacterial and archaeal locomotion and signal transduction. *J Bacteriol* 2017;199:e00203-

17.

50.   **Österberg S, Åberg A, Herrera Seitz MK, Wolf-Watz M, Shingler V**. Genetic

dissection of a motility-associated c-di-GMP signalling protein of Pseudomonas putida.

Environ Microbiol Rep 2013;5:556–65.

51.   **Xu L, Xin L, Zeng Y, Yam JKH, Ding Y, *et al.*** A cyclic di-GMP–binding adaptor

protein interacts with a chemotaxis methyltransferase to control flagellar motor switching.

Sci Signal;9. http://stke.sciencemag.org.ezlibproxy1.ntu.edu.sg/content/9/450/ra102.full

(2016, accessed 10 October 2017).

52.   **Zhulin IB**. The superfamily of chemotaxis transducers: From physiology to genomics and

back. *Adv Microb Physiol* 2001;45:157–198.

53.   **Borziak K, Fleetwood AD, Zhulin IB**. Chemoreceptor gene loss and acquisition via

horizontal gene transfer in Escherichia coli. *J Bacteriol* 2013;195:3596–602.

54.   **Qiu X, Kulasekara BR, Lory S**. Role of Horizontal Gene Transfer in the Evolution of

Pseudomonas aeruginosa Virulence. *Genome Dyn* 2009;6:126–39.

55.   **Sengeløv G, Kristensen KJ, Sørensen AH, Kroer N, Sørensen SJ**. Effect of Genomic

Location on Horizontal Transfer of a Recombinant Gene Cassette Between Pseudomonas

Strains in the Rhizosphere and Spermosphere of Barley Seedlings. *Curr Microbiol*

2001;42:160–167.

56.   **Tremaroli V, Suzzi CV, Fedi S, Ceri H, Zannoni D, *et al.*** Tolerance of Pseudomonas

pseudoalcaligenes KF707 to metals, polychlorobiphenyls and chlorobenzoates: Effects on

chemotaxis-, biofilm- and planktonic-grown cells. *FEMS Microbiol Ecol* 2010;74:291–

301.

57.   **Yao J, Allen C**. The plant pathogen Ralstonia solanacearum needs aerotaxis for normal

774   biofilm formation and interactions with its tomato host. *J Bacteriol* 2007;189:6415–6424.

775   58.   **Boin MA, Häse CC**. Characterization of Vibrio cholerae aerotaxis. *FEMS Microbiol Lett*

776   2007;276:193–201.

777   59.   **Ferrandez A, Hawkins AC, Summerfield DT, Harwood CS**. Cluster II che Genes from

778   Pseudomonas aeruginosa Are Required for an Optimal Chemotactic Response. *J Bacteriol*

779   2002;184:4374–4383.

780   60.   **Kim HE, Shitashiro M, Kuroda A, Takiguchi N, Ohtake H,** *et al.* Identification and

781   characterization of the chemotactic transducer in Pseudomonas aeruginosa PAO1 for

782   positive chemotaxis to trichloroethylene. *J Bacteriol* 2006;188:6700–6702.

783   61.   **Ortega DR, Subramanian P, Mann P, Kjær A, Chen S,** *et al.* Repurposing a

784   macromolecular machine: Architecture and evolution of the F7 chemosensory system.

785   *bioRxiv* 2019;653600.

786   62.   **Tatusova T, Ciufo S, Fedorov B, O'Neill K, Tolstoy I**. RefSeq microbial genomes

787   database: new representation and annotation strategy. *Nucleic Acids Res* 2014;42:D553-9.

788   63.   **Cochrane G, Karsch-Mizrachi I, Takagi T, International Nucleotide Sequence**

789   **Database Collaboration IN**. The International Nucleotide Sequence Database

790   Collaboration. *Nucleic Acids Res* 2016;44:D48-50.

791   64.   **Waterhouse AM, Procter JB, Martin DMA, Clamp M, Barton GJ**. Jalview Version 2-

792   -a multiple sequence alignment editor and analysis workbench. *Bioinformatics*

793   2009;25:1189–91.

65. **Edgar RC**. MUSCLE: multiple sequence alignment with high accuracy and high

throughput. *Nucleic Acids Res* 2004;32:1792–1797.

66. **R Core Team**. R: A language and environment for statistical computing. *R Found Stat*

Comput Vienna, Austria 2014;2014.

67. **Wickham H**. ggplot2. *Wiley Interdiscip Rev Comput Stat* 2011;3:180–185.

68. **Marchler-Bauer A, Derbyshire MK, Gonzales NR, Lu S, Chitsaz F, *et al.*** CDD:

NCBI's conserved domain database. *Nucleic Acids Res* 2015;43:D222-6.

69. **Crooks GE, Hon G, Chandonia JM, Brenner SE**. WebLogo: A sequence logo generator.

Genome Res 2004;14:1188–1190.

70. **Katoh K, Standley DM**. MAFFT Multiple Sequence Alignment Software Version 7:

Improvements in Performance and Usability. *Mol Biol Evol* 2013;30:772–780.

71. **Deng W, Maust BS, Nickle DC, Learn GH, Liu Y, *et al.*** DIVEIN: A web server to

analyze phylogenies, sequence divergence, diversity, and informative sites. *Biotechniques*

2010;48:405–408.

72. **Huson DH, Scornavacca C**. Dendroscope 3: An Interactive Tool for Rooted

Phylogenetic Trees and Networks. *Syst Biol* 2012;61:1061–1067.

73. **Maseda H, Sawada I, Saito K, Uchiyama H, Nakae T, *et al.*** Enhancement of the

mexAB-oprM efflux pump expression by a quorum-sensing autoinducer and its

cancellation by a regulator, MexT, of the mexEF-oprN efflux pump operon in

Pseudomonas aeruginosa. *Antimicrob Agents Chemother* 2004;48:1320–8.

74.   **Triscari-Barberi T, Simone D, Calabrese FM, Attimonelli M, Hahn KR,** *et al.*

Genome sequence of the polychlorinated-biphenyl degrader pseudomonas

pseudoalcaligenes KF707. *J Bacteriol* 2012;194:4426–4427.

75.   **Sambrook J, Fritsch EF, Maniatis T**. No Title. *Mol Biol A Lab Man*.

76.   **Ye J, Coulouris G, Zaretskaya I, Cutcutache I, Rozen S,** *et al.* Primer-BLAST: a tool

to design target-specific primers for polymerase chain reaction. *BMC Bioinformatics*

2012;13:134.

77.   **Figurski DH, Helinski DR**. Replication of an origin-containing derivative of plasmid

RK2 dependent on a plasmid function provided in trans. *Proc Natl Acad Sci U S A*

1979;76:1648–52.

78.   **Schneider CA, Rasband WS, Eliceiri KW**. NIH Image to ImageJ: 25 years of image

analysis. *Nat Methods* 2012;9:671–675.

79.   **Silva-Rocha R, Martínez-García E, Calles B, Chavarría M, Arce-Rodríguez A,** *et al.*

The Standard European Vector Architecture (SEVA): A coherent platform for the analysis

and deployment of complex prokaryotic phenotypes. *Nucleic Acids Res* 2013;41:D666–

D675.