Title: Reactive Oxygen Species Regulate Activity-Dependent Neuronal Structural Plasticity in *Drosophila*

Authors: Matthew C. W. Oswald[1][*], Paul S. Brooks[1], Maarten F. Zwart[2], Amrita Mukherjee[1],

Ryan J. H. West[3], Khomgrit Morarach[1], Sean T. Sweeney[3], Matthias Landgraf[1][*]

*Correspondence to: mo364@cam.ac.uk and ml10006@cam.ac.uk

Affiliations:

[1] University of Cambridge, Department of Zoology, Downing Street, Cambridge, CB2 3EJ, United Kingdom

[2] HHMI Janelia Research Campus, Ashburn, VA, 20147, USA

[3] Department of Biology, University of York, Heslington York YO10 5DD, United Kingdom

Abstract

Neurons are inherently plastic, adjusting their structure, connectivity and excitability in response to changes in activity. How neurons sense changes in their activity level and then transduce these to structural changes remains to be fully elucidated. Working with the *Drosophila* larval locomotor network, we show that neurons use reactive oxygen species (ROS), metabolic byproducts, to monitor their activity. ROS signals are both necessary and sufficient for activity-dependent structural adjustments of both pre- and postsynaptic terminals and for network output, as measured by larval crawling behavior. We find the highly conserved Parkinson's disease-linked protein DJ-1ß acts as a redox sensor in neurons where it regulates pre- and postsynaptic structural plasticity, in part via modulation of the PTEN-PI3Kinase pathway. Neuronal ROS thus play an important physiological role as second messengers required for neuronal and network tuning, whose dysregulation in the ageing brain and under neurodegenerative conditions may contribute to synaptic dysfunction.

Introduction

Plasticity is fundamental to neuronal development and function. Changes in activity can trigger a raft of different responses, including changes to synaptic size, strength and connectivity, as well as to neuronal excitability (Davis and Müller, 2015; Keck et al., 2017; Wefelmeyer et al., 2015). 'Hebbian' plasticity mechanisms, such as long-term potentiation or depression, lead to altered transmission strength. These are balanced by homeostatic mechanisms, which are compensatory in nature and work toward maintaining neuronal or network activity within a set range. For example, blockade of excitatory inputs in neurons can induce a range of changes that are compensatory, including enlargement of both pre- and post-synaptic specializations, increasing synapse, axonal bouton and dendritic spine number as well as reducing dendritic spine elimination (Burrone et al., 2002; Kirov and Harris, 1999; Murthy et al., 2001; Zuo et al., 2005). How neurons monitor their activity levels is not known. Models of action potential firing rate homeostasis suggest that neurons measure the frequency of action potential firing and use homeostatic plasticity mechanisms to maintain this within their set point range (Davis, 2006; Turrigiano, 2012). In support of this view, recent studies in the mammalian visual system demonstrated that following dramatic activity disturbances, such as monocular occlusion, different types of neurons do indeed return to their original cell type-specific firing rate (Hengen et al., 2013; 2016).

An important second messenger in the regulation of activity-dependent plasticity is calcium, whose intracellular concentration correlates with neuronal activity patterns. Calcium regulates cellular and synaptic changes through a raft of signaling pathways, cytoskeletal and transcriptional effector proteins, including calcium/calmodulin-dependent protein kinase II (CaMKII), the phosphatase calcineurin and cyclic nucleotide signaling pathways (Vonhoff and Keshishian, 2017).

Here, we considered whether neurons might also monitor their activity levels by utilizing other signals. Neuronal activity levels correlate with energetic demand, suggesting that signals associated with energy metabolism could in principle provide a proxy measure for activity (Attwell and Laughlin, 2001; Hallermann et al., 2012; Zhu et al., 2012). We

therefore focused on reactive oxygen species (ROS), which are formed as obligate byproducts of mitochondrial respiratory ATP synthesis; ROS form by 'leakage' of the electron transport chain, which leads to the generation of superoxide anions (O_2^-), and hydrogen peroxide (H_2O_2) (Halliwell, 1992). Intensively studied as destructive agents when at high concentration in the context of ageing and neurodegenerative conditions, in healthy cells ROS are necessary for many physiological processes, including growth factor signaling (Finkel, 2011). In cultured neurons, mitochondrial O_2^- production can be triggered by field stimulation (Hongpaisan et al., 2004; 2003). Increase in neuronal ROS levels were also reported following NMDA receptor stimulation, reportedly generated by mitochondria (Bindokas et al., 1996; Dugan et al., 1995) and NADPH oxidases (Brennan et al., 2009). In hippocampal and spinal cord slices, ROS were found sufficient and necessary for inducing 'Hebbian' forms of plasticity (LTP) (Kamsler and Segal, 2003a; 2003b; Klann, 1998; Knapp and Klann, 2002; Lee et al., 2010); and mice that over-express superoxide dismutase showed defects in hippocampal LTP and learning paradigms (Gahtan et al., 1998; Levin et al., 1998; Thiels et al., 2000).

To explore whether neurons might use metabolic ROS as signals for activity-induced structural plasticity, we used as a model the motor system of the fruitfly larva, *Drosophila melanogaster*, which is composed of individually identifiable motoneurons located in the ventral nerve cord and their specific target muscles in the periphery (Kohsaka et al., 2012). We had previously shown that ROS can regulate neuromuscular junction (NMJ) structure (Milton et al., 2011). Here, we show that the locomotor network of the *Drosophila* larva undergoes homeostatic adjustment in response to prolonged overactivation, as measured by larval crawling, its physiological output. This homeostatic adjustment of the network includes and requires structural plasticity of synaptic terminals. We show that ROS signaling, H_2O_2 in particular, is necessary and sufficient for this activity-induced synaptic terminal plasticity, of both presynaptic NMJ terminals and postsynaptic dendritic arbors within the central nervous system (CNS). As a sensor for activity-generated ROS we identified the conserved redox sensitive protein DJ-1ß, a homologue of vertebrate DJ-1

(PARK7) (Meulener et al., 2005). DJ-1ß genetically interacts with the phosphatase and

tensin homolog (PTEN) and upon ROS mediated oxidation disinhibits PI3kinase signaling at

the presynaptic NMJ, thus promoting synaptic terminal growth, as seen with activity-

induced potentiation. Strong over-activation on the other hand leads to DJ-1ß-dependent

compensatory adjustments of reduced release sites presynaptically and smaller dendritic

input arbors.

Results

Induction of prolonged activity leads to homeostatic adjustment in the larval

***Drosophila* motor network**

Neural circuits are inherently plastic and capable of adjusting their output with behavioral need. Our aim was to investigate mechanisms that allow neurons to sense changes in their activity levels and in response mediate adaptive structural adjustments. As an experimental model we used the locomotor network of the *Drosophila* larva, whose motoneurons grow dramatically over the course of larval life and are known to be plastic (Budnik et al., 1990; Davis, 2006; Schuster et al., 1996; Tripodi et al., 2008; Zito et al., 1999; Zwart et al., 2013). A simple way of manipulating activity in the locomotor network of *Drosophila* larvae is by changing ambient temperature (Dillon et al., 2009; Sigrist et al., 2003; Zhong and Wu, 2004). As previously documented (Sigrist et al., 2003; Zhong and Wu, 2004), we found that network activity increased upon *acutely* shifting animals to higher ambient temperatures (e.g. from 25°C to 29°C or 32°C), resulting in increased larval crawling speeds (Figure 1A). In contrast, following *prolonged* exposure to higher temperatures (rearing at 29°C and 32°C) we found that larvae crawled at the same speed as controls kept at 25°C (grey horizontal dotted line in Figure 1A), suggestive of homeostatic network adjustment. Mechanistically, neurons might compensate for the increased network drive that is the result of higher temperatures by reducing their individual excitability and/or synaptic input, thus returning motor output to the default crawling speed. In agreement with this model, when 29°C and 32°C-conditioned larvae were *acutely* shifted to a lower temperature of 25°C they displayed reduced crawling speed (green data in Figure 1a), with 32°C-adjusted larvae crawling significantly slower than 29°C -adjusted animals.

To test this further, we selectively over-activated the glutamatergic motoneurons by mis-expression of the warmth-gated cation channel dTrpA1 and rearing these animals at 27°C, a temperature that robustly activates dTrpA1-expressing neurons (Hamada et al., 2008; Pulver et al., 2009). Upon *acute* removal of this motoneuron over-stimulation (by

shifting to 22°C, where the dTrpA1 channel is closed) larval crawling speed reduced

significantly relative to non-dTrpA1 expressing controls (Figure 1B). Thus, prolonged

increase of network drive leads to homeostatic adjustment that returns network output, and

thus larval crawling speed, to a default set point. Cell type-selective manipulations of the

motoneurons, which constitute the output of the motor network, show that they are major

contributors to this adjustment.

Figure 1: Adaptive behavioral plasticity is response to chromic locomotor over-activation. (**A**) Larval motor network activity, assayed by crawling speed 72hrs after larval hatching (ALH), increases in response to acute temperature upshifts (blue) in wild type larvae (Oregon R = control in all conditions). In contrast, with prolonged exposure (grey) to elevated temperatures (29°C or 32°C) the motor network adapts homeostatically generating the same crawling speed as 25°C reared controls. This adaptation is further revealed by acute temperature downshifts (green). Each data point represents crawling speed from an individual uninterrupted continuous forward crawl, n = specimen replicate number, up to 3 crawls assayed for each larva. Mean +/- SEM, ANOVA, ns = not significant, *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001 (**B**) Prolonged over-activation targeted to motoneurons (*dVGlut-GAL4*; *UAS-dTrpA1*) also leads to adaptation with reduced crawling speed (dTRPA1 channels open at 27°C, closed at 22°C). Mean +/- SEM, control is *dVGlut-GAL4 / +*. ****P<0.0001 students t test, n = replicate number.

Structural plasticity of synaptic terminals is regulated by activity

We then asked what structural changes within the network might be associated and potentially responsible for these homeostatic behavioral adjustments. To this end, we focused on two well characterized motoneurons, 'aCC' and 'RP2' (Choi et al., 2004; Landgraf et al., 2003). Their presynaptic neuromuscular junction (NMJ) terminals consist of varicose compartments (boutons) strung across the muscle surface, each containing multiple synapses in the form of active zones (Van Vactor and Sigrist, 2017) (Figure 2A). We found that increases in motor network activity stimulated by higher ambient temperatures (raising animals at 29°C or 32°C) led to more, albeit smaller boutons at the NMJ by the wandering third instar larval stage, 100 hrs after larval hatching (ALH) (Figure 2C, grey data) (Sigrist et al., 2003; Zhong and Wu, 2004). To exclude non-specific effects, we targeted over-activation selectively to the aCC and RP2 motoneurons, via expression of the warmth-gated ion channel dTrpA1, and found similar presynaptic structural adjustments (Figure 2B & C, blue data; see also Figure 1 – supplement 1). While bouton number positively correlates with increases in motoneuron activity, quantification of synapse number, measured by quantification of active zones, describes a more complex relationship. As previously published, a moderate increase in activity (rearing larvae at 29°C) causes more boutons and also more active zones to be formed, thus potentiating transmission at the NMJ (Sigrist et al., 2003). In contrast, further increases in network activity, as effected by rearing larvae at 32°C, or cell-specific dTrpA1-mediated motoneuron activation led to progressive active zone reductions, consistent with a homeostatic response (Figure 2D, E).

Figure 2: Adaptive structural synaptic plasticity at motorneuron input and output terminals in response to increased neuronal activity. (**A**) Graphical illustration of a stereotypical larval motorneuron (MN), adapted from (Kohsaka et al., 2012). Pre-motor interneurons make synaptic connections with the MN dendritic arbor (input) in the larval ventral nerve cord (equivalent of mammalian spinal cord). The MN extends an axonal projection into the periphery where it connects with a target muscle via an NMJ, characterized by varicose swellings (boutons) each containing multiple individual neurotransmitter release sites (active zones). (**B** and **C**) Representative images of muscle DA1[muscle 1] NMJs from 3rd instar larvae (100hrs ALH). Dot-plot quantification shows NMJ bouton number increases in response to systemic and cell-specific activity increases. (**D** and **E**) Active zone number increases following low-level over-activation (29°C), but progressively reduces upon stronger over-activation. (**F** and **G**) Digital reconstructions and dot plots show that over-activation leads to reduced total dendritic arbor length of aCC motoneurons (24hrs ALH). '*aCC/RP2-GAL4*' expresses GAL4 in all, '*aCC-FLP-GAL4*' in single aCC and RP2 motoneurons (see Online Methods for details); 'Control' in (F + G) is *aCC-FLP-GAL4* alone. Mean +/- SEM, ANOVA, ns = not significant, *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001, n = replicate number. Comparisons with control are directly above data points.

We then turned to look at the input arbor of the motoneuron, branched dendrites on

which the pre-motor interneurons form connections (Baines et al., 1999; Schneider-Mizell et

al., 2016; Zwart et al., 2013). Using a stochastic labeling strategy within a small subset of

motoneurons (aCC-FLP-GAL4) (Ou et al., 2008), we selectively targeted GAL4 and dTrpA1

expression to individual aCC motoneurons. Following confocal imaging and digital

reconstructions of the arbors (Evers et al., 2005; Schmitt et al., 2004), morphometric

analysis revealed that the size of the postsynaptic dendritic arbor decreased with rising levels

of temperature-gated dTrpA1 activity (Figure 2F, G). Because in these motoneurons

dendritic length correlates with input synapse number and synaptic drive (Tripodi et al.,

2008; Zwart et al., 2013), we interpret this structural adjustment of the dendrites as

homeostatic.

Thus, while a moderate increase in neuronal activity can lead to potentiation of the

presynaptic motoneuron terminal by increasing bouton and synapse number (Ataman et al.,

2008; Piccioli and Littleton, 2014; Sigrist et al., 2003), stronger over-activation leads to

structural adjustments at both input and output terminals of the motorneuron that we

interpret as compensatory; smaller dendritic arbors being associated with less synaptic drive

(Zwart et al., 2013) and fewer presynaptic release sites at the NMJ reducing transmission to

the muscle.

Activity generated ROS regulate structural plasticity at synaptic terminals

Because neural activity is energetically demanding (Attwell and Laughlin, 2001; Hallermann

et al., 2012; Zhu et al., 2012) we wondered whether neurons might monitor activity levels

through metabolic signals. Others previously reported that neuronal over-activation *in vitro*

can lead to a decrease in the ATP:ADP ratio (Tantama et al., 2013) and an increase in

mitochondrial ROS (Hongpaisan et al., 2004). Moreover, at the *Drosophila* NMJ oxidative

stress has been shown to regulate NMJ structure (Milton et al., 2011). To ask whether ROS

signaling might be associated with activity-dependent synaptic terminal growth, we

expressed the mitochondrion-targeted ratiometric ROS reporter *UAS-mito-roGFP2-Orp1* in

aCC and RP2 motoneurons (Gutscher et al., 2009). This reports a titratable neuronal

activity-dependent increase in mitochondrial ROS (Figure 3A).

We therefore hypothesized that mitochondrially derived ROS might provide a

readout of neuronal activity and regulate synaptic structural plasticity. To test this

hypothesis, we selectively increased neuronal activity in the aCC and RP2 motoneurons

while at the same time over-expressing the ROS scavenging enzymes Superoxide Dismutase

2 (SOD2, which catalyses O_2^- to H_2O_2 reduction) or Catalase (H_2O_2 into H_2O and O_2).

Catalase co-expression significantly counteracted dTrpA1 activity-induced bouton addition

at the NMJ (Figure 3B) and rescued motoneuron dendritic arbor size in the CNS (Figure 3C).

SOD2 co-expression on the other hand enhanced dTrpA1-mediated NMJ elaboration,

presumably by potentiating conversion of neuronal activity-induced O_2^- into H_2O_2 (Figure

3B). We then confirmed that ROS are also sufficient to invoke structural plasticity in the

absence of neuronal activity manipulation through cell-specific RNAi knock down of SOD1,

SOD2 and Catalase (Figure 3B). All of these manipulations led to NMJ growth phenotypes

with increased bouton number that are similar to those produced by neuronal over-

activation. Based on these results, we implicate ROS, specifically H_2O_2, in the observed

activity-dependent structural plasticity.

Figure 3: Neuronal activation leads to generation of synaptic ROS that regulate structural plasticity at input and output terminals. (**A**) Elevated neuronal activity increases mitochondrial ROS production at NMJs. Dot plots of mitochondrion-targeted ratiometric H_2O_2 sensor (*UAS-mito-roGFP2-Orp1* (Gutscher et al., 2009)) in wandering 3[rd] instar larval NMJs (100hrs ALH) at 23˚C (control, dTrpA1 inactive), 25˚C (moderate activity, ~10Hz), 27˚C (strong activity, ~20Hz) and 29˚C (very strong excitation, ~40Hz). (**B**) Bouton number at the NMJ is increased by *UAS-dTrpA1*-mediated over-activation. This is exacerbated by co-expression of *UAS-SOD2* (converts O_2^- to H_2O_2) and rescued by the H_2O_2 scavenger *UAS-Catalase*. Cell-specific ROS elevation by scavenger knockdown is sufficient to induce NMJ elaboration. *aCC/RP2-GAL4*, 'Control' is *aCC/RP2-GAL4* alone. (**C**) Total dendritic arbor length is reduced by single cell over-activation, but rescued by co-expression of the H_2O_2 scavenger *UAS-Catalase* (aCC motoneurons, 24hr ALH). *aCC-FLP-GAL4*, 'Control' is *aCC-FLP-GAL4* alone. Larvae reared at 25ºC. Mean +/- SEM, ANOVA, ns = not significant, *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001, n = replicate number.

DJ-1ß acts as a ROS sensor in neurons

We then asked how neurons might sense changes in activity-induced ROS levels. ROS are known to post-transcriptionally modify many different proteins, principally on cysteine residues, including transcription factors, cell adhesion molecules and phosphatases (for review see (Milton and Sweeney, 2012)). Following a literature review, we focused on DJ-1ß, the fly ortholog of DJ-1 (PARK7), as a candidate ROS sensor. *DJ-1* codes for a highly conserved, ubiquitously expressed redox-sensitive protein, that protects against oxidative stress and regulates mitochondrial function (Ariga et al., 2013; Nagakubo et al., 1997). A mutant allele is also linked to a rare form of familial Parkinsonism (Bonifati et al., 2003). *DJ-1* null mutant adult flies had been reported to be more sensitive to oxidative stress, induced by exposure to paraquat and H_2O_2 (Meulener et al., 2005). We found that *DJ-1β* null mutant (*DJ-1β^{Δ93}*) larvae develop normally and have structurally normal NMJs (Figure 4A and Figure 4-supplement 1). What is more, systemic loss of DJ-1ß (*DJ-1β^{Δ93}* null mutants) completely rescues NMJ bouton addition phenotypes normally caused by pharmacologically induced oxidative stress (Milton et al., 2011) (Figure 4A). Loss of DJ-1ß also significantly rescues NMJ bouton phenotypes induced by cell type-specific expression of the ROS generator Duox or dTrpA1-mediated over-activation (Figure 4A). *DJ-1β^{Δ93}* mutant larvae failed to produce the presynaptic bouton and active zone addition characteristic of NMJ potentiation as seen in controls following periods of temperature-stimulated (29°C) increases in locomotor activity (Sigrist et al., 2003) (Figure 4 – supplement 2). Motoneuron-targeted expression of a dominant-acting mutant form of DJ-1ß that is non-oxidizable at the conserved cysteine 104 (DJ-1ß^{C104A}) abrogated dTrpA1 activity-mediated NMJ structural adjustment, both with respect to bouton number (Figure 4A) and active zone number (Figure 4B). Looking at structural plasticity of the postsynaptic dendritic arbor, we found that halving the *DJ-1β* copy number (in *DJ-1β^{Δ93}/+* heterozygotes) was sufficient to significantly suppress dTrpA1-mediated reductions in dendritic arbor size (Figure 4C).

In summary, these data show that DJ-1ß is necessary for sensing activity-induced

ROS and for implementing structural changes at both pre- and postsynaptic terminals in

response. At the presynaptic NMJ, DJ-1ß appears to be required in motoneurons for at least

two kinds of activity regulated structural plasticity: following mild over-activation, the

addition of both boutons and active zones, which are associated with potentiation (Sigrist et

al., 2003); and following stronger over-activation, reduction of active zone number, which

we interpret as an adaptive adjustment.

Structural plasticity of synaptic terminals is required for homeostatic

adjustment of locomotor behavior

We then asked whether this ROS - DJ-1ß mediated structural plasticity of synaptic terminal

growth was required for the homeostatic adjustments of crawling speed, which ensues after

prolonged over-activation. To answer this question, we targeted expression of the dominant

acting non-oxidizable DJ-1ß[C104A] to motoneurons (and other glutamatergic cells, using

DvGlut-T2A-GAL4). We then tested the behavior of these animals for adjustment in

response to *chronic* temperature-induced elevation of motor network activity. Our previous

experiments showed that expression of non-oxidizable DJ-1ß[C104A] in motoneurons does not

lead to abnormal bouton number or the decrease in active zone number normally caused by

over-activation (see Figures 4A & 4B). Expression of non-oxidizable DJ-1ß[C104A] in

motoneurons per se did not alter larval crawling speed at the control temperature of 25^oC.

However, when rearing these larvae at 32^oC, which is associated with elevated motor

network activation, unlike controls they failed to homeostatically adjust toward the default

crawling speed set point (Figure 5). Consequently, such larvae reared at elevated

temperatures (29^oC or 32^oC) also responded less strongly than controls to acute temperature

downshifts (Figure 5). These data suggest that activity-induced neuronal plasticity,

implemented through the ROS sensor DJ-1ß, is necessary for activity-directed homeostatic

adjustment of larval locomotor behavior.

PTEN and PI3K are downstream effectors of the DJ-1ß ROS sensor

Next, we looked for downstream effector pathways responsible for implementing activity and ROS-dependent structural plasticity. DJ-1 is a known redox-regulated inhibitor of Phosphatase and Tensin homologue (PTEN) and thus a positive regulator of PI3Kinase signaling (R. H. Kim et al., 2005; Y.-C. Kim et al., 2009); and PI3Kinase is a known positive regulator of NMJ development (Jordán-Álvarez et al., 2012; Martín-Peña et al., 2006). To test whether DJ-1ß - PTEN interactions mediate ROS-dependent NMJ adjustments, we performed genetic interaction experiments in the context of a DEM (oxidative stress) dose-response curve (Figure 6A). Focusing on the NMJ, we found that in controls bouton number

increases linearly with exposure to increased DEM concentrations, peaking at 15mM DEM

(Figure 6A). Removing one copy of *DJ-1β* (*DJ-1β^Δ93/+*) was sufficient to suppress DEM-

induced increases in bouton number. In contrast, *PTEN^CO76/+* heterozygous larvae showed

increased sensitivity to DEM, with a left-shifted dose response curve. Larvae heterozygous

for both *DJ-1β* and *PTEN* (*PTEN^CO76/+; DJ-1β^Δ93/+*) were more sensitive to DEM at higher

concentrations than *DJ-1β^Δ93/+* heterozygotes. These genetic interactions agree with

previous studies (R. H. Kim et al., 2005) and complement biochemical data that showed

H_2O_2-oxidized DJ-1ß binds to and inhibits PTEN (Y.-C. Kim et al., 2009).

Figure 6: *DJ-1β* signals via PTEN and PI3Kinase to regulate ROS and activity-induced NMJ elaboration. (**A**) *DJ-1β* and PTEN genetically interact to regulate systemic ROS-induced NMJ elaboration. NMJ bouton number varies with ROS (DEM) levels (grey data). Removal of one copy of *PTEN* sensitizes (red) while heterozygosity for *DJ-1β* desensitizes NMJs to ROS levels (yellow), partially restored in double heterozygotes (blue). Dashed boundaries indicate 95 % confidence intervals (n≥38). (**B**) Systemic ROS and activity-induced NMJ structural adjustments require PTEN and PI3Kinase signaling. Over-expression of the PI3Kinase antagonist PTEN or a dominant negative PI3Kinase form abrogates activity-induced NMJ elaboration. *aCC/RP2-GAL4*, 'Control' is *aCC/RP2-GAL4* alone. Mean +/- SEM, ANOVA, ns = not significant, **P<0.01, ***P<0.001, ****P<0.0001, n = replicate number.

To further test specificity, we manipulated PTEN and PI3Kinase activities in single

cells. Targeted knock-down of PTEN in motoneurons sensitized these neurons to ROS,

exacerbating the DEM-induced bouton addition phenotype (Figure 6B). In contrast, over-

expression of PTEN or mis-expression of a dominant negative form of PI3Kinase

significantly reduced NMJ elaboration normally caused by DEM exposure or dTrpA1-

mediated neuronal activity increase (Figure 6B). Together these interactions suggest that

PTEN and PI3Kinase act downstream of DJ-1ß and neural activity-generated ROS; and that

oxidation of DJ-1ß, through PTEN inhibition, facilitates a rise in PI3Kinase / PIP$_3$ signaling,

which in turn mediates at least part of the structural synaptic terminal plasticity (Figure 7).

Figure 7: Model summary. *DJ-1β* is a redox signaling hub that coordinates structural synaptic plasticity at motoneuron synaptic input and output terminals. Activity-induced ROS oxidize *DJ-1β*, leading to PTEN inhibition and thus to a gain in PI3Kinase signaling, which regulates activity-induced NMJ elaboration of boutons and active zones. At higher activity/ROS thresholds additional, yet to be defined, pathways downstream of *DJ-1β* are activated, implementing adaptive reductions of active zones at the NMJ and dendritic arbor length in the CNS.

Discussion

Our aim was to study mechanisms of structural plasticity in relation to neuronal activity in a motor network. To do so we used as a model the *Drosophila* larva and focused on major effectors of motor circuit output: motoneurons, their dendrites in the CNS and presynaptic terminals at peripheral neuromuscular junctions. We have identified a novel pathway by which these neurons sense changes in activity through measuring increases in activity-generated ROS, which act as second messengers for structural adjustments at both dendritic input and presynaptic output terminals. Our data further suggest that this activity-regulated structural plasticity is instrumental for the homeostatic adjustment of larval locomotor network activity and thus crawling behavior.

Homeostatic adjustment of the motor network allows animals to maintain a dynamic range of crawling speed

To investigate forms and mechanisms of structural plasticity that neurons undertake while adapting to altered activity levels, we took advantage of *Drosophila* motor network activity being readily manipulated through changes in ambient temperature. We found that while acute increases in ambient temperature from 25°C to 29°C or above quickly push larvae to their maximum crawling speed (approx. 0.8mm/sec), prolonged exposure leads to the network adjusting homeostatically toward a default crawling speed (approx. 0.65mm/sec) (Figure 1A). This adaptation might overall be energetically more favorable, and it allows larvae to stay within their dynamic range of behavioral responses to relative changes regardless of absolute ambient temperature.

Cell type-specific over-activation of the glutamatergic motoneurons (via dTrpA1) largely recapitulated the larval crawling phenotypes otherwise generated by systemic motor network overactivation. Conversely, blockade of activity-induced structural adjustment (through mis-expression of non-oxidisable DJ-1ß^{C104A}) in the glutamatergic motoneurons significantly blocked homeostatic locomotor network adjustment to higher ambient

temperatures (Figure 1B). The importance of the motoneurons in shaping much of the adaptive behavioral response is perhaps to be expected, since their dendrites are the final integrators on which all pre-motor inputs converge (Fushiki et al., 2016; Itakura et al., 2015; Kohsaka et al., 2014; Schneider-Mizell et al., 2016; Zwart et al., 2016). Systemic manipulations produce stronger behavioral phenotypes than those targeted to the glutamatergic neurons, suggesting, as would be expected, that other cells within the locomotor network are also subject to activity-regulated homeostatic plasticity (Figure 5). Interestingly, a recent study, which modeled different forms of plasticity in this motor network, concluded that during its development homeostatic mechanisms are better placed in achieving the formation of a stable balanced network producing robust activity patterns than Hebbian alternatives (Gjorgjieva et al., 2016).

Structural adjustment coordination between pre- and postsynaptic terminals

Working with identified motoneurons we attempted to take an integrated view by relating structural plasticity across its pre-synaptic and dendritic terminals. At the presynaptic NMJ low-level increases of activity lead to potentiation, associated with increased numbers of active zones, elevated levels of the Brp/CAST/ELKS family active zone protein Bruchpilot (Brp) and the addition of boutons, varicose swellings containing multiple active zones (Cho et al., 2015; Sigrist et al., 2003; Vasin et al., 2014; Weyhersmüller et al., 2011).It is known that changes to the composition of the active zone cytomatrix (Davydova et al., 2014; Lazarevic et al., 2011; Matz et al., 2010) such as elevation of Brp levels can further aid transmission by increasing release probability (Peled et al., 2014; Weyhersmüller et al., 2011).

Strong over-activation, in contrast, though triggering further addition of boutons, leads to overall reductions in active zone number at the NMJ, which correlate with the severity of over-activation (Figure 2D, E). This parallels observations in the adult *Drosophila* visual system where increased activation also triggers a reduction of Brp levels in photoreceptor terminals (Sugie et al., 2015). We interpret this negative correlation between

the degree of over-activation and the reduction in active zone number, and potentially Brp

content, as a compensatory response. In agreement, blocking activity-induced structural

adjustment in the motoneurons abrogates behavioral adaptation to prolonged over-

activation (Figure 5). Whether increased numbers of smaller boutons forms part of an

adaptive response to over-activation or is anticipatory of future synapse strengthening

remains to be established.

The input terminal of the neuron, the dendritic arbor in the CNS, responds to over-

activation with corresponding reductions in dendritic length, which we previously found to

correlate with reducing both input synapses and synaptic drive. We therefore interpret this

activity driven structural adjustment of the dendrites also as compensatory (Tripodi et al.,

2008; Zwart et al., 2013). In summary, we found that above threshold over-activation

regimes lead to structural adaptive adjustments at both the postsynaptic input and

presynaptic output terminals. We do not yet know whether these are regulated locally or

globally coordinated by transcriptional changes; though at least for the presynaptic NMJ

activity and ROS mediated structural changes have been reported as dependent on

immediate early genes (AP-1) (Milton et al., 2011; Sanyal et al., 2002). Given the available

evidence we would expect these structural adjustments to be linked to, and potentially

preceded by compensatory changes in neuronal excitability (Baines et al., 2001; Davis, 2006;

Davis et al., 1996; 1998; Driscoll et al., 2013; Frank et al., 2006; 2009; Gaviño et al., 2015;

Giachello and Baines, 2016; Lin et al., 2012; Mee et al., 2004; Müller and Davis, 2012;

O'Leary et al., 2013; Prinz, 2006; Prinz et al., 2004; Wang et al., 2014; Younger et al., 2013).

ROS and redox sensitive DJ-1ß provide a readout of neuronal activity

Neurons have cell type-specific set points of average firing ranges to which they return

following activity perturbations using a variety of homeostatic plasticity mechanisms

(Hengen et al., 2016; 2013; Keck et al., 2017; 2013). The mechanisms that determine the

homeostatic set point and allow neurons to sense when their activity deviates from such a set

point are not yet known. Intracellular concentrations of calcium correlate with neuronal

activity (Hardingham et al., 2001), making this versatile second messenger a prime

candidate for providing a readout of activity and for regulating homeostatic plasticity

(O'Leary et al., 2014). In this study we asked whether metabolically associated signals, such

as ROS, might also be involved, as these could provide a proxy measure of neuronal activity

(Attwell and Laughlin, 2001; Zhu et al., 2012). Over-activation measurably impacts on the

ADP:ATP ratio (Tantama et al., 2013) and generates mitochondrial ROS. These can

modulate calcium dependent signaling pathways, such as CamKII (Hongpaisan et al., 2004),

and potentially also other ROS sensitive pathway components including IP_3 and Ryanodine

receptor channels or the protein phosphatase Calcineurin (Hidalgo and Arias-Cavieres,

2016). We found that a mitochondrially targeted ROS sensor (Albrecht et al., 2011; Gutscher

et al., 2009) became progressively oxidized with increasing levels of neuronal over-activation

(Figure 3A). Moreover, ROS were necessary for (and their elevation was also sufficient to

mimic) implementing activity-induced structural plasticity at synaptic terminals (Figure 3B,

C). Thus, our data show that ROS, particularly H_2O_2, are both necessary and sufficient for

activity regulated structural plasticity. The source of the activity-generated ROS appears to

be mitochondrial, potentially generated as a byproduct of increased ATP metabolism or

triggered by mitochondrial calcium influx (Peng and Jou, 2010). However, other ROS

generating systems could also be involved, notably NADPH oxidases, some of which are

regulated by calcium (Kishida et al., 2006; 2005; Serrano et al., 2003; Tejada-Simon et al.,

2005).

We discovered that in neurons the highly conserved protein DJ-1ß is critical for

responding to activity-generated ROS with structural changes to synaptic terminals (Figure

4A, B, C), and propose that in neurons DJ-1ß acts as a redox sensor for activity-generated

ROS. In agreement with this idea, DJ-1ß has been shown to be oxidized at the conserved

cysteine residue C106 (C104 in *Drosophila*) by H_2O_2. Oxidation of DJ-1 leads to changes in

DJ-1 function, including translocation from the cytoplasm to the mitochondrial matrix,

which aids with protection against oxidative damage (Blackinton et al., 2009; Canet-Avilés

and Wilson, 2004; Waak et al., 2009) and maintenance of ATP levels (Calì et al., 2015). We

found that motoneurons are potently sensitive to DJ-1ß dosage with regard to their response

to altered activity and ROS levels, and expression of mutant DJ-1ß[C104A] in motoneurons

blocks their structural plasticity response to DEM and over-activation. These observations

suggest that DJ-1ß forms a critical part of the ROS sensing mechanism in neurons. It is

therefore conceivable that different sensitivities to neuronal activity could in part be

determined by cell type-specific levels of DJ-1ß and associated reducing mechanisms.

414 DJ-1ß downstream pathways implement activity-regulated plasticity

Our data suggest that DJ-1ß forms part of a signaling hub required for at least two opposing

types of structural plasticity that seem to be implemented by distinct pathways: potentiation

in response to low-level increases in neuronal activation versus compensatory changes

following stronger over-activation. In this study we demonstrated that DJ-1ß mediates the

former by activation of PI3Kinase signaling following redox-regulated binding and inhibition

of PTEN (Figure 4A) (R. H. Kim et al., 2005; Y.-C. Kim et al., 2009). PTEN and PI3Kinase

are both intermediates of metabolic signaling pathways and regulators of synaptic terminal

growth (Jordán-Álvarez et al., 2012; Martín-Peña et al., 2006). While activity and ROS levels

correlate positively with PTEN/PI3Kinase mediated bouton addition at the NMJ, active zone

number regulation is more nuanced; at a higher activity threshold other, yet to be identified

effector pathways appear to be activated that reduce presynaptic active zone number (Figure

2D, E). Because this too is dependent on DJ-1ß we would expect this pathway also to be

regulated by activity-generated ROS. Equally unresolved is the question of how postsynaptic

dendritic arbor size in the CNS is regulated. PTEN/PI3Kinase signaling is an unlikely

candidate as this normally stimulates neuronal and dendritic growth (Brierley et al., 2009).

431 ROS as gatekeepers of activity-dependent synaptic structural plasticity

Synaptic plasticity is regulated by a number of signaling pathways, including Wnts (Budnik

and Salinas, 2011), BMPs (Bayat et al., 2011; Berke et al., 2013), as well as PKA, CREB and

AP-1 (Cho et al., 2015; Davis, 2006; Davis et al., 1998; 1996; Davis and Müller, 2015; S. M.

Kim et al., 2009; Koles and Budnik, 2012; Osses and Henriquez, 2014; Sanyal et al., 2002;

2003; Sulkowski et al., 2014; Walker et al., 2013). Here we demonstrated a requirement for

ROS for activity-dependent structural plasticity at the *Drosophila* NMJ and motoneuron

dendrites. Like BMP (Berke et al., 2013) and AP-1 (Milton et al., 2011; Sanyal et al., 2002)

ROS have the potential to act as gatekeepers of neuronal plasticity. If acting as a parallel

stimulus, ROS signaling at the synapse could be synergistic with other plasticity pathways,

which could help overcome issues with biological noise and increase the fidelity of cellular

decision making. Alternatively, ROS might deliver requisite post-translational redox

modifications to components of other signaling pathways. In support of the latter, ROS

regulate by redox modification the activity of the immediate early genes Jun and Fos, which

are required for LTP in vertebrates and in *Drosophila* for activity-dependent plasticity of

motoneurons, both at the NMJ and central dendrites (Hartwig et al., 2008; Jindra et al.,

2004; Loebrich and Nedivi, 2009; Milton et al., 2011; Milton and Sweeney, 2012; Sanyal et

al., 2002). ROS modulation of BMP signaling has recently been shown in cultured

sympathetic neurons (Chandrasekaran et al., 2015) and of Wnt pathways in non-neuronal

cells (Funato et al., 2006; Love et al., 2013; Rharass et al., 2014).

Although we focused exclusively on structural plasticity in the *Drosophila* larva,

current evidence suggests synaptic ROS signaling is a conserved and central feature of

communication in the nervous system. Several studies have demonstrated a requirement for

ROS for Long Term Potentiation (Huddleston et al., 2008; Kamsler and Segal, 2003b;

2003a; Klann, 1998; Knapp and Klann, 2002; Lee et al., 2010) and also correlated defects in

learning with altered ROS signaling (Kishida et al., 2006). The underlying cellular and

molecular mechanisms though are far from clear, probably because of the complexity of ROS

signaling and because of the large number of potential redox sensitive targets. For example,

components of the presynaptic release machinery, notably SNAP25, are thought to be

modulated by ROS (Giniatullin et al., 2006). Here, we identified DJ-1ß as a ROS sensor in

neurons whose interactions with the PTEN/PI3Kinase synaptic growth pathway are

governed by redox regulation. This represents a novel mechanism by which ROS, potentially

generated as metabolic byproducts, are used to tune connectivity and communication within the nervous system through modifications of neuronal structure. Since ROS can diffuse through membranes these signals have the potential to act non-autonomously, possibly across synaptic partners. Our findings shine a new light on ROS which, when dysregulated with age or under neurodegenerative conditions, might interfere with neuronal adaptive adjustments and thus contribute to network malfunction and synapse loss.

Competing interests: The authors declare that no competing interests exist.

Author Contributions:

M.O., M.F.Z., S.T.S. & M.L. devised the project. M.O. performed most experiments and analyses. P.B., A.M., M.F.Z. & K.M. generated specimens and dendrite data on activity dependent dendrite growth. A.M. carried out experiments and analysis involving ROS measurements. R.J.H.W. processed specimens for TEM imaging and analysed TEM data. M.O. & M.L. co-wrote the manuscript and S.T.S. & M.L. co-supervised the project.

Acknowledgements:

We would like to thank Nancy Bonini, Tobias Dick, Jörg Grosshans, Karen Hibbard, Fanis Missirlis, Barret Pfeiffer and Alex Whitworth for generous fly stock and reagent donations, Other fly stocks were obtained from the Bloomington Stock Center. The monoclonal antibodies nc82 (anti-Bruchpilot), developed by E. Buchner and colleagues, was obtained from the Developmental Studies Hybridoma Bank, created by the NICHD of the NIH and maintained at The University of Iowa, Department of Biology. We thank Akinao Nose and Hiroshi Kohsaka for kindly sharing figure graphics, Benjamin Risse for his generous support with setting up a FIM Tracker and Matthew Wayland for his generous support with data analysis and imaging. We would also like to thank Jimena Berni, Alex Whitworth and members of the Landgraf lab for valuable comments on the manuscript. This work was supported by BBSRC research grants (BB/IO1179X/1, BB/M002934/1) to ML and

492 (BB/I012273/1, BB/M002322/1) to STS. This work benefited from facilities supported by a

493 Wellcome Trust Equipment Grant (WT079204) and contributions by the Sir Isaac Newton

494 Trust in Cambridge.

Materials and Methods

Fly Strains and Husbandry. Wild-type and transgenic strains were maintained on

standard yeast–agar–cornmeal medium at 25 °C. The following fly strains were used:

OregonR and *PTEN^{C076}* (Bloomington Stock Center, Indiana University), *UAS-dTrpA1*

(Hamada et al., 2008), *UAS-SOD2* (Missirlis et al., 2003), *UAS-Catalase* (Missirlis et al.,

2001), *UAS-Duox* (Ha, 2005), *DJ-1β^{Δ93}* (Meulener et al., 2005), *UAS-DJ-1β^{C104A}* (Meulener

et al., 2006), UAS-RNAi lines targeting *SOD1*, *SOD2*, *Catalase* and *PTEN* (KK collection,

Vienna Drosophila Resource Centre) (Dietzl et al., 2007), *UAS-PI3K^{DN}* (Leevers et al.,

1996), *UAS-PTEN* (Gao et al., 2000). The following two GAL4 expression lines were used to

target GAL4 to the aCC and RP2 motoneurons: *aCC-FLP-GAL4* (e*veRN2-Flippase, UAS-*

myr::mRFP1, UAS-Flp, tubulin84B-FRT-CD2-FRT-GAL4) (Roy et al., 2007) expresses

GAL4 stochastically in single aCC and RP2 motoneurons allowing the imaging of aCC

neurons in isolation, as required for dendritic arbor resolution and reconstruction.

aCC/RP2-GAL4 (*eveRN2-GAL4* (Fujioka et al., 2003)*, UAS-myr-mRFP1, UAS-Flp,*

tubulin84B-FRT-CD2-FRT-GAL4; RRaGAL4, 20xUAS-6XmCherry::HA (Shearin et al.,

2014)) was used for NMJ analysis as it expresses GAL4 in every aCC and RP2 motoneuron.

eveRN2-GAL4 expression is restricted to the embryo and FLPase-gated *tubulin84B-FRT-*

CD2-FRT-GAL4 maintains GAL4 expression at high levels during larval stages.

Dissection and Immunocytochemistry

1st Instar Ventral Nerve Cord (VNC). Flies were allowed to lay eggs on apple juice-based

agar medium for 24 hrs at 25°C. Embryos were dechorionated using bleach (3.5 minutes

room temperature) then incubated (25°C) in pre-warmed Sorensen's saline (pH 7.2, 0.075

M) whilst adhered to a petri dish. Hatched larvae (floating) were recovered hourly and

transferred to pre-warmed apple-juice agar plates supplemented with yeast paste. Larvae

were allowed to develop for a further 24 hrs (24 hrs after larval hatching, ALH) at 25°C, 27°C

or 29°C prior to dissection in Sorensen's saline. A fine hypodermic needle (30 1/2 G;

Microlance) was used as a scalpel to cut off the anterior end of each larva, allowing gut, fat

body, and trachea to be removed. The ventral nerve chord and brain lobes, extruded with

viscera upon decapitation, were dissected out and transferred to a cover glass coated with

poly-L-lysine (Sigma-Aldrich), positioned dorsal side up in Sorensen's saline. A clean cover

glass was placed on top of the preparation, with strips of double-sided sticky tape as spacers

positioned along the edges.

Wandering 3rd Instar. Flies were allowed to lay eggs on apple-juice agar based medium

overnight at 25°C, larvae were then incubated at 25°C or 27°C until the late wandering 3rd

instar stage. Larvae were reared on yeast paste colored with Bromophenol Blue sodium salt

(Sigma-Aldrich) to allow visualization of gut-clearance, an indicator of the late wandering 3rd

instar stage. For Di-ethylmaleate (DEM) (Sigma-Aldrich) and paraquat (Sigma-Aldrich)

feeding, yeast paste was made using a 5 mM – 15 mM aqueous solution. Larvae were 'fillet'

dissected in Sorensen's saline and fixed for 15 minutes at room temperature in 4%

formaldehyde (in Sorenson's saline). Specimens were then washed and stained in Sorensen's

saline containing 0.3% Triton X-100 (Sigma-Aldrich) using the following primary and

secondary antibodies; Goat-anti-HRP Alexa Fluor 488 (1:400) or Goat-anti-HRP Alexa Fluor

594 (1:400) (Jackson ImmunoResearch Cat. Nos. 123-455-021 or 123-585-021), Rabbit-

anti-DsRed (1:1200) (ClonTech Cat. No. 632496), Donkey-anti-Rabbit CF568 (1:1200)

(Biotium Cat. No. 20098), Mouse-nc82 (1:100) (Developmental Studies Hybridoma Bank),

Donkey-anti-Mouse Alex Fluor 488 (1:600) (ThermoFisher Cat. No. A-21202) or Donkey-

anti-Mouse CF633 (1:300) (Biotium Cat. No. 20124) incubated overnight at 4°C or 2 hrs at

room temperature. Specimens were mounted in EverBrite mounting medium (Biotium Cat.

No. 23001).

Image Acquisition and Analysis

1st Instar Ventral Nerve Cord (VNC). Ventral nerve cords were pre-screened for

fluorescently labeled, isolated, aCC motoneurons using a Zeiss Axiophot compound

epifluorescence microscope and a Zeiss Plan-Neofluar 40x/0.75 N.A. objective lens. Suitable

VNCs were imaged immediately with a Yokagawa CSU-22 spinning disk confocal field

scanner mounted on an Olympus BX51WI microscope, using a 60×/1.2 N.A. Olympus water

immersion objective. Images were acquired with a voxel size of 0.2 × 0.2 × 0.3 μm using a

QuantEM cooled EMCCD camera (Photometrics), operated via MetaMorph software

(Molecular Devices). Dendritic trees were digitally reconstructed using Amira Resolve RT 4.1

(FEI), supplemented with a 3D reconstruction algorithm (Evers et al., 2005; Schmitt et al.,

2004), and images were processed using Amira (FEI) and ImageJ (National Institutes of

Health).

Wandering 3rd Instar. Dissected specimens were imaged using a Leica SP5 point-scanning

confocal, and a 63x/1.3 N.A. (Leica) glycerol immersion objective lens and LAS AF (Leica

Application Suite Advanced Fluorescence) software. Confocal images were processed using

ImageJ and Photoshop (Adobe). Bouton number of the NMJ on muscle DA1 [1] from

segments A3-A5 was determined by counting every distinct spherical varicosity along the

NMJ branch. DA1 muscles were imaged using a Zeiss Axiophot compound microscope and a

Zeiss Plan-Neofluar 10x/0.3 N.A. objective lens. Muscle surface area (MSA) was determined

by multiplying muscle length by width using ImageJ. In order to correct for subtle

differences in animal size (typically 5-10%) bouton number normalization was performed

using the following calculation: (mean control MSA / mean experimental MSA) x test bouton

number = normalized experimental bouton number.

Ratiometric ROS Reporter. *aCC/RP2-Gal4* was used to drive the expression of *UAS-mito-roGFP2-Orp1* (Albrecht et al., 2011; Gutscher et al., 2009) in all aCC and RP2 motoneurons. Wandering third instar larvae were fillet dissected in PBS-NEM (137mM NaCl, 2.7mM KCl, 10mM Na_2HPO_4, 1.8mM KH_2PO_4, 20mM N-ethylmaleimide (NEM), pH 7.4). Larval fillet preparations were incubated for 5 minutes in PBS-NEM then fixed for 8 minutes in 4% formaldehyde (in PBS-NEM). Specimens were washed three times in PBS-NEM and then equilibrated in 70% glycerol. Specimens were mounted in glycerol and imaged the same day. Imaging was performed on a Leica SP5 point-scanning confocal, using a 63x/1.3 N.A. (Leica) glycerol immersion objective lens. The reporter was excited sequentially at 405nm and 488nm (Albrecht et al., 2011) with emission detected at 500–535nm. 16-bit images were acquired using Leica LAS AF software and processed using ImageJ. Z-stack images were maximally projected and converted to 32-bit. To remove fringing artefacts around bouton edges 488nm images were thresholded using the 'Intermodes' algorithm with values below threshold set to 'not a number', and ratio images were created by dividing the 405nm image by the 488nm image pixel by pixel (Albrecht et al., 2011). Regions of Interest were taken on the ratio image spanning the entire NMJ and the mean value obtained from each NMJ was used for statistical analysis.

Transmission Electron Microscopy. Third instar wandering larvae were fillet dissected in PBS and fixed overnight in 0.1M $NaPO_4$, pH 7.4, 1% glutaraldehyde, and 4% formaldehyde, pH 7.3. Fixed specimens were washed 3× in 0.1M $NaPO_4$ before incubation in OsO_4 (1% in 0.1M $NaPO_4$; 2hr). Preparations were washed 3× in distilled water, incubated in 1% uranyl acetate, then washed again 3× in distilled water and dehydrated through a graded ethanol series: 20% increments starting at 30% followed by two 100% changes and then 2× 100% propylene oxide. Specimens were incubated in a graded series of epon araldite resin (in propylene oxide): 25% increments culminating in 3× 100% changes. Individual muscles were then dissected and transferred into embedding molds, followed by polymerization at 60°C for 48 hrs. Resin mounted specimens were sectioned (60–70 nm)

using glass knives upon a microtome (Ultracut UCT; Leica). Sections were placed onto grids, incubated in uranyl acetate (50% in ethanol), washed in distilled water and incubated in lead citrate. Sections were imaged using a transmission electron microscope (TECNAI 12 G^2; FEI) with a camera (Soft Imaging Solutions MegaView; Olympus) and Tecnai user interface v2.1.8 and analySIS v3.2 (Soft Imaging Systems).

Behavior – larval crawling analysis

To record larval crawling, mid-3rd instar larvae (72hrs ALH) were briefly rinsed in water to remove any food and yeast residues, then up to 12 larvae were placed into a 24cm x 24cm arena of 0.8% agar in water, poured to 5 mm thickness. Crawling behavior was recorded in a temperature and humidity controlled incubator at temperatures ranging from 25-32^{o}C, as indicated for each experiment. Larvae were allowed to acclimatise for 5 minutes, then recorded for 15 minutes under infrared LED illumination (intensity from 14.33 nW/mm^2 in the edge to 9.12 nW/mm^2 in the center), using frustrated total internal reflection using a modified FIM tracker (Milton et al., 2011; Risse et al., 2013) https://www.uni-muenster.de/PRIA/en/FIM/index.html. Larvae were recorded with a Basler acA2040-180km CMOS camera using Pylon and StreamPix software, mounted with a 16mm KOWA IJM3sHC.SW VIS-NIR Lens. Images were acquired at 5 frames per second. For each larvae, average crawling speed was calculated from long, uninterrupted forward crawls identified manually using FIMTrack. The 15 minute recording period was partitioned into 5 minute sections with each larvae being assayed once within each section, allowing each specimen to be sampled a maximum of 3 times. We observed no change in average crawling speed within the duration of the 15-minute recording.

Data Analysis. All data handling was performed using Prism software (GraphPad). NMJ bouton number and ratiometric ROS reporter data were tested for normal / Gaussian distribution using the D'Agostino-Pearson omnibus normality test. Due to a lower replicate number, dendritic arbor reconstruction data were tested for normality using the

630   Kolmogorov-Smirnov with Dallal-Wilkinson-Lilliefor P value test. Normal distribution was

631   thus confirmed for all data presented, which were compared using one-way analysis of

632   variance (ANOVA), with Tukey's multiple comparisons test where $*P<0.05$, $**P<0.01$,

633    $***P<0.001$, $****P<0.0001$.

Figure Supplement Figures

Figure 1-Figure Supplement 1: Prolonged over-activation targeted to motoneurons (*D42-GAL4*; *UAS-dTrpA1*) also leads to adaptation with reduced crawling speed (dTRPA1 channels open at 27°C, closed at 22°C). Control is *D42-GAL4 / +*. Each data point represents the crawling speed from an individual uninterrupted continuous forward crawl, n = specimen replicate number, up to 3 crawls measured for each larva. ****P<0.0001 students t test.

OregonR   DJ-1ß$^{\Delta 93}$

500nm

Figure 4-Figure Supplement 1: *DJ-1β$^{\Delta 93}$* mutant 3rd instar larval NMJs are phenotypically normal. Representative TEM bouton cross-sectional images showing that pre-synaptic architecture is intact including active-zones with associated clustered synaptic vesicles (arrowed).

Figure 4-Figure Supplement 2: A. *DJ-1β* null mutant (*DJ-1β^{Δ93}*) larvae do not show systemic activity-dependent NMJ elaboration. Normalised bouton number dot plot showing significantly increased NMJ elaboration in *OregonR*, but not *DJ-1β^{Δ93}*, larvae reared at 29°C vs 25°C. **B.** *DJ-1β^{Δ93}* larvae also do not show systemic activity-dependent increase in active zone number. *OregonR*, but not *DJ-1β^{Δ93}*, show significantly increased NMJ active zone number when reared at 29°C vs 25°C. Mean +/- SEM, ANOVA **p<0.01, ***p<0.001, n=replicate number.

References

653
Albrecht, S.C., Barata, A.G., Grosshans, J., Teleman, A.A., Dick, T.P., 2011. In vivo mapping
of hydrogen peroxide and oxidized glutathione reveals chemical and regional specificity
of redox homeostasis. Cell Metab 14, 819–829. doi:10.1016/j.cmet.2011.10.010
Ariga, H., Takahashi-Niki, K., Kato, I., Maita, H., Niki, T., Iguchi-Ariga, S.M.M., 2013.
Neuroprotective function of DJ-1 in Parkinson's disease. Oxid Med Cell Longev 2013,
683920. doi:10.1155/2013/683920
Ataman, B., Ashley, J., Gorczyca, M., Ramachandran, P., Fouquet, W., Sigrist, S.J., Budnik,
V., 2008. Rapid activity-dependent modifications in synaptic structure and function
require bidirectional Wnt signaling. Neuron 57, 705–718.
doi:10.1016/j.neuron.2008.01.026
Attwell, D., Laughlin, S.B., 2001. An energy budget for signaling in the grey matter of the
brain. J. Cereb. Blood Flow Metab. 21, 1133–1145. doi:10.1097/00004647-200110000-
00001
Baines, R.A., Robinson, S.G., Fujioka, M., Jaynes, J.B., Bate, M., 1999. Postsynaptic
expression of tetanus toxin light chain blocks synaptogenesis in Drosophila. Curr Biol 9,
1267–1270.

Baines, R.A., Uhler, J.P., Thompson, A., Sweeney, S.T., Bate, M., 2001. Altered electrical properties in Drosophila neurons developing without synaptic transmission. J Neurosci 21, 1523–1531.

Bayat, V., Jaiswal, M., Bellen, H.J., 2011. The BMP signaling pathway at the Drosophila neuromuscular junction and its links to neurodegenerative diseases. Curr Opin Neurobiol 21, 182–188. doi:10.1016/j.conb.2010.08.014

Berke, B., Wittnam, J., McNeill, E., Van Vactor, D.L., Keshishian, H., 2013. Retrograde BMP signaling at the synapse: a permissive signal for synapse maturation and activity-dependent plasticity. J Neurosci 33, 17937–17950. doi:10.1523/JNEUROSCI.6075-11.2013

Bindokas, V.P., Jordán, J., Lee, C.C., Miller, R.J., 1996. Superoxide production in rat hippocampal neurons: selective imaging with hydroethidine. J. Neurosci. 16, 1324–1336.

Blackinton, J., Lakshminarasimhan, M., Thomas, K.J., Ahmad, R., Greggio, E., Raza, A.S., Cookson, M.R., Wilson, M.A., 2009. Formation of a stabilized cysteine sulfinic acid is critical for the mitochondrial function of the parkinsonism protein DJ-1. J Biol Chem 284, 6476–6485. doi:10.1074/jbc.M806599200

Bonifati, V., Rizzu, P., van Baren, M.J., Schaap, O., Breedveld, G.J., Krieger, E., Dekker, M.C.J., Squitieri, F., Ibanez, P., Joosse, M., van Dongen, J.W., Vanacore, N., van Swieten, J.C., Brice, A., Meco, G., van Duijn, C.M., Oostra, B.A., Heutink, P., 2003. Mutations in the DJ-1 gene associated with autosomal recessive early-onset parkinsonism. Science 299, 256–259. doi:10.1126/science.1077209

Brennan, A.M., Won Suh, S., Joon Won, S., Narasimhan, P., Kauppinen, T.M., Lee, H., Edling, Y., Chan, P.H., Swanson, R.A., 2009. NADPH oxidase is the primary source of superoxide induced by NMDA receptor activation. Nat Neurosci 12, 857–863. doi:10.1038/nn.2334

Brierley, D.J., Blanc, E., Reddy, O.V., VijayRaghavan, K., Williams, D.W., 2009. Dendritic targeting in the leg neuropil of Drosophila: the role of midline signalling molecules in generating a myotopic map. PLoS Biol 7, e1000199. doi:10.1371/journal.pbio.1000199

Budnik, V., Salinas, P.C., 2011. Wnt signaling during synaptic development and plasticity. Curr Opin Neurobiol 21, 151–159. doi:10.1016/j.conb.2010.12.002

Budnik, V., Zhong, Y., Wu, C.F., 1990. Morphological plasticity of motor axons in Drosophila mutants with altered excitability. J. Neurosci. 10, 3754–3768.

Burrone, J., O'Byrne, M., Murthy, V.N., 2002. Multiple forms of synaptic plasticity triggered by selective suppression of activity in individual neurons. Nature 420, 414–418. doi:10.1038/nature01242

Calì, T., Ottolini, D., Soriano, M.E., Brini, M., 2015. A new split-GFP-based probe reveals DJ-1 translocation into the mitochondrial matrix to sustain ATP synthesis upon nutrient deprivation. Hum Mol Genet 24, 1045–1060. doi:10.1093/hmg/ddu519

Canet-Avilés, R.M., Wilson, M.A., 2004. The Parkinson's disease protein DJ-1 is neuroprotective due to cysteine-sulfinic acid-driven mitochondrial localization, in:. Presented at the Proceedings of the ….

Chandrasekaran, V., Lea, C., Sosa, J.C., Higgins, D., Lein, P.J., 2015. Reactive oxygen species are involved in BMP-induced dendritic growth in cultured rat sympathetic neurons. Mol Cell Neurosci 67, 116–125. doi:10.1016/j.mcn.2015.06.007

Cho, R.W., Buhl, L.K., Volfson, D., Tran, A., Li, F., Akbergenova, Y., Littleton, J.T., 2015. Phosphorylation of Complexin by PKA Regulates Activity-Dependent Spontaneous Neurotransmitter Release and Structural Synaptic Plasticity. Neuron 88, 749–761. doi:10.1016/j.neuron.2015.10.011

Choi, J.C., Park, D., Griffith, L.C., 2004. Electrophysiological and morphological characterization of identified motor neurons in the Drosophila third instar larva central nervous system. J Neurophysiol 91, 2353–2365. doi:10.1152/jn.01115.2003

Crossley, A.C., 1978. Morphology and development of the Drosophila muscular system, in: Ashburner, M., Wright, T. (Eds.), Genetics and Biology of Drosophila. Genetics and biology of Drosophila, New York, pp. 499–560.

Davis, G.W., 2006. Homeostatic control of neural activity: from phenomenology to

molecular design. Annu Rev Neurosci 29, 307–323.
doi:10.1146/annurev.neuro.28.061604.135751

Davis, G.W., Davis, G.W., Schuster, C.M., Schuster, C.M., Goodman, C.S., Goodman, C.S., 1996. Genetic dissection of structural and functional components of synaptic plasticity. III. CREB is necessary for presynaptic functional plasticity. Neuron 17, 669–679.

Davis, G.W., DiAntonio, A., Petersen, S.A., Goodman, C.S., 1998. Postsynaptic PKA controls quantal size and reveals a retrograde signal that regulates presynaptic transmitter release in Drosophila. Neuron 20, 305–315.

Davis, G.W., Müller, M., 2015. Homeostatic control of presynaptic neurotransmitter release. Annu Rev Physiol 77, 251–270. doi:10.1146/annurev-physiol-021014-071740

Davydova, D., Marini, C., King, C., Klueva, J., Bischof, F., Romorini, S., Montenegro-Venegas, C., Heine, M., Schneider, R., Schröder, M.S., Altrock, W.D., Henneberger, C., Rusakov, D.A., Gundelfinger, E.D., Fejtova, A., 2014. Bassoon specifically controls presynaptic P/Q-type Ca(2+) channels via RIM-binding protein. Neuron 82, 181–194. doi:10.1016/j.neuron.2014.02.012

Dietzl, G., Chen, D., Schnorrer, F., Su, K.-C., Barinova, Y., Fellner, M., Gasser, B., Kinsey, K., Oppel, S., Scheiblauer, S., Couto, A., Marra, V., Keleman, K., Dickson, B.J., 2007. A genome-wide transgenic RNAi library for conditional gene inactivation in Drosophila. Nature 448, 151–156. doi:10.1038/nature05954

Dillon, M.E., Wang, G., Garrity, P.A., Huey, R.B., 2009. Review: Thermal preference in Drosophila. J. Therm. Biol. 34, 109–119. doi:10.1016/j.jtherbio.2008.11.007

Driscoll, H.E., Muraro, N.I., He, M., Baines, R.A., 2013. Pumilio-2 regulates translation of nav1.6 to mediate homeostasis of membrane excitability. J Neurosci 33, 9644–9654. doi:10.1523/JNEUROSCI.0921-13.2013

Dugan, L.L., Sensi, S.L., Canzoniero, L.M., Handran, S.D., Rothman, S.M., Lin, T.S., Goldberg, M.P., Choi, D.W., 1995. Mitochondrial production of reactive oxygen species in cortical neurons following exposure to N-methyl-D-aspartate. J. Neurosci. 15, 6377–6388.

Evers, J.F., Schmitt, S., Sibila, M., Duch, C., 2005. Progress in functional neuroanatomy: precise automatic geometric reconstruction of neuronal morphology from confocal image stacks. J Neurophysiol 93, 2331–2342. doi:10.1152/jn.00761.2004

Finkel, T., 2011. Signal transduction by reactive oxygen species. J Cell Biol 194, 7–15. doi:10.1083/jcb.201102095

Frank, C.A., Kennedy, M.J., Goold, C.P., Marek, K.W., Davis, G.W., 2006. Mechanisms underlying the rapid induction and sustained expression of synaptic homeostasis. Neuron 52, 663–677. doi:10.1016/j.neuron.2006.09.029

Frank, C.A., Pielage, J., Davis, G.W., 2009. A presynaptic homeostatic signaling system composed of the Eph receptor, ephexin, Cdc42, and CaV2.1 calcium channels. Neuron 61, 556–569. doi:10.1016/j.neuron.2008.12.028

Fujioka, M., Lear, B.C., Landgraf, M., Yusibova, G.L., Zhou, J., Riley, K.M., Patel, N.H., Jaynes, J.B., 2003. Even-skipped, acting as a repressor, regulates axonal projections in Drosophila. Development 130, 5385–5400. doi:10.1242/dev.00770

Funato, Y., Michiue, T., Asashima, M., Miki, H., 2006. The thioredoxin-related redox-regulating protein nucleoredoxin inhibits Wnt-beta-catenin signalling through dishevelled. Nat Cell Biol 8, 501–508. doi:10.1038/ncb1405

Fushiki, A., Zwart, M.F., Kohsaka, H., Fetter, R.D., Cardona, A., Nose, A., 2016. A circuit mechanism for the propagation of waves of muscle contraction in Drosophila. Elife 5. doi:10.7554/eLife.13253

Gahtan, E., Auerbach, J.M., Groner, Y., Segal, M., 1998. Reversible impairment of long-term potentiation in transgenic Cu/Zn-SOD mice. Eur J Neurosci 10, 538–544.

Gao, X., Neufeld, T.P., Pan, D., 2000. Drosophila PTEN regulates cell growth and proliferation through PI3K-dependent and -independent pathways. Dev Biol 221, 404–418. doi:10.1006/dbio.2000.9680

Gaviño, M.A., Ford, K.J., Archila, S., Davis, G.W., 2015. Homeostatic synaptic depression is achieved through a regulated decrease in presynaptic calcium channel abundance. Elife

4. doi:10.7554/eLife.05473

Giachello, C.N., Baines, R.A., 2016. Regulation of motoneuron excitability and the setting of homeostatic limits. Curr Opin Neurobiol 43, 1–6. doi:10.1016/j.conb.2016.09.014

Giniatullin, A.R., Darios, F., Shakirzyanova, A., Davletov, B., Giniatullin, R., 2006. SNAP25 is a pre-synaptic target for the depressant action of reactive oxygen species on transmitter release. J Neurochem 98, 1789–1797. doi:10.1111/j.1471-4159.2006.03997.x

Gjorgjieva, J., Evers, J.F., Eglen, S.J., 2016. Homeostatic Activity-Dependent Tuning of Recurrent Networks for Robust Propagation of Activity. J Neurosci 36, 3722–3734. doi:10.1523/JNEUROSCI.2511-15.2016

Gutscher, M., Sobotta, M.C., Wabnitz, G.H., Ballikaya, S., Meyer, A.J., Samstag, Y., Dick, T.P., 2009. Proximity-based protein thiol oxidation by H2O2-scavenging peroxidases. J Biol Chem 284, 31532–31540. doi:10.1074/jbc.M109.059246

Ha, E.M., 2005. A Direct Role for Dual Oxidase in Drosophila Gut Immunity. Science 310, 847–850. doi:10.1126/science.1117311

Hallermann, S., de Kock, C.P.J., Stuart, G.J., Kole, M.H.P., 2012. State and location dependence of action potential metabolic cost in cortical pyramidal neurons. Nat Neurosci 15, 1007–1014. doi:10.1038/nn.3132

Halliwell, B., 1992. Reactive oxygen species and the central nervous system. J Neurochem 59, 1609–1623.

Hamada, F.N., Hamada, F.N., Rosenzweig, M., Rosenzweig, M., Kang, K., Kang, K., Pulver, S.R., Pulver, S.R., Ghezzi, A., Ghezzi, A., Jegla, T.J., Jegla, T.J., Garrity, P.A., Garrity, P.A., 2008. An internal thermal sensor controlling temperature preference in Drosophila. Nature 454, 217–220. doi:10.1038/nature07001

Hardingham, G.E., Arnold, F.J., Bading, H., 2001. A calcium microdomain near NMDA receptors: on switch for ERK-dependent synapse-to-nucleus communication. Nat Neurosci 4, 565–566. doi:10.1038/88380

Hartwig, C.L., Worrell, J., Levine, R.B., Ramaswami, M., Sanyal, S., 2008. Normal dendrite growth in Drosophila motor neurons requires the AP-1 transcription factor. Dev Neurobiol 68, 1225–1242. doi:10.1002/dneu.20655

Hengen, K.B., Lambo, M.E., Van Hooser, S.D., Katz, D.B., Turrigiano, G.G., 2013. Firing rate homeostasis in visual cortex of freely behaving rodents. Neuron 80, 335–342. doi:10.1016/j.neuron.2013.08.038

Hengen, K.B., Torrado Pacheco, A., McGregor, J.N., Van Hooser, S.D., Turrigiano, G.G., 2016. Neuronal Firing Rate Homeostasis Is Inhibited by Sleep and Promoted by Wake. Cell 165, 180–191. doi:10.1016/j.cell.2016.01.046

Hidalgo, C., Arias-Cavieres, A., 2016. Calcium, Reactive Oxygen Species, and Synaptic Plasticity. Physiology (Bethesda) 31, 201–215. doi:10.1152/physiol.00038.2015

Hongpaisan, J., Winters, C.A., Andrews, S.B., 2004. Strong calcium entry activates mitochondrial superoxide generation, upregulating kinase signaling in hippocampal neurons. J Neurosci 24, 10878–10887. doi:10.1523/JNEUROSCI.3278-04.2004

Hongpaisan, J., Winters, C.A., Andrews, S.B., 2003. Calcium-dependent mitochondrial superoxide modulates nuclear CREB phosphorylation in hippocampal neurons. Molecular and Cellular Neuroscience 24, 1103–1115.

Huddleston, A.T., Tang, W., Takeshima, H., Hamilton, S.L., Klann, E., 2008. Superoxide-induced potentiation in the hippocampus requires activation of ryanodine receptor type 3 and ERK. J Neurophysiol 99, 1565–1571. doi:10.1152/jn.00659.2007

Itakura, Y., Kohsaka, H., Ohyama, T., Zlatic, M., Pulver, S.R., Nose, A., 2015. Identification of Inhibitory Premotor Interneurons Activated at a Late Phase in a Motor Cycle during Drosophila Larval Locomotion. PLoS ONE 10, e0136660. doi:10.1371/journal.pone.0136660

Jindra, M., Gaziova, I., Uhlirova, M., Okabe, M., Hiromi, Y., Hirose, S., 2004. Coactivator MBF1 preserves the redox-dependent AP-1 activity during oxidative stress in Drosophila. EMBO J 23, 3538–3547. doi:10.1038/sj.emboj.7600356

Jordán-Álvarez, S., Fouquet, W., Sigrist, S.J., Acebes, A., 2012. Presynaptic PI3K activity triggers the formation of glutamate receptors at neuromuscular terminals of Drosophila.

J Cell Sci 125, 3621–3629. doi:10.1242/jcs.102806

Kamsler, A., Segal, M., 2003a. Hydrogen peroxide modulation of synaptic plasticity. J
Neurosci 23, 269–276.

Kamsler, A., Segal, M., 2003b. Paradoxical actions of hydrogen peroxide on long-term
potentiation in transgenic superoxide dismutase-1 mice. J Neurosci 23, 10359–10367.

Keck, T., Hübener, M., Bonhoeffer, T., 2017. Interactions between synaptic homeostatic
mechanisms: an attempt to reconcile BCM theory, synaptic scaling, and changing
excitation/inhibition balance. Curr Opin Neurobiol 43, 87–93.
doi:10.1016/j.conb.2017.02.003

Keck, T., Keller, G.B., Jacobsen, R.I., Eysel, U.T., Bonhoeffer, T., Hübener, M., 2013.
Synaptic scaling and homeostatic plasticity in the mouse visual cortex in vivo. Neuron
80, 327–334. doi:10.1016/j.neuron.2013.08.018

Kim, R.H., Peters, M., Jang, Y., Shi, W., Pintilie, M., Fletcher, G.C., DeLuca, C., Liepa, J.,
Zhou, L., Snow, B., Binari, R.C., Manoukian, A.S., Bray, M.R., Liu, F.-F., Tsao, M.-S.,
Mak, T.W., 2005. DJ-1, a novel regulator of the tumor suppressor PTEN. Cancer Cell 7,
263–273. doi:10.1016/j.ccr.2005.02.010

Kim, S.M., Kumar, V., Lin, Y.-Q., Karunanithi, S., Ramaswami, M., 2009. Fos and Jun
potentiate individual release sites and mobilize the reserve synaptic vesicle pool at the
Drosophila larval motor synapse. Proc Natl Acad Sci USA 106, 4000–4005.
doi:10.1073/pnas.0806064106

Kim, Y.-C., Kitaura, H., Taira, T., Iguchi-Ariga, S.M.M., Ariga, H., 2009. Oxidation of DJ-1-
dependent cell transformation through direct binding of DJ-1 to PTEN. Int J Oncol 35,
1331–1341.

Kirov, S.A., Harris, K.M., 1999. Dendrites are more spiny on mature hippocampal neurons
when synapses are inactivated. Nat Neurosci 2, 878–883. doi:10.1038/13178

Kishida, K.T., Hoeffer, C.A., Hu, D., Pao, M., Holland, S.M., Klann, E., 2006. Synaptic
plasticity deficits and mild memory impairments in mouse models of chronic
granulomatous disease. Mol Cell Biol 26, 5908–5920. doi:10.1128/MCB.00269-06

Kishida, K.T., Pao, M., Holland, S.M., Klann, E., 2005. NADPH oxidase is required for
NMDA receptor-dependent activation of ERK in hippocampal area CA1. J Neurochem
94, 299–306. doi:10.1111/j.1471-4159.2005.03189.x

Klann, E., 1998. Cell-permeable scavengers of superoxide prevent long-term potentiation in
hippocampal area CA1. J Neurophysiol 80, 452–457.

Knapp, L.T., Klann, E., 2002. Potentiation of hippocampal synaptic transmission by
superoxide requires the oxidative activation of protein kinase C. J Neurosci 22, 674–683.

Kohsaka, H., Okusawa, S., Itakura, Y., Fushiki, A., Nose, A., 2012. Development of larval
motor circuits in Drosophila. Dev Growth Differ 54, 408–419. doi:10.1111/j.1440-
169X.2012.01347.x

Kohsaka, H., Takasu, E., Morimoto, T., Nose, A., 2014. A group of segmental premotor
interneurons regulates the speed of axial locomotion in Drosophila larvae. Curr Biol 24,
2632–2642. doi:10.1016/j.cub.2014.09.026

Koles, K., Budnik, V., 2012. Wnt signaling in neuromuscular junction development. Cold
Spring Harb Perspect Biol 4. doi:10.1101/cshperspect.a008045

Landgraf, M., Jeffrey, V., Fujioka, M., Jaynes, J.B., Bate, M., 2003. Embryonic origins of a
motor system: motor dendrites form a myotopic map in Drosophila. PLoS Biol 1, E41.
doi:10.1371/journal.pbio.0000041

Lazarevic, V., Schöne, C., Heine, M., Gundelfinger, E.D., Fejtova, A., 2011. Extensive
remodeling of the presynaptic cytomatrix upon homeostatic adaptation to network
activity silencing. J Neurosci 31, 10189–10200. doi:10.1523/JNEUROSCI.2088-11.2011

Lee, K.Y., Chung, K., Chung, J.M., 2010. Involvement of reactive oxygen species in long-term
potentiation in the spinal cord dorsal horn. J Neurophysiol 103, 382–391.
doi:10.1152/jn.90906.2008

Leevers, S.J., Weinkove, D., MacDougall, L.K., Hafen, E., Waterfield, M.D., 1996. The
Drosophila phosphoinositide 3-kinase Dp110 promotes cell growth. EMBO J 15, 6584–
6594.

Levin, E.D., Brady, T.C., Hochrein, E.C., Oury, T.D., Jonsson, L.M., Marklund, S.L., Crapo,
J.D., 1998. Molecular manipulations of extracellular superoxide dismutase: functional
importance for learning. Behav. Genet. 28, 381–390.
Lin, W.-H., Günay, C., Marley, R., Prinz, A.A., Baines, R.A., 2012. Activity-dependent
alternative splicing increases persistent sodium current and promotes seizure. J
Neurosci 32, 7267–7277. doi:10.1523/JNEUROSCI.6042-11.2012
Loebrich, S., Nedivi, E., 2009. The function of activity-regulated genes in the nervous
system. Physiol Rev 89, 1079–1103. doi:10.1152/physrev.00013.2009
Love, N.R., Chen, Y., Ishibashi, S., Kritsiligkou, P., Lea, R., Koh, Y., Gallop, J.L., Dorey, K.,
Amaya, E., 2013. Amputation-induced reactive oxygen species are required for
successful Xenopus tadpole tail regeneration. Nat Cell Biol. doi:10.1038/ncb2659
Martín-Peña, A., Acebes, A., Rodríguez, J.-R., Sorribes, A., de Polavieja, G.G., Fernández-
Fúnez, P., Ferrús, A., 2006. Age-independent synaptogenesis by phosphoinositide 3
kinase. J Neurosci 26, 10199–10208. doi:10.1523/JNEUROSCI.1223-06.2006
Matz, J., Gilyan, A., Kolar, A., McCarvill, T., Krueger, S.R., 2010. Rapid structural alterations
of the active zone lead to sustained changes in neurotransmitter release. Proc Natl Acad
Sci U S A 107, 8836–8841. doi:10.1073/pnas.0906087107
Mee, C.J., Pym, E.C.G., Moffat, K.G., Baines, R.A., 2004. Regulation of neuronal excitability
through pumilio-dependent control of a sodium channel gene. J Neurosci 24, 8695–
8703. doi:10.1523/JNEUROSCI.2282-04.2004
Meulener, M., Whitworth, A.J., Armstrong-Gold, C.E., Rizzu, P., Heutink, P., Wes, P.D.,
Pallanck, L.J., Bonini, N.M., 2005. Drosophila DJ-1 mutants are selectively sensitive to
environmental toxins associated with Parkinson's disease. Curr Biol 15, 1572–1577.
doi:10.1016/j.cub.2005.07.064
Meulener, M.C., Xu, K., Thomson, L., Thompson, L., Ischiropoulos, H., Bonini, N.M., 2006.
Mutational analysis of DJ-1 in Drosophila implicates functional inactivation by oxidative
damage and aging. Proc Natl Acad Sci USA 103, 12517–12522.
doi:10.1073/pnas.0601891103
Milton, V.J., Jarrett, H.E., Gowers, K., Chalak, S., Briggs, L., Robinson, I.M., Sweeney, S.T.,
2011. Oxidative stress induces overgrowth of the Drosophila neuromuscular junction.
Proc Natl Acad Sci U S A 108, 17521–17526. doi:10.1073/pnas.1014511108
Milton, V.J., Sweeney, S.T., 2012. Oxidative stress in synapse development and function. Dev
Neurobiol 72, 100–110. doi:10.1002/dneu.20957
Missirlis, F., Phillips, J.P., Jäckle, H., 2001. Cooperative action of antioxidant defense
systems in Drosophila. Current Biology 11, 1272–1277.
Missirlis, F., Rahlfs, S., Dimopoulos, N., Bauer, H., Becker, K., Hilliker, A., Phillips, J.P.,
Jäckle, H., 2003. A putative glutathione peroxidase of Drosophila encodes a thioredoxin
peroxidase that provides resistance against oxidative stress but fails to complement a
lack of catalase activity. Biol Chem 384, 463–472. doi:10.1515/BC.2003.052
Murthy, V.N., Schikorski, T., Stevens, C.F., Zhu, Y., 2001. Inactivity produces increases in
neurotransmitter release and synapse size. Neuron 32, 673–682.
Müller, M., Davis, G.W., 2012. Transsynaptic control of presynaptic Ca^{2+} influx achieves
homeostatic potentiation of neurotransmitter release. Curr Biol 22, 1102–1108.
doi:10.1016/j.cub.2012.04.018
Nagakubo, D., Taira, T., Kitaura, H., Ikeda, M., Tamai, K., Iguchi-Ariga, S.M., Ariga, H.,
1997. DJ-1, a novel oncogene which transforms mouse NIH3T3 cells in cooperation with
ras. Biochem Biophys Res Commun 231, 509–513. doi:10.1006/bbrc.1997.6132
O'Leary, T., Williams, A.H., Caplan, J.S., Marder, E., 2013. Correlations in ion channel
expression emerge from homeostatic tuning rules. Proc Natl Acad Sci U S A 110, E2645–
54. doi:10.1073/pnas.1309966110
O'Leary, T., Williams, A.H., Franci, A., Marder, E., 2014. Cell types, network homeostasis,
and pathological compensation from a biologically plausible ion channel expression
model. Neuron 82, 809–821. doi:10.1016/j.neuron.2014.04.002
Osses, N., Henriquez, J.P., 2014. Bone morphogenetic protein signaling in vertebrate motor
neurons and neuromuscular communication. Front Cell Neurosci 8, 453.

doi:10.3389/fncel.2014.00453

Ou, Y., Chwalla, B., Landgraf, M., van Meyel, D.J., 2008. Identification of genes influencing
dendrite morphogenesis in developing peripheral sensory and central motor neurons.
Neural development 3, 16. doi:10.1186/1749-8104-3-16

Peled, E.S., Newman, Z.L., Isacoff, E.Y., 2014. Evoked and spontaneous transmission
favored by distinct sets of synapses. Curr Biol 24, 484–493.
doi:10.1016/j.cub.2014.01.022

Peng, T.-I., Jou, M.-J., 2010. Oxidative stress caused by mitochondrial calcium overload.
Ann N Y Acad Sci 1201, 183–188. doi:10.1111/j.1749-6632.2010.05634.x

Piccioli, Z.D., Littleton, J.T., 2014. Retrograde BMP Signaling Modulates Rapid Activity-
Dependent Synaptic Growth via Presynaptic LIM Kinase Regulation of Cofilin. J
Neurosci 34, 4371–4381. doi:10.1523/JNEUROSCI.4943-13.2014

Prinz, A.A., 2006. Insights from models of rhythmic motor systems. Curr Opin Neurobiol 16,
615–620. doi:10.1016/j.conb.2006.10.001

Prinz, A.A., Bucher, D., Marder, E., 2004. Similar network activity from disparate circuit
parameters. Nat Neurosci 7, 1345–1352. doi:10.1038/nn1352

Pulver, S.R., Pulver, S.R., Pashkovski, S.L., Pashkovski, S.L., Hornstein, N.J., Hornstein,
N.J., Garrity, P.A., Garrity, P.A., Griffith, L.C., Griffith, L.C., 2009. Temporal dynamics
of neuronal activation by Channelrhodopsin-2 and TRPA1 determine behavioral output
in Drosophila larvae. J Neurophysiol 101, 3075–3088. doi:10.1152/jn.00071.2009

Rharass, T., Lemcke, H., Lantow, M., Kuznetsov, S.A., Weiss, D.G., Panáková, D., 2014.
Ca2+-mediated mitochondrial reactive oxygen species metabolism augments Wnt/β-
catenin pathway activation to facilitate cell differentiation. J Biol Chem 289, 27937–
27951. doi:10.1074/jbc.M114.573519

Risse, B., Thomas, S., Otto, N., Löpmeier, T., Valkov, D., Jiang, X., Klämbt, C., 2013. FIM, a
Novel FTIR-Based Imaging Method for High Throughput Locomotion Analysis. PLoS
ONE 8, e53963. doi:10.1371/journal.pone.0053963

Roy, B., Singh, A.P., Shetty, C., Chaudhary, V., North, A., Landgraf, M., VijayRaghavan, K.,
Rodrigues, V., 2007. Metamorphosis of an identified serotonergic neuron in the
Drosophila olfactory system. Neural development 2, 20. doi:10.1186/1749-8104-2-20

Sanyal, S., Narayanan, R., Consoulas, C., Ramaswami, M., 2003. Evidence for cell
autonomous AP1 function in regulation of Drosophila motor-neuron plasticity. BMC
neuroscience 4, 20. doi:10.1186/1471-2202-4-20

Sanyal, S., Sandstrom, D.J., Hoeffer, C.A., Ramaswami, M., 2002. AP-1 functions upstream
of CREB to control synaptic plasticity in Drosophila. Nature 416, 870–874.
doi:10.1038/416870a

Schmitt, S., Evers, J.F., Duch, C., Scholz, M., Obermayer, K., 2004. New methods for the
computer-assisted 3-D reconstruction of neurons from confocal image stacks.
Neuroimage 23, 1283–1298. doi:10.1016/j.neuroimage.2004.06.047

Schneider-Mizell, C.M., Gerhard, S., Longair, M., Kazimiers, T., Li, F., Zwart, M.F.,
Champion, A., Midgley, F.M., Fetter, R.D., Saalfeld, S., Cardona, A., 2016. Quantitative
neuroanatomy for connectomics in Drosophila. Elife 5. doi:10.7554/eLife.12059

Schuster, C.M., Schuster, C.M., Davis, G.W., Davis, G.W., Fetter, R.D., Fetter, R.D.,
Goodman, C.S., Goodman, C.S., 1996. Genetic dissection of structural and functional
components of synaptic plasticity. I. Fasciclin II controls synaptic stabilization and
growth. Neuron 17, 641–654.

Serrano, F., Kolluri, N.S., Wientjes, F.B., Card, J.P., Klann, E., 2003. NADPH oxidase
immunoreactivity in the mouse brain. Brain Res 988, 193–198.

Shearin, H.K., Macdonald, I.S., Spector, L.P., Stowers, R.S., 2014. Hexameric GFP and
mCherry Reporters for the Drosophila GAL4, Q, and LexA Transcription Systems.
Genetics 196, 951–960. doi:10.1534/genetics.113.161141

Sigrist, S.J., Reiff, D.F., Thiel, P.R., Steinert, J.R., Schuster, C.M., 2003. Experience-
dependent strengthening of Drosophila neuromuscular junctions. J Neurosci 23, 6546–
6556.

Sugie, A., Hakeda-Suzuki, S., Suzuki, E., Silies, M., Shimozono, M., Möhl, C., Suzuki, T.,

Tavosanis, G., 2015. Molecular Remodeling of the Presynaptic Active Zone of Drosophila Photoreceptors via Activity-Dependent Feedback. Neuron 86, 711–725. doi:10.1016/j.neuron.2015.03.046

Sulkowski, M., Kim, Y.-J., Serpe, M., 2014. Postsynaptic glutamate receptors regulate local BMP signaling at the Drosophila neuromuscular junction. Development 141, 436–447. doi:10.1242/dev.097758

Tantama, M., Martínez-François, J.R., Mongeon, R., Yellen, G., 2013. Imaging energy status in live cells with a fluorescent biosensor of the intracellular ATP-to-ADP ratio. Nat Comms 4, 2550. doi:10.1038/ncomms3550

Tejada-Simon, M.V., Serrano, F., Villasana, L.E., Kanterewicz, B.I., Wu, G.Y., Quinn, M.T., Klann, E., 2005. Synaptic localization of a functional NADPH oxidase in the mouse hippocampus. Molecular and Cellular Neuroscience 29, 97–106. doi:10.1016/j.mcn.2005.01.007

Thiels, E., Urban, N.N., Gonzalez-Burgos, G.R., Kanterewicz, B.I., Barrionuevo, G., Chu, C.T., Oury, T.D., Klann, E., 2000. Impairment of long-term potentiation and associative memory in mice that overexpress extracellular superoxide dismutase. J Neurosci 20, 7631–7639.

Tripodi, M., Evers, J.F., Mauss, A., Bate, M., Landgraf, M., 2008. Structural homeostasis: compensatory adjustments of dendritic arbor geometry in response to variations of synaptic input. PLoS Biol 6, e260. doi:10.1371/journal.pbio.0060260

Turrigiano, G., 2012. Homeostatic synaptic plasticity: local and global mechanisms for stabilizing neuronal function. Cold Spring Harb Perspect Biol 4, a005736. doi:10.1101/cshperspect.a005736

Van Vactor, D., Sigrist, S.J., 2017. Presynaptic morphogenesis, active zone organization and structural plasticity in Drosophila. Curr Opin Neurobiol 43, 119–129. doi:10.1016/j.conb.2017.03.003

Vasin, A., Zueva, L., Torrez, C., Volfson, D., Littleton, J.T., Bykhovskaia, M., 2014. Synapsin regulates activity-dependent outgrowth of synaptic boutons at the Drosophila neuromuscular junction. J Neurosci 34, 10554–10563. doi:10.1523/JNEUROSCI.5074-13.2014

Vonhoff, F., Keshishian, H., 2017. Activity-Dependent Synaptic Refinement: New Insights from Drosophila. Front Syst Neurosci 11, 23. doi:10.3389/fnsys.2017.00023

Waak, J., Weber, S.S., Görner, K., Schall, C., Ichijo, H., Stehle, T., Kahle, P.J., 2009. Oxidizable residues mediating protein stability and cytoprotective interaction of DJ-1 with apoptosis signal-regulating kinase 1. J Biol Chem 284, 14245–14257. doi:10.1074/jbc.M806902200

Walker, J.A., Gouzi, J.Y., Long, J.B., Huang, S., Maher, R.C., Xia, H., Khalil, K., Ray, A., Van Vactor, D., Bernards, R., Bernards, A., 2013. Genetic and Functional Studies Implicate Synaptic Overgrowth and Ring Gland cAMP/PKA Signaling Defects in the Drosophila melanogaster Neurofibromatosis-1 Growth Deficiency. PLoS Genet 9, e1003958. doi:10.1371/journal.pgen.1003958

Wang, T., Hauswirth, A.G., Tong, A., Dickman, D.K., Davis, G.W., 2014. Endostatin Is a Trans-Synaptic Signal for Homeostatic Synaptic Plasticity. Neuron. doi:10.1016/j.neuron.2014.07.003

Wefelmeyer, W., Cattaert, D., Burrone, J., 2015. Activity-dependent mismatch between axo-axonic synapses and the axon initial segment controls neuronal output. Proc Natl Acad Sci U S A 112, 9757–9762. doi:10.1073/pnas.1502902112

Weyhersmüller, A., Hallermann, S., Wagner, N., Eilers, J., 2011. Rapid active zone remodeling during synaptic plasticity. J Neurosci 31, 6041–6052. doi:10.1523/JNEUROSCI.6698-10.2011

Younger, M.A., Müller, M., Tong, A., Pym, E.C., Davis, G.W., 2013. A presynaptic ENaC channel drives homeostatic plasticity. Neuron 79, 1183–1196. doi:10.1016/j.neuron.2013.06.048

Zhong, Y., Wu, C.-F., 2004. Neuronal Activity and Adenylyl Cyclase in Environment-Dependent Plasticity of Axonal Outgrowth in Drosophila. J Neurosci 24, 1439–1445.

doi:10.1523/JNEUROSCI.0740-02.2004

Zhu, X.-H., Qiao, H., Du, F., Xiong, Q., Liu, X., Zhang, X., Ugurbil, K., Chen, W., 2012.
Quantitative imaging of energy expenditure in human brain. Neuroimage 60, 2107–2117.
doi:10.1016/j.neuroimage.2012.02.013

Zito, K., Parnas, D., Fetter, R.D., Isacoff, E.Y., Goodman, C.S., 1999. Watching a synapse
grow: noninvasive confocal imaging of synaptic growth in Drosophila. Neuron 22, 719–
729.

Zuo, Y., Lin, A., Chang, P., Gan, W.-B., 2005. Development of long-term dendritic spine
stability in diverse regions of cerebral cortex. Neuron 46, 181–189.
doi:10.1016/j.neuron.2005.04.001

Zwart, M.F., Pulver, S.R., Truman, J.W., Fushiki, A., Fetter, R.D., Cardona, A., Landgraf, M.,
2016. Selective Inhibition Mediates the Sequential Recruitment of Motor Pools. Neuron
91, 944. doi:10.1016/j.neuron.2016.07.041

Zwart, M.F., Randlett, O., Evers, J.F., Landgraf, M., 2013. Dendritic growth gated by a
steroid hormone receptor underlies increases in activity in the developing Drosophila
locomotor system. Proc Natl Acad Sci U S A. doi:10.1073/pnas.1311711110

1071