High-risk human papillomavirus in oral cavity squamous cell carcinoma

Vinayak Palve[1], Jamir Bagwan[1], Neeraja M Krishnan[1], Manisha Pareek[1], Udita Chandola[1], Amritha Suresh[2], Gangotri Siddappa[2], Bonney L James[2], Vikram Kekatpure[3], Moni Abraham Kuriakose[2,3] and Binay Panda[1*]

[1]Ganit Labs, Bio-IT Centre, Institute of Bioinformatics and Applied Biotechnology, Bangalore, India

[2]Mazumdar Shaw Centre for Translational Cancer Research, Bangalore, India

[3]Mazumdar Shaw Medical Centre, Bangalore, India

*corresponding author:Binay Panda, Ganit Labs, Bio-IT Centre, Institute of Bioinformatics and Applied Biotechnology, Bangalore, India, Telephone: +91-9663311188, Email: binay@ganitlabs.in

Running Head: HPV in oral cavity tumors.

ABSTRACT

Purpose: The prevalence of human papillomavirus (HPV) in oral cavity squamous cell carcinoma (OSCC) varies significantly based on assay sensitivity and patient geography. Accurate detection is essential to understand the role of HPV in disease prognosis and management of patients with OSCC.

Methods: We generated and integrated data from multiple analytes (HPV DNA, HPV RNA, and p16), assays (immunohistochemistry, PCR, qPCR and digital PCR) and molecular changes (somatic mutations and DNA methylation) from 153 OSCC patients to correlate p16 expression, HPV DNA, and HPV RNA with HPV incidence and patient survival.

Results: High prevalence (33-58%) of HPV16/18 DNA did not correlate with the presence of transcriptionally active viral genomes (15%) in tumors. Eighteen percent of the tumors were p16 positive. and only 6% were both HPV DNA and RNA positive. Most tumors with relatively high-copy HPV DNA, and/or HPV RNA, but not with HPV DNA alone (irrespective of copy number), were wild-type for *TP53* and *CASP8* genes. In our study, p16 protein, HPV DNA and HPV RNA, either alone or in combinations, did not correlate with patient survival. Nine HPV-associated genes stratified the virus +ve from the –ve tumor group with high confidence ($p<0.008$) when HPV DNA copy number and/or HPV RNA were considered to define HPV positivity and not HPV DNA alone irrespective of their copy number ($p < 0.2$).

Conclusions: In OSCC, the presence of both HPV RNA and p16 are rare. HPV DNA alone is not an accurate measure of HPV positivity and therefore not informative. Moreover, HPV DNA, RNA or p16 don't correlate with outcome.

Keywords: Human papillomavirus (HPV), HPV DNA, HPV RNA, oral cavity tumors, copy number, survival, somatic mutation, DNA methylation

INTRODUCTION

Head and neck squamous cell carcinoma (HNSCC) is the sixth most common cancer worldwide with an incidence of 550,000 cases annually[1]. Oral cavity squamous cell carcinoma (OSCC) constitute a majority of HNSCC including tumors of the oral/anterior tongue and buccal mucosa[2]. The major known risk factors of OSCC are tobacco products, alcohol, and infection with human papillomavirus (HPV)[3]. Unlike oropharyngeal tumors, where HPV incidence with HPV DNA, RNA, and HPV-proteins is reported to be very high, (up to 90%)[4,5], the prevalence of HPV in OSCC varies widely (from none to 74%) depending on the choice of analyte, detection methodology and the geography of the patient cohort[6]. Additionally, unlike oropharyngeal tumors[7-11], the role of HPV in disease prognosis and response to therapy in patients with OSCC is equivocal. Despite the fact that HPV RNA is shown to function as better screening and patient management tool[12], the presence of HPV DNA is routinely used as a measure of HPV infection in tumors. However, there is a considerable variation in the sensitivity of DNA-based assays. Although widely used, HPV DNA results does not always match with HPV RNA, especially in OSCC. Therefore, in tumors, presence of HPV E6/E7 RNA and/or their protein products are considered as gold standard tests defining HPV positivity.

HPV16 and HPV18 subtypes have been epidemiologically linked with head and neck carcinoma[13]. High-risk HPV16 and HPV18 are the most predominant subtypes in oral cavity tumors of Indian patients while the other subtypes (HPV33, HPV6, and HPV11) are rare[14,15]. HPV E6 interacts with p53 to promote its degradation via ubiquitin pathway while E7 forms a complex with Rb leading to its functional inactivation and dysregulation of the cell cycle[16]. In some HPV-related tumors, E6- and E7-mediated inactivation of p53 and Rb result in the accumulation of p16 protein[17] while in others p16 expression does not directly correlate with HPV positivity[18]. Large-scale sequencing studies have demonstrated that the majority of HPV-negative tumors harbor mutations in *TP53* and *CASP8* and a significant proportion of HPV-positive tumors in *PIK3CA*[19-21]. Additionally, past studies have identified specific mutations in potential drug targets like *FGFR2/3* and lack of *EGFR* aberrations in HPV-positive patients[21] and the role of *CASP8* in HPV-negative cell line and patients[19,22]. Despite the wealth of information, accuracy of different HPV tests and

whether HPV plays an important factor in oral cavity tumor stratification and treatment remain to

be answered.

In the present study, we addressed the following questions on HPV in a cohort of 153 patients

with oral cavity tumors. Does sensitivity matter in the detection of HPV DNA in tumors? Does the

presence of tumor p16 and HPV DNA correlate with HPV E6/E7 RNA? Does the presence of high-

copy HPV DNA accurately reflect HPV positivity? Are p16 protein, HPV DNA, HPV E6/E7 RNA,

individually or together, linked with patient survival? And finally, do somatic mutations and DNA

methylation at 5-Cytosine residues distinguish the HPV +ve from the HPV –ve tumors?

METHODS

Patient cohort

Tumor samples from patients with OSCC (Buccal mucosa, BM, including from upper and

lower gingivobuccal sulcus and retromolar trigone and Oral tongue, OT) were accumulated

consecutively. Informed consent was obtained voluntarily from each patient enrolled in the study

and ethical was obtained from the Institutional Ethics Committees.All the tissues were frozen

immediately in liquid nitrogen and stored at -80^0C until further use. Only those tumors with

squamous cell carcinoma and at least 70% tumor cells and confirmed diagnosis were included in the

current study. Patients ($n = 153$) underwent staging according to AJCC criteria, and curative intent

treatment as per NCCN guidelines involving surgery with or without postoperative adjuvant

radiation or chemo-radiation at the Mazumdar Shaw Medical Centre were accrued for the study

(Table 1). Patients who received prior investigational therapy, CT, surgery or RT within four weeks

of initiating the investigation were excluded from the current study. Post-treatment surveillance was

carried out by clinical and radiographic examinations as per the NCCN guidelines.

Cell cultures

Cell lines (UM-SCC47, from Dr.Thomas Carey, University of Michigan, USA [23]; Hep2,

National Centre for Cell Science, Pune, India; UPCI:SCC29B and UPCI:SCC040, from Dr.

Susanne Gollin, University of Pittsburg, USA [24]) were maintained in Dulbecco's Modified Eagles'

Media (DMEM) supplemented with 10% FBS, 1X MEM non-essential amino acids solution and 1X

penicillin/streptomycin mixture (Gibco) and grown in incubators at 37^0C with 5% CO2.

Immunohistochemistry (IHC)

For p16 immunohistochemistry (IHC) staining was carried out using FFPE tissue blocks and

with primary antibody from BioGenex, USA (cat: AM540-5M; Anti-p16(INK4), Clone G175-405

in the NordiQC list) and using the poly-HRP detection system (Cat: QD400-60KE, BioGenex,

Fremont, CA, USA) following manufacturers' instructions. Sections of cervical cancer were used as

the positive control.The presence of p16 protein was scored based on the proportion of

immunopositive cells with both cytoplasmic and nuclear staining, irrespective of staining intensity

(weak/moderate/strong). The scoring pattern was: 0 - no immunopositive cells, 1 - <10%

immunopositive cells; 2 – 10-40% immunopositive cells and 3 - >40% immunopositive cells.

Tumors with a score of 3 were considered to be p16 positive.

DNA extraction

The genomic DNA from tumor tissues and cell lines were extracted using DNeasy Blood and

Tissue extraction kit using the spin column method with an intermediate RNase digestion following

manufacturer's instructions (Qiagen, USA). The DNA was quantified using Qubit 2.0 fluorometer

(Invitrogen, USA). Total 300ng of genomic DNA (unless specified otherwise) from tumors was

used for the detection of HPV using PCR, qPCR, and digital droplet PCR (ddPCR) methods.

Detection of HPV DNA with PCR

We tested five sets of primers published in the literature and two newly designed ones in the

amplification reactions (Figure 1). PCR primers were either consensus or type-specific regions of

the virus (Figure 1). Five sets of primers, including 4 HPV consensus (GP5-6; CP-I/II; MY09/11

and PGMY09/11) and one type-specific (HPV16-L1) primers as described in the literature [25], in the

PCR assays. Additionally, we designed two sets of new type-specific primers (HPV16-E6 and

HPV18-L1) to detect HPV DNA. We used genomic DNA from UMSCC-47 and Hep2 cell lines to

detect HPV16 and HPV18 respectively at various dilutions to determine relative efficiency,

sensitivity, and accuracy of amplification (Figure 2B). The primer sequences and the amplification

conditions used are provided in Supplementary Table 1.

Detection of HPV DNA with qPCR

For qPCR, E6 and E7 regions from HPV16 and HPV18 respectively were cloned in pUC19

vector, and Sanger sequencing confirmed their sequences. The qPCR was carried out using KAPA

Probe Fast qPCR master mix Universal (2X) (cat: KK4701, KAPA Biosystems, USA). The primers

and probes were designed (Supplementary Table 1) within the cloned regions of the HPV16, and

HPV18 plasmids and standard quantitative PCR was performed. The standard curves were

generated using serial dilutions (from 10 to 100,000 copies) of HPV16, and HPV18 cloned pUC19

plasmids. Genomic DNA from positive and negative control cell lines was used as positive and

negative controls for HPV18 and HPV16 respectively (Supplementary Figure 1). All amplification

reactions were carried out in triplicates, using nuclease-free water (cat: AM9932, Ambion, USA) as

a negative control. The analysis for each sample was performed by using the absolute quantification

using standard curve generated with serial dilutions of cloned plasmids. Tumors and cell lines

samples were counted as positive for those having Ct values three times away from the standard

deviation from the mean of negative controls.

Detection of HPV DNA using droplet digital PCR (ddPCR)

Digital PCR was performed using the QX100 ddPCR system (Bio-Rad, USA) using primers

and probes as provided in Supplementary Table 1. The reaction mix consisted of 10 µl of 2× ddPCR

Supermix without dUTP (cat: 1863024; Bio-Rad, USA), 450 nM of both forward and reverse

primers of HPV16 and 250 nM probe and 300ng of genomic DNA in a final volume of 20 µl. The

droplets were generated following manufacturer's standard instructions and were measured and

normalized in every reaction using the Quanta soft V.6 software (Bio-Rad, USA). All the samples

were processed in triplicates. Tumors and cell lines samples were counted as positive for those

having droplet numbers three times away from the standard deviation from the mean of negative

controls.

HPV DNA copy number

We deduced the HPV absolute copy number from the qPCR standard curves using cloned

HPV16/18 (Figure 3A). We considered a tumor or cell line to have relatively high-copy of HPV

DNA when the copy number for HPV16 and HPV18 DNA were more than 3.3×10^2 and 3.3×10^3

per microgram of tumor DNA respectively. To minimize the effect of tumor cellularity, ploidy, and

heterogeneity, we expressed the HPV copy number as copies per microgram of tumor DNA used in

the reaction.

Detection of HPV16/18 E6/ E7 RNA

The total RNA was extracted from 25mg of tumor tissues using the RNeasy mini kit from

Qiagen (cat: 74104) following manufacturer's instructions. The residual DNA was digested on the

column during extraction using RNase free DNase set from Qiagen (cat: 79254). RNA (500ng) was

subjected to cDNA synthesis using Takara's Prime Script first strand cDNA synthesis kit (cat:

6110A). The qPCR was carried out using KAPA SYBR Fast qPCR master mix Universal (cat:

KK4601, KAPA Biosystems, USA) with GAPDH as an internal control. The primer sequences and

the amplification conditions are provided in Supplementary Table 1. All amplification reactions

were carried out in triplicates, using nuclease-free water (cat: AM9932, Ambion, USA) as no

template control, cDNA from UPCI:SCC29B and UPCI:SCC40 cell lines as negative controls and

cDNA from cell lines UM-SCC47 and Hep2 as positive controls for E6 and E7 amplifications

respectively. Data were normalized using relative quantification method ($2^{\Delta\Delta Ct}$) [26]. Positive tumors

were counted as ones that have Ct values three times from the standard deviation of the mean of

negative control.

Mutation analysis by Sanger sequencing

The mutation data on tumors for *TP53*, *CASP8*, and *RASA1* were retrieved from the

previously published data [22].

Statistical Analysis

The Chi-square test was used to the significance of different clinical parameters of patients.

The relationship between tumor HPV status and survival in patients was examined by Kaplan-Meier

analysis with overall survival (OS) and disease-free survival (DFS) with log-rank test for

determining significance (*p* values <0.05) Disease-free survival was calculated as the time between

surgery and relapse/recurrence (as an event) of the disease or the last follow update in the case there

is no recurrence. The median follow up time of the patients is 15 months. The KM curves were

plotted using Graphpad Prism 5.0 software.

Whole genome methylation and statistical data analyses

Production of whole genome methylation data using Illumina Infinium Methylation450

BeadChip, chip scanning and the data pre-processing are described previously [27]. Three tumors in

HPV +ve group (relatively with high-copy HPV DNA and/or E6/E7 RNA) and 18 tumors in HPV -

ve group (HPV DNA –ve and HPV RNA –ve) were considered for further analysis. Differential

methylation for probes was estimated as $\Delta\beta$ values (Tumorβ - Normalβ) resulting in values ranging

from -1 (hypo-methylation) to 1 (hypermethylation). Using a supervised clustering approach,

differentially methylated probes that discriminated HPV +ve and HPV -ve patient's samples were

identified. Hyper-methylated probes in the HPV +ve group that were hypo-methylated in the HPV -

ve group, and *vice versa*, along with other probes with $\Delta\beta$ values that differed by at least 0.5

between the two groups, with a sample frequency of at least 60% were considered further. Finally,

the list of genes consisting of average $\Delta\beta$ values across tumors for multiple probes (where present)

was drawn. Neighbourhood interaction network for these genes was inferred using PCViz from

PathwayCommons (http://www.pathwaycommons.org/pcviz/#neighborhood/). The genes and their

interacting partners were mapped to the Viral Carcinogenesis pathway from KEGG (hsa05203), and

mapped genes were linked back to respective genes found using supervised clustering. For the

linked genes, an unpaired t-test was performed to determine the significance of the difference

between the HPV +ve and HPV -ve tumor groups. The same was performed for the HPV-associated

genes between the DNA-based (irrespective of the DNA copy number) HPV +ve and HPV -ve

tumor groups.

RESULTS

p16 expression

For the p16 immunohistochemical study, the tumors ($n = 55$) were first confirmed by H and E

staining of the paraffin tissue sections followed by the immunohistochemical staining with the

surrogate marker p16. Figure 2A shows the representative p16 +ve and –ve staining in oral cavity

tumor sections (OSCC) along with the cervical tissue section (+ve control). As indicated

previously, we considered cells with cytoplasmic + nuclear staining for p16 to be counted as overall

p16 positive. In our study, 18% of the tumors were p16 positive (Table 2).

Incidence of HPV DNA

Detection of HPV using PCR:

The efficacy of HPV16/18 DNA detection using various consensus or type-specific primers is

shown in Figure 2B. The primers described in the literature (GP5+6+, MY09/11, CPI-II or

PGMY09/11, HPV16L1) could only detect HPV DNA when 0.03ng or higher amount of genomic

DNA from the cell line UMSCC-47/Hep2 was used (Figure 2B). However, the newly designed

type-specific primers (HPV16E6 and HPV18L1) were able to detect HPV DNA with as low as

0.0003ng and 0.003ng genomic DNA for HPV16 and HPV18 respectively (Figure 2B). To check

the efficiency of the newly designed primers with absolute copy numbers of HPV genome, we used

serial diluted HPV cloned plasmids DNA. As shown in Supplementary Figure 2, efficient

amplification with 100 copies of HPV genome or higher could be detected. Once we optimized the

conditions and primers for all the amplification reactions, we performed PCR in oral cavity tumors

($n = 70$). In our cohort, 58% (41/70) of the tumors were HPV DNA +ve for any of the PCR (Table

2, Supplementary Figure 3). Figure 2C shows the efficiency of the consensus and type-specific

primers in a set of representative oral cavity tumors. Widely used primers from the literature

(MY09/11, PGMY09/11, GP5+6+ and HPV16L1) yielded either least or moderate (CPI-II)

sensitivity of detection while the newly designed HPV16E6 and HPV18L1 primers showed the

optimum sensitivity of detection (Figure 2B). We observed inhibition of amplification reactions at a

high concentration of tumor genomic DNA with positive cell line spike-in experiment

(Supplementary Figure 4) and therefore avoided in the reactions. Additionally, an increase in

amplification cycles also did not aid in the detection of HPV DNA in PCR as shown in

Supplementary Figure 5.

Detection of HPV using real-time PCR:

Next, we used quantitative PCR using Taqman chemistry to detect HPV DNA. We performed

serial dilution using HPV16 E6, and HPV18 L1 cloned plasmids to obtain standard curves (Figure 3

I, IV). We used an increasing amount of genomic DNA from negative control cell lines to

demonstrate specific amplification of HPV DNA (Supplementary Figure 1). Genomic DNA from

HPV +ve and –ve cell lines were used to plot the baseline. The real-time PCR was performed on

tumors of the oral cavity (*n*=106) and were counted as HPV DNA positive for those having Ct

values three times away from the standard deviation from the mean of negative controls (Figure 3

II, V). Results from qPCR indicated that 33% (35/106) of tumors were positive for HPV DNA

(Table 2). Although we found high incidence for HPV16 (30%; 32/106) than HPV18 (18%; 19/106)

type, the HPV18 positive tumors had very high copy numbers of viral DNA as reflected in their Ct

values (Figure 3A III, VI). The prevalence of HPV DNA in oral tongue tumors was higher (44%)

compared to that of buccal mucosa (4%).

Detection of HPV using droplet digital PCR:

Finally, we used one of the most sensitive methods, digital PCR, to detect HPV DNA. Digital

PCR was recently shown to successfully to detect HPV DNA in oropharyngeal tumors in highly

specific manner [30]. We performed droplet digital PCR (ddPCR) with serially diluted plasmid

HPV16E6 clones to generate the standard curve (Figure 3B, I). The genomic DNA from positive-

(UM-SCC47) and negative control cell lines (Hep2) for HPV16 along with a no template control

(NTC) were used to plot the baseline (Figure 3B, II; Supplementary Figure 6). The digital PCR

performed on the oral cavity tumors results indicated that 43% (59/136) of oral cavity tumors were

positive for HPV16 DNA (Figure 3B, III and Table 2). Moreover, HPV16 infection was more

prevalent in the oral tongue tumors (55%) compared to the buccal mucosal ones (17%) (Table 2).

HPV E6 and E7 RNA expression

The viral mRNA, E6, and E7 (for both HPV16 and HPV18) were measured by qPCR in oral

cavity tumors and from cell lines (UMSCC-47 and Hep2). Compared to the cell lines, tumors

showed very low levels of expression of E6 and E7 mRNA (Figure 3C). Unlike HPV DNA, only

15% of the tumors showed expression of E6 and/or E7 RNA and 6% of the tumors confirmed the

presence of both HPV DNA (with all the three assays) and transcriptionally active HPV genomes

(Table 2). Similar to DNA, oral tongue tumors had a higher number (17%) of HPV E6/E7 RNA

compared to the buccal mucosa (9%, Table 2). In our cohort, younger patients (≤40yrs age) were

significantly more HPV RNA +ve than the older patients when analyzed by Chi-square analysis (p

= 0.029). When combining the results from all the assays (p16 IHC, HPV DNA, and HPV RNA),

we found 6-48% of the tumors were positive in various assay combinations with PCR (Table 2).

We found that 22% (23/106) of the tumors had relatively high-copy of HPV DNA and/or HPV

E6/E7 mRNA.

Linking tumor attributes, somatic mutations, and HPV with survival

We performed Kaplan Meir survival analysis with various tumor attributes that revealed

significant association between tumor differentiation (p=0.03) and clinical stage (p=0.0003) with

overall survival (Figure 4A)**.** None of the other tumor attributes showed significant association with

survival (Supplementary Figure 7 A-F). In patients with oral cavity tumors, p16, HPV DNA and

HPV RNA did not correlate with the either overall or disease-free survival of (Figure 4A,

Supplementary Figure 8). HPV DNA status alone measured by any of the DNA-based assays alone

or in combination did not correlate with survival (Supplementary Figure 8), except when measured

with ddPCR for overall survival (Supplementary Figure 8E, p=0.03). We tested whether tumors

with relatively high HPV DNA copy and/or HPV E6/E7 mRNA, is linked with survival. As shown

in Supplementary Figure 9 A,B, we did not find any significant association with this group of

tumors with both overall- ($p = 0.45$) and disease-free ($p = 0.68$) survival.

Further, we investigated whether somatic mutations in significantly mutated genes in OSCC

have a role to play in HPV DNA +ve tumors and patient survival. We looked at three genes (*TP53*,

CASP8, and *RASA1*) shown to be significantly mutated in oral cavity tumors [19, 22, 31]. Ninety-five

percent of the HPV +ve tumors in the group were wild-type for *TP53* and *CASP1* genes and 85% of

the HPV +ve tumors for *RASA1* gene. (Supplementary Figure 10). We tested whether the mutation

in any of the genes, alone or in combination in HPV –ve tumor group is linked with survival. As

shown in Supplementary Figure 9 C,D, we did not find any significant association with this group

of tumors with survival.

Linking methylation to HPV

Supervised clustering of the first group of patients (a group defined as one with high-copy

number HPV DNA and/or E6/E7 RNA) resulted in a list of 60 genes, out of which 9 (*FERMT3*,

GIT2, *HK3*, *PRKCZ*, *ZCCHC8*, *IRF5*, *IFFO1*, *ARID3A*, *HOXA2*) were mapped to the HPV pathway

(Figure 4B). Methylation of those genes is involved in the downstream control of the expression of

different target genes. For example, *ZCCHC8* methylation is linked with the expression of RB1,

PRKCZ methylation controls-state-change-of *DLG1*, methylation in *ARID3A*, *IRF5*, *IFFO1*, and

HOXA2 are connected with the expression of *TP53*, and *FERMT3*, *HK3*, and *GIT2* genes control

the expression of *AP1* (*JUN*) (Figure 4B). All the genes, except for *HOXA2*, were significantly

hypermethylated in the HPV +ve group of tumors compared to the HPV –ve group (Figure 4B).

The linked four genes, obtained from the nine significantly methylated genes, were mapped to the

pathways involving HPV E6 and E7 proteins (Figure 4C). To test significance, we performed

unpaired t-tests between the two groups, group 1 that has relatively high-copy HPV DNA and/or

HPV RNA and group 2 that were negative for both HPV DNA and HPV RNA. All of the eight

hypermethylated genes showed very significance (p < 0.00001) and for *HOXA2* that was

hypomethylated the *p*-value was 0.007 (Supplementary Table 2). However, when the patients were

grouped based on HPV DNA positivity alone (irrespective of their copy number), most of these 9

HPV-linked genes did not show significant association.

DISCUSSION

HPV plays a vital role in the prognosis of patients with oropharyngeal tumors [32, 33]. Unlike

oropharynx, HPV incidence and its role in disease prognosis in oral cavity tumors are not well

established. Past results on HPV DNA incidence in oral cavity tumors were widely variable (from

very low to very high) depending on the assay sensitivity, analyte and patient cohort used [34]. We

performed literature survery and found that the rate of incidence, the role of HPV in disease

outcome varies significantly in different groups (Supplementary Table 3). The accuracy of the HPV

tests employed is, and HPV positivity needs to be answered to make treatment decisions in patients

with head and neck tumors confidently [35]. There are fewer studies that used multiple analytes

(protein, DNA, and RNA) and various molecular tests (IHC, PCR, qPCR and digital PCR) to

establish HPV positivity in oral cavity tumors and that correlated HPV with tumor attributes

(including somatic mutations and methylation) and survival. In the present study, we attempted to

do this in 153 patients with oral cavity tumors.

While p16 expression, as measured by immunohistochemistry (IHC), is a commonly used

proxy for HPV in HNSCC, its expression is not specific in HPV associated tumors [33]. Although a

number of past studies correlated p16 expression with HPV [36-38], p16 IHC has shortcomings,

especially when relating its expression with patient survival. Limitations, such as variation in

staining intensities [39, 40], nonspecific binding of antibodies and the lack of scoring and interpretive

criteria used for p16 staining [41] make the method less reliable. In our study, although we found an

unusually high number (51%) of tumor cells showing immunopositive staining, only a small

percentage (18%) had both cytoplasmic+nuclear staining, an accurate reflection of HPV positivity

as described earlier [28, 29]. Unlike some of the previous studies [24, 42], we could not find any correlation

(either positive or negative) between p16 expression and survival (Figure 4). Like previous reports

[36, 43], we found p16 expression not to be a useful surrogate marker for HPV in oral cavity tumors. A

weak correlation between p16 IHC and HPV *in situ* hybridization (ISH) is reported in the past [44].

Unlike antibody-based methods, nucleic acids-based methods detect HPV with high

sensitivity and therefore used widely [45]. Meta-analysis of 5478 oral cavity tumors suggested the

overall HPV DNA prevalence to be 24.2% with 11% tumors being positive for both HPV DNA and

E6/E7 RNA [46]. India has one of the highest incidence rates of oral cavity cancers with a significant

difference in the trend of incidence between oropharyngeal and oral cavity cancer [47]. Previously,

PCR coupled with Mass Array is shown to provide high sensitivity of detection with a low amount

of input genomic DNA [48]. From our results, we found 38% tumors to be positive, and 13% were

negative in all three DNA-based assays (PCR, qPCR, and ddPCR) respectively. Overall, the

prevalence (33-58%) of HPV DNA was dependent on the type of test used with PCR yielding the

highest incidence over the more sensitive methods like qPCR and ddPCR assays (Table 2). This

was possibly due to the use of consensus primers in PCR but not in qPCR and ddPCR, in addition

to the type-specific ones, resulting in the detection of non-HPV16/18 subtypes. As expected, digital

PCR, being the most-sensitive method out of the three DNA-based assays, resulted in more number

of tumors being HPV16-DNA positive resulting in the detection of very low copy viral genomes in

tumor samples. Based on several levels of evidence as noted below, we conclude that the presence

of low copy HPV DNA alone may not be a reflection of functionally active HPV. First, in the

tumors positive for HPV DNA, we found only a fraction (15%) with HPV E6/E7 RNA. Second,

only 6% of tumors were positive for both the presence of HPV genome and E6/E7 RNA. Third,

almost all the tumors with relatively high-copy HPV genome and/or HPV RNA had wild-type *TP53*

and *CASP8* genes, which was not the case with tumors with the low-copy HPV DNA. Both *TP53*

and *CASP8* are known to be wild-type primarily in HPV positive tumors 19, and in our study, we

also found this corresponds to the tumors with high-copy HPV genome and/or transcriptionally

active genome only (Table 2). We believe that the high prevalence of HPV DNA in the tumor

tissues might have been a result of highly sensitive assays used in our study and might suggest

either the presence of passenger HPV genomes coming from adjacent normal cells, as earlier

reported [48, 49] and/or might have been a reflection of inactive or passenger virus in oral cavity

tumors. Although very few (*n*=3), we cannot explain why some tumors in our study with HPV

E6/E7 RNA did not show the presence of HPV DNA. It is possible that the genomic DNA for those

tumors were degraded and therefore, could not serve as ideal templates for DNA-based assays. An

additional factor could be the presence of inhibitors for DNA-based assays in those tumors.

Although HPV RNA is considered to be gold standard analyte to test for HPV in tumors, RNA is a

difficult analyte to handle and is more labile than DNA. Additionally, studying RNA in archival

samples may pose an additional set of challenges.

The fact that there were only two tumors, which were p16 positive and HPV RNA negative, a

definitive conclusion on the lack of correlation between p16 and HPV RNA can't be made from our

study. Similarly, there were two tumors, which were positive for HPV RNA and negative for p16.

In HNSCC, p16 is often mutated/silenced resulting in its loss of expression. This could have led to

the lack of p16 expression in those two tumors. We did not find any significant correlation between

p16, HPV-DNA and/or HPV RNA and disease outcome (Figure 4A and Supplementary Figure 8).

Even the tumors with relatively high-copy HPV genomes and/or E6/E7 RNA did not support the

role of HPV in patient survival (Supplementary Figure 9 A, B). These aspects need further studies

and analysis.

There are several limitations of our study. Not all the tumors were assayed with all the

analytes, making the sample number different for different methods. We could not perform

additional survival analyses in the tumors that were HPV RNA positive given the small sample size.

In our study, we did not perform HPV ISH, which could have provided additional information on

p16 positivity and HPV prevalence. It is possible that HPV has a different mechanism of action in

oral cavity tumors. Indirect support to linking high HPV genome copy number with HPV positivity

in our study comes from the fact that we did not find significance in HPV-linked genes stratifying

HPV +ve from HPV –ve group when HPV DNA irrespective of their copy number is taken into

account in defining the HPV positivity (Figure 4B). It is possible that the presence of high-copy

HPV genome in those tumors does not correlate with the presence of the biologically active virus.

Further studies may help answer this question.

ACKNOWLEDGEMENTS

Research presented in this article is funded by Department of Electronics and Information

Technology, Government of India (Ref No: 18(4)/2010-E-Infra., 31-03-2010) and Department of

IT, BT and ST, Government of Karnataka, India (Ref No: 3451-00-090-2-22). We thank Mr.

Osama Mansour of Bio-Rad, India in helping us with ddPCR.

CONFLICT OF INTEREST

None of the authors declare any conflicts of interest.

AUTHOR CONTRIBUTIONS

Study design: Vinayak Palve and Binay Panda

Data analysis: Vinayak Palve, Neeraja M Krishnan and Binay Panda

Manuscript writing: Vinayak Palve and Binay Panda

Conducting experiments: Vinayak Palve, Jamir Bagwan, Udita Chandola, Manisha Pareek, Amritha

Suresh, Gangotri Siddappa and Bonne L James

Collection of patient samples: Amritha Suresh, Vikram Kekatpure and Moni A Kuriakose

Overall study supervision: Binay Panda

ABBREVIATIONS

HPV, human papillomavirus; HNSCC, head and neck squamous cell carcinoma; OSCC, oral cavity

squamous cell carcinoma; IHC, immunohistochemistry; PCR, polymerase chain reaction; qPCR,

quantitative polymerase chain reaction; ddPCR, droplet digital polymerase chain reaction

REFERENCES
438

439	1.	Ferlay J, Shin HR, Bray F, Forman D, Mathers C, Parkin DM. Estimates of worldwide
440		burden of cancer in 2008: GLOBOCAN 2008. *International journal of cancer Journal*
441		*international du cancer*. 2010;127:2893-2917.
442	2.	Bhat SP, Bhat VS, Permi H, Shetty JK, Aroor R. Oral and oropharyngeal malignancy: A
443		clinicopathological study. *Internet Journal of Pathology and Laboratory Medicine*. 2015;1.
444	3.	Mishra A, Meherotra R. Head and neck cancer: global burden and regional trends in India.
445		*Asian Pac J Cancer Prev*. 2014;15:537-550.
446	4.	Nasman A, Attner P, Hammarstedt L, et al. Incidence of human papillomavirus (HPV)
447		positive tonsillar carcinoma in Stockholm, Sweden: an epidemic of viral-induced
448		carcinoma? *International journal of cancer Journal international du cancer*. 2009;125:362-
449		366.
450	5.	Walline HM, Komarck C, McHugh JB, et al. High-risk human papillomavirus detection in
451		oropharyngeal, nasopharyngeal, and oral cavity cancers: comparison of multiple methods.
452		*JAMA Otolaryngology–Head & Neck Surgery*. 2013;139:1320-1327.
453	6.	Isayeva T, Li Y, Maswahu D, Brandwein-Gensler M. Human papillomavirus in non-
454		oropharyngeal head and neck cancers: a systematic literature review. *Head Neck Pathol*.
455		2012;6 Suppl 1:S104-120.
456	7.	Ang KK, Harris J, Wheeler R, et al. Human papillomavirus and survival of patients with
457		oropharyngeal cancer. *New England Journal of Medicine*. 2010;363:24-35.
458	8.	Fakhry C, Westra WH, Li S, et al. Improved survival of patients with human
459		papillomavirus-positive head and neck squamous cell carcinoma in a prospective clinical
460		trial. *Journal of the National Cancer Institute*. 2008;100:261-269.
461	9.	Fakhry C, Zhang Q, Nguyen-Tan PF, et al. Human papillomavirus and overall survival after
462		progression of oropharyngeal squamous cell carcinoma. *J Clin Oncol*. 2014;32:3365-3373.
463	10.	Lassen P, Eriksen JG, Hamilton-Dutoit S, Tramm T, Alsner J, Overgaard J. Effect of HPV-
464		associated p16INK4A expression on response to radiotherapy and survival in squamous cell
465		carcinoma of the head and neck. *J Clin Oncol*. 2009;27:1992-1998.
466	11.	Rischin D, Young RJ, Fisher R, et al. Prognostic significance of p16INK4A and human
467		papillomavirus in patients with oropharyngeal cancer treated on TROG 02.02 phase III trial.
468		*J Clin Oncol*. 2010;28:4142-4148.
469	12.	Cattani P, Siddu A, D'Onghia S, et al. RNA (E6 and E7) assays versus DNA (E6 and E7)
470		assays for risk evaluation for women infected with human papillomavirus. *J Clin Microbiol*.
471		2009;47:2136-2141.
472	13.	Huang C-G, Lee L-A, Tsao K-C, et al. Human papillomavirus 16/18 E7 viral loads predict
473		distant metastasis in oral cavity squamous cell carcinoma. *Journal of Clinical Virology*.
474		2014;61:230-236.
475	14.	Balaram P, Nalinakumari KR, Abraham E, et al. Human papillomaviruses in 91 oral cancers
476		from Indian betel quid chewers--high prevalence and multiplicity of infections.
477		*International journal of cancer Journal international du cancer*. 1995;61:450-454.
478	15.	D'Costa J, Saranath D, Dedhia P, Sanghvi V, Mehta AR. Detection of HPV-16 genome in
479		human oral cancers and potentially malignant lesions from India. *Oral oncology*.
480		1998;34:413-420.
481	16.	zur Hausen H. Papillomaviruses and cancer: from basic studies to clinical application. *Nat*
482		*Rev Cancer*. 2002;2:342-350.
483	17.	Lu DW, El-Mofty SK, Wang HL. Expression of p16, Rb, and p53 proteins in squamous cell
484		carcinomas of the anorectal region harboring human papillomavirus DNA. *Mod Pathol*.
485		2003;16:692-699.
486	18.	Wang H, Sun R, Lin H, Hu WH. P16INK4A as a surrogate biomarker for human
487		papillomavirus-associated oropharyngeal carcinoma: consideration of some aspects. *Cancer*
488		*Sci*. 2013;104:1553-1559.
489	19.	Cancer Genome Atlas N. Comprehensive genomic characterization of head and neck
490		squamous cell carcinomas. *Nature*. 2015;517:576-582.
491	20.	Lechner M, Frampton GM, Fenton T, et al. Targeted next-generation sequencing of head
492		and neck squamous cell carcinoma identifies novel genetic alterations in HPV+ and HPV-
493		tumors. *Genome Med*. 2013;5:49.

21. Seiwert TY, Zuo Z, Keck MK, et al. Integrative and comparative genomic analysis of HPV-
positive and HPV-negative head and neck squamous cell carcinomas. *Clin Cancer Res*.
2015;21:632-641.
22. Krishnan N, Gupta S, Palve V, et al. Integrated analysis of oral tongue squamous cell
carcinoma identifies key variants and pathways linked to risk habits, HPV, clinical
parameters and tumor recurrence. *F1000Res*. 2015;4:1215.
23. Brenner JC, Graham MP, Kumar B, et al. Genotyping of 73 UM-SCC head and neck
squamous cell carcinoma cell lines. *Head & neck*. 2010;32:417-426.
24. Auw-Haedrich C, Martin G, Spelsberg H, et al. Expression of p16 in conjunctival
intraepithelial neoplasia does not correlate with HPV-infection. *The open ophthalmology
journal*. 2008;2:48.
25. Romero-Pastrana F. Detection and typing of human papilloma virus by multiplex PCR with
type-specific primers. *ISRN microbiology*. 2012;2012.
26. Schmittgen TD, Livak KJ. Analyzing real-time PCR data by the comparative CT method.
Nature protocols. 2008;3:1101-1108.
27. Krishnan NM, Dhas K, Nair J, et al. A Minimal DNA Methylation Signature in Oral Tongue
Squamous Cell Carcinoma Links Altered Methylation with Tumor Attributes. *Mol Cancer
Res*. 2016;14:805-819.
28. Chen ZW, Weinreb I, Kamel-Reid S, Perez-Ordonez B. Equivocal p16 immunostaining in
squamous cell carcinoma of the head and neck: staining patterns are suggestive of HPV
status. *Head Neck Pathol*. 2012;6:422-429.
29. Lewis JS, Jr., Chernock RD, Ma XJ, et al. Partial p16 staining in oropharyngeal squamous
cell carcinoma: extent and pattern correlate with human papillomavirus RNA status. *Mod
Pathol*. 2012;25:1212-1220.
30. Biron VL, Kostiuk M, Isaac A, et al. Detection of human papillomavirus type 16 in
oropharyngeal squamous cell carcinoma using droplet digital polymerase chain reaction.
Cancer. 2016;122:1544-1551.
31. India Project Team of the International Cancer Genome C. Mutational landscape of gingivo-
buccal oral squamous cell carcinoma reveals new recurrently-mutated genes and molecular
subgroups. *Nature communications*. 2013;4:2873.
32. Maxwell JH, Grandis JR, Ferris RL. HPV-Associated Head and Neck Cancer: Unique
Features of Epidemiology and Clinical Management. *Annu Rev Med*. 2016;67:91-101.
33. Vokes EE, Agrawal N, Seiwert TY. HPV-Associated Head and Neck Cancer. *Journal of the
National Cancer Institute*. 2015;107:djv344.
34. Bruni L, Barrionuevo-Rosas L, Serrano B, et al. Human papillomavirus and related diseases
report. *L'Hospitalet de Llobregat: ICO Information Centre on HPV and Cancer*. 2014.
35. Seiwert T. Accurate HPV testing: a requirement for precision medicine for head and neck
cancer. *Annals of oncology : official journal of the European Society for Medical Oncology /
ESMO*. 2013;24:2711-2713.
36. Chung CH, Zhang Q, Kong CS, et al. p16 protein expression and human papillomavirus
status as prognostic biomarkers of nonoropharyngeal head and neck squamous cell
carcinoma. *Journal of Clinical Oncology*. 2014:JCO. 2013.2054. 5228.
37. Stephen JK, Divine G, Chen KM, Chitale D, Havard S, Worsham MJ. Significance of p16
in site-specific HPV positive and HPV negative head and neck squamous cell carcinoma.
Cancer and clinical oncology. 2013;2:51.
38. Gröbe A, Hanken H, Kluwe L, et al. Immunohistochemical analysis of p16 expression, HPV
infection and its prognostic utility in oral squamous cell carcinoma. *Journal of Oral
Pathology & Medicine*. 2013;42:676-681.
39. Larsen CG, Gyldenløve M, Jensen D, et al. Correlation between human papillomavirus and
p16 overexpression in oropharyngeal tumours: a systematic review. *British journal of
cancer*. 2014;110:1587-1594.
40. Seiwert TY. Ties That Bind: p16 As a Prognostic Biomarker and the Need for High-
Accuracy Human Papillomavirus Testing. *Journal of Clinical Oncology*. 2014:JCO.
2014.2057. 9268.

41. El-Naggar AK, Westra WH. p16 expression as a surrogate marker for HPV-related oropharyngeal carcinoma: A guide for interpretative relevance and consistency. *Head & neck*. 2012;34:459-461.

42. Alexander RE, Hu Y, Kum JB, et al. p16 expression is not associated with human papillomavirus in urinary bladder squamous cell carcinoma. *Modern Pathology*. 2012;25:1526-1533.

43. Lingen MW, Xiao W, Schmitt A, et al. Low etiologic fraction for high-risk human papillomavirus in oral cavity squamous cell carcinomas. *Oral oncology*. 2013;49:1-8.

44. Chung CH, Zhang Q, Kong CS, et al. p16 protein expression and human papillomavirus status as prognostic biomarkers of nonoropharyngeal head and neck squamous cell carcinoma. *J Clin Oncol*. 2014;32:3930-3938.

45. Gibson JS. Nucleic acid-based assays for the detection of high-risk human papillomavirus: A technical review. *Cancer cytopathology*. 2014;122:639-645.

46. Ndiaye C, Mena M, Alemany L, et al. HPV DNA, E6/E7 mRNA, and p16INK4a detection in head and neck cancers: a systematic review and meta-analysis. *Lancet Oncol*. 2014;15:1319-1331.

47. Chaturvedi AK, Anderson WF, Lortet-Tieulent J, et al. Worldwide trends in incidence rates for oral cavity and oropharyngeal cancers. *J Clin Oncol*. 2013;31:4550-4559.

48. Terai M, Hashimoto K, Yoda K, Sata T. High prevalence of human papillomaviruses in the normal oral cavity of adults. *Oral Microbiol Immunol*. 1999;14:201-205.

49. Gillison ML, Broutian T, Pickard RK, et al. Prevalence of oral HPV infection in the United States, 2009-2010. *JAMA*. 2012;307:693-703.

605 **Table 1:** Patient details and characteristics of tumor tissues used in the study.
606

Characteristics		Number
Primary site		
	Buccal mucosa	41
	Oral tongue	112
Gender		
	Male	114
	Female	39
Age		
	>40	112
	<=40	40
	NA	1
Risk habits		
	Alcohol	4
	Chewing	43
	Smoking	6
	Alcohol+ Chewing	14
	Smoking + Alcohol	16
	Smoking + Chewing	9
	Smoking + Alcohol+ Chewing	10
	No habits	44
	NA	7
Tumor classification/stage		
	I+II	43
	III+IV	109
	NA	1
Differentiation		
	Well	48
	Moderate	73
	Poor	20
	NA	12
Abbreviations: NA = Not Available		

Table 2. Summary of HPV assays in oral cavity tumors. p16 was measured by the presence of

immuopositive cells with both nulcear and cytoplasmic staining in IHC, PCR results indicate

presence of any HPV subtype with consensus primers or HPV16/18 type specific primers, qPCR

and ddPCR results indicate Taqman assay results with primers and probes for HPV 16/18 and

HPV16 respectively and HPV RNA results indicate presence of E6 and/or E7 mRNA for

HPV16/HPV18.

614
615

Detection	Method	% Positivity		
		Oral tongue	Buccal mucosa	Together (Oral cavity)
p16 IHC	p16	18% (10/55)	NA	18% (10/55)
DNA based	PCR	59% (39/66)	50% (2/4)	58% (41/70)
	qPCR	44% (34/78)	3.6% (1/28)	33% (35/106)
	ddPCR	55% (52/95)	17% (7/41)	43% (59/136)
RNA based	qPCR	17% (5/30)	9% (1/11)	15% (6/41)
Combinations	PCR + qPCR	38% (23/60)		
	PCR + ddPCR	48.3%(29/60)		
	qPCR + ddPCR	27.2% (27/99)		
	PCR +qPCR + ddPCR	37.7% (20/53)		
	p16 + 3/3 Methods	5.5% (2/36)		
	RNA + DNA	6% (1/17)		

Figure legends

Figure 1: HPV genome organization and locations of different primer/probes used in the study to

detect HPV DNA (either consensus or type specific) and RNA. The numbers represent the

corresponding nucleotide number in the HPV genome.

Figure 2: p16 and HPV DNA in OSCC. Representative images of immunohistochemical staining

of p16 in positive, negative OSCC tissue sections and cervical tissue as positive control (A), the

relative PCR amplification efficiency and sensitivityof consensus /type specific primers for

detection of HPV using HPV16 or HPV18 individual positive control cell lines (UMSCC-47 and

Hep2) (B) and representative HPV DNA PCR in oral cavity tumors with both consensus and type-

specific primers (C).

Figure 3: Detection of HPV DNA and RNA. HPV16/18 assays using qPCR (A I-VI) and digital

PCR (B I-III) in OSCCS is shown. Standard curves were obtained using cloned HPV16/18 plasmids

(A-I,IV, B -I) and data was subsequently obtained using the positive (UM-SCC-47 and Hep2) and

negative (UPCI:SCC29B & UPCI:SCC40) cell line DNA (A-II,V, B-II) to count HPV DNA in oral

cavity tumors (A-III,VI, B-III). Figure 3C indicates HPV16 (top panel) and HPV18 (bottom panel)

E6/E7 mRNA expression in tumors using qPCR..

Figure 4: Kaplan Meier survival plots. Survival plots linking tumors with various attributes like

grade, stage, p16 IHC, HPV DNA and HPV RNA (A) is shown. Figure 4B and 4C depict clustering

of nine methylated genes stratifying HPV positive from negative group of tumors (B) and HPV-

associated pathways (C) in HPV positive tumors.

Figure 1

Figure 2

Figure 3

Figure 4