p53 dynamically directs TFIID assembly on target gene promoters

Coleman, R. A.[1#], Qiao, Z.[1], Singh, S. K.[1], Peng, C. S.[1], Cianfrocco, M.[2], Zhang, Z.[3], Piasecka, A.[1], Aldeborgh, H.[1], Basishvili, G.[1], and Liu, W. L.[1]

[1]Gruss-Lipper Biophotonics Center, Department of Anatomy and Structural Biology, Albert Einstein College of Medicine, Bronx, NY 10461, USA.

[2]Department of Cellular and Molecular Medicine, University of California, San Diego, La Jolla, CA 92093, USA.

[3]Surrozen, 240 East Grand Avenue, 2[nd] Floor, South San Francisco, CA 94080, USA.

#To whom correspondence should be addressed. Department of Anatomy and Structural Biology, Albert Einstein College of Medicine, Bronx, NY 10461 Tel: 718-430-8623; Email: robert.coleman2@einstein.yu.edu

Abstract: 200 words, 1,195 characters
Introduction+Results+Discussion: 5,481 words, 30,987 characters
Figure Legends: 1029 words, 5,460 characters
Materials and Methods: 2,231 words, 11,857 characters

Running title: Binding dynamics of p53 and TFIID on DNA

Keywords: p53; TFIID; transcription; structure; single molecule microscopy

Abstract

The p53 tumor suppressor protein is a central regulator that turns on vast gene networks to maintain cellular integrity upon various stimuli. p53 activates transcription initiation in part by aiding recruitment of TFIID to the promoter. However, the precise means by which p53 dynamically interacts with TFIID to facilitate assembly on target gene promoters remains elusive. To address this key question, we have undertaken an integrated approach involving single molecule fluorescence microscopy, single particle cryo-electron microscopy, and biochemistry. Our real-time single molecule imaging demonstrates that TFIID alone binds poorly to native p53 target promoters. p53 unlocks TFIID's ability to bind DNA by increasing TFIID contacts with both the core promoter and a region surrounding p53's response element (RE). Analysis of single molecule dissociation kinetics reveals that TFIID interacts with promoters via transient and prolonged DNA binding modes that are each regulated by p53. Importantly, our structural work reveals that TFIID's conversion from a canonical form to a rearranged DNA-binding conformation is enhanced in the presence of DNA and p53. Notably, TFIID's interaction with DNA induces p53 to rapidly dissociate, effectively liberating the RE on the promoter. Collectively, these findings indicate that p53 dynamically escorts and loads the basal transcription machinery onto its target promoters.

66 **Introduction**
67
More than 50% of cancer patients harbor p53 mutations, highlighting the essential role of

this protein in tumor suppression (1). To properly maintain genomic stability, p53 induces vast

gene networks involved in diverse cellular pathways including cell cycle arrest, apoptosis and

DNA repair (2). In response to various stress stimuli, p53 acts as a transcriptional activator that

specifically binds consensus response elements (REs; two ten base-pair half sites,

RRRCWWGYYY) within its target genes to directly stimulate gene expression (2). p53 utilizes

its ability to non-specifically bind and slide along the DNA to expedite the search for target REs.

(3). Upon recognition of its response genes, p53 facilitates transcription at least in part by

targeting the TFIID-mediated transcription machinery to the promoter (4-7). TFIID is composed

of TBP and 14 TBP-associated factors (TAFs). TFIID recognizes and binds multiple core

promoter elements (e.g. the TATA box, Initiator, and downstream core promoter elements

[DPE]) surrounding the transcription start site (TSS) (8). Once bound, TFIID serves as a central

scaffold for six basal factors, including RNA Polymerase II, to form a pre-initiation complex

(PIC) directing transcription initation. Importantly, without the assistance of activators such as

p53, TFIID's core promoter recognition is weak and rate limiting for transcription initiation due

to inefficient PIC assembly (9-16). As TFIID is responsible for transcription of at least ~90% of

protein-coding genes (17), unraveling how p53 intervenes in this central step of the PIC

assembly (i.e. TFIID's binding of promoter DNA) is essential for understanding mechanisms

controlling gene-specific transcription initiation throughout the cell.

Previous biochemical studies of different activators (e.g. p53 and Zta) suggested a

conserved albeit poorly characterized mechanism for facilitating transcription (10, 18, 19). It has

been shown that p53 binds to its REs and enhances TFIID promoter recruitment via direct

contact with TBP and select TAFs (19, 20). However, p53 target genes have diverse spatial

arrangements of core promoter elements and p53 REs. Each target promoter also exhibits varied

affinities for p53 and TFIID (1). Therefore, it is unclear how distinct combinations of p53 REs

and core promoter elements affect the dynamics and the global architecture of TFIID assemblies

on target genes. Additional evidence has reported that p53 and other activators introduce

structural changes within TFIID when associated with TFIIA and DNA (10, 18, 19). This

activator-induced isomerization leads to enhanced TFIID binding to core promoter elements

downstream of the TSS (10, 18, 19). However, the structural mechanism of p53-mediated

isomerization of TFIID and its impact on PIC assembly is currently unknown.

Single molecule-based imaging has emerged as an advanced tool to discern complex

dynamic behavior of large multi-subunit protein assemblies involved in regulating gene

expression. A recent study utilizing real-time single molecule fluorescence microscopy provided

insight that TFIID's interaction with a synthetic core promoter DNA tightly correlates with

productive transcription (21). Another single molecule report uncovered TFIIB's dynamic

binding to TFIID at the promoter (22). Additional live cell studies also suggest that the PIC

composition is extraordinarily dynamic with the loading and promoter escape of RNA

Polymerase II occurring every ~ 4 to 9 seconds during bursts of transcription (23). How

activators such as p53 regulate TFIID's dynamic binding to native target DNA remains elusive.

Advanced single particle transmission electron microscopy (EM) has unraveled the three-

dimensional (3D) structures of TFIID and its co-complexes. Recent EM studies unmasked a

structural plasticity of TFIID, likely attributed to its promoter recognition activity (20, 24-27). In

particular, human TFIID features a "horseshoe" shape containing three major lobes (A, B & C)

and a well-defined central cavity (28, 29). Interestingly, we demonstrated that, in the absence of

DNA, three different activators, including p53, introduce a common set of local structural

changes in TFIID, such as movement of lobe A towards lobe B (20). Intriguingly, human TFIID

undergoes a significant structural rearrangement involving movement of Lobe A when bound to

a specific promoter DNA optimized for TFIID recognition (26, 27). It remains to be determined

if p53's facilitated movement of Lobe A results in a rearranged conformation of TFIID that

contributes to stable interaction between TFIID and DNA.

To uncover the mechanistic underpinnings of how p53 facilitates TFIID assembly on

promoter DNA, we exploited a combination of single molecule fluorescence microscopy,

biochemistry, and single particle cryo-EM analysis. These studies illuminate dynamic

interactions and static global molecular architectures of p53-induced TFIID assemblies on two

representative target genes (i.e. *hdm2* and *bax*). Our real-time single molecule studies show that

TFIID alone infrequently assembles onto these native p53 target promoters. TFIID/DNA binding

primarily occurs on the order of seconds. p53 facilitates TFIID's promoter recruitment by

enhancing multiple TFIID contacts throughout the core promoter and RE. Notably, p53 also

increases the percentage of TFIID molecules displaying long-lived DNA binding. Consistent

with our single molecule studies, EM structural analysis reveals that conversion of canonical

TFIID to a rearranged DNA binding conformation is enhanced in the presence of p53 and DNA.

TFIID/DNA binding induces p53 dissociation, effectively "liberating" the RE to allow p53-

mediated recruitment of additional factors involved in PIC formation. Therefore, proper response

to divergent stress signals involves conserved structural mechanisms relating to p53's ability to

dynamically stimulate TFIID's interaction with various target promoters.

Results

TFIID association with native target genes in the presence of p53

We sought to further understand the molecular mechanism pertaining to activator-

mediated enhancement of TFIID's ability to bind DNA. Thus, dynamic interactions between

p53, TFIID and promoter DNA were characterized via a single molecule co-localization assay

using Total Internal Reflection Fluorescence Microscopy (TIRF) (Figure 1A)(21, 22). Over the

first 15' of binding, Quantum dot-labeled TFIID (QD-TFIID) associated with only 1.5% of the

Cy3/Cy5-labeled native wild-type *hdm2* promoter DNA in the absence of p53 (Figure 1B and

Supplemental Figure S1, left panel), revealing a low intrinsic affinity of TFIID for DNA.

Increasing the time between addition of TFIID to the DNA and the imaging of binding did not

improve co-localization efficiencies, suggesting that reactions are at equilibrium (data not

shown). This finding indicates that, at low concentrations (~1nM), TFIID inefficiently

recognizes DNA. On the other hand, increased binding of TFIID to DNA was observed in our

bulk biochemical assembly assays when much higher concentrations (100-600nM) were used as

in Supplemental Figure S6B.

Upon addition of p53, a clear enhancement of TFIID's association with the *bax* or *hdm2*

promoter DNA was observed (Figure 1B). This result is consistent with a general increase of

DNA retained by TFIID in the presence of p53 in our bulk biochemical assays (Supplemental

Figure S6B). A control experiment detected no p53-mediated co-localization of DNA with the

Quantum dot-labeled TAF1 antibody ($0.2 \pm 0.07\%$ co-localization) in the absence of TFIID (data

not shown). Therefore p53's significant enhancement TFIID/DNA association is likely due to the

direct interaction between p53 and TFIID.

To directly address the role of the p53 RE and core promoter elements in p53-mediated

TFIID/DNA binding, three mutant *hdm2* DNA templates were analyzed with the same real-time

co-localization assay (Figure 1B). The first mutant *hdm2* DNA template (i.e. Mu RE) harbors

mutations in the response elements known to impair p53's sequence-specific interactions (30). A

mild but significant decrease of TFIID's association with DNA was observed on the mutant p53

RE template (Figure 1B). This result suggests that enhancement of TFIID/DNA binding can

occur via a mechanism that is independent of p53/RE contacts.

To assess the impact of the core promoter in p53-mediated TFIID/DNA binding,

mutations (i.e. Mutant Core) known to weaken TFIID binding and inhibit transcription (22, 31)

were exploited. Upon addition of p53, TFIID's association with the mutant core template was

further reduced compared to the wild-type promoter (Figure 1B). As expected, when both the

p53 RE and core promoter elements were altered (i.e. Mu RE/Co), TFIID/DNA co-localization

was further decreased (Figure 1B). These results indicate that the p53 RE and core promoter

elements were both required for optimal association of TFIID with p53 target gene promoters.

p53 regulates TFIID/DNA contacts within the core promoter and p53 response elements

As an alternative approach to study p53-mediated TFIID/DNA binding, we developed an

in vitro formaldehyde-crosslinking/exonuclease mapping assay (xlink-Exo). Our strategy was

adapted from a well-established strategy (ChIP-exo) to map out protein/DNA contacts of

transcription factors in vivo at near base-pair resolution (32). Based on this analysis, TFIID

makes extensive contacts with well-characterized core promoter elements (i.e. TATA, InR, MTE

and DPE) on the wild-type *hdm2* DNA (Figure 1C, top panels). Addition of p53 led to elevated

crosslinking of TFIID to the wild-type core promoter with select elements showing much larger

changes (i.e. TATA, InR) than others (i.e. MTE/DPE) (top panels). Consistent with our single

molecule data, p53 increases the specific binding of TFIID to the core promoter. Importantly,

there was a clear reduction in p53's ability to enhance contacts within mutated core promoter

elements (Figure 1C, bottom panels and Supplemental Figure S2A). This finding is consistent

with our single molecule assay showing reduced TFIID/DNA co-localization on the mutant core

promoter (Figure 1B). Interestingly, p53 slightly increases TFIID's contacts within the -4 and

+11 regions on these mutant core promoters (Figure 1C, bottom panel and Supplementary Figure

S2A). Thus our mutantions don't completely eliminate TFIID/core promoter contacts and may

explain why a small percentage of TFIID is still bound to the mutant promoters in our single

molecule assays (Figure 1B).

In addition to these well-established contacts, our xlink-Exo assays also mapped

TFIID/DNA interactions far upstream of TATA (-55 to -137) and downstream of the DPE (~

+37 to +68) (Figure 1C, top panel). This result is consistent with previous DNAse I footprinting

studies reporting TFIID/DNA interactions upstream of TATA (-40 to -140) and downstream of

the DPE (+32 to +68) at similar positions on different promoters (26, 33, 34). Many of these

upstream and downstream TFIID contacts were enhanced when p53 was co-present on the wild-

type *hdm2* promoter (Figure 1C). Moreover, some of TFIID's upstream contacts (i.e. -55 to -88)

reside within the p53 RE on the wild-type promoter even in the absence of p53 (Figure 1C, top

panels). Mutations of the p53 RE led to a loss of TFIID's contacts at -61 and -88, suggesting that

TFIID recognizes specific sequences within the p53 RE (Figure 1C, bottom panels and

Supplemental Figure S2B). Furthermore, a number of upstream TFIID contacts (-61, -79, and -

88/-90) were also abrogated on the mutant core promoter (Supplementary Figure S2). This

suggests that these novel upstream DNA contacts arise from TFIID's interaction with the TATA,

InR and DPE sequences. Therefore, TFIID may utilize additional uncharacterized elements to

recognize promoters.

p53 modulates the prolonged association of TFIID with DNA

In addition to p53's ability to stimulate TFIID/DNA association, we sought to test if p53

could also stabilize TFIID's binding to DNA. Therefore, the amount of time that QD-TFIID

remains bound to a single promoter DNA before dissociating (i.e. residence time) with and

without p53 was examined (Figure 2A). In the absence of p53, two populations of DNA-bound

TFIID complexes displaying distinct residence times were detected on the wild-type *hdm2*

template (Figure 2B). The majority (78%) of TFIID/DNA binding events were short-lived

(average residence time of 6.6 ± 0.4 seconds) in the absence of p53 (Figure 2B, left panel).

Notably, p53 significantly increases the percentage of long-lived TFIID/DNA bound complexes

from 22% to 64% (Figure 2B, left versus right panel). Furthermore, p53 prolongs the residence

time of both the short and long-lived TFIID/DNA binding events (Figure 2C). TFIID also

displayed a similar stability profile on the *bax* promoter in the presence of p53 (Supplemental

Figure S3). These results suggest that TFIID interacts with DNA via two intrinsic binding modes

that can be modulated by p53.

We next defined how the RE and core promoter elements might regulate the relative

proportion of these two TFIID/DNA binding populations in the presence of p53. Disruption of

the p53 binding site on the *hdm2* promoter (Mu RE) had little effect on the distribution between

these two populations (Figure 2D). This indicates that a p53-mediated increase in the percentage

of TFIID molecules displaying long-lived DNA binding does not solely require contacts with the

RE. In contrast, mutation of the core promoter region (Mu Core) reduced the percentage of

TFIID molecules in the long-lived DNA binding population (Figure 2D). Alteration of the p53

binding site and the core promoter elements (Mu RE/Co) further lowered the percentage of the

TFIID's long-lived DNA binding population (Figure 2D). These data suggest that the TFIID's

long-lived DNA binding mode was related primarily to recognition of the core promoter with

p53/RE interactions providing additional stability post-recruitment to the DNA.

Our results also reveal that p53 RE and core promoter mutations also impacted TFIID's

residence time on DNA. Mutation of the p53 RE (Mu RE) predominantly decreased the

residence time of TFIID's long-lived DNA binding without significantly affecting the short-lived

events (Supplementary Figure S4B). Therefore, interactions between p53 and its RE are required

for maximal stability of TFIID/DNA contacts on native target promoters. On the mutant core

promoter (Mu Core) residence times for both the long- and short-lived TFIID/DNA binding

events were also significantly reduced (Supplementary Figure S4). Thus, both DNA binding

modes were related to core promoter recognition. Mutation of both the p53 RE and core

promoter were changed (Mu RE/Co) had little effect of TFIID's residence time on DNA

(Supplementary Figure S4B). These results suggest that mutation of both the RE and core

promoter elements primarily effects TFIID's distribution between short and long-lived DNA

binding modes (Figure 2D). Together with the single molecule data, our xlink-Exo studies

suggest that TFIID's remaining minimal contacts on our mutant promoter are sufficient to anchor

short-lived TFIID binding to DNA.

p53/target promoter interactions in the presence of TFIID

Our previous report documented that a TFIID variant could cooperatively increase the

binding of the activator c-Jun to its binding site on promoter DNA (24). To quantitatively assess

if this was the case for p53, the dissociation dynamics of fluorescently labeled-p53/DNA

interactions with or without TFIID were examined (Figure 3). In the absence of TFIID, p53

primarily displayed two populations of distinct binding events on the wild-type *hdm2* template

(i.e. long-lived [avg = 33.1 ± 3.9"] and short-lived [avg = 7.4 ± 0.7"], Figure 3A, left panel).

Since earlier reports showed that p53 can bind DNA non-specifically (35, 36), we hypothesized

that these short-lived p53/DNA binding events were non-specific interactions. To test our

hypothesis, the same experiments were performed using the mutant p53 RE template (right

panel). Indeed, disruption of p53's sequence-specific binding sties on DNA led to a single short-

lived population (avg = 6.8 ± 0.7"). Based upon this result, the long-lived p53/DNA binding

events most likely represent sequence-specific interactions (left panel), whereas the short-lived

p53/DNA binding events are non-specific in nature. Alternatively, these short-lived events could

represent interactions with an unknown lower affinity p53 binding site in the promoter or binding

of dimeric p53 to concensus half-sites within the mutant RE. It is of note that photobleaching of

the fluorescently labeled p53 did not affect our results, since similar residence times were

observed with multiple dyes of differing photostability (data not shown). In addition, a previous

study detected similar short residence times of ~10-12" for p53 bound to a consensus site (37).

Collectively, these findings indicate that p53 dynamically interacts with DNA via both specific

and non-specific interactions.

Importantly, addition of TFIID resulted in a single population of p53 bound to wild-type

DNA with an "intermediate-lived" residence time distinct from the two populations observed

with p53 alone (Figure 3B). Furthermore, we noticed that fitting of the data showed slight

deviations from an exponential decay model. A previous study indicated that the existence of

complex multi-step dissociation events could be discerned using a gamma distribution model to

fit single molecule data (38). Indeed, we found that a gamma distribution model better represents

the data (Figure 3B, right panel). This analysis shows that in the presence of TFIID, p53

dissociates from the DNA in ~2 steps. Each step contains roughly equivalent residence times of

8.2 seconds per step (average total time = 14.8 ± 1.5"). Moreover, single step dissociation

kinetics was observed for p53 alone based upon gamma distribution fitting (data not shown).

Therefore, TFIID alters both the rate and number of kinetic steps involved in p53's dissociation

from promoter DNA.

Mutation of the p53 RE further reduces p53's residence time on DNA in the presence of

TFIID (Supplemental Figure S5A, left panel versus Figure 3B, right panel). Disruption of

TFIID's interaction with the TATA, InR and DPE elements on the mutant p53 RE/Co DNA

template does not lead to a further decrease in p53's residence time (Supplemental Figure S5A

versus S5B). Fitting of the dissociation data on the mutant templates to a gamma distribution

yielded poor fits, suggesting single step dissociation kinetics for p53 on these mutant promoters

(Supplemental Figures S5A and S5B, right panels).

Overall, these data suggest that TFIID binding to DNA can influence p53's stability at

the RE. TFIID's interaction with p53 likley promotes the dissociation of p53 bound to specific

sequences in the promoter (Figure 3C, left panel). Importantly, the average time of p53 bound to

the wild-type *hdm2* template (~15") is significantly shorter than TFIID's long-lived DNA

binding time (~100") (Figure 3C, right panel). These dynamic studies strongly suggest that p53

transiently escorts TFIID to the promoter DNA. Importantly, p53 is not required to remain bound

for TFIID's maximum stability on promoter DNA.

The conformational status of TFIID when co-present with p53 and promoter DNA

A recent cryo-EM study discovered that human TFIID switches from a canonical to a

rearranged conformation upon binding to the Super Core Promoter (SCP), a DNA fragment

optimized for interaction with TFIID (26, 31). Thus far our single molecule data indicates that

p53 stimulates binding of TFIID to DNA. Therefore, we hypothesized that TFIID could form a

rearranged DNA binding conformation with p53 and promoter DNA. Therefore, we set out to

determine TFIID's global structural architecture when p53 and target promoter DNA were

present. As the first step, unbiased reference-free 2D classification experiments were performed

to examine conformational states of TFIID in the presence of p53 or p53/*bax* DNA (Figure 4A).

The movement of Lobe A from Lobe C to Lobe B is the hallmark of the rearranged DNA

binding conformation of TFIID (26). Therefore, three conformational states of TFIID

represented by corresponding views of 2D class averages were defined based upon the position

of Lobe A relative to Lobes C and B (top panels, i.e. canonical, transition, and rearranged state).

We found that addition of p53 led to an elevated percentage of particles within the corresponding

view displaying the rearranged TFIID form compared to TFIID alone (Figure 4A, bottom panel).

This finding implies that the formation of TFIID's rearranged state is affected by p53. This result

is consistent with conformational changes observed in our previous negative stained structure of

a TFIID/p53 co-complex (20). Furthermore, the 2D classification analysis showed that the

populations of both the transition and rearranged forms increased when p53 and DNA was added

(bottom panel). Therefore, these findings suggest that TFIID's conformational switches can be

regulated via target gene promoters and/or its interaction with p53.

A previous biochemical study reported that p53 stimulated the assembly of TFIIA and

TFIID on a synthetic promoter DNA (19). Interestingly, unlike p53, TFIIA alone promotes the

canonical form of TFIID (26). Thus, to assess the structural framework of TFIID in the presence

of p53 and TFIIA, two TFIID/p53/TFIIA assemblies on the native *hdm2* and *bax* promoter

fragments (488 bp and 500 bp) were obtained (Supplemental Figure S6). To visualize both

conformations and the DNA, cryo-EM data of these assemblies were obtained for reference-free

2D classification analysis. Some class averages showing the presence of promoter DNA

spanning the central cavity of TFIID were observed (Figure 4B). Moreover, both assemblies

displayed a similar rearranged TFIID architecture, despite the different organization of their

promoters. Each template harbors non-consensus core promoter elements with different spatial

arrangements between TATA and the Initiator (Figure 1B, top panel) (8, 39, 40). Combined with

our 2D classification analysis (Figure 4A), these structural studies indicate that TFIID binds

DNA in a common rearranged conformation in the presence of p53 and TFIIA.

Next, to identify the relative DNA orientation and any distinct structural features, we

compared the class averages of our two assemblies with the previous structural analysis of

TFIID/TFIIA/SCP DNA containing a mapped DNA orientation (26). Our co-complexes

displayed clear views of TFIID stably bound to the *bax* DNA downstream of the TSS (Figure

4C, top two rows). In contrast, upstream promoter DNA was weakly visible in our averages

(middle panel). These results suggest that the p53 RE upstream of the TATA box element is

flexible within the co-complex. To define promoter DNA upstream of TATA, focused

classification on a class average displaying flexible upstream DNA (middle panel) was

performed as previously described (26). This analysis revealed a clear density for the upstream

promoter DNA (highlighted by yellow arrow, bottom panel). Given the strong density for

downstream DNA within the assembly, these results imply that TFIID serves as a strong anchor

for downstream promoter DNA during PIC assembly.

The 2D class averages of TFIID/DNA/p53/TFIIA assemblies exhibit a similar topology

to the averages from TFIID/TFIIA bound to the SCP DNA (refer to Figure 3 in (26)) raising two

major points. First, TFIID adapts a common DNA binding conformation to bind various p53

target gene promoters. Second, structural rearrangements within TFIID related to DNA binding

may cause p53 to dissociate, as supported by our single molecule dynamics results (Figure 3). To

test these hypotheses, single particle 3D reconstruction was conducted to determine the 3D

structure of the TFIID/TFIIA/p53/*bax* DNA co-complex (Supplemental Figure S7A). The overall

structural architecture of our co-complex is comparable to the 3D structure of TFIID/TFIIA/SCP

DNA. Our 3D reconstruction was also verified by analyzing its 3D projections with their

matching reference-free 2D averages (Supplemental Figure S7B). The same observation was also

obtained with the 3D structure of the TFIID/TFIIA/p53/*hdm2* promoter DNA (Supplemental

Figure S8). These results suggest that TFIID utilizes one common rearranged form to bind

various promoters.

Our 3D reconstruction did not reveal any prominent extra densities that would most

likely represent the p53 protein. This indeed suggests that p53 dissociates from the

TFIID/TFIIA/*bax* DNA co-complex. Alternatively, p53 could have a transient interaction with

the assembly, which led to the EM density being averaged out during analysis. Since we

previously determined the 3D structure of the p53 bound TFIID co-complex without DNA (20),

we attempted to capture the p53-bound state of TFIID in the cryo-EM dataset of our

TFIID/TFIIA/p53/*bax* DNA assembly. Thus, 3D reconstructions were performed using the

structure of p53/TFIID as an initial reference volume. A poor resolution 3D reconstruction of

p53/TFIID was obtained, indicating a small subset of the single cryo-EM particles (Supplemental

Figure S9). This observation suggests that the majority of TFIID complexes have switched to

the p53-free, DNA-bound state when p53 and DNA were co-present. Collectively, these

structural studies support the observations from our single molecule analyses, illustrating that

p53 dissociates from TFIID and DNA when recruited to target promoters. Our work also implies

that when p53 stimulates TFIID binding to DNA, a common rearranged DNA binding scaffold is

formed on various target genes.

Discussion

TFIID infrequently interacts with native p53 target genes in the absence of additional factors

Our single molecule assays reveal that TFIID poorly associates with native p53 target

promoters in the absence of p53 (Figure 1B). TFIID's inefficient DNA binding activity may be

one way to prevent spurious basal expression of these stress response genes. TFIID/promoter

interactions are likely repressed in part due to the TAF1 N-terminal domain (TAND) which

occupies TBP's concave DNA binding surface (41-44). This TAF1/TBP interaction is also

thought to control the TFIID's global conformation to regulate promoter interactions. TFIID's

canonical conformation is non-permissive to DNA binding (26, 27) (Figure 5 top panel, left

side). Researchers postulate that the TAF1 TAND acts to tether TBP to TFIID's lobe C in the

canonical form (27). Large-scale structural rearrangement of TFIID dramtically shifts TBP's

positon within TFIID to allow simultaneous contact of TBP with the upstream TATA and TAF1

with downstream core promoter elements (Figure 5 top panel, right side). Thus TAF1/TBP

interactions repress the large-scale structural rearrangement within TFIID required for TBP to

bind the TATA element in a core promoter.

TFIID's infrequent association with p53 target gene promoters occurs via distinct short

and long-lived DNA binding modes (Figures 2B, 2C and Supplementary Figure S4B). The

majority (78%) of these TFIID/DNA binding events last just a few seconds (~7") (Figures 1B,

2B and 2D). This short-lived DNA binding population may represent a transient non-productive

TFIID encounter complex in the canonical form where TAF1 prevents stable TBP/TATA

interactions (Figure 5). In such a scenario, TAF contacts with the initiator and DPE may be

sufficient to temporarily anchor TFIID to DNA. Due to this transient dynamic interaction

behavior, we were likely unable to obtain the 3D structure of those short-lived DNA-bound

TFIID complexes via cryo-EM.

DNA binding and structural features of TFIID are regulated by p53

A number of factors, including activators (i.e. c-Jun and VP16) and TFIIA, can stimulate

TFIID/promoter binding by de-repressing the TAF1-mediated inhibition of TBP (19, 45-48).

TFIIA has also been found to aid conversion of canonical TFIID to the rearranged DNA binding

form likely via disruption of TBP/TAF1 TAND interactions (26, 27). It was previously unclear if

activators could also promote structural rearrangement of TFIID to stimulate promoter

association. Our past work has shown that p53 binds stably to TFIID and targets TAF1 in the

absence of DNA (20). We also found that p53's interaction with TFIID also results in movement

of Lobe A towards Lobe B in a manner analogous to a structurally rearranged DNA binding form

of TFIID (20). Therefore we postulated that DNA binding may be increased in this structurally

rearranged pre-assembled p53/TFIID complex. Indeed our single molecule data along with

previous bulk biochemical experiments indicate that p53 can dramatically stimulate the binding

of TFIID to DNA (Figure 1B and (19)). We also find that conversion of canonical TFIID to the

structurally rearranged form is enhanced in the presence of p53 and DNA (Figure 4A). Thus we

speculate that p53, whose acidic activation domain is highly homologous to VP16 (49), may

potentially disrupt TAF1/TBP interactions to aid in TFIID structural rearrangement and DNA

binding (Figure 5, bottom panel).

p53 promotes isomerization of TFIID to stimulate long-lived binding to DNA

Cryo-EM of a TFIID/TFIIA/DNA complex has revealed a number of TBP/TAF contacts

with core promoter elements spanning from ~ -33 to +32bp surrounding the transcription start

site (27). Many of these contacts, especially TBP/TATA, are only possible due to TFIID's

structural rearrangement or isomerization upon DNA binding. Therefore we anticipate that the

p53-mediated binding of TFIID to DNA is also likely a multi-step isomerization process

involving contacts with multiple core promoter elements (Figure 5, bottom panel). Indeed our x-

link Exo analysis revealed numerous TFIID crosslinks throughout the promoter that are

enhanced by p53 (Figure 1C). Many of these TFIID/*hdm2* DNA contacts were strictly conserved

in the TFIID/TFIIA/super core promoter DNA cryo-EM structure (27). Consistently, the

architectures of TFIID bound to the *hdm2* and supercore promoters are equivalent (Supplemental

Figures S7 and S8). This suggests that TFIID in the presence of p53 can specifically bind our

native *hdm2* promoter to adopt a common DNA binding conformation.

In the presence of p53, TFIID associates with DNA robustly. Analysis of single molecule

dissociation kinetics also revealed two TFIID/core promoter DNA binding modes with

interactions predominantly being long-lived (~100") (Figures 2B and 2D). Therefore, p53 shifts

the TFIID population from short to long-lived DNA binding, likely acting to facilitate TFIID's

known isomerization on promoters (Figure 5, bottom panel). This isomerization step may

possibly permit stable interaction of TFIID with the upstream DNA, TATA box and downstream

DNA resulting in long-lived DNA binding (Figure 5, bottom panel). We speculate that the p53-

mediated long-lived DNA-bound TFIID complex is structurally related to the stable

TFIID/TFIIA/promoter DNA complex depicted in our cryo-EM studies. In support of this model,

previous biochemical assays have shown that activators induce conformational changes within

TFIID that enhance multiple contacts with promoter sequences downstream of the TSS (10).

Mutation of the p53 RE only affected the residence time of TFIID's long-lived DNA

binding population (Supplementary Figure S3). Importantly, the percentage of long-lived DNA

binding events was essentially unchanged on the mutant RE template (Figure 2D). This suggests

that isomerization of TFIID does not depend on p53 contacts with the RE. Given that distinct

TFIID contacts surrounding the p53 RE were identified via our xlink-Exo assays (Figure 1C), we

speculate that this upstream DNA is involved in both binding p53 and stabilizing interaction of

an isomerized form of TFIID with the promoter. These upstream contacts may be possibly

mediated by the TAF4 subunit of TFIID, which was previously shown to bind ~70 bp of DNA

with a weak sequence preference (50).

TFIID isomerization upon DNA binding liberates p53 from the RE

Our single molecule data indicate that TFIID reduces p53's residence time on DNA

(Figures 3A and 3B). This result is consistent with previous DNAse I footprinting studies

showing decreased binding of p53 to REs in the presence of TFIID (19). More importantly,

p53's residence time on DNA is 6.7 fold shorter (~15") than TFIID's (~100") (Figure 3C, right

panel). Therefore p53 disengages the promoter before TFIID during long-lived DNA binding

events. In our EM structures, we are also unable to detect density associated with p53 in the

rearranged DNA bound TFIID further confirming our single molecule findings (Supplemental

Figures S7, S8 and S9). Previous structural analysis shows that p53 bridges lobes A and C of

TFIID in the absence of DNA (20). Thus, we speculate lobe A's movement away from lobe C

upon DNA binding disrupts p53/TFIID contacts inducing p53's dissociation from the assembly

(Figure 5, bottom panel).

While p53 dissociates from the TFIID/DNA scaffold, it is highly likely that p53 will

rapidly re-associate with the complex, particularly at the high concentrations (600nM) used in

Supplementary Figure S6. We suspect that p53's rapid removal could consequently lead to a

faster "repurposing" of the RE for additional rounds of p53-mediated recruitment of other basal

transcription factors such as TFIIB (51, 52), RNA Pol II (53), and TFIIH (49, 54, 55). In this

scenario, stable prolonged binding of p53 to a TFIID/DNA complex would negatively regulate

PIC formation, because p53 would occupy the RE and inhibit subsequent recruitment of a

p53/basal factor co-complex.

TFIID/promoter binding dynamics in relation to transcriptional burst kinetics

Once p53 loads TFIID onto our native p53 target gene promoters, TFIID remains bound

for only approximately 100 seconds likely due to interactions with suboptimal or non-consensus

core promoter elements (Figure 2). Notably, our residence times for TFIID at the promoter were

measured in the absence of additional PIC components. Most likely, additional factors such as

TFIIA, TFIIF, and RNA Polymerase II will alter the stability of TFIID and p53 as the PIC forms

on promoter DNA. However, recent in vivo work suggests that transcription complex formation

is highly dynamic. Consistent with our data, yeast TBP occupies the HSC82 heat shock promoter

for 60 seconds in vivo (56). In human tissue culture cells, RNA Polymerase II loads onto

promoters every ~4 to 9 seconds to form convoys on an active gene (23). These bursts of RNA

Polymerase II convoys last for approximately 100 seconds before the promoter becomes inactive

again (23). Therefore it is tempting to speculate that the long-lived TFIID residency (~100

seconds) on our native target promoters could be related to RNA Polymerase II convoy

formation (~100 seconds) and transcriptional bursting. TFIID's dynamic binding to promoters

could be a mechanism to allow rapid shut down of stress response gene expression after re-

establishment of cellular homeostasis.

TFIID can also direct infrequent loading of RNA Polymerase II convoys on mutant

TATA containing promoters leading to bursts of transcription (23). Interestingly, the duration of

RNA Polymerase II convoys on the mutant TATA promoter is only slightly less than that on a

wild-type TATA template (~60-80 seconds) (23). Our study also revealed infrequent p53

mediated binding of TFIID on our mutant core promoter DNA lasting for a significantly long

time (~64 seconds) (Figure 2D and Supplementary Figure S3). Therefore we suspect that TFIID

stably loaded onto our mutant core DNA is fully competent to direct low levels of infrequent

transcription. Overall, our studies indicate that p53, and potentially activators in general, serve as

escorts to dynamically recruit and load the basal transcription machinery onto DNA via complex

interactions between the RE, the core promoter and bound basal factors (i.e. TFIID). Our work

also reveals how p53 stimulates TFIID binding to different promoters that comprise variable

arrangements of p53 binding sites and core promoter elements. More importantly, our combined

functional and structural studies are crucial for understanding how eukaryotic transcription

complexes dynamically assemble on different protein-coding gene promoters.

Materials and Methods

Reagents and protein purification

Experimental details, including reagents, protein purification of the p53/TFIID/TFIIA/DNA co-complexes and fluorescent-labeled proteins/DNA used in this study along with our custom-built TIRF microscope setup, are described in the Supplemental Materials.

In vitro immuno-assembly assay

HeLa cells (32 liters) were grown in suspension culture with 1X DMEM plus 5% newborn calf serum for each assay. Nuclear extract was prepared and fractionated with phosphocellulose P11 (P-Cell) resins exactly as previously described (20). P-Cell column fractions eluting at 1M KCl/HEMG buffer (pH 7.9, [20 mM Hepes, 0.2 mM EDTA, 2 mM $MgCl_2$, 10% Glycerol, 1 mM DTT and 0.5 mM PMSF]) were pooled, dialyzed to 0.3 M KCl/HEMG buffer, added with final concentration of 0.1% NP40, and then immunoprecipitated overnight at 4°C with an anti-TAF4 mAb covalently conjugated to Protein G Sepharose beads (GE Healthcare Life Sciences). TAF4-immunoprecipitates were extensively washed with 0.65 M KCl/HEMG buffer, 0.3 M KCl/HEMG buffer and 0.1M KCl/HEMG buffer (containing 0.05% NP40 buffer and 10 μM Leupeptin) and split into four assembly reactions as indicated in Figure 1. Three reactions of TFIID-bound resins were incubated with 6-fold molar excess of promoter DNA. Next, two TFIID/DNA reactions were further added with 10-fold molar excess of p53 alone and/or TFIIA. Four total reactions were incubated at 4°C for 2 hours prior to elution of the co-complex from the TAF4 mAb affinity resin. TFIID/factor/DNA-bound resins were washed 5 times with 0.1 M KCl/HEMG buffer (containing 0.1% NP40 and 10 μM Leupeptin) followed by elution with a peptide (1 mg/ml) in 0.1 M KCl/HEMG buffer (plus 0.1% NP40 and 10 μM Leupeptin) recognized by the TAF4 mAb. Eluates were analyzed with 4-12% SDS-PAGE (NuPage gel system, Thermo Fisher) and visualized by silver staining analysis.

Xlink-Exo assay

Exonuclease assays were carried out in 10µl reactions containing a modified 50 mM KCl/HEMG buffer (pH 7.9, [12.5 mM Hepes, 0.05 mM EDTA, 6.25 mM $MgCl_2$, 5% glycerol] plus 1% DTT, 0.05 M PMSF, and 0.01% NP-40), IRdye800 (IDT) antisense strand labeled *hdm2* promoter DNA (9nM) , p53 (90nM, monomer), and TFIID (4nM). After incubation at room temperature for 20', reactions were crosslinked with formaldehyde (1% final) for 5' and stopped by addition of Tris-HCl (100mM final, pH 8.0). Next, each reaction was mixed with Exonuclease III (50 Units, NEB diluted in a buffer containing $MgCl_2$ [16.25 mM final]), and incubated at $37^{o}C$ for 1 hour. Proteinase K (10µg) was then added into each reaction and incubated at $70^{o}C$ for 3 hours. Reactions were mixed in formamide loading dye, heated at $95^{o}C$ and resolved on a 8% Polyacrylamide-TBE gel containing 7M urea. A series of known 5' truncated IRdye800 (IDT) antisense strand labeled wild-type *hdm2* DNA fragments were generated and used as molecular standards to determine the size of Exonuclease digested bands seen on our mapping gels.

TIRF Single Molecule Imaging

Passivated glass coverslips (see Supplemental materials) were assembled into a multichannel flow cell using double sided tape (3M). Flow cells were mounted on the microscope and streptavidin was added to the flow cell to coat surfaces. Each flow cell holds ~50 ul of sample. Reference beads that serve as fiducial marks were immobilized on the bottom surface of the flow cell. Biotinylated Cy3/Cy5 promoter DNA in 1X PBS was then loaded into the flow cell and bound over time specifically to the streptavidin that was non-covalently attached through biotin linkages on the top and bottom surfaces. Promoter DNA templates were briefly imaged at 3HZ to determine surface attached DNA densities that were sufficient for localization of single DNA molecules. Unbound biotinylated Cy3/Cy5 promoter DNA was washed away. Promoter DNA

templates were then imaged at 3HZ. The surface attached Cy3/Cy5 promoter DNA was then

photobleached. Reactivation of the Cy3/Cy5-labeled promoter DNA was not evident over the

timescale of our experiments (typically 15-20 minutes). Qdot 565-labeled TFIID (1 nM) in the

absence and presence of unlabeled p53 (8 nM, monomer) was then added to the flow cell in a

modified 50 mM KCl/HEMG buffer (pH 7.9, [12.5 mM Hepes, 0.05 mM EDTA, 6.25 mM

$MgCl_2$, 5% glycerol] plus 1% DTT, 0.05 M PMSF, and 0.01% NP-40) and immediately imaged

at 3Hz at 23^oC. An oxygen scavenging system consisting of 0.4% b-D-Glucose, 1 mg/ml glucose

oxidase (Sigma), and 0.4 mg/ml catalase (Sigma) was used for imaging. For experiments to

measure p53's interaction with promoter DNA, AlexaFluor 555, Cy3- or Cy5-labeled p53 (8nM,

monomer) in the absence or presence of unlabeled TFIID (1 nM) was added to the flow cell

containing surfaced attached mapped promoter DNA and immediately imaged at 3Hz.

AlexaFluor 555, Cy3- and Cy5-labeled p53 had equivalent residence times on promoter DNA

indicating that photobleaching of molecules didn't significantly affect results.

Single Molecule Co-localization Analysis

Fluorescent spots in each frame for the movie of the promoter DNA, TFIID or p53 were mapped

using 2D Gaussian fitting methods (21, 57, 58) to determine the location (x, y) at subpixel

resolution. Co-localization analysis was performed as in (21) with the following modifications.

Coordinates of fluorescent spots were adjusted based upon movement of the fiducial marks (i.e.

reference beads) over time. Stage movement/drift was typically less than 200 nm over the time of

imaging (~20 minutes). Fluorescent spots that were located within 84 nm (1 pixel) throughout

the movie were grouped together in a cluster. The positions of all spots within a cluster were

averaged to obtain the overall location of an individual cluster of molecules. Position maps of the

clustered surface attached DNA molecules were then compared with position maps of either the

TFIID or p53 clusters. Clusters of TFIID or p53 molecules within 3 pixels (252 nm) of a DNA

cluster were found. Displacements of TFIID or p53 clustered molecules with a maximum equal

to (0,0) in the x and y positions are indicative of non-random co-localization between DNA and

TFIID or p53 clusters. Displacements between the x,y positions of the TFIID or p53 clusters and

the DNA clusters were then plotted as a 2D histogram with a bin size of 50 nm. 2D Gaussian

fitting of the 2D histogram yields σ which is equivalent to the variation in the displacement of

TFIID or p53 within 252 nm of a DNA molecule. A co-localization threshold was set to 1.8σ (84

nm).

Single Molecule Residence Time Analysis of TFIID and p53 on promoter DNA

For each TFIID or p53 cluster that co-localized with a DNA cluster (i.e. within 84 nm), we the

plotted the amplitude from the 2D Gaussian fit of every fluorescent spot within the TFIID or p53

cluster as a function of time. Homemade MATLAB software was then used to analyze the rise

and the fall of the amplitude of the fluorescent signal to determine the appearance and

corresponding disappearance of each spot over time. The residence time for each DNA binding

event is defined as the difference in time between the appearance and disappearance of a TFIID

or p53 fluorescent spot. The fluorescent signal from TFIID or p53 had to last for at least 1.5" to

be considered as a discrete binding event in our analysis. Due to rapid stochastic blinking of the

quantum dot, a dissociation event was only considered to have occurred if the quantum dot signal

remained in the off state longer than 4.5". At least 300 binding events of co-localized TFIID or

100 binding events of p53 clusters were analyzed for each particular promoter DNA construct.

For a subset of molecules (~100), the data was processed by hand to determine the appearance

and disappearance of fluorescent spots. Comparison of the residence times generated by our

automated software and by hand yielded negligible differences. Histograms of residence times

for TFIID or p53 bound to promoter DNA were fit to a two-component exponential decay model

to determine the average residence time of TFIID or p53 bound to promoter DNA (58). If the

difference between the fitted residence times for each individual component in the two-

component model varied less than 20%, histograms were re-fitted using a single-component

exponential decay model (58). For Figures 3 and S5 data was additionally fit to a gamma

distribution (38) via custom MATLAB software.

2D Classification Analysis of TFIID co-complexes via negative stain EM

4 µL (10-20 ng in total amount) of fresh TFIID/p53/bax DNA assemblies in 0.1M

KCl/HEMG buffer (20mM HEPES, 0.2 mM EDTA, 2mM $MgCl_2$, 10% Glucose [final pH 7.9])

was applied directly onto a thin carbon film supported by holey carbon on a 400-mesh copper

grid (Pacific Grid tech.) for 3', which was freshly glow-discharged. After incubating the sample

on the grid for 3', the sample grid was stained with five successive 75 µL drops of 1% Uranyl

Formate.

The negative stain image data were collected with a Tecnai F20 TWIN transmission

electron microscope operating at 120 keV at a calibrated magnification of 62,000x with a

defocus range of -0.5µm to -2.5µm using a Tietz F416 (4K x 4K) CCD camera (resulting in a

2.82 A/pixel) under the dosage 39.98 e^-/Å. A total of 14,000 particles (for TFIID alone), 13,855

particles (for TFIID/p53), and 19262 particles (for TFIID/p53/bax DNA) were manually picked

using Boxer (EMAN; (59)). These particles were aligned and classified in a reference-free

fashion as previously described (24) for 8 iterations. The similar measurement of Lobe's A

position relative to the lobes B and C was employed as in (26). Based upon the position of Lobe

A in relation to Lobe C v.s. Lobe B (26), three conformational groups of TFIID were defined

(i.e. canonical, transition, and rearranged state) in this study. The canonical form is represented

by Lobe A retaining its attachment to Lobe C. The transition state is represented by Lobe A

beginning to detach and move in between Lobe C and B. The rearranged form is represented by

Lobe A reattaching to Lobe B. Next, individual classums from the 5^{th}, 6^{th}, 7^{th} and 8^{th} iteration

for each TFIID/p53 prep with or without the *bax* DNA were examined carefully to select 2D

class averages displaying views of these three conformational groups.   The number of 2D

averages in each conformational form v.s. total averages was calculated to yield the percentage

of TFIID's three conformations in each TFIID/p53 prep with or without the *bax* DNA. Three

independent protein purifications and EM data acquisition/analysis were performed to obtain the

standard deviation for the figure. A two sided student's T-test was used to determine statistical

significance.

Cryo-EM data collection and image analysis

For cryo-EM sample grid preparation, a thin carbon film supported by a 400-mesh carbon-

thickened C-flat holey grid with hole diameter 1.2 μm/spacing 1.3 μm (CF-1.2/1.3-4C,

Protochips) was plasma cleaned for 2 min using Denton high-vacuum evaporator. The grid was

loaded into a Vitrobot (FEI) that was preset at 100% humidity at 4ºC for vitrification of our

samples. 3 μl (504~900 ng in total amount) of the p53/TFIID/TFIIA/DNA assemblies was

applied directly onto the grid, incubated for 2 min and then blotted for 6.5 sec prior to plunge-

frozen in liquid ethane.

Cryo-EM data were collected with a JEM-2100F transmission electron microscope (JEOL)

operated at 120 KeV and at a magnification of 50,000x with a defocus range of –2.5 μm to –4.5

633   μm. Digital micrographs were collected using a TEM 2048x 2048 CCD camera (TemCam-

634   F224HD, TVIPS) with a pixel size of 24 μm and a calibrated magnification of 84,037x (2.86

635   Å/pixel) under low-dose conditions (~13 e$^-$ /Å) via the semi-automatic image acquisition

636   SerialEM software (60). Digital micrographs showed homogeneous particles in terms of lack of

637   aggregation and apparent integrity of the intact complexes.

638   Particles were manually selected using boxer (EMAN) (59) for an initial total of 32,275

639   particles from the p53/TFIID/TFIIA/*bax* DNA dataset and 22,070 particles from the

640   p53/TFIID/TFIIA/*hdm2* DNA dataset. The particles were then phase flipped using CTF-

641   estimated values determined by CTFFIND3(61) and extracted using SPIDER (<u>S</u>ystem for

642   <u>P</u>rocessing of <u>I</u>mage <u>D</u>ata from <u>E</u>lectron microscopy and <u>R</u>elated fields (62)) to a particle

643   window size of 128 x 128 pixels (5.72 Å/pixel), following the protocol as described in (26). The

644   particles were then normalized prior to the following data analyses. Reference-free 2D class

645   averaging image analysis was performed using IMAGIC (63) following the procedure precisely

646   as described (20). Using these 2D class averages as templates, we automatically picked particles

647   from a larger cryo-EM dataset using Signature (64) embedded in the Appion image processing

648   platform (65). A final total of 59,632 particles for the p53/TFIID/TFIIA/*bax* DNA co-complex

649   was obtained, normalized, and low-pass band filtered to perform the 3D reconstruction using

650   SPIDER's multi-reference projection matching approach as previously described (29). Both

651   rearranged and canonical forms of TFIID/TFIIA/SCP DNA from Eletron Microscopy Data Bank

652   (accession number 10832 and 10834, respectively) were filtered to 60 Å as a reference for the

653   first cycle of projection matching. Different initial reference models including TFIID alone, p53-

654   bound TFIID or TFIIA-bound TFIID were also tested for model bias refinement. New 3D

655   volumes reconstructed from the data set were used as references for all subsequent cycles of

alignment (angular step sizes ranging from an initial 20° to a final 2°). The resolution throughout

the refinement was determined by the 0.5 cut-off in the Fourier shell correlation (FSC) curve.

The 3D structure of the p53/TFIID/TFIIA/bax assembly was filtered at a final resolution ~35 Å.

All the 3D reconstructions were represented as isodensity surfaces using the UCSF Chimera

package (66) developed by the Resource for Biocomputing, Visualization, and Informatics at the

University of California, San Francisco (supported by NIH NIGMS P41-GM103311).

663
664
665
666

Funding information

This study was supported by start up funds (Albert Einstein College of Medicine) and

NIH/NIBIB 1U01EB021236-02. Some of this work was performed at the Simons Electron

Microscopy Center at the New York Structural Biology Center that is supported by a grant from

the Simons Foundation (Grant Number: 349247) with additional support from: NIH S10

OD019994-01, the Agouron Institute (Grant Number: F00316), NIH S10 RR029300-01, NIH

S10 RR017291-01, NYSTAR, and NIH C06 RR017528-01-CEM.

Acknowledgement

We specifically thank R. Tjian, S. Chu, and E. Nogales for their initial support of this study,

S. Zheng for generating TAF4 mAb supernatant and HeLa cells, D. King for providing the TAF4

mAb elution peptide, A. Revyakin, A. Pertsinidis and S.R. Park for their help on initial stages of

the single molecule microscopy, and R. Henderson for his technical advice critical for the cryo-

EM work. We also thank S.M. Shenoy, J. Hargitai, J. Wang, S. Lenny, J. Greally and E. Eng for

high performance computing cluster and cryo-EM technical support. We appreciate assistance

from Einstein's AIF facility, especially F. P. Macaluso, L. Cummins and G. S. Perumal. We are

grateful to M. Keogh, Charles Kenworthy and E. Nogales for critical comments of the

manuscript. W. Liu is an affiliated member of the New York Structural Biology Center. M.

Cianfrocco is an HHMI fellow of the Damon Runyon Cancer Research Foundation.

Figure legends

Figure 1. Single Molecule Total Internal Reflection Fluorescence (TIRF) Microscopy

analysis of Quantum dot-labeled TFIID binding of promoter DNA in vitro. (A) Schematic

representation of an in vitro single molecule TIRF assay to monitor binding of quantum dot

(QD)-labeled TFIID to fluorescently labeled *bax* or *hdm2* promoter DNA in real-time. (B)

Dependence of TFIID/promoter DNA co-localization on p53, the p53 RE, and core promoter

elements. Promoter DNA binding is plotted as a percentage of total DNA molecules in a field of

view that are co-localized with TFIID over 15' of binding based on TFIID−promoter DNA

displacement histograms. A co-localization threshold of 84 nm was set based upon our previous

study (21). TFIID binding to wild-type *bax* DNA is compared with and without p53. Additional

comparisons were made for TFIID binding to wild-type (WT), Mutant p53 RE (Mu RE), Mutant

Core (Mu Core), or combined mutant RE and core (Mu RE/Co) *hdm2* DNA templates. (C) xlink-

Exo analysis of TFIID binding to wild-type (top panel) and Mu RE/Co (bottom panel) *hdm2*

DNA. Basepair annotations (+/- 2bp) above individual peaks in the line traces (right panel)

represent a -4bp adjustment to reflect the known inability of Exonuclease to cut immediately

adjacent to protein/DNA crosslink sites (32).

704
Figure 2. Residence time analysis of TFIID bound to promoter DNA. (A) Traces of

individual co-localized TFIID/promoter DNA binding events were further analyzed to determine

TFIID's average residence time. TFIID displays a mixture of short (left panel) and long-lived

(right panel) binding events. (B) 1−Cumulative Distribution Function Plots (1−CDF) of TFIID

bound to DNA in the absence (left panel) and presence (right panel) of p53 were fitted to a single

(gray dashed) or two-component (red solid) exponential decay model. Fitting analysis reveals

that TFIID complexes bound to promoter DNA display residence times comprised of a long- (τ1)

and a short- ($\tau 2$) lived component. Percentages of the long- ($\tau 1$) and short- ($\tau 2$) lived

components are listed next to the residence time. (C) Bar graph comparing TFIID's average

short- and long-binding in the absence and presence of p53. (D) Percentage of short and long-

lived TFIID binding events on the wild-type (RE), mutant RE (Mu RE), mutant Core (Mu Core)

and mutant RE/Core (Mu RE/Co) *hdm2* promoter DNA. * = p-value < 0.05, *** = p-value <

0.001, N.S. = Not significant.

Figure 3. Residence time analysis of p53 bound to promoter DNA with and without TFIID.

(A) Individual co-localized p53/promoter DNA binding events were further analyzed to

determine the average residence time for p53 bound to wild-type (left panel) and Mutant p53 RE

(right panel) *hdm2* promoter DNA. 1−Cumulative Distribution Function Plots (1−CDF) of p53

bound to promoter DNA were fit to a single (gray dashed) or two (red solid)-component

exponential decay model. Based on comparisons of the *hdm2* promoter DNAs containing wild-

type and mutant p53 REs, $\tau 1$ represents specific binding of p53 to the wild-type *hdm2* DNA

while τns equals non-specific interactions between p53 and promoter DNA. (B) 1−CDF plots of

p53 bound to the wild-type *hdm2* DNA template in the presence of TFIID were fit to a single

(gray dashed) and two (red solid)-component exponential decay (left panel) or gamma

distribution (right panel) model. (C) Bar graph of p53's residence time in the absence and

presence of TFIID bound to the wild-type *hdm2* DNA (left panel). Residence time analysis of

TFIID and p53 bound to the wild-type *hdm2* DNA (right panel).

Figure 4. Structural EM analysis on p53's impact on TFIID bound to native target gene

promoters. (A) TFIID's structural plasticity as defined by the position of its Lobe A (26)(top

panels, the major lobes of TFIID are labeled as A, B, and C). TFIID's conformations are

classified into three groups: Canonical, Transition and Rearranged state as shown. Reference-

free 2D classification analysis on negative stain EM datasets showing populations of different

TFIID conformations in the absence of p53 (gray), presence of p53 (white), and with p53 + *bax*

DNA (black). The scale bar represents 100Å. (B) Three representative reference-free 2D class

averages of p53/TFIID/TFIIA bound to either the *bax* or *hdm2* DNA are shown based upon cryo-

EM particle datasets. The scale bar is 200Å. (C) Distinct 2D class averages showing views of the

assemblies bound to the *bax* promoter DNA fragment downstream (top panel, DNA highlighted

in green arrows) and upstream (middle panel, DNA highlighted in yellow arrows) of the

transcription start site (TSS, +1) are presented respectively. The orientation of the core promoter

elements bound by TFIID were previously mapped in the 3D structure of TFIID/TFIIA bound to

the super-core promoter (SCP) DNA (26). The corresponding views of the 3D structure of

TFIID/TFIIA/SCP DNA are shown (far right) to indicate the DNA orientation. The strong 3D

density highlighted in green represents the promoter DNA comprising the initiator, TSS and

downstream promoter elements. Since the upstream promoter DNA was weakly visible, focused

classification analysis (bottom panel) were performed using those corresponding class averages

(middle panel) to clarify the presence of the *bax* promoter DNA upstream of the TATA box

(highlighted by yellow arrow). The scale bar is 200Å.

Figure 5. A representative model of how TFIID engages the core promoter at various p53-

responsive promoters. A proposed model of how p53 aids TFIID in core promoter recognition

is presented based upon our current and previous studies (20). In the absence of p53, TFIID

primarily exists in the canonical conformation which is weakly permissive for DNA binding due

in part to interactions between the TAF1 TAND and TBP. The canonical form of TFIID can

infrequently and transiently interact with DNA. In rare occurances, canonical TFIID isomerizes

to a structurally rearranged form that involves severance of TAF1 TAND/TBP contacts and

movement of TBP from lobe A/C to lobe B. In the presence of p53, the TBP/TAF1 TAND

interaction is disrupted leading to a TFIID complex that is highly permissive to isomerization

and stable DNA binding. Once TFIID structurally rearranges and binds DNA via movement of

TBP from lobe A/C to lobe B, p53 disengages TFIID through loss of its contacts with Lobe A

and C. Dissociation of p53 from the TFIID/promoter DNA scaffold results in "recycling" of the

p53 RE for further assembly of the PIC components.

References

1.   **Menendez D, Inga A, Resnick MA.** 2009. The expanding universe of p53 targets.
Nat Rev Cancer **9:**724-737.
2.   **Bieging KT, Mello SS, Attardi LD.** 2014. Unravelling mechanisms of p53-mediated
tumour suppression. Nat Rev Cancer **14:**359-370.
3.   **Tafvizi A, Huang F, Fersht AR, Mirny LA, van Oijen AM.** 2011. A single-molecule
characterization of p53 search on DNA. Proc Natl Acad Sci U S A **108:**563-568.
4.   **Lu H, Levine AJ.** 1995. Human TAFII31 protein is a transcriptional coactivator of
the p53 protein. Proc Natl Acad Sci U S A **92:**5154-5158.
5.   **Thut CJ, Chen JL, Klemm R, Tjian R.** 1995. p53 transcriptional activation mediated
by coactivators TAFII40 and TAFII60. Science **267:**100-104.
6.   **Buschmann T, Lin Y, Aithmitti N, Fuchs SY, Lu H, Resnick-Silverman L,**
Manfredi JJ, Ronai Z, Wu X. 2001. Stabilization and activation of p53 by the
coactivator protein TAFII31. J Biol Chem **276:**13852-13857.
7.   **Li AG, Piluso LG, Cai X, Gadd BJ, Ladurner AG, Liu X.** 2007. An acetylation switch
in p53 mediates holo-TFIID recruitment. Mol Cell **28:**408-421.
8.   **Juven-Gershon T, Hsu JY, Kadonaga JT.** 2006. Perspectives on the RNA
polymerase II core promoter. Biochem Soc Trans **34:**1047-1050.
9.   **Colgan J, Manley JL.** 1992. TFIID can be rate limiting in vivo for TATA-containing,
but not TATA-lacking, RNA polymerase II promoters. Genes Dev **6:**304-315.
10.   **Chi T, Carey M.** 1996. Assembly of the isomerized TFIIA--TFIID--TATA ternary
complex is necessary and sufficient for gene activation. Genes Dev **10:**2540-2550.
11.   **Wang W, Gralla JD, Carey M.** 1992. The acidic activator GAL4-AH can stimulate
polymerase II transcription by promoting assembly of a closed complex requiring
TFIID and TFIIA. Genes Dev **6:**1716-1727.
12.   **White J, Brou C, Wu J, Lutz Y, Moncollin V, Chambon P.** 1992. The acidic
transcriptional activator GAL-VP16 acts on preformed template-committed
complexes. Embo J **11:**2229-2240.
13.   **Chi T, Carey M.** 1993. The ZEBRA activation domain: modular organization and
mechanism of action. Mol Cell Biol **13:**7045-7055.
14.   **Lieberman PM, Berk AJ.** 1994. A mechanism for TAFs in transcriptional activation:
activation domain enhancement of TFIID-TFIIA--promoter DNA complex formation.
Genes Dev **8:**995-1006.
15.   **Chi T, Lieberman P, Ellwood K, Carey M.** 1995. A general mechanism for
transcriptional synergy by eukaryotic activators. Nature **377:**254-257.
16.   **Kobayashi N, Boyer TG, Berk AJ.** 1995. A class of activation domains interacts
directly with TFIIA and stimulates TFIIA-TFIID-promoter complex assembly. Mol
Cell Biol **15:**6465-6473.
17.   **Rhee HS, Pugh BF.** 2012. Genome-wide structure and organization of eukaryotic
pre-initiation complexes. Nature **483:**295-301.
18.   **Lieberman PM, Ozer J, Gursel DB.** 1997. Requirement for transcription factor IIA
(TFIIA)-TFIID recruitment by an activator depends on promoter structure and
template competition. Mol Cell Biol **17:**6624-6632.

19.   **Xing J, Sheppard HM, Corneillie SI, Liu X.** 2001. p53 Stimulates TFIID-TFIIA-
814   promoter complex assembly, and p53-T antigen complex inhibits TATA binding
815   protein-TATA interaction. Mol Cell Biol **21:**3652-3661.
816   20.   **Liu WL, Coleman RA, Ma E, Grob P, Yang JL, Zhang Y, Dailey G, Nogales E, Tjian
817   R.** 2009. Structures of three distinct activator-TFIID complexes. Genes Dev **23:**1510-
818   1521.
819   21.   **Revyakin A, Zhang Z, Coleman RA, Li Y, Inouye C, Lucas JK, Park SR, Chu S, Tjian
820   R.** 2012. Transcription initiation by human RNA polymerase II visualized at single-
821   molecule resolution. Genes Dev **26:**1691-1702.
822   22.   **Zhang Z, English BP, Grimm JB, Kazane SA, Hu W, Tsai A, Inouye C, You C,
823   Piehler J, Schultz PG, Lavis LD, Revyakin A, Tjian R.** 2016. Rapid dynamics of
824   general transcription factor TFIIB binding during preinitiation complex assembly
825   revealed by single-molecule analysis. Genes Dev **30:**2106-2118.
826   23.   **Tantale K, Mueller F, Kozulic-Pirher A, Lesne A, Victor JM, Robert MC, Capozi S,
827   Chouaib R, Backer V, Mateos-Langerak J, Darzacq X, Zimmer C, Basyuk E,
828   Bertrand E.** 2016. A single-molecule view of transcription reveals convoys of RNA
829   polymerases and multi-scale bursting. Nat Commun **7:**12248.
830   24.   **Liu WL, Coleman RA, Grob P, King DS, Florens L, Washburn MP, Geles KG, Yang
831   JL, Ramey V, Nogales E, Tjian R.** 2008. Structural changes in TAF4b-TFIID correlate
832   with promoter selectivity. Mol Cell **29:**81-91.
833   25.   **Papai G, Tripathi MK, Ruhlmann C, Layer JH, Weil PA, Schultz P.** 2010. TFIIA and
834   the transactivator Rap1 cooperate to commit TFIID for transcription initiation.
835   Nature **465:**956-960.
836   26.   **Cianfrocco MA, Kassavetis GA, Grob P, Fang J, Juven-Gershon T, Kadonaga JT,
837   Nogales E.** 2013. Human TFIID binds to core promoter DNA in a reorganized
838   structural state. Cell **152:**120-131.
839   27.   **Louder RK, He Y, Lopez-Blanco JR, Fang J, Chacon P, Nogales E.** 2016. Structure
840   of promoter-bound TFIID and model of human pre-initiation complex assembly.
841   Nature **531:**604-609.
842   28.   **Andel F, 3rd, Ladurner AG, Inouye C, Tjian R, Nogales E.** 1999. Three-
843   dimensional structure of the human TFIID-IIA-IIB complex. Science **286:**2153-2156.
844   29.   **Grob P, Cruse MJ, Inouye C, Peris M, Penczek PA, Tjian R, Nogales E.** 2006. Cryo-
845   electron microscopy studies of human TFIID: conformational breathing in the
846   integration of gene regulatory cues. Structure **14:**511-520.
847   30.   **Bourdon JC, Deguin-Chambon V, Lelong JC, Dessen P, May P, Debuire B, May E.**
848   1997. Further characterisation of the p53 responsive element--identification of new
849   candidate genes for trans-activation by p53. Oncogene **14:**85-94.
850   31.   **Juven-Gershon T, Cheng S, Kadonaga JT.** 2006. Rational design of a super core
851   promoter that enhances gene expression. Nat Methods **3:**917-922.
852   32.   **Chang GS, Chen XA, Park B, Rhee HS, Li P, Han KH, Mishra T, Chan-Salis KY, Li Y,
853   Hardison RC, Wang Y, Pugh BF.** 2014. A Comprehensive and High-Resolution
854   Genome-wide Response of p53 to Stress. Cell Rep **8:**514-527.
855   33.   **Galasinski SK, Lively TN, Grebe De Barron A, Goodrich JA.** 2000. Acetyl
856   coenzyme A stimulates RNA polymerase II transcription and promoter binding by
857   transcription factor IID in the absence of histones. Mol Cell Biol **20:**1923-1930.

34. **Zhang Z, Boskovic Z, Hussain MM, Hu W, Inouye C, Kim HJ, Abole AK, Doud MK, Lewis TA, Koehler AN, Schreiber SL, Tjian R.** 2015. Chemical perturbation of an intrinsically disordered region of TFIID distinguishes two modes of transcription initiation. Elife **4**.

35. **Mazur SJ, Sakaguchi K, Appella E, Wang XW, Harris CC, Bohr VA.** 1999. Preferential binding of tumor suppressor p53 to positively or negatively supercoiled DNA involves the C-terminal domain. J Mol Biol **292:**241-249.

36. **Weinberg RL, Freund SM, Veprintsev DB, Bycroft M, Fersht AR.** 2004. Regulation of DNA binding of p53 by its C-terminal domain. J Mol Biol **342:**801-811.

37. **Emamzadah S, Tropia L, Vincenti I, Falquet B, Halazonetis TD.** 2014. Reversal of the DNA-binding-induced loop L1 conformational switch in an engineered human p53 protein. J Mol Biol **426:**936-944.

38. **Floyd DL, Harrison SC, van Oijen AM.** 2010. Analysis of kinetic intermediates in single-particle dwell-time distributions. Biophys J **99:**360-366.

39. **Miyashita T, Reed JC.** 1995. Tumor suppressor p53 is a direct transcriptional activator of the human bax gene. Cell **80:**293-299.

40. **Phelps M, Darley M, Primrose JN, Blaydes JP.** 2003. p53-independent activation of the hdm2-P2 promoter through multiple transcription factor response elements results in elevated hdm2 expression in estrogen receptor alpha-positive breast cancer cells. Cancer Res **63:**2616-2623.

41. **Liu D, Ishima R, Tong KI, Bagby S, Kokubo T, Muhandiram DR, Kay LE, Nakatani Y, Ikura M.** 1998. Solution structure of a TBP-TAF(II)230 complex: protein mimicry of the minor groove surface of the TATA box unwound by TBP. Cell **94:**573-583.

42. **Anandapadamanaban M, Andresen C, Helander S, Ohyama Y, Siponen MI, Lundstrom P, Kokubo T, Ikura M, Moche M, Sunnerhagen M.** 2013. High-resolution structure of TBP with TAF1 reveals anchoring patterns in transcriptional regulation. Nat Struct Mol Biol **20:**1008-1014.

43. **Kokubo T, Gong DW, Yamashita S, Horikoshi M, Roeder RG, Nakatani Y.** 1993. Drosophila 230-kD TFIID subunit, a functional homolog of the human cell cycle gene product, negatively regulates DNA binding of the TATA box-binding subunit of TFIID. Genes Dev **7:**1033-1046.

44. **Kokubo T, Yamashita S, Horikoshi M, Roeder RG, Nakatani Y.** 1994. Interaction between the N-terminal domain of the 230-kDa subunit and the TATA box-binding subunit of TFIID negatively regulates TATA-box binding. Proc Natl Acad Sci U S A **91:**3520-3524.

45. **Lively TN, Ferguson HA, Galasinski SK, Seto AG, Goodrich JA.** 2001. c-Jun binds the N terminus of human TAF(II)250 to derepress RNA polymerase II transcription in vitro. J Biol Chem **276:**25582-25588.

46. **Lively TN, Nguyen TN, Galasinski SK, Goodrich JA.** 2004. The basic leucine zipper domain of c-Jun functions in transcriptional activation through interaction with the N terminus of human TATA-binding protein-associated factor-1 (human TAF(II)250). J Biol Chem **279:**26257-26265.

47. **Kokubo T, Swanson MJ, Nishikawa JI, Hinnebusch AG, Nakatani Y.** 1998. The yeast TAF145 inhibitory domain and TFIIA competitively bind to TATA-binding protein. Mol Cell Biol **18:**1003-1012.

904   48.   **Ozer J, Mitsouras K, Zerby D, Carey M, Lieberman PM.** 1998. Transcription factor
905   IIA derepresses TATA-binding protein (TBP)-associated factor inhibition of TBP-
906   DNA binding. J Biol Chem **273:**14293-14300.
907   49.   **Xiao H, Pearson A, Coulombe B, Truant R, Zhang S, Regier JL, Triezenberg SJ,**
908   **Reinberg D, Flores O, Ingles CJ, et al.** 1994. Binding of basal transcription factor
909   TFIIH to the acidic activation domains of VP16 and p53. Mol Cell Biol **14:**7013-7024.
910   50.   **Gazit K, Moshonov S, Elfakess R, Sharon M, Mengus G, Davidson I, Dikstein R.**
911   2009. TAF4/4b x TAF12 displays a unique mode of DNA binding and is required for
912   core promoter function of a subset of genes. J Biol Chem **284:**26286-26296.
913   51.   **Liu X, Berk AJ.** 1995. Reversal of in vitro p53 squelching by both TFIIB and TFIID.
914   Mol Cell Biol **15:**6474-6478.
915   52.   **Espinosa JM, Verdun RE, Emerson BM.** 2003. p53 functions through stress- and
916   promoter-specific recruitment of transcription initiation components before and
917   after DNA damage. Mol Cell **12:**1015-1027.
918   53.   **Singh SK, Qiao Z, Song L, Jani V, Rice W, Eng E, Coleman RA, Liu WL.** 2016.
919   Structural visualization of the p53/RNA polymerase II assembly. Genes Dev
920   **30:**2527-2537.
921   54.   **Okuda M, Nishimura Y.** 2015. Real-time and simultaneous monitoring of the
922   phosphorylation and enhanced interaction of p53 and XPC acidic domains with the
923   TFIIH p62 subunit. Oncogenesis **4:**e150.
924   55.   **Leveillard T, Andera L, Bissonnette N, Schaeffer L, Bracco L, Egly JM, Wasylyk**
925   **B.** 1996. Functional interactions between p53 and the TFIIH complex are affected by
926   tumour-associated mutations. EMBO J **15:**1615-1624.
927   56.   **Poorey K, Viswanathan R, Carver MN, Karpova TS, Cirimotich SM, McNally JG,**
928   **Bekiranov S, Auble DT.** 2013. Measuring chromatin interaction dynamics on the
929   second time scale at single-copy genes. Science **342:**369-372.
930   57.   **Huang B, Jones SA, Brandenburg B, Zhuang X.** 2008. Whole-cell 3D STORM
931   reveals interactions between cellular structures with nanometer-scale resolution.
932   Nat Methods **5:**1047-1052.
933   58.   **Chen J, Zhang Z, Li L, Chen BC, Revyakin A, Hajj B, Legant W, Dahan M, Lionnet**
934   **T, Betzig E, Tjian R, Liu Z.** 2014. Single-molecule dynamics of enhanceosome
935   assembly in embryonic stem cells. Cell **156:**1274-1285.
936   59.   **Ludtke SJ, Baldwin PR, Chiu W.** 1999. EMAN: semiautomated software for high-
937   resolution single-particle reconstructions. J Struct Biol **128:**82-97.
938   60.   **Mastronarde DN.** 2005. Automated electron microscope tomography using robust
939   prediction of specimen movements. J Struct Biol **152:**36-51.
940   61.   **Mindell JA, Grigorieff N.** 2003. Accurate determination of local defocus and
941   specimen tilt in electron microscopy. J Struct Biol **142:**334-347.
942   62.   **Shaikh TR, Gao H, Baxter WT, Asturias FJ, Boisset N, Leith A, Frank J.** 2008.
943   SPIDER image processing for single-particle reconstruction of biological
944   macromolecules from electron micrographs. Nat Protoc **3:**1941-1974.
945   63.   **van Heel M, Harauz G, Orlova EV, Schmidt R, Schatz M.** 1996. A new generation of
946   the IMAGIC image processing system. J Struct Biol **116:**17-24.
947   64.   **Chen JZ, Grigorieff N.** 2007. SIGNATURE: a single-particle selection system for
948   molecular electron microscopy. J Struct Biol **157:**168-173.

949   65.   **Lander GC, Stagg SM, Voss NR, Cheng A, Fellmann D, Pulokas J, Yoshioka C,**
950   **Irving C, Mulder A, Lau PW, Lyumkis D, Potter CS, Carragher B.** 2009. Appion: an
951   integrated, database-driven pipeline to facilitate EM image processing. J Struct Biol
952   **166:**95-102.
953   66.   **Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC,**
954   **Ferrin TE.** 2004. UCSF Chimera--a visualization system for exploratory research
955   and analysis. J Comput Chem **25:**1605-1612.
956

Figure 1

Figure 2

A

B

C

D

A TFIID form

Canonical ⇌ Transition ⇌ Rearranged

C strong downstream DNA view

+1

weak upstream DNA view

Focused classification: upstream DNA

B

TFIID/p53/TFIIA/promoter DNA

bax promoter

hdm2 promoter

TFIID
TFIID/p53
TFIID/p53/*bax* DNA

holo-TFIID

Canonical

TAF6
TAF2
B
C
TAF1-
TAND
TAF1
TBP
A

**Weakly-permissive for
DNA binding**

Downstream
DNA
B
C
TATA
RE
A

7 sec.

**Transient TFIID/DNA
"encounter complex"**

Isomerization

Rearranged

B
TATA
TBP
RE
A
C

p53/TFIID

Canonical

TAF6
TAF2
B
C
p53
TAF1-
TAND
TAF1
TBP
A

**Strongly-permissive for
DNA binding**

Downstream
DNA
B
C
p53
TATA
RE
A

**Transient p53/TFIID/DNA
"encounter complex"**

Isomerization

Rearranged

B
TATA
TBP
RE
p53
A
C

**Transient isomerized
p53/TFIID/DNA complex**

p53

TAF6
TAF2
B
C
TAF1-
TAND
TAF1
TBP
A

Weakly-permissive for DNA binding

RE
TATA
Downstream
DNA

100 sec.

**p53 dissociates
to liberate the RE**

p53

B
TATA
TBP
RE
A
C

Stable isomerized TFIID/DNA complex