p53 dynamically directs TFIID assembly on target gene promoters

Coleman, R. A.[1#], Qiao, Z.[1], Singh, S. K, [1]Peng, C. S.[1], Cianfrocco, M.[2], Zhang, Z.[3], Song, L.[1], Piasecka, A.[1], Aldeborgh, H.[1], Basishvili, G.[1], Rice, W.[4], and Liu, W. L.[1]

[1]Gruss-Lipper Biophotonics Center, Department of Anatomy and Structural Biology, Albert Einstein College of Medicine, Bronx, NY 10461, USA.

[2]Department of Cellular and Molecular Medicine, University of California, San Diego, La Jolla, CA 92093.

[3]Howard Hughes Medical Institute, Janelia Research Campus, Ashburn, VA 20147, USA.

[4]New York Structural Biology Center, 89 Convent Avenue, Manhattan, NY 10027

#To whom correspondence should be addressed. Department of Anatomy and Structural Biology, Albert Einstein College of Medicine, Bronx, NY 10461 Tel: 718-430-8623; Email: robert.coleman2@einstein.yu.edu

Running title: Dynamics and structure of p53-induced TFIID assembly

Keywords: p53; TFIID; transcription; structure; single molecule microscopy

Abstract

p53 is a central regulator in tumor suppression. In response to various signals, p53 turns on vast gene networks to maintain cellular integrity, in part by aiding promoter recruitment of the TFIID-mediated transcription machinery to stimulate transcription initiation. However, the precise means by which p53 dynamically interacts with TFIID to facilitate assembly on target gene promoters remains elusive. To address this key question, we have undertaken an integrated approach involving single molecule fluorescence microscopy, single particle cryo-electron microscopy, and biochemistry. Our real-time single molecule imaging demonstrated that p53 both delivers and promotes the stable binding of TFIID to DNA. Furthermore, TFIID binds core promoter sequences via distinct modes displaying transient and prolonged stability. Importantly, TFIID's interaction with DNA induces p53 to rapidly dissociate, effectively recycling p53's binding site on the promoter. This notion is further supported by our structural studies. In addition, upstream promoter DNA comprising p53's response elements is highly mobile when bound to TFIID compared to the core promoter DNA. Collectively, these findings indicate that p53 serves as an escort to dynamically recruit and load the basal transcription machinery onto its target promoters by regulating the structural architecture of TFIID.

Introduction

More than 50% of cancer patients harbor p53 mutations, highlighting the essential role of this protein in tumor suppression (1). To properly maintain genomic stability, p53 induces vast gene networks involved in diverse cellular pathways including cell cycle arrest, apoptosis and DNA repair (2). In response to various stress stimuli, p53 acts as a transcriptional activator that specifically binds consensus response elements (REs; two ten base-pair half sites, RRRCWWGYYY) within its target genes to directly stimulate gene expression (2). p53's search for target REs is potentially expedited by its ability to non-specifically bind and slide along the DNA (3). Upon recognition of its response genes, p53 facilitates transcription at least in part by targeting the TFIID-mediated transcription machinery to the promoter (4-7). This large TFIID complex, composed of TBP and 14 TBP-associated factors (TAFs), is responsible for recognizing and binding multiple specific regions of the promoter (i.e. core promoter elements) surrounding the transcription start site (TSS) (8). Once bound, TFIID serves as a central scaffold for basal factors, which include RNA Polymerase II, to form a pre-initiation complex (PIC) that culminates with active transcription. Importantly, without the assistance of activators such as p53, TFIID's recognition of core promoter DNA sequences is weak and rate limiting for transcription initiation due to inefficient PIC assembly (9-16). As TFIID is responsible for transcription of at least ~90% of protein-coding genes (17), unraveling how p53 intervenes in this central step of the PIC assembly (i.e. TFIID's binding of promoter DNA) is essential for understanding mechanisms controlling gene-specific transcription initiation throughout the cell.

Previous biochemical studies of different activators (e.g. p53 and Zta) suggested a conserved albeit poorly characterized mechanism for facilitating transcription (10, 18, 19). It has been shown that p53 binds to its REs and enhances TFIID promoter recruitment via direct

contact with TBP and select TAFs (19, 20). Each of the various p53 target genes has a unique

arrangement of REs and core promoter elements (e.g. the TATA box, Initiator, and downstream

core promoter elements [DPE] (8)), which exhibit varied affinities for p53 and TFIID,

respectively (1). It is unclear how the variable spatial arrangement of REs with respect to core

promoter elements leads to distinct PIC assembly dynamics and gene-specific

p53/TFIID/promoter DNA structures. Additional evidence has shown that p53 and other

activators also stimulate structural changes within TFIID when associated with TFIIA and DNA

(10, 18, 19). This activator-induced isomerization results in the enhanced ability of TFIID to

bind core promoter element sequences downstream of the TSS (10, 18, 19). However, the

structural mechanism of p53-induced isomerization of TFIID and its impact on PIC assembly is

currently unknown.

Single molecule/particle-based imaging has emerged as an advanced tool to study

complex dynamic behavior and static structures of large multi-subunit protein assemblies

involved in regulating gene expression. A recent study utilizing real-time single molecule

fluorescence microscopy provided insight that TFIID's interaction with a synthetic core

promoter DNA tightly correlates with productive transcription (21). Despite years of intense

studies, how activators regulate TFIID's dynamic binding to native target DNA remains elusive.

In addition, advanced single particle transmission electron microscopy (EM) has unraveled the

three-dimensional (3D) structures of TFIID and its co-complexes. Recent EM studies unmasked

a structural plasticity of TFIID, likely attributed to its promoter recognition activity (20, 22-25).

In particular, human TFIID features a "horseshoe" shape containing three major lobes (A, B &

C) and a well-defined central cavity (26, 27). Interestingly, we demonstrated that, in the absence

of DNA, three different activators (e.g. p53) introduce a common set of local structural changes

in TFIID, including movement of lobe A towards lobe B resulting in a more closed central cavity

(20). Intriguingly, it was recently discovered that human TFIID undergoes a significant structural

rearrangement involving movement of Lobe A when bound to a specific promoter DNA

optimized for TFIID recognition (24). Whether p53's induced movement of Lobe A results in a

rearranged conformation of TFIID that contributes to stable interaction between TFIID and DNA

remains to be determined.

To uncover the mechanistic underpinnings of how p53 dynamically facilitates TFIID

assembly on promoter DNA, we exploited a combination of biochemistry, single molecule

fluorescence microscopy and single particle cryo-EM analysis. These studies illuminate dynamic

interactions and static global molecular architectures of p53-induced TFIID assemblies on two

representative target genes (i.e. *hdm2* and *bax*). Our real-time single molecule studies reveal that

p53 transiently interacts with DNA to help load TFIID onto promoters. To assist recruitment,

p53 stimulates structural rearrangement of TFIID to allow anchoring of downstream core

promoter DNA sequences. Notably, binding of TFIID to DNA induces dissociation of p53 from

the promoter, thereby effectively "recycling" the RE to allow recruitment of additional factors

involved in PIC formation. Our cryo-EM analyses further support this finding. In addition, the

upstream promoter that contains p53's response elements exerts structural plasticity when bound

to TFIID while the core promoter is stably bound. Collectively, these studies suggest that proper

response to divergent stress signals involves conserved structural mechanisms relating to p53's

ability to dynamically stimulate TFIID's interaction with various target promoters.

Results

Characterization of p53's impact on TFIID/DNA binding

We sought to decipher the molecular mechanism of how p53 targets TFIID to stimulate

transcription. To achieve this goal, we chose two representative p53 target genes, *bax* and *hdm2*

(28, 29). We previously established unique purification strategies to isolate stoichiometric

activator/TFIID assemblies with high purity and determined their 3D structures (20). Here our

protocol was adapted to include the addition of native promoter DNA fragments (i.e. *bax* or

hdm2, Figure 1). In brief, we conducted highly specific immunoprecipitations (IP) from

fractionated HeLa nuclear extracts using a monoclonal antibody (mAb) against the TAF4 subunit

of TFIID (Figure 1A). Resins containing the immunoprecipitated TFIID were washed

extensively and split into four reactions (as indicated in Figure 1A). Next, while TFIID was still

bound to the TAF4 affinity resin, three of these reactions were incubated with 1.) a 6-fold molar

excess of the native promoter DNA alone; 2.) promoter DNA and purified human p53; or 3.)

promoter DNA, p53 and TFIIA, respectively. Unbound factors and DNA were removed by

extensive washing of the TAF4 affinity resin. Co-complexes containing DNA and/or factors

bound to TFIID were eluted from the resin using specific peptides recognized by the TAF4 mAb.

The TFIID assemblies were analyzed by SDS-PAGE and silver staining (Figure 1B).

We found that addition of p53 increased the amount of promoter DNA fragments co-

purifying with TFIID (Figure 1B, lanes 3, 4, 9 and 10) compared to TFIID incubated with DNA

alone (lanes 2 and 8). These results suggest that p53 aids in TFIID's binding to native target

gene promoters. A previous biochemical report documented that p53 stimulated the assembly of

TFIIA and TFIID on a synthetic promoter DNA (19). Consistent with these observations,

addition of TFIIA along with p53 (lanes 4 and 10) further enhanced the amount of DNA co-

present with TFIID compared to incubation of p53 alone (lanes 3 and 9). Collectively, these

results suggest that p53 promotes TFIID binding to DNA.

p53 facilitates binding of TFIID to DNA via the response elements and core promoter

We sought to further understand the molecular mechanism pertaining to TFIID's

enhanced DNA binding (Figure 1B). Thus, dynamic interactions between p53, TFIID and

promoter DNA were characterized via a single molecule co-localization assay using Total

Internal Reflection Fluorescence Microscopy (TIRF) (21) along with Quantum dot-labeled

TFIID (QD-TFIID) and Cy3/Cy5-labeled native promoter DNA templates (Figure 2A). 2D

Projection maps indicated weak binding of TFIID to DNA within the first 15' with only 1.3% of

the wild-type _hdm2_ DNA occupied by TFIID (Figure 2B, left two panels). The majority of bound

complexes lasted ~7" (left two panels and Figure 3A), revealing a low intrinsic affinity of TFIID

for DNA. Increasing the time between addition of TFIID to the DNA and the imaging of binding

did not improve co-localization efficiencies (data not shown), suggesting that reactions are at

equilibrium. This finding suggests that, at low concentrations (~1nM), TFIID inefficiently

recognizes DNA. On the other hand, increased binding of TFIID to DNA was observed in our

bulk biochemical assembly assays when much higher concentrations (100-600nM) were used as

in Figure 1B.

Upon addition of p53, a clear enhancement of _bax_ or _hdm2_ promoter DNA bound by

TFIID was observed (Figure 2B, right panel and Figure 2C), which is consistent with the general

increase of DNA retained in our bulk biochemical assays (Figure 1B). This significant

enhancement is due to the direct interaction between p53 and TFIID, because a control

experiment detected no p53-mediated co-localization of DNA with the Quantum dot-labeled

TAF1 antibody ($0.2 \pm 0.07\%$ co-localization) in the absence of TFIID (data not shown). These

results confirm that p53 greatly facilitates TFIID binding to DNA.

To address the role of the p53 RE and core promoter elements in p53-mediated

TFIID/DNA binding, three mutant *hdm2* DNA templates were analyzed with the same real-time

co-localization assay (Figure 2C). The first mutant *hdm2* DNA template (i.e. Mu RE) harbors

mutations in the response elements known to impair p53's sequence-specific interactions (30). A

mild decrease of TFIID/DNA binding was observed on the mutant p53 RE template (Figure 2C).

This result suggests that p53's direct contacts with its RE are not critical for enhanced

TFIID/promoter DNA binding and other mechanisms may be utilized. Next, to assess the impact

of the core promoter in p53-mediated TFIID/DNA binding, mutations (i.e. Mutant Core) known

to inhibit transcription (31) were exploited. Upon addition of p53, TFIID's binding to the mutant

core promoter was reduced (Figure 2C). When both the p53 RE and core promoter elements

were altered (i.e. Mu p53 RE/Core), TFIID/DNA co-localization was significantly decreased

compared to the wild-type promoter (Figure 2C). These results indicate that the p53 RE and core

promoter elements were both required for optimal interaction of TFIID with p53 target gene

promoters.

As an alternative approach to study p53-mediated TFIID/DNA binding, we developed an

in vitro formaldehyde-crosslinking/exonuclease mapping assay (xlink-Exo), adapted from a well-

established strategy (ChIP-exo) to map out protein/DNA contacts of transcription factors in vivo

at near base-pair resolution (32). This analysis revealed that TFIID made extensive contacts with

well-characterized core promoter elements (i.e. TATA, InR, MTE and DPE) on the wild-type

hdm2 DNA (Figure 2D, top panels). Notably, addition of p53 led to elevated crosslinking of

TFIID to the wild-type core promoter with select elements showing much larger changes (i.e.

TATA, InR) than others (i.e. MTE/DPE) (top panels). This result is consistent with our single

molecule data showing that p53 increases the specific binding of TFIID to the core promoter.

Importantly, when the Mu RE/Co and Mu Core templates were examined, a clear reduction in

p53's ability to selectively enhance contacts within mutated core promoter elements was

observed (Figure 2D, bottom panels and Supplemental Figure S1). This finding suggests that a

reduction of p53-mediated TFIID contacts within TATA, InR and the DPE leads to a decrease

co-localization of TFIID with mutant promoters (Figure 2C). TFIID's contacts within the -4 and

+11 regions were still slightly elevated in the presence of p53 on these mutant core promoters.

This result may explain why some percentage of TFIID/mutant DNA co-localization was

retained in our single molecule assays (Figure 2C).

In addition to these well-established contacts, previous DNAse I footprinting studies

reported TFIID/DNA interactions downstream of the DPE (+32 to +68) and upstream of TATA

(-40 to -140) (24, 33, 34). Our xlink-Exo assays also mapped TFIID/DNA interactions far

upstream of TATA (-55 to -137) and downstream of the DPE (~ +37 to +68), which were

enhanced when p53 was co-present on the wild-type *hdm2* promoter (Figure 2D). Therefore,

TFIID may utilize additional uncharacterized elements to recognize promoters. Surprisingly,

some of TFIID's upstream contacts (i.e. -55, -61, -80, and -88) reside within the p53 RE on the

wild-type promoter even in the absence of p53 (Figure 2D, top panels). Mutations of the p53 RE

led to a loss of TFIID's contacts at -61 and -88, indicating that TFIID recognizes specific

sequences within the p53 RE (Figure 2D bottom panels and Supplemental Figure S1).

p53 stabilizes TFIID binding at the core promoter

Thus far, our studies demonstrate that p53 stimulates the association of TFIID with

promoter DNA. To test if p53 could also stabilize TFIID's binding to DNA, the amount of time

that QD-TFIID remains bound to a single promoter DNA before dissociating (i.e. residence time)

was determined (Figures 3 and Supplemental Figure S2). In the absence of p53, two populations

of DNA-bound TFIID complexes displaying distinct residence times were detected on the wild-

type *hdm2* template (i.e. long- and short-lived binding events) (Figure 3A, left panel).

TFIID/DNA binding events were predominantly short-lived (average residence time of 7.1 ± 0.6

seconds), with a minority of long-lived events (68.4 ± 8.0 seconds). Importantly, when p53 was

present, a majority of TFIID complexes on the wild-type *hdm2* DNA switched to long-lived with

an average residence time of 99.8 ± 5.0 seconds, while a minority exhibited a short-lived

residence time of 12.2 ± 0.8 seconds (right panel). TFIID also displayed a similar stability on the

bax promoter in the presence of p53 (Supplemental Figure S3). These results suggest that TFIID

alone weakly interacts with the DNA via two intrinsic binding modes that are both stabilized

when p53 is added.

We next assessed the role of the RE and core promoter elements in p53-mediated

stabilization of TFIID on DNA. Interestingly, disruption of the p53 binding site on the *hdm2*

promoter (Mu p53 RE) predominantly reduced the long-lived residence time of TFIID (avg =

75.1 ± 5.6") without significantly affecting the short-lived events (avg = 11.6 ± 1.0") (Figure 3B,

left panel). This suggests that interactions between p53 and its RE are required for maximal

stability of TFIID/DNA contacts on native target promoters. On the mutant core promoter (Mu

Core), a slight majority of TFIID/DNA binding events was now short-lived (54%). Residence

times for both the long- and short-lived TFIID/DNA binding events were also reduced (i.e. avg =

63.3 ± 4.7" and 5.8 ± 0.4", center panel), suggesting that both DNA binding modes were related

to core promoter recognition. When both the p53 RE and core promoter were altered (Mu

RE/Co), TFIID's promoter binding events were predominantly short-lived (72%) with reduced

short- and long-lived residence times relative to wild-type DNA (compare Figure 3B, right panel

v.s. Figure 3A, right panel). These experiments demonstrate that mutations of select core

promoter elements result in reduced stability of TFIID on DNA with additional changes in the

p53 RE further weakening its binding. Interestingly, our xlink-Exo studies showed that only a

few TFIID contacts on the mutant Core and RE/Core templates were slightly enhanced by p53

(Figures 2D and S1). Together with the single molecule data, our studies suggest that these

remaining enhanced contacts are sufficient to anchor TFIID, albeit weakly, to DNA. Overall,

these studies indicate that p53 generally enhances the stability of TFIID on DNA, which could be

important for recognition of weak non-consensus core promoter elements within p53 target genes.

Optimal stability requires p53's stimulation of TFIID's interaction with both the core promoter

elements and p53's binding site.

TFIID alters the binding dynamics of p53 to its cognate sites

Our previous study revealed that a variant form of TFIID could cooperatively increase

the binding of the activator c-Jun to its binding site on promoter DNA (22). To quantitatively

assess if this was the case for p53, the dissociation dynamics of fluorescently labeled-p53/DNA

interactions plus or minus TFIID were examined (Figure 4). In the absence of TFIID, p53

primarily displayed two populations of distinct binding events on the wild-type *hdm2* template

(i.e. long-lived [avg = 33.1 ± 3.9"] and short-lived [avg = 7.4 ± 0.7"], Figure 4A, left panel).

Since earlier reports showed that p53 can bind DNA non-specifically (35, 36), we hypothesized

that these short-lived p53/DNA binding events were non-specific interactions. To test our

hypothesis, the same experiments were performed using the mutant p53 RE template (right

panel). Indeed, disruption of p53's sequence-specific binding sties on DNA led to a single short-

lived population (avg = 6.8 ± 0.7"). Based upon this result, the long-lived p53/DNA binding

events most likely represent sequence-specific interactions (left panel), whereas the short-lived

p53/DNA binding events are non-specific in nature. Alternatively, these short-lived events could

represent interactions with an unknown lower affinity p53 binding site in the promoter or binding

of dimeric p53 to consensus half-sites within the mutant RE. Notably, photobleaching of the

fluorescently labeled p53 did not affect our results, since similar residence times were observed

with multiple dyes of differing photostability (data not shown). In addition, a previous study

detected similar short residence times of ~10-12" for p53 bound to a consensus site (37).

Collectively, these results indicate that p53 dynamically interacts with DNA via both specific

and non-specific interactions.

Intriguingly, addition of TFIID resulted in a single population of p53 bound to wild-type

DNA with an "intermediate-lived" residence time distinct from the two populations observed

with p53 alone (Figure 4B). We noticed that fitting of the data showed slight deviations from an

exponential decay model. Importantly, a previous study indicated that the existence of complex

multi-step dissociation events could be discerned using a gamma distribution model to fit single

molecule data (38). Indeed, the data was better represented by a gamma distribution model (right

panel). Parameters from the gamma distribution fit reveal that, in the presence of TFIID, p53

dissociates from the DNA in ~2 steps with roughly equivalent residence times of 8.2" per step

(average total time = 14.8 ± 1.5"). Furthermore, gamma distribution fitting of p53's dissociation

from wild-type DNA in the absence of TFIID indicated a single step dissociation pathway (data

not shown). These findings demonstrate that TFIID alters both the rate and mechanism of p53's

dissociation from promoter DNA.

Given that TFIID directly contacts the RE as revealed via our xlink-Exo assays (Figure

2D, top panels), it is possible that TFIID could help stabilize p53 binding to weak REs. Indeed,

in the presence of TFIID, there was a slight enhancement of p53's residence time on the Mu RE

DNA compared to p53 alone (Supplemental Figure S4A left panel versus Figure 4A right panel),

consistent with TFIID stabilizing p53's binding to a non-canonical mutant RE. This subtle

stabilization does not require TFIID's interaction with the TATA, InR and DPE elements, since

we also observed an increase in p53's residence time on the Mu RE/Co DNA (Supplemental

Figure S4B). Fitting of the dissociation data on the mutant templates to a gamma distribution

yielded poor fits, suggesting single step dissociation kinetics despite fitting parameters that

indicated a multi-step dissociation (Figures S4, right panels). Overall, these data indicate that

TFIID binding to DNA can influence p53's stability at the RE. TFIID's interaction with p53

likely promotes the dissociation of p53 bound to specific sequences in the promoter. Importantly,

the average time of p53 bound to the wild-type *hdm2* template (~15") is significantly shorter

than TFIID's long-lived DNA binding time (~100") (Figure 4C). These dynamic studies strongly

suggest that p53 transiently escorts TFIID to the promoter DNA. Importantly, p53 is not required

to remain bound for TFIID's maximum stability on promoter DNA.

p53 promotes a common DNA binding conformation of TFIID

Since p53 enhances TFIID's binding to DNA, we speculated that p53 regulates the

structural architecture of TFIID to aid recognition of various target genes. Our previous

structural studies demonstrated that, in the absence of DNA, p53 binds TFIID via its direct

contact with Lobe A and C (20). Intriguingly, this p53-bound TFIID did not undergo drastic

conformational changes. Rather, the p53/TFIID interaction led to unique local structural changes

within TFIID including the movement of Lobe A towards Lobe B (20). A recent cryo-EM study

discovered that human TFIID switches from a canonical to a rearranged conformation that

permits binding to the super core promoter (SCP) DNA optimized for interaction with TFIID (24,

31). This rearranged DNA binding conformation of TFIID was formed by the movement of Lobe

A from Lobe C to Lobe B (24). It was previously unclear how this p53-induced Lobe A

movement toward Lobe B was related to TFIID's rearranged DNA binding conformation.

Therefore, we hypothesized that p53-induced local structural changes aid in formation of the

rearranged state to facilitate TFIID's interaction with DNA. To test this idea, we set out to reveal

TFIID's global structural architecture when p53 and target promoter DNA were present.

As the first step, unbiased reference-free 2D classification experiments were performed to

examine conformational states of TFIID when p53 alone and p53/*bax* DNA were present (Figure

5A). Three conformational states of TFIID represented by corresponding views of 2D class

averages were classified based upon the position of Lobe A relative to Lobes C and B (top panels,

i.e. canonical, transition, and rearranged state). Strikingly, in the presence of p53, an elevated

population of TFIID displayed the rearranged form, while the percentage of transition state

remained similar compared to TFIID alone (bottom panel). This finding suggests that p53 can

specifically impact formation of TFIID's rearranged state. To assess the effect when p53 and the

bax DNA were added, the same 2D classification analysis was also conducted. Interestingly, the

populations of both the transition and rearranged forms increased when p53 and DNA was co-

present (bottom panel). This observation indicates that target gene promoters and/or its

interaction with p53 also play a role in TFIID's conformational switches.

A previous biochemical study reported that p53 stimulated the assembly of TFIIA and

TFIID on a synthetic promoter DNA (19). Interestingly, unlike p53, TFIIA alone promotes the

canonical form of TFIID (24). Thus, to assess p53's impact on the structural framework of

TFIID in the presence of TFIIA, two TFIID/p53/TFIIA assemblies on the native *hdm2* and *bax*

promoter fragments (488 bp and 500 bp) were obtained as described in Figure 1 (Supplemental

Figure S5). To visualize both conformations and the DNA, cryo-EM data of these assemblies

were collected for reference-free 2D classification analysis. Some class averages showing the

presence of promoter DNA spanning the central cavity of TFIID were observed (Figure 5B).

Surprisingly, both assemblies displayed a similar rearranged TFIID architecture, despite the

different organization of their promoters. Each template harbors non-canonical core promoter

elements with different spatial arrangements between TATA and the Initiator (Figure 1B) (8, 28,

39). Combined with our 2D classification analysis (Figure 5A), these structural studies suggest

that p53 promotes formation of a common rearranged TFIID/DNA binding conformation.

Next, to identify the relative DNA orientation and any distinct structural features, we

compared the class averages of our two assemblies with the previous structural analysis of

TFIID/TFIIA/SCP DNA containing a mapped DNA orientation (24). Our co-complexes

displayed clear views of TFIID stably bound to the *bax* DNA downstream of the TSS (Figure 5C,

top two panels). In contrast, upstream promoter DNA was weakly visible in our averages

(middle panel). These results suggest that the p53 RE upstream of the TATA box element is

flexible within the co-complex. To define the presence of the upstream promoter, focused

classification on a class average displaying flexible upstream DNA (middle panel) was

performed as previously described (24). This analysis revealed a clear density for the upstream

promoter DNA (highlighted by yellow arrow, bottom panel). Given the strong density for

downstream DNA within the assembly, these results indicate that TFIID serves as a strong

anchor for downstream promoter DNA during PIC assembly.

Another observation is that the 2D class averages of p53/TFIID/TFIIA/DNA assemblies

exhibit a similar topology to the averages from TFIID/TFIIA bound to the SCP DNA (refer to

Figure 3 in (24)) raising two major points. First, these data suggest that TFIID adapts a common

DNA binding conformation to bind various p53 target gene promoters. Second, p53 may

dissociate from TFIID, as supported by our single molecule dynamics results (Figure 4), in order

to complete structural rearrangements associated with the DNA binding form of TFIID. To test

these hypotheses, single particle 3D reconstruction was conducted to determine the 3D structure

of the TFIID/TFIIA/p53/*bax* DNA co-complex (Supplemental Figure S6A). The overall

structural architecture of our co-complex is comparable to the 3D structure of TFIID/TFIIA/SCP

DNA. Our 3D reconstruction was also verified by analyzing its 3D projections with their

matching reference-free 2D averages (Supplemental Figure S6B). This result suggests that TFIID

utilizes one common rearranged form to bind various promoters.

Our 3D reconstruction did not reveal any prominent extra densities that would most

likely represent the p53 protein. This indeed suggests that either p53 dissociates from the

TFIID/TFIIA/*bax* DNA co-complex or p53 has a transient interaction with the assembly, which

led to the EM density being averaged out during analysis. The same observation was also

obtained with the 3D structure of the TFIID/TFIIA/p53/*hdm2* promoter DNA (Supplemental

Figure S7). Since we previously determined the 3D structure of the p53 bound TFIID co-

complex without DNA (20), we attempted to capture the p53-bound state of TFIID in the cryo-

EM dataset of our TFIID/TFIIA/p53/*bax* DNA assembly. Thus, 3D reconstructions were

performed using the structure of p53/TFIID as an initial reference volume. A poor resolution 3D

reconstruction of p53/TFIID was obtained (Supplemental Figure S8), indicating that the majority

of TFIID complexes have switched to the p53-free DNA-bound state. Collectively, these

structural studies support the observations from our single molecule analyses, illustrating that

p53 dissociates from TFIID and DNA when recruited to target promoters. More importantly, our

work suggest that p53 facilitates the formation of a common rearranged DNA binding

conformation of TFIID to stimulate promoter recruitment on various target genes.

Discussion

p53 promotes distinct DNA binding modes within TFIID

p53-mediated binding of TFIID to promoter DNA is likely a multi-step process involving

the formation of multiple coordinate contacts between p53, the RE, TFIID and the individual

core promoter elements. Our single molecule analyses strongly suggest that p53/TFIID interacts

with target gene promoter DNA via two distinct DNA binding modes with a strong preference

for stable binding over transient interaction. Remarkably, TFIID's DNA binding mechanism

appears to be similar on the *hdm2* and *bax* DNA, despite different spatial arrangements and

sequences in the p53 RE and core promoter elements. It would be interesting to determine if

TFIID's residence time on DNA is dependent on the type of activator utilized to load TFIID onto

the promoter and how this correlates with the frequency of transcription initiation/re-initiation.

To potentially address how TFIID engages promoter DNA differentially, we studied how

these two populations of DNA-bound TFIID complexes responded to mutations in the core

promoter elements and the p53 RE. Intriguingly, the majority of DNA binding events on the

mutant core promoters were short-lived (Figure 3B). Residence times for both the long- and

short-lived TFIID complexes were also reduced. TFIID's decreased stability is likely due to

changes in contacts with the TATA, InR, and DPE on the Mu Core DNA as revealed in our

xlink-Exo assays (Figures 2D and S1). Strikingly, mutation of the p53 RE primarily affected the

residence time of TFIID's long-lived DNA binding population (Figure 3B). This finding suggest

that contacts between p53 and the RE are somehow required to increase the stability of TFIID on

promoter DNA via an unknown mechanism. Given that distinct TFIID contacts surrounding the

p53 RE were identified via our xlink-Exo assays (Figure 2D), we speculate that this upstream

DNA is involved in both binding p53 and stabilizing TFIID to the promoter. These upstream

contacts may be possibly mediated by the TAF4 subunit of TFIID, which was previously shown

to bind ~70 bp of DNA with a weak sequence preference (40).

Thus far, we surmise that TFIID/DNA binding occurs through at least two discernable

regulatory steps involving recognition of multiple different core promoter elements. The short-

lived DNA binding population could likely represent a transient non-productive TFIID complex,

which initially encounters the promoter DNA and is only partially engaged with the core

promoter elements (Figure 6). In such a scenario, interaction between p53 and its RE promotes

efficient isomerization within the initial TFIID/DNA encounter complex. This p53-mediated

isomerization step may possibly permit stable interaction of TFIID with the upstream DNA,

TATA box and downstream DNA resulting in long-lived DNA binding (Figure 6), which is

consistent with our xlink-Exo data (Figure 2D). In support of this model, previous biochemical

assays have shown that activators induce conformational changes within TFIID that enhance

multiple contacts with promoter sequences downstream of the TSS (10).

TFIID switches p53 to a factor-loading mode to "recycle" the p53 RE

In the absence of TFIID, p53 displays both specific and non-specific DNA binding modes

(i.e. average residence time = ~33" and ~7"). Remarkably, TFIID's presence switches p53 to a

single "factor-loading" mode with a residence time on DNA (~15") that is intermediate to its

specific (~33") and non-specific rates (~7") (Figure 4). A previous study documented that p53

binds non-specific DNA via select residues in its lysine rich C-terminal domain (CTD) (36).

Interestingly, TFIID's TAF1 bromodomains interact with acetylated marks on many of these

same residues in p53's CTD (41). Therefore, it is possible that TFIID's interaction could

eliminate p53's short-lived (7") population by sterically hindering its ability to non-specifically

bind DNA. Strikingly, deletion of the CTD also allosterically reduces p53's binding to specific

DNA targets (42, 43). Thus, we speculate that the interaction between TFIID and the CTD of

p53 would also explain the reduction in the long-lived residence time of p53 on specific DNA

sites. Overall, TFIID potentially alters p53's DNA binding dynamics by eliminating non-specific

DNA binding, reducing the affinity of p53 for specific target sites via contacts within its CTD,

and sterically blocking p53's contacts with specific DNA residues via direct TAF contacts within

the RE.

Our results show that TFIID's DNA binding induces dismounting of p53 from the

promoter, consistent with previous DNAse I footprinting studies showing decreased binding of

p53 to REs in the presence of TFIID (19). While p53 dissociates from the TFIID/DNA scaffold,

it is highly likely that p53 can rapidly re-associate with the complex, particularly at the high

concentrations (600nM) used in Figure 1B. We suspect that this rapid removal could

consequently lead to a faster "recycling" of p53 from the RE for its subsequent recruitment of

other basal transcription factors such as TFIIB (44, 45) and TFIIH (46-48). In this scenario,

stable prolonged binding of p53 to a TFIID/DNA complex would negatively regulate PIC

formation, because p53 would occupy the RE and inhibit subsequent recruitment of a p53/basal

factor co-complex. Notably, the residence times for p53 and TFIID at the promoter were

measured in the absence of additional PIC components. Most likely, additional factors such as

TFIIA, TFIIF, and RNA Polymerase II will alter the stability of TFIID and p53 as the PIC forms

on promoter DNA. Indeed, our previous single molecule studies showed prolonged binding of

QD-TFIID to DNA in the presence of PIC components (21). This suggests that our current work

here measures TFIID dissociation events and not simply dissociation of the QD labeled TAF1

antibody. Overall, our studies indicate that p53, and potentially activators in general, serve as

escorts to dynamically recruit and load the basal transcription machinery onto DNA via complex

interactions between the RE, the core promoter and bound basal factors (i.e. TFIID).

p53 facilitates formation of a common rearranged DNA binding framework of TFIID

Our previous structural studies demonstrated that, in the absence of DNA, p53 bridges

TFIID's Lobe A/C while introducing movement of Lobe A towards Lobe B (20). Interestingly,

repositioning of Lobe A to Lobe B is a hallmark feature associated with a rearranged form of

TFIID that binds promoter DNA (24). Here we found that p53 and DNA enhances formation of

the rearranged TFIID via 2D classification analysis (Figure 5A). In addition, the majority of our

TFIID assemblies in the cryo-EM datasets display this rearranged DNA binding form (Figures 5

and S6). Therefore, these results strongly suggest that the p53-induced specific movement of

Lobe A within TFIID likely helps formation of a TFIID/DNA co-complex by shortening the

distance required for repositioning of lobe A to lobe B (24). In addition, we have determined that

the stable TFIID/DNA interaction does not require p53 to remain bound to TFIID and DNA

(Figures 4, 5 and S6). It is thereby reasonable to propose that p53 stimulates the transition of

canonical TFIID to the rearranged DNA binding state, followed by dissociation of p53 once

TFIID stably loads onto DNA (Figure 6).

Our single molecule studies indicate that p53 promotes both transient and stable binding

of TFIID to DNA. Due to this transient dynamic interaction behavior, we were likely unable to

obtain the 3D structure of those short-lived DNA-bound TFIID complexes via cryo-EM. It is

possible that these short-lived DNA-bound TFIID complexes could be a unstable "encounter

complex" (Figure 6) or reminiscent of the "recruitment" form of DNA-bound yeast TFIID

induced by the Rap1 activator in the absence of TFIIA (23). We speculate that the long-lived

DNA-bound TFIID complex induced by p53 is structurally related to the stable

TFIID/TFIIA/promoter DNA complex depicted in our cryo-EM studies.

Our cryo-EM analysis uncovers the global similarity of the 3D reconstructions between

p53/TFIID/TFIIA/*bax* DNA, p53/TFIID/TFIIA/*hdm2* DNA and TFIID/TFIIA/SCP DNA

(Figures 5B, S6 and S7). This significant observation in combination with our 2D classification

analysis (Figure 5A) clearly suggests that p53 aids in formation of a common DNA binding

mode of TFIID, which effectively allows TFIID to bind diverse target gene promoters.

Interestingly, p53 alone enhances the rearranged form of TFIID (Figure 5A), while TFIIA alone

promotes the canonical form (24). These observations indicate that different transcription factors

use distinct modes of structural regulation to stimulate TFIID recruitment. Nevertheless, it is

difficult to clearly define the individual roles of p53 and TFIIA in our 3D structures, because

each factor plus DNA is capable of promoting TFIID's rearranged DNA binding conformation

(this study and (24)).

Intriguingly, the proximal promoter DNA upstream of the TATA box is highly mobile

when TFIID binds to p53 target gene promoters as revealed in our 2D class average analysis

(Figure 5B). Notably, our sensitive xlink-exo assays were able to detect TFIID contacts with

upstream sequences (Figure 2D). These results suggest a dynamic transient interaction between

TFIID and upstream DNA. Since a number of activator binding sites reside in a promoter

proximal region upstream of the TATA box (49), our findings indicate the potential for TFIID to

also influence binding of other activators with their cognate sites as observed previously (22).

Collectively, our work reveals the molecular mechanism pertaining to how p53 stimulates TFIID

binding to different promoters that comprise variable arrangements of p53 binding sites and core

promoter elements. More importantly, our combined functional and structural studies are crucial

for understanding how eukaryotic transcription complexes dynamically assemble on different

protein-coding gene promoters.

501
502

Materials and Methods

Reagents and protein purification

Experimental details, including reagents, protein purification of the p53/TFIID/TFIIA/DNA

co-complexes and fluorescent-labeled proteins/DNA used in this study along with our custom-

built TIRF microscope setup, are described in the Supplemental Materials.

In vitro immuno-assembly assay

HeLa cells (32 liters) were grown in suspension culture with 1X DMEM plus 5% newborn calf

serum for each assay. Nuclear extract was prepared and fractionated with phosphocellulose P11

(P-Cell) resins exactly as previously described (20). P-Cell column fractions eluting at 1M

KCl/HEMG buffer (pH 7.9, [20 mM Hepes, 0.2 mM EDTA, 2 mM $MgCl_2$, 10% Glycerol, 1 mM

DTT and 0.5 mM PMSF]) were pooled, dialyzed to 0.3 M KCl/HEMG buffer, added with final

concentration of 0.1% NP40, and then immunoprecipitated overnight at 4°C with an anti-TAF4

mAb covalently conjugated to Protein G Sepharose beads (GE Healthcare Life Sciences). TAF4-

immunoprecipitates were extensively washed with 0.65 M KCl/HEMG buffer, 0.3 M

KCl/HEMG buffer and 0.1M KCl/HEMG buffer (containing 0.05% NP40 buffer and 10 μM

Leupeptin) and split into four assembly reactions as indicated in Figure 1. Three reactions of

TFIID-bound resins were incubated with 6-fold molar excess of promoter DNA. Next, two

TFIID/DNA reactions were further added with 10-fold molar excess of p53 alone and/or TFIIA.

Total four reactions were incubated at 4°C for 2 hours prior to elution of the co-complex from

the TAF4 mAb affinity resin. TFIID/factor/DNA-bound resins were washed 5 times with 0.1 M

KCl/HEMG buffer (containing 0.1% NP40 and 10 μM Leupeptin) followed by elution with a

peptide (1 mg/ml) in 0.1 M KCl/HEMG buffer (plus 0.1% NP40 and 10 μM Leupeptin)

recognized by the TAF4 mAb. Eluates were analyzed with 4-12% SDS-PAGE (NuPage gel

system, Thermo Fisher) and visualized with silver staining analysis.

Xlink-Exo assay

Xlink-Exonuclease assays were carried out in 10 µl reactions containing a modified 50 mM

KCl/HEMG buffer (pH 7.9, [12.5 mM Hepes, 0.05 mM EDTA, 6.25 mM $MgCl_2$, 5% glycerol]

plus 1% DTT, 0.05 M PMSF, and 0.01% NP-40), IRdye800 (IDT) antisense strand labeled *hdm2*

promoter DNA (9nM) , p53 (90nM, monomer), and TFIID (4nM). After incubation at room

temperature for 20', reactions were crosslinked with formaldehyde (1% final) for 5' and stopped

by addition of Tris-HCl (100 mM final, pH 8.0). Next, each reaction was mixed with

Exonuclease III (50 Units, NEB diluted in a buffer containing $MgCl_2$ [16.25 mM final]), and

incubated at $37^{o}C$ for 1 hour. Proteinase K (10 µg) was then added into each reaction and

incubated at $70^{o}C$ for 3 hours. Reactions were mixed in formamide loading dye, heated at $95^{o}C$

and resolved on a 8% Polyacrylamide-TBE gel containing 7M urea. A series of known 5'

truncated IRdye800 (IDT) antisense strand labeled wild-type *hdm2* DNA fragments were

generated and used as molecular standards to determine the size of Exonuclease III digested

bands seen on our mapping gels.

TIRF Single Molecule Imaging

Passivated glass coverslips (see Supplemental materials) were assembled into a multichannel

flow cell using double sided tape (3M). Flow cells were mounted on the microscope and

streptavidin was added to the flow cell to coat surfaces. Each flow cell holds ~50 ul of sample.

Reference beads that serve as fiducial marks were immobilized on the bottom surface of the flow

cell. Biotinylated Cy3/Cy5 promoter DNA in 1X PBS was then loaded into the flow cell and

bound over time specifically to the streptavidin that was non-covalently attached through biotin

linkages on the top and bottom surfaces. Promoter DNA templates were briefly imaged at 3HZ to

determine surface attached DNA densities that were sufficient for localization of single DNA

molecules. Unbound biotinylated Cy3/Cy5 promoter DNA was washed away. Promoter DNA

templates were then imaged at 3HZ. The surface attached Cy3/Cy5 promoter DNA was then

photobleached. Reactivation of the Cy3/Cy5-labeled promoter DNA was not evident over the

timescale of our experiments (typically 15-20 minutes). Qdot 565-labeled TFIID (1 nM) in the

absence and presence of unlabeled p53 (8 nM, monomer) was then added to the flow cell in a

modified 50 mM KCl/HEMG buffer (pH 7.9, [12.5 mM Hepes, 0.05 mM EDTA, 6.25 mM

MgCl$_2$, 5% glycerol] plus 1% DTT, 0.05 M PMSF, and 0.01% NP-40) and immediately imaged

at 3Hz at 23^{o}C. An oxygen scavenging system consisting of 0.4% b-D-Glucose, 1 mg/ml glucose

oxidase (Sigma), and 0.4 mg/ml catalase (Sigma) was used for imaging. For experiments to

measure p53's interaction with promoter DNA, AlexaFluor 555, Cy3- or Cy5-labeled p53 (8nM,

monomer) in the absence or presence of unlabeled TFIID (1 nM) was added to the flow cell

containing surfaced attached mapped promoter DNA and immediately imaged at 3Hz.

AlexaFluor 555, Cy3- and Cy5-labeled p53 had equivalent residence times on promoter DNA

indicating that photobleaching of molecules didn't significantly affect results.

567
Single Molecule Co-localization Analysis

Fluorescent spots in each frame for the movie of the promoter DNA, TFIID or p53 were mapped

using 2D Gaussian fitting methods (21, 50, 51) to determine the location (x, y) at subpixel

resolution. Co-localization analysis was performed as in (21) with the following modifications.

Coordinates of fluorescent spots were adjusted based upon movement of the fiducial marks (i.e.

reference beads) over time. Stage movement/drift was typically less than 200 nm over the time of

imaging (~20 minutes). Fluorescent spots that were located within 84 nm (1 pixel) throughout

the movie were grouped together in a cluster. The positions of all spots within a cluster were

averaged to obtain the overall location of an individual cluster of molecules. Position maps of the

clustered surface attached DNA molecules were then compared with position maps of either the

TFIID or p53 clusters. Clusters of TFIID or p53 molecules within 3 pixels (252 nm) of a DNA

cluster were found. Displacements of TFIID or p53 clustered molecules with a maximum equal

to (0,0) in the x and y positions are indicative of non-random co-localization between DNA and

TFIID or p53 clusters. Displacements between the x,y positions of the TFIID or p53 clusters and

the DNA clusters were then plotted as a 2D histogram with a bin size of 50 nm. 2D Gaussian

fitting of the 2D histogram yields σ which is equivalent to the variation in the displacement of

TFIID or p53 within 252 nm of a DNA molecule. A co-localization threshold was set to 1.8σ (84

nm).

Single Molecule Residence Time Analysis of TFIID and p53 on promoter DNA

For each TFIID or p53 cluster that co-localized with a DNA cluster (i.e. within 84 nm), we the

plotted the amplitude from the 2D Gaussian fit of every fluorescent spot within the TFIID or p53

cluster as a function of time. Homemade MATLAB software was then used to analyze the rise

and the fall of the amplitude of the fluorescent signal to determine the appearance and

corresponding disappearance of each spot over time. The residence time for each DNA binding

event is defined as the difference in time between the appearance and disappearance of a TFIID

or p53 fluorescent spot. The fluorescent signal from TFIID or p53 had to last for at least 1.5" to

be considered as a discrete binding event in our analysis. Due to rapid stochastic blinking of the

quantum dot, a dissociation event was only considered to have occurred if the quantum dot signal

remained in the off state longer than 4.5". At least 300 binding events of co-localized TFIID or

100 binding events of p53 clusters were analyzed for each particular promoter DNA construct.

For a subset of molecules (~100), the data was processed by hand to determine the appearance

and disappearance of fluorescent spots. Comparison of the residence times generated by our

automated software and by hand yielded negligible differences. Histograms of residence times

for TFIID or p53 bound to promoter DNA were fit to a two-component exponential decay model

to determine the average residence time of TFIID or p53 bound to promoter DNA (51). If the

difference between the fitted residence times for each individual component in the two-

component model varied less than 20%, histograms were re-fitted using a single-component

exponential decay model (51). For Figures 4 and S3 data was additionally fit to a gamma

distribution (38) via custom MATLAB software.

2D Classification Analysis of TFIID co-complexes via negative stain EM

4 µL (10-20 ng in total amount) of fresh TFIID/p53/bax DNA assemblies in 0.1M

KCl/HEMG buffer (20mM HEPES, 0.2 mM EDTA, 2mM $MgCl_2$, 10% Glucose [final pH 7.9])

was applied directly onto a thin carbon film supported by holey carbon on a 400-mesh copper

grid (Pacific Grid tech.) for 3', which was freshly glow-discharged. After incubating the sample

on the grid for 3', the sample grid was stained in five successive 75 µL drops of 1% Uranyl

Formate.

The negative stain image data were collected with a Tecnai F20 TWIN transmission

electron microscope operating at 120 keV at a calibrated magnification of 62,000x with a

defocus range of -0.5µm to -2.5µm using a Tietz F416 (4K x 4K) CCD camera (resulting in a

2.82 A/pixel) under the dosage 39.98 e$^-$/Å. A total of 14,000 particles (for TFIID alone), 13,855

particles (for TFIID/p53), and 19262 particles (for TFIID/p53/bax DNA) were manually picked

using Boxer (EMAN; (52)). These particles were aligned and classified in a reference-free

fashion as previously described (22) for 8 iterations. The similar measurement of Lobe's A

position relative to the lobes B and C was employed as in (24). Based upon the position of Lobe

A in relation to Lobe C v.s. Lobe B (24), three conformational groups of TFIID were defined (i.e.

canonical, transition, and rearranged state) in this study. The canonical form is represented by

Lobe A retaining its attachment to Lobe C. The transition state is represented by Lobe A

beginning to detach and move in between Lobe C and B. The rearranged form is represented by

Lobe A reattaching to Lobe B. Next, individual classums from the 5th , 6th, 7th and 8th iteration

for each TFIID/p53 prep with or without the *bax* DNA were examined carefully to select 2D

class averages displaying views of these three conformational groups. The number of 2D

averages in each conformational form v.s. total averages was calculated to yield the percentage

of TFIID's three conformations in each TFIID/p53 prep with or without the *bax* DNA. Three

independent protein purifications and EM data acquisition/analysis were performed to obtain the

statistically significant standard deviation for the figure.

Cryo-EM data collection and image analysis

For cryo-EM sample grid preparation, a thin carbon film supported by a 400-mesh carbon-

thickened C-flat holey grid with hole diameter 1.2 μm/spacing 1.3 μm (CF-1.2/1.3-4C,

Protochips) was plasma cleaned for 2 min using Denton high-vacuum evaporator. The grid was

loaded into a Vitrobot (FEI) that was preset at 100% humidity at 4ºC for vitrification of our

samples. 3 μl (504~900 ng in total amount) of the p53/TFIID/TFIIA/DNA assemblies was

applied directly onto the grid, incubated for 2 min and then blotted for 6.5 sec prior to plunge-

frozen in liquid ethane.

Cryo-EM data were collected with a JEM-2100F transmission electron microscope (JEOL)

operated at 120 KeV and at a magnification of 50,000x with a defocus range of –2.5 μm to –4.5

μm. Digital micrographs were collected using a TEM 2048x 2048 CCD camera (TemCam-

F224HD, TVIPS) with a pixel size of 24 μm and a calibrated magnification of 84,037x (2.86

Å/pixel) under low-dose conditions (~13 e$^-$ /Å) via the semi-automatic image acquisition

SerialEM software (53). Digital micrographs showed homogeneous particles in terms of lack of

aggregation and apparent integrity of the intact complexes.

Particles were manually selected using boxer (EMAN) (52) for an initial total of 32,275

particles from the p53/TFIID/TFIIA/*bax* DNA dataset and 22,070 particles from the

p53/TFIID/TFIIA/*hdm2* DNA dataset. The particles were then phase flipped using CTF-

estimated values determined by CTFFIND3(54) and extracted using SPIDER (System for

Processing of Image Data from Electron microscopy and Related fields (55)) to a particle

window size of 128 x 128 pixels (5.72 Å/pixel), following the protocol as described in (24). The

particles were then normalized prior to the following data analyses. Reference-free 2D class

averaging image analysis was performed using IMAGIC (56) following the procedure precisely

as described (20). Using these 2D class averages as templates, we automatically picked particles

from a larger cryo-EM dataset using Signature (57) embedded in the Appion image processing

platform (58). A final total of 59,632 particles for the p53/TFIID/TFIIA/*bax* DNA co-complex

was obtained, normalized, and low-pass band filtered to perform the 3D reconstruction using

SPIDER's multi-reference projection matching approach as previously described (27). Both

rearranged and canonical forms of TFIID/TFIIA/SCP DNA from Eletron Microscopy Data Bank

(accession number 10832 and 10834, respectively) were filtered to 60 Å as a reference for the

first cycle of projection matching. Different initial reference models including TFIID alone, p53-

bound TFIID or TFIIA-bound TFIID were also tested for model bias refinement. New 3D

volumes reconstructed from the data set were used as references for all subsequent cycles of

alignment (angular step sizes ranging from an initial 20° to a final 2°). The resolution throughout

the refinement was determined by the 0.5 cut-off in the Fourier shell correlation (FSC) curve.

The 3D structure of the p53/TFIID/TFIIA/bax assembly was filtered at a final resolution ~35 Å.

All the 3D reconstructions were represented as isodensity surfaces using the UCSF Chimera

package (59) developed by the Resource for Biocomputing, Visualization, and Informatics at the

University of California, San Francisco (supported by NIH NIGMS P41-GM103311).

675
676

Funding information

This study was supported by Einstein start up fund (Albert College of Medicine). Some of this work was performed at the Simons Electron Microscopy Center at the New York Structural Biology Center that is supported by a grant from the Simons Foundation (Grant Number: 349247) with additional support from: NIH S10 OD019994-01, the Agouron Institute (Grant Number: F00316), NIH S10 RR029300-01, NIH S10 RR017291-01, NYSTAR, and NIH C06 RR017528-01-CEM.

Acknowledgement

We specifically thank R. Tjian, S. Chu, and E. Nogales for their initial support of this study, S. Zheng for generating TAF4 mAb supernatant and HeLa cells, D. King for providing the TAF4 mAb elution peptide, A. Revyakin, A. Pertsinidis and S.R. Park for their help on initial stages of the single molecule microscopy, along with R. Henderson and D. Stokes for their technical advice critical for the cryo-EM work. We also thank S.M. Shenoy, J. Hargitai, J. Wang, S. Lenny, J. Greally and E. Eng for high performance computing cluster and cryo-EM technical support. We appreciate assistance from Einstein's AIF facility, especially F. P. Macaluso, L. Cummins and G. S. Perumal. We are grateful to M. Keogh and E. Nogales for critical comments of the manuscript. W. Liu is an affiliated member of the New York Structural Biology Center. M. Cianfrocco is an HHMI fellow of the Damon Runyon Cancer Research Foundation.

Figure legends

Figure 1. In vitro assembly of p53/TFIID/TFIIA/promoter DNA complexes. (A) Schematic

representation of procedures for assembly assays to study TFIID bound to p53, TFIIA and native

p53 target gene promoter DNAs in vitro. TFIID was immunoprecipiated from the

phosphocellulose 1M KCl fraction using a highly specific anti-TAF4 antibody. After extensive

washing, buffer alone (ctl) or a 6-fold molar excess amount of the promoter DNA (i.e. *bax* or

hdm2 DNA) along with either p53 or p53 plus TFIIA was loaded onto the resins containing

TFIID. TFIID co-complexes were obtained after extensive washes followed by an elution step

using a specific peptide against TAF4. (B) Schematic representation of the *bax* and *hdm2*

promoter DNA templates used in the assembly reactions. Relative positions of p53 response

elements (RE), the core promoter element TATA box, and transcription start site (TSS) within

both p53 target gene promoters are indicated respectively. Eluates of TFIID co-complexes

obtained from the assembly reactions as described in (A) were analyzed by 4-12% SDS-PAGE

(Invitrogen) and visualized by silver staining analysis.

712
Figure 2. Single Molecule <u>T</u>otal <u>I</u>nternal <u>R</u>eflection <u>F</u>luorescence (TIRF) Microscopy

analysis of Quantum dot-labeled TFIID binding of promoter DNA in vitro. (A) Schematic

representation of an in vitro single molecule TIRF assay to monitor binding of quantum dot

(QD)-labeled TFIID to fluorescently labeled *bax* or *hdm2* promoter DNA in real-time. (B) Co-

localization analysis of Cy3/Cy5-labeled promoter DNA and QD-labeled TFIID. Representative

2D projection maps of single promoter DNA molecules (red) and single TFIID complexes

(green) are shown. The intensity of the green spot reflects the time TFIID binds that specific

location on the slide. Transient binding (~1-2") results in a dim green spot. A co-localization

threshold of 84 nm was set based upon our previous study (21). Yellow boxes represent co-

localized molecules. The numbers below the projection maps represent the percentage of DNA

molecules co-localizing with TFIID (i.e. number of DNA molecules co-localizing with

TFIID/Total number of DNA molecules in a field of view). Due to the minute co-localization

efficiency (1.3%) in the absence of p53, higher DNA densities were necessary to yield a

statistically significant number of binding events. 2D histograms of TFIID–promoter DNA

displacement define the number of TFIID complexes mapped within 200 nm of a DNA molecule

in a particular field of view. σ equals the standard deviation (in nm) of the 2D Gaussian fits of

the TFIID–promoter DNA displacement histograms. N_{DNA}= Total number of DNA molecules in

a field of view. (C) Dependence of TFIID/promoter DNA co-localization on p53, the p53 RE,

and core promoter elements. Promoter DNA binding is plotted as a percentage of total DNA

molecules in a field of view that are co-localized with TFIID over 15' of binding based on

TFIID–promoter DNA displacement histograms. TFIID binding to wild-type *bax* DNA is

compared with and without p53. Additional comparisons were made for TFIID binding to wild-

type (WT), Mutant p53 RE (Mu RE), Mutant Core (Mu Core), or combined mutant RE and core

(Mu RE/Co) *hdm2* DNA templates. (D) xlink-Exo analysis of TFIID binding to wild-type (top

panel) and Mu RE/Co (bottom panel) *hdm2* DNA. Basepair annotations (+/- 2bp) above

individual peaks in the line traces (right panel) represent a -4bp adjustment to reflect the known

inability of Exonuclease III to cut immediately adjacent to protein/DNA crosslink sites (32).

740
Figure 3. Residence time analysis of TFIID bound to promoter DNA. (A) Individual co-

localized TFIID/promoter DNA binding events were further analyzed to determine the average

residence time for TFIID bound to wild-type *hdm2* promoter DNA in the absence (left panel) and

presence (right panel) of p53. 1–Cumulative Distribution Function Plots (1–CDF) of TFIID

bound to DNA were fitted to a single (gray dashed) or two-component (red solid) exponential

decay model. Fitting analysis reveals that TFIID complexes bound to promoter DNA display

residence times comprised of a long- (τ1) and a short- (τ2) lived component. Percentages of the

long- (τ1) and short- (τ2) lived components are listed next to the residence time. (B) Average

residence time analysis for TFIID bound to Mu RE(left panel), Mu Core (middle panel), and Mu

RE/Co (right panel) *hdm2* promoter DNA in the presence of p53 as described in (A).

Figure 4. Residence time analysis of p53 bound to promoter DNA with and without TFIID.

(A) Individual co-localized p53/promoter DNA binding events were further analyzed to

determine the average residence time for p53 bound to wild-type (left panel) and Mutant p53 RE

(right panel) *hdm2* promoter DNA. 1−Cumulative Distribution Function Plots (1−CDF) of p53

bound to promoter DNA were fit to a single (gray dashed) or two (red solid)-component

exponential decay model. Based on comparisons of the *hdm2* promoter DNAs containing wild-

type and mutant p53 REs, τ1 represents specific binding of p53 to the wild-type *hdm2* DNA

while τns equals non-specific interactions between p53 and promoter DNA. (B) 1−Cumulative

Distribution Function Plot (1−CDF) of p53 bound to the wild-type *hdm2* DNA template in the

presence of TFIID was fit to a single (gray dashed) and two (red solid)-component exponential

decay (left panel) or gamma distribution (right panel) model. (C) Comparison of p53's residence

time (τ1) on the wild-type *hdm2* DNA with and without TFIID based upon the results shown in

(A) and (B) (left panel). Comparison of TFIID's and p53's residence time (τ1) on the wild-type

hdm2 DNA based upon results shown in Figures 3 and 4.

Figure 5. Structural EM analysis on p53's impact on TFIID bound to native target gene

promoters. (A) TFIID's structural plasticity as defined by the position of its Lobe A (24)(top

panels, the major lobes of TFIID are labeled as A, B, and C). TFIID's conformations are

classified into three groups: Canonical, Transition and Rearranged state as shown. Reference-

free 2D classification analysis on negative stain EM datasets showing populations of different

TFIID conformations in the absence of p53 (gray), presence of p53 (white), and with p53 + *bax*

DNA (black). The scale bar represents 100Å. (B) Three representative reference-free 2D class

averages of p53/TFIID/TFIIA bound to either the *bax* or *hdm2* DNA are shown based upon cryo-

EM particle datasets. The scale bar is 200Å. (C) Distinct 2D class averages showing views of the

assemblies bound to the *bax* promoter DNA fragment downstream (top panel, DNA highlighted

in green arrows) and upstream (middle panel, DNA highlighted in yellow arrows) of the

transcription start site (TSS, +1) are presented respectively. The orientation of the core promoter

elements bound by TFIID were previously mapped in the 3D structure of TFIID/TFIIA bound to

the super-core promoter (SCP) DNA (24). The corresponding views of the 3D structure of

TFIID/TFIIA/SCP DNA are shown (far right) to indicate the DNA orientation. The strong 3D

density highlighted in green represents the promoter DNA comprising the initiator, TSS and

downstream promoter elements. Since the upstream promoter DNA was weakly visible, focused

classification analysis (bottom panel) were performed using those corresponding class averages

(middle panel) to clarify the presence of the *bax* promoter DNA upstream of the TATA box

(highlighted by yellow arrow). The scale bar is 200Å.

Figure 6.   A representative model of how p53 induces a common DNA binding mode

within TFIID that targets various p53-responsive promoters.

A proposed model of how p53 facilitates a common structural module within TFIID for core

promoter recognition is presented based upon our current and previous studies (20).   In the

absence of DNA, p53 binds to lobe A/C, which results in a shortening of the distance between

lobes A and B in addition to an extension of lobe D (20). This structural repositioning of lobe A

to lobe B results in the formation of a common DNA-binding mode within TFIID that allows

recognition of various p53 target promoters. Once TFIID structurally rearranges, p53 disengages

via loss of its contacts with Lobe A/C to allow TFIID to stably bind the promoter. Dissociation

of p53 from the TFIID/promoter DNA scaffold results in "recycling" of the p53 RE for further

assembly of the PIC components. The scale bar represents 100Å.

References

1.   **Menendez D, Inga A, Resnick MA.** 2009. The expanding universe of p53 targets. Nat
803   Rev Cancer **9:**724-737.
804   2.   **Bieging KT, Mello SS, Attardi LD.** 2014. Unravelling mechanisms of p53-mediated
805   tumour suppression. Nat Rev Cancer **14:**359-370.
806   3.   **Tafvizi A, Huang F, Fersht AR, Mirny LA, van Oijen AM.** 2011. A single-molecule
807   characterization of p53 search on DNA. Proc Natl Acad Sci U S A **108:**563-568.
808   4.   **Lu H, Levine AJ.** 1995. Human TAFII31 protein is a transcriptional coactivator of the
809   p53 protein. Proc Natl Acad Sci U S A **92:**5154-5158.
810   5.   **Thut CJ, Chen JL, Klemm R, Tjian R.** 1995. p53 transcriptional activation mediated
811   by coactivators TAFII40 and TAFII60. Science **267:**100-104.
812   6.   **Buschmann T, Lin Y, Aithmitti N, Fuchs SY, Lu H, Resnick-Silverman L, Manfredi
813   JJ, Ronai Z, Wu X.** 2001. Stabilization and activation of p53 by the coactivator protein
814   TAFII31. J Biol Chem **276:**13852-13857.
815   7.   **Li AG, Piluso LG, Cai X, Gadd BJ, Ladurner AG, Liu X.** 2007. An acetylation switch
816   in p53 mediates holo-TFIID recruitment. Mol Cell **28:**408-421.
817   8.   **Juven-Gershon T, Hsu JY, Kadonaga JT.** 2006. Perspectives on the RNA polymerase
818   II core promoter. Biochem Soc Trans **34:**1047-1050.
819   9.   **Colgan J, Manley JL.** 1992. TFIID can be rate limiting in vivo for TATA-containing,
820   but not TATA-lacking, RNA polymerase II promoters. Genes Dev **6:**304-315.
821   10.   **Chi T, Carey M.** 1996. Assembly of the isomerized TFIIA--TFIID--TATA ternary
822   complex is necessary and sufficient for gene activation. Genes Dev **10:**2540-2550.
823   11.   **Wang W, Gralla JD, Carey M.** 1992. The acidic activator GAL4-AH can stimulate
824   polymerase II transcription by promoting assembly of a closed complex requiring TFIID
825   and TFIIA. Genes Dev **6:**1716-1727.
826   12.   **White J, Brou C, Wu J, Lutz Y, Moncollin V, Chambon P.** 1992. The acidic
827   transcriptional activator GAL-VP16 acts on preformed template-committed complexes.
828   Embo J **11:**2229-2240.
829   13.   **Chi T, Carey M.** 1993. The ZEBRA activation domain: modular organization and
830   mechanism of action. Mol Cell Biol **13:**7045-7055.
831   14.   **Lieberman PM, Berk AJ.** 1994. A mechanism for TAFs in transcriptional activation:
832   activation domain enhancement of TFIID-TFIIA--promoter DNA complex formation.
833   Genes Dev **8:**995-1006.
834   15.   **Chi T, Lieberman P, Ellwood K, Carey M.** 1995. A general mechanism for
835   transcriptional synergy by eukaryotic activators. Nature **377:**254-257.
836   16.   **Kobayashi N, Boyer TG, Berk AJ.** 1995. A class of activation domains interacts
837   directly with TFIIA and stimulates TFIIA-TFIID-promoter complex assembly. Mol Cell
838   Biol **15:**6465-6473.
839   17.   **Rhee HS, Pugh BF.** 2012. Genome-wide structure and organization of eukaryotic pre-
840   initiation complexes. Nature **483:**295-301.
841   18.   **Lieberman PM, Ozer J, Gursel DB.** 1997. Requirement for transcription factor IIA
842   (TFIIA)-TFIID recruitment by an activator depends on promoter structure and template
843   competition. Mol Cell Biol **17:**6624-6632.

19. **Xing J, Sheppard HM, Corneillie SI, Liu X.** 2001. p53 Stimulates TFIID-TFIIA-
promoter complex assembly, and p53-T antigen complex inhibits TATA binding protein-
TATA interaction. Mol Cell Biol **21:**3652-3661.
20. **Liu WL, Coleman RA, Ma E, Grob P, Yang JL, Zhang Y, Dailey G, Nogales E,**
Tjian R. 2009. Structures of three distinct activator-TFIID complexes. Genes Dev
23:1510-1521.
21. **Revyakin A, Zhang Z, Coleman RA, Li Y, Inouye C, Lucas JK, Park SR, Chu S,**
Tjian R. 2012. Transcription initiation by human RNA polymerase II visualized at
single-molecule resolution. Genes Dev **26:**1691-1702.
22. **Liu WL, Coleman RA, Grob P, King DS, Florens L, Washburn MP, Geles KG,**
Yang JL, Ramey V, Nogales E, Tjian R. 2008. Structural changes in TAF4b-TFIID
correlate with promoter selectivity. Mol Cell **29:**81-91.
23. **Papai G, Tripathi MK, Ruhlmann C, Layer JH, Weil PA, Schultz P.** 2010. TFIIA
and the transactivator Rap1 cooperate to commit TFIID for transcription initiation.
Nature **465:**956-960.
24. **Cianfrocco MA, Kassavetis GA, Grob P, Fang J, Juven-Gershon T, Kadonaga JT,**
Nogales E. 2013. Human TFIID binds to core promoter DNA in a reorganized structural
state. Cell **152:**120-131.
25. **Louder RK, He Y, Lopez-Blanco JR, Fang J, Chacon P, Nogales E.** 2016. Structure
of promoter-bound TFIID and model of human pre-initiation complex assembly. Nature
531:604-609.
26. **Andel F, 3rd, Ladurner AG, Inouye C, Tjian R, Nogales E.** 1999. Three-dimensional
structure of the human TFIID-IIA-IIB complex. Science **286:**2153-2156.
27. **Grob P, Cruse MJ, Inouye C, Peris M, Penczek PA, Tjian R, Nogales E.** 2006. Cryo-
electron microscopy studies of human TFIID: conformational breathing in the integration
of gene regulatory cues. Structure **14:**511-520.
28. **Miyashita T, Reed JC.** 1995. Tumor suppressor p53 is a direct transcriptional activator
of the human bax gene. Cell **80:**293-299.
29. **Manfredi JJ.** 2010. The Mdm2-p53 relationship evolves: Mdm2 swings both ways as an
oncogene and a tumor suppressor. Genes Dev **24:**1580-1589.
30. **Bourdon JC, Deguin-Chambon V, Lelong JC, Dessen P, May P, Debuire B, May E.**
1997. Further characterisation of the p53 responsive element--identification of new
candidate genes for trans-activation by p53. Oncogene **14:**85-94.
31. **Juven-Gershon T, Cheng S, Kadonaga JT.** 2006. Rational design of a super core
promoter that enhances gene expression. Nat Methods **3:**917-922.
32. **Chang GS, Chen XA, Park B, Rhee HS, Li P, Han KH, Mishra T, Chan-Salis KY,**
Li Y, Hardison RC, Wang Y, Pugh BF. 2014. A comprehensive and high-resolution
genome-wide response of p53 to stress. Cell Rep **8:**514-527.
33. **Galasinski SK, Lively TN, Grebe De Barron A, Goodrich JA.** 2000. Acetyl coenzyme
A stimulates RNA polymerase II transcription and promoter binding by transcription
factor IID in the absence of histones. Mol Cell Biol **20:**1923-1930.
34. **Zhang Z, Boskovic Z, Hussain MM, Hu W, Inouye C, Kim HJ, Abole AK, Doud**
MK, Lewis TA, Koehler AN, Schreiber SL, Tjian R. 2015. Chemical perturbation of
an intrinsically disordered region of TFIID distinguishes two modes of transcription
initiation. Elife **4**.

35. **Mazur SJ, Sakaguchi K, Appella E, Wang XW, Harris CC, Bohr VA.** 1999.
Preferential binding of tumor suppressor p53 to positively or negatively supercoiled DNA
involves the C-terminal domain. J Mol Biol **292:**241-249.
36. **Weinberg RL, Freund SM, Veprintsev DB, Bycroft M, Fersht AR.** 2004. Regulation
of DNA binding of p53 by its C-terminal domain. J Mol Biol **342:**801-811.
37. **Emamzadah S, Tropia L, Vincenti I, Falquet B, Halazonetis TD.** 2014. Reversal of
the DNA-binding-induced loop L1 conformational switch in an engineered human p53
protein. J Mol Biol **426:**936-944.
38. **Floyd DL, Harrison SC, van Oijen AM.** 2010. Analysis of kinetic intermediates in
single-particle dwell-time distributions. Biophys J **99:**360-366.
39. **Phelps M, Darley M, Primrose JN, Blaydes JP.** 2003. p53-independent activation of
the hdm2-P2 promoter through multiple transcription factor response elements results in
elevated hdm2 expression in estrogen receptor alpha-positive breast cancer cells. Cancer
Res **63:**2616-2623.
40. **Gazit K, Moshonov S, Elfakess R, Sharon M, Mengus G, Davidson I, Dikstein R.**
2009. TAF4/4b x TAF12 displays a unique mode of DNA binding and is required for
core promoter function of a subset of genes. J Biol Chem **284:**26286-26296.
41. **Li AG, Piluso LG, Cai X, Gadd BJ, Ladurner AG, Liu X.** 2007. An acetylation switch
in p53 mediates holo-TFIID recruitment. Mol Cell **28:**408-421.
42. **Espinosa JM, Emerson BM.** 2001. Transcriptional regulation by p53 through intrinsic
DNA/chromatin binding and site-directed cofactor recruitment. Mol Cell **8:**57-69.
43. **Laptenko O, Shiff I, Freed-Pastor W, Zupnick A, Mattia M, Freulich E, Shamir I,**
Kadouri N, Kahan T, Manfredi J, Simon I, Prives C. 2015. The p53 C terminus
controls site-specific DNA binding and promotes structural changes within the central
DNA binding domain. Mol Cell **57:**1034-1046.
44. **Liu X, Berk AJ.** 1995. Reversal of in vitro p53 squelching by both TFIIB and TFIID.
Mol Cell Biol **15:**6474-6478.
45. **Espinosa JM, Verdun RE, Emerson BM.** 2003. p53 functions through stress- and
promoter-specific recruitment of transcription initiation components before and after
DNA damage. Mol Cell **12:**1015-1027.
46. **Okuda M, Nishimura Y.** 2015. Real-time and simultaneous monitoring of the
phosphorylation and enhanced interaction of p53 and XPC acidic domains with the
TFIIH p62 subunit. Oncogenesis **4:**e150.
47. **Xiao H, Pearson A, Coulombe B, Truant R, Zhang S, Regier JL, Triezenberg SJ,**
Reinberg D, Flores O, Ingles CJ, et al. 1994. Binding of basal transcription factor
TFIIH to the acidic activation domains of VP16 and p53. Mol Cell Biol **14:**7013-7024.
48. **Leveillard T, Andera L, Bissonnette N, Schaeffer L, Bracco L, Egly JM, Wasylyk B.**
1996. Functional interactions between p53 and the TFIIH complex are affected by
tumour-associated mutations. EMBO J **15:**1615-1624.
49. **Maston GA, Evans SK, Green MR.** 2006. Transcriptional regulatory elements in the
human genome. Annu Rev Genomics Hum Genet **7:**29-59.
50. **Huang B, Jones SA, Brandenburg B, Zhuang X.** 2008. Whole-cell 3D STORM reveals
interactions between cellular structures with nanometer-scale resolution. Nat Methods
5:1047-1052.

51.   **Chen J, Zhang Z, Li L, Chen BC, Revyakin A, Hajj B, Legant W, Dahan M,**
Lionnet T, Betzig E, Tjian R, Liu Z. 2014. Single-molecule dynamics of enhanceosome
assembly in embryonic stem cells. Cell **156:**1274-1285.
52.   **Ludtke SJ, Baldwin PR, Chiu W.** 1999. EMAN: semiautomated software for high-
resolution single-particle reconstructions. J Struct Biol **128:**82-97.
53.   **Mastronarde DN.** 2005. Automated electron microscope tomography using robust
prediction of specimen movements. J Struct Biol **152:**36-51.
54.   **Mindell JA, Grigorieff N.** 2003. Accurate determination of local defocus and specimen
tilt in electron microscopy. J Struct Biol **142:**334-347.
55.   **Shaikh TR, Gao H, Baxter WT, Asturias FJ, Boisset N, Leith A, Frank J.** 2008.
SPIDER image processing for single-particle reconstruction of biological
macromolecules from electron micrographs. Nat Protoc **3:**1941-1974.
56.   **van Heel M, Harauz G, Orlova EV, Schmidt R, Schatz M.** 1996. A new generation of
the IMAGIC image processing system. J Struct Biol **116:**17-24.
57.   **Chen JZ, Grigorieff N.** 2007. SIGNATURE: a single-particle selection system for
molecular electron microscopy. J Struct Biol **157:**168-173.
58.   **Lander GC, Stagg SM, Voss NR, Cheng A, Fellmann D, Pulokas J, Yoshioka C,**
Irving C, Mulder A, Lau PW, Lyumkis D, Potter CS, Carragher B. 2009. Appion: an
integrated, database-driven pipeline to facilitate EM image processing. J Struct Biol
166:95-102.
59.   **Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC,**
Ferrin TE. 2004. UCSF Chimera--a visualization system for exploratory research and
analysis. J Comput Chem **25:**1605-1612.
956

A

HeLa
Nuclear Extract

Phosphocellulose
P11 column

1M

0.5M

0.3M

IP
α−TAF4 mAb

Ctl | + DNA | + DNA + p53 | + DNA + p53 + TFIIA

Wash

Peptide eluates

TFIID co-complexes

B

bax promoter template

-675 ⎯⎯ [poly A stretch] ⎯⎯ [p53 REs] ⎯⎯ [TATA] ⎯→ TSS ⎯ -317

hdm2 promoter template

2415 ⎯⎯ [Sp1][Sp1][Sp1][Sp1] [ETS][AP1][p53 REs][TATA] ⎯→ TSS ⎯ 2853

Left gel labels (left to right lanes: IID | *bax*+IID | p53+*bax*+IID | IIA+p53+*bax*+IID | *bax* DNA):

TAF1
TAF2/3/4
TAF5
TAF6
← *bax* DNA
TAF7
← p53
TAF8
TBP
TAF9a/9b
TAF10
TAF11
TAF12
TAF13

Lanes: 1 2 3 4 5

Right gel labels (lanes: IIA+p53+*hdm2* DNA | IID | *hdm2*+IID | p53+*hdm2*+IID | IIA+p53+*hdm2*+IID):

hdm2 → DNA
p53 →
IIA-α/β

IIA-γ →

TAF1
TAF2/3/4
TAF5
TAF6
TAF7
TAF8/TBP
TAF9a/9b/TAF10
TAF11
TAF12
TAF13

Lanes: 6 7 8 9 10

Figure 2

A

QD-TFIID
QD-565
+
p53

Promoter DNA
Cy3
p53 REs
TATA
InR
Cy5
Streptavidin
Glass slide
532nm Laser
640nm Laser
Total Internal Reflection

Promoter DNA Channel

QD-TFIID Channel

B

- p53

Promoter DNA
QD-TFIID

σ = 129nm

DNA-TFIID Displacement
(N_DNA=4154)

1.3% DNA co-localized
(58/4154)

+ p53

Promoter DNA
QD-TFIID

σ = 48nm

DNA-TFIID Displacement
(N_DNA=919)

20.7% DNA co-localized
(191/919)

C

☐ TFIID   ☐ TFIID + p53

% DNA Co-localized

bax   hdm2

WT | WT | Mu RE | Mu Core | Mu RE/Co

D

WT hdm2

-   IID   p53 IID

p53
RE
TATA
InR
MTE
DPE

TFIID+p53+DNA
TFIID+DNA
DNA

-127 -88 -80 -55 -27 10 68
-112 -61 -32 -4 31 37 44 51 54 59 62
-15 22

p53 RE TATA InR MTE DPE

Mu RE/Co hdm2

-   IID   p53 IID

p53
RE
TATA
InR
MTE
DPE

TFIID+p53+DNA
TFIID+DNA
DNA

-100 -55 11 68
-137 -112 -78 -33 -4 37 44 51
-28 -17 22 31 53 62

p53 RE TATA InR MTE DPE

Figure 3

A

B

Figure 4

A

TFIID form

Canonical ⇌ Transition ⇌ Rearranged

Frequency

- TFIID
- TFIID/p53
- TFIID/p53/*bax* DNA

0.65 0.6 0.5 0.4 0.3 0.2 0.1 0

Canonical   Transition   Rearranged

C strong downstream DNA view

+1

weak upstream DNA view

Focused classification: upstream DNA

B

TFIID/p53/TFIIA/promoter DNA

bax promoter

hdm2 promoter

Figure 6