Natural Selection has Shaped Coding and Non-coding Transcription in Primate CD4+ T-cells

Charles G. Danko[1,2,*], Lauren A. Choate[1], Brooke A. Marks[1], Edward J. Rice[1], Zhong Wang[1], Tinyi Chu[1,3], Andre L. Martins[1,3], Noah Dukler[4], Scott A. Coonrod[1,2], Elia D. Tait Wojno[1,5], John T. Lis[6], W. Lee Kraus[7,8], & Adam Siepel[4,*]

[1] Baker Institute for Animal Health, College of Veterinary Medicine, Cornell University, Ithaca, NY 14853.

[2] Department of Biomedical Sciences, College of Veterinary Medicine, Cornell University, Ithaca, NY 14853.

[3] Graduate field of Computational Biology, Cornell University, Ithaca, NY 14853.

[4] Simons Center for Quantitative Biology, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY 11724.

[5] Department of Microbiology & Immunology, College of Veterinary Medicine, Cornell University, Ithaca, NY 14853.

[6] Department of Molecular Biology and Genetics, Cornell University, Ithaca, NY 14853.

[7] Laboratory of Signaling and Gene Regulation, Cecil H. and Ida Green Center for Reproductive Biology Sciences, University of Texas Southwestern Medical Center, Dallas, TX 75390.

[8] Division of Basic Research, Department of Obstetrics and Gynecology, University of Texas Southwestern Medical Center, Dallas, TX 75390.

*__Address correspondence to__:

Charles G. Danko, Ph.D.	Adam Siepel, Ph.D.
Baker Institute for Animal Health	Simons Center for Quantitative Biology
Cornell University	Cold Spring Harbor Laboratory
Hungerford Hill Rd.	1 Bungtown Rd.
Ithaca, NY 14853	Cold Spring Harbor, NY 11724
Phone: 607-256-5620	Phone: 516-367-6922
E-mail: dankoc@gmail.com	E-mail: asiepel@cshl.edu

Abstract:

Transcriptional regulatory changes have been shown to contribute to phenotypic differences between species, but many questions remain about how gene expression evolves. Here we report the first comparative study of nascent transcription in primates. We used PRO-seq to map actively transcribing RNA polymerases in resting and activated CD4+ T-cells in multiple human, chimpanzee, and rhesus macaque individuals, with rodents as outgroups. This approach allowed us to measure transcription separately from post-transcriptional processes. We observed general conservation in coding and non-coding transcription, punctuated by numerous differences between species, particularly at distal enhancers and non-coding RNAs. We found evidence that transcription factor binding sites are a primary determinant of transcriptional differences between species, that stabilizing selection maintains gene expression levels despite frequent changes at distal enhancers, and that adaptive substitutions have driven lineage-specific transcription. Finally, we found strong correlations between evolutionary rates and long-range chromatin interactions. These observations clarify the role of primary transcription in regulatory evolution.

Following decades of speculation that changes in the regulation of genes could be a potent
force in the evolution of form and function[1-3], investigators have now empirically demonstrated the
evolutionary importance of gene regulation across the tree of life[4-12]. Changes in gene expression
are primarily driven by mutations to non-coding DNA sequences, particularly those that bind
sequence-specific transcription factors[13]. Accordingly, adaptive nucleotide substitutions at
transcription factor binding sites (TFBSs)[9,10,14-16] and gains and losses of TFBSs[17-25] both appear to
make major contributions to the evolution of gene expression. These events are believed to modify
a variety of rate-limiting steps early in transcriptional activation[26]. In addition, transcriptional
activity is generally correlated with various epigenomic and structural features, including post-
translational modifications to core histones, the locations of architectural proteins such as CTCF,
and the organization of topological associated domains. Like TFBSs, these features display general
conservation across species, yet do exhibit some variation, with differences between species
roughly proportional to evolutionary distance[27]. Moreover, differences between species in these
features correlate with differences in gene expression[8,24,28-30].

Nevertheless, many open questions remain about the roles of TFBSs, chromatin
organization, and posttranscriptional regulation in the evolution of gene expression. For example,
there is a surprisingly limited correlation between differences in binding events and differences in
mRNA expression levels[31-33]. Possible reasons for this discordance include non-functional TF
binding[31,32,34], compensatory gains and losses of TFBSs[20,35-38], difficulties associating distal
enhancers with target genes[39], and a dependency of TF function on chromatin or chromosomal
organization[40]. In addition, some changes in mRNA expression appear to be "buffered" at the post-
transcriptional level[41-43]. Finally, it remains unclear to what degree epigenomic differences
between species are causes and to what degree they are effects of differences in gene expression.

One reason why it has been difficult to disentangle these contributions to gene expression is
that expression is typically measured in terms of the abundance of mRNA, which is subject to post-
transcriptional processing[44] and therefore is an indirect measure of the transcription of genes by
RNA polymerase II. An alternative and complementary approach is to measure the production of
nascent RNAs using Precision Run-On and sequencing (PRO-seq) and related technologies[45-50].
These nascent RNA sequencing methods directly measure active transcription and are highly
sensitive to immediate and transient transcriptional responses to stimuli[51]. In addition, they can
detect active regulatory elements as well as target genes, because these elements themselves
display distinctive patterns of transcription, which are highly underrepresented in RNA-seq data
owing to their rapid degradation[34,52,53]. Indeed, the latest nascent RNA sequencing methods, such as
PRO-seq[46], in combination with new computational tools for regulatory element prediction[54], serve
as powerful single-assay systems for both identifying regulatory elements and measuring
transcription levels.

With these advantages in mind, we undertook a genome-wide comparative analysis of
transcription in primates using PRO-seq. Our comparison of PRO-seq data across species revealed
overall conservation in the transcription of both coding and non-coding elements, but also
uncovered numerous differences between species. Together, our observations provide new
insights into the evolution of transcription in primates.

Patterns of transcription in resting and activated CD4+ T-cells

We developed nucleotide-resolution maps of RNA polymerase in CD4+ T-cells isolated from five mammalian species. Samples were collected under resting and activated conditions from three unrelated individuals representing each of three primate species—humans, chimpanzees, and rhesus macaques—spanning ~25-30 million years of evolution (MYR) (**Fig. 1a**). To compare with studies that focus on longer evolutionary branch lengths, we also collected resting samples from a single individual in each of two rodent species—mouse and rat—which together serve as an outgroup to the primates (~80 MYR divergence). We used flow cytometry to validate the purity of isolated CD4+ cells (**Supplementary Fig. 1**). In addition, we used measurements of transcriptional activity of T-cell subset markers for T-helper type 1 (Th1), Th2, Th17, T-regulatory, and T-follicular helper cells to demonstrate that the population of CD4+ T-cell subsets within the total CD4+ population was largely similar across species (**Supplementary Fig. 2**).

PRO-seq[46,49] libraries were sequenced to a combined depth of ~1 billion uniquely mapped reads (~78-274 million per species) (**Supplementary Table 1**). According to a principal component analysis, the first, second, and third sources of variation in the complete dataset were, respectively, the rodent vs. primate species, variation across the primate species, and the treatment condition (**Supplementary Fig. 3**). Similarly, hierarchical clustering of these data grouped the primate samples first by cell type or treatment condition and subsequently by species, with the rodent samples as outgroups that were more similar to untreated primate CD4+ T-cells than to other samples included in the comparison (**Fig. 1b**). The correlation between untreated samples decreased linearly with evolutionary time (**Supplementary Fig. 4**), consistent with reports that differences between species arise, on average, at a roughly constant evolutionary rate[27,55].

To gain further insight into the evolution of the response to CD4+ T-cell stimulation, we compared transcriptional activity under resting and activated conditions within and between species. Here we focused on 42,556 GENCODE-annotated transcription units (TUs) best supported by PRO-seq data for human CD4+ T-cells[56]. In humans, we found that PMA and ionomycin (π) significantly altered the transcription levels of 6,940 (13%) of these TUs ($p < 0.01$, deSeq2[57]; **Fig. 1c**). Parallel analyses in chimpanzee and rhesus macaque revealed many similarities in transcriptional changes following π treatment (**Supplementary Fig. 5a-b**). We identified a core set of 3,157 TUs that undergo evolutionarily conserved transcriptional changes in all three species following 30 min. of π-treatment, including many of the classical response genes (e.g., IFNG, TNFα, IL2, and IL2RA), as well as numerous novel genes and lincRNAs (**Supplementary Fig. 5a-c**). Active transcriptional regulatory elements (TREs) undergoing changes following π-treatment were enriched for a similar set of transcription factor binding motifs across species, including those for NF-kB and the AP-1 heterodimers FOS and JUN, which are known to be activated by canonical T-cell receptor signaling (**Fig. 1d; Supplementary Note 1**). Thus, the core regulatory principles responsible for T-cell signaling and activation appear to remain broadly conserved across primate evolution.

Rapid evolutionary changes in transcribed enhancers

We used dREG[54] to identify 30,357 active TREs in human CD4+ T-cells, based on patterns of enhancer-templated RNA (eRNA) or upstream antisense (uaRNA) transcription evident from PRO-seq data (**Online Methods**). We classified these predicted TREs as either protein-coding promoters

or candidate enhancers based on their proximity to gene annotations. The predicted TREs in each
group were highly concordant with other marks of regulatory function in human CD4+ T-cells used
previously to define groups of candidate enhancers, including acetylation of histone 3 lysine 27
(H3K27ac), mono- and trimethylation of histone 3 lysine 4 (H3K4me1 and H3K4me3), and DNase-I-
seq signal[58] (**Fig. 1e**). Notably, dREG identified >83% of DNase-I hypersensitive sites (DHS) marked
by H3K27 acetylation in human CD4+ T-cells, consistent with prior work suggesting that
transcription patterns alone can recover the majority of active enhancers defined by independent
criteria[34,52,54]. Furthermore, we identified 88% and 91% of DHSs marked, respectively, by H3K9ac
and H4K16ac, two other markers of regulatory function. Taken together, these data suggest that
PRO-seq patterns reveal the locations of TREs with high sensitivity. In the analysis that follows, we
will refer to these dREG-identified distal TREs as "enhancers" for simplicity.
Extending our dREG analysis to untreated CD4+ T-cells from additional species revealed
71,748 TREs that were active in untreated T-cells in at least one species (ranging between 27,581
and 39,387 TREs in each species). We defined two types of changes between species: (1) changes
in the abundance of Pol II at TREs that were present across all species, and (2) complete gains or
losses in at least one species (see **Supplementary Note 2, Supplementary Fig. 6,** and **Online
Methods**). We found that 52% of enhancers showed evidence of changes in Pol II transcriptional
activity in at least one of the three primate species and 81% showed changes at the longer
evolutionary distance between primates and rodents (**Fig. 2a**), similar to recent observations in
other systems[10,24] (**Supplementary Note 3**). Enhancers were predicted to be completely gained or
lost at nearly eight times the rate of promoters (35% of enhancers; 12% of promoters; p < 2.2e-16
by Fisher's exact test), consistent with recent observations based on H3K27ac and H3K4me3[24]. By
contrast, TREs induced by π treatment were much more likely to be conserved, and showed similar
conservation at promoters and enhancers (**Fig. 2a**).
Next we tested whether evolutionary changes in transcriptional activity correlate with the
enrichment of other marks of active enhancers. Predicted lineage-specific human enhancers were
enriched for both active and repressive enhancer marks (**Fig. 2b; Supplementary Fig. 7**). Whereas
apparent human gains were enriched for high levels of the active enhancer marker H3K27ac, sites
with reduced transcriptional activity in humans showed much lower enrichments of H3K27ac.
Furthermore, locations at which the dREG signal was completely lost in a human-specific fashion
displayed levels of H3K27ac approaching those of randomly selected background sites (**Fig. 2b**).
Intriguingly, many of the losses on the human branch retained H3K4me1, which marks both active
and inactive enhancers[59,60], and these losses displayed higher levels of chromatin marks associated
with transcriptional repression than a random background (**Fig. 2b**), indicating that, at least in
some cases, an active ancestral primate enhancer retains a 'poised' chromatin state in human,
despite losing both transcriptional activity and H3K27ac. Thus, evolutionary changes in poised and
active marks may commonly occur as distinct events.
163
Transcriptional changes correlate with DNA sequence differences
To investigate whether changes in TRE activity are accompanied by changes in DNA
sequence, we compared phyloP sequence conservation scores[61] at transcriptionally conserved TREs
with phyloP scores at TREs that display evolutionary changes in transcription. Because signatures
of sequence conservation in TREs are likely to be most pronounced in transcription factors binding

sites (TFBS), we restricted our sequence conservation analyses to matches to 567 clusters of TF
binding motifs selected based on their distinct DNA binding specificities (see **Online Methods**)[62,63].
We required that motif matches were present in dREG sites, and adopted a threshold at which
nearly half of the motifs discovered were bona fide TFBSs, as measured by ChIP-seq (positive
predictive value [PPV]= 0.47).

TFBSs found in transcriptionally conserved dREG sites showed a marked enrichment for
higher phyloP scores relative to surrounding regions, indicating local sequence conservation (**Fig.**
3a). By contrast, TFBSs in lineage-specific dREG sites had substantially reduced enrichments in
phyloP scores (**Fig. 3a, cyan/blue**). Notably, TFBSs in dREG sites lost on the human lineage
showed enhanced conservation compared with those in human-specific gains. This observation is
consistent with losses evolving under conservation in other mammalian species (which contribute
to the phyloP scores) and gains emerging relatively recently. Each of these patterns was robust to
corrections for potentially confounding differences in the distribution of sites, as well to choices of
motif score thresholds (**Supplementary Fig. 8a**). Relaxing the motif score threshold to provide
sensitivity for larger numbers of TFBSs at the expense of specificity, revealed patterns of
conservation that correlate with the information content of positions within the DNA sequence
motif (**Supplementary Fig. 8b**), further supporting TF binding as the functional property driving
sequence conservation at these sites. Together, these analyses support the hypothesis that the
sequences in TFBSs are a primary driver of transcriptional differences between species.

We searched for examples of DNA sequence differences that might be responsible for
transcriptional changes following π treatment, hypothesizing that they might be characterized
more easily than sequences responsible for transcription in the untreated condition, because these
transcriptional changes were likely driven by a limited number of master TFs[64] (**Fig. 1e**). In one
example, we found nucleotide substitutions in three apparent NF-kB binding sites in the proximal
promoter and an internal enhancer of *SGPP2* that correlate with differences in *SGPP2* expression
(**Fig. 3b; Supplementary Fig. 9**). Two of these putative binding sites were bound by NF-kB in
human cell lines according to ChIP-seq data from ENCODE. Moreover, substitutions observed in
human were found to disrupt the same position in the motif as NF-kB binding QTLs[65] (see **Online**
Methods), and showed a general trend toward higher NF-kB binding in the human alleles
(**Supplementary Fig. 9**). To test the hypothesis that observed DNA sequence changes produced
differential transcriptional activity, we cloned DNA from each primate species into a reporter
vector driving luciferase activity in an MCF-7 cell model, which recapitulates the primary
transcriptional features of the *SGPP2* locus[66] (**Supplementary Fig. 9**). Differences in basal
luciferase activity were generally concordant with those observed between species
(**Supplementary Fig. 10**). Moreover, both the proximal promoter of *SGPP2* and the internal
enhancer both activated luciferase expression more strongly following NF-kB activation when
human DNA was cloned, but not with orthologous DNA from the other primates (**Fig. 3c**).

To determine whether these TREs affect the expression of *SGPP2* in its native genomic
context, we silenced each TRE by using CRISPRi, which targets a catalytically dead CAS9 fused to
the Krüppel-associated box repressor (dCAS9-KRAB), to specifically tri-methylate lysine 9 of
histone 3 (H3K9me3)[67]. Three independent single-guide RNAs (sgRNAs) targeting the internal
enhancer and two designed for the proximal promoter reduced *SGPP2* transcription to 50-60% of
its resting level (p = 1.5e-3 and 2.6e-2, respectively, by a two-tailed t-test; **Fig. 3d**), consistent with

these TREs directly contributing to *SGPP2* expression in MCF-7 cells. Three sgRNAs targeting the
upstream enhancer also had a significant effect on *SGPP2* expression (p = 1.8e-4). Notably, the
genome assemblies for chimpanzee and rhesus macaque harbor deletions of this upstream TRE that
appear likely to affect its activity (**Supplementary Fig. 9**). However, although silencing individual
enhancers reduced the transcription level of *SGPP2* following NF-kB activation, silencing individual
enhancers was insufficient to completely abolish induction of *SGPP2* (**Fig. 3d**)*.* Taken together, our
findings show that at least two of the three TREs regulating *SGPP2* drove expression patterns
matching PRO-seq data in a reporter assay, but none completely explained *SGPP2* activation *in situ*.
These observations suggest that that multiple causal substitutions in NF-kB binding sites may work
in concert to achieve human-specific activation of *SGPP2*.

In several cases, as with *SGPP2*, we observed numerous nucleotide substitutions within
individual or clustered TFBSs. These clusters of substitutions are highly unlikely to occur by chance
and suggest that positive selection may have driven adaptation of these binding sites. Indeed,
SGPP2 falls in a region recently identified as having an excess of derived alleles in modern humans
compared with the sequenced Neanderthal (**Fig. 3d**)[68], potentially consistent with recent positive
selection driving evolutionary changes in *SGPP2* transcription. To more directly measure the
impact of positive selection, we used INSIGHT[69] to compare patterns of within-species
polymorphism and between-species sequence divergence in TREs that had undergone human
lineage-specific transcriptional changes. This analysis indicated that dREG sites are most strongly
influenced by weak negative selection (**Fig. 3f**), based on an excess of low-frequency derived alleles
in human populations, as has been reported previously for regulatory sequences[9]. Nevertheless,
TREs with lineage-specific transcriptional changes in human CD4+ T-cells showed reduced weak
negative selection and were strikingly enriched for adaptive nucleotide substitutions (p < 0.01
INSIGHT likelihood ratio test; **Fig. 3f**), consistent with positive selection at these sites. We estimate
a total of at least 121 adaptive substitutions since the human/chimpanzee divergence within TFBSs
that undergo transcriptional changes in human CD4+ T-cells. Despite limited power to detect the
specific contributions of many individual TFs at our stringent motif match score threshold, we did
note significant excesses of putatively adaptive substitutions in the predicted binding sites of
several TFs, including motifs recognized by forkhead box family, POU-domain containing, and ELF/
ETS family (**Supplementary Fig. 11**; p < 0.01, INSIGHT likelihood ratio test). These estimates
highlight the substantial contribution of adaptive evolutionary changes in TFBSs that may influence
the transcriptional activity of TREs.

Correlation between protein-coding and non-coding transcription

We noticed that evolutionary changes in protein-coding gene transcription frequently
correlate with changes in non-coding transcription units (TU) located nearby. To examine this
pattern more generally, we adapted our recently reported hidden Markov model (HMM)[70] to
estimate the location of TUs genome-wide, based on patterns of aligned PRO-seq reads and the
location of TREs. Using this method, we annotated 54,793 TUs active in CD4+ T-cells of at least one
of the primate species, approximately half of which overlap existing GENCODE annotations or their
associated upstream antisense RNAs (**Supplementary Fig. 12a**). A cross-species comparison of
the transcription levels for various TU classes (**Fig. 4a**) revealed that non-coding RNAs evolve in
expression most rapidly and protein-coding genes evolve most slowly. GENCODE-annotated

lincRNAs undergo evolutionary changes in expression about as frequently as the unannotated non-
coding RNAs predicted by our HMM, which are likely enriched for bi-directionally unstable eRNA
species. The broad similarity in evolutionary conservation between these two non-coding RNA
classes may be consistent with observations that some lincRNAs function as enhancers for nearby
protein-coding genes, and that stable accumulation of the transcript is dispensable for this
biological function[71].

We next measured the extent to which non-coding and protein-coding transcriptional
activities are correlated through evolutionary time. We found that evolutionary changes in protein-
coding gene expression among any of the primate species were highly correlated with those at both
upstream (Pearson's $R = 0.85$, $p < 2.2e-16$) and internal ($R = 0.66$, $p < 2.2e-16$) antisense transcripts
of the same genes. Moreover, changes in the transcriptional activity of gene promoters correlate
with changes in the activity of matched distal enhancers defined by various criteria: namely,
enhancers to which the promoters loop according to cell-type matched ChIA-PET data ($R = 0.45$-
0.61, $p < 2.2e-16$; depending on analysis assumptions)[72], enhancers located nearby the promoters
($R = 0.69$, $p < 2.2e-16$), or enhancers that share the same topological associated domain as the
promoter ($R = 0.62$, $p < 2.2e-16$)[73]. Using a generalized linear model to integrate expression
changes in multiple types of TUs, we can explain 74% of the variance in gene transcription levels
when we observe differences between species ($R^2 = 0.74$ in a held-out set of sites, $p < 2.2e-16$; **Fig.**
4b) based on the activities of looped TREs, nearby TREs in the same topological associated domain,
internal antisense TUs, and the upstream antisense TU. Thus evolutionary changes that result in
differences in Pol II recruitment to protein-coding genes are well correlated across all interacting
TREs, indicating a shared evolutionary pressure at proximal and distal TREs.

Rates of Enhancer Evolution Vary with Evidence for Gene Interactions
Despite this overall positive correlation, transcription at enhancers evolves rapidly and is
frequently unaccompanied by transcriptional changes at nearby protein-coding genes. For example,
CCR7 transcription is highly conserved among both primate and rodent species (**Fig. 5a;**
Supplementary Fig. 2) in spite of several apparent changes in enhancer activity within the same
locus (gray vertical bars). These findings are consistent with recent observations that changes in
enhancers within densely populated loci often do not have appreciable effects on the transcription
of genes within the locus[36,37].

To explain this effect, we searched for genomic features correlated with conservation of
transcription at enhancers, focusing on untreated CD4+ T-cells in order to leverage the large
amount of public data available for this cell type. Not surprisingly, one of the features most strongly
correlated with transcriptional conservation at enhancers is the distance from the nearest
transcription start site of a protein-coding gene (**Fig. 5b**). In particular, more than half of
enhancers located within 10 kbp of an annotated TSS are shared across all three primate species,
whereas for distal enhancers located between 100 kbp to 1 Mbp from a TSS that fraction drops to
roughly a third. This relationship is driven by lineage-specific gains or losses of enhancer activity,
and to a lesser extent by changes in TRE activity levels, rather than by differences in the alignability
of orthologous DNA (**Supplementary Fig. 13**).

These simple distance-based observations, however, ignore the critical issue of chromatin
interactions between enhancers and promoters. To account for such loop interactions, we extracted

6,520 putative TRE interactions from Chromatin Interaction Analysis with Paired End Tag
sequencing (ChIA-PET) data recognizing loops marked with H3K4me2 in human CD4+ T-cells[72].
We found that 55% of enhancers that participate in these loops were conserved between primate
species compared to only 47% of non-looped enhancers (**Fig. 5c**; p = 5.6e-4, Fisher's exact test).
Moreover, higher transcriptional conservation at looped enhancers does not depend on the
distance to the transcription start site. Parallel analysis of promoter-capture Hi-C data[74] revealed
that the strength of chromatin interaction was correlated with evolutionary conservation of distal
TREs, corroborating the result obtained using ChIA-PET (p < 1e-3, bootstrap test). We observed
similar levels of conservation at recently defined super-enhancers[75], although this conservation
may simply reflect an enrichment for loop interactions (48% of TREs in super-enhancers loop
according to ChIA-PET, compared to 15% of all TREs). Looped enhancers were also enriched for
elevated phyloP scores relative to either non-looped enhancers or randomly selected DNA
sequences (**Supplementary Fig. 14**; phyloP > 0.75; p < 2.2e-16, Wilcoxon Rank Sum Test). That the
enhancers participating in loop interactions are more highly conserved at both the transcription
and DNA-sequence levels indicates that these enhancers have a disproportionately large effect on
fitness, presumably owing to a more direct role in transcriptional regulation.

Enhancer-Promoter Interactions Contribute to Constraint on Gene Transcription
Levels
Distal loop interactions do not fully account for the disparity between enhancers and
promoters in evolutionary rates—even looped enhancers still evolve significantly faster than
promoters (p = 3e-3, Fisher's exact test). We hypothesized that redundancy in enhancers may help
to explain rapid enhancer evolution. More specifically, we asked whether redundancy makes
protein-coding genes regulated by multiple distal TREs, such as *CCR7* (**Fig. 5a**), more robust to
enhancer turnover than those influenced by fewer distal TREs. Indeed, we found that evolutionary
conservation of promoter TRE transcription is remarkably strongly correlated with the number of
loop interactions a promoter has with distal sites (**Fig. 6a**, weighted Pearson's correlation = 0.87; p
< 1e-3 by a bootstrap test). A similar trend was observed between the number of loop interactions
made by a target promoter and DNA sequence conservation in transcription factor binding motifs at
the promoter, although the effect was weaker and did not meet our criteria for statistical
significance (**Fig. 6b**, weighted Pearson's correlation = 0.65; p = 0.07 by a bootstrap test).
But how does redundancy in distal TREs relate to the evolutionary conservation of the
distal TREs themselves? If redundant distal TREs compensate for one another in some way,
perhaps each one will be less, rather than more, conserved when their associated promoters have
larger numbers of loop interactions. To address this question, we examined the rate of
conservation of looped distal TREs as a function of the number of loops in which their gene-
proximal partners participated. We found that DNA sequence conservation in putative TFBSs
negatively correlates with the number of loops at the proximal end (**Fig. 6d;** weighted Pearson's
correlation = –0.80; p = 2e-3 by a bootstrap test). We noted a similar trend toward a negative
correlation between the conservation of distal TRE transcription and the number of loop
interactions (weighted Pearson's correlation = –0.67; p = 0.059, two-sided bootstrap test, **Fig. 6c**).
These results suggest that each associated distal TFBS is individually less essential at genes having
larger numbers of loop interactions with distal sites, and they are therefore consistent with a model

in which such TFBSs are more freely gained and lost during evolution. Taken together, our results
imply that distance, looping, and redundancy of enhancers all contribute to constraints on the
evolutionary rates of changes in gene transcription.
344
Discussion:
We have carried out the first comparative analysis of primary transcription in any
phylogenetic group, focusing on CD4+ T-cells in primates. Using PRO-seq and various
computational tools, we estimated the locations and abundance of transcription units with high
resolution and accuracy. In comparison to previous studies in primates[29,33,76–78], this approach
separated primary transcription from post-transcriptional processing, allowing us to study eRNAs,
lincRNAs, and other rapidly degraded non-coding RNAs, as well as protein-coding genes. We found
clear relationships between the DNA sequences of TFBSs and differential transcription across
species and treatment conditions. We also found evidence that some transcriptional changes in
humans were driven by adaptive evolution in nearby binding sites. Overall, our study provides new
insights into the mode and tempo of recent evolutionary changes in transcription in primates.
Perhaps our most striking observation is that many non-coding transcription units,
particularly eRNAs and lincRNAs, have undergone rapid evolutionary changes in comparison to
protein-coding genes. Similar observations have been reported previously for lincRNAs[79], but, to
our knowledge, the observation for eRNAs is new, and it raises a number of questions. First, why
are some enhancers more conserved than others? In particular, we find that enhancers proximal to,
or that loop to, annotated promoters tend to be most constrained (**Fig 5b-c**). These enhancers may
simply be most crucial for activating their target genes, but other factors may also contribute to
their constraint. For example, perhaps these enhancers are enriched for tissue-specific functions,
and are less constrained due to reduced pleiotropy[80]. Or perhaps many of them simply are not
functional at all, and are transcribed as a by-product of other processes.
Second, how do protein-coding genes maintain stable transcription levels across species
despite the rapid turnover of associated enhancers? One possibility is that many rapidly evolving
enhancers are either not functional or act on targets other than the ones we have identified.
However, several of our findings argue against this possibility; for example, we find that even
looped enhancers, for which we have direct evidence of a promoter interaction, evolve significantly
faster than promoters, and that eRNA conservation is strongly correlated with the number of loop
interactions at associated promoters (**Fig. 6**). An alternative explanation, which appears more
plausible to us, is that stabilizing selection on transcription levels drives enhancers to compensate
for one another as they undergo evolutionary flux. This explanation would be compatible with
reports from model systems[36,38]. Our finding that sequence conservation at distal enhancers is
negatively correlated with the number of loop interactions at associated promoters
(**Supplementary Fig. 14**) is also consistent with this explanation. The possibility of pervasive
stabilizing selection on transcription levels in primates has been noted previously based on RNA-
seq data[81], but our data allow for more direct observations of both active transcription and
associated regulatory elements.
A third question is, if most transcribed enhancers do indeed influence gene expression, then
why are so many of them weakly maintained by natural selection? At present, we can only
speculate on the answer to this question. One possibility is that some of the apparent turnover

events we have observed actually represent enhancers that have simply switched cell types in
activity, as has been reported in some cases[19]. But it is also possible that selection tends to act
diffusely on enhancers across an entire locus, rather than strongly on individual enhancers, as has
been proposed in cancer evolution[82]. Our observation that multiple DNA sequence changes at the
SGPP2 locus appear functional provides some initial support this this hypothesis. It will be possible
to evaluate these hypotheses more rigorously as better data describing enhancers and enhancer-
promoter interactions across many cell types become available for these and other groups of
species.

Methods:
393
Multiple species PRO-seq library generation. *Isolation of primate CD4+ T-cells.* All human and
animal experiments were done in compliance with Cornell University IRB and IACUC guidelines.
We obtained peripheral blood samples (60-80 mL) from healthy adult male humans, chimpanzees,
and rhesus macaques. Informed consent was obtained from all human subjects. To account for
within-species variation in gene transcription we used three individuals to represent each primate
species. Blood was collected into purple top EDTA tubes. Human samples were maintained
overnight at 4C to mimic shipping non-human primate blood samples. Blood was mixed 50:50 with
phosphate buffered saline (PBS). Peripheral blood mononuclear cells (PBMCs) were isolated by
centrifugation (750x g) of 35 mL of blood:PBS over 15 mL Ficoll-Paque for 30 minutes at 20C. Cells
were washed three times in ice cold PBS. CD4+ T-cells were isolated using CD4 microbeads
(Miltenyi Biotech, 130-045-101 [human and chimp], 130-091-102 [rhesus macaque]). Up to 10^8
PBMCs were resuspended in binding buffer (PBS with 0.5% BSA and 2mM EDTA). Cells were
bound to CD4 microbeads (20uL of microbeads/ 10^7 cells) for 15 minutes at 4C in the dark. Cells
were washed with 1-2 mL of PBS/BSA solution, resuspended in 500uL of binding buffer, and passed
over a MACS LS column (Miltenyi Biotech, 130-042-401) on a neodymium magnet. The MACS LS
column was washed three times with 2mL PBS/BSA solution, before being eluted off the
neodymium magnet. Cells were counted in a hemocytometer.
411
Isolation of CD4+ T-cells from mouse and rat. Spleen samples were collected from one male mouse
(FVB) and one male rat (Albino Oxford) that had been sacrificed for IACUC-approved research not
related to the present study. Dissected spleen was mashed through a cell strainer using a sterile
glass pestle and suspended in 20 mL RPMI-1640. Cells were pelleted at 800xg for 3 minutes and
resuspended in 1-5mL of ACK lysis buffer for 10 minutes at room temperature to lyse red blood
cells. RPMI-1640 was added to a final volume 10 times that used for ACK lysis (10-40 mL). Cells
were pelleted at 800xg for 3 minutes, counted in a hemocytometer, and resuspended in RPMI-1640
to a final concentration of 250,000 cells per ml. CD4+ T-cells were isolated from splenocytes using
products specific for mouse and rat (Miltenyi Biotech, 130-104-453 [mouse], 130-090-319 [rat])
following instructions from Miltenyi Biotech, and as described above.
422
T-cell treatment and PRO-seq library generation. CD4+ T-cells were allowed to equilibrate in RPMI-
1640 supplemented with 10% FBS for 2-4 hours before starting experiments. Primate CD4+ T-cells
were stimulated with 25ng/mL PMA and 1mM Ionomycin (P/I or π) or vehicle control (2.5uL EtOH
and 1.66uL DMSO in 10mL of culture media). We selected the minimum concentrations which
saturate the production of IL2 and IFNG mRNA after 3 hours of treatment (data not shown). A 30
min. treatment duration was selected after observing a sharp increase in ChIP-qPCR signal for RNA
Pol II phosphorylated at serine 5 on the C-terminal domain on the IFNG promoter at 30 min. (data
not shown). To isolate nuclei, we resuspended cells in 1 mL lysis buffer (10 mM Tris-Cl, pH
8, 300 mM sucrose, 10 mM NaCl, 2 mM MgAc2, 3 mM CaCl2 and 0.1% NP-40). Nuclei were
washed in 10 mL of wash buffer (10 mM Tris-Cl, pH 8, 300 mM sucrose, 10 mM NaCl and 2 mM
MgAc2) to dilute free NTPs. Nuclei were washed in 1 mL, and subsequently resuspended in 50
µL, of storage buffer (50 mL Tris-Cl, pH 8.3, 40% glycerol, 5 mM MgCl2 and 0.1 mM EDTA), snap

frozen in liquid nitrogen and kept for up to 6 months before making PRO-seq libraries. PRO-
seq libraries were created exactly as described previously[46]. In most cases, we completed library
preps with one member of each species (usually one human, chimpanzee, and rhesus macaque) to
prevent batch effects from confounding differences between species. Samples were sequenced on
an Illumina Hi-Seq 2000 or NextSeq500 at the Cornell University Biotechnology Resource Center.
440
Mapping PRO-seq reads. We mapped PRO-seq reads using standard informatics tools. Our PRO-seq
mapping pipeline begins by removing reads that fail Illumina quality filters and trimming adapters
using cutadapt with a 10% error rate[83]. Reads were mapped with BWA[84] to the appropriate
reference genome (either hg19, panTro4, rheMac3, mm10, or rn6) and a single copy of the Pol I
ribosomal RNA transcription unit (GenBank ID# U13369.1). Mapped reads were converted to
bigWig format for analysis using BedTools[85] and the bedGraphToBigWig program in the Kent
Source software package[86]. The location of the RNA polymerase active site was represented by the
single base, the 3' end of the nascent RNA, which is the position on the 5' end of each sequenced
read. After mapping reads to the reference genome, three samples (one human, U and PI, one
chimpanzee, U and PI, and one rhesus macaque, U and PI) were identified as having poor data
quality on the basis of the number of uniquely mapped reads, and were excluded from downstream
analysis.
453
Mapping 1:1 orthologs between different species. During all comparative analyses, the genomic
coordinates of mapped reads, dREG scores, and other parameters of interest were converted to the
human assembly (hg19) using CrossMap[87]. We converted genomic coordinates between genome
assemblies using reciprocal-best (rbest) nets[88]. Reciprocal-best nets have the advantage that
comparisons between species are constrained to 1:1 orthologues. This constraint on mapping is
enforced by requiring each position to map uniquely in a reciprocal alignment between the human
reference and the other species in the comparison. We downloaded rbest nets for hg19-mm10,
hg19-panTro4, hg19-rn6 from the UCSC Genome Browser. We created rbest nets for hg19-rheMac3
using the doRecipBets.pl script provided as part of the Kent Source software package.
463
Analysis of transcriptional regulatory elements. *Defining a consensus set of transcriptional*
regulatory elements. We predicted TREs using dREG[54] separately in each species' reference
genome. dREG uses a support vector regression model to score each site covered in a PRO-seq
dataset based on its resemblance to features associated with transcription start sites in a reference
training dataset. The dREG model was trained to recognize DNase-I-hypersensitive sites that also
show substantial evidence of GRO-cap data in six PRO-seq or GRO-seq datasets measuring
transcription in resting K562 cells. dREG scores were computed in the reference genome of each
species in order to provide as much information as possible on the native context of each locus. In
all cases, we combined the reads from all individuals for each species in order to maximize power
for the discovery of TREs. In the primate species, treated and untreated CD4+ T-cells were
analyzed separately.
We then defined a consensus set of TRE annotations, each of which bore the signature of an
active TRE in at least one species and treatment condition. To define such a set, dREG scores were
first converted to human reference genome (hg19) coordinates using CrossMap and the reciprocal-

best nets. The advantage of converting dREG scores between the reference genome is that
individual bases transfer more completely than genomic intervals using CrossMap and related
tools. We then identified TREs in each species separately by thresholding the dREG scores. In all
analyses, we selected a threshold of 0.3, which corresponds to a predicted false discovery rate of
<7% compared with other sources of genomic data in human CD4+ T-cells. In addition, parallel
analyses at separate thresholds (0.25 and 0.35) provided results that were in all cases consistent
with those reported in the main manuscript (**Supplementary Table 2**). The set of overlapping
TREs from each species were reduced to a single element containing the union of all positions
covered by the set using bedops, and sites within 500 bp of each other were further merged. We
assigned each putative TRE the maximum dREG score for each species and for each treatment
condition.

Identifying differences in TREs between species. Differences in TRE transcription in 3-way (human-
chimp-rhesus macaque) or 5-way (human-chimp-rhesus macaque-mouse-rat) species comparisons
were identified using a combination of heuristics and statistical tests. Starting with the consensus
set of TREs in hg19 coordinates, we first excluded potential one-to-many orthologs, by eliminating
TREs that overlapped gaps in the reciprocal-best nets that were not classified as gaps in the
standard nets. The remaining TREs were classified as unmappable when no orthologous position
was defined in the rbest nets. Complete gains and losses were defined as TREs that were mappable
in all species and for which the dREG score was less than 0.05 in at least one species and greater
than 0.30 in at least one other species (see **Supplementary Note 1**). Gains and losses were
assigned to a lineage based on an assumption of maximum parsimony under the known species
phylogeny. We defined a set of TREs that displayed high-confidence changes in activity by
comparing differences in PRO-seq read counts between species using deSeq2 and thresholding at a
1% false discovery rate (as described below). Changes in TRE activities were compared to histone
modification ChIP-seq, DNase-I-seq, and DNA methyl immunoprecipitation data from the
Epigenome Roadmap project[58].

TRE classification. For some analyses, TREs were classified as likely promoters or enhancers on the
basis of their distance from known protein-coding gene annotations (GENCODE v.19). TRE classes
of primary interest include (see also **Supplementary Fig. 7**): (1) Promoters: near an annotated
transcription start site (<100 bp); (2) Enhancers: distal to an annotated transcription start site
(>5,000 bp)

Covariates that correlate with TRE changes. We compared the frequency at which evolutionary
changes in transcription occur at TREs in a variety of different genomic contexts. We examined the
rate of change as a function of distance from the nearest annotated transcription start site in
GenCode v.19. TREs were binned by distance in increments of 0.02 on a log10 scale and we
evaluated the mean rate at which evolutionary changes in TRE transcription arise in each bin. We
also compared the rate of changes in TRE transcription across a variety of functional associations,
including loop interactions, within the same topological associated domain, and in super-enhancers.
H3K4me2 ChIA-PET data describing loop interactions were downloaded from the Gene Expression
Omnibus (GEO) database (GSE32677) and the genomic locations of loops were converted from

hg18 to hg19 coordinates using the liftOver tool. We also analyzed a separate dataset profiling loop
interactions based on promoter capture Hi-C data in human CD4+ T-cells taken from the
supplementary materials of ref. [74]. Topological associated domains (TADs) based on Hi-C data for
GM12878 cells were also downloaded from GEO (GSE63525). Super-enhancers in CD4+ T-cells
were taken from the supplementary data for ref. [75]. In all cases we excluded sites with potential
one-to-many orthology in any of the species included in the comparison (typically just the three
primates). Potential one-to-many orthologs were defined based on differences in the standard and
reciprocal-best nets for each species pair.
529
Refining the location of active TREs using dREG-HD. During analyses of transcription factor binding
motifs we further refined the location of TREs to the region between divergent paused RNA
polymerase using a strategy that we call dREG-HD (manuscript in preparation, preliminary version
available at https://github.com/Danko-Lab/dREG.HD). Briefly, we used an epsilon-support vector
regression (SVR) with a Gaussian kernel to map the distribution of PRO-seq reads to smoothed
DNase-I signal intensities. Training was conducted on randomly chosen positions within dREG
peaks extended by 200bp on either side. Selection of feature vectors was optimized based on
Pearson correlation coefficients between the imputed and experimental DNase-I score over the
validation set. PRO-seq data was normalized by sequencing depth and further scaled such that the
maximum value of any prediction dataset is within 90 percentile of the training examples. We
chose a step size to be 60bp and extending 30 steps on each direction. The final model was trained
using matched DNase-I and PRO-seq data in K562 cells.
Next we identified peaks in the imputed DNase-I hypersensitivity profile by fitting the
imputed DNase-I signal using a cubic spline and identifying local maxima. We optimized two free
parameters that control the (1) smoothness of spline curve fitting, and (2) threshold on the
imputed DNase-I signal intensity. Parameters were optimized to achieve an appropriate trade-off
between FDR and sensitivity on the testing K562 dataset. Parameters were tuned using a grid
optimization over free parameters.
548
DNA sequence analysis. *Finding candidate transcription factor binding motifs.* All motif analyses
focused on 1,964 human TF binding motifs from the CisBP database[62] clustered using an affinity
propagation algorithm into 567 maximally distinct DNA binding specificities (see ref [63]). Scores,
which reflect a $\log_e$-odds ratio comparing each candidate motif model to a third-order Markov
background model, were calculated using the RTFBSDB package[63]. We selected two separate motif
thresholds for different analyses. Scores >10 were used in analyses which mix multiple TF binding
motifs, and strike a tradeoff that focuses on minimizing false positives at the expense of sensitivity.
We dropped the cutoff score to motifs >8 in analyses that use individual motifs in order to increase
statistical power. For each of these thresholds, we estimated the mean genome-wide positive
predictive values to be 0.47 and 0.38, respectively, for motif cutoffs of 10 and 8, by comparing
motifs to ChIP-seq peak calls in K562 cells. During comparative analyses we scanned each primate
reference genome separately with each motif to allow the detection of a putative binding site in any
of the species included in the analysis, and then moved scores to a human (hg19) reference genome
using the CrossMap tool. We chose this strategy because changes in TRE activity may reflect
changes in binding in any of the primate species. For example, human gains may be explained by

either a new binding site for a transcriptional activator in the human genome, or a loss in binding of
a transcriptional repressor that was bound in both primate species.

Motif enrichment in TREs that change during CD4+ T-cell activation. Motifs enriched in up- or down-
regulated dREG-HD TREs during CD4+ T-cell activation ($p < 0.01$) were selected using Fisher's exact
test with a Bonferroni correction for multiple hypothesis testing. Up- or down-regulated TREs
were compared to a background set of >2,500 GC-content matched TREs that do not change
transcription levels following π treatment (log-2 fold change <0.5-fold in magnitude and $p > 0.25$)
using the *enrichmentTest* function in RTFBSDB[63]. To test for motif robustness, the background
resampling was repeated 100 times and motifs were selected that achieve a significant result in
>90%.

DNA sequence conservation analysis. For our evolutionary conservation analysis, we used phyloP
scores[61] based on the 100-way genome alignments available in the UCSC Genome Browser (hg19).
In all cases, bigWig files were obtained from the UCSC Genome Browser and processed using the
bigWig package in R. We represented evolutionary conservation as the mean phyloP score in each
identified TFBS in the indicated set of dREG-HD sites.

Enrichment of DNA sequence changes in motifs. We identified single-nucleotide DNA sequence
differences at sites at which two of three primate species share one base and the third species
diverges.   We intersected these species-specific divergences with matches to transcription factor
binding motifs found within dREG-HD sites that undergo transcriptional changes between primate
species. Because many motifs in Cis-BP are similar to one another, we first partitioned the motifs
using clustering (as described above), and examined enrichments at the level of these clusters.
Motifs were ranked by the Fisher's exact test p-value of the enrichment of species divergences in
dREG-HD sites that change transcription status (where changes in DNA sequence and transcription
occur on the same branch) to dREG-HD sites that do not change. We also compute the enrichment
ratio, which we define as the number of species divergences in each TF binding motif in dREG-HD
sites that change on the same branch normalized to the same statistic in sites that do not change.

INSIGHT analysis. We examined the modes by which DNA sequences evolve in human lineage-
specific dREG-HD sites or DHSs using INSIGHT[69]. We passed INSIGHT either complete DHSs, dREG-
HD sites, or TFBS within dREG-HD sites that undergo the changes (see *Identifying differences in*
TREs between species) indicated in the comparison. Human gains and losses, for example, were
comprised of 4,384 dREG-HD sites with 9,924 separate regions (median length of 16 bp) after
merging overlapping TFBSs with a log-odds score greater than 10.   We also analyzed 24
transcription factors each of which has more than 900 occurrences in dREG-HD sites that change on
the human branch (log-odds score >8). All analyses were conducted using the INSIGHT web server
(http://compgen.cshl.edu/INSIGHT/) with the default settings enabled.

bQTL analysis. Frequency shift estimates for all variants in Teranchi et al. (2016) were provided by
the Frasier lab and converted to a queryable database filtered to include only variants with
coverage by 25 reads (75th percentile) or more to avoid noise at low read counts. For each

sequence/variant query, a set of four equivalent sequences/alternate allele pairs was constructed
by swapping which allele was the reference and getting the reverse complement for both alleles.
For example, given a sequence:variant:position combination of AATCGAA:C:3, the other queries
produced were AACCGAA:T:3 (allele swap), TTCGATT:G:5 (reverse complement), and TTCGGTT:A:5
(reverse complement allele swap). Frequency shifts were computed by taking the post-ChIP
frequency minus the pre-ChIP frequency for the human reference allele. Since k-mers longer than 7
had few hits, we allowed for wildcards (N) in longer sequences that would match any base.
Wildcards were introduced into a k-mer by matching the k-mer sequence to the NF-kB motif and
replacing the 3 lowest information content positions with N(s). Systematic shifts from 0 were
tested using a one-tailed t-test. P-values for systematic differences at multiple sites were combined
using Fisher's method.
618
De novo discovery of transcription units. *Identification of transcription units (TU) using a three-*
state hidden Markov model. We inferred transcription units (TU) using a three-state hidden Markov
model (HMM) similar to those we have recently published[51,70]. Each TU begins at a TRE identified
using dREG and continues through the entire region inferred to be transcribed, which can covers
tens- to hundreds- of kilobases. Three states were used to represent background (i.e., outside of a
transcription unit), the TU body, and a post-polyA decay region. The HMM transition structure is
shown in **Supplementary Fig. 13a**. We allow skipping over the post-polyA state, as unstable
transcripts do not have these two-phase profiles. We took advantage of dREG as a potential signal
for transcription initiation by incorporating the dREG score (maximum value in the interval from a
given positive read-count position until the next, clamped to the zero-one interval) as a transition
probability from the background to the transcription body state. PRO-seq data is generally sparse,
so we applied a transformation that encoded only non-zero positions and the distance between
such consecutive positions (**Supplementary Fig. 13a**). Our model described this transformed data
using emissions distributions based on two types of variables. The first type of emission variable
defines the PRO-seq read counts in non-zero positions. These counts were modeled using Poisson
distributions in the background and post-polyA states, and using a Negative Binomial distribution
in the transcription body state. The negative binomial distribution can be seen as a mixture of
Poisson distributions with gamma-distributed rates and therefore allows for variation in TU
expression levels across the genome. The second type of emission variable describes the
distribution of distances in base pairs between positions having non-zero read counts. This
distribution was modeled using a separate geometric distribution for each of the three states.
Maximum likelihood estimates of all free parameters were obtained via Expectation Maximization,
on a per-chromosome basis. TU predictions were then obtained using the Viterbi algorithm with
parameters fixed at their maximum-likelihood values. Finally these predictions were mapped from
the transformed coordinates back to genomic coordinates. Source code for our implementation is
publicly available on GitHub: https://github.com/andrelmartins/tunits.nhp.
645
Inferring TU boundaries in the common great ape ancestor. We identified the most likely TU
locations in the great ape ancestor by maximum parsimony. TUs were identified and compared in
human reference coordinates (hg19) for all species. We used the bedops package to find the
intersection between the predicted TU intervals in each pair of species (i.e., human-chimp, human-

rhesus macaque, and chimp-rhesus macaque). Intersections (>= 1bp) between pairs of species
were merged, resulting in a collection of TUs shared by any two pairs of species, and therefore
likely to be a TU in the human-chimp ancestor. All steps were applied independently on the plus
and minus strands. These steps identified 37,626 putative TUs active in CD4+ T-cells of the primate
ancestor. We added 17,167 TUs that did not overlap ancestral TUs but were found in any one of the
three primate species.

Transcription unit classification. TUs were classified by annotation type using a pipeline similar to
ones that we have described recently[51,70,89]. Before classifying TUs we applied a heuristic to refine
TUs on the basis of known annotations. TUs that completely overlap multiple gene annotations
were broken at the transcription start site provided that a dREG site overlapped that transcription
start site. Classification was completed using a set of rules to iteratively refine existing annotations,
as shown in **Supplementary Fig. 13A**. Unless otherwise stated, overlap between a TU and a
transcript annotation was defined such that >50% of a TU matched a gene annotation and covers at
least 50% of the same annotation. TUs overlapping GENCODE annotations (>50% overlap, defined
as above) were classified using the biotype in the GENCODE database into protein coding, lincRNA
(lincRNA or processed transcript), or pseudogene. The remaining transcripts were classified as
annotated RNA genes using GENCODE annotations, the rnaGenes UCSC Genome Browser track
(converted from hg18 to hg19 coordinates), and miRBase v20[90]. As many RNA genes are processed
from much longer TUs, we required no specific degree of overlap for RNA genes. Upstream
antisense (i.e., divergent) TUs were classified as those within 500bp of the transcription start site of
any GENCODE or higher level TU annotation (including lincRNAs). Antisense transcripts were
defined as those with a high degree of overlap (>50%) with annotated protein coding genes in the
opposite orientation. The remaining transcripts with a high degree of overlap (>50%) to annotated
repeats in the repeatmasker database (rmsk) were classified as repeat transcription. Finally, any
TUs still remaining were classified as unannotated, and were further divided into those which are
intergenic or that partially overlapping existing annotations.

Comparing transcription between conditions and species. *Comparing transcription before and*
after CD4+ T-cell activation. We compared π treated and untreated CD4+ T-cells within each of the
primate species using gene annotations (GENCODE v19). We counted reads in the interval between
500 bp downstream of the annotated transcription start site and either the end of the gene or
60,000 bp into the gene body (whichever was shorter). This window was selected to avoid (1)
counting reads in the pause peak near the transcription start site, and (2) to focus on the 5' end of
the gene body affected by changes in transcription during 30 minutes of π treatment assuming a
median elongation rate of 2 kb/ minute[51,91]. We limited analyses to gene annotations longer than
500 bp in length. To quantify transcription at enhancers, we counted reads in the window covered
by each dREG-HD site plus an additional 250 bp on each end. Differential expression analysis was
conducted using deSeq2[57].

Comparing transcription between species. Read counts were compared between different species in
hg19 coordinates. In all analyses, reads were transferred to the hg19 reference genome using
CrossMap with rbest nets. Our analysis focused on transcription units or on the union of dREG sites

across species. We focused our analysis of transcription units on the interval between 250 bp
downstream of the annotated transcription start site and either the end of the gene or 60,000 bp
into the gene body (whichever was shorter). We limited our analyses to TUs longer than 500 bp in
length. Reads counts were obtained within each transcription unit, gene annotation, or enhancer,
abbreviated here as a 'region of interest' (ROI), that has confident one-to-one orthology in all
species examined in the analysis. This strategy of focusing on blocks of one-to-one orthology avoids
errors caused by systematic differences in mappability or repeat content of species-specific
genomic segments. We broke each each ROI into segments that have conserved orthology between
hg19 and all species examined in the analysis, which included either a three-way (human-chimp-
rhesus macaque) or five-way (human-chimp-rhesus macaque-mouse-rat) species comparison. We
defined intervals of one-to-one orthology as those represented in levels 1, 3, and 5 of the reciprocal
best nets (with gaps defined in levels 2, 4, and 6)[88]. Reads that map to regions that have orthology
defined in all species were counted using the bigWig package in R using reads mapped to hg19
coordinates. Final counts for each ROI were defined as the sum of read counts within the regions of
orthology that intersect that ROI. ROIs without confident one-to-one orthologs in all species
analyzed were discarded. Our pipeline makes extensive use of the bigWig R package, Kent source
tools, as well as the bedops and bedtools software packages[85,92]. Differential expression was
conducted between species using the deSeq2 package for R, as described above.
711

712   **MCF-7 G11 cell culture.** MCF7 G11 tamoxifen resistant cells, were a gift from Dr. Joshua LaBaer.
713   Cells were maintained in DMEM with 5% FBS, antibiotics, and 1uM tamoxifen. MCF-7 G11 dCas9-
714   KRAB stable cell lines were made (as described below) and were maintained in DMEM with 5%
715   FBS, antibiotics, and 1uM tamoxifen. MCF-7 G11 dCas9-KRAB sgRNA stable cell lines were
716   maintained in DMEM with 5% FBS, antibiotics, 1uM tamoxifen, and 150ug/ul Hygromycin B.
717

Luciferase assays. Genomic DNA was isolated from human, chimp, and rhesus macaque PBMCs
depleted for CD4+ cells using a Quick-DNA Miniprep Plus Kit (#D4068S; Zymo research) following
the manufacturer's instructions. Putative enhancer regions were amplified from the genomic DNA,
restriction digested with KpnI and MluI, and cloned into the pGL3-promoter vector (Promega). The
same orthologous regions were amplified from all three species with identical primers where
possible or species-specific primers covering orthologous DNA in diverged regions. Vectors were
co-transfected with pRL-SV40 Renilla (Promega) in a 10:1 ratio (500ng pGL3 to 50ng pRL-SV40) in
MCF7 G11 cells cultured in 1uM tamoxifen. Transfected cells were treated with either 25ng/ml
TNFa or water 21 hours after transfection. 24 hours post-transfection, luminescence was measured
in triplicate using the Dual-Luciferase® Reporter Assay System (Promega).
728

Silencing endogenous TREs using dCAS9-KRAB. *Cloning signle-guide RNAs (sgRNAs).* Single-
guide RNAs (sgRNAs) were designed using the CRISPR design tool (http://crispr.mit.edu) and
sequences are shown in **Supplementary Table 3**. Forward and reverse sgRNAs were synthesized
separately by IDT and annealed. T4 Polynucleotide Kinase (NEB) was used to phosphorylate the
forward and reverse sgRNA during the annealing. 10x T4 DNA Ligase Buffer, which contains 1mM
ATP, was incubated for 30 minutes at 37°C and then at 95C for 5 minutes, decreasing by 5°C every 1
minute until 25°C. Oligos were diluted 1:200 using Molecular grade water. sgRNAs were inserted

into the pLenti SpBsmBI sgRNA Hygro plasmid from addgene (#62205) by following the authors
protocol[93]. The plasmid was linearized using BsmBI digestion (NEB) and purified using gel
extraction (QIAquick Gel Extraction Kit). The purified linear plasmid was then dephosphorylated
using Alkaline Phosphatase Calf Intestinal (CIP) (NEB) to ensure the linear plasmid did not ligate
with itself. A second gel extraction was used as before to purify the linearized plasmid. The purified
dephosphorylated linear plasmid and phosphorylated annealed oligos were ligated together using
the Quick Ligation Kit (NEB). The ligated product was transformed into One Shot Stbl3 Chemically
Competent E. coli (ThermoFisher Scientific). 100ul of the transformed bacteria were plated on
Ampicillin (200ug/ml) plates. Single colonies were picked, sequenced, and the plasmid was isolated
using endo free midi-preps from Omega.

Transfection of MCF-7 G11 cell lines. We used lentivirus to transfect MCF-7 cells. Lentivirus was
made using lipofectamine 3000 from Invitrogen. Phoenix Hek cells (grown in DMEM with 10% FBS
and antibiotics) were seeded in a 6-well plate at 400,000 cells/plate. Cells were grown until ~90%
confluent. 1ug of pHAGE_EF1a_dCas9-KRAB plasmid from addgene (#50919) plasmid or the pLenti
SpBsmBI sgRNA Hygro (addgene #62205) containing each sgRNA, 0.5ug of psPAX (addgene
#12260), and 0.25ug pMD2.G (addgene #12259) were mixed.
MCF-7 G11 cells were plated at ~200,000 cells/well in a 6-well plate. 24 hours later
3ml/well of virus was mixed with 10ug/ml polybrene and incubated for 5 minutes at room
temperature. This mix was added to the cells and centrifuged for 40 minutes at 800g at 32C (Viju
Vijayan Pillai, personal communication). 12-24 hours later the virus was removed and fresh media
was added. 24-48 hours later the cells were selected with 2ug/ml puromycin for 2 weeks. The MCF-
7 G11 dCas9-KRAB stable cell lines was grown and maintained in puromycin. A second lentiviral
infection was done using the stable MCF-7 G11 dCas9-KRAB cells. The same protocol was used. 24-
48 hours later the cells were selected with 150ug/ml Hygromycin B. New stable cell lines were
grown and maintained in hygromycin B.

TNFa treatment. Prior to TNFa treatment, cells were grown for 3 days in DMEM with 5% FBS,
antibiotics, tamoxifen and hygromycin. Cells were then left untreated or treated for 40 min with
25ng/ml TNFa. RNA was extracted using TRIzol Reagent (Invitrogen). We reverse transcribed 1ug
of RNA and used this as input for real-time quantitative PCR (RT-PCR) to analyze *SGPP2* expression.
Primers for *SGPP2* were designed targeting a sequence in intron 1, upstream of the intronic
enhancer. Raw *Cp* values were transferred to units of expression using a standard dilution curve
comprised of a mixture of cDNA from each sample within the biological replicate. We included four
serial dilutions, each of which covered a two-fold difference in expression. Each sample was further
normalized for differences in RNA content by primers recognizing the 18S rRNA control. The ratio
between normalized *SGPP2* expression in each sgRNA-transfected MCF-7 cell line and the empty
vector control was log-2 transformed and tested for differences from 0 using a two-sided t-test.

Data availability. PRO-seq data was deposited into the Gene Expression Omnibus database under
accession number GSE85337.

Code availability. All data analysis scripts and software are publicly available on GitHub:
https://github.com/Danko-Lab/CD4-Cell-Evolution.
780

References:

1. Jacob, F. & Monod, J. Genetic regulatory mechanisms in the synthesis of proteins. *J. Mol. Biol.* **3,**
318–356 (1961).
2. Britten, R. J. & Davidson, E. H. Gene regulation for higher cells: a theory. *Science* **165,** 349–357
(1969).
3. King, M. C. & Wilson, A. C. Evolution at two levels in humans and chimpanzees. *Science* **188,**
107–116 (1975).
4. Rockman, M. V. *et al.* Ancient and recent positive selection transformed opioid cis-regulation in
humans. *PLoS Biol.* **3,** e387 (2005).
5. Prabhakar, S. *et al.* Human-specific gain of function in a developmental enhancer. *Science* **321,**
1346–1350 (2008).
6. Capra, J. A., Erwin, G. D., McKinsey, G., Rubenstein, J. L. R. & Pollard, K. S. Many human
accelerated regions are developmental enhancers. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **368,**
20130025 (2013).
7. McLean, C. Y. *et al.* Human-specific loss of regulatory DNA and the evolution of human-specific
traits. *Nature* **471,** 216–219 (2011).
8. Cotney, J. *et al.* The evolution of lineage-specific regulatory activities in the human embryonic
limb. *Cell* **154,** 185–196 (2013).
9. Arbiza, L. *et al.* Genome-wide inference of natural selection on human transcription factor
binding sites. *Nat. Genet.* **45,** 723–729 (2013).
10. Prescott, S. L. *et al.* Enhancer Divergence and cis-Regulatory Evolution in the Human and
Chimp Neural Crest. *Cell* **163,** 68–83 (2015).
11. Siepel, A. & Arbiza, L. Cis-regulatory elements and human evolution. *Curr. Opin. Genet. Dev.* **29,**
81–89 (2014).
12. Wray, G. A. The evolutionary significance of cis-regulatory mutations. *Nat. Rev. Genet.* **8,** 206–
216 (2007).
13. Wilson, M. D. *et al.* Species-specific transcription in mice carrying human chromosome 21.
Science **322,** 434–438 (2008).
14. Khurana, E. *et al.* Integrative annotation of variants from 1092 humans: application to cancer
genomics. *Science* **342,** 1235587 (2013).
15. Haygood, R., Fedrigo, O., Hanson, B., Yokoyama, K.-D. & Wray, G. A. Promoter regions of many
neural- and nutrition-related genes have experienced positive selection during human
evolution. *Nat. Genet.* **39,** 1140–1144 (2007).
16. Torgerson, D. G. *et al.* Evolutionary processes acting on candidate cis-regulatory regions in
humans inferred from patterns of polymorphism and divergence. *PLoS Genet.* **5,** e1000592
(2009).
17. Schmidt, D. *et al.* Five-vertebrate ChIP-seq reveals the evolutionary dynamics of transcription
factor binding. *Science* **328,** 1036–1040 (2010).
18. Ballester, B. *et al.* Multi-species, multi-transcription factor binding highlights conserved control
of tissue-specific biological pathways. *Elife* **3,** e02626 (2014).
19. Vierstra, J. *et al.* Mouse regulatory DNA landscapes reveal global principles of cis-regulatory
evolution. *Science* **346,** 1007–1012 (2014).
20. Arnold, C. D. *et al.* Quantitative genome-wide enhancer activity maps for five Drosophila
species show functional enhancer conservation and turnover during cis-regulatory evolution.
Nat. Genet. **46,** 685–692 (2014).
21. Doniger, S. W. & Fay, J. C. Frequent gain and loss of functional transcription factor binding sites.
PLoS Comput. Biol. **3,** e99 (2007).
22. Zheng, W., Zhao, H., Mancera, E., Steinmetz, L. M. & Snyder, M. Genetic analysis of variation in
transcription factor binding in yeast. *Nature* **464,** 1187–1191 (2010).

23. Bradley, R. K. *et al.* Binding site turnover produces pervasive quantitative changes in
831   transcription factor binding between closely related Drosophila species. *PLoS Biol.* **8,**
832   e1000343 (2010).
833   24. Villar, D. *et al.* Enhancer evolution across 20 mammalian species. *Cell* **160,** 554–566 (2015).
834   25. Chuong, E. B., Elde, N. C. & Feschotte, C. Regulatory evolution of innate immunity through co-
835   option of endogenous retroviruses. *Science* **351,** 1083–1087 (2016).
836   26. Fuda, N. J., Ardehali, M. B. & Lis, J. T. Defining mechanisms that regulate RNA polymerase II
837   transcription in vivo. *Nature* **461,** 186–192 (2009).
838   27. Carvunis, A.-R. *et al.* Evidence for a common evolutionary rate in metazoan transcriptional
839   networks. *Elife* **4,** (2015).
840   28. Zhou, X. *et al.* Epigenetic modifications are associated with inter-species gene expression
841   variation in primates. *Genome Biol.* **15,** 547 (2014).
842   29. Cain, C. E., Blekhman, R., Marioni, J. C. & Gilad, Y. Gene expression differences among primates
843   are associated with changes in a histone epigenetic modification. *Genetics* **187,** 1225–1234
844   (2011).
845   30. Xiao, S. *et al.* Comparative epigenomic annotation of regulatory DNA. *Cell* **149,** 1381–1392
846   (2012).
847   31. Paris, M. *et al.* Extensive divergence of transcription factor binding in Drosophila embryos with
848   highly conserved gene expression. *PLoS Genet.* **9,** e1003748 (2013).
849   32. Cusanovich, D. A., Pavlovic, B., Pritchard, J. K. & Gilad, Y. The functional consequences of
850   variation in transcription factor binding. *PLoS Genet.* **10,** e1004226 (2014).
851   33. Wong, E. S. *et al.* Decoupling of evolutionary changes in transcription factor binding and gene
852   expression in mammals. *Genome Res.* **25,** 167–178 (2015).
853   34. Hah, N., Murakami, S., Nagari, A., Danko, C. G. & Kraus, W. L. Enhancer transcripts mark active
854   estrogen receptor binding sites. *Genome Res.* **23,** 1210–1223 (2013).
855   35. Domené, S. *et al.* Enhancer turnover and conserved regulatory function in vertebrate evolution.
856   *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **368,** 20130027 (2013).
857   36. Wunderlich, Z. *et al.* Krüppel Expression Levels Are Maintained through Compensatory
858   Evolution of Shadow Enhancers. *Cell Rep.* **12,** 1740–1747 (2015).
859   37. Cannavò, E. *et al.* Shadow Enhancers Are Pervasive Features of Developmental Regulatory
860   Networks. *Curr. Biol.* **26,** 38–51 (2016).
861   38. Ludwig, M. Z., Bergman, C., Patel, N. H. & Kreitman, M. Evidence for stabilizing selection in a
862   eukaryotic enhancer element. *Nature* **403,** 564–567 (2000).
863   39. Sanyal, A., Lajoie, B. R., Jain, G. & Dekker, J. The long-range interaction landscape of gene
864   promoters. *Nature* **489,** 109–113 (2012).
865   40. Vietri Rudan, M. *et al.* Comparative Hi-C reveals that CTCF underlies evolution of chromosomal
866   domain architecture. *Cell Rep.* **10,** 1297–1309 (2015).
867   41. Khan, Z. *et al.* Primate transcript and protein expression levels evolve under compensatory
868   selection pressures. *Science* **342,** 1100–1104 (2013).
869   42. Bauernfeind, A. L. *et al.* Evolutionary Divergence of Gene and Protein Expression in the Brains
870   of Humans and Chimpanzees. *Genome Biol. Evol.* **7,** 2276–2288 (2015).
871   43. Battle, A. *et al.* Genomic variation. Impact of regulatory variation from RNA to protein. *Science*
872   **347,** 664–667 (2015).
873   44. Pai, A. A. *et al.* The contribution of RNA decay quantitative trait loci to inter-individual
874   variation in steady-state gene expression levels. *PLoS Genet.* **8,** e1003000 (2012).
875   45. Core, L. J., Waterfall, J. J. & Lis, J. T. Nascent RNA sequencing reveals widespread pausing and
876   divergent initiation at human promoters. *Science* **322,** 1845–1848 (2008).
877   46. Kwak, H., Fuda, N. J., Core, L. J. & Lis, J. T. Precise maps of RNA polymerase reveal how
878   promoters direct initiation and pausing. *Science* **339,** 950–953 (2013).
47. Churchman, L. S. & Weissman, J. S. Nascent transcript sequencing visualizes transcription at

nucleotide resolution. *Nature* **469,** 368–373 (2011).

48.   Nojima, T. *et al.* Mammalian NET-Seq Reveals Genome-wide Nascent Transcription Coupled to
RNA Processing. *Cell* **161,** 526–540 (2015).

49.   Mahat, D. B. *et al.* Base-pair-resolution genome-wide mapping of active RNA polymerases using
precision nuclear run-on (PRO-seq). *Nat. Protoc.* **11,** 1455–1476 (2016).

50.   Schwalb, B. *et al.* TT-seq maps the human transient transcriptome. *Science* **352,** 1225–1228
(2016).

51.   Hah, N. *et al.* A rapid, extensive, and transient transcriptional response to estrogen signaling in
breast cancer cells. *Cell* **145,** 622–634 (2011).

52.   Core, L. J. *et al.* Analysis of nascent RNA identifies a unified architecture of initiation regions at
mammalian promoters and enhancers. *Nat. Genet.* **46,** 1311–1320 (2014).

53.   Andersson, R. *et al.* Nuclear stability and transcriptional directionality separate functionally
distinct RNA species. *Nat. Commun.* **5,** 5336 (2014).

54.   Danko, C. G. *et al.* Identification of active transcriptional regulatory elements from GRO-seq
data. *Nat. Methods* **12,** 433–438 (2015).

55.   Khaitovich, P. *et al.* A Neutral Model of Transcriptome Evolution. *PLoS Biol.* **2,** e132 (2004).

56.   Dukler, N. *et al.* Nascent RNA sequencing reveals a dynamic global transcriptional response at
genes and enhancers to the natural medicinal compound celastrol. *bioRxiv* 117689 (2017).
doi:10.1101/117689

57.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol.* **15,** 550 (2014).

58.   Roadmap Epigenomics Consortium *et al.* Integrative analysis of 111 reference human
epigenomes. *Nature* **518,** 317–330 (2015).

59.   Bonn, S. *et al.* Tissue-specific analysis of chromatin state identifies temporal signatures of
enhancer activity during embryonic development. *Nat. Genet.* **44,** 148–156 (2012).

60.   Zentner, G. E., Tesar, P. J. & Scacheri, P. C. Epigenetic signatures distinguish multiple classes of
enhancers with distinct cellular functions. *Genome Res.* **21,** 1273–1283 (2011).

61.   Pollard, K. S., Hubisz, M. J., Rosenbloom, K. R. & Siepel, A. Detection of nonneutral substitution
rates on mammalian phylogenies. *Genome Res.* **20,** 110–121 (2010).

62.   Weirauch, M. T. *et al.* Determination and inference of eukaryotic transcription factor sequence
specificity. *Cell* **158,** 1431–1443 (2014).

63.   Wang, Z., Martins, A. L. & Danko, C. G. RTFBSDB: an integrated framework for transcription
factor binding site analysis. *Bioinformatics* (2016). doi:10.1093/bioinformatics/btw338

64.   Macian, F. NFAT proteins: key regulators of T-cell development and function. *Nat. Rev.*
Immunol. **5,** 472–484 (2005).

65.   Tehranchi, A. K. *et al.* Pooled ChIP-Seq Links Variation in Transcription Factor Binding to
Complex Disease Risk. *Cell* **165,** 730–741 (2016).

66.   Franco, H. L., Nagari, A. & Kraus, W. L. TNFα signaling exposes latent estrogen receptor binding
sites to alter the breast cancer cell transcriptome. *Mol. Cell* **58,** 21–34 (2015).

67.   Thakore, P. I. *et al.* Highly specific epigenome editing by CRISPR-Cas9 repressors for silencing
of distal regulatory elements. *Nat. Methods* **12,** 1143–1149 (2015).

68.   Green, R. E. *et al.* A draft sequence of the Neandertal genome. *Science* **328,** 710–722 (2010).

69.   Gronau, I., Arbiza, L., Mohammed, J. & Siepel, A. Inference of natural selection from
interspersed genomic elements based on polymorphism and divergence. *Mol. Biol. Evol.* **30,**
1159–1171 (2013).

70.   Chae, M., Danko, C. G. & Kraus, W. L. groHMM: a computational tool for identifying unannotated
and cell type-specific transcription units from global run-on sequencing data. *BMC*
Bioinformatics **16,** 222 (2015).

71.   Engreitz, J. M. *et al.* Local regulation of gene expression by lncRNA promoters, transcription
and splicing. *Nature* **539,** 452–455 (2016).

72. Chepelev, I., Wei, G., Wangsa, D., Tang, Q. & Zhao, K. Characterization of genome-wide
enhancer-promoter interactions reveals co-expression of interacting genes and modes of
higher order chromatin organization. *Cell Res.* **22,** 490–503 (2012).

73. Rao, S. S. P. *et al.* A 3D map of the human genome at kilobase resolution reveals principles of
chromatin looping. *Cell* **159,** 1665–1680 (2014).

74. Javierre, B. M. *et al.* Lineage-Specific Genome Architecture Links Enhancers and Non-coding
Disease Variants to Target Gene Promoters. *Cell* **167,** 1369–1384.e19 (2016).

75. Hnisz, D. *et al.* Super-enhancers in the control of cell identity and disease. *Cell* **155,** 934–947
(2013).

76. Barreiro, L. B., Marioni, J. C., Blekhman, R., Stephens, M. & Gilad, Y. Functional comparison of
innate immune signaling pathways in primates. *PLoS Genet.* **6,** e1001249 (2010).

77. Gilad, Y., Oshlack, A., Smyth, G. K., Speed, T. P. & White, K. P. Expression profiling in primates
reveals a rapid evolution of human transcription factors. *Nature* **440,** 242–245 (2006).

78. Blekhman, R., Oshlack, A., Chabot, A. E., Smyth, G. K. & Gilad, Y. Gene regulation in primates
evolves under tissue-specific selection pressures. *PLoS Genet.* **4,** e1000271 (2008).

79. Kutter, C. *et al.* Rapid turnover of long noncoding RNAs and the evolution of gene expression.
PLoS Genet. **8,** e1002841 (2012).

80. Lewis, J. J., van der Burg, K. R. L., Mazo-Vargas, A. & Reed, R. D. ChIP-Seq-Annotated Heliconius
erato Genome Highlights Patterns of cis-Regulatory Evolution in Lepidoptera. *Cell Rep.* **16,**
2855–2863 (2016).

81. Gilad, Y., Oshlack, A. & Rifkin, S. A. Natural selection on gene expression. *Trends Genet.* **22,** 456–
461 (2006).

82. Bailey, S. D. *et al.* Noncoding somatic and inherited single-nucleotide variants converge to
promote ESR1 expression in breast cancer. *Nat. Genet.* **48,** 1260–1266 (2016).

83. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal **17,** 10–12 (2011).

84. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform.
Bioinformatics **25,** 1754–1760 (2009).

85. Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic features.
Bioinformatics **26,** 841–842 (2010).

86. Kuhn, R. M., Haussler, D. & Kent, W. J. The UCSC genome browser and associated tools. *Brief.*
Bioinform. **14,** 144–161 (2013).

87. Zhao, H. *et al.* CrossMap: a versatile tool for coordinate conversion between genome
assemblies. *Bioinformatics* **30,** 1006–1007 (2014).

88. Kent, W. J., Baertsch, R., Hinrichs, A., Miller, W. & Haussler, D. Evolution's cauldron: duplication,
deletion, and rearrangement in the mouse and human genomes. *Proc. Natl. Acad. Sci. U. S. A.*
100, 11484–11489 (2003).

89. Luo, X., Chae, M., Krishnakumar, R., Danko, C. G. & Kraus, W. L. Dynamic reorganization of the
AC16 cardiomyocyte transcriptome in response to TNFα signaling revealed by integrated
genomic analyses. *BMC Genomics* **15,** 155 (2014).

90. Kozomara, A. & Griffiths-Jones, S. miRBase: annotating high confidence microRNAs using deep
sequencing data. *Nucleic Acids Res.* **42,** D68–73 (2014).

91. Danko, C. G. *et al.* Signaling pathways differentially affect RNA polymerase II initiation, pausing,
and elongation rate in cells. *Mol. Cell* **50,** 212–222 (2013).

92. Neph, S. *et al.* BEDOPS: high-performance genomic feature operations. *Bioinformatics* **28,**
1919–1920 (2012).

93. Pham, H., Kearns, N. A. & Maehr, R. Transcriptional Regulation with CRISPR/Cas9 Effectors in
Mammalian Cells. *Methods Mol. Biol.* **1358,** 43–57 (2016).

Acknowledgements: We thank M. Jin for assistance in establishing the magnetic separation of
CD4+ T-cells, L. Core, H. Kwak, N. Fuda, and I. Jonkers for assistance troubleshooting the PRO-seq
library prep, and A. Wetterau for preparing nuclei for mouse and rat CD4+ T-cells. Work in this
publication was supported by generous seed grants from the Cornell University Center for
Vertebrate Genomics (CVG), the Center for Comparative and Population Genetics (3CPG), NHGRI
(National Human Genome Research Institute) grant HG009309 to CGD, NHLBI (National Heart,
Lung, and Blood Institute) grant UHL129958A to CGD and JTL, NIGMS (National Institute of General
Medical Sciences) grant GM102192 to AS, NHGRI (National Human Genome Research Institute)
grant HG0070707 to AS and JTL, NIH/NIDDK DK058110 to WLK, and CPRIT RP160319 to WLK.
The content is solely the responsibility of the authors and does not necessarily represent the official
views of the US National Institutes of Health. Finally, we would like to thank the anonymous human
and non-human primate donors who gave blood in support of this study.

Author contributions: LAC, BAM, CGD, EJR, and ETW performed CD4+ T-cell extraction, validation,
and PRO-seq experiments. CGD, ZW, TC, ALM, LAC, and ND analyzed the data. CGD, AS, JTL, WLK,
and SAC supervised data collection and analysis. CGD and AS wrote the paper with input from the
other authors.

Competing financial interests: The authors declare no competing financial interests.

Author information: PRO-seq data was deposited into the Gene Expression Omnibus database
under accession number GSE85337. All data analysis scripts and software are publicly available on
GitHub: https://github.com/Danko-Lab/CD4-Cell-Evolution.

  Figure Legends:

  Fig. 1 | Maps of primary transcription in CD4+ T-cells. (a) CD4+ T-cells were isolated from the
blood or spleen of individuals from five vertebrate species, including human, chimpanzee, rhesus
macaque, mouse, and rat. **(b)** Hierarchical clustering of PRO-seq signal intensities in gene bodies
groups CD4+ T-cell samples first by treatment condition and second by species. The color scale
represents Spearman's rank correlation between normalized transcription levels in active gene
bodies. Colored boxes (top) represents the species and treatment condition of each sample. **(c)** MA
plot shows the $\log_2$ fold-change following π treatment in human CD4+ T-cells (y-axis) as a function
of the mean transcription level in GENCODE annotated genes (x-axis). Red points indicate
statistically significant changes ($p < 0.01$). Several classical response genes that undergo well-
documented changes in transcript abundance following CD4+ T-cell activation (e.g., *IL2*, *IFNG*,
TNFα, and *EGR3*) are marked. **(d)** Enrichment of TF binding motifs in TREs that increase
transcription levels following π treatment in the indicated species compared to TREs whose
transcription abundance does not change. Table shows the Bonferroni corrected p-value, based on
a Fisher's exact test (circle size), and the fold-enrichment over a group of unchanged background
sequences (color scale). Motif logos and the candidate transcription factor or Cis-BP motif ID are
shown. **(e)** Heatmaps show the distribution of PRO-seq (red and blue indicate transcription on the
plus and minus strand, respectively), ChIP-seq for H3K27ac, H3K4me1, and H3K4me3, and DNase-
I-seq signal intensity. Plots are centered on transcriptional regulatory elements (TREs) predicted
in untreated human CD4+ T-cells using dREG-HD (see Online Methods). All plots are ordered based
on the maximum dREG score in the window.

Fig. 2 | Frequency of changes in TRE transcription. (a) The fractions of TREs active in untreated
CD4+ T-cells that are present in the human reference genome and are conserved across all species
(blue), are not detectable and are therefore inferred as gains or losses (teal-white) or undergo
significant changes (green) in at least one species, or fall in regions for which no ortholog occurs in
at least one of the indicated genomes (pink). Inferred gains or losses are colored according to the
FDR corrected p-value associated with changes in RNA polymerase abundance (deSeq2). Plots
labeled "Primate" illustrate frequency of changes in a three-way comparison of human, chimpanzee,
and rhesus macaque focusing on the untreated condition, whereas those labeled "Mammal"
summarize a five-way comparison also including rat and mouse. π treatment denotes a comparison
between human untreated and PMA+Ionomycin treated CD4+ T-cell samples. **(b)** Boxplots show
the ChIP-seq signal near dREG sites classified as conserved, gains, losses, or complete losses for the
indicated chromatin or DNA modification in units of reads per kilobase. The box represents the
25th and 75th percentile. Whiskers represent 1.5 times the interquartile range, and points outside
of this range are not shown.

Fig. 3 | Evolutionary changes in TRE transcription correlate with DNA sequence
conservation. (a) Mean phyloP scores near TFBSs that are conserved (red), gained (blue), or lost
(cyan) on the human branch. Motifs (score > 10) are at least 100 bp from the nearest annotated
exon. **(b)** UCSC Genome Browser track shows transcription near *SGPP2* and *FARSB* in untreated (U)
and PMA+ionomycin (π) treated CD4+ T-cells isolated from the indicated primate species. PRO-

seq tracks show transcription on the plus (red) and minus (blue) strands. Axes for the PRO-seq
data are in units of reads per kilobase per million mapped (RPKM). Transcription units inferred
from the PRO-seq data are shown above the plot. dREG tracks show the distribution of dREG signal.
The Green et. al. (ref[64]) selective sweep scan track (top) represents the enrichment of derived
alleles in modern human where Neanderthal has the ancestral allele. Points below the line
represent a statistically significant number of derived alleles in modern human (line indicates a Z-
score of -2). Net synteny tracks show the position of regions that have one-to-one orthologs in the
chimpanzee and rhesus macaque genomes. **(c)** Luciferase signal driven by the *SGPP2* promoter or
the internal enhancer in MCF-7 cells using DNA from each primate species. Bars show the mean
fold-induction following 3 hours of stimulation with TNFα. Error bars represent the standard error
of the mean. Red ** denotes $p < 1e-3$ by a two-tailed t-test. **(d)** Transcription of *SGPP2* using
primers targeting intron 1 following 0 or 40 min. of TNFα treatment after silencing the indicated
TRE using dCAS9-KRAB. Bars represent the median of three independent biological replicates of
two gRNAs targeting the promoter, three targeting the internal enhancer, and four targeting the
upstream enhancer. Error bars represent the standard error. Red ***** denotes $p < 5e-2$ and ****** $p < 5e-$
3 by a two-tailed t-test. **(e)** INSIGHT estimates of the fraction of nucleotides under selection (ρ), the
expected density of segregating polymorphisms under weak negative selection (E[Pw]/kbp), or the
expected density of human nucleotide substitutions driven by positive selection (E[Dp]/kbp) in
human populations in the indicated class of sites. Red ***** denotes conditions significantly enriched
over random background sequences ($p < 0.01$; two-tailed X^2-test).
64
Fig. 4 | Changes in non-coding RNA transcription predict changes in gene transcription. (a)
The fraction of each indicated class of RNAs that undergo changes in transcription in human CD4+
T-cells (see Online Methods). The relationships among the indicated classes of transcription units
are depicted at top. **(b)** Scatterplot shows the magnitude of changes in transcription predicted for
protein-coding genes using changes in the transcription of nearby non-coding RNAs (y-axis) as a
function of changes observed (x-axis). The line has a slope of 1 and an intercept of 0.
71
Fig. 5 | TRE conservation correlates with loop interactions and distance to gene promoters.
(a) UCSC Genome Browser tracks show transcription, dREG signal, and ChIA-PET loop interactions
near the *CCR7* superenhancer in the human genome.   PRO-seq tracks show transcription on the
plus (red) and minus (blue) strands in units of RPKM. Net synteny tracks show regions of one-to-
one orthology with the chimpanzee and rhesus macaque genomes. **(b)** Scatterplot shows the
percentage of TREs conserved among all three primate species (y-axis) as a function of distance,
either upstream or downstream, from the nearest annotated protein-coding transcription start site
(x-axis). The size of each point represents the amount of data in the corresponding distance bin.
(c) The percentage of all dREG sites that are conserved in each indicated class of TRE. TREs are
separated into three bins based on the distance relative to the nearest transcription start site.
Error bars reflect a 1,000-sample bootstrap.
83
Fig. 6 | Stabilizing selection on protein coding gene transcription. (a-b) Scatterplot shows
promoter conservation **(a)** or DNA sequence conservation **(b)** as a function of the number of loop
interactions made by that site to distal sites across the genome (x-axis). **(c-d)** TRE conservation **(c)**

or DNA sequence conservation **(d)** as a function of the number of loop interactions made by the
sequence at the distal end of the loop interaction (x-axis). In all panels the size of each point is
proportional to the number of examples in the corresponding bin, following the scale shown in the
center.
91
92
93   **<u>Supplementary Figure Legends:</u>**
94
95   **Supplementary Fig. 1 | Validation of CD4+ cell enrichment by flow cytometry.** Representative
96   plots of CD4 expression in human, chimpanzee, and rhesus macaque PBMC, before (left) and after
97   (right) CD4 microbead enrichment. Percentage of total live lymphocytes shown.
98
Supplementary Fig. 2 | Transcription abundance in the gene bodies of T-cell lineage specific
markers. Plots show normalized expression (log2 scale) of transcription factors and cytokines that
mark specific subsets of CD4+ T-cell population in the species indicated below the plot. Each point
represents the transcription of the indicated gene in a different untreated T-cell sample. The bar
indicates the mean in each species. In all cases read counts were limited to regions of orthology in
the bodies of genes indicated on each plot.
105
Supplementary Fig. 3 | Principal component analysis (PCA) of CD4+ T-cell PRO-seq libraries.
Scatterplots show the first five principal components (PC) from CD4+ T-cell PRO-seq libraries. PCA
was constructed using regions of orthology in all five species in the bodies of transcription units
identified by a three state hidden Markov model. The fraction of the variance explained by each PC
is shown in parentheses. The key shown below the plot indicates the species and treatment
condition of each point.
112
Supplementary Fig. 4 | Sample correlation plotted as a function of estimated evolutionary
divergence time between species. Scatterplot shows the evolutionary divergence time (X-axis) as
a function of Spearman's correlation in gene body transcription between each sample collected in
the untreated condition and the mean gene expression in untreated human CD4+ T-cells (Y-axis).
The red line shows the best linear fit and dotted lines indicate the 99% confidence interval. We
assume the following evolutionary divergence estimates for each species pair with respect to
human, 12 MYR for chimp-human [Moorjani et. al. (2016)], 25 MYR for human-rhesus [Rogers
(2013)], and 75 MYR for human-rodent [Chinwalla et. al. (2002)].
121
Supplementary Fig. 5 | Changes in gene transcription following PMA+Ionomycin treatment in
chimpanzee and rhesus macaque CD4+ T-cells. (a-b) MA plot shows the log-2 fold change
following π treatment (y-axis) as a function of the mean transcription level in GENCODE annotated
genes (x-axis) in data from chimpanzee (left) and rhesus macaque (right) CD4+ T-cells. Red points
indicate statistically significant changes (p < 0.01). Several classical response genes that undergo
well-documented changes in transcript abundance following CD4+ T-cell activation (e.g., *IL2*, *IFNG*,
TNF, and *EGR3*) are marked. **(c)** UCSC genome browser track shows transcription in the *IFNG* locus
in untreated (U) and PMA+ionomycin (π) treated CD4+ T-cells isolated from the primate species

indicated at left.   PRO-seq tracks show transcription on the plus (red) and minus (blue) strands.
dREG tracks show the distribution of dREG signal. The net-synteny tracks show the fraction of the
genomic area that is mappable in the indicated species. The location of transcription units inferred
in the common ancestor of human and chimpanzee, and the location of RefSeq gene annotations,
are shown at the bottom. **(d-f)** Scatterplots show the correlation between changes in gene
expression (log-2 scale) following π treatment in the species indicated on the axes. Color scale
indicates the density of points in the region.
137
138   **Supplementary Fig. 6 | Evolutionary changes in TREs. (a)** Venn diagram illustrating raw
139   changes in TREs among primate species. In all cases, TREs were discovered in untreated CD4+ T-
140   cells using dREG (threshold > 0.3). **(b)** Q-Q plot showing observed p-values (deSeq2 in human
141   compared to the other two primate species) among TREs that were not identified by dREG in at
142   least one species (red), all TREs identified (black), and a set of conserved TREs (gray).
143
144   **Supplementary Fig. 7 | Evolutionary changes in TREs correlate with chromatin and DNA**
145   **modifications.** ChIP-seq signal for H3K27ac and H3K4me1 near dREG sites classified as gains,
146   losses, or complete losses of TRE signal (dREG score < 0.05) on the human branch.
147
Supplementary Fig. 8 | PhyloP scores in transcription factor (TF) binding motifs. (a)
Evolutionary conservation centered on matches to a TF binding motif at the indicated cut off score
(left), or adjusted for distance to the nearest annotated transcription start site by subsampling
(right) **(b)** PhyloP scores that fall within the binding motifs recognized by STAT2 (M6494_1.02),
YY1 (M4490_1.02), CREB1 (M6180_1.02), and ELF1 (M6203_1.02). In all cases motifs fall in dREG-
HD that are gained (blue) or lost (cyan) on the human branch, or are conserved among all primate
species (red). **(c)** The distribution of human derived alleles near dREG sites that are gained (blue)
or lost (cyan) on the human branch, or are conserved among all primate species (red).
156
Supplementary Fig. 9 | Candidate causal DNA sequence differences underlying changes in
SGPP2 **transcription.** UCSC genome browser track shows transcription near *SGPP2* and *FARSB* in
untreated (U) and PMA+ionomycin (π) treated human CD4+ T-cells or in human MCF-7 cells. PRO-
seq tracks show transcription on the plus (red) and minus (blue) strands. Axes for the PRO-seq
data are in units of reads per kilobase per million mapped (RPKM) or in raw reads (MCF-7). dREG
tracks show the distribution of dREG signal. Heatmap (top) shows Hi-C signal in GM12878
lymphoblastoid cell lines. Insert (bottom) shows lack of orthology in chimpanzee and rhesus
macaque in an active TRE (human) that binds a number of TFs in ENCODE cell lines (left) and
substitutions in NF-kB binding motifs near *SGPP2*. Two motif occurrences in the proximal
promoter were bound by RELA, a subunit of NF-kB, based on human ChIP-seq data in ENCODE cell
lines (green boxes). Positions where human carries a derived allele are indicated by yellow
highlights. PRO-seq reads matched the human reference allele in all positions (15/ 15 reads match
C and 26/ 26 match the reference T allele in the NF-kB binding site in the promoter; 11/ 11 reads
match the G and 11/ 11 match the T reference allele in the NF-kB binding site in the promoter; and
24/ 24 reads support the TG human reference sequence in the internal enhancer). Scatterplots
show the relative frequencies of the human allele in RELA (NF-kB) ChIP-seq data matching NF-kB

binding QTLs that mimic the human and ancestral alleles, while controlling for the flanking
sequence indicated below the plot. The red dot denotes the mean. All four human-specific DNA
sequence changes in NF-kB binding motifs in the proximal promoter together show trend toward
higher NF-kB binding in human (p = 0.017, using Fisher's method to combine p-values).
177

178   **Supplementary Fig. 10 | Luciferase assays for TREs identified near *SGPP2*.** The Y-axis shows
179   the luciferase signal driven by the *SGPP2* promoter or the internal enhancer in MCF-7 cells using
180   DNA from each primate species following 3 hours of stimulation with TNFα or vehicle control. Bars
181   show the mean luciferase activity in each species, over the empty vector and renilla controls. Error
182   bars represent the standard error of the mean.
183

Supplementary Fig. 11 | Adaptive substitutions in specific TF binding motifs. Adaptive
substations in TF binding motifs (TFBM) occurring commonly (>900 times) in human lineage-
specific dREG-HD sites. Columns denote the TF name annotated in CisBP (TF), number of sites
(Sites), the number of bases (Bases), the expected number of adaptive substitutes per kilobase
(E[A]), the standard error in the expected substitutions per kilobase (E[A]_stderr), and the
estimated number of adaptive substitutions (# Adaptive Substitutions). TFBSs may be bound by
any TF that recognizes a similar motif. TFBM in which E[A] is significantly larger than 0 are
highlighted in bold fold. The estimated number of adaptive substitutions for each of these sites is
shown.
193

Supplementary Fig. 12 | Discovery of transcription units (TU) in primate T-cells. (a) A novel
three-state hidden Markov model (HMM) was used to discover transcription units. States
correspond to non-transcribed background sequence, transcribed sequence, and post polyA
transcription. TUs were classified into one of seven classes as indicated in the cartoon. **(b)** The
number and fraction of transcription units that fall into each TU classification. **(c)** Example of the
hidden Markov model (HMM) in a typical region. TUs largely agree with RefSeq gene annotations
when available.
201

Supplementary Fig. 13 | DNA sequence conservation as a function of genomic distance to the
nearest start site. Scatterplot shows the percentage of TREs undergoing complete gains and
losses (left), undergoing a partial change in the abundance of Pol II (center), or that are not
alignable between species (right) as a function of distance from the nearest annotated transcription
start site (x-axis). The size of each point represents the amount of data in the corresponding
distance bin.
208

209   **Supplementary Fig. 14 | Evolutionary conservation of DNA sequence mirrors functional**
210   **conservation at looped- and un-looped enhancers.** Cumulative distribution function of phyloP
211   scores from the 100-way alignments in the indicated class of dREG site. The insert shows the
212   fraction of sites in each class exceeding a phyloP score cutoff of 0.75.

Fig. 1 | Maps of primary transcription in CD4+ T-cells. **(a)** CD4+ T-cells were isolated from the blood or spleen of individuals from five vertebrate species, including human, chimpanzee, rhesus macaque, mouse, and rat. **(b)** Hierarchical clustering of PRO-seq signal intensities in gene bodies groups CD4+ T-cell samples first by treatment condition and second by species. The color scale represents Spearman's rank correlation between normalized transcription levels in active gene bodies. Colored boxes (top) represents the species and treatment condition of each sample. **(c)** MA plot shows the log2 fold-change following π treatment in human CD4+ T-cells (y-axis) as a function of the mean transcription level in GENCODE annotated genes (x-axis). Red points indicate statistically significant changes (p < 0.01). Several classical response genes that undergo well-documented changes in transcript abundance following CD4+ T-cell activation (e.g., IL2, IFNG, TNFa, and EGR3) are marked. **(d)** Enrichment of TF binding motifs in TREs that increase transcription levels following π treatment in the indicated species compared to TREs whose transcription abundance does not change. Table shows the Bonferroni corrected p-value, based on a Fisher's exact test (circle size), and the fold-enrichment over a group of unchanged background sequences (color scale). Motif logos and the candidate transcription factor or Cis-BP ID are shown. **(e)** Heatmaps show the distribution of PRO-seq (red and blue indicate transcription on the plus and minus strand, respectively), ChIP-seq for H3K27ac, H3K4me1, and H3K4me3, and DNase-I-seq signal intensity. Plots are centered on transcriptional regulatory elements (TREs) predicted in untreated human CD4+ T-cells using dREG-HD (see Online Methods). All plots are ordered based on the maximum dREG score in the window.

Fig. 2 | Frequency of changes in TRE transcription. (a) The fractions of TREs active in untreated CD4+ T-cells that are present in the human reference genome and are conserved across all species (blue), are not detectable and are therefore inferred as gains or losses (teal-white) or undergo significant changes (green) in at least one species, or fall in regions for which no ortholog occurs in at least one of the indicated genomes (pink). Inferred gains or losses are colored according to the FDR corrected p-value associated with changes in RNA polymerase abundance (deSeq2). Plots labeled "Primate" illustrate frequency of changes in a three-way comparison of human, chimpanzee, and rhesus macaque focusing on the untreated condition, whereas those labeled "Mammal" summarize a five-way comparison also including rat and mouse. π treatment denotes a comparison between human untreated and PMA+Ionomycin treated CD4+ T-cell samples. **(b)** Boxplots show the ChIP-seq signal near dREG sites classified as conserved, gains, losses, or complete losses for the indicated chromatin or DNA modification in units of reads per kilobase. The box represents the 25th and 75th percentile. Whiskers represent 1.5 times the interquartile range, and points outside of this range are not shown.

Fig. 3 | Evolutionary changes in TRE transcription correlate with DNA sequence conservation. (a) Mean phyloP scores near TFBSs that are conserved (red), gained (blue), or lost (cyan) on the human branch. Motifs (score > 10) are at least 100 bp from the nearest annotated exon. **(b)** UCSC Genome Browser track shows transcription near SGPP2 and FARSB in untreated (U) and PMA+ionomycin (π) treated CD4+ T-cells isolated from the indicated primate species. PRO-seq tracks show transcription on the plus (red) and minus (blue) strands. Axes for the PRO-seq data are in units of reads per kilobase per million mapped (RPKM). Transcription units inferred from the PRO-seq data are shown above the plot. dREG tracks show the distribution of dREG signal. The Green et. al. (ref64) selective sweep scan track (top) represents the enrichment of derived alleles in modern human where Neanderthal has the ancestral allele. Points below the line represent a statistically significant number of derived alleles in modern human (line indicates a Z-score of -2). Net synteny tracks show the position of regions that have one-to-one orthologs in the chimpanzee and rhesus macaque genomes. **(c)** Luciferase signal driven by the SGPP2 promoter or the internal enhancer in MCF-7 cells using DNA from each primate species. Bars show the mean fold-induction following 3 hours of stimulation with TNFa. Error bars represent the standard error of the mean. Red ** denotes p < 1e-3 by a two-tailed t-test. **(d)** Transcription of SGPP2 using primers targeting intron 1 following 0 or 40 min. of TNFa treatment after silencing the indicated TRE using dCAS9-KRAB. Bars represent the median of three independent biological replicates of two gRNAs targeting the promoter, three targeting the internal enhancer, and four targeting the upstream enhancer. Error bars represent the standard error. Red * denotes p < 5e-2 and ** p < 5e-3 by a two-tailed t-test. **(e)** INSIGHT estimates of the fraction of nucleotides under selection (ρ), the expected density of segregating polymorphisms under weak negative selection (E[Pw]/kbp), or the expected density of human nucleotide substitutions driven by positive selection (E[Dp]/kbp) in human populations in the indicated class of sites. Red * denotes conditions significantly enriched over random background sequences (p < 0.01; two-tailed Chi-squared-test).

Fig. 4 | Changes in non-coding RNA transcription predict changes in gene transcription. (a) The fraction of each indicated class of RNAs that undergo changes in transcription in human CD4+ T-cells (see Online Methods). The relationships among the indicated classes of transcription units are depicted at top. **(b)** Scatterplot shows the magnitude of changes in transcription predicted for protein-coding genes using changes in the transcription of nearby non-coding RNAs (y-axis) as a function of changes observed (x-axis). The line has a slope of 1 and an intercept of 0.

Fig. 5 | TRE conservation correlates with loop interactions and distance to gene promoters. (a) UCSC Genome Browser tracks show transcription, dREG signal, and ChIA-PET loop interactions near the CCR7 superenhancer in the human genome. PRO-seq tracks show transcription on the plus (red) and minus (blue) strands in units of RPKM. Net synteny tracks show regions of one-to-one orthology with the chimpanzee and rhesus macaque genomes. **(b)** Scatterplot shows the percentage of TREs conserved among all three primate species (y-axis) as a function of distance, either upstream or downstream, from the nearest annotated protein-coding transcription start site (x-axis). The size of each point represents the amount of data in the corresponding distance bin. **(c)** The percentage of all dREG sites that are conserved in each indicated class of TRE. TREs are separated into three bins based on the distance relative to the nearest transcription start site. Error bars reflect a 1,000-sample bootstrap.

Fig. 6 | Stabilizing selection on protein coding gene transcription. (a-b) Scatterplot shows promoter conservation **(a)** or DNA sequence conservation **(b)** as a function of the number of loop interactions made by that site to distal sites across the genome (x-axis). **(c-d)** TRE conservation **(c)** or DNA sequence conservation **(d)** as a function of the number of loop interactions made by the sequence at the distal end of the loop interaction (x-axis). In all panels the size of each point is proportional to the number of examples in the corresponding bin, following the scale shown in the center.