Natural Selection has Shaped Coding and Non-coding Transcription in Primate CD4+ T-cells

Charles G. Danko[1,2,*], Zhong Wang[1], Edward J. Rice[1], Tinyi Chu[1,3], Andre L. Martins[1],
Elia Tait Wojno[1,4], John T. Lis[5], W. Lee Kraus[6,7], & Adam Siepel[8,*]

[1] Baker Institute for Animal Health, College of Veterinary Medicine, Cornell University, Ithaca, NY 14853.

[2] Department of Biomedical Sciences, College of Veterinary Medicine, Cornell University, Ithaca, NY 14853.

[3] Graduate field of Computational Biology, Cornell University, Ithaca, NY 14853.

[4] Department of Microbiology & Immunology, College of Veterinary Medicine, Cornell University, Ithaca, NY 14853.

[5] Department of Molecular Biology and Genetics, Cornell University, Ithaca, NY 14853.

[6] Laboratory of Signaling and Gene Regulation, Cecil H. and Ida Green Center for Reproductive Biology Sciences, University of Texas Southwestern Medical Center, Dallas, TX 75390.

[7] Division of Basic Research, Department of Obstetrics and Gynecology, University of Texas Southwestern Medical Center, Dallas, TX 75390.

[8] Simons Center for Quantitative Biology, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY 11724.

[*] **Address correspondence to:**

Charles G. Danko, Ph.D.   Adam Siepel, Ph.D.
Baker Institute for Animal Health   Simons Center for Quantitative Biology
Cornell University   Cold Spring Harbor Laboratory
Hungerford Hill Rd.   1 Bungtown Rd.
Ithaca, NY 14853   Cold Spring Harbor, NY 11724
Phone: 607-256-5620   Phone: 516-367-6922
E-mail: dankoc@gmail.com   E-mail: asiepel@cshl.edu

Abstract:
Transcriptional regulatory changes have been shown to contribute to phenotypic differences between species, but many questions remain about how gene expression evolves. Here we report the first comparative study of nascent transcription in primates. We used PRO-seq to map actively transcribing RNA polymerases in resting and activated CD4+ T-cells in multiple human, chimpanzee, and rhesus macaque individuals, with rodents as outgroups. This approach allowed us to directly measure active transcription separately from post-transcriptional processes. We observed general conservation in coding and non-coding transcription, punctuated by numerous differences between species, particularly at distal enhancers and non-coding RNAs. Transcription factor binding sites are a primary determinant of transcriptional differences between species. We found evidence for stabilizing selection on gene expression levels and adaptive substitutions associated with lineage-specific transcription. Finally, rates of evolutionary change are strongly correlated with long-range chromatin interactions. These observations clarify the role of primary transcription in regulatory evolution.

Following decades of speculation that changes in the regulation of genes could be a potent force in the evolution of form and function[1-3], investigators have now empirically demonstrated the evolutionary importance of gene regulation across the tree of life[4-12]. Changes in gene expression are primarily driven by mutations to non-coding DNA sequences, particularly those that bind sequence-specific transcription factors[13]. Accordingly, adaptive nucleotide substitutions at transcription factor binding sites (TFBSs)[9,10,14-16] and gains and losses of TFBSs[17-25] both appear to make major contributions to the evolution of gene expression. These events are believed to modify a variety of rate-limiting steps early in transcriptional activation[26]. Transcriptional activity is generally correlated with epigenomic and structural features such as post-translational modifications to core histones, the locations of architectural proteins such as CTCF, and the organization of topological associated domains. Like TFBSs, these features display general conservation across species, yet do exhibit some variation, which correlates with differences in gene expression[8,24,27-29].

Nevertheless, many open questions remain about the relative roles of TFBSs, chromatin organization, and posttranscriptional regulation in the evolution of gene expression. There is a surprisingly limited correlation between differences in binding events and differences in mRNA expression levels[30-32]. Possible reasons for this discordance include non-functional TF binding[30,31,33], compensatory gains and losses of TFBSs[20,34-37], difficulties associating distal enhancers with target genes[38], and a dependency of TF function on chromatin or chromosomal organization[39]. In addition, it remains unclear to what degree epigenomic differences between species are causes and to what degree they are effects of differences in gene expression. Finally, some changes in mRNA expression appear to be "buffered" at the post-transcriptional level[40-42].

One reason why it has been difficult to disentangle these contributions is that gene expression is typically measured in terms of the abundance of mRNA, which is subject to post-transcriptional processing[43] and therefore is an indirect measure of the transcription of genes by RNA polymerase II. An alternative and complementary approach is to measure the production of nascent RNAs using Precision Run-On and sequencing (PRO-seq) and related technologies[44-48]. Nascent RNA sequencing methods directly measure active transcription and are highly sensitive to immediate and transient transcriptional responses to stimuli[49]. They can detect active regulatory elements as well as target genes, because these elements themselves display distinctive patterns of transcription, which are obscured in RNA-seq data owing to rapid degradation[33,50,51]. Indeed, the latest nascent RNA sequencing methods, such as PRO-seq[45], in combination with new computational tools for regulatory element prediction[52], serve as powerful single-assay systems for both identifying regulatory elements and measuring transcription levels.

With these advantages in mind, we undertook a genome-wide comparative analysis of transcription in primates using PRO-seq. Our comparison of PRO-seq data across species revealed overall conservation in the transcription of both coding and non-coding elements, but also uncovered numerous differences between species. Together, our observations provide new insights into the evolution of transcription in primates.

Patterns of transcription in resting and activated CD4+ T-cells

We developed nucleotide-resolution maps of RNA polymerase for CD4+ T-cells isolated from five mammalian species. Samples were collected under resting and activated conditions from

three unrelated individuals representing each of three primate species, humans, chimpanzees, and
rhesus macaques, spanning ~25-30 million years of evolution (MYR) (**Fig. 1a**). Resting samples
were also collected from a single individual in each of two rodent species, mouse and rat, which
together serve as an outgroup to the primates (~80 MYR divergence). PRO-seq[45,48] libraries were
sequenced to a combined depth of 873 million uniquely mapped reads (~78-274 million per
species) (**Supplementary Table 1**). Flow cytometry was used to validate the purity of isolated
CD4+ cells (**Supplementary Fig. 1**), and measurements of transcriptional activity of T-cell subset
markers for T-helper type 1 (Th1), Th2, Th17, T-regulatory, and T-follicular helper cells were used
to demonstrate that the population of CD4+ T-cell subsets within the total CD4+ population is
largely similar among these mammalian species (**Supplementary Fig. 2**). Principal component
analysis ranked the rodent vs. primate species, variation within the primate species, and the
treatment condition as the first, second, and third sources of variation, respectively, in the complete
dataset (**Supplementary Fig. 3**). Similarly, hierarchical clustering of these data together with data
from other cell types grouped primate samples first by cell type or treatment condition and
subsequently by species (**Fig. 1b**). These results demonstrate that genome-wide patterns of
transcription remain generally concordant in CD4+ T-cells in the species we examined, especially
within the primates.
Before comparing species, we evaluated differences in transcription between resting and
activated conditions. Human CD4+ T-cells activated using PMA and ionomycin (π) underwent
significant changes in transcription levels at 5,945 genes ($p < 0.01$, deSeq2[53]), including classical
early response genes such as *IL2*, *IL2RA*, and *EGR3* (**Fig. 1c**). Parallel analyses in chimpanzee and
rhesus macaque revealed comparable transcription programs both before and following π
treatment (**Supplementary Fig. 4a-b**). We identified a core set of 2,953 genes that undergo
evolutionarily conserved transcriptional changes following 30 min. of π-treatment in all three
species, including many of the classical response genes (e.g., IFNG, TNFα, IL2, and IL2RA), as well as
numerous novel genes and lincRNAs (**Supplementary Fig. 4a-c**). Fold-changes of these core genes
following π treatment are highly correlated and similar in magnitude among all three primate
species (R > 0.92; **Supplementary Fig. 4d-f**), suggesting that species-specific differences in T-cell
activation magnitude[54] may occur primarily during the initial signaling events and are largely
bypassed by π-treatment.
To shed light on the mechanisms underlying transcriptional changes immediately following
π treatment, we used dREG[52] to identify 25,513 active transcriptional regulatory elements (TREs),
including promoters and enhancers, in human CD4+ T-cells, based on patterns of enhancer-
templated RNA (eRNA) transcription evident from PRO-seq data (**Fig. 1d**). These predicted TREs
displayed many of the expected marks of regulatory function in human CD4+ T-cells, including
acetylation of histone 3 lysine 27 (H3K27ac), and mono- and trimethylation of histone 3 lysine 4
(H3K4me1 and H3K4me3)[55], consistent with previous observations[33,50,52]. We identified 7,340
TREs with π-dependent changes in RNA polymerase abundance ($p < 0.01$, deSeq2). TREs with
transcriptional increases following π-treatment were enriched for DNA sequence motifs recognized
by NF-kB, NFAT, and the AP-1 heterodimers FOS and JUN (**Fig. 1e**), all transcription factors
activated by canonical T-cell receptor signaling[56]. Motifs for other calcium-responsive factors such
as MEF2A and CREB were also enriched in up-regulated TREs, consistent with the activation of
calcium signaling by Ionomycin, a calcium ionophore. Downregulated TREs were enriched for

different motifs, including those for FOXO1, ELK/ELF/ETS, and SMARCC2 (**Supplementary Fig. 5**).
The enrichment of the forkhead box (FOXO1) binding motif is concordant with findings that FOXO1
is required for naive T-cell homeostasis[57], and suggests that this TF is down-regulated as naive T-
cells differentiate. Motif analysis in the chimpanzee and rhesus macaque genomes revealed a
similar composition of motifs putatively involved in mediating transcriptional changes following π
treatment (**Supplementary Fig. 6**). Thus, the core regulatory principles responsible for T-cell
signaling and activation appear to remain broadly conserved across primate evolution.
135

Rapid evolutionary changes in transcribed enhancers
We focused an initial comparative analysis on evolutionary changes in active enhancers.
We used dREG to identify 53,476 TREs that are active in either treated or untreated CD4+ T-cells of
at least one primate species (ranging between 25,269-35,343 TREs in each species). TREs were
classified as promoters (n = 7,114) or enhancers (n = 21,121) based on their proximity to gene
annotations and the stability of the associated transcription units predicted using the order and
orientation of U1 and polyadenylation site motifs[50] (**Supplementary Fig. 7**, see Methods). We
found that 62% of enhancers changed Pol II transcriptional activity in at least one of the three
primate species and nearly 80% at the longer evolutionary distance between primates and rodents
(**Fig. 2a**). Conservation was highest between human and chimpanzee (10-20% of TREs change),
roughly consistent with recent estimates based on the distribution of H3K27ac[10]. Enhancers are
completely gained or lost at nearly eight times the rate of promoters (38% of enhancers; 7% of
promoters; p < 2.2e-16 by Fisher's exact test), consistent with recent observations based on
H3K27ac and H3K4me1[24]. By contrast, π treatment is rarely associated with complete differences
in the location of transcribed TREs and the rate of differences is similar at promoters and enhancers
(**Fig. 2a**).
Next we tested whether evolutionary changes in transcriptional activity correlate with the
enrichment of other active enhancer marks. We used the known phylogeny for these species to
identify likely lineage-specific gains and losses of enhancer activity in human CD4+ T-cells, and
examined independent sources of genomic data[55] at the corresponding genomic locations. Overall,
predicted lineage-specific human enhancers were enriched for both active and repressive enhancer
marks (**Fig. 2b; Supplementary Fig. 8**). Whereas human gains were enriched for high levels of
H3K27ac, sites with reduced transcriptional activity in humans showed much lower enrichments of
this mark of active enhancers. Furthermore, locations of complete human lineage-specific loss of
dREG signal displayed levels of H3K27ac approaching the background, consistent with a complete
loss of enhancer activity (**Fig. 2b**). Intriguingly, H3K4me1, which marks active and some inactive
enhancers[58,59], was enriched at predicted human lineage-specific losses in enhancer activity (**Fig.
2b**), indicating that, at least in some cases, an active ancestral primate enhancer retains a 'poised'
chromatin state in human, despite losing transcriptional activity and H3K27ac. This result suggests
a possible model of TRE assembly and activation in which changes in poised and active marks are
evolutionarily distinct events.
167

168 **Transcriptional changes correlate with DNA sequence differences**
169 To investigate whether changes in TRE activity are accompanied by changes in DNA
170 sequence, we compared phyloP sequence conservation scores[60] at conserved TREs with scores at

TREs that display evolutionary changes in transcription. Because signatures of sequence conservation in TREs are likely to be most pronounced in transcription factors binding sites (TFBS), we restricted our sequence conservation analyses to strong matches to 1,964 human TF binding motifs. We grouped these binding motifs by their DNA sequence similarity into 567 clusters and we excluded motifs for TFs that are transcriptionally inactive in CD4+ T-cells[61,62].

TFBSs found in transcriptionally conserved dREG sites showed a marked enrichment for higher phyloP scores relative to surrounding regions, indicating that these TFBSs are enriched for evolutionary conservation in their DNA sequences (**Fig. 3a**). By contrast, TFBSs in lineage-specific dREG sites had lower enrichments in phyloP scores (**Fig. 3a, red/blue**). Notably, TFBSs in dREG sites lost on the human lineage show enhanced conservation compared with those in human-specific gains, consistent with losses evolving under conservation in other mammalian species (which contribute to the phyloP scores) and gains emerging relatively recently. Restricting these analyses to binding sites for specific TFs revealed patterns of conservation that correlate with the information content of the DNA sequence motif (**Fig 3b; Supplementary Fig. 9**), further supporting TF binding as the functional property driving evolutionary conservation at these sites. This analysis suggests that the sequences in TFBSs are a primary driver of transcriptional differences between species.

To estimate the specific contribution of each TF to transcriptional changes, we compared lineage-specific single-nucleotide substitutions within TFBSs in lineage-specific TREs to the same statistic for TFBSs in conserved TREs. This analysis identified 363 TFs with statistically significant associations between lineage-specific transcription and nucleotide substitutions in their TFBSs in resting CD4+ T-cells (**Fig. 3c**). The most strongly enriched of these TFs include several with known or predicted roles in CD4+ T-cell biology, such as ELF1, YY1, and CREB1. ELF1 bindings sites, for example, are enriched in non-coding GWAS SNPs putatively affecting human autoimmune phenotypes in CD4+ T-cell TREs[63]. We applied the same analysis to TREs displaying lineage-specific changes in π-dependent activation, and identified enrichments for the motifs binding NFAT, AP-1 heterodimers (FOS and JUN), and NF-kB (p = 0.03), consistent with our motif discovery results within each individual primate species. In one example at the *SGPP2* locus (**Fig. 3d**), two NF-kB binding motifs found in the proximal promoter and an internal enhancer of *SGPP2* (**Fig. 3e; Supplementary Fig. 10**) are candidates for changes that cause humans to activate *SGPP2* transcription following π treatment.

In some cases, we observed numerous nucleotide substitutions within short TFBSs, either individually or in clusters of nearby TFBSs, which is unlikely to occur by chance and suggests a possible role for positive selection in the evolution of these binding sites. For example, upstream of *SGPP2* we noted an excess of derived alleles in modern humans in cases where the sequenced Neanderthal has the ancestral allele (**Fig. 3d**)[64], potentially consistent with positive selection driving evolutionary changes in *SGPP2* transcription since the divergence of modern humans and Neanderthals. To more directly gauge the impact of positive selection, we used INSIGHT[65] to compare patterns of within-species polymorphism and between-species sequence divergence in TREs that had undergone human lineage-specific transcriptional changes. As has been reported previously for regulatory sequences[9], dREG sites appear to be most strongly influenced by weak negative selection, which is reflected in an excess of low frequency derived alleles in human populations (**Fig. 3f**). Nevertheless, TREs with lineage-specific transcriptional changes in human

CD4+ T-cells showed reduced weak negative selection and were strikingly enriched for adaptive
nucleotide substitutions ($p < 0.01$ INSIGHT likelihood ratio test; **Fig. 3f**), consistent with positive
selection at these sites. We estimate that at least 160 adaptive substitutions have occurred since
the human/chimpanzee divergence within TFBSs that undergo transcriptional changes in human
CD4+ T-cells. Although we are underpowered to detect the specific contribution of many TFs in this
analysis, we did note statistically significant excesses of predicted adaptive substitutions in the
bindings sites of several TFs, including FOXO1, GATA3, IRF4, RARG, and ZEB1 (**Supplementary Fig.
11**; $p < 0.01$, INSIGHT likelihood ratio test). These estimates highlight the substantial contribution
of adaptive evolutionary changes in TFBSs that influence the transcriptional activity of TREs.
223

Correlation between protein-coding and non-coding transcription

We noticed that evolutionary changes in protein-coding gene transcription frequently
correlate with changes in non-coding transcription units (TU) located nearby. *SGPP2*, for instance,
undergoes a ~60-fold increase in basal transcription in humans, which broadly correlates with
changes in nearby non-coding TUs (**Fig. 3d; Supplementary Fig. 10**). To examine this pattern
more generally, we designed a hidden Markov model (HMM) to estimate the boundaries of TUs
genome-wide, based on patterns of aligned PRO-seq reads. Using this method, we annotated
32,602 TUs active in CD4+ T-cells of at least two of the three primates, likely indicating a TU in the
human-chimpanzee ancestor, as well as an additional 13,085 TUs that are specific to one of the
primate species (**Supplementary Fig. 12a**). Approximately half of the TUs identified using this
approach overlap either annotated protein-coding genes or their associated upstream antisense
RNAs, a small fraction overlap annotated lincRNAs, and approximately half are completely un-
annotated (**Supplementary Fig. 12b**).
A cross-species comparison of the transcription levels for various TU classes (**Fig. 4a**)
revealed that enhancer RNAs evolve in expression most rapidly and protein-coding genes evolve
most slowly. Upstream antisense RNAs are relatively conserved compared to more distal TUs,
perhaps because they share regulatory sequences with protein-coding genes. Intriguingly,
lincRNAs—including both those annotated in GENCODE and novel lincRNAs predicted by our
HMM—undergo evolutionary changes in expression about as frequently as enhancer RNAs. Since
patterns of nascent transcription at active lincRNA promoters are highly similar to those at active
enhancers[50,66], this finding suggests that, on average, lincRNA transcription is no more conserved
than can be explained by the activity of the lincRNA promoter.
We measured the extent to which non-coding and protein-coding transcriptional activities
are correlated through evolutionary time. Evolutionary changes in protein-coding gene expression
were highly correlated with those at upstream (Pearson's $R = 0.81$, $p < 2.2e-16$) and internal ($R = $
0.66, $p < 2.2e-16$) antisense transcripts of the same genes. Moreover, changes in the transcriptional
activity of gene promoters correlate with the activities of distal enhancers to which they loop
according to cell-type matched ChIA-PET data ($R = 0.40-0.69$, p < 2.4e-6; depending on analysis
assumptions)[67], which are located nearby ($R = 0.58$, p < 2.2e-16), or which share the same
topological associated domain ($R = 0.53$, $p < 2.2e-16$)[68]. Using a generalized linear model to
integrate expression changes in multiple types of TUs, we can explain 65% of the variance in gene
transcription levels ($R^2 = 0.65$ in a heldout set of sites, $p < 2.2e-16$; **Fig. 4b**) based on the activities of
looped TREs, nearby TREs, internal antisense TUs, and the upstream antisense TU. Thus

evolutionary changes that result in differences in Pol II recruitment to protein-coding genes are
remarkably correlated across all interacting TREs, indicating a shared evolutionary pressure at
proximal and distal TREs.
260

Rates of Enhancer Evolution Vary with Evidence for Gene Interactions

Transcription at enhancers evolves rapidly and is frequently unaccompanied by changes at
nearby protein-coding genes. For example, *CCR7* transcription is highly conserved among both
primate and rodent species (**Fig. 5a; Supplementary Fig. 2**) in spite of several apparent changes in
enhancer activity within the same locus (gray vertical bars). These findings are consistent with
recent observations that changes in enhancers within densely populated loci often do not have
appreciable effects on the transcription of genes within the locus[35,36].
To explain this effect, we searched for genomic features correlated with conservation of
transcription at enhancers. We found that one of the most strongly correlated features is the
distance between enhancers and the nearest annotated RefSeq transcription start site (**Fig. 5b**). In
particular, about half of enhancers located within 10 kbp of an annotated TSS are shared across all
three primate species, whereas for distal enhancers located between 100 kbp to 1 Mbp from a TSS
that fraction drops to roughly a third. DNA sequences of proximal sites were also more highly
conserved (**Supplementary Fig. 13**), correlating with the higher transcriptional conservation,
though the effect was limited to the area highly proximal to the transcription start site.
These simple distance-based observations, however, ignore the critical issue of chromatin
interactions between enhancers and promoters. To account for such loop interactions, we extracted
6,520 putative TRE interactions from chromatin interaction analysis with paired end tag
sequencing (ChIA-pet) data recognizing loops marked with H3K4me2 in human CD4+ T-cells[67].
Strikingly, we found that, overall, 51% of enhancers that participate in these loops were conserved
between primate species compared to 37% of non-looped enhancers (**Fig. 5c**;, p = 2.1e-13, Fisher's
exact test). Moreover, this high degree of transcriptional conservation at looped enhancers does
not depend on the distance to the transcription start site. We observed similar levels of
conservation at recently defined superenhancers[69], which may simply reflect an enrichment for
loop interactions (48% of TREs in superenhancers loop according to ChIA-PET, compared to 15% of
all TREs). Looped enhancers were also enriched for elevated phyloP scores relative to either non-
looped enhancers or randomly selected DNA sequences (**Fig. 5d**; phyloP > 0.75; p < 2.2e-16,
Wilcoxon Rank Sum Test). That the subset of enhancers which form loop interactions with distal
sites is more highly conserved at both the transcription and DNA sequence levels suggests this
subset has a disproportionately large effect on fitness, perhaps because it is more directly involved
in transcriptional regulation.
292

Enhancer-Promoter Interactions Contribute to Constraint on Gene Transcription Levels

Distal loop interactions do not fully account for the disparity in evolutionary rates between
enhancer and promoter transcription. Looped enhancers still evolve significantly faster than
promoters (**Supplementary Fig. 14**; p = 3e-5, Fisher's exact test). We hypothesized that a higher
redundancy in cis-regulatory signals makes protein-coding genes controlled by multiple TREs, such
as *CCR7* (**Fig. 5a**), more robust to enhancer turnover. Indeed, we found that evolutionary

conservation of promoter TRE transcription is remarkably strongly correlated with the number of
loop interactions with distal sites (Fig. 5e, weighted Pearson's correlation = 0.91; p < 1e-3 by a
bootstrap test). Promoters involved in one or more chromatin loops change expression 15% more
slowly than non-looped promoters (p = 4.7e-4; Fisher's exact test), and the probably of
transcription conservation increases by ~4% with each additional loop interaction. We observed a
similar correlation between the number of loop interactions made by a target promoter and DNA
sequence conservation in transcription factor binding motifs at the promoter (Fig. 5e;
Supplementary Fig. 15a).

We hypothesized that compensation among TREs may permit changes to distal TREs that
loop to promoters which themselves have many other loop interactions. We therefore examined
the conservation of looped TREs as a function of the number of loops in which their gene-proximal
partners participate. We found that DNA sequence conservation in putative TFBSs is significantly
reduced at the distal end of loop interactions and negatively correlates with the number of loops at
the proximal end (Fig. 5f; Supplementary Fig. 15b). This result suggests that each associated
distal TFBS is individually less essential at genes having multiple loop interactions with distal sites,
and it is therefore consistent with a model in which such TFBSs are more freely gained and lost
during evolution. This result may also explain why superenhancers have lower DNA sequence
conservation than typical looped enhancers, in spite of their strong enrichment for loop
interactions (Fig. 5d). Taken together, our results imply that distance, looping, and redundancy of
enhancers all contribute to constraints on the evolutionary rates of changes in gene transcription.
320
Discussion:
We describe the first comparative analysis of primary transcription in any phylogenetic
group, focusing on CD4+ T-cells in primates. Using PRO-seq and several new bioinformatic tools we
estimated the locations and abundance of transcription units with high resolution and accuracy. In
comparison to previous studies in primates[28,32,70–72], this approach separated primary transcription
from post-transcriptional processing, allowing us to study eRNAs, lincRNAs, and other rapidly
degraded non-coding RNAs, as well as protein-coding genes. We found clear relationships between
the DNA sequences of TFBSs and differential transcription across species and treatment conditions.
We also found evidence that some transcriptional changes in humans were driven by adaptive
evolution in nearby binding sites. Overall, our study provides new insights into the mode and
tempo of recent evolutionary changes in transcription in primates.

Perhaps our most striking observation is that many non-coding transcription units,
particularly eRNAs and lincRNAs, have undergone rapid evolutionary changes in comparison to
protein-coding genes. Similar observations have been reported previously for lincRNAs[73], but, to
our knowledge, the observation for eRNAs is new, and it raises a number of questions. First, why
are some enhancers more conserved than others? We find that enhancers proximal to, or that loop
to, annotated promoters tend to be constrained (Fig 5b-c). These enhancers may simply be most
crucial for activating their target genes, but other factors may also contribute to their constraint.
For example, perhaps these enhancers are enriched for tissue-specific functions, and are less
constrained due to reduced pleiotropy[74]. Or perhaps many of them are simply not functional at all,
and are transcribed as a by-product of other processes.

Second, how do protein-coding genes maintain stable transcription levels across species
despite the rapid turnover of associated enhancers? One possibility is that many rapidly evolving
enhancers are either not functional or act on targets other than the ones we have identified.
However, several of our findings argue against this possibility; for example, we find that even
looped enhancers evolve significantly faster than promoters (**Supplementary Fig. 14**), that eRNA
conservation is strongly correlated with the number of loop interactions at associated promoters
(**Fig. 5e**), and that sequence conservation at distal enhancers is negatively correlated with the
number of loop interactions at associated promoters (**Fig. 5f**). An alternative explanation, which
appears more plausible to us, is that stabilizing selection on transcription levels drives enhancers to
compensate for one another as they undergo evolutionary flux. This observation would be
compatible with reports from model systems[35,37]. The possibility of pervasive stabilizing selection
on transcription levels in primates has been noted previously based on RNA-seq data[75], but our
data allow for more direct observations of both active transcription and associated regulatory
elements.
Third, if most transcribed enhancers do indeed influence gene expression, then why are so many of
them weakly maintained by natural selection? One possibility is that some of these enhancers have
simply switched cell types, as has been reported in some cases[19]. Another possibility is that
selection acts diffusely on enhancers across an entire locus, rather than strongly on individual
enhancers, as has been proposed in cancer evolution[76]. A related idea is that, over long
evolutionary time periods, it may be useful to maintain a collection of enhancers at each locus, even
if all of them are not being used at any given time—they may, in a sense, serve as "spare parts" that
can be mixed and matched, as opportunities allow, for cell-type and developmental-stage-specific
functions. Similar suggestions have been made for noncoding transcription units such as eRNAs,
upstream antisense RNAs, and lincRNAs[50,77], but it may hold as well for regulatory function. It will
be possible to begin to evaluate some of these hypotheses as better data describing enhancers and
enhancer-promoter interactions across many cell types become available for these and other
groups of species.

Methods:

Multiple species PRO-seq library generation. *Isolation of primate CD4+ T-cells.* All human and animal experiments were done in compliance with Cornell University IRB and IACUC guidelines. We obtained peripheral blood samples (60-80 mL) from three healthy male humans, chimpanzees, and rhesus macaques. Blood was collected into purple top EDTA tubes. Human samples were maintained overnight at 4C to mimic shipping non-human primate blood samples. Blood was mixed 50:50 with phosphate buffered saline (PBS). Peripheral blood mononuclear cells (PBMCs) were isolated by centrifugation (750x g) of 35 mL of blood:PBS over 15 mL Ficoll-Paque for 30 minutes at 20C. Cells were washed three times in ice cold PBS. CD4+ T-cells were isolated using CD4 microbeads (Miltenyi Biotech, 130-045-101 [human and chimp], 130-091-102 [rhesus macaque]). Up to 10^8 PBMCs were resuspended in binding buffer (PBS with 0.5% BSA and 2mM EDTA). Cells were bound to CD4 microbeads (20uL of microbeads/ 10^7 cells) for 15 minutes at 4C in the dark. Cells were washed with 1-2 mL of PBS/BSA solution, resuspended in 500uL of binding buffer, and passed over a MACS LS column (Miltenyi Biotech, 130-042-401) on a neodymium magnet. The MACS LS column was washed three times with 2mL PBS/BSA solution, before being eluted off the neodymium magnet. Cells were counted in a hemocytometer.

Isolation of CD4+ T-cells from mouse and rat. Spleen samples were collected from one male mouse (FVB) and one male rat (Albino Oxford) that had been sacrificed for IACUC-approved research not related to the present study. Dissected spleen was mashed through a cell strainer using a sterile glass pestle and suspended in 20 mL RPMI-1640. Cells were pelleted at 800xg for 3 minutes and resuspended in 1-5mL of ACK lysis buffer for 10 minutes at room temperature to lyse red blood cells. RPMI-1640 was added to a final volume 10 times that used for ACK lysis (10-40 mL). Cells were pelleted at 800xg for 3 minutes, counted in a hemocytometer, and resuspended in RPMI-1640 to a final concentration of 250,000 cells per ml. CD4+ T-cells were isolated from splenocytes using products specific for mouse and rat (Miltenyi Biotech, 130-104-453 [mouse], 130-090-319 [rat]) following instructions from Miltenyi Biotech, and as described above.

T-cell treatment and PRO-seq library generation. CD4+ T-cells were allowed to equilibrate in RPMI-1640 supplemented with 10% FBS for 2-4 hours before starting experiments. Primate CD4+ T-cells were stimulated with 25ng/mL PMA and 1mM Ionomycin (P/I or π) or vehicle control (2.5uL EtOH and 1.66uL DMSO in 10mL of culture media). We selected the minimum concentrations which saturate the production of IL2 and IFNG mRNA after 3 hours of treatment (data not shown). A 30 min. treatment duration was selected after observing a sharp increase in ChIP-qPCR signal for RNA Pol II phosphorylated at serine 5 on the C-terminal domain on the IFNG promoter at 30 min. (data not shown). To isolate nuclei, we resuspended cells in 1 mL lysis buffer (10 mM Tris-Cl, pH 8, 300 mM sucrose, 10 mM NaCl, 2 mM MgAc2, 3 mM CaCl2 and 0.1% NP-40). Nuclei were washed in 10 mL of wash buffer (10 mM Tris-Cl, pH 8, 300 mM sucrose, 10 mM NaCl and 2 mM MgAc2) to dilute free NTPs. Nuclei were washed in 1 mL, and subsequently resuspended in 50 µL, of storage buffer (50 mL Tris-Cl, pH 8.3, 40% glycerol, 5 mM MgCl2 and 0.1 mM EDTA), snap frozen in liquid nitrogen and kept for up to 6 months before making PRO-seq libraries. PRO-

412   seq libraries were created exactly as described previously[45]. In most cases, we completed library
413   preps with one member of each species (usually one human, chimpanzee, and rhesus macaque) to
414   prevent batch effects from confounding differences between species. Samples were sequenced on
415   an Illumina Hi-Seq 2000 or NextSeq500 at the Cornell University Biotechnology Resource Center.
416
Mapping PRO-seq reads. We mapped PRO-seq reads using standard informatics tools. Our PRO-seq
mapping pipeline begins by removing reads that fail Illumina quality filters and trimming adapters
using cutadapt with a 10% error rate[78]. Reads were mapped with BWA [79] to the appropriate
reference genome (either hg19, panTro4, rheMac3, mm10, or rn6) and a single copy of the Pol I
ribosomal RNA transcription unit (GenBank ID# U13369.1). Mapped reads were converted to
bigWig format for analysis using BedTools[80] and the bedGraphToBigWig program in the Kent
Source software package[81]. The location of the RNA polymerase active site was represented by the
single base, the 3' end of the nascent RNA, which is the position on the 5' end of each sequenced
read. During comparative analyses, the genomic coordinates of mapped reads were converted to
the human assembly (hg19) using CrossMap[82]. We converted genomic coordinates between
genome assemblies using reciprocal-best (rbest) nets[83]. We downloaded rbest nets for hg19-
mm10, hg19-panTro4, hg19-rn6 from the UCSC genome browser. We created rbest nets for hg19-
rheMac3 using the doRecipBets.pl script provided as part of the Kent Source software package.
430
Analysis of transcriptional regulatory elements. *Defining a consensus set of transcriptional*
regulatory elements. We predicted TREs in each species' reference genome using dREG[52]. In all
cases, we combined the reads from all individuals for each species and T-cell treatment condition in
order to maximize power for the discovery of TREs.   We then defined a consensus set of TRE
annotations, each of which bore the signature of an active TRE in at least one species and treatment
condition.   To define such a set, dREG scores were first converted to human reference genome
(hg19) coordinates using CrossMap and the reciprocal-best nets. We then identified TREs in each
species separately by thresholding the dREG scores. Instead of simply merging overlapping TREs,
which tends to create large elements, we applied the thresholding procedure several times, taking
scores greater than 0.9, 0.8, and 0.7, and each time taking only TREs that did not intersect a
previously selected TRE at a higher threshold. Finally, the set of overlapping TREs from each
species were reduced to a single element containing the union of all positions covered by the set
using bedops, and sites within 500 bp of each other were further merged. We assigned each
putative TRE the maximum dREG score for each species and for each treatment condition.
445
Identifying differences in TREs between species. Differences in TRE transcription in 3-way (human-
chimp-rhesus macaque) or 5-way (human-chimp-rhesus macaque-mouse-rat) species comparisons
were identified using a combination of heuristics and statistical tests. Starting with the consensus
set of TREs in hg19 coordinates, we first excluded potential one-to-many orthologs, by eliminating
TREs that overlapped gaps in the reciprocal-best nets that were not classified as gaps in the
standard nets. The remaining TREs were classified as unmappable when no orthologous position
was defined in the rbest nets. Complete gains and losses were defined as TREs that were mappable
in all species and for which the dREG score was less than 0.1 in at least one species and greater than
0.7 in at least one other species. Gains and losses were assigned to a lineage based on an

assumption of maximum parsimony under the known species phylogeny. We defined a set of TREs
that displayed high-confidence changes in activity by comparing differences in PRO-seq read counts
between species using the Limma package[84] and thresholding at a 5% false discovery rate (as
described below). These TREs were often active in all species. Changes in TRE activities were
compared to histone modification ChIP-seq and DNase-I-seq data from the Epigenome Roadmap
project [55].

TRE classification. For some analyses, TREs were classified into likely promoters and enhancers on
the basis of their distance from known gene annotations (RefGene) and the predicted stability of
the resulting transcription unit (TU). TRE classes of primary interest include (see also
Supplementary Fig. 7): (1)promoters: near an annotated transcription start site (<500 bp) and
producing a stable TU (instability score <0.1); (2)enhancers: distal to an annotated transcription
start site (>10,000 bp) and producing an unstable TU (instability score >0.1). TU stability was
defined using the posterior probability that a TRE yields an unstable transcription unit using the
forward-backward tables in a hidden Markov model we described recently[50]. This approach
produced a score for both the forward and the upstream divergent TU of each TRE. We took the
minimum of these instability scores to represent the stability of that TRE.

Covariates that correlate with TRE changes. We compared the frequency at which evolutionary
changes in transcription occur at TREs in a variety of different genomic context. We compared
changes as a function of distance from the nearest annotated transcription start site in RefSeq.
TREs were binned by distance in increments of 0.02 on a log10 scale and we evaluated the mean
rate at which evolutionary changes in TRE transcription arise in each bin. We also compared the
rate of changes in TRE transcription in a variety of functional associations, including loop
interactions, within the same topological associated domain, and in superenhancers. H3K4me2
ChIA-PET data describing loop interactions were downloaded from the Gene Expression Omnibus
(GEO) database (GSE32677) and the genomic locations of loops were converted from hg18 to hg19
coordinates using the liftOver tool. Looped enhancers were defined as those within 5,000 bp of a
loop center. Topological associated domains (TADs) based on Hi-C data for GM12878 cells were
also downloaded from GEO (GSE63525). Superenhancers in CD4+ T-cells were taken from the
supplementary data for ref. [69].

Refining dREG peak calls using dREG-HD. During analyses on transcription factor binding motifs we
further refined the location of TREs to the region between divergent paused RNA polymerase using
a strategy that we call dREG-HD (manuscript in preparation, preliminary version available at
https://github.com/Danko-Lab/dREG.HD). Briefly, we used an epsilon-support vector regression
(SVR) with a Gaussian kernel to map the distribution of PRO-seq reads to smoothed DNase-I signal
intensities. Training was conducted on randomly chosen positions within dREG peaks extended by
200bp on either side. Selection of feature vectors was optimized based on Pearson correlation
coefficients between the imputed and experimental DNase-I score over the validation set. PRO-seq
data was normalized by sequencing depth and further scaled such that the maximum value of any
prediction dataset is within 90 percentile of the training examples. We chose a step size to be 60bp

and extending 30 steps on each direction. The final model was trained using matched DNase-I and
PRO-seq data in K562 cells.
Next we identified peaks in the imputed DNase-I hypersensitivity profile by fitting the
imputed DNase-I signal using a cubic spline and identifying local maxima. We optimized two free
parameters that control the (1) smoothness of spline curve fitting, and (2) threshold on the
imputed DNase-I signal intensity. Parameters were optimized to achieve an appropriate trade-off
between FDR and sensitivity on the testing K562 dataset. Parameters were tuned using a grid
optimization over free parameters. Testing the optimized dREG-HD (including both DNase-I
imputation and peak calling) on GM12878, a GRO-seq dataset completely held out from model
training and parameter optimization, revealed 82% sensitivity for DNase-I peaks within dREG sites
at a 10% false discovery rate (FDR).
508
DNA sequence analysis. *DNA sequence conservation analysis.* For our evolutionary conservation
analysis, we used phyloP scores[60] based on either (1) the primate species in the 46-way alignments,
or (2) the 100-way genome alignments, both available in the UCSC Genome Browser (hg19). In all
cases, bigWig files were obtained from the UCSC Genome Browser and processed using the bigWig
package in R. We represented evolutionary conservation as the mean phyloP score in each
identified TFBS in the indicated set of dREG-HD sites. We focused on 1,964 human TF binding
motifs from the CisBP database[61] and clustered motifs using an affinity propagation algorithm into
567 maximally distinct DNA binding specificities (see ref[62]). We used TFBSs having a $\log_e$-odds
score >10 in any of the primate reference genomes, with scores obtained by comparing each
candidate motif model to a third-order Markov background model using the RTFBSDB package[62].
519
Motif enrichment in TREs that change during CD4+ T-cell activation. Motif enrichment analyses were
completed using RTFBSDB[62] as described above, except that motifs were clustered into 621
maximally distinct DNA binding specificities (see ref[62]). We selected the motif whose canonical
transcription factor is most highly transcribed in human CD4+ T-cells to represent each cluster. We
defined a motif cutoff $\log_e$ odds ratio of 7.5 in a sequence compared with a third-order Markov
model as background. Motifs enriched in up- or down-regulated dREG-HD TREs during CD4+ T-cell
activation (>8-fold in magnitude and $p < 0.01$) were selected using Fisher's exact test with a
Bonferroni correction for multiple hypothesis testing. Up- or down-regulated TREs were compared
to a background set of >2,500 GC-content matched TREs that do not change transcription levels
following π treatment (<2-fold in magnitude and $p > 0.1$) using the *enrichmentTest* function in
RTFBSDB[62].
531
Enrichment of DNA sequence changes in motifs. We identified single nucleotide DNA sequence
differences at sites at which two of three primate species share one base and the third species
diverges. We intersected these species-specific divergences with matches to transcription factor
binding motifs found within dREG-HD sites that undergo transcriptional changes between primate
species. Because many motifs in Cis-BP are similar to one another, we first partitioned the motifs
into 567 clusters having highly similar binding preferences, and examined enrichments at the level
of these clusters. Motifs were ranked by the Fisher's exact test p-value of the enrichment of species
divergences in dREG-HD sites that change transcription status (where changes in DNA sequence

and transcription occur on the same branch) to dREG-HD sites that do not change. We also
compute the enrichment ratio, which we define as the number of species divergences in each TF
binding motif in dREG-HD sites that change on the same branch normalized to the same statistic in
sites that do not change.
544
INSIGHT analysis. We examined the modes by which DNA sequences evolve in human lineage
specific dREG-HD sites or DHSs using INSIGHT[65]. We passed INSIGHT either complete DHSs, dREG-
HD sites, or TFBS (log-odds score >7) within dREG-HD sites that undergo the changes (see
Identifying differences in TREs between species) indicated in the comparison. Human gains and
losses, for example, were comprised of 2,975 dREG-HD sites with 13,843 separate regions (median
length of 29 bp) after merging overlapping TFBSs with a log-odds score greater than 7. We also
analyzed 27 transcription factors each of which has more than 1,000 occurrences in dREG-HD sites
that change on the human branch. All analyses were conducted using the INSIGHT web server
(http://compgen.cshl.edu/INSIGHT/) with the default settings enabled.
554
De novo discovery of transcription units. *Identification of transcription units (TU) using a three-*
state hidden Markov model. We inferred transcription units (TU) using a three-state hidden Markov
model (HMM) to capture the distribution of PRO-seq read densities similar to those we have
recently published[49,85]. Three states were used to represent background (i.e., outside of a
transcription unit), TU body, and a post-polyA decay region. The HMM transition structure is shown
in **Supplementary Fig. 13a**. We allow skipping over the post-polyA state, as unstable transcripts
do not have these two-phase profiles. We took advantage of dREG as a potential signal for
transcription initiation by incorporating the dREG score (maximum value in the interval from a
given positive read-count position until the next, clamped to the zero-one interval) as a transition
probability from the background to the transcription body state. PRO-seq data is generally sparse,
so we applied a transformation that encoded only non-zero positions and the distance between
such consecutive positions (**Supplementary Fig. 13a**). Our model described this transformed data
using emissions distribution based on two types of variables. The first type of emission variable
defines the PRO-seq read counts in non-zero positions. These counts were modeled using Poisson
distributions in the background and post-polyA states, and using a Negative Binomial distribution
in the transcription body state. The negative binomial distribution can be seen as a mixture of
Poisson distributions with gamma-distributed rates and therefore allows for variation in TU
expression levels across the genome. The second type of emission variable describes the
distribution of distances in base pairs between positions having non-zero read counts. This
distribution was modeled using a separate geometric distribution for each of the three states.
Maximum likelihood estimates of all free parameters were obtained via Expectation Maximization,
on a per-chromosome basis. TU predictions were then obtained using the Viterbi algorithm with
parameters fixed at their maximum-likelihood values. Finally these predictions were mapped from
the transformed coordinates back to genomic coordinates. Source code for our implementation is
publicly available on GitHub: https://github.com/andrelmartins/tunits.nhp.
580
Inferring TU boundaries in the common great ape ancestor. We identified the most likely TU
boundaries in the great ape ancestor by maximum parsimony. TUs were identified and compared

in human reference coordinates (hg19) for all species. We used the bedops package to mark the
intersection between each pair of species (i.e., human-chimp, human-rhesus macaque, and chimp-
rhesus macaque). Intersections (>= 1bp) between pairs of species were merged, resulting in a
collection of TUs shared by any two pairs of species, and therefore likely to be a TU in the human-
chimp ancestor. All steps were applied independently on the plus and minus strands. These steps
identified 32,602 putative TUs active in CD4+ T-cells of the primate ancestor. We added 13,085
TUs that did not overlap ancestral TUs but were found in any one of the three primate species.
590
Transcription unit classification. TUs were classified into by annotation type using a pipeline similar
to ones that we have described recently[49,85,86]. Before classifying TUs we applied a heuristic to
refine TUs on the basis of known annotations. TUs that completely overlap multiple gene
annotations were broken at the transcription start site provided that a dREG site overlapped that
transcription start site. Classification was completed using a set of rules to iteratively refine
existing annotations, as shown in **Supplementary Fig. 13A**. Unless otherwise stated, overlap
between a TU and a transcript annotation was defined such that >50% of a TU matched a gene
annotation and covers at least 50% of the same annotation. TUs overlapping GENCODE annotations
(>50% overlap, defined as above) were classified using the biotype in the GENCODE database into
protein coding, lincRNA (lincRNA or processed transcript), or pseudogene. The remaining
transcripts were classified as annotated RNA genes using GENCODE annotations, the rnaGenes
UCSC genome browser track (converted from hg18 to hg19 coordinates), and miRBase v20 [87]. As
many RNA genes are processed from much longer TUs, we required no specific degree of overlap
for RNA genes. Upstream antisense (i.e., divergent) TUs were classified as those within 500bp of
the transcription start site of any GENCODE or higher level TU annotation (including lincRNAs).
Antisense transcripts were defined as those with a high degree of overlap (>50%) with annotated
protein coding genes in the opposite orientation. The remaining transcripts with a high degree of
overlap (>50%) to annotated repeats in the repeatmasker database (rmsk) were classified as
repeat transcription. Finally, any TUs still remaining were classified as unannotated, and were
further divided into those which are intergenic or that partially overlapping existing annotations.
611
Comparing transcription between conditions and species. *Comparing transcription before and*
after CD4+ T-cell activation. We compared π treated and untreated CD4+ T-cells within each of the
primate species using gene annotations (GENCODE v19). We counted reads in the interval between
500 bp downstream of the annotated transcription start site and either the end of the gene or
60,000 bp into the gene body (whichever was shorter). This window was selected to avoid (1)
counting reads in the pause peak near the transcription start site, and (2) to focus on the 5' end of
the gene body affected by changes in transcription during 30 minutes of π treatment assuming a
median elongation rate of 2 kb/ minute[49,88]. We limited analyses to gene annotations longer than
1,000 bp in length. To quantify transcription at enhancers, we counted reads in the window
covered by each dREG-HD site plus an additional 250 bp on each end. Differential expression
analysis was conducted using deSeq2[53].
623
Comparing transcription between species. Read counts were compared between different species in
hg19 coordinates. In all analyses reads were transferred to the hg19 reference genome using

CrossMap with rbest nets. Our analysis focused on transcription units or on the union of dREG sites
across species. We focused our analysis of transcription units on the interval between 250 bp
downstream of the annotated transcription start site and either the end of the gene or 60,000 bp
into the gene body (whichever was shorter). We limited our analyses to TUs longer than 500 bp in
length. Reads counts were obtained within each transcription unit, gene annotation, or enhancer,
abbreviated here as a 'region of interest' (ROI), that has confident one-to-one orthology in all
species examined in the analysis. We broke each each ROI into segments that have conserved
orthology between hg19 and all species examined in the analysis, which included either a three-
way (human-chimp-rhesus macaque) or five-way (human-chimp-rhesus macaque-mouse-rat)
species comparison. We defined intervals of one-to-one orthology as those represented in levels 1,
3, and 5 of the reciprocal best nets (with gaps defined in levels 2, 4, and 6)[83]. Reads that map to
regions that have orthology defined in all species were counted using the bigWig package in R using
reads mapped to hg19 coordinates. Final counts for each ROI were defined as the sum of read
counts within the regions of orthology that intersect that ROI. ROIs without confident one-to-one
orthologs in all species analyzed were discarded. Our pipeline makes extensive use of the bigWig R
package, Kent source tools, as well as the bedops and bedtools software packages[80,89]. Differential
expression was conducted between species using the LIMMA package for R[84].
643
644

References (Paperpile):

1. Jacob, F. & Monod, J. Genetic regulatory mechanisms in the synthesis of proteins. *J. Mol. Biol.* **3,** 318–356 (1961).
2. Britten, R. J. & Davidson, E. H. Gene regulation for higher cells: a theory. *Science* **165,** 349–357 (1969).
3. King, M. C. & Wilson, A. C. Evolution at two levels in humans and chimpanzees. *Science* **188,** 107–116 (1975).
4. Rockman, M. V. *et al.* Ancient and recent positive selection transformed opioid cis-regulation in humans. *PLoS Biol.* **3,** e387 (2005).
5. Prabhakar, S. *et al.* Human-specific gain of function in a developmental enhancer. *Science* **321,** 1346–1350 (2008).
6. Capra, J. A., Erwin, G. D., McKinsey, G., Rubenstein, J. L. R. & Pollard, K. S. Many human accelerated regions are developmental enhancers. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **368,** 20130025 (2013).
7. McLean, C. Y. *et al.* Human-specific loss of regulatory DNA and the evolution of human-specific traits. *Nature* **471,** 216–219 (2011).
8. Cotney, J. *et al.* The evolution of lineage-specific regulatory activities in the human embryonic limb. *Cell* **154,** 185–196 (2013).
9. Arbiza, L. *et al.* Genome-wide inference of natural selection on human transcription factor binding sites. *Nat. Genet.* **45,** 723–729 (2013).
10. Prescott, S. L. *et al.* Enhancer Divergence and cis-Regulatory Evolution in the Human and Chimp Neural Crest. *Cell* **163,** 68–83 (2015).
11. Siepel, A. & Arbiza, L. Cis-regulatory elements and human evolution. *Curr. Opin. Genet. Dev.* **29,** 81–89 (2014).
12. Wray, G. A. The evolutionary significance of cis-regulatory mutations. *Nat. Rev. Genet.* **8,** 206–216 (2007).
13. Wilson, M. D. *et al.* Species-specific transcription in mice carrying human chromosome 21. *Science* **322,** 434–438 (2008).
14. Khurana, E. *et al.* Integrative annotation of variants from 1092 humans: application to cancer genomics. *Science* **342,** 1235587 (2013).
15. Haygood, R., Fedrigo, O., Hanson, B., Yokoyama, K.-D. & Wray, G. A. Promoter regions of many neural- and nutrition-related genes have experienced positive selection during human evolution. *Nat. Genet.* **39,** 1140–1144 (2007).
16. Torgerson, D. G. *et al.* Evolutionary processes acting on candidate cis-regulatory regions in humans inferred from patterns of polymorphism and divergence. *PLoS Genet.* **5,** e1000592 (2009).
17. Schmidt, D. *et al.* Five-vertebrate ChIP-seq reveals the evolutionary dynamics of transcription factor binding. *Science* **328,** 1036–1040 (2010).
18. Ballester, B. *et al.* Multi-species, multi-transcription factor binding highlights conserved control of tissue-specific biological pathways. *Elife* **3,** e02626 (2014).
19. Vierstra, J. *et al.* Mouse regulatory DNA landscapes reveal global principles of cis-regulatory evolution. *Science* **346,** 1007–1012 (2014).
20. Arnold, C. D. *et al.* Quantitative genome-wide enhancer activity maps for five Drosophila species show functional enhancer conservation and turnover during cis-regulatory evolution. *Nat. Genet.* **46,** 685–692 (2014).
21. Doniger, S. W. & Fay, J. C. Frequent gain and loss of functional transcription factor binding sites. *PLoS Comput. Biol.* **3,** e99 (2007).
22. Zheng, W., Zhao, H., Mancera, E., Steinmetz, L. M. & Snyder, M. Genetic analysis of variation in transcription factor binding in yeast. *Nature* **464,** 1187–1191 (2010).

694   23. Bradley, R. K. *et al.* Binding site turnover produces pervasive quantitative changes in
695   transcription factor binding between closely related Drosophila species. *PLoS Biol.* **8,**
696   e1000343 (2010).
697   24. Villar, D. *et al.* Enhancer evolution across 20 mammalian species. *Cell* **160,** 554–566 (2015).
698   25. Chuong, E. B., Elde, N. C. & Feschotte, C. Regulatory evolution of innate immunity through co-
699   option of endogenous retroviruses. *Science* **351,** 1083–1087 (2016).
700   26. Fuda, N. J., Ardehali, M. B. & Lis, J. T. Defining mechanisms that regulate RNA polymerase II
701   transcription in vivo. *Nature* **461,** 186–192 (2009).
702   27. Zhou, X. *et al.* Epigenetic modifications are associated with inter-species gene expression
703   variation in primates. *Genome Biol.* **15,** 547 (2014).
704   28. Cain, C. E., Blekhman, R., Marioni, J. C. & Gilad, Y. Gene expression differences among primates
705   are associated with changes in a histone epigenetic modification. *Genetics* **187,** 1225–1234
706   (2011).
707   29. Xiao, S. *et al.* Comparative epigenomic annotation of regulatory DNA. *Cell* **149,** 1381–1392
708   (2012).
709   30. Paris, M. *et al.* Extensive divergence of transcription factor binding in Drosophila embryos with
710   highly conserved gene expression. *PLoS Genet.* **9,** e1003748 (2013).
711   31. Cusanovich, D. A., Pavlovic, B., Pritchard, J. K. & Gilad, Y. The functional consequences of
712   variation in transcription factor binding. *PLoS Genet.* **10,** e1004226 (2014).
713   32. Wong, E. S. *et al.* Decoupling of evolutionary changes in transcription factor binding and gene
714   expression in mammals. *Genome Res.* **25,** 167–178 (2015).
715   33. Hah, N., Murakami, S., Nagari, A., Danko, C. G. & Kraus, W. L. Enhancer transcripts mark active
716   estrogen receptor binding sites. *Genome Res.* **23,** 1210–1223 (2013).
717   34. Domené, S. *et al.* Enhancer turnover and conserved regulatory function in vertebrate evolution.
718   *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **368,** 20130027 (2013).
719   35. Wunderlich, Z. *et al.* Krüppel Expression Levels Are Maintained through Compensatory
720   Evolution of Shadow Enhancers. *Cell Rep.* **12,** 1740–1747 (2015).
721   36. Cannavò, E. *et al.* Shadow Enhancers Are Pervasive Features of Developmental Regulatory
722   Networks. *Curr. Biol.* **26,** 38–51 (2016).
723   37. Ludwig, M. Z., Bergman, C., Patel, N. H. & Kreitman, M. Evidence for stabilizing selection in a
724   eukaryotic enhancer element. *Nature* **403,** 564–567 (2000).
725   38. Sanyal, A., Lajoie, B. R., Jain, G. & Dekker, J. The long-range interaction landscape of gene
726   promoters. *Nature* **489,** 109–113 (2012).
727   39. Vietri Rudan, M. *et al.* Comparative Hi-C reveals that CTCF underlies evolution of chromosomal
728   domain architecture. *Cell Rep.* **10,** 1297–1309 (2015).
729   40. Khan, Z. *et al.* Primate transcript and protein expression levels evolve under compensatory
730   selection pressures. *Science* **342,** 1100–1104 (2013).
731   41. Bauernfeind, A. L. *et al.* Evolutionary Divergence of Gene and Protein Expression in the Brains
732   of Humans and Chimpanzees. *Genome Biol. Evol.* **7,** 2276–2288 (2015).
733   42. Battle, A. *et al.* Genomic variation. Impact of regulatory variation from RNA to protein. *Science*
734   **347,** 664–667 (2015).
735   43. Pai, A. A. *et al.* The contribution of RNA decay quantitative trait loci to inter-individual
736   variation in steady-state gene expression levels. *PLoS Genet.* **8,** e1003000 (2012).
737   44. Core, L. J., Waterfall, J. J. & Lis, J. T. Nascent RNA sequencing reveals widespread pausing and
738   divergent initiation at human promoters. *Science* **322,** 1845–1848 (2008).
739   45. Kwak, H., Fuda, N. J., Core, L. J. & Lis, J. T. Precise maps of RNA polymerase reveal how
740   promoters direct initiation and pausing. *Science* **339,** 950–953 (2013).
741   46. Churchman, L. S. & Weissman, J. S. Nascent transcript sequencing visualizes transcription at
742   nucleotide resolution. *Nature* **469,** 368–373 (2011).
47. Nojima, T. *et al.* Mammalian NET-Seq Reveals Genome-wide Nascent Transcription Coupled to

RNA Processing. *Cell* **161,** 526–540 (2015).

48. Mahat, D. B. *et al.* Base-pair-resolution genome-wide mapping of active RNA polymerases using
precision nuclear run-on (PRO-seq). *Nat. Protoc.* **11,** 1455–1476 (2016).

49. Hah, N. *et al.* A rapid, extensive, and transient transcriptional response to estrogen signaling in
breast cancer cells. *Cell* **145,** 622–634 (2011).

50. Core, L. J. *et al.* Analysis of nascent RNA identifies a unified architecture of initiation regions at
mammalian promoters and enhancers. *Nat. Genet.* **46,** 1311–1320 (2014).

51. Andersson, R. *et al.* Nuclear stability and transcriptional directionality separate functionally
distinct RNA species. *Nat. Commun.* **5,** 5336 (2014).

52. Danko, C. G. *et al.* Identification of active transcriptional regulatory elements from GRO-seq
data. *Nat. Methods* **12,** 433–438 (2015).

53. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol.* **15,** 550 (2014).

54. Nguyen, D. H., Hurtado-Ziola, N., Gagneux, P. & Varki, A. Loss of Siglec expression on T
lymphocytes during human evolution. *Proc. Natl. Acad. Sci. U. S. A.* **103,** 7765–7770 (2006).

55. Roadmap Epigenomics Consortium *et al.* Integrative analysis of 111 reference human
epigenomes. *Nature* **518,** 317–330 (2015).

56. Macian, F. NFAT proteins: key regulators of T-cell development and function. *Nat. Rev.*
Immunol. **5,** 472–484 (2005).

57. Ouyang, W., Beckett, O., Flavell, R. A. & Li, M. O. An essential role of the Forkhead-box
transcription factor Foxo1 in control of T cell homeostasis and tolerance. *Immunity* **30,** 358–
371 (2009).

58. Bonn, S. *et al.* Tissue-specific analysis of chromatin state identifies temporal signatures of
enhancer activity during embryonic development. *Nat. Genet.* **44,** 148–156 (2012).

59. Zentner, G. E., Tesar, P. J. & Scacheri, P. C. Epigenetic signatures distinguish multiple classes of
enhancers with distinct cellular functions. *Genome Res.* **21,** 1273–1283 (2011).

60. Pollard, K. S., Hubisz, M. J., Rosenbloom, K. R. & Siepel, A. Detection of nonneutral substitution
rates on mammalian phylogenies. *Genome Res.* **20,** 110–121 (2010).

61. Weirauch, M. T. *et al.* Determination and inference of eukaryotic transcription factor sequence
specificity. *Cell* **158,** 1431–1443 (2014).

62. Wang, Z., Martins, A. L. & Danko, C. G. RTFBSDB: an integrated framework for transcription
factor binding site analysis. *Bioinformatics* (2016). doi:10.1093/bioinformatics/btw338

63. Farh, K. K.-H. *et al.* Genetic and epigenetic fine mapping of causal autoimmune disease variants.
Nature **518,** 337–343 (2015).

64. Green, R. E. *et al.* A draft sequence of the Neandertal genome. *Science* **328,** 710–722 (2010).

65. Gronau, I., Arbiza, L., Mohammed, J. & Siepel, A. Inference of natural selection from
interspersed genomic elements based on polymorphism and divergence. *Mol. Biol. Evol.* **30,**
1159–1171 (2013).

66. Sun, M., Gadad, S. S., Kim, D.-S. & Kraus, W. L. Discovery, Annotation, and Functional Analysis of
Long Noncoding RNAs Controlling Cell-Cycle Gene Expression and Proliferation in Breast
Cancer Cells. *Mol. Cell* **59,** 698–711 (2015).

67. Chepelev, I., Wei, G., Wangsa, D., Tang, Q. & Zhao, K. Characterization of genome-wide
enhancer-promoter interactions reveals co-expression of interacting genes and modes of
higher order chromatin organization. *Cell Res.* **22,** 490–503 (2012).

68. Rao, S. S. P. *et al.* A 3D map of the human genome at kilobase resolution reveals principles of
chromatin looping. *Cell* **159,** 1665–1680 (2014).

69. Hnisz, D. *et al.* Super-enhancers in the control of cell identity and disease. *Cell* **155,** 934–947
(2013).

70. Barreiro, L. B., Marioni, J. C., Blekhman, R., Stephens, M. & Gilad, Y. Functional comparison of
innate immune signaling pathways in primates. *PLoS Genet.* **6,** e1001249 (2010).

71. Gilad, Y., Oshlack, A., Smyth, G. K., Speed, T. P. & White, K. P. Expression profiling in primates
reveals a rapid evolution of human transcription factors. *Nature* **440,** 242–245 (2006).
72. Blekhman, R., Oshlack, A., Chabot, A. E., Smyth, G. K. & Gilad, Y. Gene regulation in primates
evolves under tissue-specific selection pressures. *PLoS Genet.* **4,** e1000271 (2008).
73. Kutter, C. *et al.* Rapid turnover of long noncoding RNAs and the evolution of gene expression.
PLoS Genet. **8,** e1002841 (2012).
74. Lewis, J. J., van der Burg, K. R. L., Mazo-Vargas, A. & Reed, R. D. ChIP-Seq-Annotated Heliconius
erato Genome Highlights Patterns of cis-Regulatory Evolution in Lepidoptera. *Cell Rep.* **16,**
2855–2863 (2016).
75. Gilad, Y., Oshlack, A. & Rifkin, S. A. Natural selection on gene expression. *Trends Genet.* **22,** 456–
461 (2006).
76. Bailey, S. D. *et al.* Noncoding somatic and inherited single-nucleotide variants converge to
promote ESR1 expression in breast cancer. *Nat. Genet.* **48,** 1260–1266 (2016).
77. Wu, X. & Sharp, P. A. Divergent transcription: a driving force for new gene origination? *Cell*
155, 990–996 (2013).
78. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal **17,** 10–12 (2011).
79. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform.
Bioinformatics **25,** 1754–1760 (2009).
80. Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic features.
Bioinformatics **26,** 841–842 (2010).
81. Kuhn, R. M., Haussler, D. & Kent, W. J. The UCSC genome browser and associated tools. *Brief.*
Bioinform. **14,** 144–161 (2013).
82. Zhao, H. *et al.* CrossMap: a versatile tool for coordinate conversion between genome
assemblies. *Bioinformatics* **30,** 1006–1007 (2014).
83. Kent, W. J., Baertsch, R., Hinrichs, A., Miller, W. & Haussler, D. Evolution's cauldron: duplication,
deletion, and rearrangement in the mouse and human genomes. *Proc. Natl. Acad. Sci. U. S. A.*
100, 11484–11489 (2003).
84. Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing and
microarray studies. *Nucleic Acids Res.* **43,** e47 (2015).
85. Chae, M., Danko, C. G. & Kraus, W. L. groHMM: a computational tool for identifying unannotated
and cell type-specific transcription units from global run-on sequencing data. *BMC*
Bioinformatics **16,** 222 (2015).
86. Luo, X., Chae, M., Krishnakumar, R., Danko, C. G. & Kraus, W. L. Dynamic reorganization of the
AC16 cardiomyocyte transcriptome in response to TNFα signaling revealed by integrated
genomic analyses. *BMC Genomics* **15,** 155 (2014).
87. Kozomara, A. & Griffiths-Jones, S. miRBase: annotating high confidence microRNAs using deep
sequencing data. *Nucleic Acids Res.* **42,** D68–73 (2014).
88. Danko, C. G. *et al.* Signaling pathways differentially affect RNA polymerase II initiation, pausing,
and elongation rate in cells. *Mol. Cell* **50,** 212–222 (2013).
89. Neph, S. *et al.* BEDOPS: high-performance genomic feature operations. *Bioinformatics* **28,**
1919–1920 (2012).

836
837

838
839
Acknowledgements: We thank M. Jin for assistance in establishing the magnetic separation of
CD4+ T-cells, L. Core, H. Kwak, N. Fuda, and I. Jonkers for assistance troubleshooting the PRO-seq
library prep, and A. Wetterau for preparing nuclei for mouse and rat CD4+ T-cells. Work in this
publication was supported by generous seed grants from the Cornell University Center for
Vertebrate Genomics (CVG), the Center for Comparative and Population Genetics (3CPG), NHLBI
(National Heart, Lung, and Blood Institute) grant UHL129958A to CGD and JTL, NIGMS (National
Institute of General Medical Sciences) grant GM102192 to AS, and NHGRI (National Human Genome
Research Institute) grant HG0070707 to AS and JTL. The content is solely the responsibility of the
authors and does not necessarily represent the official views of the US National Institutes of Health.
849
850 **Author contributions:** CGD, EJR, and ETW performed CD4+ T-cell extraction, validation, and PRO-
851 seq experiments. CGD, ZW, TC, and ALM analyzed the data. CGD, AS, JTL, and WLK supervised data
852 collection and analysis. CGD and AS wrote the paper with input from the other authors.
853
854 **Competing financial interests:** The authors declare no competing financial interests.
855
856 **Author information:** PRO-seq data was deposited into the Gene Expression Omnibus database
857 under accession number GSE85337. All data analysis scripts and software are publicly availiable on
858 GitHub: https://github.com/Danko-Lab/CD4-Cell-Evolution.

Fig. 1 | Maps of primary transcription in CD4+ T-cells. **(a)** CD4+ T-cells were isolated from the blood or spleen of individuals from five vertebrate species, including human, chimpanzee, rhesus macaque, mouse, and rat. **(b)** Hierarchical clustering of PRO-seq signal intensities in gene bodies groups 19 CD4+ T-cell samples first by treatment condition and second by species. The color scale represents Spearman's rank correlation between normalized transcription levels in active gene bodies. Colored boxes (top) represents the species and treatment condition of each sample. **(c)** MA plot shows the log2 fold-change following π treatment (y-axis) as a function of the mean transcription level in GENCODE annotated genes (x-axis). Red points indicate statistically significant changes ($p < 0.01$). Several classical response genes that undergo well-documented changes in transcript abundance following CD4+ T-cell activation (e.g., IL2, IFNG, TNF, and EGR3) are marked. **(d)** Heatmaps show the distribution of PRO-seq (red and blue indicate transcription on the plus and minus strand, respectively), H3K27ac, H3K4me1, and H3K4me3 ChIP-seq signal intensity. Plots are centered on transcriptional regulatory elements (TREs) predicted in untreated human CD4+ T-cells using dREG-HD (see Methods), including both promoters and enhancers. All plots are ordered based on H3K27ac ChIP-seq intensity. **(e)** Enrichment of TF binding motifs in TREs that increase transcription levels following π treatment compared to TREs transcribed in both conditions. Plot shows the -log 10 p-value, based on a Fisher's exact test and Bonferroni correction (y-axis), as a function of the rank order of enriched motifs (x-axis). The color scale denotes the degree of enrichment. Motifs, the Cis-BP ID number, and the target transcription factor for several outliers are shown in the plot.

Fig. 2 | Frequency of changes in TRE transcription. **(a)** The fractions of TREs that are present in the human reference genome and are conserved across all species (blue), are not detectable and are therefore inferred as gains or losses (teal) or undergo significant changes (green) in at least one species, or fall in regions for which no ortholog occurs in at least one of the indicated genomes (pink). Plots labeled "Primate" illustrate frequency of changes in a three-way comparison of human, chimpanzee, and rhesus macaque, whereas those labeled "Mammal" summarize a five-way comparison also including rat and mouse. p treatment denotes a similar comparison between human untreated and PMA+Ionomycin treated CD4+ T-cell samples. A yellow box shows the complete fraction of TREs that undergo statistically significant changes in Pol II abundance (FDR-corrected p-value < 0.05). **(b)** ChIP-seq signal for H3K27ac (green) and H3K4me1 (orange) near dREG sites classified as gains, losses, or complete losses of TRE signal (dREG score < 0.1) on the human branch.

Fig. 3 | Evolutionary changes in TRE transcription correlate with DNA sequence conservation. (a) Mean phyloP scores near transcription factor binding motifs in dREG-HD sites that are conserved (green), gained (red), or lost (blue) on the human branch. PhyloP scores were computed based on the 100-way species alignments in the UCSC genome browser. **(b)** PhyloP scores that fall within the binding motif recognized by STAT2 (M6494_1.02). **(c)** Motifs were ranked based on the enrichment of DNA sequence differences found inside of dREG-HD sites that change in primate species compared to those which do not change. Plot shows the Bonferroni-corrected -log 10 p-value (y-axis) as a function of motif rank order (x-axis). **(d)** UCSC Genome Browser track shows transcription near SGPP2 and FARSB in untreated (U) and PMA+ionomycin (p) treated CD4+ T-cells isolated from the indicated primate species. PRO-seq tracks show transcription on the plus (red) and minus (blue) strands. Axes for the PRO-seq data are in units of reads per kilobase per million mapped (RPKM). dREG tracks show the distribution of dREG signal. The Green et. al. (ref63) selective sweep scan track (top) represents the enrichment of derived alleles in modern human where Neanderthal has the ancestral allele. Points below the line represent a statistically significant number of derived alleles in modern human (line indicates a Z-score of -2). Net synteny tracks show the position of regions that have one-to-one orthologs in the chimpanzee and rhesus macaque genomes. **(e)** Multiple species alignment shows DNA sequence near one dREG site that contains a strong match to the NF-κB binding motif in human. This motif occurrence is bound by NF-κB subunit RELA based on ChIP-seq data in multiple ENCODE cell lines (green box). **(f)** INSIGHT estimates of the expected number of segregating polymorphisms under weak negative selection (E[Pw]/kbp) or the expected number of human nucleotide substitutions driven by positive selection (E[Dp]/kbp) in human populations in the indicated class of dREG-HD sites. All estimates are in units of nucleotides per kilobase.

Fig. 4 | Changes in non-coding RNA transcription predict changes in gene transcription. (a) The fraction of each indicated class of RNAs that undergo changes in transcription in human CD4+ T-cells (see Methods). The relationships among the indicated classes of transcription units are depicted at top. **(b)** Scatterplot shows the magnitude of changes in transcription predicted for 928 protein-coding genes using changes in the transcription of nearby non-coding RNAs (y-axis) as a function of changes observed (x-axis). The line has a slope of 1 and an intercept of 0.

Fig. 5 | Stabilizing selection on protein-coding gene transcription. **(a)** UCSC Genome Browser tracks show transcription, dREG signal, and ChIA-PET loop interactions near the CCR7 superenhancer in the human genome. PRO-seq tracks show transcription on the plus (red) and minus (blue) strands in units of RPKM. Net synteny tracks show regions of one-to-one orthology with the chimpanzee and rhesus macaque genomes. **(b)** Scatterplot shows the percentage of TREs conserved among all three primate species (y-axis) as a function of distance from the nearest annotated transcription start site (x-axis). The size of each point represents the amount of data in the corresponding distance bin. **(c)** The percentage of all dREG sites that are conserved in each indicated class of TRE. TREs are separated into three bins based on the distance relative to the nearest transcription start site. Error bars reflect a 1,000-sample bootstrap. **(d)** Cumulative distribution function of phyloP scores from the 100-way alignments in the indicated class of dREG site. The insert shows the fraction of sites in each class exceeding a phyloP score cutoff of 0.75. **(e)** Scatterplot shows promoter conservation (y-axis, black) or DNA sequence conservation (y-axis, red) as a function of the number of loop interactions made by that site to distal sites across the genome (x-axis). **(f)** TRE conservation (y-axis) or DNA sequence conservation (y-axis, red) as a function of the number of loop interactions made by the distal sequence element (x-axis). In panels (e-f) the size of each point is proportional to the number of examples in the corresponding bin.