Stochastic Model of Neuronal Outgrowth Movement: UNC-5 (UNC5) Regulates the Length and Number of Neuronal Processes in _Caenorhabditis elegans_

Gerard Limerick[*, †], Xia Tang[*, †], Won Suk Lee[*, †], Ahmed Mohamed[*], Aseel Al-Aamiri[*], and William G. Wadsworth[*]

[*]Department of Pathology and Laboratory Medicine, Rutgers Robert Wood Johnson Medical School, Piscataway, NJ 08854

[†]These authors contributed equally to this work

running title: neuronal outgrowth patterning

corresponding author:

     William G. Wadsworth

     Department of Pathology and Laboratory Medicine

     Rutgers Robert Wood Johnson Medical School

     675 Hoes Lane West

     Piscataway, NJ 08854-5635

     732-235-5768

     william.wadsworth@rwjms.rutgers.edu

Abstract

Neurons extend processes that vary in number, length, and direction of outgrowth. Extracellular cues help determine the patterning of outgrowth. In *Caenorhabditis elegans,* neurons respond to the extracellular UNC-6 (netrin) cue via UNC-40 (DCC) and UNC-5 (UNC5) receptors. We have postulated that UNC-40 undergoes stochastically orientated asymmetric localization (SOAL) within neurons. Extracellular cues govern the probability of UNC-40 localizing and mediating outgrowth at points along the surface of the neuron. For each instance of time there is a probability that UNC-40-mediated outgrowth will occur in a specific direction and so over time the direction of outgrowth fluctuates. Random walk modeling predicts that the degree of fluctuation affects the extent of outgrowth movement. Therefore, different patterns of outgrowth could be caused by regulating UNC-40 SOAL. Here we present evidence that UNC-5 (UNC5) receptor activity regulates UNC-40 SOAL and affects the length and number of processes that neurons develop. We find that loss of UNC-5 function increases the probability of UNC-40-mediated outgrowth in different directions, thereby increasing the degree of fluctuation. Consistent with the model, in *unc-5* loss-of-function mutants neurons fail to extend processes to full length or fail to develop multiple processes. We further show genetic interactions that suggest the UNC-5 and SAX-3 (Robo) receptors, and the cytoplasmic proteins, UNC-53 (NAV2), MIG-15 (NIK kinase), and MADD-2 (TRIM), function through specific signaling pathways to regulate UNC-40 SOAL in response to the UNC-6 and EGL-20 (wnt) extracellular cues. We propose genes influence the patterning of neuronal outgrowth by regulating the SOAL process.

Introduction

During development, an intricate network of neuronal connections is established. As processes extend from the neuronal cell bodies, distinct patterns of outgrowth emerge. Some extensions remain as a single process, whereas others branch and form multiple processes. If they branch, the extensions can travel in the same or in different directions. Processes vary in length. Extracellular cues are known to influence this patterning, but the underlying logic that governs the formation of patterns remains a mystery.

The secreted extracellular UNC-6 (netrin) molecule and its receptors, UNC-5 (UNC5) and UNC-40 (DCC) are highly conserved in invertebrates and vertebrates, and are known to play key roles in cell and axon migrations. In *Caenorhabditis elegans*, UNC-6 is produced by ventral cells in the midbody and by glia cells at the nerve ring in the head (WADSWORTH *et al.* 1996; WADSWORTH AND HEDGECOCK 1996; ASAKURA *et al.* 2007). It's been observed that neurons that express the receptor UNC-40 (DCC) extend axons ventrally, towards the UNC-6 sources; whereas neurons that express the receptor UNC-5 (UNC5) alone or in combination with UNC-40 extend axons dorsally, away from the UNC-6 sources (HEDGECOCK *et al.* 1990; LEUNG-HAGESTEIJN *et al.* 1992; CHAN *et al.* 1996; WADSWORTH *et al.* 1996).

It is commonly proposed that axons are guided by attractive and repulsive mechanisms (Tessier-Lavigne and Goodman 1996). According to this model, extracellular cues act as attractants or repellants to direct neuronal outgrowth towards or away from sources of the cues. UNC-5 (UNC5) has been described as a "repulsive" netrin receptor because it mediates guidance away from netrin sources (Leung-Hagesteijn et al. 1992; Hong et al. 1999; Keleman and Dickson 2001; Moore et al. 2007). The attraction and repulsion model is deterministic. That is, given the same

conditions, the response of the neuron, attractive or repulsive, will always be the same. As

such, the neuronal response to a particular cue is thought to be mediated by attractive or

repulsive signaling pathways. In genetic studies, a mutation that disrupts movement towards

the cue source denotes gene function within an attractive pathway, whereas mutations that

disrupt movement away from a source denotes gene function within a repulsive pathway.

Furthermore, the model predicts that the responsiveness of a neuron must switch from

attractive to repulsive if the axonal growth cone move towards and then away from the source

of a cue.

We have proposed an alternative model in which the movement of neuronal outgrowth is

established through a stochastic process. This model is based on evidence indicating that the

asymmetrical localization of UNC-40 in the neuron is self-organizing and that the surface to

which UNC-40 localizes and mediates outgrowth is stochastically determined (Figure 1A) (XU *et*

al. 2009; KULKARNI *et al.* 2013).   Because of this process, which we name "stochastically

orientated asymmetric localization" (SOAL), there is a probability that UNC-40-mediated

outgrowth will take place at each point along the surface of the neuron. At any instance of time,

extracellular cues govern the probability of UNC-40-mediated outgrowth at each point

(KULKARNI *et al.* 2013; TANG AND WADSWORTH 2014; YANG *et al.* 2014). Outgrowth extension can

be envisioned as a series of steps, where at each step extracellular cues control the probability

for outgrowth in each direction from the point on the surface. This means that over time, the

direction of outgrowth movement fluctuates. Such movement can be mathematically described

as a random walk, *i.e.* a succession of randomly directed steps (Figure 1B). Random walk

movement is diffusive and one property of diffusive motion is that the mean square

displacement grows proportionate to the time traveled. Consequently, the more the direction

of movement fluctuates, the shorter the distance of travel is in a given amount of time (Figure

1B). In this paper, we propose that UNC-5 helps regulate this fluctuation and, therefore, the

extent of outgrowth movement (Figure 1A). In the neuron, this diffusive motion occurs at the

micro-scale as innumerable forces act upon the membrane. An increase in diffusive motion at

the micro-scale might be observed at the macro-scale as a decrease in the rate of outgrowth.

In this model, guidance cues promote or inhibit outgrowth but they do not intrinsically cause

an attractive or repulsive directional response. This is because the direction of outgrowth is

determined by a directional bias that is created over time by the combined effect of

extracellular cues. In the example illustrated in Figure 2A, a neuronal protrusion experiences

an extracellular guidance cue(s) (orange) that is present in a posterior (right) to anterior (left)

increasing concentration gradient. This cue acts together with other cues (purple and blue)

that set the probability of outgrowth in the dorsal and ventral directions. As long as the

probabilities of dorsal and ventral outgrowth are equal, a bias for anterior movement over time

is created. For example, at time 1 (Figure 2A top) the probability of anterior outgrowth is 0.8,

whereas the probability of dorsal outgrowth is 0.1 and probability of ventral outgrowth is 0.1.

As the protrusion travels in the anterior direction it encounters higher levels of the cue. Even if

the cue inhibits outgrowth, an anterior directional bias towards the source of that cue can be

maintained. For example, at time 2 (Figure 2B bottom), the probability of anterior outgrowth

may have been reduced to 0.4 because of the inhibitory effect of the cue. Assuming that

signaling mechanisms within the neuron remain relatively constant as the protrusion

transverses this environment, the probabilities of dorsal and ventral outgrowth will increase

(the sum of the probabilities must equal one because of SOAL). Therefore, the probability of

dorsal outgrowth will be 0.3 and the probability of ventral outgrowth will be 0.3 in this

example.   While this increases the back and forth fluctuation that occurs perpendicular to the

direction of outgrowth, it does not change the directional bias that occurs over time and

movement will still be towards the inhibitory cue.

The model makes predictions about patterns of outgrowth in response to extracellular cues. In

this example, the neuron's response to the extracellular cues does not have an effect on the

directional bias. However, random walk modeling (shown above each diagram and explained

in detail later) predicts that a change in the degree to which the direction of outgrowth

movement fluctuates will alter the extent of outgrowth.   The pattern of outgrowth can be

influenced by how great the fluctuation becomes as outgrowth moves towards a source of a

cue. For example, if the fluctuation becomes significant as a protrusion moves towards the

source, outgrowth will stall before reaching the source. However, if the fluctuation is not great

enough at the source of the cue, outgrowth could continue to move forward. This is because the

probability of movement in the opposite, inward, direction is low. For instance, a probability of

0.33 anterior, 0.33 dorsal, and 0.33 ventral outgrowth still creates an anterior directional bias.

Therefore, in contrast to the attraction and repulsion model, this model does not predict that

movement across a source of a cue requires the neuron to switch its responsiveness to the cue.

In the discussion above, a protrusion moves toward the source of an inhibitory cue. However, a

similar prediction can be made for movement towards the source of a cue that promotes

outgrowth. Assuming that receptor levels at each surface and signaling mechanisms remain

relatively constant as the protrusion transverses this environment, the probabilities of dorsal

and ventral outgrowth will increase as more receptors at these surfaces are stimulated due to

an increasing concentration of the cue. Correspondingly, the probability of outgrowth at the

anterior surface will decrease because of SOAL. As with the case of the inhibitory cue, the

result is an increase in the degree to which the direction of outgrowth fluctuates.

The SOAL model makes predictions about how guidance genes might affect outgrowth

patterning. The SOAL process allows the direction of outgrowth activity to fluctuate. In Figure

2, the relative degree to which UNC-40-mediated (red) and nonUNC-40-mediated (green)

outgrowth activity fluctuates under different circumstances is depicted by the extent to which

the colored areas cover the neuron's surface. Extracellular cues can influence the degree to

which the direction of outgrowth fluctuates (Figure 2A). Likewise, mutations that disrupt the

UNC-40 SOAL process could also change the degree to which the direction of UNC-40-mediated

outgrowth fluctuates. By altering the extent of outgrowth movement, these genes could

regulate the length of a neuron's extension (Figure 2B). In addition to length, the model

predicts that the number of processes may be affected. Since extracellular cues are not

uniformly distributed across a neuron's surface, the extent of outgrowth movement could vary

along the leading-edge surface of a neuron, leading to multiple outgrowths extending in the

same direction. Such phenotypes might not have been associated with guidance genes since

most genetic studies have focused on only whether movement is disrupted towards or away

from the source of a cue. UNC-5, in particular, has been regarded as the quintessential

repulsive guidance receptor and *unc-5* mutations are most often interpreted using this

perspective.

Because we observed that UNC-5 affects UNC-40 SOAL (KULKARNI *et al.* 2013), we hypothesized

that *unc-5* might regulate the length and number of neuronal processes. Therefore, we decided

to look at neuronal outgrowth patterns in *unc-5* mutants for changes in the length and number

of processes. In particular, we focused on outgrowth that was not directed away from UNC-6

sources since the SOAL model predicts that these outgrowth patterns are not caused by a

repulsion mechanism and that they should occur independently of the direction of outgrowth.

We further reasoned that if *unc-5* regulates this type of patterning then UNC-40 SOAL might be

regulated by specific genetic pathways. We therefore asked whether genetic interactions

between *unc-5* and other guidance genes can affect UNC-40 SOAL.

In this paper, we present genetic evidence that UNC-5 regulates the length and number of

processes that neurons development. We argue that these results can be interpreted using the

SOAL model. We also show genetic interactions that suggest the UNC-5 and SAX-3 (Robo)

receptors, and the cytoplasmic proteins, UNC-53 (NAV2), MIG-15 (NIK kinase), and MADD-2

(TRIM), function through specific signaling pathways to regulate UNC-40 SOAL in response to

the UNC-6 and EGL-20 (wnt) extracellular cues. Together these results suggest that the SOAL

model is useful for understanding how genes regulate patterns of outgrowth.

Materials and Methods

Strains

Strains were handled at 20 ℃ using standard methods (Brenner, 1974) unless stated

otherwise. A Bristol strain N2 was used as wild type. The following alleles were used: **LGI**, *unc-40(e1430), unc-40(ur304), zdIs5[mec-4::GFP]*; **LGII**, *unc-53(n152)*; **LGIV,** *unc-5(e152), unc-5(e53), unc-5(ev480), unc-5(ev585),egl-20(n585), kyIs262[unc-86::myr-GFP;odr-1::dsRed]*; **LGIV,** *madd-2(ky592), madd-2(tr103)*; **LGX**, *mig-15(rh148), unc-6(ev400), sax-3(ky123), sax-3(ky200).*

Transgenes maintained as extrachromosomal arrays included: *kyEx1212 [unc-86::unc-40-GFP;odr-1::dsRed].*

Analysis of axon outgrowth and cell body position

HSN neurons were visualized using expression of the transgene *kyIs262[unc-86::myr-GFP].* The mechanosensory neurons, AVM, ALM, and PLM, were visualized using the expression of the transgene *zdIs5[Pmec-4::GFP].* Synchronized worms were obtained by allowing eggs to hatch overnight in M9 buffer without food. The larval stage was determined by using differential interference contrast (DIC) microscopy to examine the gonad cell number and the gonad size. Staged larvae were mounted on a 5% agarose pad with 10 mM levamisole buffer. Images were taken using epifluorescent microscopy with a Zeiss 63X water immersion objective.

The number of processes during early L1 larval stage was scored by counting the number of processes that extended for a distance greater than the length of one cell body. We report instances in which there were no such processes, one process or more than one processes. In the L2 larval stage, a single early process was scored if there was only one major extension from the ventral leading edge. The HSN cell body in L2 stage larvae was scored as dorsal if the

cell body had failed to migrate ventrally and was not positioned near the PLM axon. In L4 stage

larvae, a multiple ventral processes phenotype was scored if more than one major extension

protruded from the ventral side of cell body.

Extension into the nerve ring was scored as defective if the axon did not extend further than

approximately half the width of the nerve ring. Anterior extension was scored as defective if

the axon did not extend further anteriorly than the nerve ring. PLM axons are scored as over-

extending if they extended further anterior than the position of the ALM cell body.

Analysis of the direction of HSN outgrowth

HSN was visualized using the transgene *kyIs262[unc-86::myr-GFP]*. L4 stage larvae were

mounted on a 5% agarose pad with 10 mM levamisole buffer. An anterior protrusion was

scored if the axon extended from the anterior side of the cell body for a distance greater than

the length of three cell bodies. A dorsal or posterior protrusion was scored if the axon extended

dorsally or posteriorly for a distance greater than two cell body lengths. HSN was considered

multipolar if more than one process extended a length greater than one cell body. Images were

taken using epifluorescent microscopy with a Zeiss 40X objective.

Analysis of the UNC-40::GFP localization in L2 stage animal

For analysis of UNC-40::GFP localization, L2 stage larvae with the transgenic marker

kyEx1212[unc-86::unc-40::GFP; odr-1::dsRed] were mounted on a 5% agarose pad with 10 mM

levamisole buffer. Staging was determined by examining the gonad cell number and the gonad

size under differential interference contrast (DIC) microscopy. Images were taken using

epifluorescent microscopy with a Zeiss 63X water immersion objective. The UNC-40::GFP

localization was determined by measuring the average intensity under lines drawn along the dorsal and ventral edges of each HSN cell body by using ImageJ software. For analysis of the anterior–posterior orientation of UNC-40::GFP, the dorsal segment was geometrically divided into three equal lengths (dorsal anterior, dorsal central and dorsal posterior segments). The line-scan intensity plots of each of these segments were recorded. ANOVA test was used to determine if there is a significant difference between intensities of the three segments. The dorsal distribution was considered uniform if $p \geq 0.05$ and was considered asymmetrical if $p \leq 0.05$. Within an asymmetric population, the highest percent intensity was considered to localize UNC-40::GFP to either anterior, posterior or central domain of the dorsal surface.

Computations

A program to simulate a two-dimensional lattice random walk based on the probability of dorsal, ventral, anterior, and posterior outgrowth for a mutant (Table 1) was created using MATLAB. (The directions of the axons from multipolar neurons were not scored. These axons appear to behave in the same manner as the axons from monopolar neurons, but this has not yet been tested.) The probability of dorsal, ventral, anterior, or posterior outgrowth was assigned for the direction of each step of a random walk moving up, down, left or right, respectively. Each variable is considered independent and identically distributed. Simulations of 500 equal size steps (size =1) were plotted for 50 tracks (Figure 1B, 5B and 6B inserts). A Gaussian distribution for the final positions of the tracks was generated using Matlab's random function (Figure 6).

The mean squared displacement (MSD) is used to provide a quantitative characteristic of the

motion that would be created by the outgrowth activity undergoing the random walk. Using

the random walks generated for a mutant the MSD can be calculated:

$$msd(\tau) = <[r(t + \tau) - r(t)]^2>$$

Here, r(t) is the position at time t and τ is the lag time between two positions used to calculate

the displacement, $\Delta r(\tau) = r(t+\tau) - r(t)$. The time-average over t and the ensemble-average over

the 50 trajectories were calculated. This yields the MSD as a function of the lag time. A

coefficient giving the relative rate of diffusion was derived from a linear fit of the curve. The

first two lag time points were not considered, as the paths often approximate a straight line at

short intervals.

Results

UNC-5 regulates the pattern of outgrowth from the HSN neuron

To investigate whether UNC-5 activity can regulate the length or number of processes that a neuron can develop when outgrowth is towards an UNC-6 source, we examined the development of the HSN axon in *unc-5* mutations. The HSN neuron sends a single axon to the ventral nerve cord, which is a source of the UNC-6 cue (WADSWORTH *et al.* 1996; ADLER *et al.* 2006; ASAKURA *et al.* 2007). Axon formation is dynamic (ADLER *et al.* 2006). Shortly after hatching, HSN extends short neurites in different directions. These neurites, which dynamically extend and retract filopodia, become restricted to the ventral side of the neuron where a leading edge forms. Multiple neurites extend from this surface until one develops into a single axon extending to the ventral nerve cord. Measurements of growth cone size, maximal length, and duration of growth cone filopodia indicate that UNC-6, UNC-40, and UNC-5 control the dynamics of protrusion (NORRIS AND LUNDQUIST 2011).

We observe that in *unc-5* mutants, the patterns of extension are altered. In wild-type animals at the L1 stage of development most HSN neurons extends more than one short neurite, however in *unc-5(e53)* mutants nearly half the neurons do not extend a process (Figures 3A and 3B). During the L2 stage in wild-type animals a prominent ventral leading edge forms and the cell body undergoes a short ventral migration that is completed by the L3 stage. By comparison, in *unc-5* mutants the cell body may fail to migrate and instead a single large ventral process may form early during the L2 stage (Figures 3A, 3C and 3E). It may be that the ventral migration of the HSN cell body requires the development of a large leading edge with multiple extensions. Together the observations indicate that loss of *unc-5* function affects the patterning of outgrowth, *i.e.* the timing, length, and number of extensions that form. Loss of *unc-5* function does not prevent movement, in fact, a single large

ventral extension can form in the mutant at a time that is even earlier than when a single ventral

extension can be observed in wildtype.

We tested four different *unc-5* alleles in these experiments. The *unc-5(e53)* allele is a putative

molecular null allele, *unc-5(ev480)* is predicted to truncate UNC-5 after the cytoplasmic ZU-5

domain and before the Death Domain, *unc-5(e152)* is predicted to truncate UNC-5 before the

ZU-5 domain and Death Domain, and *unc-5(ev585)* is a missense allele that affects a predicted

disulfide bond in the extracellular Ig(C) domain (KILLEEN *et al.* 2002). Although both the *unc-*

5(ev480) and *unc-5(e152)* are predicted to cause premature termination of protein translation

in the cytodomain, the *unc-5(e152)* product retains the signaling activity that prevents these

phenotypes. Based on other phenotypes, previous studies reported that the *unc-5(e152)* allele

retains UNC-40-dependent signaling functions (MERZ *et al.* 2001; KILLEEN *et al.* 2002).

UNC-5 is required for the induction of multiple HSN axons by UNC-6ΔC and a *mig-15*

mutation

The results above suggest that UNC-5 activity can regulate the number of HSN extensions that

form. To further test this hypothesis, we checked whether loss of UNC-5 function can suppress

the development of additional processes that can be induced. Previously we reported that

expression of the N-terminal fragment of UNC-6, UNC-6ΔC, induces excessive branching of

ventral nerve cord motor neurons and that UNC-5 can suppress this branching (LIM *et al.* 1999).

We now report that HSN develops an extra process in response to UNC-6ΔC and that UNC-5

suppresses the development of this extra process (Figures 3D and 3F).

To investigate whether this UNC-5 activity might involve known effectors of asymmetric neuronal outgrowth, we tested for genetic interactions between *unc-5* and both *mig-10* and *mig-15*. MIG-10 (lamellipodin) is a cytoplasmic adaptor protein that can act cell-autonomously to promote UNC-40-mediated asymmetric outgrowth (ADLER *et al.* 2006; CHANG *et al.* 2006; QUINN *et al.* 2006; QUINN *et al.* 2008; MCSHEA *et al.* 2013). MIG-15 (NIK kinase) is a cytoplasmic protein and evidence indicates that *mig-15* functions cell-autonomously to mediate a response to UNC-6 (POINAT *et al.* 2002; TEULIÈRE *et al.* 2011). It's proposed that *mig-15* acts with *unc-5* to polarize the growth cone's response and that it controls the asymmetric localization of MIG-10 and UNC-40 (TEULIÈRE *et al.* 2011; YANG *et al.* 2014). We previously noted that HSN neurons often become bipolar in *mig-15* mutants and frequently UNC-40::GFP is localized to multiple surfaces in a single neuron, suggesting that loss of MIG-15 enhances the ability of UNC-40::GFP to cluster (YANG *et al.* 2014). In our experiments we used the *mig-10 (ct141)* loss-of-function allele (MANSER AND WOOD 1990; MANSER *et al.* 1997) and the *mig-15(rh148)* allele, which causes a missense mutation in the ATP-binding pocket of the kinase domain and is a weak allele of *mig-15* (SHAKIR *et al.* 2006; CHAPMAN *et al.* 2008).

We find that the extra processes induced by UNC-6ΔC expression are suppressed by *mig-10(ct141)* (Figures 3F). We also find that the *mig-15* mutation causes extra HSN processes and that the loss of UNC-5 function suppresses these extra HSN processes (Figures 3F and 3G). These results support the hypothesis that the ability of UNC-5 to regulate the development of multiple protrusions involves the molecular machinery that controls UNC-40-mediated asymmetric neuronal outgrowth.

UNC-5 is required for PLM overextension

The SOAL model predicts that the ability of UNC-5 to regulate the length and number of neural

protrusions is independent of the direction of outgrowth. HSN sends a single axon ventrally, while

PLM sends an axon anteriorly from a posteriorly positioned cell body. The HSN axon travels

towards UNC-6 sources, whereas the PLM axon pathway is perpendicular to UNC-6 sources.   To

investigate whether UNC-5 activity can regulate the length or number of processes that develop

perpendicular to UNC-6 sources we examined the development of the PLM axon. We also chose

PLM because UNC-5 is already known to affect the length of the PLM axon. It was previously

reported that in *rpm-1* mutants the PLM axon will overextend in the anterior direction and that

this phenotype can be suppressed by *sax-3* and *unc-5* loss-of-function mutations (LI *et al.* 2008).

In *rpm-1* loss-of-function mutations there is an increase in the level of UNC-5::GFP expression,

suggesting that the level of UNC-5 within PLM is important for controlling overextension.

However, the overextension in the *rpm-1* mutant is not suppressed by a loss-of-function *unc-40*

mutation (LI *et al.* 2008), suggesting that the overextension does not involve UNC-40 SOAL, but

rather another outgrowth activity that is nonUNC-40-mediated.

Given that UNC-5 activity is involved in the overextension of the PLM axon, and that the *mig-15*

mutation affects HSN outgrowth in an UNC-40 dependent fashion, we decided to test whether

PLM overextension might be induced by the *mig-15* mutation in an UNC-40-dependent fashion.

The HSN results suggest that altering *mig-15* function creates a sensitized genetic background.

That is, the *unc-5(ev480)* mutation suppresses HSN outgrowth extension in both the wild-type

and *mig-15(rh148)* backgrounds, but the *mig-15* mutation creates a stronger patterning

phenotype.   This idea is supported by the evidence that the *mig-15* mutation enhances the

ability of UNC-40 to localize at surfaces (YANG *et al.* 2014).

We find that in *mig-15(rh148)* mutants the PLM axon often fails to terminate at its normal

position and instead extends beyond the ALM cell body. This overextension is suppressed in

unc-5(e53);mig-15(rh148) and *unc-40(e1430);mig-15(rh148)* mutants (Figures 4A and 4B).

The results are consistent with the idea that UNC-5 is required for the UNC-40-mediated

outgrowth activitythat causes overextension in *mig-15(rh148)* mutants.

UNC-5 is required for ALM and AVM branching and extension

We also investigated the effect of UNC-5 activity on patterning where sources of UNC-6 and

other cues are in a more complex arrangement. Specifically, we examined whether UNC-5 plays

a role in the outgrowth of AVM and ALM processes at the nerve ring. During larval

development, processes from the AVM neuron and the two ALM neurons (one on each side of

the animal) migrate anteriorly to the nerve ring at dorsal and ventral positions respectively

(Figure 4C). At the nerve ring each axon branches; one branch extends further anteriorly and the

other extends into the nerve ring. Evidence suggests that at the midbody of the animal the

positioning of these axons along the dorsal-ventral axis requires UNC-6, UNC-40, and UNC-5

activity.   In *unc-6*, *unc-40,* and *unc-5* null mutants, or when the UNC-6 expression pattern is altered,

the longitudinal nerves are mispositioned (REN *et al.* 1999). Glia cells and neurons at the nerve ring

are sources of UNC-6 (WADSWORTH *et al.* 1996). The guidance of some axons in the nerve ring are

disrupted in *unc-6* and *unc-40* mutants (HAO *et al.* 2001; YOSHIMURA *et al.* 2008).   The precise

spatial and temporal arrangement of the UNC-6 cue in relationship to the position of the

migrating growth cones is not fully understood.   Nevertheless, the anteriorly migrating growth

cones appear to use the UNC-6 cue from the ventral sources to help maintain the correct dorsal-

ventral position, even while moving towards the nerve ring, which is a new source of UNC-6

that is perpendicular to the ventral source.   At the nerve ring the axons branch. One process

continues anteriorly, moving past the new UNC-6 source, whereas the other projects at a right

angle and moves parallel to the new source.

We find genetic interactions involving *unc-5, unc-40,* and *mig-15* that affect outgrowth

patterning of the ALM and AVM extensions at the nerve ring (Figures 4C, 4D, and 4E). In *mig-15(rh148);unc-5(e53)* mutants, the AVM axon often fails to extend anteriorly from the branch

point and only extends into the nerve ring, or it fails to extend into the nerve ring and only

extends anteriorly, or it fails to do both and terminates at this point. In *unc-40(e1430)* mutants,

the axon often fails to branch into the nerve ring, although it extends anteriorly. In comparison,

in *unc-40(e1430);mig-15(rh148)* mutants more axons extend into the nerve ring. These results

suggest that UNC-5 helps regulate UNC-40-mediated outgrowth to pattern the outgrowth at the

nerve ring.

Interactions between *unc-5* and other genes affect the probability of HSN extension in each direction

We hypothesize that there are interactions between *unc-5* and other genes that control the

degree to which the direction of outgrowth fluctuates. In the SOAL model, the collective effect

of all the cues that promote and inhibit outgrowth set a probability of outgrowth at each

surface at each instance of time. As the direction of outgrowth fluctuates over time, a bias is

created that determines directionality. Since the direction of outgrowth is stochastically

determined, gene activity is not inherently associated with movement in a specific direction.

This model differs from the attraction and repulsion model where gene activity is assayed

based on direction, *i.e.* the direction of movement relative to the source of a cue. The attraction

and repulsion model places genes into attractive or repulsive pathways. We hypothesize that

unc-5 and other genes might instead be ordered into pathways based on how they affect the

fluctuation of outgrowth activity.

Probability distributions for the direction of outgrowth are used to study how genes affect the

fluctuation of outgrowth activity. By comparing the distributions created from wild-type and

mutant animals, the relative effect that genes have on the fluctuation can be determined. To

accomplish this, the direction that the HSN axon initially projects from the cell body is scored

(Figure 5A). We reason that the initial development of an axon is the result of the collective

impact of all the individual outgrowth movements that take place during the initial formation of

the axon. In wildtype, there is a high probability for outgrowth in the ventral direction at each

instance of time. As such, there is little fluctuation over time and a strong bias for ventrally

directed outgrowth is created. A mutation can cause a lower probability of outgrowth in the

ventral direction and a higher probability for outgrowth in other directions. This creates

greater fluctuation over time and a weaker bias of ventrally directed outgrowth. For this assay,

all the outgrowth movement during the initial axon extension is treated as a single event, *i.e.* the

movement of the membrane occurs in a single step from the cell body to a position that is at a

set length from the cell body (see Materials and Methods). This step is in the anterior,

posterior, dorsal, or ventral direction. Thus, direction is the random variable which takes a

value of anterior, posterior, dorsal, or ventral. We can determine the probability of occurrence

for each value by scoring many individual animals for each strain.

Using this assay, we tested for genetic interactions between *unc-5* and four other genes; *elg-20,*

sax-3, madd-2, or *unc-6.* We have chosen these particular genes because previous observations

suggest interactions. 1) EGL-20 (Wnt) is a secreted cue expressed from posterior sources (PAN

et al. 2006) and it affects to which surface of the HSN neuron the UNC-40 receptor localizes and mediates outgrowth (KULKARNI *et al.* 2013). Based on a directional phenotype, a synergistic interaction between *unc-5* and *egl-20* has been observed. In either *unc-5* or *egl-20* mutants the ventral extension of AVM and PVM axons is only slightly impaired, whereas in the double mutants there is much greater penetrance (LEVY-STRUMPF AND CULOTTI 2014). 2) SAX-3(Robo) is a receptor that regulates axon guidance and is required for the asymmetric localization of UNC-40 in HSN (TANG AND WADSWORTH 2014). Based on a directional phenotype, SAX-3 and UNC-40 appear to act in parallel to guide the HSN towards the ventral nerve cord (XU *et al.* 2015). 3) MADD-2 is a cytoplasmic protein of the tripartite motif (TRIM) family that potentiates UNC-40 activity in response to UNC-6 (ALEXANDER *et al.* 2009; ALEXANDER *et al.* 2010; HAO *et al.* 2010; MORIKAWA *et al.* 2011; SONG *et al.* 2011; WANG *et al.* 2014). MADD-2::GFP and F-actin colocalize with UNC-40::GFP clusters in the anchor cell (WANG *et al.* 2014). 4) Of course, UNC-6 is an UNC-5 ligand. DCC (UNC-40) and UNC5 (UNC-5) are thought to act independently or in a complex to mediate responses to netrin (UNC-6) (COLAVITA AND CULOTTI 1998; HONG *et al.* 1999; MACNEIL *et al.* 2009; LAI WING SUN *et al.* 2011).

In a test for interaction with *egl-20*, we find that in comparison to *unc-5(e53)* or *egl-20(n585)* mutants, the *unc-5(e53);egl-20(n585)* double mutant have a lower probability for ventral outgrowth and higher probability for outgrowth in other directions (Table 1). This suggests that *unc-5* and *egl-20* may act in parallel to achieve the highest probability for HSN ventral outgrowth, *i.e.* they act to prevent UNC-40-mediated outgrowth from fluctuating in other directions.

In a test for interaction with *sax-3,* we find that the probability of outgrowth in each direction in *unc-5(e53);sax-3(ky200)* mutants is similar to the probabilities in *sax-3(ky200)*or *sax-3(ky123)* mutants (Table 1). Given the results with *unc-5* and *egl-20*, we further tested the probability of outgrowth in each direction in *egl-20(n585);sax-3(ky123)* mutants. We find that it is similar to the probabilities in *sax-3(ky200)*or *sax-3(ky123)* mutants (Table 1). The *sax-3(ky123)* allele results in a deletion of the signal sequence and first exon of the gene, whereas sax-3(ky200) contains a missense mutation which is thought to cause protein misfolding and mislocalization at the restrictive temperature (25°C) (ZALLEN *et al.* 1998; WANG *et al.* 2013). The *egl-20(n585);sax-3(ky123)* mutants do not grow well and so it is easier to use the temperature sensitive *sax-3* allele. Together, the results suggest that *sax-3* may be required for both the *unc-5-* and the *egl-20*-mediated activities that allow the highest probability for HSN ventral outgrowth.

In a test for interaction with *madd-2,* we find that the probability of outgrowth in each direction in *unc-5(e53);madd-2(tr103)* mutants is similar to the probabilities in *madd-2(tr103)* mutants (Table 1). There is a higher probability for anterior HSN outgrowth, similar to what is observed in *unc-40(e1430)* mutants. These results suggest that *madd-2* might be required for the *unc-40* outgrowth activity. The probability of outgrowth in each direction in *madd-2(tr103);sax-3(ky123)* mutants is similar to the probabilities in *sax-3(ky200)*or *sax-3(ky123)* mutants (Table 1). The *madd-2(tr103)* allele appears to act as a genetic null (ALEXANDER *et al.* 2010).

In a test for interaction with *unc-6,* we find that the probability of outgrowth in each direction in *unc-5(e53);unc-6(ev400)* and *unc-40(e1430);unc-5(e53)* mutants is similar to the probabilities in *unc-6(ev400)* mutants insofar as there is a lower probability for ventral outgrowth and a higher probability for anterior outgrowth (Table 1). However, the probabilities in each

direction are closer to those obtained from the *unc-40(e1430)* mutants because the probability

of anterior outgrowth is lower in these mutants than in *unc-6* mutants. This suggest that UNC-5

and UNC-40 might help increase the probability of anterior outgrowth in the absence of UNC-6.

Taken together, these results support the hypothesis that there are genetic interactions that

regulate the degree to which the direction of outgrowth fluctuates. Rather than defining the

function of a gene by its ability to control a deterministic event, *i.e.* outgrowth in one direction,

gene function can be defined by its ability to control a stochastic process, *i.e.* the probability of

outgrowth in different directions.

***unc-5* is a member of a class of genes that has a similar effect on the spatial extent of movement**

How does *unc-5* affect outgrowth movement? The results above show that *unc-5* and its

interactions with other genes affect the degree to which the direction of outgrowth fluctuates.

The degree of fluctuation differs depending on the genes involved. The SOAL model predicts

that these differences alter the extent of outgrowth; the more the direction of movement

fluctuates, the shorter the distance of travel is in a given amount of time.

To depict how *unc-5* and other genes differentially regulate outgrowth movement, we use

random walk modeling. Random walks describe movement that occurs as a series of steps in

which the angles and the distances between each step is decided according to some probability

distribution. By using the probability distribution obtained from a mutant for each step of a

random walk, and by keeping the distance of each step equal, a random walk can be

constructed (Figure 5A). In effect, this method applies the probability distribution to discrete

particles having idealized random walk movement on a lattice. By plotting random walks

derived from wild-type animals and different mutants, the relative effect that mutations have

on random walk movement can be visualized. For example, Figure 5B shows 50 tracks of 500

steps for wildtype and two mutants (mutant A is *unc-5(e53)* and mutant B is *egl-20(n585);sax-*

3(ky123)). This reveals the effect that a mutation has on the displacement of movement. After

500 steps the displacement from the origin (0,0) is on average less for mutant A than for

wildtype, and less for mutant B than for wildtype or mutant A.

The random walk models show the relative effect that a mutation has on a property of

outgrowth movement. It is worth noting that this is not modeling the actual trajectory of

migrating axons. Neuronal outgrowth is essentially a mass transport process in which mass

(the molecular species of the membrane) is sustained at the leading edge and moves outward.

Our assay compares the effect that different mutations would have on the movement of mass at

the leading edge of an extension if the conditions of the system were kept constant. Of course,

in vivo the conditions are not constant. For one, as an extension moves it will encounter new

environments where the cues may be new or at different concentrations, all of which affect the

probability distribution. The actual patterns of outgrowth observed are the result of all the

probabilities for outgrowth that occur at each instance of time. It has recently been suggested

that our description might be more accurately described as neuro-percolation, a superposition

of random-walks (AIELLO 2016).

Our random walk analysis compares the effect that different mutations have on the properties

of movement. In wild-type animals, there is a high probability for outgrowth in the ventral

direction. The analysis shows that conditions in wildtype create nearly straight-line movement,

i.e. if the same random walk is repeatedly done for the same number of steps, starting at the same origin, the final position of the walk along the x axis does not vary a great amount. In comparison, we find that a mutation can create random walk movement in which the final position is more varied. This variation occurs because the mutation increases the probability of outgrowth in other directions. For each mutation, we simulate 50 random walks of 500 steps and derive the mean and standard deviation of the final position along the X-axis. To compare strains, we plot the normal distribution, setting the mean at the same value for each. The difference between the curve for a mutant and wildtype shows the degree to which the mutation caused the direction of outgrowth to fluctuate (Figure 5C).

The results reveal four different distribution patterns (Figure 6). The first class is the wild-type distribution, which has the distribution curve with the highest peak. The second class comprises *unc-5, egl-20, unc-53,* and *unc-6* in which the distribution curve is flatter than the wild-type curve. We included *unc-53* because our previous study showed that it has genetic interactions with *unc-5* and *unc-6* (KULKARNI *et al.* 2013). The *unc-53* gene encodes a cytoskeletal regulator related to the mammalian NAV proteins and *unc-53* mutations cause guidance defects (MAES *et al.* 2002; STRINGHAM *et al.* 2002; STRINGHAM AND SCHMIDT 2009). The third class has a distribution curve which is flatter than the second and comprises *sax-3, mig-15,* and several double mutation combinations (Figure 6). The fourth class has the flattest distribution curve and comprises *egl-20;sax-3*, *unc-40;sax-3,* and *unc-53;sax-3;unc-6*. This class indicates the greatest degree of fluctuation. The ability to cause the direction of movement to fluctuate is not associated with a specific direction of HSN movement. For example, *unc-5;sax-3, unc-53;unc-6, unc-40;egl-20,* and *madd-2;sax-3* each show a widely dispersed pattern, but the direction is ventral, dorsal, anterior, and posterior, respectively (Figure 6).

The distribution patterns indicate that genes have different effects on the extent that outgrowth

movement can travel through the environment. Mean squared displacement (MSD) is a

measure of the spatial extent of random motion. The MSD can be calculated from the random

walk data. Plotting MSD as a function of the time interval shows how much an object displaces,

on average, in a given interval of time, squared (Figure 7A). For normal molecular diffusion, the

slope of the MSD curve is directly related to the diffusion coefficient. In cell migration models

this value is referred to as the random motility coefficient. Coefficients are experimentally

determined; they describe how long it takes a particular substance to move through a

particular medium. We determine this value in order to numerically and graphically compare

how mutations can alter displacement relative to wildtype (Figure 7B). The four classes of

genes are apparent by comparing the height of the bars in Figure 7B.

The results of this modeling suggest that the activities of certain genes, and combinations of

genes, have distinct effects on the rate of outgrowth movement. In theory, these differences

could be an important means by which genes cause different outgrowth patterns.

unc-5 and other genes regulate UNC-40 receptor clustering, which is a consequence of the

SOAL process

How does unc-5 affect the localization of the UNC-40 receptor and how does this localization

affect UNC-40-mediated outgrowth activity and the pattern of outgrowth? Beginning in the

early L2 stage, UNC-40::GFP becomes localized to the ventral side of HSN in wildtype (ADLER *et*

al. 2006; KULKARNI *et al.* 2013). Reflecting the dynamic morphological changes that occur as the

HSN axon forms, the site of asymmetric UNC-40::GFP localization alternates in the neurites and

along the ventral surface of the neuron (KULKARNI *et al.* 2013). Dynamic UNC-40::GFP

localization patterns have also been reported for the anchor cell, in which UNC-40 and UNC-6

are also key regulators of extension (ZIEL *et al.* 2009; HAGEDORN *et al.* 2013). Live imaging of the

anchor cell reveals that UNC-40::GFP "clusters" form, disassemble, and reform along the

membrane (WANG *et al.* 2014). However, live imaging can't directly ascertain whether the

position of a cluster is randomly determined since a movement event cannot be repeatedly

observed to determine a probability distribution. Mathematical modeling of cluster movement

as a stochastic process has not been done. Nevertheless, the results are consistent with the fact

that a probability distribution for the direction of UNC-40-mediated HSN outgrowth movement

can be determined and, therefore, the UNC-40 localization that gives rise to the outgrowth

activity can be defined as a stochastic process.

The UNC-40::GFP clustering phenomena raises questions about the relationship between

robust UNC-40 clustering (*i.e.,* sites of distinct UNC-40 localization observable by UNC-40::GFP)

and UNC-40-mediated outgrowth activity. Two models are presented in Figure 8. In the first

model, the SOAL mechanism causes UNC-40 clustering, which is required for UNC-40-mediated

outgrowth activity (Figure 8A). In the second model, SOAL and outgrowth activities are

independent and can happen concurrently (Figure 8B). In the first model, the output of the

SOAL process is receptor clustering. This is next followed by outgrowth activity from the

locations at the membrane where the clusters become stable. This model renders UNC-40-

mediated outgrowth a deterministic event occurring at the macro-scale. In the second model,

the SOAL process and outgrowth activity are concurrent activities which can occur

stochastically along the membrane at the micro-scale. UNC-40 clustering in this model might be

an observable consequence of the micro-scale stochastic process that arises over time.

The models make specific predictions that can be tested. In the first model, UNC-40-mediated

outgrowth will not happen if UNC-40 does not cluster. In the second model, the loss of UNC-40

clustering does not lead to a loss of UNC-40-mediated outgrowth. We favor the second model.

In the *sax-3* mutant there is a large fluctuation in the direction of outgrowth; it is in the third

class of mutants (Figures 6 and 7). We previously reported that *sax-3* is required for robust

UNC-40::GFP asymmetric localization; in *sax-3* mutants UNC-40::GFP remains uniformly

dispersed around the periphery of HSN ((TANG AND WADSWORTH 2014) and Figure 9). Whereas

in the *sax-3* mutant there is a ventral bias for outgrowth, in the *unc-40;sax-3* mutant there is not

(Figure 6). This suggests that in the *sax-3* mutant there is UNC-40-mediated outgrowth activity

that helps create a ventral bias. This is consistent with the second model because UNC-40-

mediated outgrowth activity is occurring even when robust UNC-40::GFP is not observed.

We hypothesize that a consequence of the micro-scale SOAL process over time is macro-scale

UNC-40 clustering. If so, then *unc-5* activity should affect UNC-40::GFP clustering because it

affects the degree to which the direction of UNC-40 receptor localization fluctuates. However,

even though there is a higher probability that localization occurs at surfaces other than at the

ventral surface, we observe robust asymmetrically localized UNC-40::GFP clustering in *unc-*

5(e53) mutants (KULKARNI *et al.* 2013). We speculate that *unc-5(e53),* as well as other gene

mutations, do not cause the direction of UNC-40 localization to fluctuate enough to prevent

observable UNC-40::GFP clustering. We decided to examine UNC-40::GFP clustering in double

mutants to see whether there is an additive effect. In double mutants will the direction of UNC-

40 localization fluctuate enough so that UNC-40::GFP clustering cannot be observed in our

assay?

We made double mutant combinations between *unc-5*, and *egl-20* or *unc-53*. In *egl-20* and *unc-53* single mutants there is fluctuation in the direction of outgrowth (Figures 6 and 7) and robust asymmetrical UNC-40::GFP localization (Figure 9, *unc-53* results were previously reported (KULKARNI *et al.* 2013)). In comparison to the single mutants, the double mutants all show an increase in the degree to which the direction of outgrowth fluctuates (Figures 6 and 7). Further, in contrast to the single mutants, UNC-40::GFP remains uniformly dispersed around the periphery of HSN in the double mutants (Figure 9). The results suggest a correlation between the degree to which UNC-40-mediated outgrowth activity fluctuates and the ability to detect UNC-40::GFP clustering. This is consistent with the second model (Figure 8). We also observe that in *madd-2(tr103)* mutants the direction of outgrowth fluctuates (Table 1), but unlike *egl-20* and *unc-53* single mutants, there is not robust asymmetrical UNC-40::GFP localization and UNC-40::GFP remains uniformly dispersed (Figure 9). The double mutants, *unc-5;madd-2,* are similar to the single *madd-2* mutant. Similar results are observed with *sax-3* and *unc-5;sax-3* mutants (Figure 9). We hypothesize that in the *madd-2* and *sax-3* mutations the degree to which the direction of UNC-40 localization fluctuates is so great that the *unc-5* mutation makes no difference on the UNC-40::GFP clustering phenotype.

We suggest these results support our model in which UNC-40-mediated outgrowth is an independent process that is coupled to the SOAL process in the HSN neuron. Previously we hypothesized that UNC-40 SOAL is a general cellular process that polarizes and orients cellular activities to the surrounding environment (YANG *et al.* 2014). In neurons during the development of axons, UNC-40 SOAL is coupled to the outgrowth machinery. However, UNC-40 SOAL may also become coupled to other cellular activities. For example, UNC-6, UNC-5, and

UNC-40 are also known to play a role in localizing synapses in neurons (CoLON-RAMOS *et al.* 2007; Poon *et al.* 2008; KILLEEN 2009).

Discussion

In this paper, we show that UNC-5 regulates the number and length of neuronal processes that a neuron can develop. We observe that *unc-5* loss-of-function mutations inhibit the development of multiple neurites during early outgrowth from HSN. They also suppress the development of extra HSN processes that are induced by a *mig-15* mutation or by expression of the N-terminal fragment of UNC-6. We also observe that *unc-5* mutations suppress the anterior overextension of the PLM axon that occurs in the *mig-15* mutant.   Finally, in combination with the *mig-15* mutation, *unc-5* loss-of-function mutations affect the branching and extension of ALM and AVM axons at the nerve ring. In each of these cases, the pattern of outgrowth is altered by loss of UNC-5 function. We further show that UNC-5 acts together with UNC-6, EGL-20, SAX-3, UNC-53, MIG-15, and MADD-2 to regulate UNC-40 asymmetric localization at a surface of HSN. Here we discuss a model to explain how UNC-5 could control patterns of outgrowth by regulating UNC-40 stochastically orientated asymmetric localization (SOAL).

UNC-40 SOAL; a model to explain *unc-5* phenotypes

We hypothesize that UNC-40 undergoes stochastically orientated asymmetric localization (SOAL) within neurons.   Because this is a stochastic process, there is a probability of UNC-40 localizing and mediating outgrowth at each surface of the neuron. The probability of outgrowth at each point along a surface may vary. This means that at one instance of time there is a probability that UNC-40-mediated outgrowth will occur in a specific direction from a point along the surface of the neuron. At the next instance of time there is a probability that

outgrowth will not occur at this point, but will instead occur at another point and in a different

direction. When considered over time, the direction of outgrowth fluctuates. This type of

movement can be modeled as a random walk, and random walks predict that the displacement

of the membrane will depend on the degree to which the direction of outgrowth movement

fluctuates. Extracellular cues, and the neuron's response to these cues, set the probability of

outgrowth happening at each point along the surface and therefore regulate the amount of

fluctuation. The process occurs at the micro-scale and it controls the direction and rate of

outgrowth movement. Below we describe how this model can explain the patterns of

outgrowth extension seen in the *unc-5* mutants.

Figure 2 depicts the essential concepts underlying the explanations. The same depiction is used

to explain the mutant phenotypes in Figures 10-15. Four important concepts are presented by

these figures. (1) There are numerous extracellular cues acting on the neuron to set the

probability of outgrowth in each direction. These cues include secreted molecules, such as

UNC-6 and EGL-20, as well as many other cues, such as extracellular matrix components (YANG

et al. 2014). For movement in the anterior and posterior directions, the neuron's response to

the cues sets an equal probability for movement in the dorsal and ventral direction. For

movement in the dorsal and ventral directions, the neuron's response to the cues sets an equal

probability for movement in the anterior and posterior direction. These cues are depicted in

the figures as purple, blue, orange, and red color gradients. (2) Besides UNC-40-mediated

outgrowth activity, cues also set the probability of nonUNC-40-mediated outgrowth activity at

each surface of the neuron. For example, there is a high probability for anteriorly directed

outgrowth in *unc-40* mutants, indicating that the cues set a high probability for anterior

nonUNC-40-mediated activity in the mutants. In the figures, a high relative probability for

UNC-40-mediated or nonUNC-40-mediated outgrowth activity at a surface is depicted as red

and green areas, respectively. The colored area is concentrated when there is a high

probability for outgrowth in one direction, and the area is more diffuse when the probability

for outgrowth in directions perpendicular to the directional bias are greater.   (3) As the

probability of outgrowth decreases in the direction of outgrowth, and the probability of

outgrowth in directions perpendicular to the directional bias become greater, the extent of

movement forward decreases. This idea is based on random walk modeling.   Representative

modeling is shown above or to the side of each diagram. 50 random walks from a point on a

line were calculated based on the given probabilities. As the probability for movement in

directions perpendicular to the directional bias become greater, the extent of movement from

the origin decreases. Similarly, we hypothesize that the forward displacement of the

membrane decreases as the probability for perpendicular outgrowth movement become

greater at the surface of the neuronal membrane.   (4) Loss of UNC-5 function causes a higher

probability of UNC-40 mediating outgrowth in directions perpendicular to the directional bias.

Loss of MIG-15 reduces the probability of UNC-40-mediated outgrowth in directions

perpendicular to the directional bias. These ideas are based on our observations of the effects

that *unc-5* and *mig-15* mutations have on HSN localization of UNC-40::GFP and the direction of

HSN outgrowth.   We further speculate that loss of MIG-15 function increases the probability of

nonUNC-40-mediated outgrowth in directions perpendicular to the directional bias. This is

based on our observation that HSN tends to be bipolar in the *mig-15* mutant (YANG *et al.* 2014),

suggesting that nonUNC-40-mediated outgrowth activity, which drives outgrowth anteriorly in

other mutants, becomes ineffectual.

PLM extension phenotype: We hypothesize that there is high probability for anteriorly

directed outgrowth from the PLM cell body because of the strong effect of a cue(s) which

inhibits outgrowth activity and is present in high concentration posterior to the cell body

(Figure 10A, position 1). Dorsal and ventral cues create an equal probability for dorsal and

ventral outgrowth, allowing the UNC-40 SOAL process to cause a high probability for anterior

outgrowth activity (Figure 10B, position 1).   At a position that is further anterior (Figure 10A,

position 2), the inhibitory effect of another cue reduces the probability for anterior outgrowth

activity, while increasing the probability of outgrowth in the dorsal and ventral directions. This

does not affect the anterior bias for outgrowth, although it may affect the rate of outgrowth

(Figure 10B, position 2). Cues at an anterior position (Figure 10A, position 3), strongly inhibit

outgrowth and cause a greater probability of outgrowth in the dorsal and ventral directions

(Figure 10B, position 3). This arrests forward outgrowth. However, in some mutants the

inhibitory effect of these cues is not as great, allowing a stronger anterior bias at position 3. As

the outgrowth passes this position (Figure 10A, position 4), a situation similar to position 1 is

reestablished (Figure 10B, position 4).

We propose that PLM termination and overextension is influenced by the degree to which UNC-

40-mediated outgrowth fluctuates at position 3 (Figure11). In wildtype, the cues cause the

direction of both UNC-40-mediated and nonUNC-40-mediated outgrowth to fluctuate, resulting

in outgrowth stalling. Loss of UNC-40 function will not alter this. Loss of MIG-15 function

increases the probability of anterior UNC-40-mediated outgrowth by decreasing the degree to

which the direction of UNC-40-mediated outgrowth fluctuates. This allows a stronger anterior

directional bias, which results in the overextension phenotype. Loss of UNC-5 suppresses the

overextension caused by loss of MIG-15 function by increasing the degree to which the

direction of UNC-40-mediated outgrowth fluctuates, reverting the state back towards

wildtype.

ALM and AVM nerve ring branching and extension phenotype: We hypothesize that at the nerve ring there are cues, including UNC-6, that are arranged along the dorsal/ventral axis. These cues can promote ALM and AVM axon outgrowth. As an extension nears the nerve ring, these cues help create a high probability for anterior outgrowth (Figure 12A and 12B, position 1). At the nerve ring, the cues create an equal probability for anterior outgrowth and outgrowth into the ring (Figure 12A and 12B, position 2). Outgrowth extends into the nerve ring because the cues of the nerve ring promote outgrowth and because the cues anterior and posterior to the ring set an equal probability of outgrowth in those directions (Figure 12A and 12B, position 3). Outgrowth can also extend anteriorly from the nerve ring if cues anterior to position 2 also promote outgrowth and cues dorsal and ventral create an equal probability of outgrowth in those directions (Figure 12A and 12B, position 4).

We propose that ALM and AVM nerve ring branching and extension at the nerve ring is influenced by the degree to which UNC-40-mediated outgrowth fluctuates at position 2 (Figure 13). In wildtype, the nerve ring cues create a higher probability for UNC-40-mediated outgrowth into the nerve ring. In UNC-40 mutants, the axons often fail to branch into the nerve ring but they still project further anterior, suggesting that the probability for anterior nonUNC-40-mediated outgrowth is high in the *unc-40* mutant. Loss of MIG-15 function does not affect the branching and extension because the higher probability for UNC-40-mediated outgrowth activity remains. In *unc-40;mig-15* mutants, there is extension in both directions because loss of MIG-15 function increases the probability of nonUNC-40-mediated outgrowth in other directions and the nonUNC-40-mediated outgrowth allows outgrowth into the nerve ring. In *unc-5;mig-15* mutants, the direction of both UNC-40-mediated and nonUNC-40-mediated outgrowth fluctuate, causing outgrowth stalling in each direction. In these mutants, there is often an absence of extension both anteriorly and into the nerve ring.

HSN extension phenotypes: We hypothesize that there is high probability for ventrally

directed outgrowth from the HSN cell body because of the strong outgrowth-promoting effect

of the UNC-6 cue, which is in a higher concentration ventral of the cell body (Figure 14A,

position 1). Anterior and posterior cues create an equal probability for anterior and posterior

outgrowth, and there is a high probability for ventral outgrowth activity. As an extension

moves ventrally it encounters higher levels of the UNC-6 cue. The probability of outgrowth

towards the UNC-6 source increases, while the probabilities of outgrowth in the anterior and

posterior decrease (Figure 14B, positions 2). However, as the level of UNC-6 saturates the

receptors the probability for outgrowth in different directions increases, causing outward

movement to stall (Figure 14B, positions 3). The distance from the source of the cue at which

this happens is a function of the concentration of the cues, and the cell's ability to interact and

respond to the cues. If the cell's response to the UNC-6 cue causes a lower probability of

ventral outgrowth than in wildtype, the probabilities of anterior and posterior outgrowth will

be greater (Figure 14B, positions 1, right column). The greater fluctuation will slow the rate of

outgrowth as extension proceeds (Figure 14B, positions 2, right column). Again, at the source

of the cue the response may become saturated and outward movement will stall (Figure 14B,

positions 3, right column). In this scenario, the rate of outgrowth decreases, while the direction

of outgrowth is the same.

We propose that HSN extension is influenced by the degree to which UNC-40-mediated

outgrowth fluctuates. In wildtype, UNC-40-mediated outgrowth creates a high probability for

the initial ventral outgrowth (Figure 15A, top). Loss of UNC-5 function causes a higher

probability of UNC-40 mediating outgrowth in the anterior and posterior directions, and

consequently a decrease in the rate of outgrowth (Figure 15A, second from top). We suggest

this underlies the observation that extensions fail to form in the L1 stage in *unc-5* mutants, but

do form over time, and in the same direction, as in wild-type animals.

We also propose that controlling the degree to which the direction of UNC-40-mediated

outgrowth fluctuates at points along the surface can regulate the number of processes that

form. We suggest that UNC-40 (DCC), UNC-5 (UNC5), and UNC-6 (netrin) are part of an activator-

inhibitor system (GIERER AND MEINHARDT 1972; MEINHARDT AND GIERER 2000). In an activator-

inhibitor system two substances act on each other. The activator stimulates its own production as

well as produces an inhibitor that can repress the production of the activator. The inhibitor is able to

diffuse more rapidly than the activator, which causes patterns of varying concentrations of activator

and inhibitor. Pattern formation through local symmetry breaking, signal amplification, and long-

range inhibition may involve chemical, mechanical, or coupled mechanochemical processes

(GOEHRING AND GRILL 2013).

We suggest an activator-inhibitor system in which UNC-40 and UNC-5 are the two substances

that interact with each other (Figure 15B). This is consistent with evidence indicating that

Netrin-1 binds simultaneously to two DCC (UNC-40) molecules or an UNC5/DCC complex (FINCI

et al. 2014). In the context of an activator-inhibitor system, UNC-5 signaling acts as an activator

and UNC-40 signaling acts as an inhibitor. The activator and inhibitor activities "diffuse" at

different rates because recruitment of UNC-40 and UNC-5 to the plasma membrane occurs at

different rates. This idea is consistent with imaging experiments of cells in culture which

suggest that netrin-1 (UNC-6) regulates the distribution of DCC (UNC-40) and UNC5B (UNC-5)

at the plasma membrane (GOPAL *et al.* 2016). In these studies, netrin-1 (UNC-6) was shown to

stimulate translocation of DCC (UNC-40) and UNC5B (UNC-5) receptors from intracellular

vesicles to the plasma membrane and, further, the transported receptors were shown to

localize at the plasma membrane (GOPAL *et al.* 2016).

We propose that signaling by UNC-6-ligated UNC-40 causes outgrowth activity and stimulates

translocation of UNC-40 to the site (Figure 15B, step 1). If the outgrowth reaches a critical

level of UNC-6 (Figure 15B, step 2), changes to the UNC-6/UNC-40/UNC-5 receptor complexes

increase UNC-5 signaling. UNC-5 signaling inhibits UNC-40 translocation to the site and

increases the relative levels of UNC-5 (short-range autocatalytic). Locally, UNC-40-mediated

outgrowth activity becomes more dispersed, resulting in a slower rate of outgrowth. As UNC-

40 translocation to the leading edge is inhibited, the relative levels of UNC-40 at flanking

surfaces increases (Figure 15B, step 3). These UNC-40 levels inhibit UNC-5 signaling (long-

range inhibition) and prevent the dispersion of UNC-40-mediated outgrowth activity, resulting

in a faster rate of outgrowth at the flanking surfaces. In the context of an activator-inhibitor

system, UNC-5 signaling functions as an activator because it stimulates its own activity by

decreasing the ratio of UNC-40 to UNC-5 at the site of UNC-5 signaling. It also stimulates

production of UNC-40 UNC-5-inhibitory activity by increasing the ratio of UNC-40 to UNC-5 at

the flanking regions. As the flanking regions move further outward due to the UNC-40-

mediated outgrowth activity, UNC-40 translocation to the surface is further stimulated by

increasing levels of the extracellular UNC-6 cue.

In this type of model, the reaction and diffusion rates of the substances influence the pattern

that emerges. In the above example, whether the flanking areas branch further depends on

whether the critical level of receptors and extracellular ligands is realized at the flanking

region. The *mig-15(rh148)* mutation and the expression of UNC-6ΔC could alter the rate by

which components of the UNC-6/UNC-40/UNC-5 receptor complexes interact or could alter the

rate of translocation of the components to the membrane, or both. In fact, cell culture

experiments suggest Netrin VI-V (UNC-6ΔC) induces DCC and UNC5B colocalization, but not

DCC recruitment (GOPAL *et al.* 2016). Moreover, mutations that suppress ectopic branching

induced by UNC-6ΔC expression affect members of second messenger systems which could

influence the rates of UNC-40 and UNC-5 interactions and trafficking (WANG AND WADSWORTH

2002).

A genetic pathway for UNC-40 asymmetric localization

We present a genetic pathway for the asymmetrical localization of UNC-40 based on the phenotype

of robust UNC-40::GFP clustering in HSN. It is worth noting that the SOAL model of axon

outgrowth does not depend on knowledge of the molecular mechanisms of outgrowth. Rather,

the SOAL model uses a statistical approach to understand how a gene influences the collective

impact of all the underlying molecular mechanisms that drives outgrowth. It can be envisioned

that the movement of an extension occurs through innumerable forces acting at the molecular

level. The effect that each molecular event has on movement is not easily observed or

measured. In fact, it may be that patterns of outgrowth can't be fully understood by only

knowing the molecular mechanisms of outgrowth. Instead, the effects that molecular events

collectively have on movement must be understood through a statistical model.

Nevertheless, a full understanding of the molecular mechanisms underlying the SOAL process is

an important long-term goal. Since we believe that UNC-40::GFP clustering is a readout of that

process, constructing genetic pathways for the clustering of UNC-40::GFP is a step toward this

goal. We wish to know how UNC-5 mediates signaling within HSN that controls the UNC-40

asymmetric localization process. However, a role for UNC-5 in HSN is paradoxical given the

widespread idea that UNC-5 mediates a repulsive response to UNC-6 and that HSN outgrowth is towards the source of UNC-6. All the same, we suggest a cell-autonomous role for UNC-5 in HSN is the most parsimonious model. First, UNC-5 is an UNC-6 receptor that can mediate neuronal responses when in complex with UNC-40 (HONG *et al.* 1999; GEISBRECHT *et al.* 2003; KRUGER *et al.* 2004; FINCI *et al.* 2014). We previously showed that UNC-40 conformational changes regulate HSN asymmetric localization in HSN (XU *et al.* 2009) and we now show that UNC-5 regulates UNC-40 asymmetric localization in HSN. It is therefore plausible that UNC-5 affects UNC-40 conformational changes that regulate UNC-40 asymmetric localization. Second, UNC-5 can alter the number to HSN outgrowths in response to UNC-6 and to the UNC-6ΔC ligand. Directional guidance by UNC-6 and UNC-6ΔC is generally normal in an *unc-5* mutant, suggesting that the ability of UNC-5 to regulate the number of outgrowth is not due to an alteration in the extracellular distribution of its UNC-6 ligand. Further, the UNC-6ΔC ligand and the *mig-15* mutation create the same outgrowth phenotype, which can be suppressed by loss of UNC-5 function, and we have shown that MIG-15 acts cell autonomously in HSN to regulate UNC-40 asymmetric localization (YANG *et al.* 2014). Further, we have shown that the UNC-5-mediated response that regulates UNC-40 asymmetric localization also depends on UNC-53 (NAV2) (KULKARNI *et al.* 2013), a cytoplasmic protein that functions cell-autonomously for cell migration and axon guidance (STRINGHAM *et al.* 2002). Together, these observations strongly suggest that UNC-5 directly regulates signaling within HSN. Third, a role for UNC-5 in the guidance of AVM and PVM axons towards UNC-6 sources has also been suggested. A synergistic interaction between *unc-5* and *egl-20* is observed; in either *unc-5* or *egl-20* mutants the ventral extension of AVM and PVM axons is only slightly impaired, whereas in the double mutants there is a much greater penetrance (LEVY-STRUMPF AND CULOTTI 2014). The expression of an *unc-5* transgene in AVM and PVM can rescue the AVM and PVM axon guidance defects of the *unc-5;egl-20* double mutant (LEVY-STRUMPF AND CULOTTI 2014). We note that for HSN, transgenic

rescue using *unc-5* constructs have not been successful and in wild-type animals UNC-5

expression in HSN has not been reported. As well, expression has not been reported in AVM,

PVM, and PLM wild-type neurons. We suspect there may be technical difficulties or that UNC-5

expression might be low in these cells.   UNC-5 is detected in PLM in *rpm-1* mutants, which is

consistent with evidence that UNC-5 activity is required for PLM overextension in these

mutants (LI *et al.* 2008).

To construct genetic pathways, we use the readout of whether UNC-40::GFP is clearly and

consistently localized to any side of the HSN neuron in different mutants (Figure 9). A

summary of the results is presented (Figure 16A). UNC-6 is required for robust asymmetric

UNC-40 localization; in the absence of UNC-6 function UNC-40 remains uniformly distributed

along the surface of the plasma membrane. The loss of both UNC-53 and UNC-5 function also

results in a uniform distribution, however loss of either one alone does not. This suggests that

UNC-53 and UNC-5 pathways act redundantly downstream of UNC-6 (Figure 16B). Moreover,

we observe there is robust asymmetric UNC-40 localization when there is a loss of UNC-6

activity in addition to the loss of UNC-53 and UNC-5.   This suggests a third pathway that is

suppressed by UNC-6 when UNC-53 and UNC-5 activity are missing.   Loss of both UNC-5 and

UNC-6 does not allow UNC-40 localization, whereas loss of both UNC-53 and UNC-6 does,

therefore UNC-53, rather than UNC-5, acts with UNC-6 to suppress the third pathway.

UNC-40 becomes localized when EGL-20 activity is lost. As well, UNC-40 becomes localized

when both EGL-20 and UNC-53 activities are lost. This is consistent with UNC-6 promoting

UNC-40 localization via the UNC-5 pathway. Loss of EGL-20 and UNC-5 prevents UNC-40

localization. In these animals, the UNC-5 pathway is absent and UNC-6 is present to blocks the

third pathway, therefore the UNC-53 pathway that leads to UNC-40 localization must require

EGL-20, as well as UNC-6.

Loss of UNC-6 activity or loss of both UNC-6 and EGL-20 activity prevents localization, whereas

loss of only EGL-20 does not. To explain this, we propose that when UNC-6 is lost, the third

pathway, which would otherwise be activated by the loss of UNC-6, remains suppressed

because EGL-20 activity promotes suppression via UNC-53 activity.   This suppression also

explains why loss of UNC-6 and UNC-5 activity does not cause localization.

Importantly, this genetic analysis indicates that netrin (UNC-6) and wnt (EGL-20) signaling are

integrated to regulate self-organizing UNC-40 asymmetric localization. An implication of this

result is that the extracellular concentrations of UNC-6 and EGL-20 could control the activation

or inhibition of UNC-40-mediated outgrowth.   This could be important for generating patterns

of outgrowth when neurons move to new locations within the animal. The picture is

complicated by the evidence that both UNC-6 and EGL-20 affect the SOAL of both UNC-40-

mediated and nonUNC-40-mediated outgrowth activity. It is possible that overlapping sets of

extracellular cues and their receptors are involved in setting the probability of outgrowth for

each activity.   SAX-3 and MADD-2 are required for UNC-40::GFP localization, but also affect

nonUNC-40-mediated outgrowth. The *egl-20;sax-3* and *unc-40;sax-3* double mutations have the

greatest effect on restricting the extent of outgrowth movement in any direction (Figures 6 and

7). Moreover, the number of HSN neurites is reduced in *unc-5* mutants, whereas the number in

unc-5;sax-3 double mutants appears normal (Figure 3B). Understanding the interdependence

of these outgrowth activities will be necessary for better understand how extracellular cues

affect the patterns of outgrowth *in vivo*.

Acknowledgments

We thank Caenorhabditis Genetics Center, J. Culotti, and C. Bargmann for strains; we thank

Martha Soto and members of the Soto laboratory for support and helpful discussions; we thank

Bhumi Patel and Leely Rezvani for comments on the manuscript. This work was supported by

grants NS033156 and NS061805 from the National Institutes of Health, National Institute of

Neurological Disorders and Stroke and grant 07-3060-SCR-E-0 from the New Jersey

Commission on Spinal Cord to WGW. This work was also supported by grant DFHS13PPCO28

from the New Jersey Commission on Cancer Research to AM.

944   Table

Table 1. Direction of Axon Formation from the HSN Cell Body							
	direction of axon protrusion						
	dorsal	ventral	anterior	posterior	multipolar		
	%	%	%	%	%	*n*	reference
wildtype	0	96±2	3±2	0	1±1	221	(Kulkarni *et al.* 2013)
unc-6(ev400)	2±2	3±2	81±2	8±2	6±1	218	(Kulkarni *et al.* 2013)
unc-40(e1430)	2±1	6±2	67±2	19±1	6±1	183	(Kulkarni *et al.* 2013)
unc-5(e53)	0	75±3	19±2	1±1	5±1	245	(Yang *et al.* 2014)
unc-53(n152)	0	67±3	22±2	5±1	6±1	238	(Kulkarni *et al.* 2013)
sax-3(ky123)	2±1	31±1	21±1	37±2	9±2	232	(Tang and Wadsworth 2014)
*sax-3(ky200)**	2±1	32±1	19±2	42±3	5±2	198	(Tang and Wadsworth 2014)
unc-5(e53);sax-3(ky200)	2±1	40±3	24±2	28±2	6±1	120	
unc-5(e53);unc-6(ev400)	4±2	5±3	59±4	22±4	9±1	201	
unc-5(e53);egl-20(n585)	3±1	28±4	22±4	35±5	11±2	114	
unc-53(n152);unc-5(e53)	0	19±1	62±2	17±1	3±1	224	(Kulkarni *et al.* 2013)
unc-53(n152);unc-6(ev400)	24±2	0	19±2	22±2	34±3	144	(Kulkarni *et al.* 2013)
unc-53(n152);sax-3(ky123)	1±1	47±3	24±2	23±5	6±3	207	(Tang and Wadsworth 2014)
unc-40(e1430);unc-5(e53)	5±1	6±1	55±2	19±2	14±1	196	(Kulkarni *et al.* 2013)
*unc-40(e1430);sax-3(ky200)**	14±3	2±1	40±2	35±3	9±4	191	(Tang and Wadsworth 2014)
sax-3(ky200); unc-6(ev400)*	8±1	8±2	49±3	20±5	14±2	211	(Tang and Wadsworth 2014)
unc-53(n152);unc-5(e53);unc-6(ev400)	23±2	0	34±2	15±2	28±2	148	(Kulkarni *et al.* 2013)
unc-53(n152);sax-3(ky200);unc-6(ev400)*	11±2	2±1	33±4	30±3	25±5	189	
egl-20(n585)	0	64±2	21±2	7±1	8±1	304	(Tang and Wadsworth 2014)
egl-20(n585); unc-6(ev400)	18±2	0	43±2	15±2	24±2	205	(Tang and Wadsworth 2014)
unc-40(e1430); egl-20(n585)	6±2	17±2	45±5	15±2	16±2	173	(Tang and Wadsworth 2014)
egl-20(n585);sax-3(ky123)	1±1	12±2	39±2	39±1	8±3	177	(Tang and Wadsworth 2014)
madd-2(tr103)	0	19±2	55±5	17±4	8±2	179	
madd-2(ky592)	0	52±2	43±2	5±1	0	95	
unc-5(e53);madd-2(tr103)	3±1	15±2	52±4	17±4	13±1	197	
madd-2(tr103);sax-3(ky123)	2	24±3	19±4	47±1	7±2	171	
unc-53(n152);madd-2(tr103)	1±1	15±2	43±2	17±1	24±4	148	
mig-15(rh326)	2±1	15±1	24±3	11±3	48±8	131	(Yang *et al.* 2014)

Numbers represent percentage value ± SEM.

*Animals grown at the *sax-3(ky200)* restrictive temperature (25°C).

References

Adler, C. E., R. D. Fetter and C. I. Bargmann, 2006 UNC-6/Netrin induces neuronal asymmetry and defines the site of axon formation. Nat Neurosci 9: 511-518.

Aiello, G. L., 2016 Neuro-Percolation as a Superposition of Random-Walks, pp. in *European Symposium on Artificial Neural Networks, Computational Intelligence and Machine Learning*. i6doc.com Bruges, Belgium.

Alexander, M., K. K. Chan, A. B. Byrne, G. Selman, T. Lee *et al.*, 2009 An UNC-40 pathway directs postsynaptic membrane extension in Caenorhabditis elegans. Development 136: 911-922.

Alexander, M., G. Selman, A. Seetharaman, K. K. Chan, S. A. D'Souza *et al.*, 2010 MADD-2, a homolog of the Opitz syndrome protein MID1, regulates guidance to the midline through UNC-40 in Caenorhabditis elegans. Developmental Cell 18: 961-972.

Asakura, T., K. Ogura and Y. Goshima, 2007 UNC-6 expression by the vulval precursor cells of Caenorhabditis elegans is required for the complex axon guidance of the HSN neurons. Dev Biol 304: 800-810.

Chan, S. S., H. Zheng, M. W. Su, R. Wilk, M. T. Killeen *et al.*, 1996 UNC-40, a C. elegans homolog of DCC (Deleted in Colorectal Cancer), is required in motile cells responding to UNC-6 netrin cues. Cell 87: 187-195.

Chang, C., C. E. Adler, M. Krause, S. G. Clark, F. B. Gertler *et al.*, 2006 MIG-10/Lamellipodin and AGE-1/PI3K Promote Axon Guidance and Outgrowth in Response to Slit and Netrin. Curr Biol 16: 854-862.

Chapman, J. O., H. Li and E. A. Lundquist, 2008 The MIG-15 NIK kinase acts cell-autonomously in neuroblast polarization and migration in C. elegans. Dev Biol 324: 245-257.

Colavita, A., and J. G. Culotti, 1998 Suppressors of ectopic UNC-5 growth cone steering identify eight genes involved in axon guidance in Caenorhabditis elegans. Dev Biol 194: 72-85.

Colon-Ramos, D. A., M. A. Margeta and K. Shen, 2007 Glia promote local synaptogenesis through UNC-6 (netrin) signaling in C. elegans. Science 318: 103-106.

Finci, L. I., N. Krüger, X. Sun, J. Zhang, M. Chegkazi *et al.*, 2014 The crystal structure of netrin-1 in complex with DCC reveals the bifunctionality of netrin-1 as a guidance cue. Neuron 83: 839-849.

Geisbrecht, B. V., K. A. Dowd, R. W. Barfield, P. A. Longo and D. J. Leahy, 2003 Netrin binds discrete subdomains of DCC and UNC5 and mediates interactions between DCC and heparin. J Biol Chem 278**:** 32561-32568.

Gierer, A., and H. Meinhardt, 1972 A theory of biological pattern formation. Kybernetik 12**:** 30-39.

Goehring, N. W., and S. W. Grill, 2013 Cell polarity: mechanochemical patterning. Trends Cell Biol 23**:** 72-80.

Gopal, A. A., B. Rappaz, V. Rouger, I. B. Martyn, P. D. Dahlberg *et al.*, 2016 Netrin-1-Regulated Distribution of UNC5B and DCC in Live Cells Revealed by TICCS. Biophys J 110**:** 623-634.

Hagedorn, E. J., J. W. Ziel, M. A. Morrissey, L. M. Linden, Z. Wang *et al.*, 2013 The netrin receptor DCC focuses invadopodia-driven basement membrane transmigration in vivo. J Cell Biol 201**:** 903-913.

Hao, J. C., C. E. Adler, L. Mebane, F. B. Gertler, C. I. Bargmann *et al.*, 2010 The tripartite motif protein MADD-2 functions with the receptor UNC-40 (DCC) in Netrin-mediated axon attraction and branching. Dev Cell 18**:** 950-960.

Hao, J. C., T. W. Yu, K. Fujisawa, J. G. Culotti, K. Gengyo-Ando *et al.*, 2001 C. elegans Slit Acts in Midline, Dorsal-Ventral, and Anterior-Posterior Guidance via the SAX-3/Robo Receptor. Neuron 32**:** 25-38.

Hedgecock, E. M., J. G. Culotti and D. H. Hall, 1990 The unc-5, unc-6, and unc-40 genes guide circumferential migrations of pioneer axons and mesodermal cells on the epidermis in C. elegans. Neuron 4**:** 61-85.

Hong, K., L. Hinck, M. Nishiyama, M. M. Poo, M. Tessier-Lavigne *et al.*, 1999 A ligand-gated association between cytoplasmic domains of UNC5 and DCC family receptors converts netrin-induced growth cone attraction to repulsion. Cell 97**:** 927-941.

Keleman, K., and B. J. Dickson, 2001 Short- and long-range repulsion by the Drosophila Unc5 netrin receptor. Neuron 32**:** 605-617.

Killeen, M., J. Tong, A. Krizus, R. Steven, I. Scott *et al.*, 2002 UNC-5 function requires phosphorylation of cytoplasmic tyrosine 482, but its UNC-40-independent functions also require a region between the ZU-5 and death domains. Dev Biol 251**:** 348-366.

Killeen, M. T., 2009 The dual role of the ligand UNC-6/Netrin in both axon guidance and synaptogenesis in C. elegans. Cell Adh Migr 3**:** 268-271.

Kruger, R. P., J. Lee, W. Li and K. L. Guan, 2004 Mapping netrin receptor binding reveals domains of Unc5 regulating its tyrosine phosphorylation. J Neurosci 24**:** 10826-10834.

Kulkarni, G., Z. Xu, A. M. Mohamed, H. Li, X. Tang *et al.*, 2013 Experimental evidence for UNC-6 (netrin) axon guidance by stochastic fluctuations of intracellular UNC-40 (DCC) outgrowth activity. Biol Open 2**:** 1300-1312.

Lai Wing Sun, K., J. P. Correia and T. E. Kennedy, 2011 Netrins: versatile extracellular cues with diverse functions. Development 138**:** 2153-2169.

Leung-Hagesteijn, C., A. M. Spence, B. D. Stern, Y. Zhou, M. W. Su *et al.*, 1992 UNC-5, a transmembrane protein with immunoglobulin and thrombospondin type 1 domains, guides cell and pioneer axon migrations in C. elegans. Cell 71**:** 289-299.

Levy-Strumpf, N., and J. G. Culotti, 2014 Netrins and Wnts function redundantly to regulate antero-posterior and dorso-ventral guidance in C. elegans. PLoS Genet 10**:** e1004381.

Li, H., G. Kulkarni and W. G. Wadsworth, 2008 RPM-1, a Caenorhabditis elegans protein that functions in presynaptic differentiation, negatively regulates axon outgrowth by controlling SAX-3/robo and UNC-5/UNC5 activity. J Neurosci 28**:** 3595-3603.

Lim, Y. S., S. Mallapur, G. Kao, X. C. Ren and W. G. Wadsworth, 1999 Netrin UNC-6 and the regulation of branching and extension of motoneuron axons from the ventral nerve cord of Caenorhabditis elegans. J Neurosci 19**:** 7048-7056.

MacNeil, L. T., W. R. Hardy, T. Pawson, J. L. Wrana and J. G. Culotti, 2009 UNC-129 regulates the balance between UNC-40 dependent and independent UNC-5 signaling pathways. Nat Neurosci 12**:** 150-155.

Maes, T., A. Barcelo and C. Buesa, 2002 Neuron navigator: a human gene family with homology to unc-53, a cell guidance gene from Caenorhabditis elegans. Genomics 80**:** 21-30.

Manser, J., C. Roonprapunt and B. Margolis, 1997 C. elegans cell migration gene mig-10 shares similarities with a family of SH2 domain proteins and acts cell nonautonomously in excretory canal development. Dev Biol 184**:** 150-164.

Manser, J., and W. B. Wood, 1990 Mutations affecting embryonic cell migrations in Caenorhabditis elegans. Dev Genet 11**:** 49-64.

McShea, M. A., K. L. Schmidt, M. L. Dubuke, C. E. Baldiga, M. E. Sullender *et al.*, 2013 Abelson interactor-1 (ABI-1) interacts with MRL adaptor protein MIG-10 and is required in guided cell migrations and process outgrowth in C. elegans. Dev Biol 373**:** 1-13.

Meinhardt, H., and A. Gierer, 2000 Pattern formation by local self-activation and lateral inhibition. Bioessays 22**:** 753-760.

Merz, D. C., H. Zheng, M. T. Killeen, A. Krizus and J. G. Culotti, 2001 Multiple signaling mechanisms of the unc-6/netrin receptors unc-5 and unc-40/dcc in vivo. Genetics 158**:** 1071-1080.

Moore, S. W., M. Tessier-Lavigne and T. E. Kennedy, 2007 Netrins and their receptors. Adv Exp Med Biol 621**:** 17-31.

Morikawa, R. K., T. Kanamori, K. Yasunaga and K. Emoto, 2011 Different levels of the Tripartite motif protein, Anomalies in sensory axon patterning (Asap), regulate distinct axonal projections of Drosophila sensory neurons. Proc Natl Acad Sci U S A 108**:** 19389-19394.

Norris, A. D., and E. A. Lundquist, 2011 UNC-6/netrin and its receptors UNC-5 and UNC-40/DCC modulate growth cone protrusion in vivo in C. elegans. Development.

Pan, C. L., J. E. Howell, S. G. Clark, M. Hilliard, S. Cordes *et al.*, 2006 Multiple Wnts and frizzled receptors regulate anteriorly directed cell and growth cone migrations in Caenorhabditis elegans. Dev Cell 10**:** 367-377.

Poinat, P., A. De Arcangelis, S. Sookhareea, X. Zhu, E. M. Hedgecock *et al.*, 2002 A conserved interaction between beta1 integrin/PAT-3 and Nck-interacting kinase/MIG-15 that mediates commissural axon navigation in C. elegans. Curr Biol 12**:** 622-631.

Poon, V. Y., M. P. Klassen and K. Shen, 2008 UNC-6/netrin and its receptor UNC-5 locally exclude presynaptic components from dendrites. Nature.

Quinn, C. C., D. S. Pfeil, E. Chen, E. L. Stovall, M. V. Harden *et al.*, 2006 UNC-6/netrin and SLT-1/slit guidance cues orient axon outgrowth mediated by MIG-10/RIAM/lamellipodin. Curr Biol 16**:** 845-853.

Quinn, C. C., D. S. Pfeil and W. G. Wadsworth, 2008 CED-10/Rac1 mediates axon guidance by regulating the asymmetric distribution of MIG-10/lamellipodin. Curr Biol 18**:** 808-813.

Ren, X. C., S. Kim, E. Fox, E. M. Hedgecock and W. G. Wadsworth, 1999 Role of netrin UNC-6 in patterning the longitudinal nerves of Caenorhabditis elegans. J Neurobiol 39**:** 107-118.

Shakir, M. A., J. S. Gill and E. A. Lundquist, 2006 Interactions of UNC-34 Enabled with Rac GTPases and the NIK kinase MIG-15 in Caenorhabditis elegans axon pathfinding and neuronal migration. Genetics 172**:** 893-913.

Song, S., Q. Ge, J. Wang, H. Chen, S. Tang *et al.*, 2011 TRIM-9 functions in the UNC-6/UNC-40 pathway to regulate ventral guidance. Journal of genetics and genomics = Yi chuan xue bao 38**:** 1-11.

Stringham, E., N. Pujol, J. Vandekerckhove and T. Bogaert, 2002 unc-53 controls longitudinal migration in C. elegans. Development 129**:** 3367-3379.

Stringham, E. G., and K. L. Schmidt, 2009 Navigating the cell: UNC-53 and the navigators, a family of cytoskeletal regulators with multiple roles in cell migration, outgrowth and trafficking. Cell adhesion & migration 3**:** 342-346.

Tang, X., and W. G. Wadsworth, 2014 SAX-3 (Robo) and UNC-40 (DCC) Regulate a Directional Bias for Axon Guidance in Response to Multiple Extracellular Cues. PLoS One 9**:** e110031.

Tessier-Lavigne, M., and C. S. Goodman, 1996 The molecular biology of axon guidance. Science 274**:** 1123-1133.

Teulière, J., C. Gally, G. Garriga, M. Labouesse and E. Georges-Labouesse, 2011 MIG-15 and ERM-1 promote growth cone directional migration in parallel to UNC-116 and WVE-1. Development 138**:** 4475-4485.

Wadsworth, W. G., H. Bhatt and E. M. Hedgecock, 1996 Neuroglia and pioneer neurons express UNC-6 to provide global and local netrin cues for guiding migrations in C. elegans. Neuron 16**:** 35-46.

Wadsworth, W. G., and E. M. Hedgecock, 1996 Hierarchical guidance cues in the developing nervous system of C. elegans. Bioessays 18**:** 355-362.

Wang, Q., and W. G. Wadsworth, 2002 The C domain of netrin UNC-6 silences calcium/calmodulin-dependent protein kinase- and diacylglycerol-dependent axon branching in Caenorhabditis elegans. J Neurosci 22**:** 2274-2282.

Wang, Z., Y. Hou, X. Guo, M. van der Voet, M. Boxem *et al.*, 2013 The EBAX-type Cullin-RING E3 ligase and Hsp90 guard the protein quality of the SAX-3/Robo receptor in developing neurons. Neuron 79**:** 903-916.

Wang, Z., L. M. Linden, K. M. Naegeli, J. W. Ziel, Q. Chi *et al.*, 2014 UNC-6 (netrin) stabilizes oscillatory clustering of the UNC-40 (DCC) receptor to orient polarity. J Cell Biol 206**:** 619-633.

Xu, Y., H. Taru, Y. Jin and C. C. Quinn, 2015 SYD-1C, UNC-40 (DCC) and SAX-3 (Robo) function interdependently to promote axon guidance by regulating the MIG-2 GTPase. PLoS Genet 11**:** e1005185.

Xu, Z., H. Li and W. G. Wadsworth, 2009 The roles of multiple UNC-40 (DCC) receptor-mediated
signals in determining neuronal asymmetry induced by the UNC-6 (netrin) ligand. Genetics 183**:**
941-949.

Yang, Y., W. S. Lee, X. Tang and W. G. Wadsworth, 2014 Extracellular Matrix Regulates UNC-6
(Netrin) Axon Guidance by Controlling the Direction of Intracellular UNC-40 (DCC) Outgrowth
Activity. PLoS One 9**:** e97258.

Yoshimura, S., J. I. Murray, Y. Lu, R. H. Waterston and S. Shaham, 2008 mls-2 and vab-3 Control
glia development, hlh-17/Olig expression and glia-dependent neurite extension in C. elegans.
Development 135**:** 2263-2275.

Zallen, J. A., B. A. Yi and C. I. Bargmann, 1998 The conserved immunoglobulin superfamily
member SAX-3/Robo directs multiple aspects of axon guidance in C. elegans. Cell 92**:** 217-227.

Ziel, J. W., E. J. Hagedorn, A. Audhya and D. R. Sherwood, 2009 UNC-6 (netrin) orients the
invasive membrane of the anchor cell in C. elegans. Nat Cell Biol 11**:** 183-189.

1109

Figure 1. Model for UNC-40 self-organizing polarization and random walk movement.

(**A**) Along the plasma membrane, complexes comprising UNC-40 and UNC-5 interact with extracellular cues. A self-organizing process is triggered. In wild-type animals (top) there is a high probability that complexes associated with specific levels of the extracellular cues will mediate outgrowth activity. Compared to wildtype, in animals with loss of UNC-5 function (bottom) the probability of these same complexes mediating outgrowth activity is lower and the probability that the other complexes instead will mediate outgrowth is higher. Direction is labeled left, down, and right, in reference to the orientation of the figure and to emphasize that the model is not referring to any specific cues or their sources in the animal. (**B**) Two sets of tracks are shown to compare random walk models derived from representative wildtype and mutant experimental results. Compared to wildtype, the mutant has a lower probability of outgrowth in one specific direction (down) and a higher probability of outgrowth in other directions (left and right). Using the probability distributions shown below the tracks, 50 simulated random walks of 500 steps were plotted from an origin. This comparison emphasizes that even though the direction of movement is the same, there is a difference in the average displacement. We propose this property of movement is manifested during neuronal extension in wildtype and mutant animals as a difference in the rate of outgrowth. In statistical physics, net movement through the action of random motion is diffusion. Here, each track models the trajectory that mass of the plasma membrane could move during extension. The probability density function of the position of the mass as a function of space and time is described by an advection-diffusion equation. In summary, we suggest that the advection-diffusion model can be used to describe the process by which mass at the leading edge of an extension is transported and that our results describe how genes regulate this transport.

A

B

Example:

	P(right)	=	0.1	0.33	=	P(right')
	P(left)	=	0.1	0.33	=	P(left')
	P(down)	=	0.8	0.33	=	P(down')

1134

1135

Figure 2. Model for outgrowth movement based on the SOAL model. Schematic diagrams

of the outgrowth of a neuron (gray) through an environment of multiple extracellular cues.

These cues may be molecules presented at the surfaces of surrounding cells and extracellular

matrix. The extracellular cues are represented here as color gradients of purple, blue, and

orange. Outgrowth activity is mediated at the surface of the neuron through receptor activity.

UNC-40-mediated activity is depicted in red, and nonUNC-40-mediated in green. The relative

probability of activity along the surface is represented by the extent to which the colored areas

cover the neuron's surface. Increasing the degree to which the direction of activity fluctuates

over time decreases the extent of outward movement. Random walk models depicting this

property are shown above each diagram. See text for details. **(A)** Depicted is the movement of

a neuronal extension towards a source of a cue(s) (orange). At time 1 (top) the probability of

outgrowth towards the source is high, however at time 2 (bottom) the probability of outgrowth

in other directions increases as more receptors away from the leading edge are stimulated due

to the higher concentration of the cue. This increases the degree to which the direction of

activity fluctuates over time and decreases the extent of outward movement. **(B)** Depicted is

the movement of a neuronal extension in wildtype (top) and an *unc-5* mutant (bottom). Loss of

UNC-5 affects the UNC-40 SOAL process and causes more receptor activity away from the

leading edge, thus decreasing the probability of outgrowth towards the source and increasing

the probability of outgrowth in other directions. This increases the degree to which the

direction of activity fluctuates over time and decreases the extent of outward movement.

Under the same conditions, the rate of outward movement is reduced in *unc-5* mutants as

compared to wildtype.

1158

1159

1160

1161

1162

1163

 Figure 3. UNC-5 regulates the patterning of outgrowth extensions from HSN. (**A**)

Photomicrographs of HSN at the L1, L2, and adult stages in wildtype and *unc-5(e53)* mutants.

In L1 and L2 animals neurite extensions (arrows) are often observed in wild-type animals but

are more rare in *unc-5* mutants. The short ventral migration of the cell body that occurs in

wild-type animal sometimes fails in *unc-5* mutants, leaving the cell body farther from the PLM

axon (arrowhead) with a single longer ventral extension. The position of the cell body remains

dorsal. Scale bar: 10 μm. (**B**) The percentage of HSN neuron with 0, 1, or more than 1 neurite

extension at the L1 stage. In *unc-5* mutants nearly half of the neurons do not extend a process.

Error bars indicated the standard error mean; n values are indicated above each column.

Significant differences (two-tailed t-test), *P<0.001. (**C**) The percentage of HSN neurons with a

single long extension at the L2 stage. Several *unc-5* alleles were tested as described in the text.

In mutants with loss-of-function there is more often a single extension from the cell body and

the cell body is dorsally mispositioned.   (**D**) Photomicrographs of HSN at the early L4, late L4,

and adult stages in wildtype and in animals expressing UNC-6ΔC. The expression of UNC-6ΔC

induces multiple processes, most often two major extensions, that are guided ventrally. (**E**) The

percentage of HSN neurons with a cell body mispositioned dorsally at the L2 stage. In loss-of-

function mutants the cell body often fails to undertake a short ventral migration during the L2

stage. The migration is not delayed, but rather it remains dorsal. (**F**) The percentage of HSN

neurons with multiple ventral extensions at the L4 stage. The additional processes induced by

UNC-6ΔC can be suppressed by *unc-5* and *mig-10* mutations. Additional processes induced by

mig-15(rh148) can also be suppressed by the *unc-5* mutation. (**G**) Photomicrographs of HSN at

adult stages in a *mig-15* mutant. Similar to UNC-6ΔC expression, *mig-15* mutations can also

cause additional processes that are guided ventrally (Yang *et al.* 2014).

1188

Figure 4 UNC-5 regulates the patterning of extension from ALM, AVM, and PLM. (**A**) Photomicrographs of the ALM, AVM, and PLM neurons at the L4 stage in wild-type animals and *mig-15* mutants. In wildtype (top) a single PLM axon travels anteriorly from the posterior cell body (not shown). Near the vulva (arrow) the axon branches; one branch extends to the ventral nerve chord and another extends anteriorly. The anterior extension terminates before reaching the area of the ALM cell body. In *mig-15* mutants the PLM can extend anteriorly past the ALM cell body (bottom). (**B**) The percentage of PLM neurons where the PLM neuron extend anteriorly past the ALM cell body. The anterior extension often over-extends in *mig-15* mutants. Loss of *unc-5* or *unc-40* function can suppress this phenotype. (**C**) Photomicrographs of the ALM and AVM neurons at the L4 stage in wild-type animals and mutants showing different patterns of outgrowth extension. In wildtype (top) a single axon travels anteriorly to the nerve ring (arrowheads). At the nerve ring the axon branches; one branch extends further anteriorly and the other extends into the nerve ring. In mutants, one or both axons may only extend anteriorly and will not extend into the nerve ring (second from top). Or one or both axons will only extend into the nerve ring and will not extend anteriorly (third from top). Or one or both axons will fail to extend into either the nerve ring or anteriorly (bottom). Scale bar: 20 µm. (**D**) The percentage of AVM neurons where the AVM neuron failed to extend into the nerve ring. The neuron often fails to extend in the *unc-40* and *mig-15;unc-5* mutants, whereas it does extend in the *mig-15, unc-5,* and *mig-15;unc-40* mutants. Error bars indicated the standard error mean; n values are indicated above each column. Significant differences (two-tailed t-test), *P<0.001. (**E**) The percentage of AVM neurons where the AVM neuron failed to extend anteriorly, past the nerve ring. The neuron often fails to extend anteriorly in the *mig-15;unc-5* mutants, whereas it does extend in the *mig-15, unc-5, unc-40,* and *unc-40;mig-15* mutants. There is a significant difference between the *unc-40* and *unc-40;mig-15* mutants.

1217
1218

Figure 5 Assay to measure the effects a mutation has on movement. (**A**) The direction of outgrowth extension from the HSN cell body can vary and whether the axon developed in the dorsal, anterior, posterior, or ventral direction in L4 stage animals is scored (left panel). This creates a probability distribution in which the direction (*X*) is a random variable (center panel). A simple random walk is generated by using the same probability distribution for a succession of steps with an equal time interval (right panel). (**B**) For wildtype and two mutants, 50 simulated random walks of 500 steps were plotted from an origin (0,0). The results graphically indicate the directional bias for movement. For random walk movement created in mutant A (red, results from *unc-5(e53)*), the directional bias is shifted anteriorly (left) relative to wildtype. The results also graphically show the displacement of movement. For random walk movement created in mutant B (blue, results from *egl-20(n585);sax-3(ky123)*), the average of the final position (displacement) from the origin is a much shorter distance than wildtype. (**C**) Plots of the normal distribution of the final position along the x axis of the random walk tracks shown in B. The mean position for each is set at 0. The plots graphically illustrate how random walks constructed from the probability distribution for the direction of outgrowth extensions can reveal a diffusion process.

A direction (X)   probability distribution   simple random walk

dorsal (d)

anterior (a)

posterior (p)

ventral (v)

$P(X = d_1)$

$P(X = a_1)$ $P(X = p_1)$

$P(X = v_1)$

$\Delta t = 1$

$\Delta t = 1$

$\Delta t = 1$

stochastic sequence $\{S_n\}$, with $S_0 = 0$,

$$S_n = \sum_{k=1}^{n} X_k$$

B

−500

−200   0   100

500 steps

mutant B

mutant A

wild type

C

0.12

0.1

0.08

0.06

0.04

0.02

concentration(x,t)

−100   0   80

X

wild type

displacement of outgrowth

+++

++

+

mutant A

mutant B

Figure 6. Mutations have different effects on movement. Examples of random walk

analyses using the direction of axon development from the HSN neuron in different mutants

(Table 1). The graphs were created as described in the figure legend of Figure 3. For each

panel, plots are shown for the normal distribution of the final position along the x axis for the

random walk tracks plotted in the inserts. The inserts depict the random walk movement that

would be produced by the probability distribution for the direction of outgrowth in the mutant.

Plots derived from the same data are colored alike. Each panel depicts the analyses of four

different mutants and wildtype. Three different distribution patterns are observed: (1) the

wild-type distribution, which has the distribution curve with the highest peak; (2) the *unc-5,*

egl-20, unc-53, and *unc-6* (not shown) distribution, which is flatter than the wild-type curve; (3)

the *madd-2, sax-3, mig-15,* and double combinations, which have the flattest distribution curve.

1249

Figure 7. Mutations alter the spatial extent of movement. (**A**) Plotted are the mean

squared displacement (MSD) curves as a function of time interval (d*t*). The values are in

arbitrary units, since the time scale was arbitrarily set at 1. The curves show the extent that

different mutations can alter the MSD relative to wildtype and the MSD caused by an unbiased

random walk. For each time interval, mean and s.e.m. are plotted. (**B**) From the slope of MSD

curves a coefficient can be derived that gives the relative rate of diffusion. Colored bars

correspond to the like-colored curves given in panel A. The coefficients for *unc-5, egl-20, unc-*

53, and *unc-6* form a class that is distinct from that derived from wildtype and from the double

mutants.

Figure 8. Models for the relationship between UNC-40-mediated outgrowth activity and UNC-40 receptor clustering. (A) In this model the self-organizing UNC-40 SOAL process causes observable UNC-40 receptor clustering. Receptor clusters become stable and exit the SOAL process, at which point they recruit the outgrowth machinery. Whereas the direction of asymmetric receptor localization is determined stochastically, the direction of outgrowth is determined by the site of stabilization. **(B)** In this model the self-organizing UNC-40 SOAL process is coupled to the outgrowth machinery. The direction of both asymmetric receptor localization and outgrowth activity are stochastically determined. Observable receptor clustering may arise over time and these sites may become paramount for outgrowth movement, but cluster formation is not a prerequisite for outgrowth activity. Conceptually, in both models the site of outgrowth activity is determined by where UNC-40 receptor complexes recruit the outgrowth machinery, however the second model postulates situations where there are innumerable fluctuating sites that create outgrowth movement in various directions. Because the effect at each instance of time of every outgrowth event is immeasurable, outgrowth in this model is considered as a continuous stochastic process that evolves over time.

A

B

1286

1287

Figure 9. Mutations affect asymmetric intracellular UNC-40::GFP localization. (**A**) Graph indicating the dorsal-ventral localization of UNC-40::GFP in HSN. The graph shows the average ratio of dorsal-to-ventral intensity from linescan intensity plots of the UNC-40::GFP signal around the periphery of the HSN cell. UNC-40::GFP is ventrally localized in wildtype, but the ratio is different in the mutants. Error bars represent standard error of mean. Below is a graphic representation of the possible UNC-40 localization patterns when the intensity ratio is ≥ 1 or is <1. (**B**) Graph indicating the anterior-posterior localization of UNC-40::GFP. To determine orientation, line-scan intensity plots of the UNC-40::GFP signal across the dorsal periphery of the HSN cell were taken, the dorsal surface was geometrically divided into three equal segments, and the total intensity of each was recorded. The percent intensity was calculated for each segment and ANOVA was used to determine if there is a significant difference between the three segments (see Material and Methods). Whereas in the *unc-5* and *egl-20* mutants there is a bias for anterior or posterior localization, there is a uniform distribution in *unc-5;egl-20* double mutants. Uniform distribution is also observed in strong loss-of-function *sax-3* and *madd-2* mutants. (*) Animals grown at the *sax-3(ky200)* restrictive temperature (25°C). Below is a graphic representation of the possible UNC-40 localization patterns.

A. Dorsal-ventral orientation of UNC-40 asymmetric localization

B. Anterior-posterior orientation of UNC-40 asymmetric localization

1306

1307

Figure 10. Model for the outgrowth movement of PLM. Schematic diagrams of the anteriorly directed outgrowth of PLM. The elements of the schematics are described in Figure 2. **(A)** At each of the four positions an extension encounters different levels of extracellular cues. At position 3, a cue(s) is present that inhibits outgrowth activity. PLM extension terminates in this area in wildtype, but in some mutants there is an overextension to position 4. **(B)** Depicted are the effects that extracellular cues have on the relative probabilities of UNC-40-mediated and nonUNC-40-mediated outgrowth activity at the different positions. At position 3 in wildtype the degree to which the direction of outgrowth activity fluctuates increases and the extent of outgrowth movement decreases, stalling outgrowth. Because the SOAL process is perturbed in the mutants, at position 3 the probability of outgrowth activity perpendicular to the direction bias does not increase as much as in wildtype and the probability of anterior outgrowth activity remains greater. This allows a stronger directional bias to persist, allowing forward extension. To illustrate the concept, random walk models are shown which were constructed using the indicated values for outgrowth in the anterior, dorsal, and ventral directions. See text for details.

A

extracellular cues set a lower probability for outgrowth

4 ← 3 ← 2 ← 1 ← neuron

extracellular cues set equal probability for dorsally and ventrally directed outgrowth

B

axon
termination

axon
overextension

1325

1326

1327 **Figure 11. Model for the effects that mutations have on the outgrowth movement of PLM.**

1328 Schematic diagrams of PLM outgrowth at position 3 of Figure 10 in wildtype and mutants. The

1329 elements of the schematics are described in Figure 2. PLM outgrowth terminates in wildtype,

1330 *unc-40*, and *unc-5;mig-15* mutants, but overextends in *mig-15* mutants. See text for details.

1331

1332

figure 10, position 3

genotype	phenotype	random walk modeling
wildtype		
unc-40(-)		
mig-15(rh148)		
unc-5(-); mig-15(rh148)		

1333

1334

Figure 12. Model for the outgrowth movement of AVM at the nerve ring. Schematic

diagrams of the outgrowth of AVM at the nerve ring. The elements of the schematics are

described in Figure 2. **(A)** At each of the four positions an extension encounters different

levels of extracellular cues. At position 2, a cue(s) at the nerve ring promotes UNC-40-mediated

outgrowth activity. From this position in wildtype, AVM extends further anteriorly and also

dorsally along the nerve ring. However, in some mutants either the anterior or dorsal

extension is absent. In other mutants, extension in both directions is absent. **(B)** Depicted are

the effects that extracellular cues have on the relative probabilities of UNC-40-mediated and

nonUNC-40-mediated outgrowth activity at the different positions. At position 2 in wildtype

the extracellular cues create a weak bias for outgrowth in both the anterior and dorsal

directions. Further dorsally at position 3, cues create an equal probability for outgrowth in the

anterior and posterior directions, allowing the nerve ring cue that promotes UNC-40-mediated

outgrowth to maintain a dorsal directional bias. Further anteriorly at position 4, cues create an

equal probability for outgrowth in the dorsal and ventral directions, allowing anterior cues that

promote outgrowth to create an anterior directional bias. To illustrate the concept, random

walk models are shown which were constructed using the indicated values for outgrowth in the

indicated directions. See text for details.

A

anterior extracellular
cues allow a higher
probability for
anteriorly directed
outgrowth

4 ← 2 ← 1 ←

neuron

nerve ring

UNC-6 and other extracellular cues set equal
probability for dorsally and ventrally directed outgrowth

B

.4   .3

.3

.4

.3   .3

.33

.33   .33

.33   .33

.33

.6   .2

.2

1357

1358

Figure 13. Model for the effects that mutations have on the outgrowth movement of AVM at the nerve ring. Schematic diagrams of AVM outgrowth at position 2 of Figure 12 in wildtype and mutants. The elements of the schematics are described in Figure 2. AVM outgrowth extends both anteriorly and dorsally from this position in wildtype, *mig-15,* and *unc-40;mig-15* mutants. Dorsal extensions are often absent in *unc-40* mutants, whereas both anterior and dorsal extensions are absent in *unc-5;mig-15* mutants. See text for details.

figure 12, position 2

genotype   phenotype   random walk modeling

wildtype

unc-40(-)

mig-15(rh148)

unc-40(-);
mig-15(rh148)

unc-5(-);
mig-15(rh148)

1370

1371

1372

Figure 14. Model for the outgrowth movement of HSN. Schematic diagrams of the ventrally directed outgrowth of HSN. The elements of the schematics are described in Figure 2. **(A)** At each of the three positions an extension encounters different levels of extracellular cues. At position 3, a cue(s) is present that promotes outgrowth activity. At the time when extension is first observed in wildtype, extension is absent in some mutants. Nevertheless, ventral extension will be complete in the mutants at about the same time as in wildtype. **(B)** Depicted are the effects that extracellular cues have on the relative probabilities of UNC-40-mediated and nonUNC-40-mediated outgrowth activity at the different positions. Compared to wildtype, at position 1 in mutants with delayed protrusion the degree to which the direction of outgrowth activity fluctuates is greater and the extent of outgrowth movement is less, resulting in a reduced rate of outgrowth. A weak ventral directional bias develops in the mutant that allows initial ventral extension and by position 2 this ventral bias is stronger. At position 3 in both wildtype and mutant, the degree to which the direction of outgrowth activity fluctuates increases and the extent of outgrowth movement decreases. To illustrate the concept, random walk models are shown which were constructed using the indicated values for outgrowth in the anterior, posterior, and ventral directions. See text for details.

A

extracellular cues set an equal probability for anteriorly and posteriorly directed outgrowth

neuron

1
2
3

UNC-6 and other extracellular cues set higher probability for ventrally directed outgrowth

B   wildtype   delayed protrusion

1   1
.3   .3   .4   .4
                                    .2
    .4

2   2
.2   .2   .3   .3
    .6   .4

3   3
.33   .33   .33   .33
    .33   .33

1392

Figure 15. Model for the effects that mutations have on the outgrowth movement of HSN.

(A) Schematic diagrams of HSN outgrowth at position 1 of Figure 14 in wildtype and mutants.

The elements of the schematics are described in Figure 2. HSN outgrowth is delayed in *unc-5*

mutants and can develop multiple processes in *mig-15* mutants. The multiple process

phenotype is suppressed in *unc-5;mig-15* mutants. See text for details. **(B)** Schematic diagrams

of the development of multiple process that maintain the same directional bias. The process is

shown in three steps. At the first step, UNC-40 receptors (blue rectangles) are trafficked to the

membrane as part of the UNC-40 SOAL process (blue arrows). At the second step, UNC-5

signaling alters the trafficking, increasing the rates anterior (orange arrows) and posterior (red

arrows). By the third step, the rate of outgrowth at the flanking regions is greater. We propose

that UNC-40, UNC-5, and UNC-6 can be considered as components of an activator-inhibitor

system which allows outgrowth patterns to self-organize. See text for details.

1405

1406

1407

1408

1409

1410

1411

1412

Figure 16. Genetic pathways for self-organizing UNC-40 asymmetric localization. (**A**)

Table summarizing the results of experiments previously reported and described in Figure 9 of

this paper. (**B**) The genetic data support a model whereby the UNC-6 and EGL-20 extracellular

cues regulate at least three pathways leading to robust asymmetric UNC-40 localization.

Robust asymmetric UNC-40 localization refers to the ability to observe UNC-40::GFP clustering

at the surface of the neuron. Arrows represent activation; bars represent repression. See text

for the logic used to construct the pathways.

1420

A

loss-of-function			robust asymmetric UNC-40 localization	reference
unc-6			no	(Kulkarni et al., 2013)
unc-5			yes	(Kulkarni et al., 2013)
unc-5	unc-6		no	(Kulkarni et al., 2013)
unc-5	egl-20		no	
unc-5	sax-3		no	
unc-5	unc-53		no	(Kulkarni et al., 2013)
unc-5	unc-53	unc-6	yes	(Kulkarni et al., 2013)
unc-5	unc-53	sax-3	no	
unc-5	madd-2		no	
unc-6	unc-53		yes	(Kulkarni et al., 2013)
unc-6	sax-3		no	(Tang and Wadsworth, 2014)
unc-6	unc-53	sax-3	no	(Tang and Wadsworth, 2014)
egl-20			yes	(Kulkarni et al., 2013)
egl-20	unc-53		yes	(Kulkarni et al., 2013)
egl-20	unc-6		no	(Kulkarni et al., 2013)
egl-20	sax-3		no	(Tang and Wadsworth, 2014)
sax-3			no	(Tang and Wadsworth, 2014)
sax-3	unc-53		no	(Tang and Wadsworth, 2014)
unc-53			yes	(Kulkarni et al., 2013)
madd-2			no	
mig-15			yes	(Yang et al., 2014)

1421

B

81