1 **An assessment of phylogenetic tools for analyzing the interplay between interspecific**
2 **interactions and phenotypic evolution**
3

4 Drury, J.P.[1,2*], Grether, G.F.[2], Garland, Jr., T.[3], & Morlon, H.[1]
5
6 [1]Institut de Biologie, École Normale Supérieure, CNRS UMR 8197, 75005 Paris, France
7 [2]Department of Ecology & Evolutionary Biology, University of California, 621 Charles E.
8 Young Dr. S, Los Angeles, CA 90095-1606, USA
9 [3]Department of Biology, University of California, Riverside, CA 92521, USA
10
11 *Corresponding author:*
12 druryj@ucla.edu
13 (424) 362-6230
14 621 Charles E. Young Dr. S
15 Los Angeles, CA 90095-1606, USA
16

17 *Running head*: INTERSPECIFIC INTERACTIONS & TRAIT EVOLUTION

Abstract

Much ecological and evolutionary theory predicts that interspecific interactions often

drive phenotypic diversification and that species phenotypes in turn influence species

interactions. Several phylogenetic comparative methods have been developed to assess the

importance of such processes in nature; however, the statistical properties of these methods have

gone largely untested. Focusing mainly on scenarios of competition between closely-related

species, we assess the performance of available comparative approaches for analyzing the

interplay between interspecific interactions and species phenotypes. We find that currently used

statistical methods largely fail to detect the impact of interspecific interactions on trait evolution,

that sister taxa analyses often erroneously detect character displacement where it does not exist,

and that recently developed process-based models have more satisfactory statistical properties. In

weighing the strengths and weaknesses of different approaches, we hope to provide a clear guide

for empiricists testing hypotheses about the reciprocal effect of interspecific interactions and

species phenotypes and to inspire further development of process-based models.

Keywords: character displacement, competition, interspecific interactions, phylogenetic

comparative methods, trait evolution

Interactions between species are a fundamental aspect of life on earth, and understanding

the evolutionary and ecological consequences of such interactions are a central goal of many

classical theoretical frameworks in ecology and evolutionary biology. Identifying both the

predictors of interspecific interactions and the consequences of such interactions for

diversification and coexistence is thus an important contemporary research area, with strong

implications for conservation biology.

Several phylogenetic comparative methods have been deployed with the goal of

elucidating how interspecific interactions drive (or are driven by) character evolution, but the

reliability and efficacy of these methods remain largely untested. Here we focus on methods used

to study interactions between closely related species (e.g., members of the same family) that

arise from similarity in morphology, signaling traits or habitat (Brown and Wilson 1956;

Schluter 2000; Pfennig and Pfennig 2009), rather than on community-wide interactions and

interaction networks (Webb et al. 2002; Rezende et al. 2007; Cavender-Bares et al. 2009;

Cadotte et al. 2013).

Classical character displacement theory (Brown and Wilson 1956; Grether et al. 2009;

Pfennig and Pfennig 2009) predicts that, where heterospecifics compete, selection should favor

divergence in the traits responsible for competition, until lineages in sympatry no longer compete

intensely. In a seminal example, selection resulting from exploitative competition between

medium and large ground finches (*Geospiza fortis* & *G. magnirostris*) has driven bill size

divergence on Daphne Major in the Galápagos (Grant and Grant 2006). Investigators who

conduct comparative studies of divergent character displacement often test for a relationship

between biogeographic overlap and trait dissimilarity, predicting that coexisting species will be

more phenotypically divergent than non-coexisting ones. Recent studies on *Bicyclus* butterflies

and *Euglossa* bees, for example, show that male chemical cues are more distinct between

sympatric species than allopatric species, suggesting that reproductive character displacement

has driven signal divergence in these taxa (Bacquet et al. 2015; Weber et al. 2016).

Interspecific interactions can also lead to convergent, rather than divergent, character

displacement (Cody 1969, 1973; Grant 1972; Grether et al. 2013). Agonistic character

displacement theory (Grether et al. 2013) predicts convergence in traits mediating interspecific

aggression when species compete strongly for the same resources. In other words, between-

species similarity in resource use may make interspecific territoriality adaptive, resulting in

subsequent convergence in signaling traits involved in mediating territorial interactions (e.g.,

song in ovenbirds, Tobias *et al.* 2014). Therefore, tests of convergent character displacement

typically test the prediction that sympatric lineages are more phenotypically similar than

allopatric ones. Because sympatric similarity can also result from convergence to local

conditions (e.g., habitat, climate), it is important for empiricists to account for abiotic factors in

tests of character convergence.

In some instances, rather than identifying the effect of species interactions on trait

evolution, empiricists aim to identify traits that mediate particular pairwise interactions, such as

hybridization or interspecific aggression. In this case, investigators test for a relationship

between the measured interactions and trait similarity. Recent studies on New World warblers

(Parulidae), for example, show that hybridization occurs more often between species with similar

songs and that interspecific territoriality occurs more often between species that share similar

plumage and territorial song phenotypes (Willis et al. 2014; Losin et al. 2016).

Although the examples presented here largely represent scenarios where interactions

between species are competitive, empiricists apply methods discussed here to other non-

competitive interactions as well (e.g., predicting links in plant/pollinator networks, identifying

Müllerian mimicry rings: Elias et al. 2008; Eklöf and Stouffer 2016). Regardless of the

biological question, a particularity of comparative tests aimed at understanding the interplay

between interspecific interactions and species phenotypes is that they largely involve testing

correlations between pairwise data (e.g. range overlap, phenotypic similarity, frequency of

hybridization). In contrast, most phylogenetic comparative methods have been developed and

tested on tip data (e.g. range size, morphological trait values), and the statistical properties of

methods adapted to handle pairwise data (Box 1) have gone untested (but see Harmon & Glor

2010). Furthermore, species interactions are inherently affected by the biogeographic history of

dispersal and speciation in an evolving clade and the resulting patterns of range overlap. Patterns

of trait dissimilarity between sympatric lineages—the classic test of character displacement—

may actually be the null expectation if allopatric speciation is the norm, because then sympatric

species pairs will tend to share more distant common ancestors than allopatric species pairs do

(Weir and Price 2011; Tobias et al. 2014).

Here, we apply the main phylogenetic comparative methods that investigators use to test

hypotheses about interactions between closely related lineages and phenotypes (Box 1, Fig. 1) to

datasets simulated under different evolutionary histories of speciation, dispersal, species

interactions, and trait evolution. We then compare the efficacy of these methods, discuss the

relative merits of each, and outline directions for future research.

115
116

117 *Figure 1*. Schematic examples of the processes examined in our simulation study. A. Phylogeny

118 along which the trait evolves. B. A trait evolving via divergent character displacement, C. A trait

119 evolving via convergent character displacement, and D. A species interaction that exists at

120 present due to pairwise trait similarity. For simulation details, see the main text and

121 *Supplementary Methods*.

METHODS

We compared the performance of different phylogenetic comparative methods by measuring their statistical power (e.g., probability of detecting divergence when divergence is simulated) and Type I error rate (e.g., probability of detecting an effect of species interactions when such an effect is not simulated) across three scenarios.

Phylogeny and Range Simulations

We jointly simulated trees {# spp. = 20, 50, 100, 150, 200, 250} and biogeographies under the dispersal-extinction-cladogenesis model of biogeographical evolution (i.e., DEC+J, with the inclusion of founder event speciation) in BioGeoBEARS (Ree and Smith 2008; Matzke 2014). Briefly, the DEC+J model is a model of range evolution in which species ranges change along the branches of a phylogeny as a function of dispersal and local extinction and are inherited by daughter taxa at speciation according to several possible cladogenetic scenarios (see more details in Supplementary Methods). For each tree, we started with a single ancestral species occupying one of ten equidistant regions, and simulated trees with constant rates of speciation and local extinction. We considered different biogeographic scenarios by varying the rate of dispersal events between ranges ("high" and "low" dispersal; see details in Supplementary Methods) and the probability that speciation events occur in sympatry versus allopatry ("high" and "low" sympatric speciation; Supplementary Methods). Each of these simulations resulted in a phylogeny (the tree of extant species) and its associated biogeography (the set of regions in which each lineage occurred throughout the history of the clade). Lineages were identified as

146   sympatric if they co-occurred in at least one of the ten geographic regions, and allopatric if they

147   did not co-occur in any.

148   We simulated four biogeographic scenarios (combinations of low or high dispersal and

low or high sympatric speciation) for each tree size. The resulting biogeographies span scenarios

where sympatric speciation is common and dispersal is low (e.g., lizards on islands) to scenarios

where allopatric speciation is the main mode of speciation and dispersal between regions is high

(e.g., birds on continents). These parameter combinations produced a range of realistic

proportions of sister taxa that are sympatric (Fig. S1A) and a range of realistic differences in age

between sympatric and allopatric sister taxa (Pigot and Tobias 2014; Fig. S1B). In defining

sympatry as any overlap, the mean magnitude of range overlap fell between 33-42% across all

tree sizes and simulation parameters (Fig. S1C,D), which falls well within the range of overlap of

sympatric taxa defined under commonly used minimum threshold values applied to continuous

indices of range overlap (e.g. Pigot and Tobias 2014; Tobias et al. 2014).

159   For each combination of tree sizes and DEC parameter combinations ($n = 24$), we

160   performed 100 simulations, resulting in a bank of 2,400 trees with associated biogeographies.

162   *Character Displacement*

163   *The model.*—To simulate both divergent and convergent character displacement, we

simulated a continuous trait z under a model in which trait values of sympatric species in an

evolving clade are repelled from (or drawn toward) one another. In divergent character

displacement, trait divergence is driven by pairwise similarity in that same trait z; in convergent

character displacement however, convergence in trait z (e.g. a signaling trait) is driven by

pairwise similarity in another trait y (e.g. a resource use trait). To create a generic model of

character displacement, we thus modified the matching competition model (Nuismer & Harmon

2015; Drury *et al*. 2016) by describing the mean value for trait *z* in lineage *i* after an

infinitesimally small time step *dt* by:

$$z_i(t + dt) = z_i(t) + \psi(\theta - z_i(t))dt + m\left[\sum_{j \neq i}^{n} \mathbf{A}_{i,j} \times sign\left(z_i(t) - z_j(t)\right) \times e^{-\alpha\left(y_j(t) - y_i(t)\right)^2}\right]dt + \delta$$

(Eq. 1)

where $y = z$ in the case of divergent character displacement and $y \neq z$ in the case of divergent

character displacement, $\psi(\theta - z_i(t))$ describes attraction to a single stationary peak (i.e., the

Ornstein-Uhlenbeck [OU] process, Felsenstein 1988; Garland et al. 1993; Hansen and Martins

1996), *n* is the number of species, δ is a random variable with mean 0 and variance $= \sigma^2 dt$ (the

Brownian motion [BM] rate parameter, describing the stochastic component of trait evolution),

and $\mathbf{A}$ is a piecewise-constant matrix representing biogeographical overlap such that $\mathbf{A}_{i,j}$ equals 1

if species *i* and *j* are sympatric at time *t*, and 0 otherwise. The "sign" portion determines the

relative position of each species in trait space (i.e. it equals +1 if z_i is larger than z_j, and -1

otherwise). The α value ($\alpha > 0$) determines the effect of pairwise similarity in trait *y* on

competition: if α is close to zero, all lineages sympatric with lineage *i* have the same competitive

effect on *i*, regardless of their similarity in trait *y*; conversely, if *α* is large, sympatric lineages

similar to i in terms of the *y* trait will have a much stronger competitive effect on *i* than

sympatric lineages dissimilar to i in terms of the *y* trait. The parameter *m* represents the

magnitude of the effect of competition when two lineages have identical *y* values (i.e., it provides

an upper bound for the deterministic effect of competition). When $m = 0$, this equation reduces to

an OU model, whereas positive *m* values result in pairwise divergence and negative values result

in pairwise convergence. When both *m* and $\psi = 0$, this model reduces to Brownian motion. For

additional simulation details, see *Supplementary Methods*.

We use a lineage-based "phenomenological" model for our simulations rather than an

individual-based model to have the computational ability to produce datasets of a size

comparable to the maximum sometimes reached in empirical comparative phylogenetic studies

(i.e. often reaching several hundreds of species). Models derived from microevolutionary first

principles (e.g., Grether et al. 2009; Nuismer and Harmon 2015) generate similar patterns of

sympatric shifts resulting from character displacement, and using such a model here would be

much more computationally intensive, therefore restricting the range of parameter values that

can be studied. For simplicity, this model also omits the effect of a species' geographic structure

and the effect of gene flow between distinct populations on the evolution of the mean species

phenotype. This simplification is reasonable in the context of our study because there is no

reason to expect that it will systematically bias the patterns generated in such a way as to yield

different conclusions regarding the performance of the various analytical approaches that we use

here. Finally, in all of our simulations, we considered sympatry to be a binomial variable, so $A_{i,j}$

equaled either 1 (if species *i* and *j* are sympatric) or 0 (if species *i* and *j* are allopatric). This index

of sympatry is similar to commonly used indices (Pigot and Tobias 2014; Tobias et al. 2014), but

other formulations of sympatry, such as continuous measurements of range overlap (Bothwell et

al. 2015; Martin et al. 2015) are also possible. We did not explore continuous measurements of

range overlap here, but have uploaded our simulation scripts to RPANDA (Morlon et al. 2016;

https://github.com/hmorlon/PANDA), which could easily be modified to do so.

Divergent character displacement.—We simulated datasets with divergent character

displacement by setting $y = z$ in Eq. 1 such that trait divergence is driven by pairwise similarity

in that trait. Biologically, this could represent a feeding trait that co-varies with resource use

(e.g., bill shape in Galápagos finches, Grant & Grant 2011) and which would be directly affected

by interspecific competition. To assess whether each method could detect divergent character

displacement when it occurred and did not erroneously detect character displacement when it

was absent, we simulated datasets both with repulsion $\{m = 2\}$ and without repulsion $\{m = 0\}$

(see Supplementary Methods). We also simulated datasets with $\{\psi = 2\}$ and without $\{\psi = 0\}$ the

OU process. In all simulations, we held σ^2 constant at 0.5, α constant at 1, and both the state at

the root (z_0) and the OU optimum (θ) constant at 0.

In additional simulations run only on 100-species trees, we analyzed the effect of both the

maximum strength of repulsion $\{m = 0, 1, 2, 10\}$ and, to understand how the opposing forces of

repulsion and attraction to an optimum influence analyses, the ratio of attraction to the maximum

effect of competition $\{\psi:m = 0, 0.2, 0.5, 1\}$ on inferences. To achieve these ratios of $\psi:m$, we

varied ψ while holding m constant (e.g., for the case where $m = 2$, we simulated datasets where ψ

$= 0, 0.4, 1,$ and 2, respectively). As above, these values were arbitrarily chosen based on visual

inspection of realized simulations.

For each parameter combination, we simulated 10 datasets for each tree, resulting in

1,000 simulations for each tree size / biogeographic scenario combination.

Convergent character displacement.—We simulated datasets with convergent character

displacement under Eq. 1, where the term y represents a trait determining resource use or niche

occupation evolving via BM or OU. A species' trait z in this model—a trait used as a territorial

signal—is thus attracted most strongly to the signal trait values of sympatric lineages with the

most similar resource-use traits. Biologically, this represents a scenario where selection favors

interspecific territoriality—mediated by similarity in territorial signals—because the benefits of

excluding heterospecifics are similar to the benefits of excluding conspecifics (Grether et al.

2009). As a species' resource-use trait becomes less similar to that of sympatric species, the

strength of attraction decreases to zero.

We simulated resource-use traits under both BM ($\sigma^2_{resource} = 0.5$, $\psi_{resource} = 0$) and OU

($\sigma^2_{resource} = 0.5$, $\psi_{resource} = 2$, $\theta_{resource} = 0$) models. For the signal trait, we simulated datasets both

with convergence $\{m = -0.25\}$ and without convergence $\{m = 0\}$. We did not include attraction

toward a stable peak for the signal trait (i.e. ψ was held constant at 0). As above, we held $\sigma^2 = 0.5$

and $z_0 = 0$, though we held α constant at 10, since smaller values result in rapid, cladewise

convergence in traits. To analyze the effect of the maximum strength of convergence, we ran

another set of simulations on 100-species trees varying m $\{m = 0, -0.1, -0.25, -0.5\}$ (see

Supplementary Methods). The resource trait (y) and signal trait (z) were modeled as unlinked and

genetically uncorrelated.

As above, we simulated 10 datasets for each tree, resulting in 1,000 simulations for each

tree size / biogeographic scenario combination.

Predictors of Interspecific Interactions

In some cases, investigators wish to identify which factors explain the occurrence of

particular interspecific interactions. For example, investigators may want to understand which

traits cause species to hybridize (e.g., Willis et al. 2014). In this scenario, species interactions

vary according to phenotypic similarity between sympatric species pairs (i.e., species pairs that

could potentially interact). Additionally, and unlike character displacement analyses, predicting

the occurrence of interspecific interactions requires treating trait similarity as a predictor variable

rather than a response variable. Thus, we generated datasets where the presence of interactions

between sympatric taxa depends on pairwise similarity in traits.

264   Under this scenario, we first evolved a trait along the phylogeny under a BM ($\sigma^2 = 0.5$, ψ

265     $= 0$) or OU ($\sigma^2 = 0.5$, $\psi = 2$, $\theta = 0$) model. Next, we simulated a second, independently evolving

266   trait ($\sigma^2_{unmeasured} = 1$, $\psi_{unmeasured} = 0$) to represent an unmeasured trait that could cause the

267   interaction of interest. To generate datasets where species interactions depend on similarity in

268   trait space at the present, we created species interactions in the form of a binomial variable by

269   sampling from a binomial distribution with the probability of interaction equal to:

271
$$P = \frac{e^{(b_1 Dx_1 + b_2 Dx_2)}}{1 + e^{(b_1 Dx_1 + b_2 Dx_2)}}$$

272   (Eq. 2)

(e.g., Hilbe 2009) where Dx_n is the distance between species at the present (e.g., distance

between tip values) in simulated trait n (simulated using fastBM in phytools, Revell 2012), and

b_n is the coefficient determining the magnitude of the relationship between the species

interaction and similarity in trait n. Trait 1 is the measured, focal trait and trait 2 represents the

independently evolving, unmeasured trait. As the effect of b_n on the species interaction depends

on the Dx_n distribution, which in turn depends on the total height of the tree, we scaled the trees

to a height of one prior to simulating datasets to facilitate comparison of results across trees and

parameter space.

To determine the statistical power of each analytical method, we generated species

interactions based on similarity in the measured trait ($b_1 = -4$, $b_2 = 0$); to assess the Type I error

rate, we simulated species interactions based on similarity in the unmeasured trait ($b_1 = 0$, $b_2 = -$

4). To determine the effect of the magnitude of the coefficient determining the relationship

between the measured trait and the interactions, we ran another set of simulations on 100-species

trees varying b_1 {$b_1 = 0$, -2, -4, -6, -8} and holding $b_2 = -4$. As above, we ran 1,000 simulations

for each tree size / biogeographic scenario combination.

Phylogenetic Tests

Among our tests of character displacement (both divergent and convergent), the

"correlation" tests involved assessing the significance of the relationship between phenotypic

similarity and coexistence, using either the "full" dataset (all species pairs) or the "sister taxa"

subset obtained by culling sister taxa from trees with ≥150 tips (Box 1, Diagram S1). To the full

datasets, we applied standard non-phylogenetic regression analyses that ignore phylogenetic non-

independence (Box 1.1), the raw and phylogenetically permuted partial Mantel tests (Box 1.2,

1.3), phylogenetic linear mixed models (PLMMs, Box 1.4), and the simulation approach (Box

1.5, Supplementary Methods). To the sister-taxa datasets, we applied non-phylogenetic

regression analyses (Box 1.1), PLMMs (Box 1.4), the simulation approach (Box 1.5), sister-taxa

GLMs (Box 1.7), and fit process based models in EvoRAG (Box 1.8, Supplementary Methods).

We did not perform Mantel tests on the sister-taxa data because such tests require complete

matrices and distance matrices with data for only sister taxa would mostly contain empty cells

(i.e. all those cells that correspond to non sister taxa species pairs). We compared the fit of

process-based phenotypic models with and without species interactions (Brownian motion,

Ornstein-Uhlenbeck, diversity dependent, and matching competition models; see Box 1.6 and

Supplementary Methods) to the full datasets from divergence scenarios using the R packages

geiger (Pennell et al. 2014) and RPANDA (Morlon et al. 2016). We acknowledge that diversity-

dependent models were not designed to analyze character displacement *per se*, but because they

incorporate interspecific interactions, we hypothesized that (and wanted to test if) they could be

useful in doing so. We did not apply process-based models to convergence scenarios because the

necessary model fitting tools have yet to be developed (see Discussion).

Our tests of predictors of species interactions involved assessing the significance of the

relationship between phenotypic similarity and species interactions (i.e., whether the species

interact where they occur in sympatry). Since the response variable is binary, we fit non-

phylogenetic logistic regressions, logistic PLMMs, and employed the simulation approach (see

Supplementary Methods). We did not perform Mantel tests or sister-taxa analyses because the

species pair matrix was incomplete (species that do not coexist cannot interact) and typically too

few sister taxa occurred in sympatry for regression analysis.

RESULTS

Divergent Character Displacement

When all possible pairwise comparisons are included in analyses, the ability of most

methods to detect divergent character displacement depends on the presence of the OU process.

As expected, non-phylogenetic regression analyses have a high Type I error rate in either

scenario (Figs. 2Ai,iv, S2Ai,iv [NB: throughout, results for low sympatric speciation

biogeographies are plotted in the main text and high sympatric speciation biogeographies in the

supplement]). When the OU process is present ($\psi = 2$), all methods generally have low Type I

error rates and high power (Figs. 2Aiv-vi, S2iv-vi, Supplementary Tables). However, when there

is no pull toward a peak ($\psi = 0$), the Type I error rate is higher for Mantel tests (Figs. 2Ai-ii, S2i-

ii), and the power is much lower for all methods, though the pppMantel and raw Mantel perform

better than the simulation and PLMM methods (Figs. 2Aiii, Fig. S2iii). Repulsion is easier to

detect against an OU background of traits converging toward a common optimum than against a

background of traits diverging under BM, likely because the repulsion process is more active

when species occupy similar trait space (Figs. S3, S4). High rates of sympatric speciation and

dispersal tend to slightly decrease the power of all methods (Fig. S2ii,iv, Supplementary Tables).

The ability to detect divergence was relatively similar for $m = 1$ and $m = 2$, but declined

for $m = 10$ (Fig. S5). This is due to a positive relationship between the ability to detect character

displacement and the ratio of ψ:m (Fig. S6), resulting from a higher absolute magnitude of

repulsion when both processes are present (Figs. S4, S6), indicating that this ratio impacts the

ability to detect divergence more than the raw value of m.

For sister-taxa analyses, there is a high probability of falsely concluding that character

displacement occurred in datasets simulated under BM and, to a lesser extent, OU, whether

analyzed with simple linear regressions, sister-taxa GLMs, or PLMMs (Figs. 2Bi,iii, S2Bi,iii).

As with the whole-tree approach, the power tends to increase and Type I error rate tends to

decrease in datasets with attraction toward a single-stationary peak (Figs. 2Biv-vi, S2Biv-vi).

However, the overall power to infer the presence of divergence was low with sister-taxa analyses

(Figs. 2Biii,vi, S2Biii,vi). Inferences were generally better when dispersal was high, which may

reflect the elevated observed divergence in high dispersal scenarios (Fig. S3). Allopatric

speciation scenarios increased the probability of Type I error (Fig. 2Bi-ii).

For the phylogenetic trait model-fitting analyses, BM and OU were generally correctly

chosen when they were the generating models (i.e., when $m = 0$ and when $\psi = 0$ or 2,

respectively, Figs. 3, S8). When $\psi = 0$ and $m > 0$, the matching competition (MC) model with

biogeography is consistently the best-fit model (Figs. 3A, S8A). When $m > 0$ and $\psi = 2$, the

diversity dependent exponential (DD_{exp}) model with biogeography was favored over other

models in most scenarios (Figs. 3B, S8B), with positive rate parameters estimated in the

maximum likelihood solution (Fig. S9). The biogeographic scenario did not greatly affect the

outcome of model fitting, though correct models were slightly more supported when dispersal

was high (Fig. S10), again in agreement with the observed magnitude of repulsion (Fig. S3).

Although the models are less identifiable when $m = 10$ and $\psi = 2$ (Figs. 3, S8), this results from

variation in the ψ:m ratio— there is a ratio of ψ:m around which these models cannot be

distinguished (Fig. S11).

Process-based models fit to sister-taxa datasets in EvoRAG did not mistakenly identify an

effect of species interactions when they were absent (Fig. S4A, C, Table 2), but they were unable

to identify the effect of competition when $\psi = 0$ (Fig. S4B, Table 2). However, as with process-

based models fit to the whole phylogeny, when data were simulated with both repulsion and a

pull toward a stable peak, a model where evolutionary rates vary linearly with the number of

sympatric taxa is often the best-fit model, though generally with only a marginally lower AICc

value (i.e., ΔAICc < 2) than BM (Fig. S4, Table 2).

Figure 2. Proportion of statistically significant analyses in datasets simulated under divergent

character displacement in biogeographic scenarios with low sympatric speciation rates. A.

Results from approaches using data from all pairwise comparisons in a clade, plotted as a

function of the phylogeny size and dispersal rate when i-ii. $m = 0$ and $\psi = 0$ (i. all analyses and ii.

only analyses returning divergence in sympatry), iii. $m = 2$ and $\psi = 0$, iv-v. $m = 0$ and $\psi = 2$ (iv.

all analyses and v. only analyses returning divergence in sympatry), and vi. $m = 2$ and $\psi = 2$. B.

Results from analyses of sister-taxa culled from complete phylogenies binned by the number of

resulting species pairs, plotted as a function of the number of sister taxa comparisons and

dispersal rate when i-ii. $m = 0$ and $\psi = 0$ (i. all analyses and ii. only analyses returning

divergence in sympatry), iii. $m = 2$ and $\psi = 0$, iv-v. $m = 0$ and $\psi = 2$ (iv. all analyses and v. only

analyses returning divergence in sympatry), and vi. $m = 2$ and $\psi = 2$. For scenarios where $m = 2$,

only the proportion of significant results showing divergence are plotted. Dashed horizontal lines

represent a Type I error rate of 5%.

386
387

388
Figure 3. Boxplots of Akaike weights for each trait model fit to simulated datasets in

biogeographic scenarios with low sympatric speciation rates as a function of *m* in trees with 100

species. A. When OU is absent, BM is the best-fit model when $m = 0$, and the matching

competition model with biogeography is the best model when competitive divergence is present.

B. When OU is present, OU is the best-fit model when $m = 0$, and the diversity-dependent

exponential model with biogeography is the best model when competitive divergence is present

and $\psi{:}m$ is relatively high.

Convergent Character Displacement

397
As with divergent character displacement, with all pairwise species combinations, the

ability of most methods to detect convergent character displacement depends on the presence of

the OU process on the resource-use trait: datasets simulated under an OU model were more

likely to be statistically significant (Figs. 4A.iv-vi, S12A.iv-vi) across all methods than those

with BM simulated resource-use traits (Figs. 4A.i-iii, S12A.i-iii). Again, this is likely because

the presence of the OU process in the resource-use trait amplifies the magnitude of convergence

(Fig. S13, S14). Overall, however, only the simulation approach had substantial power (> 0.80)

to detect convergent character displacement (Table 1), and only in trees with 100 or more tips

and datasets with the OU process in the simulated resource-use trait. Indeed, the non-

phylogenetic regressions often (spuriously) detected divergence rather than the simulated

convergence, especially in smaller trees (Supplementary Tables). Both types of Mantel tests were

unable to detect convergence, in fact having a higher Type I error rate (detecting divergence in

BM simulated datasets, Supplementary Tables) than power. As with divergent character

displacement, there was a tendency for higher power in lower dispersal scenarios.

The power to detect convergence generally increased with increasingly negative values of

m, the maximum strength of attraction in the signal trait when species are identical in the

resource-use trait (Fig. S15), though as *m* gets large, the probability that all species converge on

the same trait value increases, especially when $\psi_{resource} = 2$ (data not shown).

Regardless of whether resource-use traits are simulated under OU or BM, when there is

no convergence, most methods used for sister-taxa analyses tend to have high Type I error rates,

though these analyses return an erroneous inference of divergence, rather than convergence,

between sister taxa (Figs. 4B.i,ii,iv,v, S12B.i,ii,iv,v, Supplementary Tables). Sister-taxa analyses

had overall low power to detect convergence when it did exist, and non-phylogenetic regressions

often detected divergence, rather than convergence (Supplementary Tables). When convergence

was detected, it tended to be in biogeographic scenarios with high dispersal, likely reflecting the

overall magnitude of convergence achieved (Fig. S13). As with divergent character displacement

simulations, the allopatric speciation biogeographic scenarios were more likely to lead to higher

Type I error rates (Figs. 4B.i,iv). Process-based models fit to sister-taxa datasets in EvoRAG did

not erroneously detect divergence or convergence (i.e., BM was the best-fit model when $m = 0$,

Fig. S14 A, C, Table 2), but they could not detect an effect of species interactions when

convergence was present, at least for the number of sister taxa in this study, as OU was the best-

fit model when $m = -0.25$ (Fig. S14 B, C, Table 2).

Figure 4. Proportion of statistically significant analyses in datasets simulated under convergent

character displacement in biogeographic scenarios with low sympatric speciation rates. A.

Results from approaches using data from all pairwise comparisons in a clade, plotted as a

function of the phylogeny size and dispersal rate when i-ii. $m = 0$ and $\psi_{resource} = 0$ (i. all analyses

and ii. only analyses returning convergence in sympatry), iii. $m = -0.25$ and $\psi_{resource} = 0$, iv-v. m

$= 0$ and $\psi_{resource} = 2$ (iv. all analyses and v. only analyses returning convergence in sympatry),

and vi. $m = -0.25$ and $\psi_{resource} = 2$. B. Results from analyses of sister-taxa culled from complete

phylogenies binned by the number of resulting species pairs, plotted as a function of the number

of sister taxa comparisons and dispersal rate when i-ii. $m = 0$ and $\psi_{resource} = 0$ (i. all analyses and

ii. only analyses returning convergence in sympatry), iii. $m = -0.25$ and $\psi_{resource} = 0$, iv-v. $m = 0$

and $\psi_{resource} = 2$ (iv. all analyses and v. only analyses returning convergence in sympatry), and vi.

$m = -0.25$ and $\psi_{resource} = 2$. For scenarios where $m = -0.25$, only the proportion of significant

results showing convergence are plotted. Dashed horizontal lines represent a Type I error rate of

5%.

Predicting Interspecific Interactions

Although all three methods used to identify traits that are causally related to interspecific

interactions had high power (>>0.8, Table 1, Supplementary Tables) to do so in the parameter

space explored here (Figs. 5ii,iv, S16ii,iv), only the simulation approach had both high power

and a low Type I error rate (Table 1), whereas non-phylogenetic regressions and PLMMs had

fairly high Type I error rates (Table 1) when interactions were simulated based on similarity in a

trait other than the measured one (Fig. 5i,iii, S16i,iii). The power to detect an interaction was not

greatly affected by the coefficients used to simulate datasets (Fig. S17). Biogeography did not

have a large impact on analyses, though there were slightly higher Type I error in low-dispersal

scenarios (Fig. 5i,iii).

461
Figure 5. Proportion of statistically significant analyses in datasets with interactions simulated

under a simple phenotype matching process in biogeographic scenarios with low sympatric

speciation rates. Results from analyses where the measured trait was simulated under BM (i, ii)

or OU (iii, iv), plotted as a function of the phylogeny size and dispersal rate when i. b_1 (the

simulation coefficient determining the relationship between the interaction and the measured

trait) = 0, b_2 (the simulation coefficient for an unmeasured trait) = -4, and ψ = 2, ii. b_1 = -4, b_2 =

0, and ψ = 2, iii. b_1 = 0, b_2 = -4, and ψ = 0, and iv. b_1 = -4, b_2 = 0, and ψ = 0.

DISCUSSION

As open-access databases with species range, trait, and phylogenetic data rapidly expand, investigators are able to test hypotheses about the relationships between interspecific interactions and phenotypic evolution at an unprecedented scale. Understanding the relative strengths and weaknesses of phylogenetic comparative methods available for testing such hypotheses is thus paramount. We found that currently used methods for detecting causal relationships between interspecific interactions and species phenotypes suffer from severe limitations (Tables 1,2).

Overall, standard methods are better at detecting divergent character displacement when divergence does not drive unbounded trait evolution (i.e., when selection acts against extreme phenotypes, as can be modeled by the OU process). Consistent with previous reports (Harmon and Glor 2010; Guillot and Rousset 2013), Mantel tests had high Type I error rates and both standard and pppMantel tests have low power (Table 1, Figs. 2Ai, S2Ai). Alarmingly, we found that commonly used sister taxa approaches have high Type I error rates (Table 2, Figs. 2Bi,iv, S2Bi,iv, 4Bi,iv, S12Bi,iv, Supplementary Tables), which would lead investigators to conclude that divergent character displacement had occurred when, in fact, it had not, and none have a reasonable combination of Type I error and power. Thus, we discourage empiricists from using sister-taxa approaches to study character displacement. If no other data are available for testing for character displacement on the whole tree, then we recommend phylogenetic simulations, as they are the only method with reasonably low type I error rates, even though they suffer from low power (Tables 1, 2).

Fitting process-based phylogenetic trait models to datasets simulated with divergent character displacement yielded more consistent patterns (Fig. 3). Without attraction toward a single stationary peak to bound trait evolution, the matching competition (MC) model with

493   biogeography was predominantly the best-fit model. For datasets simulated with the OU process,

494   the diversity-dependent exponential (DD_{exp}, see Box 1) model with biogeography was the best-fit

495   model, and similarly a model with a linear relationship between evolutionary rates and the

496   number of sympatric taxa often fit sister-taxa datasets, though with much lower power overall

Table 1. Summary of the statistical properties of the analytical approaches tested under scenarios using data from all tips (i.e., with sister-taxa analyses excluded). Values refer to the range of type I error rates and power levels for each tree size $\geq$50, averaged across biogeographic scenarios and scenarios where ψ or ψ_{nf} = 0 or 2. Power refers to only those statistically significant tests in the appropriate tail (i.e., in the lower tail for divergent character displacement and upper tail for convergent character displacement). For each analytical scenario, the cell with the method with the best trade-off between Type I error and power is highlighted.

Analysis	non-phylogenetic regression		Mantel test		pppMantel test		PLMM		simulation test		process-based models	
	type I	*power*	*type I*	*power*	*type I*	*power*	*type I*	*power*	*type I*	*power*	*type I**	*power†*
divergent char. displacement	0.37-0.61	0.51-1	0.05-0.10	0.29-1	0.04-0.06	0.19-1	0.05-0.06	0.12-1	0.05-0.07	0.08-1	0.04-0.05	0.91-0.94
convergent char. displacement	0.40-0.60	0.31-0.99	0.08-0.09	0-0.02	0.05-0.06	0-0.01	0.05-0.07	0.07-0.26	0.04-0.05	0.12-0.91	--	--
predicting spp. interactions	0.08-0.3	1	--	--	--	--	0.07-18	1	0.03-0.04	1	--	--

*Type I error rate calculated as the proportion of datasets simulated without divergence for which a model that includes species interaction— DD_{exp}, DD_{lin}, or MC—was chosen by model selection (i.e., for which ΔAICc = 0 and ΔAICc for all other models > 2).

† Power calculated as the proportion of datasets simulated with divergence for which either DD_{exp}, DD_{lin}, or MC was chosen by model selection.

508 **Table 2.** Summary of the statistical properties of the analytical approaches tested under scenarios using sister-taxa analyses. Values

509 refer to the range of type I error rates and power levels, averaged across biogeographic scenarios and scenarios where ψ or $\psi_{nf} = 0$ or

510 2. Power refers to only those statistically significant tests in the appropriate tail (i.e., in the upper tail for divergent character

511 displacement and lower tail for convergent character displacement). We caution against using sister-taxa approaches to test for

512 character displacement.

Analysis	non-phylogenetic regression		sister-taxa GLM		PLMM		simulation test		process-based models in EvoRAG	
	type I	power	type I	power	type I	power	type I	power	type I*	power[†]
divergent char. displacement	0.07-0.42	0.68-0.75	0.05-0.07	0.20-0.34	0.08-0.50	0.70-0.78	0.01-0.02	0.18-0.31	0.04-0.07	0.03-0.36
convergent char. displacement	0.33-0.43	0.01-0.2	0.07	0.04-0.06	0.41-0.5	0.02-0.21	0.03	0.01-0.2	0.04	0.09-0.55

514 *Type I error rate calculated as the proportion of datasets simulated without divergence for which a model that includes a linear dependency on the number of sympatric lineages—

515 BM_{linear} or OU_{linear_beta}—was chosen by model selection (i.e., for which $\Delta AICc = 0$ and $\Delta AICc$ for all other models > 2).

516 [†] Power calculated as the proportion of datasets simulated with divergence for which either BM_{linear} or OU_{linear_beta} was chosen by model selection.

517
518

(Fig S4, Table 2). In the DD_{exp} model, rates of trait evolution vary exponentially with the

number of sympatric lineages through time, so incorporating the effect of interspecific

interactions on the rate of trait of evolution but not explicitly modeling the process of character

displacement acting on the mean trait values. It may nonetheless provide a useful proxy for

detecting patterns that are similar to those left by character displacement, in the absence of a

process-based model that incorporates both attraction toward an optimum trait value and

divergent character displacement. We emphasize, however, that statistical support for the DD_{exp}

model does not in itself constitute decisive evidence that character displacement has occurred, as

other processes may generate increasing evolutionary rates with increasing lineage diversity.

Given that the DD_{exp} model is the best-fit model in parameter space where other methods also

perform well, combined evidence from model-fitting and other, non-process based methods

would constitute a strong case for the presence of character displacement. In the absence of tip

data (e.g., due to incomplete sampling or traits that are inherently measured as pairwise

properties), process-based models are unsuitable and we recommend using data from as many

species pairs as possible—not just sister taxa—and using simulation approaches or PLMMs. In

other words, to detect divergent character displacement, we recommend that empiricists fit the

MC model to their dataset when possible. High support for the MC model would constitute

evidence that character displacement has acted on a trait. If the MC model does not provide a

good fit for the data, this could be because character displacement proceeds in the presence of

bounded trait evolution, in which case a signature of the DD_{exp} model with a positive rate

parameter and/or a signature of sympatric divergence in phylogenetic simulations or PLMMs

would constitute evidence consistent with divergent character displacement.

Interestingly, even though most previous investigators have used the DD_{exp} model to

represent a decline in ecological opportunity with increasing species richness (Mahler et al.

2010; Weir and Mursleen 2013), the maximum likelihood estimates of the rate parameters for

this model were positive, rather than negative, when both divergence and the OU process were

present (Fig. S9). This is consistent with our finding of increasing evolutionary rates with

increasing species richness (Figs. S3, S4, S7) in this scenario. An increase in the rate of

evolutionary changes in trait values toward the present likely results from selection not only

restricting species to certain trait space but also partitioning that space. The resulting adaptive

landscape is therefore rapidly changing, causing accelerating evolutionary rates as lineages fill

this increasingly constrained space.

The MC model (Box 1) is similar to the model used to simulate data (Eq. 1), with the

assumption that α is very small ($\ll 1$) and consequently, competitive interactions are affected by

the mean trait values of all sympatric species, rather than by pairwise similarity (Nuismer and

Harmon 2015; Drury et al. 2016). Biologists, however, generally assume that competition is

stronger between phenotypically similar species (Brown and Wilson 1956). Our results show that

the assumption of a small α does not render the MC model useless for studying character

displacement, as the MC model is the best-fit model for many datasets simulated under the

character displacement model used here. Nevertheless, the finding that the DD_{exp} model is the

best-fit model in datasets simulated under character displacement including OU indicates that the

MC model is not a perfect model of character displacement. Recently, approximate Bayesian

computational (ABC) tools have been published to fit a model of character displacement in

which, like in our simulation model, the strength of competition depends on similarity in trait

space (Clarke et al. 2017). This model provides an alternative tool for detecting character

displacement in comparative datasets, and we hope that further development of methods such as

this ABC method will help ameliorate the statistical issues shown here.

For datasets simulated including the OU process, the ratio of the pull-parameter in the

OU portion of the model to the maximum amount of repulsion (ψ:m) had a consistent impact

across all methods, which results from the overall larger magnitude of evolutionary changes in

traits in scenarios with a high ψ:m ratio (Figs. S3, S4, S7). As ψ:m approached 1, all methods

were better at detecting character displacement. Currently, there are no analytical approaches that

can disentangle the simultaneous impact of attraction toward a peak and divergence due to

competition, though we hope our results will inspire development of such tools.

Unlike for divergent character displacement, available statistical methods for detecting

convergence in comparative datasets generally do a poor job of detecting convergence, with the

simulation method outperforming others (Table 1). With whole-dataset approaches, Type I error

rates are acceptable for phylogenetic analyses (~5%), however, so although detecting

convergence is difficult, the risk of mistakenly detecting convergence is low. In sister-taxa

analyses, although Type I error rates are high for PLMMs (Table 2), these largely return

erroneous divergence results, rather than erroneous convergence (Figs. 4Bii,v, S12Bii,v). In

short, if an empiricist detects convergence in their dataset, they can be fairly confident in the

result. Yet if empiricists do not detect convergence, this could simply be a result of lower power

of the available analytical tools. Currently, there are no tools to fit phylogenetic trait models of

convergence between species (e.g., Nuismer & Harmon 2015); such tools might more

successfully identify convergent character displacement in comparative datasets than the

available statistical methods.

For both divergent and convergent character displacement scenarios, we found that sister-

taxa GLMs and the simulation approach had a mean Type I error rate near 5% (Table 2).

However, in some scenarios, the Type I error for sister taxa GLMs was much higher than for the

simulation approach (Figs. 2Bi, 4Bi, Supplementary Tables), which suggests that including a

model-based estimate of the rate of trait evolution more properly accounts for the effect of

divergence than simply including the branch lengths separating sister taxa as a covariate in

analyses to control for variation in the amount of time sister taxa have had to diverge from one

another. The high overall Type I error rate for sister-taxa analyses may also result from the

unrealistic assumption that transitions between allopatry and sympatry are uncommon along

branches connecting sister taxa (Weir and Price 2011; Tobias et al. 2014). Supporting this

explanation, we found that biogeographic scenarios with high levels of sympatric speciation and

low dispersal tended to have overall lower Type I error rates (cf. Figs. 2,S2; Figs. 4,S12).

The outlook for identifying which traits drive species interaction is brighter. The

statistical methods available to test for causal relationships between phenotypic similarity and

interactions between species have very high power. The simulation approach has a low Type I

error rate when causal relationships are simulated based on an unmeasured trait, although non-

phylogenetic regressions and PLMMs suffer from relatively high Type I error rates (Table 1).

Thus, we recommend that empiricists interested in predicting pairwise species interactions based

on trait data use phylogenetic simulations. While we did not simulate interactions between

clades, our results are likely applicable to other empirical questions, such as identifying traits that

predict links in ecological networks (Rafferty and Ives 2013; Hadfield et al. 2014; Eklöf and

Stouffer 2016).

By simulating datasets with various types of interactions between species across different

modes of speciation and dispersal rates, we have shown that many of the methods that

investigators use to analyze empirical datasets have low power to detect such patterns (Table 1).

Worse still, widely-used sister taxa approaches, including standard regressions and sister-taxa

GLMs, often detected character displacement in datasets that were simulated under a simple BM

model (Figs. 2Bi-ii, 4Bi-ii). We therefore urge investigators to use caution when interpreting the

results of such analyses, even in cases when sympatry is delineated using other criteria than the

one considered here. When process-based models could be fit to these datasets, they tended to

consistently identify patterns of divergence (i.e., either the matching competition model or a

diversity-dependent model is the best fit model >91% of the time). Thus, when possible,

empiricists should employ such methods. Statistical tools to fit process-based models of

phenotypic evolution including species interactions are in their infancy (Drury et al. 2016;

Manceau et al. 2017) and many possible models are not yet available (e.g., convergent character

displacement, character divergence in the presence of an adaptive pull towards a peak). We hope

that our results encourage the continued development of such tools.

In closing, we note that divergent character displacement is erroneously detected with

many statistical approaches, indicating that there may be an overrepresentation of empirical

studies that imply that divergence has occurred. In particular, studies that have used sister-taxa

methods to document character displacement may have falsely interpreted a null expectation—

larger trait differences between sympatric lineages owing to allopatric speciation—as evidence

for divergent character displacement. Conversely, convergent character displacement is often

hard to detect with existing methods, suggesting that convergence in signal traits (e.g., Cody

1969, 1973; Tobias et al. 2014; Losin et al. 2016) might be more prevalent than previously

thought.

ACKNOWLEDGEMENTS

This research was funded by the European Research Council (grant 616419-PANDA to HM) and

the National Science Foundation (grant DEB-1457844 to GFG). We thank M. Manceau, J.

Clavel, E. Lewitus, O. Maliet, and O. Missa for feedback and F. Hartig for assistance

streamlining our simulation script.

REFERENCES

Allen W.L., Stevens M., Higham J.P. 2014. Character displacement of Cercopithecini primate

visual signals. Nat. Commun. 5:4266.

Bacquet P.M.B., Brattström O., Wang H.L., Allen C.E., Löfstedt C., Brakefield P.M.,

Nieberding C.M. 2015. Selection on male sex pheromone composition contributes to

butterfly reproductive isolation. Proc. R. Soc. London B Biol. Sci. 282:20142734.

Bothwell E., Montgomerie R., Lougheed S.C., Martin P.R. 2015. Closely related species of birds

differ more in body size when their ranges overlap — in warm , but not cool , climates.

Evolution. 69:1701–1712.

Brown W.L.L., Wilson E.O. 1956. Character displacement. Syst. Zool. 5:49–64.

Cadotte M., Albert C.H., Walker S.C. 2013. The ecology of differences: Assessing community

assembly with trait and evolutionary distances. Ecol. Lett. 16:1234–1244.

Cavender-Bares J., Kozak K.H., Fine P.V.A., Kembel S.W. 2009. The merging of community

ecology and phylogenetic biology. Ecol. Lett. 12:693–715.

Clarke M., Thomas G.H., Freckleton R.P. 2017. Trait evolution in adaptive radiations: modelling

and measuring interspecific competition on phylogenies. Am. Nat. 189:doi:

10.1086/689819.

Cody M.L. 1969. Convergent characteristics in sympatric species: a possible relation to

interspecific competition and aggression. Condor. 71:223–239.

Cody M.L. 1973. Character Convergence. Annu. Rev. Ecol. Syst. 4:189–211.

Davies J.T., Meiri S., Barraclough T.G., Gittleman J.L. 2007. Species co-existence and character

divergence across carnivores. Ecol. Lett. 10:146–152.

Drury J., Clavel J., Manceau M., Morlon H. 2016. Estimating the effect of competition on trait

evolution using maximum likelihood inference. Syst. Biol. 65:700–710.

Drury J.P., Okamoto K.W., Anderson C.N., Grether G.F. 2015. Reproductive interference

explains persistence of aggression between species. Proc. R. Soc. B. 282:20142256.

Eklöf A., Stouffer D.B. 2016. The phylogenetic component of food web structure and intervality.

Theor. Ecol. 9:107–115.

Elias M., Gompert Z., Jiggins C., Willmott K. 2008. Mutualistic interactions drive ecological

niche convergence in a diverse butterfly community. PLoS Biol. 6:e300.

Felsenstein J. 1985. Phylogenies and the comparative method. Am. Nat. 125:1–15.

Felsenstein J. 1988. Phylogenies and quantitative characters. Annu. Rev. Ecol. Syst. 19:445–471.

Garland T., Dickerman A.W., Janis C.M., Jones J.A. 1993. Phylogenetic analysis of covariance

by computer simulation. Syst. Biol. 42:265–292.

Grant P.R. 1972. Convergent and divergent character displacement. Biol. J. Linn. Soc. 4:39–68.

Grant P.R., Grant B.R. 2006. Evolution of character displacement in Darwin's finches. Science.

313:224–226.

Grant P.R., Grant B.R. 2011. How and Why Species Multiply: the Radiation of Darwin's

Finches. Princeton, NJ: Princeton University Press.

Grether G.F., Anderson C.N., Drury J.P., Kirschel A.N.G., Losin N., Okamoto K., Peiman K.S.

2013. The evolutionary consequences of interspecific aggression. Ann. N. Y. Acad. Sci.

1289:48–68.

Grether G.F., Losin N., Anderson C.N., Okamoto K. 2009. The role of interspecific interference

competition in character displacement and the evolution of competitor recognition. Biol.

Rev. 84:617–635.

Guillot G., Rousset F. 2013. Dismantling the Mantel tests. Methods Ecol. Evol. 4:336–344.

Hadfield J.D., Krasnov B.R., Poulin R., Nakagawa S. 2014. A tale of two phylogenies:

comparative analyses of ecological interactions. Am. Nat. 183:174–187.

Hadfield J.D., Nakagawa S. 2010. General quantitative genetic methods for comparative biology:

phylogenies, taxonomies and multi-trait models for continuous and categorical characters. J.

Evol. Biol. 23:494–508.

Hansen T.F., Martins E.P. 1996. Translating between microevolutionary process and

macroevolutionary patterns: The correlation structure of interspecific data. Evolution.

50:1404–1417.

Harmon L.J., Glor R.E. 2010. Poor statistical performance of the Mantel test in phylogenetic

comparative analyses. Evolution. 64:2173–2178.

Harmon L.J., Losos J.B., Jonathan Davies T., Gillespie R.G., Gittleman J.L., Bryan Jennings W.,

Kozak K.H., McPeek M.A., Moreno-Roark F., Near T.J., Purvis A., Ricklefs R.E., Schluter

D., Schulte J.A., Seehausen O., Sidlauskas B.L., Torres-Carvajal O., Weir J.T., Mooers

A.T. 2010. Early bursts of body size and shape evolution are rare in comparative data.

Evolution. 64:2385–2396.

Hilbe J.M. 2009. Logistic regression models. CRC press.

Lapointe F.J., Garland T. 2001. A generalized permutation model for the analysis of cross-

species data. J. Classif. 18:109–127.

Losin N., Drury J.P., Peiman K.S., Storch C., Grether G.F. 2016. The ecological and

evolutionary stability of interspecific territoriality. Ecol. Lett. 19:260–267.

Mahler D.L., Revell L.J., Glor R.E., Losos J.B. 2010. Ecological opportunity and the rate of

morphological evolution in the diversification of greater Antillean anoles. Evolution.

64:2731–2745.

Manceau M., Lambert A., Morlon H. 2017. A unifying comparative phylogenetic framework

including traits coevolving across interacting lineages. Syst.

Biol.:https://doi.org/10.1093/sysbio/syw115.

Mantel N. 1967. The detection of disease clustering and a generalized regression approach.

Cancer Res. 27:209–220.

Martin P.R., Montgomerie R., Lougheed S.C. 2010. Rapid sympatry explains greater color

pattern divergence in high latitude birds. Evolution. 64:336–347.

Martin P.R., Montgomerie R., Lougheed S.C. 2015. Color patterns of closely related bird species

are more divergent at intermediate levels of breeding-range sympatry. Am. Nat. 185:443–

451.

Martins E.P., Garland Jr T. 1991. Phylogenetic analyses of the correlated evolution of continuous

characters: a simulation study. Evolution. 45:534–557.

Matzke N.J. 2014. Model selection in historical biogeography reveals that founder-event

speciation is a crucial process in island clades. Syst. Biol. 63:951–970.

Medina-García A., Araya-Salas M., Wright T.F. 2015. Does vocal learning accelerate acoustic

diversification? Evolution of contact calls in Neotropical parrots. J. Evol. Biol. 28:1782–

1792.

Morlon H., Lewitus E., Condamine F.L., Manceau M., Clavel J., Drury J. 2016. RPANDA: an R

package for macroevolutionary analyses on phylogenetic trees. Methods Ecol. Evol. 7:589–

597.

Nuismer S.L., Harmon L.J. 2015. Predicting rates of interspecific interaction from phylogenetic

trees. Ecol. Lett. 18:17–27.

Pennell M.W., Eastman J.M., Slater G.J., Brown J.W., Uyeda J.C., FitzJohn R.G., Alfaro M.E.,

Harmon L.J. 2014. geiger v2. 0: an expanded suite of methods for fitting macroevolutionary

models to phylogenetic trees. Bioinformatics. 30:2216–2218.

Pfennig K.S., Pfennig D.W. 2009. Character displacement: Ecological and reproductive

responses to a common evolutionary problem. Q. Rev. Biol. 84:253–276.

Pigot A.L., Tobias J.A. 2014. Dispersal and the transition to sympatry in vertebrates. Proc. R.

Soc. B Biol. Sci. 282:20141929.

Rafferty N.E., Ives A.R. 2013. Phylogenetic trait-based analyses of ecological networks.

Ecology. 94:2321–2333.

Ree R.H., Smith S.A. 2008. Maximum likelihood inference of geographic range evolution by

dispersal, local extinction, and cladogenesis. Syst. Biol. 57:4–14.

Rezende E.L., Diniz-Filho J.A.F. 2012. Phylogenetic analyses: Comparing species to infer

adaptations and physiological mechanisms. Compr. Physiol. 2:639–674.

Rezende E.L., Lavabre J.E., Guimaraes Jr. P.R., Jordano P., Bascompte J. 2007. Non-random

coextinctions in phylogenetically structured mutualistic networks. Nature. 448:925–929.

Roncal J., Henderson A., Borchsenius F., Cardoso S.R.S., Balslev H. 2012. Can phylogenetic

signal, character displacement, or random phenotypic drift explain the morphological

variation in the genus Geonoma (Arecaceae)? Biol. J. Linn. Soc. 106:528–539.

Schluter D. 2000. The Ecology of Adaptive Radiation. Oxford, UK: Oxford University Press.

Seddon N., Botero C.A., Tobias J.A., Dunn P.O., Macgregor H.E., Rubenstein D.R., Uy J.A.,

Weir J.T., Whittingham L.A., Safran R.J. 2013. Sexual selection accelerates signal

evolution during speciation in birds. Proc. R. Soc. B Biol. Sci. 280:20131065.

Tobias J.A., Cornwallis C.K., Derryberry E.P., Claramunt S., Brumfield R.T., Seddon N. 2014.

Species coexistence and the dynamics of phenotypic evolution in adaptive radiation. Nature.

506:359–363.

de Villemereuil P., Nakagawa S. 2014. General quantitative genetic methods for comparative

biology. Modern Phylogenetic Comparative Methods and Their Application in Evolutionary

Biology. Springer. p. 287–303.

Webb C.O., Ackerly D.D., McPeek M.A., Donoghue M.J. 2002. Phylogenies and community

ecology. Annu. Rev. Ecol. Syst.:475–505.

Weber M.G., Mitko L., Eltz T., Ramírez S.R. 2016. Macroevolution of perfume signalling in

orchid bees. Ecol. Lett.

Weir J.T., Lawson A. 2015. Evolutionary rates across gradients. Methods Ecol. Evol. 6:1278–

1286.

Weir J.T., Mursleen S. 2013. Diversity-dependent cladogenesis and trait evolution in the

adaptive radiation of the auks (Aves: Alcidae). Evolution. 67:403–416.

Weir J.T., Price T.D. 2011. Limits to speciation inferred from times to secondary sympatry and

ages of hybridizing species along a latitudinal gradient. Am. Nat. 177:462–469.

Weir J.T., Wheatcroft D. 2011. A latitudinal gradient in rates of evolution of avian syllable

diversity and song length. Proc. R. Soc. B Biol. Sci. 278:1713–20.

Willis P.M., Symula R.E., Lovette I.J. 2014. Ecology, song similarity and phylogeny predict

natural hybridization in an avian family. Evol. Ecol. 28:299–322.

Box 1. Methods for assessing the interplay between interspecific interactions and species

phenotypes

777   Comparative analyses of the interplay between interspecific interactions and species

phenotypes can either be conducted on entire clades, or, commonly, on sister taxa—species pairs

that share a most recent common ancestor—that are culled from larger phylogenies. Such

analyses generally consist of testing the statistical significance of correlations between either

phenotypic similarity and geographic overlap (to test for divergent or convergent character

displacement) or species interactions and phenotypic similarity (to find predictors of species

interactions). As we are looking for correlations between pairwise comparisons (e.g., trait

similarity, biogeographical overlap, hybridization, magnitude of pre-zygotic isolation), rather

than "tip values" belonging to a single species, phylogenetically independent contrasts and

extensions of PGLS analyses (Felsenstein 1985; Rezende and Diniz-Filho 2012) cannot be used,

and alternative tests have been developed.

1. Non-phylogenetic regressions

790   "Non-phylogenetic regressions" refers to Generalized Linear Models (GLMs) that ignore

phylogenetic structure. Though less commonly applied to whole-clade analyses, investigators

sometimes use non-phylogenetic regressions for sister-taxa analyses, on the basis that branches

connecting sister taxa represent independent evolutionary histories (Felsenstein 1985).

2. Mantel tests

Several previous investigators have implemented Mantel tests (Mantel 1967) in analyses

of species-pair comparisons (e.g., Roncal *et al.* 2012). These tests are designed to assess

correlations between matrices, which here comprise interspecific trait distances or differences.

Existing accounts of Mantel tests describe procedures only for complete matrices, so they cannot

be used in many cases, including sister-taxa analyses (for which most off-diagonal elements of

distances matrices are by definition excluded) and in identifying predictors of species

interactions (e.g., hybridization), as only sympatric lineages can interact and setting values for

allopatric comparisons to zero would not make biological sense.

3. Phylogenetically permuted partial Mantel tests

Phylogenetically permuted partial Mantel (pppMantel) tests (Lapointe and Garland 2001)

account for phylogenetic non-independence by permuting null datasets that are structured

phylogenetically, and are popular among investigators studying species interactions (e.g., Allen

et al. 2014; Willis *et al.* 2014; Medina-García *et al.* 2015). Like Mantel tests, pppMantel tests

also require complete interaction matrices.

4. Phylogenetic linear mixed models

In recent years, researchers have adapted animal models from quantitative genetics to

incorporate phylogenies as random effects in mixed-effect regressions on comparative datasets

(Hadfield & Nakagawa 2010). Such phylogenetic linear mixed models (PLMMs) have been

modified to accommodate pairwise species data (Tobias et al. 2014), wherein the identity of the

species being compared and the node connecting them in the phylogeny are included as random

effects. PLMMs are promising new tools, as they are not limited to sister-taxa data and model

predictions can be generated and plotted.

5. Phylogenetic simulations

Simulation approaches are widely used to control for phylogenetic non-independence in

tip data (Martins & Garland Jr 1991; Garland *et al.* 1993), and have been applied to pairwise

species comparisons (Elias et al. 2008; Drury et al. 2015; Losin et al. 2016). In these approaches,

trait evolution is simulated along phylogenies, often scaled such that the simulated tip data

resemble real data. Pairwise comparisons are then calculated on many simulated datasets and

used to generate a phylogenetically informed null distribution of test statistics against which to

compare test statistics calculated from the real data.

6. Process-based models of phenotypic evolution

In the statistical approaches outlined thus far, the data analyzed are measurements of

pairwise differences between species, and the statistical tests for the effect of species interactions

on trait evolution consist of testing for significant correlations between either phenotypic

similarity and geographic overlap or species interactions and trait similarity. However, it is also

possible to detect a signature of interspecific competition in the distributions of continuous trait

values across the tips of a phylogeny by fitting process-based models of phenotypic evolution to

the data. These models allow testing hypotheses about which processes are most likely to have

generated the observed distribution of traits in a clade (Hansen & Martins 1996).

Interspecific interactions have recently been incorporated into such models in two ways.

First, in diversity-dependent (DD) models, evolutionary rates change as a function (either linear

[DD$_{lin}$] or exponential [DD$_{exp}$]) of the number of extant lineages through time (e.g., Weir &

Mursleen 2013). Secondly, in the 'matching competition' (MC) model, trait evolution in an

evolving lineage varies as a function of the values of traits in other evolving lineages (Nuismer &

Harmon 2015, Drury *et al*. 2016). Comparing the fit of these models to other models that exclude

interspecific interactions (e.g., Brownian motion and Ornstein-Uhlenbeck models) tests whether

there is evidence that interspecific interactions have influenced the trajectory of trait evolution in

a clade.

7. Sister-taxa GLMs

If allopatric speciation is common, then sympatry occurs after a period of initial isolation,

resulting in a pattern where sympatric sister taxa are older than allopatric sister taxa. Thus, even

random genetic drift can generate a pattern in which sympatric lineages have more divergent

traits compared to allopatric lineages, simply because divergence has had more time to evolve

(Weir and Price 2011; Tobias et al. 2014). To control for variation in the evolutionary distance

between sister taxa, "sister taxa GLMs" include patristic distance as a predictor in non-

phylogenetic regressions (e.g., Davies *et al.* 2007; Martin *et al.* 2010).

8. Sister-taxa model fitting

Recently, tools have been described for fitting process-based models to sister taxa

datasets using maximum likelihood (Weir and Wheatcroft 2011; Weir and Lawson 2015). With

these tools, it is possible to test whether models that allow evolutionary rates to vary as a linear

function of a gradient (e.g., whether male plumage coloration varies as a function of the strength

of sexual selection, Seddon et al. 2013) better fit sister-taxa datasets than constant rates models.

When the gradient is the number of sympatric lineages, these models are conceptually similar to

the linear diversity dependent models described above.

Supporting Information for "An assessment of phylogenetic tools for analyzing the interplay between interspecific interactions and phenotypic evolution"

Drury, J.P., Grether, G.F., Garland, Jr., T., & Morlon, H.

1. Supplementary Methods

2. Supplementary Diagram 1

3. Supplementary Figures 1-17

SUPPLEMENTARY METHODS

Simulating Phylogenies Under Varying Biogeographic Scenarios

Joint simulation of trees and biogeographies requires parameterizing (1) the diversification process (the rules defining how new species appear), (2) rates of anagenetic range gains/losses (the rules defining how lineages' ranges change), and (3) cladogenetic range inheritance (the rules defining how two sister lineages divide their ancestral range upon speciation). In each scenario, we evolved species ranges under the DEC+J model (Matzke 2014) across a grid of ten possible regions with equal probability of transitions between regions and only one region occupied at the root of the tree. We simulated the diversification process along each tree with a rate of 0.25 (speciation/lineage/time unit). Anagenetic changes were simulated under two dispersal rates {d (transition/lineage/unit time) = 0.03, 0.06}, chosen from within a range of dispersal values estimated from empirical datasets. When anagenetic dispersal happens, a lineage occupies an additional region chosen at random. Across these two values of "d", we held local extirpation rates constant at e = 0.03 (local extirpation/lineage/unit time), the median value used for simulations in Matzke 2014. In cases where a lineage occupying only one region goes locally extinct, that lineage goes extinct (Matzke 2014).

The cladogenetic events possible in the DEC+J model are—briefly (for a detailed explanation of each of these processes, see Matzke 2014)— sympatric speciation (both daughter lineages keep the one-region ancestral range, parameter "y"), subset sympatric speciation (one of the daughter lineages keeps the ancestral range, the other one inherits a subset of the ancestral range, parameter "s"), vicariance (the two daughter lineages split the ancestral range, parameter "v"), and founder event speciation (one daughter lineages keeps the ancestral range, the other occupies a new region, parameter "j").

As the pool of possible daughter ranges changes depending on the ancestral range (e.g., there are fewer ways in which the ancestral two-region range "AB" can be propagated to daughter lineages than the three-range region "ABC"), cladogenetic range changes are sampled by first assigning each type of cladogenetic change (y,s,v, and j from above) a particular weight. From these weighted ranges, the probability that a daughter inherits a specific range is calculated by dividing the weight assigned to that range by the sum of the weights of each possible daughter range (Matzke 2014). By default, the relative weight of sympatric speciation (range copying), subset sympatric speciation, and vicariance are equal in most implementations of the DEC model (i.e., y=s=v). To generate biogeographic scenarios where recently diverged sister taxa are more likely to be allopatric, we implemented a scenario where the weight of either type of sympatric speciation event is very low {y = s = 0.005*v} as well as under default parameter values {y = s = v}. Across both scenarios, we held the relative weight of founder event speciation constant (j = 0.1, y + s + v = 2.9).

Simulating Character Displacement

Datasets were simulated under Eq. 1 in the main text as follows:
1. At the first time step (the root), the trait value was set to z_0 (= 0 in all cases).
2. Each time step *dt* was set to the total tree height divided by 2500. To complete the simulation along any branch not divisible by this value of *dt*, we set the time step equal to the remainder of the branch length divided by *dt*.

3. During each time step from the root to the tip of the tree, the trait value of lineage *i* is calculated according to Eq 1. For the component of Eq.1 dictating the magnitude of divergence or convergence, between-lineage distances in trait space were calculated based on similarity at time *t-dt*. In simulations of divergent character displacement, if species have identical trait values, the "sign" value is overridden so that species move in the opposite direction of one another in trait space.
4. At a branching event, the trait values of both daughter lineages are set to equal the value of the parent lineage.

For divergent character displacement scenarios, we arbitrarily chose simulation parameter values based on visual inspection of simulated trajectories under different combinations of parameter values. We chose to focus on *m* = 2, and also explored the effect of varying this parameter between 1 & 10, because we could visualize divergence in the realized simulations. For example, for a 20 tip tree simulated with high dispersal and low sympatric speciation:

In convergent character displacement simulations, we likewise chose parameter values based on visual inspection of simulated trajectories under different combinations of parameter values. We chose to focus on $m = -0.25$, and also explored the effect of varying this parameter between -0.1 & -0.50, because we could visualize convergence in the realized simulations and the simulated trait values did not converge across the entire clade. For example, on the same tree as above:

Analytical Methods

Non-phylogenetic regressions: General Linear Models (GLMs) were fitted to simulated datasets using the glm function in R. For character displacement analyses, linear models were fit to pairwise differences in trait data, with sympatry/allopatry as the predictor variable. For interaction analyses, logistic regressions were fit to the simulated species interaction variable, with pairwise difference in the focal trait as the predictor variable.

Mantel tests: We computed raw Mantel tests using mantel.rtest in ade4 (Dray and Dufour 2007), specifying 1000 permutations to assess statistical significance. We computed phylogenetically permuted partial Mantel (pppMantel) tests (Lapointe and Garland 2001) using the script phyloMantel.R from Harmon & Glor (2010), again using 1000 permutations to test for significance.

Phylogenetic linear mixed models (PLMMs): For the character displacement datasets, we fitted PLMMs using asreml-R (Butler et al. 2009), including both sympatry and the patristic distance (calculated using the cophenetic.phylo function in ape [Paradis 2011]) between each pairwise species comparison as fixed effects and including the identity of each lineage and the phylogeny as random effects, following Tobias *et al.* (2014). To assess statistical significance of the fixed effects, we used Wald-type F-tests with Kenward-Rogers adjustments to the denominator degrees of freedom using the wald.asreml function with the option denDF = "numeric" in asreml-R, again following Tobias *et al*. (2014)

For the scenario using traits to predict species interactions, we fit PLMMs using MCMCglmm in R (Hadfield 2010), because the residual maximum likelihood approach used in asreml-R may bias estimates for logistic regressions (Bolker et al. 2009). We used standard inverse-Gamma priors for the fixed and random effects, using the code:

```
prior<-list(G=list(G1=list(V=1,nu=0.002),G2=list(V=1,nu=0.002),G3=list(V=1,nu=0.002)),R=list(V=1,nu=0.02))
```

We ran each fit for between 2 million and 20 million chains based on preliminary assessment of convergence for different tree sizes, varying the burn-in and thinning periods to result in approximately 2000 runs. We visually inspected convergence for a large sample of model fits to make sure that our MCMC parameterization was working well. MCMC fits were very computationally expensive, so we fit the models to a subset of simulated datasets (*n* = 100 per tree size, biogeography, and simulation parameter combination).

For all PLMMs, we randomized the order in which the identity of the lineages was passed to the random effects.

Simulation approach: BM models were first fit to simulated trait data using mvBM in the mvMORPH package (Clavel et al. 2015). Then, 5,000 datasets were simulated using the maximum likelihood estimate of the BM rate parameter and state at the root using fastBM in phytools (Revell 2012). A GLM (as outlined in "non-phylogenetic regressions", the predictor variable for character displacement analyses was biogeographical overlap and the response variable was pairwise trait similarity, and for analyses of species interactions, the pairwise trait similarity was the predictor variable and a binomial variable of species interaction was the response variable) was fit to each of these simulated datasets, and the resulting analysis was considered statistically significant if both (a) the non-phylogenetic test on the raw data was statistically significant and (b) the test statistic from the raw analysis was outside of the 2.5-97.5% quantile interval of test statistics estimated on the simulated datasets.

Sister taxa analyses: for trees of size 150 and larger, we culled sister taxa from the full trees. We then ran non-phylogenetic regressions, PLMMs, the simulation approach and sister taxa GLMs on the culled dataset.

Phenotypic models: We fit all phenotypic models by maximum likelihood. We fit BM and OU models using geiger (Pennell et al. 2014) and the matching competition and two diversity dependent models (with either linear or exponential dependence of sigma on the number of species) using RPANDA (Morlon et al. 2016) in R. For the latter models, we included the biogeography used to simulate the datasets in the model fits, as described in Drury *et al.* (2016). We then compared the relative support of these models using Akaike weights (Burnham and Anderson 2002), since the models are not nested. Since model-fitting is computationally expensive, we fit the process-based models to a subset of datasets simulated on 100 tip trees ($n =$ 100 biogeography and simulation parameter combination).

Additionally, to analyze the effect of m and ψ on rates of evolution, we fit four trait models to the sister-taxa datasets for both divergent and convergent character displacement scenarios using the R package EvoRAG (Weir and Lawson 2015): BM, OU, plus versions of BM and OU that allow the rate of trait evolution to change with the number of sympatric lineages (i.e., "BM_null", "OU_null", "BM_linear", and "OU_linear_beta").

SUPPLEMENTARY REFERENCES

Bolker, B. M., M. E. Brooks, C. J. Clark, S. W. Geange, J. R. Poulsen, M. H. H. Stevens, and J.-S. S. White. 2009. Generalized linear mixed models: a practical guide for ecology and evolution. Trends Ecol. Evol. 24:127–135. Elsevier.

Burnham, K. P., and D. R. Anderson. 2002. Model selection and multimodel inference: a practical information-theoretic approach. 2nd editio. Springer, New York, NY.

Butler, D., B. Cullis, A. Gilmour, and B. Gogel. 2009. ASReml user guide release 3.0. VSN Int. Ltd, Hemel Hempstead, UK.

Clavel, J., G. Escarguel, and G. Merceron. 2015. mvMORPH: an R package for fitting multivariate evolutionary models to morphometric data. Methods Ecol. Evol. 6:1311–1319. Wiley Online Library.

Dray, S., and A. B. Dufour. 2007. The ade4 package: implementing the duality diagram for ecologists. J. Stat. Softw. 22:1–20.

Drury, J., J. Clavel, M. Manceau, and H. Morlon. 2016. Estimating the effect of competition on trait evolution using maximum likelihood inference. Syst. Biol. 65:700–710.

Hadfield, J. D. 2010. MCMC methods for multi-response generalized linear mixed models: the MCMCglmm R package. J. Stat. Softw. 33:1–22.

Harmon, L. J., and R. E. Glor. 2010. Poor statistical performance of the Mantel test in phylogenetic comparative analyses. Evolution (N. Y). 64:2173–2178.

Lapointe, F. J., and T. Garland. 2001. A generalized permutation model for the analysis of cross-species data. J. Classif. 18:109–127. Springer.

Matzke, N. J. 2014. Model selection in historical biogeography reveals that founder-event speciation is a crucial process in island clades. Syst. Biol. 63:951–970.

Morlon, H., E. Lewitus, F. L. Condamine, M. Manceau, J. Clavel, and J. Drury. 2016. RPANDA: an R package for macroevolutionary analyses on phylogenetic trees. Methods Ecol. Evol. 7:589–597. Wiley Online Library.

Paradis, E. 2011. Analysis of Phylogenetics and Evolution with R. Springer, New York, NY.

Pennell, M. W., J. M. Eastman, G. J. Slater, J. W. Brown, J. C. Uyeda, R. G. FitzJohn, M. E. Alfaro, and L. J. Harmon. 2014. geiger v2. 0: an expanded suite of methods for fitting macroevolutionary models to phylogenetic trees. Bioinformatics btu181. Oxford Univ Press.

Revell, L. J. 2012. phytools: An R package for phylogenetic comparative biology (and other things). Methods Ecol. Evol. 3:217–223.

Tobias, J. A., C. K. Cornwallis, E. P. Derryberry, S. Claramunt, R. T. Brumfield, and N. Seddon. 2014. Species coexistence and the dynamics of phenotypic evolution in adaptive radiation. Nature 506:359–363. Nature Publishing Group.

Weir, J. T., and A. Lawson. 2015. Evolutionary rates across gradients. Methods Ecol. Evol. 6:1278–1286.

Supplementary Diagram 1. Analytical methods used for each simulation scenario (see Box 1 for more details).

analytical method	divergent character displacement		convergent character displacement		predicting spp. interactions
	whole tree	sister taxa	whole tree	sister taxa	sympatric spp.
(1) non-phylogenetic regression	✓	✓	✓	✓	✓
(2) Mantel test	✓		✓		
(3) phylogenetically permuted partial Mantel tests	✓		✓		
(4) phylogenetic linear mixed models	✓	✓	✓	✓	✓
(5) phylogenetic simulations	✓	✓	✓	✓	✓
(6) process-based models using maximum likelihood	✓				
(7) sister taxa GLMs		✓		✓	
(8) sister taxa process-based models in EvoRAG		✓		✓	

Supplementary Figure 1. The resulting biogeographic landscape under each of the four biogeographic scenarios. A. The proportion of sister taxa that are living in sympatry varies as a function of both the dispersal rate and the level of sympatric speciation. B. The patristic distances (the branch lengths separating sister taxa) are larger for sympatric taxa, although the magnitude of this difference depends on the biogeographic scenario. C. The mean number of ranges occupied at the tips is higher under high dispersal rates. D. The mean range overlap for sympatric species pairs, calculated as the number of areas two species both occur in, divided by the total number of areas occupied by the lineage with the smaller range.

Supplementary Figure 2. Proportion of statistically significant analyses in datasets simulated under divergent character displacement in biogeographic scenarios with high sympatric speciation rates. A. Results from approaches using data from all pairwise comparisons in a clade, plotted as a function of the phylogeny size and dispersal rate when i-ii. $m = 0$ and $\psi = 0$ (i. all analyses and ii. only analyses returning divergence in sympatry), iii. $m = 2$ and $\psi = 0$, iv-v. $m = 0$ and $\psi = 2$ (iv. all analyses and v. only analyses returning divergence in sympatry), and vi. $m = 2$ and $\psi = 2$. B. Results from analyses of sister-taxa culled from complete phylogenies binned by the number of resulting species pairs, plotted as a function of the number of sister taxa comparisons and dispersal rate when i-ii. $m = 0$ and $\psi = 0$ (i. all analyses and ii. only analyses returning divergence in sympatry), iii. $m = 2$ and $\psi = 0$, iv-v. $m = 0$ and $\psi = 2$ (iv. all analyses and v. only analyses returning divergence in sympatry), and vi. $m = 2$ and $\psi = 2$. For scenarios where $m = 2$, only the proportion of significant results showing divergence are plotted. Dashed horizontal lines represent a Type I error rate of 5%.

Supplementary Figure 3. The effect of repulsion on the magnitude of changes in traits from the root to the tip of phylogenies under different simulation scenarios. To calculate the absolute value of divergence steps (A & B), we recorded the absolute value of the magnitude of the step size resulting from the deterministic repulsion step of Eq. 1 (i.e., $m\left[\sum_{j\neq i}^{n}\mathbf{A}_{i,j} \times sign\left(z_i(t) - z_j(t)\right) \times e^{-\alpha\left(y_j(t) - y_i(t)\right)^2}\right]dt$) at each time step, took the mean during each internode interval for all extant lineages within trees, and then averaged each internode interval across all simulations. Each point thus represents the mean of means and the standard error across 100 phylogenies. The absolute value of the steps pulling trait values toward a stable peak (C & D) similarly represents the absolute value of the magnitude of the step size resulting from the deterministic component of the OU attraction to the peak in Eq. 1 (i.e., $\psi(\theta - z_i(t))dt$).

Supplementary Figure 4. Models fit to sister-taxa datasets generated under divergent character displacement scenarios in EvoRAG corroborate the influence of the number of sympatric lineages on the rate of evolution. A. When $\psi = 0$, BM is the best fitting model, whether A. $m = 0$ or B. $m = 2$. When $\psi = 2$, OU is the best fitting model when C. $m = 0$, whereas a BM model that allows the rate of trait evolution to vary linearly with the number of sympatric lineages is a best fitting model when D. $m = 2$. Plots depict the Akaike weight of each model as a function of both the biogeographic scenario and the number of sister taxa comparisons.

Supplementary Figure 5. The proportion of statistically significant analyses for datasets with divergent character displacement varies as a function of both *m* and *ψ*.

Supplementary Figure 6. As *ψ:m* increases, the ability of all methods to detect character displacement, when present, increases under both low sympatric speciation (A) and high sympatric speciation (B) scenarios.

Supplementary Figure 7. The effect of repulsion on the magnitude of changes in traits from the root to the tip of phylogenies as a function of the ratio ψ:m. As ψ increases relative to m, both the magnitude of divergence steps (panels A-C) and the magnitude of the steps pulling trait values toward the peak (panels D-F) increase. This effect becomes increasingly apparent at higher m values and in biogeographic scenarios generated under high dispersal rates. For a description of the data, see the legend of Fig. S3. In this plot, data are averaged within 10 time bins representing each 10% of the total height of the phylogeny.

Supplementary Figure 8. Proportion of simulated datasets for which a given phenotypic model was chosen using model selection as a function of *m*. We considered a model to be chosen if it had the lowest AICc score among all of the models, and if the AICc score was > 2 units away from BM model. A. Results for datasets simulated without the OU process. B. Results from datasets simulated with $\psi = 2$.

Supplementary Figure 9. The rate parameter *r* of the DDexp+GEO model is increasingly negative with increasing *m* values when $\psi = 0$, yet is positive when $\psi > 0$.

Supplementary Figure 10. Akaike weights for each trait model fit to simulated datasets in biogeographic scenarios with high sympatric speciation rates as a function of *m*. A. When OU is absent, BM is the best-fit model when *m* = 0, and the matching competition model with biogeography is the best model when competitive divergence is present. B. When OU is present, OU is the best-fit model when *m* = 0, and the diversity-dependent exponential model with biogeography is the best model when competitive divergence is present and *ψ:m* is relatively high.

Supplementary Figure 11. As ψ:m increases, the DDexp + GEO model is increasingly better fit to datasets simulated with character displacement.

Supplementary Figure 12. Proportion of statistically significant analyses in datasets simulated under convergent character displacement in biogeographic scenarios with high sympatric speciation rates. A. Results from approaches using data from all pairwise comparisons in a clade, plotted as a function of the phylogeny size and dispersal rate when i-ii. $m = 0$ and $\psi_{resource} = 0$ (i. all analyses and ii. only analyses returning convergence in sympatry), iii. $m = -0.25$ and $\psi_{resource} = 0$, iv-v. $m = 0$ and $\psi_{resource} = 2$ (iv. all analyses and v. only analyses returning convergence in sympatry), and vi. $m = -0.25$ and $\psi_{resource} = 2$. B. Results from analyses of sister-taxa culled from complete phylogenies binned by the number of resulting species pairs, plotted as a function of the number of sister taxa comparisons and dispersal rate when i-ii. $m = 0$ and $\psi_{resource} = 0$ (i. all analyses and ii. only analyses returning convergence in sympatry), iii. $m = -0.25$ and $\psi_{resource} = 0$, iv-v. $m = 0$ and $\psi_{resource} = 2$ (iv. all analyses and v. only analyses returning convergence in sympatry), and vi. $m = -0.25$ and $\psi_{resource} = 2$. For scenarios where $m = -0.25$, only the proportion of significant results showing convergence are plotted. Dashed horizontal lines represent a Type I error rate of 5%.

Supplementary Figure 13. The effect of convergence on the magnitude of changes in traits in simulations, plotted as a function of stabilizing selection in the resource-use trait ($\psi_{resource}$) and biogeography. With $\psi_{resource} = 2$, the magnitude of convergence steps is higher, and this is particularly true for high dispersal biogeographies. Data generated as described in the legend of Fig. S3.

Supplementary Figure 14. Models fit to sister-taxa datasets generated under convergent character displacement scenarios in EvoRAG. A. When $m = 0$, BM is the best fitting model, whether A. $\psi_{resource} = 0$ or C. $\psi_{resource} = 2$. When $m = -0.25$, OU is the best fitting model when C. $\psi_{resource} = 0$ and D. $\psi_{resource} = 2$, but an OU model with evolutionary rates that change linearly as a function of the number of sympatric lineages (OU linear) is also a strong model. Plots depict the Akaike weight of each model as a function of both the biogeographic scenario and the number of sister taxa comparisons.

Supplementary Figure 15. The proportion of statistically significant analyses for datasets with convergent character displacement varies as a function of both *m*, and is generally lower in the absence of a pull toward a peak (i,iii) compared to cases where data were simulated with a pull toward a peak (ii,iv).

Supplementary Figure 16. Proportion of statistically significant analyses in datasets with interactions simulated under a simple phenotype matching process in biogeographic scenarios with high sympatric speciation rates. Results from analyses where the focal trait was simulated under A. BM or B. OU, plotted as a function of the dispersal rate when i. b_1 (the simulation coefficient determining the relationship between the interaction and the measured trait) = 0, b_2 (the simulation coefficient for an unmeasured trait) = -4, and ψ = 2, ii. b_1 = -4, b_2 = 0, and ψ = 2, iii. b_1 = 0, b_2 = -4, and ψ = 0, and iv. b_1 = -4, b_2 = 0, and ψ = 0.

Supplementary Figure 17. The power to detect a causal relationship between a species interaction and similarity in a trait value is not affected greatly by varying the coefficient determining the strength of this relationship, regardless of whether the OU process is absent (i, iii) or present (ii, iv) in the measured trait. In these simulations, tree size was held constant at 100 species.