Microsaccadic information sampling provides *Drosophila* hyperacute vision

Mikko Juusola[1,2]*‡, An Dau[2]‡, Zhuoyi Song[2]‡, Narendra Solanki[2], Diana Rien[1,2], David Jaciuch[2], Sidhartha Dongre[2], Florence Blanchard[2], Gonzalo G. de Polavieja[3], Roger C. Hardie[4] and Jouni Takalo[2]

[1]National Key laboratory of Cognitive Neuroscience and Learning, Beijing, Beijing Normal University, Beijing 100875, China

[2]Department of Biomedical Science, University of Sheffield, Sheffield S10 T2N, UK

[3]Champalimaud Neuroscience Programme, Champalimaud Center for the Unknown, Lisbon, Portugal

[4]Department of Physiology Development and Neuroscience, Cambridge University, Cambridge CB2 3EG, UK

*Correspondence to: m.juusola@sheffield.ac.uk

‡ Equal contribution

Abstract: Small fly eyes should not see fine image details. Because flies exhibit saccadic visual behaviors and their compound eyes have relatively few ommatidia (sampling points), their photoreceptors would be expected to generate blurry and coarse retinal images of the world. Here we demonstrate that *Drosophila* see the world far better than predicted from the classic theories. By using electrophysiological, optical and behavioral assays, we found that R1-R6 photoreceptors' encoding capacity *in time* is maximized to high-frequency saccadic light changes. Whilst *over space*, R1-R6s resolve moving objects at saccadic speeds beyond the predicted motion-blur-limit. Our results show how refractory phototransduction and rapid photomechanical photoreceptor contractions jointly sharpen retinal images *in space-time*, enabling hyperacute vision, and explain how such microsaccadic information sampling exceeds the compound eyes' optical limits.

One Sentence Summary: Microsaccadic information sampling expands compound eyes' visual capabilities by encoding space in time.

The spatial vision of insects is thought to be poor, limited by their compound eyes' low optical resolution and motion blur during fast self-motion (*1-4*). Here, we challenge these classic ideas by showing how fast microsaccadic information sampling, combined with rapid autonomous photoreceptor contractions, improves vision in *Drosophila* R1-R6 photoreceptors. Our intracellular recordings and biophysically-realistic model simulations reveal that their responses to high-contrast bursts (periods of rapid light changes followed by quiescent periods), which resemble high-frequency saccadic modulation, capture 2-to-4-times more information in time than the previous maximum estimates (*5-7*). This improvement largely results from interspersed "fixation" intervals, which relieve photoreceptors' refractory microvilli (sampling units) to

sample more information from phasic light changes. Remarkably, over space, our intracellular recordings, high-speed microscopy and modeling further reveal how photomechanical photoreceptor contractions (*8*) work together with refractory sampling to improve spatial acuity. We discover that by actively modulating light input and photoreceptor output, these processes reduce motion blur during saccades and adaptation during gaze stabilization, which otherwise could fade vision (*9, 10*). The resulting phasic responses sharpen retinal images by highlighting the times when visual objects cross a photoreceptor's receptive field, thereby encoding space in time (*cf.* (*11-14*)). Incorporation of this novel opto-mechano-electric mechanism into our 'microsaccadic sampling' model predicts that *Drosophila* can see >4-fold finer details than their eyes' spatial sampling limit – a prediction directly confirmed by optomotor behavior experiments. By demonstrating how fly photoreceptors' fast microsaccadic information sampling provides hyperacute vision, these results revolutionize our understanding of spatial vision.

A photoreceptor counts photons and integrates these counts to an estimate, a macroscopic voltage response, of light changes within its receptive field. In a *Drosophila* R1-R6 photoreceptor, this estimate is counted by 30,000 microvilli, which form its light-sensor, the rhabdomere (**Supplementary Text 1**). Each microvillus is essentially a photon sampling unit, which contains all the elements of the phototransduction cascade and can transduce a photon's energy to a unitary response (quantum bump or sample) (*5-7*, *15*). But following each bump, each light-activated microvillus becomes refractory (*5, 6*) for 50–300 ms. Therefore, with brightening light, a photoreceptor's sample rate gradually saturates, as fewer microvilli are available to generate bumps. Although refractory sampling makes photoreceptors imperfect photon counters, it benefits vision by representing a fast automatic adaptation mechanism, reducing sensitivity in proportion to background intensity (*5, 6*), whilst accentuating responses to contrast changes (*5*).

A well-known trade-off of fast adaptation is that it causes perceptual fading during fixation (*9, 10*), and to see the world requires motion or self-motion: body, head and eye movements (*16-21*), which remove adaptation. But it remains unclear whether or how the fly phototransduction biophysics is tuned to visual behaviors to see the world better. To start unravelling these questions, we first surveyed what kind of stimuli drove their information transfer maximally, ranging from high-frequency bursts (>10x briefer than *Drosophila*'s normal head/body-saccades (*22, 23*)) to Gaussian white-noise (GWN). These stimuli tested systematically different contrast and bandwidth patterns over R1-R6s' diurnal encoding gamut.

High-frequency "saccades" maximize encoding

Intracellular recordings (**Fig. 1A**) revealed that photoreceptors responded most vigorously to high-contrast bursts, which contained high-frequency saccade-like events with darker intervals. **Figure 1B** shows the averages (signals; thick) and individual responses (thin) of a typical R1-R6, grouped by the stimulus bandwidth and mean contrast. For all the bandwidths (columns), the responses increased with contrast, while for all the contrasts (rows), the responses decreased with the increasing bandwidth (left). Therefore, the slowest high-contrast bursts (red; top-left), with the longest darker intervals relieving most refractory microvilli, evoked the largest peak-to-peak responses (43.4 ± 5.6 mV; mean ± SD, n = 16 cells; **fig. S1**). Whereas the fastest low-contrast GWN (blue; bottom-right), which would keep more microvilli refractory, evoked the smallest responses (3.7 ± 1.1 mV; n = 4).

Fig. 1. **Photoreceptors respond best to high-frequency saccadic bursts**. (**A**) Schematic of intracellular recordings to repeated bursty light intensity time series (20 Hz bandwidth). Responses changed little (minimal adaptation) during bursts. (**B**) Testing a R1-R6 photoreceptor's diurnal encoding gamut. Means (thick traces) and 20 individual responses (thin; near-perfectly overlapped) to 20 different stimuli; each with a specific bandwidth (columns: from 20 Hz, red, to 500 Hz, blue) and mean contrast (rows). By reducing the background (BG) of Gaussian white-noise stimuli (GWN; 2-unit peak-to-peak modulation) from bright (1.5-unit, bottom) to dark (0-unit, top), halved their modulation, generating bursts of increasing contrast:

the lower the BG, the higher the contrast. Left-top: responses from (**A**). Yellow box: maximum information responses. Arrows: dark intervals. Because of half-Gaussian waveforms, bursts carried fewer photons (**fig. S4**). Their larger responses resulted from dark intervals, which rescued refractory microvilli for transducing high-frequency (1-20 ms) saccadic photon surges into elementary responses efficiently. Recordings are from the same photoreceptor. Vertical dotted rectangle (orange square) and horizontal rectangle (black circle): responses for contrast and bandwidth analyses in **Fig. 2A**. Similar R1-R6 population data is in **fig. S1**.

Information theoretical analysis (**Fig. 2** and **fig. S2**) further indicated that the response differences largely reflected differences in their quantum bump counts. The maximum signal power spectra to bursty stimuli could be up to ~6,000-times larger than that of the noise, which was effectively stimulus-invariant (**fig. S3A**). Because the noise power spectrum largely represents the average quantum bump's frequency composition (*5, 7, 24*), the bumps adapted to a similar size. Thereby, the larger responses simply comprised more bumps. Moreover, with Poisson light statistics, the response precision - how well it estimated photon flux changes - should increase with the square root of bump count until saturation; when more microvilli remained refractory (*5*). Accordingly, signaling performance (**Fig. 2A** and **C**) increased both with the stimulus bandwidth (left) and contrast (right), until these became too fast to follow. Information transfer peaked at 100 Hz bursts, which allocated the R1-R6's limited bandwidth and amplitude range near-optimally, generating the broadest frequency (**Fig. 2A** and **fig. S2B**) and (Gaussian) voltage distributions (**Fig. 2B** and **fig. S2B**).

Thus, with the right mixture of high-frequency bright "saccades" (to maximally activate microvilli) and darker "fixation" intervals (to recover from refractoriness), information transfer approached the capacity (*25*), the theoretical maximum, where every symbol of a message is transmitted equally often (**Fig. 2C** and **fig. S2C**). Remarkably, this performance (610-850 bits/s) was 2-to-4-times of that for GWN (200-350 bits/s), which has often been used for characterizing maximal encoding (*7, 26*), and twice of that for rich naturalistic stimuli (380-510 bits/s) (*5*) (**fig. S4**). GWN, especially, lacks longer darker events, making microvilli refractory (*5*) with fewer sampled photons limiting information transfer (**Supplementary Texts 2-3**).

Fig. 2. **High-frequency saccadic bursts drive maximal encoding**. A R1-R6's information transfer to high-frequency 100 Hz bursts exceeded 2-to-4-times the previous estimates. (**A**) Response signal-to-noise ratio (SNR, left) to 20 (red), 100 (yellow) and 500 Hz (blue) bursts, and to 50 Hz bandwidth stimuli of different contrasts (right); data from **Fig. 1**. SNR increased with contrast (right), reaching the maximum (~6,000) for 20 Hz bursts (left, red) and the broadest frequency range for 100 Hz bursts (yellow). (**B**) Skewed bursts drove largely Gaussian responses (exception: 20 Hz bursts, red), with 100 Hz bursts evoking the broadest amplitude range (yellow). (**C**) Information transfer peaked for 100 Hz stimuli, irrespective of contrast (or BG; left), having the global maximum of ~850 bits/s (capacity, infomax) for the high-frequency saccades. (**D**) Encoding efficiency, the ratio between input and output information (R_{output}/R_{input}), was >100% for 20 Hz bursts. Extra information came from the neighboring cells. R_{input} at each BG was determined for the optimal mean light intensity, which maximized a biophysically realistic photoreceptor model's information transfer (**Supplementary Text 2**). Encoding efficiency fell with stimulus bandwidth but remained more constant with contrast. Population dynamics are in **figs S2** and **S7**.

Simulations reveal network contribution

These findings were largely replicated by stochastic simulations (**fig. S5**). A biophysically realistic photoreceptor model, which contains 30,000 microvilli (*6*), sampled light information much like a real R1-R6, generating authentic responses to all the test stimuli. Yet, markedly, the model lacked the intracellular pupil (or any adaptation), which protects microvilli from saturation (*5, 27*), and network connections (*28-30*). In real photoreceptors, the pupil screens off excess light to maximize information transfer (*5, 6*). Similarly, in the simulations, the mean light intensity of each stimulus was optimized (**Supplementary Text 2**) for maximum information (**fig. S6**). At its peak, the model transferred 633 ± 20 bits/s (mean ± SD) for 100 Hz bursts of $8{\times}10^5$ photons/s, with further brightening reducing information as more microvilli became refractory. This performance matches that of many real R1-R6s (**fig. S2C**), but is ~200 bits/s less than in some recordings (**Fig. 2c**). The real R1-R6s, on balance, receive extra information from

their neighbors (*28*, *29*), which through superposition (*30*) sample information from overlapping receptive fields. Moreover, as their rhabdomere sizes and connectivity vary systematically (*28*), each R1-R6 receives different amounts of information (*29*) (**fig. S7**). Here, R6s, with large rhabdomeres and gap-junctions to R7/R8, should receive the most (29), suggesting that the best performing cells (*e.g.* **Figs 1** and **2**) might be of the R6-type.

Encoding efficiency for the different stimuli (**Fig. 2d** and **fig. S2D**) was determined as the ratio between the related photoreceptor and light information rates (R_{output} / R_{input}); with R_{input} estimated from the simulated Poisson stimulus repeats, which maximized information in R1-R6 model output (**Supplementary Text 2**). The estimates for both the recordings (**Fig. 2d** and **fig. S2D**) and simulations (**fig. S6D**) indicated that encoding weakened with the increasing stimulus bandwidth (left) but less so with contrast (right). However, as predicted, some recordings showed >100% efficiency for 20 Hz bursts, presumably due to their extra network information.

A locomoting *Drosophila* generates ~1-5 head/body-saccades/s, which direct its gaze in high velocities to the surroundings (*22*, *23*). Here, our recordings and simulations suggest that the refractoriness in R1-R6s' phototransduction, together with network inputs, might be tuned for capturing information during such fast light changes in time. We next asked how R1-R6s see saccadic light changes over space.

Testing acuity at saccadic velocities

A *Drosophila*'s head/body-saccades generate fast phasic photoreceptor movements, which ought to blur retinal images (*22*, *23*). Moreover, saccades provide no distance information (*31*) because objects, near and far, would move across the retina with the same speed. Therefore, it has been long thought that visual information is mostly captured during translational motion and gaze-fixation, and less during saccades.

To test this hypothesis, we reasoned that object motion and self-motion shape a photoreceptor's light input the same way. Thus, the influence of eye movements (and motion blur) on a R1-R6's ability to resolve objects could be measured in experiments, where, instead of moving the eye, the objects were moved over its stationary receptive field (**Fig. 3A**; **Supplementary Texts 4-6**).

Using this approach, we recorded individual R1-R6s' voltage responses (**Fig. 3B**; black traces) to a pair of bright dots (each 1.7° in size and 6.8° apart, as seen by the fly), moving at constant speed across their receptive field in front-to-back direction. The movement was either fast (205 °/s) or double-fast (409 °/s), in terms of a walking *Drosophila*'s head/body saccadic velocity range (200-800 °/s) (*23*), and it was presented against a dark or lit background. Importantly, the dots' angular separation was less than the half-width of a R1-R6's receptive field (**Fig. 3C**) at the two backgrounds ($\Delta\rho_{dark}$ = 9.47 ± 1.57°, n = 19 cells; $\Delta\rho_{light}$ = 7.7 ± 1.27°, n = 6; mean ± SD) and 1.5-times the average interommatidial angle ($\Delta\varphi$ ~4.5°), which should determine *Drosophila*'s visual acuity (*32*). Thus, these fast-moving point objects tested the theoretical limit of what a R1-R6 should be able to resolve. We further estimated each cell's respective impulse response (**Supplementary Text 6**). Then following the classic theory of compound eyes' resolving power (*2-4*), we calculated each R1-R6's expected voltage output to the moving dots by convolving its impulse response with its measured dark- or light-adapted receptive field. These Volterra-model (*3*) predictions (**Figs 3B-C**; blue) were then compared to the actual recordings (black).

Fig. 3. **Photoreceptors resolve dots at saccadic velocities far better than the classic models**. (**A**) 25-light-point stimulus array centered at a R1-R6's receptive field. Each tested photoreceptor saw two bright dots, 6.8^o apart, travelling fast (205 ^{o}/s) or double-fast (409 ^{o}/s) in front-to-back direction. (**B**) Responses (black), both at dark (left) or illuminated backgrounds (right), characteristically showed two peaks. In contrast, the corresponding classic model simulations (blue) rarely resolved the dots. (**C**) In the simulations, each photoreceptor's receptive field (or its Gaussian fit) was convolved with its impulse response (1st Volterra kernel). The resolvability, *D*, of the recordings and simulations, was determined by Raleigh criterion. (**D**) Recordings outperformed simulations. (**E**) hdc^{JK910} R1-R6s (red), which lacked the neurotransmitter histamine, and so network modulation, resolved the dots as well as the wild-type, indicating that the recordings' higher resolvability was intrinsic and unpredictable by classic deterministic models (**Supplementary Text 1**). (**F**) Normalized responses of a typical R1-R6 to a bright dot, crossing its receptive field in front-to-back or back-to-front at different speeds. Responses to back-to-front motions rose and decayed earlier, suggesting direction-selective encoding. (**G**) This lead at the half-maximal values was 2-10 ms. Details in **Supplementary Texts 4** and **6**.

Eyesight beyond the motion blur-limit

Remarkably in all these tests, the recordings showed distinctive responses to the two dots (Fig. 3B), as two peaks separated by a trough. The relative magnitude of this amplitude separation was quantified as resolvability, using the Raleigh criterion (*3*) (**Fig. 3C**). However, in marked contrast, the model predictions failed to resolve the double-fast dots, instead blurring into one broad response in both adapting states (**Fig. 3D**; blue vs. black bars, respectively). The

predictions for the fast dots were also poorer than the measured responses. Thus, a photoreceptor's real resolving power was significantly better and less affected by motion blur than predicted by classic theory (**Supplementary Text 6**).

We next asked whether this better-than-expected resolving power resulted from synaptic interactions (*30*, *33*) by using hdc^{JK910} mutants (**Fig. 3E**, red traces), in which photoreceptors lacked their neurotransmitter, histamine (*34*) (**Supplementary Texts 4-6**). Because hdc^{JK910} R1-R6s cannot transmit information to their post-synaptic targets (*35*) (LMCs, which initiate the motion detection pathways (*36*), and the amacrine cells), neither could these photoreceptors receive any light-driven interneurone feedback modulation (*35*). Therefore, if the synaptic interactions improved the wild-type output to the moving dots, then hdc^{JK910} R1-R6s, which lacked these interactions, should show diminished resolvability. But this was never observed. Instead, we found that hdc^{JK910} R1-R6s resolved the dots at least equally well as the wild-type (**Fig. 3D**, red). Thus, high acuity did not result from synaptic inputs but was intrinsic to photoreceptors.

We further discovered that R1-R6 recordings often showed phasic directional selectivity (**Fig. 3F**), with the responses rising and decaying faster to back-to-front than to front-to-back moving dots (**Fig. 3G**). We asked whether these lag-time differences originated from asymmetric photomechanical photoreceptor contractions. Namely, atomic-force microscopy has revealed minute (<275 nm) vertical movements on the surface of dissected *Drosophila* eyes, generated by contraction of individual microvilli as PIP_2 is hydrolyzed from the inner leaflet of the lipid bilayer (*8*). Here, instead, we reasoned that if the ommatidium lenses were effectively fixed and R1-R8s levered to the retinal matrix, the contractions (**Movie S1**) might be larger *in situ*, moving and shaping the photoreceptors' receptive fields along some preferred direction. Such mechanical feedback could then reduce light input to R1-R8s, making it more transient and directional.

Microsaccadic sampling of retinal images

To probe this idea, we recorded *in vivo* high-speed videos of photoreceptor rhabdomeres (viewed by optical neutralization of the cornea) inside the eyes reacting to blue-green light flashes (470 + 560 nm) (**Fig. 4A**). The recordings were performed under far-red (>720 nm) illumination, which is nearly invisible to *Drosophila* (**Supplementary Text 7**).

We found that 8-20 ms after a flash the rhabdomeres, which directly faced the light source at the image center, suddenly surged frontwards (**Figs 4B**). These local movements were faster and larger the brighter the flash (**Figs 4C**), and reached their intensity-dependent maxima (0.2-1.2 μm; **Fig 4D**) in 70-200 ms, before returning slower to their original positions (**Supplementary Text** 7 analyses hdc^{JK910}-rhabdomere responses). Because the mean R1-R6 rhabdomere tip diameter is ~1.7 μm (**fig. S7B**), a bright flash could shift it more than its half-width sideways. Consequently, the fast rhabdomere movements, whilst still ~3-times slower than their voltage responses (**Fig. 4C**, wine), adapted photoreceptors photomechanically by shifting their receptive fields by 0.5-4.0^o, away from directly pointing to the light source.

Fig. 4. **Microsaccadic rhabdomere contractions and refractory photon sampling improve visual resolution of moving objects**. (**A**) High-speed videos showed fast lateral R1-R7 rhabdomere movements to blue/green flashes, recorded under far-red illumination that *Drosophila* barely saw (*29*). (**B**) Rhabdomeres moved inside those 7 ommatidia (up-right: their pseudopupil), which faced and absorbed the incident blue/green light, while the others reflected it. Rhabdomeres moved frontwards 8-20 ms after a flash onset, being maximally displaced 70-200 ms later, before returning. (**C**) Movements were larger and faster the brighter the flash, but slower than R1-R6s' voltage responses. (**D**) Movements followed R1-R6s' logarithmic light-sensitivity relationship. Concurrently, given the ommatidium optics (*32*, *39*), R1-R6s' receptive fields (RFs) shifted by 0.5-4.0°. (**E**) Rhabdomeres moved along the eye's horizontal (red) axis, with little vertical components (black), adapting to ~30% contractions in ~10 s during 1 s repetitive flashing. (**F**) Moving ommatidium structures. Cone and pigment cells, linking to the rhabdomeres by adherens-junctions (*37*), formed an aperture smaller than the rhabdomeres' pseudopupil pattern. Rhabdomeres moved ~2-times more than this aperture, and ~10-times more than the lens. (**G-H**) Simulated light inputs and photoreceptor outputs for the classic theory and new "microsaccadic sampling"-hypothesis when two dots cross a R1-R6's RF (i) front-to-back at

Video footage at different eye locations indicated that light-activated rhabdomeres moved in back-to-front direction along the eye's equatorial plane (**Fig. 4E**, red; **Movie S2**), with little up-down components (black). Therefore, as each ommatidium lens inverts projected images, the photoreceptors' receptive fields should follow front-to-back image motion. This global motion direction, which corresponds to a forward locomoting fly's dominant horizontal optic flow field, most probably explains the phasic directional selectivity we found to opposing image motions (**Fig. 3F**; **Supplementary Text 8**). Thus, the responses to back-to-front moving dots were faster because the dots entered and exited each contracting photoreceptor's front-to-back moving receptive field earlier; whereas the dots moving in the opposite direction stayed slightly longer inside each receptive field. Video analyses further revealed that the first rhabdomere movement was the largest (**Fig. 4E**), but 1 s dark intervals, as used in **Figure 3**, could resensitise the photoreceptors for the next (~0.5 μm) movements. Even <100 ms dark periods rescued noticeable contractility (**fig. S3E**).

To inspect how rhabdomere contractions affected the cornea lens system's image projection, we scanned ommatidia by z-axis piezo steps, with the imaged focal plane travelling down from the lens surface into rhabdomeres (**Fig. 4F**; **Movie S3**), delivering flashes at predetermined depths. Crucially, we found that the ommatidium lens stayed nearly still, while specific pigment and cone cells, which are connected to the rhabdomere tips by adherens junctions (*37*), formed a narrow aperture that moved with the rhabdomeres but only half as much. Thus, as the lens system was immobile but the aperture and sensors (rhabdomeres) underneath swung differentially, the light input to the moving rhabdomeres was shaped dynamically. This implied that, during saccadic image motion, R1-R6s' receptive fields might not only move but also narrow (**Supplementary Texts 7-8**).

Essentially, light input to a R1-R6 was modulated by the photoreceptor itself (**Fig. 4F**). To estimate how these photomechanics influenced encoding, we implemented them in stochastic model simulations. We then compared how the predicted light inputs of the classic theory (**Fig. 4G**) and the new "microsaccadic sampling"-hypothesis (**Fig. 4H**) would drive R1-R6 output during the saccadic dot stimulation. In the classic theory, the rhabdomere is immobile (ii). Therefore, light input of two moving dots was a convolution of two broad Gaussians (i) that fused together (iii), making them irresolvable to phototransduction (iv); this also flawed the Volterra-models (**Fig. 3**). In the new hypothesis, instead, as microvilli became light-activated (ii), the rhabdomere contracted away from the focal point, and then returned back more slowly, recovering from refractoriness. And because its receptive field moved and narrowed concurrently (its acceptance angle, $\Delta\rho$, halved to 4.0^o), the light input of two moving dots transformed into two intensity peaks (iii), in which time-separation was enhanced by the rhabdomere's asymmetric motion. Crucially, with such input driving the refractory photon sampling model, its output (iv) closely predicted the responses to the two moving dots (**Fig. 4I** and **fig. S8**). Interestingly, early behavioral experiments in bright illumination (*38*) suggested similarly narrow R1-R6 acceptance angles (~3.5^o).

Fig. 5. "**Microsaccadic sampling" hypothesis predicts visual hyperacuity**. (**A**) Simulated R1-R6 output to two dots, at different distances apart, crossing the photoreceptor's receptive field at different speeds. The yellow backgrounds indicate those inter-dot-distances and speeds, which evoked two-peaked responses. The prediction is that the real R1-R6s could resolve (and *Drosophila* distinguish) these dots as two separate objects, whereas those on the white backgrounds would be seen as one object. The simulations were generated with our biophysically realistic R1-R6 model (*5*, *6*, *40*), which now included the estimated light input modulation by photomechanical rhabdomere movements (**Fig. 4H**). (**B**) The resulting object resolution/speed heat-map, using the Raleigh criterion, *D* (**Fig. 3C**), shows the stimulus/behavioral speed regime where *Drosophila* should have hyperacute vision. Thus, by adjusting its behavior (from gaze-fixation to saccadic turns) to changing surroundings, *Drosophila* should see the world better than its compound eye's optical resolution. (**C**) Intracellular R1-R6 responses resolved the two dots, which were less that the interommatidial angle ($\Delta\varphi = 4.5^{\circ}$) apart when these crossed the cell's receptive field at the predicted speed range. Arrows indicate the two response peaks corresponding to the dot separation. *Cf.* **fig. S9**; details in **Supplementary Texts 7-8**.

From microsaccades to hyperacuity

Because of the close correspondence between R1-R6 recordings and the new hypothesis (**Supplementary Texts 6-9**), we used it further to predict whether *Drosophila* possessed hyperacute vision (**Fig. 5**). We asked whether "saccade-fixation-saccade"-like behaviors, when linked to refractory photon sampling and photomechanical photoreceptor contractions, allowed

encoding in time finer spatial details than the compound eye's optical limit ($\Delta\varphi$ ~4.5°). R1-R6 output was simulated to two bright dots 1-4° apart, crossing its receptive field at different speeds at 25°C.

We found that if the dots, or a *Drosophila*, moved at suitable speed, a photoreceptor should resolve them well (**Fig. 5A**), with this performance depending upon the inter-dot-distance. When the dots/eye moved at 10 °/s, a R1-R6 may capture image details at 1° resolution. But with slower movement (≤2.5 °/s), adaptation should fuse the dots together, making them neurally unresolvable. Conversely, 3°-apart-dots should be seen at 5-100 °/s speeds and 4°-apart-dots even during fast saccades (200-300 °/s). Thus, the "microsaccadic sampling"-hypothesis implied that *Drosophila* had hyperacute vision over a broad speed range (**Fig. 5B**), and through its own self-motion, could adjust the resolution of its neural images. Further comparisons of model outputs with and without refractoriness indicated that it extends the speed range of hyperacute vision (**Supplementary Text 8**). Again, intracellular recordings corroborated these predictions (**Fig. 5C** and **fig. S9**), demonstrating how acuity could be enhanced by encoding space in time.

Fig. 6. **Optomotor behavior in a flight simulator system confirms hyperacute vision**. Classic open-loop experiments using high-resolution panoramas. (**A**) 360° hyperacute black-and-white bar panorama, with 1.16° or 2.88° wavelengths (= 0.58° and 1.44° inter-bar-distance), rotated counterclockwise and clockwise (grey, arrows) around a tethered fly, with a torque meter measuring its optomotor responses. (**B**) Controls: the same flies' optomotor responses to white (no bars) and wide-bar (14.4° wavelength) rotating panoramas. (**C**) Every *Drosophila* responded to the hyperacute panoramas (wavelength < interommatidial angle, $\Delta\varphi$, yellow area; **fig. S9**), but not to the white panorama (orange), which thus provided the recording noise level. The flies optomotor responses were the strongest to the wide-bar panorama (perception index = 1). As the flies' optomotor responses followed the rotation directions consistently, irrespective of the tested bar wavelengths, the hyperacute visual panorama did not generate perceptual aliasing. Details in **Supplementary Text 10**.

Lastly, we investigated the spatial resolution of *Drosophila* vision through their optomotor behavior in a conventional flight simulator system, which used brightly-lit high-resolution prints for panoramic scenes (**Fig. 6**; **Supplementary Text 10**). We asked whether tethered *Drosophila* possessed motion vision hyperacuity by recording their yaw torque (optomotor response) to vertical black-and-white bar panoramas with <4.5^o wavelengths, which slowly rotated (45 ^{o}/s) to clockwise and counterclockwise. We found that every tested fly responded down to ~1^o panoramic bar resolution (**Fig. 6A** and **fig. S10**) with their responses becoming smaller the finer its bars. Importantly, because these responses consistently followed the rotation direction changes, they were not caused by aliasing. Thus, optomotor behavior verified that *Drosophila* sees the world at least in 4-fold finer detail than what was previously thought.

Discussion

We have provided deep new insight into spatiotemporal information processing in *Drosophila* R1-R6 photoreceptors and animal perception in general. Our results indicate that the dynamic interplay between saccades and gaze-fixation is important for both the maintenance and enhancement of vision. This interplay, which is commonly observed in locomoting *Drosophila* (*23*), makes light input to photoreceptors bursty. We showed that high-frequency saccadic bursts maximize photoreceptors' information capture in time, and provided evidence that such encoding involves four interlinked mechanisms. Light input is first regulated by two processes inside photoreceptors: slower screening pigment migration (intracellular pupil, 1-10 s) and much faster photomechanical rhabdomere contractions (0.01-1s). Their light input modulation has low noise (**fig. S3**), enabling refractory photon sampling by microvilli to enhance information intake in phasic stimulus components. Finally, asymmetric synaptic inputs from the network differentiate individual R1-R6 outputs. Remarkably, over space, these mechanisms further sharpen neural resolvability by ~4-fold below the theoretical limit of the compound eye optics, providing hyperacute vision. Further analyses imply that these mechanisms with systematic rhabdomere size variations combat aliasing (**Supplementary Texts 2** and **5**). Thus, with microsaccadic sampling, a fly's behavioral decisions govern its visual information/acuity trade-off. To see the finest image details it should scan the world slowly, which probably happens during normal gaze fixation. But gaze fixation cannot be too slow; otherwise, adaptation would fade vision. Conversely, by locomoting faster, in a saccadic or bursty fashion, visual information capture in time is increased, while surprisingly little spatial details about its surroundings would be lost.

Our results make a general prediction about the optimal viewing strategy for maximizing information capture from the world. Animals should fixate gaze on darker features, as this resensitises photoreceptors by relieving their refractory sampling units (*e.g.* microvilli). And then, rapidly move gaze across to brighter image areas, as saccadic crossings over high-contrast boundaries enhance information intake by increasing photoreceptors' sample (quantum bump) rate changes/time (**Supplementary Text 9**). Given the high occurrence of eye/head-saccades in animals with good vision (*31*), it seems plausible that their photoreceptors could also have adapted encoding dynamics to quicken response modulation, reducing motion blur. Therefore, if information sampling biophysics in rods and cones were matched to microsaccadic eye movements, this could provide a mechanistic explanation to the old paradox: how saccadic gaze fixation provides stable perception of the world, while curtailing motion blur effects.

References:

1.   H. B. Barlow, The Size of Ommatidia in Apposition Eyes. *J Exp Biol* **29**, 667 (1952).

2.   M. F. Land, Visual acuity in insects. *Annu Rev Entomol* **42**, 147 (1997).

3.   M. Juusola, A. S. French, Visual acuity for moving objects in first- and second-order neurons of the fly compound eye. *J Neurophysiol* **77**, 1487 (Mar, 1997).

4.   M. V. Srinivasan, G. D. Bernard, Effect of Motion on Visual-Acuity of Compound Eye - Theoretical-Analysis. *Vision Res* **15**, 515 (1975).

5.   Z. Song, M. Juusola, Refractory sampling links efficiency and costs of sensory encoding to stimulus statistics. *J Neurosci* **34**, 7216 (May 21, 2014).

6.   Z. Song et al., Stochastic, Adaptive Sampling of Information by Microvilli in Fly Photoreceptors. *Curr Biol* **22**, 1371 (Aug 7, 2012).

7.   M. Juusola, R. C. Hardie, Light adaptation in *Drosophila* photoreceptors: I. Response dynamics and signaling efficiency at 25 degrees C. *J Gen Physiol* **117**, 3 (Jan, 2001).

8.   R. C. Hardie, K. Franze, Photomechanical Responses in *Drosophila* Photoreceptors. *Science* **338**, 260 (Oct 12, 2012).

9.   R. W. Ditchburn, B. L. Ginsborg, Vision with a Stabilized Retinal Image. *Nature* **170**, 36 (1952).

10.   L. A. Riggs, F. Ratliff, The Effects of Counteracting the Normal Movements of the Eye. *J Opt Soc Am* **42**, 872 (1952).

11.   E. Ahissar, A. Arieli, Figuring space by time. *Neuron* **32**, 185 (Oct 25, 2001).

12.   S. Viollet, Vibrating makes for better seeing: from the fly's micro-eye movements to hyperacute visual sensors. *Frontiers in Bioengineering and Biotechnology* **2**, 1 (2014).

13.   K. Donner, S. Hemilä, Modelling the effect of microsaccades on retinal responses to stationary contrast patterns. *Vision Res* **47**, 1166 (Apr, 2007).

14.   R. Juston, L. Kerhuel, N. Franceschini, S. Viollet, Hyperacute edge and bar detection in a bioinspired optical position sensing device. *IEEE/ASME Transactions on Mechatronics* **19**, 1025 (2014).

15.   S. R. Henderson, H. Reuss, R. C. Hardie, Single photon responses in *Drosophila* photoreceptors and their regulation by Ca^{2+}. *J Physiol-London* **524**, 179 (Apr 1, 2000).

16.   G. Blaj, J. H. van Hateren, Saccadic head and thorax movements in freely walking blowflies. *J Comp Physiol A* **190**, 861 (Nov, 2004).

17.   M. F. Land, Head Movement of Flies during Visually Guided Flight. *Nature* **243**, 299 (1973).

18.   C. Schilstra, J. H. van Hateren, Stabilizing gaze in flying blowflies. *Nature* **395**, 654 (Oct 15, 1998).

19.   N. Franceschini, R. Chagneux, paper presented at the Gottingen Neurobiology Report, Stuttgart, 1997.

20.   R. Hengstenberg, Eye Muscle System of Housefly Musca-Domestica .1. Analysis of Clock Spikes and Their Source. *Kybernetik* **9**, 56 (1971).

21.   S. Martinez-Conde, J. Otero-Millan, S. L. Macknik, The impact of microsaccades on vision: towards a unified theory of saccadic function. *Nat Rev Neurosci* **14**, 83 (Feb, 2013).

22.   S. N. Fry, R. Sayaman, M. H. Dickinson, The aerodynamics of free-flight maneuvers in *Drosophila*. *Science* **300**, 495 (Apr 18, 2003).

23.   B. R. H. Geurten, P. Jahde, K. Corthals, M. C. Gopfert, Saccadic body turns in walking *Drosophila*. *Front Behav Neurosci* **8**, (Oct 22, 2014).

24.   F. Wong, B. W. Knight, F. A. Dodge, Adapting Bump Model for Ventral Photoreceptors of *Limulus*. *J Gen Physiol* **79**, 1089 (1982).

25.   C. E. Shannon, A Mathematical Theory of Communication. *AT&T Tech J* **27**, 379 (1948).

26.   R. R. D. van Steveninck, S. B. Laughlin, The rate of information transfer at graded-potential synapses. *Nature* **379**, 642 (Feb 15, 1996).

27.   J. Howard, B. Blakeslee, S. B. Laughlin, The Intracellular Pupil Mechanism and Photoreceptor Signal - Noise Ratios in the Fly Lucilia-Cuprina. *Proc R Soc Ser B-Bio* **231**, 415 (Sep 22, 1987).

28.   M. Rivera-Alba et al., Wiring Economy and Volume Exclusion Determine Neuronal Placement in the *Drosophila* Brain. *Curr Biol* **21**, 2000 (Dec 6, 2011).

29.   T. J. Wardill et al., Multiple Spectral Inputs Improve Motion Discrimination in the *Drosophila* Visual System. *Science* **336**, 925 (May 18, 2012).

30.   L. Zheng et al., Feedback network controls photoreceptor output at the layer of first visual synapses in *Drosophila*. *J Gen Physiol* **127**, 495 (May, 2006).

31.   M. F. Land, Motion and vision: why animals move their eyes. *J Comp Physiol A* **185**, 341 (Oct, 1999).

32.   P. T. Gonzalez-Bellido, T. J. Wardill, M. Juusola, Compound eyes and retinal information processing in miniature dipteran species match their specific ecological demands. *P Natl Acad Sci USA* **108**, 4224 (Mar 8, 2011).

33.   L. Freifeld, D. A. Clark, M. J. Schnitzer, M. A. Horowitz, T. R. Clandinin, GABAergic Lateral Interactions Tune the Early Stages of Visual Processing in *Drosophila*. *Neuron* **78**, 1075 (Jun 19, 2013).

34.   M. G. Burg, P. V. Sarthy, G. Koliantz, W. L. Pak, Genetic and Molecular-Identification of a *Drosophila* Histidine-Decarboxylase Gene Required in Photoreceptor Transmitter Synthesis. *Embo J* **12**, 911 (Mar, 1993).

35.   A. Dau et al., Evidence for Dynamic Network Regulation of *Drosophila* Photoreceptor Function from Mutants Lacking the Neurotransmitter Histamine. *Frontiers in Neural Circuits* **10**, (2016).

36.   M. Joesch, B. Schnell, S. V. Raghu, D. F. Reiff, A. Borst, ON and OFF pathways in *Drosophila* motion vision. *Nature* **468**, 300 (Nov 11, 2010).

37.   U. Tepass, K. P. Harris, Adherens junctions in *Drosophila* retinal morphogenesis. *Trends Cell Biol* **17**, 26 (Jan, 2007).

38.   K. G. Götz, Optomotory Investigation of the Visual-System of Some Eye Mutations of the *Drosophila* Fruit-Fly. *Kybernetik* **2**, 77 (1964).

39.   D. G. Stavenga, Angular and spectral sensitivity of fly photoreceptors. II. Dependence on facet lens F-number and rhabdomere type in *Drosophila*. *Journal of Comparative Physiology a-Neuroethology Sensory Neural and Behavioral Physiology* **189**, 189 (Mar, 2003).

40.   M. Juusola, Z. Song, R. C. Hardie, in Encyclopedia of Computational Neuroscience, D. Jaeger, R. Jung, Eds. (Springer, New York, 2015), pp. 2359-2376.

Acknowledgments: We thank C-H. Lee, E. Chiappe, R. Strauss and E. Buchner for flies, S. Tang for help with the flight simulator, O. List for some illustrations and A. Nikolaev, A. Lin for discussions and comments. This work was supported by the open research fund of the State Key Laboratory of Cognitive Neuroscience and Learning (M.J.), NSFC project (30810103906: MJ), Jane and Aatos Erkko Foundation Fellowships (M.J. and J.T.), The Leverhulme Trust grant (RPG-2012-567: M.J.) and the BBSRC grants (BB/F012071/1, BB/D001900/1 and BB/H013849/1: M.J.). The authors declare no competing financial interest. Supplementary materials accompany the paper on Science Online. Author Contributions: M.J. designed the study and wrote the main paper; M.J. and A.D. wrote the Supplementary Texts; M.J., A.D., J.T., N.S., D.R., D.J., S.D., F.B. and R.H. performed or participated in the experiments; M.J., A.D., Z.S., S.D., G.d.P. and J.T. analyzed the data. Z.S., AD, J.T. G.d.P. and M.J. developed the models, Z.S., AD, G.d.P. and J.T. ran the simulations.

Supplementary Materials:

Materials and Methods

Supplementary Texts S1-S10

Figures S1-S10

Movies S1-S3

References

Supplementary Materials for

Microsaccadic information sampling provides *Drosophila* hyperacute vision

Mikko Juusola[1,2]*‡, An Dau[2]‡, Zhuoyi Song[2]‡, Narendra Solanki[2], Diana Rien[1,2], David Jaciuch[2], Sidhartha Dongre[2], Florence Blanchard[2], Gonzalo G. de Polavieja[3], Roger C. Hardie[4] and Jouni Takalo[2]

correspondence to: m.juusola@sheffield.ac.uk

This PDF file includes:

> Materials and Methods
> Supplementary Texts 1-10
> Figs. S1 to S10
> Captions for Movies S1 to S3
> References

Other Supplementary Materials for this manuscript includes the following:

> Movies S1 to S3

Materials and Methods
Flies
2-10 day old wild-type red-eyed (Canton-S and Berlin) fruit flies (*Drosophila melanogaster*) and *hdc^{JK910}*-mutants were used in the experiments. Other transgenic and mutant *Drosophila* tests and controls are explained in specific **Supplementary Texts**. *Drosophila* were raised at 18°C in a 12-hour/12-hour dark/light cycle and fed on standard medium in our laboratory culture.

Electrophysiology
Sharp microelectrode recordings from *Drosophila* R1-R6 photoreceptors were detailed before (*7, 41*), and we only list the key steps here. Flies were immobilized to a conical holder by beeswax (*7*) (**Fig. 1A**). A small hole, the size of a few ommatidia, was cut in the dorsal cornea for the recording electrode and sealed with Vaseline to prevent tissue from drying. R1-R6s' intracellular voltage responses were recorded to different spatiotemporal light patterns (see below) using sharp filamented quartz or borosilicate microelectrodes (120–220 MΩ), filled with 3 M KCl. A blunt reference electrode, filled with fly ringer, was inserted in the head capsule. The flies' temperature was kept either at 19 ± 1 or 25 ± 1°C by a feedback-controlled Peltier device, as indicated in the figures. The recordings were performed after 1-2 minutes of dark adaptation, using the discontinuous clamp method with a switching frequency 20-40 kHz. The electrode capacitance was compensated using the head-stage output voltage. To minimize effects of damage and external noise on the analysis, only stable recordings of low-noise and high sensitivity were chosen for this study (sometimes lasting several hours). Such photoreceptors typically had resting potentials <-60 mV in darkness and >45 mV responses to saturating test light pulses (*7*).

Light stimulation. We used a high power "white" LED (Seoul Z-Power P4 star, white, 100 Lumens) to test individual R1-R6 photoreceptors' encoding dynamics (**Fig. 1**). It was connected to a randomized quartz fiber optic bundle (transmission range: 180-1,200 nm), fitted with a lens (providing ~3° homogeneous light disk as seen by the flies), and attached onto a Cardan arm system for accurate positioning at the center of each tested cell's receptive field. Its light output was driven by an OptoLED (Cairn Research Ltd, UK), which utilizes a feedback circuitry with a light-sensor. This LED has red component wavelengths, which minimizes prolonged depolarizing afterpotential (PDA) effects (*42*). Because long recordings can show sensitivity drifts, attributable to muscle activity (**Supplementary Text 4**), the stimulus XY-position was regularly tested and, if needed, re-centered between long stimulus runs.

We used a bespoke 25 light-point array to measure individual R1-R6 photoreceptors' receptive fields and responses to moving point objects (bright dots, **Fig. 3**; dark dots, **Supplementary Text 9**). Again, a custom-made Cardan arm system was used to accurately position the array's center light-point (no. 13) at the center of each tested cell's receptive field. The dot size and the minimum inter-dot-distance, as seen by *Drosophila*, was 1.7°. Details of this device and the recording procedures are given in **Supplementary Texts 4** and **6**.

Stimulus patterns. Single photoreceptors' diurnal temporal encoding gamut was tested systematically over different bandwidth and contrast distributions; using 20 distinct light intensity time series stimuli, which were presented at the center of their receptive fields. The used test stimuli was based upon 5 different 2 s long Gaussian white-noise light intensity time series patterns, which had "flat" power spectrum up to 20, 50, 100, 200, or 500 Hz (**Fig. 1B**), as

low-pass filtered by MATLAB's filter toolbox, and the same peak-to-peak modulation (2 units). These were then superimposed on four backgrounds: BG0 (0 units, dark), BG0.5 (0.5 units), BG1 (1 unit) or BG1.5 (1.5 units, bright) on a linear intensity scale, giving altogether 20 unique stimulus patterns. As the two lowest backgrounds clipped downwards-modulation, prolonging dark intervals, the resulting stimuli ranged from high-contrast bursts ($c = \Delta I/I \sim 1.46$ at BG0) to low-contrast Gaussian ($c \sim 0.22$ at BG1.5).

As further controls, we tested how well R1-R6 photoreceptors responded to dark saccadic contrast changes (**Supplementary Text 9**) and to their bright counterparts. In these experiments, R1-R6s were adapted for 10 s to BG0.5 and BG1 before repeated stimulation. In addition, we recorded the tested cells' responses to naturalistic light intensity time series (*5*, *43*) (NS), selected from van Hateren natural stimulus collection (*43*) (**fig. S4**).

In all these experiments, ≥25 consecutive responses to each repeated stimulus were recorded.

Data acquisition. Both the stimuli and responses were filtered at 500 Hz (KEMO VBF/23 low-pass elliptic filter, UK), and sampled together at 1-10 kHz using a 12-bit A/D converter (National Instruments, USA), controlled by a custom-written software system, Biosyst in Matlab (Mathworks, USA) environment. For signal analyses, if need, the data was down-sampled to 1 kHz.

Analyses
Because of short-term adaptive trends, we removed the first 3–10 responses to repeated stimulation from the analysis and used the most stable continuous segment of the recordings. Information theoretical methods for quantifying responses of approximately steady-state-adapted fly photoreceptors to different stimuli were described in detail before (*5*, *6*, *44*, *45*). Below we list the key approaches used here.

Signal-to-noise ratio (SNR) and information transfer rate estimates. In each recording, simulation or Poisson light stimulus series (see below), the signal was the mean, and the noise was the difference between individual traces and the signal (*5*, *7*). Therefore, for a dataset of 20 responses (n = 20 traces), there was one signal trace and 20 noise traces. The signal and noise traces were divided into 50% overlapping stretches and windowed with a Blackman–Harris 4-term window, each giving three 500-points-long samples. Because each trace was 2 s long, we obtained 60 spectral samples for the noise and 7 for the signal. These were averaged, respectively, to improve the estimates.

$SNR(f)$ of the recording, simulation, or Poisson light stimulus series was calculated from their signal and noise power spectra, $<|S(f)|^2>$ and $<|N(f)|^2>$, respectively, as their ratio, where $|\ |$ denotes the norm and $<>$ the average over the different stretches (*45*). To eliminate data size and processing bias, the same number of traces (n = 20) of equal length (2,000 points) and sampling rate (1 kHz; 1 ms bin size) were used for calculating the $SNR(f)$ estimates for the corresponding real recordings, photoreceptor model simulations and the simulated Poisson stimuli.

Information transfer rates, *R*, for each recording, simulation, or Poisson light stimulus series were estimated by using the Shannon formula (*25*), which has been shown to obtain robust estimates for these types of continuous signals (*5*, *44*). We analyzed steady-state-adapted recordings and simulations, in which each response (or stimulus trace) is expected to be equally

representative of the underlying encoding (or statistical) process. From *SNR(f)*, the information transfer rate estimates were calculated as follows:

$$R = \int_0^\infty (\log_2[SNR(f) + 1])df \qquad (1)$$

We used minimum = 2 Hz and maximum = 500 Hz (resulting from 1 kHz sampling rate and 500 points window size). The underlying assumptions of this method and how the number and resolution of spectral signal and noise estimates and the finite size of the used data can affect the resulting Information transfer rate estimates have been analyzed before (*5*, *44*, *46*, *47*) and are further discussed in **Supplementary Text 2**. All data analyses were performed with Matlab (MathWorks).

Measuring photoreceptors' visual acuity. We measured dark- and light-adapted wild-type R1-R6 photoreceptors' receptive fields by their acceptance angles, Δρ, using intracellular voltage responses to random light-points in a stimulation array. These measurements were compared to those of *hdc^{JK910}*-mutants (*34*, *48*), in which first-order interneurones receive no neurotransmitter (histamine) from photoreceptors and so are incapable of feedback-modulating the photoreceptor output. Both the wild-type and mutant R1-R6 photoreceptors' mean Δρ was about twice the mean interommatidial angle, Δφ. The stimulus apparatus, the method and result details and theoretical electron micrograph comparisons of their mean rhabdomere sizes are explained in **Supplementary Texts 4-5**.

Spatiotemporal analyses using the classic conventional models. Voltage responses of wild-type and *hdc^{JK910}* R1-R6s to moving bright dots were evaluated against their respective classic model simulations, in which each recorded receptive field was convoluted by the same cell's impulse response (*3*, *4*) (1st order Volterra kernels). The motion blur effects were quantified by comparing the real R1-R6 outputs to their deterministic model predictions. Details of the analysis are given in **Supplementary Text 6**.

Biophysical modeling
Time series analyses. We used our recently published biophysically-realistic stochastic photon sampling model (*6*) of a *Drosophila* R1-R6 photoreceptor to simulate macroscopic voltage response to different repeated light intensity time series patterns from a point source (**figs S5-6**). The model has no free parameters. Its design and the general aims and details of these simulations are given in **Supplementary Texts 1** and **2**. To eliminate data size bias, the signaling properties and performance of the simulations were quantified and compared to the corresponding recordings by using the same analytical routines on the same-sized data-chunks. The models were run using Matlab in the University of Sheffield computer cluster.

For each stimulus, its mean was adjusted to maximize information of the simulated photoreceptor outputs, mimicking the action of the photomechanical adaptations (intracellular pupil mechanism and rhabdomere contractions; **Supplementary Texts 2** and **7**). This optimization set the effective mean photon rates from 8×10^4 at BG1.5 to 8×10^5 photons/s at BG0 (**fig. S5**). Thus, each of these light levels was considered to represent the optimal daylight input (through the photomechanical adaptations) to a rhabdomere, in which modulation enabled the largest sample (bump) rate changes. Otherwise, more of its sampling units (30,000 microvilli) would be either underutilized or refractory (saturation). The maximum information rates of the simulated photoreceptor outputs closely followed the corresponding mean information transfer rates of the real recordings over the whole tested encoding range (**Supplementary Text 2**). This

implies that the central function of the photoreceptors' combined photomechanical adaptations is to maximize their information transfer, and that the resulting estimates represent realistic maxima.

Encoding efficiency. A photoreceptor's encoding efficiency, η, was the ratio between the information rates of its voltage output, R_{output}, and the corresponding effective light input, R_{input}:

$$\eta = \frac{R_{output}}{R_{input}} \tag{2}$$

with R_{output} and R_{input} estimated by the Shannon formula (Eq. 1). Details are in **Supplementary Text 2.**

Modeling R1-R6 output to moving dots. We developed a new "microsaccadic sampling"-model to predict how photomechanical rhabdomere contractions (microsaccades) move and narrow *Drosophila* R1-R6 photoreceptors' receptive fields to resolve fast-moving objects. **Supplementary Text 8** gives the details of this modeling approach, which combines the stochastic photon sampling model (*6*) with additional fixed ommatidium optics and photomechanical rhabdomere contraction parameters. The same supplement shows examples of how refractory photon sampling and rhabdomere contractions jointly improve visual acuity.

High-speed video of the light-induced rhabdomere movements
Cornea-neutralization method with antidromic far-red (>720 nm) illumination was used to observe deep pseudopupils (*47*) (photoreceptor rhabdomeres) in the *Drosophila* eye at 21 °C. A high-speed camera (Andor Zyla, UK; 500 frames/s), connected to a purpose-built microscope system, recorded fast rhabdomere movements *in vivo* to blue-green light stimuli (470 + 535 nm peaks), which were delivered orthodromically into the eye. The method details, mutant and transgenic *Drosophila* used and the related image analyses are explained in **Supplementary Text 7**.

Flight simulator experiments
Open-loop configuration was used to test hyperacute motion vision. Wild-type flies were tethered in a classic torque meter (*50*) with heads fixed, and lowered by a manipulator into the center of a black and white cylinder (spectral full-width: 380–900 nm). A flying fly saw a continuous panoramic scene (360°), which in the tests contained multiple vertical stripes (black and white bars of equal width). The control was a diffuse white background. After viewing the still scene for 1 s, it was spun counterclockwise by a linear stepping motor for 2 s, stopped for 2 s before rotating clockwise for 2 s, and stopped again for 1 s. This 8 s stimulus was repeated 10 times and each trial, together with the fly's yaw torque responses, was sampled at 1 kHz (*29*). Flies followed the stripe scene rotations, generating yaw torque responses (optomotor responses to right and left), the strength of which reflected the strength of their motion perception. The flies did not follow the white control scene rotations. The panoramic scenes had ±360° azimuth and ±45° elevation, as seen by the fly. The stripe scenes had 1.0 contrast and their full-wavelength resolutions were either hyperacute (1.16° or 2.88°) or coarse (14.40°), giving the inter-bar-distances of 0.58°, 1.44° and 7.20°, respectively. The white scene has zero contrast. The scene rotation velocity was 45 °/s.

Transmission Electron Microscopy

The fly eye dissection, fixation embedding, sectioning and imaging protocols for EM (**fig. S7A**) are described in **Supplementary Text 5**.

Statistics

Test responses were compared with their controls by performing two-tailed t-tests to evaluate the difference in the compared datasets. Welch's t-test was used to accommodate groups with different variances for the unpaired comparisons. In the figures, asterisks are used to mark the statistical significance: ns indicates $p > 0.05$, $*$ indicates $p \leq 0.05$, $**$ indicates $p \leq 0.01$, and $***$ indicates $p \leq 0.001$.

Supplementary Texts 1-10

Table of Contents   Pages: 23-25

Text S1: "Stochastic adaptive visual information sampling"-theory in brief

Overview

This supplement describes the basic theoretical principles of how fly photoreceptors sample photons, providing central background information for the results presented in **Figures 1-5** in the main paper.

Stochastic adaptive sampling of information by R1-R6 photoreceptors

Many lines of evidence imply that in a *Drosophila* R1-R6 photoreceptor 30,000 individual refractory sampling units (microvilli) integrate exponential photon flux changes (~10^6-fold) from the environment into macroscopic voltage responses of biophysically limited amplitude range (~60 mV) and bandwidth (~200 Hz) (*5-7, 51*). In essence, a light-adapted R1-R6 counts photons imperfectly, which, nonetheless, adds up highly reproducible neural representations of light changes within its receptive field (*40*).

In this publication, we quantify such quantal information processing through large-scale experimental and theoretical analyses. Our overriding aim is to analyze R1-R6 photoreceptors' diurnal encoding range systematically; from light bursts to Gaussian white-noise stimulation, to point-objects moving across their receptive fields at saccadic speeds. Because of the outstanding stability and signal-to-noise ratio of the intracellular recordings from an *in vivo Drosophila* preparation (*5, 7, 30*), providing apparent ergodicity, we can directly compare experimentally recorded macroscopic voltage responses to those of biophysically realistic photoreceptor model simulations (*5, 6, 40, 52*), in which stochastically operating microvilli sampled similar stimuli (**Box S1.1**). The mechanistic knowledge so obtained about the dynamics and limitations of quantal visual information processing provides us with deep new understanding of how well *Drosophila*, and other insect eyes, can see the world.

Box S1.1. Schematic of the biophysically realistic *Drosophila* R1-R6 model, which mimics phototransduction by transducing light input (a dynamic flux of photons) into macroscopic output, light-induced current (LIC). The model is modular, containing four parts, three of which are shown here. Phototransduction occurs within a photoreceptor's light-sensitive part, the rhabdomere, which contains 30,000 photon sampling units, microvilli (blue bristles). Each microvillus contains a full phototransduction cascade reactions, and can transduce single photon energies (green dots) into unitary responses, quantum bumps (QB or samples) of variable amplitudes. (**A**) In the 1st module, 30,000 microvilli sample incoming photons. The light input, as photons/s, is randomly distributed over them (each row of open circles indicate photons hitting - being absorbed by - a single microvillus). (**B**) In the 2nd module, the successfully absorbed photons in each microvillus are transduced into QBs (a row of unitary events). In each microvillus, the success of transducing a photon into a QB depends upon the refractoriness of its phototransduction reactions. This means that a microvillus cannot respond to the next photons until its phototransduction reactions have recovered from the previous photon absorption, which takes about 50-300 ms. The photons hitting a refractory microvillus cannot evoke QBs, but will be lost. (**C**) In the 3rd module, QBs from all the microvilli then integrate the dynamic

macroscopic LIC. Conversely, the light input (green trace) can be reconstructed by adding up all the photons distributed across the 30,000 microvilli.

Main framework

We used our previously published biophysical *Drosophila* R1-R6 model (*5, 6, 40, 52*) to simulate voltage responses to time series of light intensities. The model contains four modules:

- *Random Photon Absorption Model*: regulates photon hits (absorptions) in each microvillus, following Poisson statistics (*52*) (**Box S1.1A**).
- *Stochastic Bump Model*: stochastic biochemical reactions inside a microvillus capture and transduce the energy of photons to variable quantum bumps or failures (**Box S1.1B**). Here, Gillespie algorithm provides discrete and stochastic phototransduction cascade simulations with few reactants as every reaction is explicitly simulated.
- *Summation Model*: bumps from 30,000 microvilli integrate to the macroscopic light-induced current (LIC) response (**Box S1.1C**).
- *Hodgkin-Huxley (HH) Model of the photoreceptor plasma membrane* (*53, 54*): transduces LIC into a voltage response (**Box S1.2**).

Box S1.2. *Drosophila* **R1-R6 photoreceptor membrane's electrical circuit (HH-model)**. A photoreceptor's membrane potential, V_m, is the difference between the negative inside (intracellular) and positive outside (extracellular) voltages. V_m can be calculated, using Hodgkin-Huxley formalism, whereupon, a photoreceptor membrane is modelled as a capacitor, C_m, voltage-gated channels as voltage-regulated conductances, g, leak channels as fixed conductances, reversal potentials for different ion species as DC-batteries. Abbreviations: I_{ksh}: Shaker; I_{dr}, delayed rectifier; I_{novel}, novel K^+; K_{leak}: K^+ leak; I_{cl}, chloride leak currents. The used deterministic *Drosophila* photoreceptor HH-model is adapted from (*53, 54*).

The formalism, assumptions and many tests of the biophysical photoreceptor model, which has no free parameters, are given in our previous publications (*5, 6, 40*). All its parameter values can be found and downloaded from: http://www.sciencedirect.com/science/article/pii/S0960982212006343

Stochastic quantal models supersede empirical black-box approaches

Stochastic photon sampling fly photoreceptor model:

- Has no free parameters
- Is general - predicts realistic responses to any light stimulus pattern
- Is transferable - predicts realistic responses of different fly photoreceptors
- Provides deep mechanistic understanding to light information sampling

We have shown before that this quantal stochastic modeling approach is general and transferable, and therefore directly applicable to quantify photoreceptor functions in different light conditions

and fly species (*5, 6, 40*). Importantly, it does not require full knowledge of all molecular players and dynamics in the phototransduction to generate realistic responses (*5, 6, 40*). From a computational viewpoint, the exactness of the simulated molecular interactions is not critical. As long as the photoreceptor model contains the right number of microvilli (*e.g.* 30,000 in a *Drosophila* and 90,000 in a *Calliphora* R1-R6 photoreceptor), each of which is a semiautonomous photon sampling unit, and the dynamics of their photon-triggered unitary responses (quantum bumps [QBs] or samples) approximate those in the real recordings, it will sample and process information much like a real photoreceptor. Conversely, this further means that by knowing the number of microvilli and their average QB waveform, latency distribution, and refractory period distributions, we can closely predict a fly photoreceptor's macroscopic response to any given light intensity time series stimulus (*cf.* **figs S5** and **S6**). The same model can then be applied to estimate how well the real photoreceptor output resolves moving objects (see **Figs 4G-I** and **5** in the main paper; **Text S8** gives the details of this approach).

Deterministic empirical fly photoreceptor models:

- Fit parameters to specific stimulus sets
- Predict less accurately responses to new stimuli of different input statistics (to which the models have not been tuned to before)
- Cannot provide deep mechanistic understanding of how photoreceptors sample light information

In the conventional empirical "black-box" approaches, the photoreceptor models' filters, such as linear and nonlinear kernels, and static nonlinearities are adjusted to minimize the difference between the recorded responses and the model output to a specific stimulus set (light condition) (*3, 55-57*). However, because such *models are not built upon the real cells' physical quantal information sampling constraints*, which change from one stimulus statistics to another (*5*), they struggle to respond accurately to new stimulus statistics. Explicitly, the models lack intrinsic structural information of how quantum bump dynamics and microvilli refractoriness must differ during different stimuli. For example, Volterra (*55*) kernels estimated for Gaussian white-noise stimulation will predict less accurately responses to bursty light inputs. This is because during bursty light stimulation the fly photoreceptors' quantal information sampling dynamics rapidly adapt to a different regime, where their microvilli (sampling units) are less refractory. Hence, the real photoreceptors now integrate macroscopic responses from larger sample (QB) rate changes of enhanced rise and decay dynamics. To appropriately approximate these new dynamics, the empirical models would need to generate new kernels of different temporal profiles, which is impossible without retuning the model parameters. Accordingly, without the biophysical knowledge being implemented in their mathematical structure, the classic dynamic photoreceptor models fail to predict how well the real photoreceptors resolve moving objects (see **Fig. 3**, **Text S6** and **Text S8**).

Text S2: Information maximization by photomechanical adaptations & connectivity

Overview
This supplement describes how the "stochastic adaptive visual information sampling"-theory (**Text S1**) predicts and explains the roles of R1-R6 photoreceptors' photomechanical adaptations and network connections in information maximization at different stimulus conditions, as shown in **Figures 1-2** in the main paper.

In this supplement:
- We test the hypotheses that photomechanical adaptations (intracellular pupil and rhabdomere contractions; see **Text S7**) and network connections in the *Drosophila* eye contribute importantly to optimizing the capture and representation of visual information.
- We first estimate through simulations how a R1-R6 photoreceptor's intracellular pupil and rhabdomere contractions are jointly optimized for maximal information sampling by its 30,000 microvilli. The simulations predict that these mechanisms' optimal combined light throughput in bright conditions should be different for different stimuli.
- We then compare the model predictions to corresponding intracellular recordings and find a comprehensive agreement between the theory and mean experiments for all the tested stimuli.
- This striking correspondence enables us to further estimate how the lamina network shapes information transfer of individual R1-R6 photoreceptors.
- Remarkably, our data and analyses strongly suggest that voltage output is different in each R1-R6, which are brought together in neural superposition during development to sample light changes from a small local visual area.
- These results are consistent with the hypothesis that the variability in the retinal sampling matrix dynamics and topology minimizes aliasing and noise, enabling its parallel processing to generate reliable and maximally informative neural estimates of the variable world (*5, 40, 58, 59*).
- Finally, we explain how to calculate a photoreceptor's encoding efficiency for different light stimuli, highlighting the assumptions and limits of this method.

Fly photoreceptors' pupil mechanism
In a fly photoreceptor, intracellular screening pigments form its pupil mechanism (**Box S2.1**). The pupil protects a photoreceptor's sampling units (30,000 microvilli in a *Drosophila* R1-R6) from saturation (*5, 27*). At bright light exposure, screening pigments migrate to narrow the aperture they form collectively (*49*), shielding off excess light from reaching the microvilli. This is important because midday sunshine on a photoreceptor may contain 10^{6-8} photons/s, and without the pupil mechanism would deteriorate the encoding function of its finite microvillus population (*5, 27*). The pupil opening and closing seem modulated by light-driven intracellular Ca^{2+}-concentration changes (*60*), and show reasonably fast dynamics (from fully open to fully closed within 15 s) (*61*) for adapting its light throughput to ambient changes. Although our biophysical (stochastically operating) *Drosophila* photoreceptor model (*6, 40*) lacks the pupil mechanism and any other photomechanical adaptations (*cf.* **Text S7**), their joint effects can be predicted through simulations; by assuming that their objective function is to maximize the photoreceptor's information capture.

Box S2.1. Compound eye and photoreceptors' intracellular pupil mechanism. (**A**) *Drosophila* eyes are composed of about 800 modular units, ommmadia. (**B**) Each ommatidium contains a lens system and underneath it 8 photoreceptor cells: the outer receptors, R1-R6, and the inner receptors, R7/R8. In the electron micrograph, which numbers each cell's rhabdomere (light sensitive part), R8 is not shown because it lies directly below R7. (**C**) Schematic of the intracellular pupil mechanism. Left: During dark-adaptation, screening pigments (small dots) are scattered in the R1-R7 somata. Middle: R1-R6 light-adapted. Blue-green bright light drives the screening pigment migration towards the R1-R6 rhabdomeres (central discs, containing 30,000 microvilli, photon sampling units, depicted as stripes in the discs), which express blue-green-sensitive Rh1-rhodopsin, as their phototransduction rises intracellular Ca^{2+}-concentration. With the pupil closing (seen as the dark rims around the rhabdomeres), light input to the microvilli reduces. Note that R7, which expresses UV-rhodopsin, is not light-adapted and its screening pigments remain scattered. Right: All photoreceptors light-adapted. Bright UV-light closes all pupils because R7s express UV-sensitive Rh3- and Rh4-rhodopsins, and in R1-R6s' Rh1-rhodopsin is electrochemically coupled to UV-sensitive sensitizing pigment. Redrawn and modified from (*61, 62*).

Photomechanical light-screening hypothesis

We hypothesize that the intracellular pupil, besides affecting a photoreceptor's angular and spectral sensitivity (*63*) (see **Text S4**), participates in maximizing a photoreceptor's information sampling by optimizing light input intensity to its microvilli in time. Specifically in this context, it works together with all other photomechanical adaptations within an ommatidium, including the much faster light-induced rhabdomere contractions (**Videos S1-3**), in protecting the microvilli from saturation (see **Text S7**). Thus, collectively, we consider the photomechanical adaptations (**Box S2.2**) as a biological manifestation of a mathematical information maximization function.

Box S2.2. Different temporal ranges of a *Drosophila* R1-R6 photoreceptor's intrinsic light adaptation mechanisms. Photomechanical adaptations, such as light-induced intracellular screening pigment migration (pupil mechanism, black; see **Box S2.1**) and rhabdomere contractions (red; see **Text S7**), operate with refractory photon sampling by 30,000 microvilli (blue) and their quantum bump dynamics (sample duration and jitter, green; see **Text S1**) in modulating light input to a photoreceptor, and consequently its voltage output. Together these mechanisms enable efficient encoding of behaviorally important visual information at daylight conditions (**Figs 1-2**) by covering a broad range of temporal light changes; with the pupil and rhabdomere contractions being slower than the photon sampling dynamics.

To function optimally, the photomechanical adaptations need to regulate the ambient illumination so that the temporal light changes they let through would cause maximal sample

(quantum bump) rate changes (*5*). The higher the photoreceptor's sample rate changes, the higher its rate of information transfer (*5*). Too bright light would saturate microvilli because most of them would be rendered refractory, reducing their dynamic sample counts. Conversely, too dim light would not utilize microvilli population fully, producing low (suboptimal) sample counts. Therefore, the optimal light intensity throughput for maximum information capture is somewhere between. This value is expected to depend upon five factors:

- Light intensity time series structure (we consider all the 20 light patterns tested in **Fig. 1B**)
- Number of microvilli (~30,000 in a R1-R6 *Drosophila* photoreceptor)
- Refractory period distribution of the microvilli (50-300 ms in a R1-R6 *Drosophila* photoreceptor)
- Quantum bump waveform (sample duration)
- Quantum bump latency distribution ("sample jitter")

Detailed tests and descriptions of why and how these factors contribute to encoding in fly photoreceptors are given in our previous publications (*5-7, 15, 40, 44, 45, 64*).

Hypothesis testing and verification

To test the photomechanical light-screening hypothesis, we simulated voltage responses to 20 different light patterns (**Fig. 1B**) at 15 different light intensity (or brightness) levels, which ranged from 5 x 10^4 to 1 x 10^6 photons/s (5, 6, …, 9 x10^4; 1, 2, ..., 9 x10^5; 1 x 10^6). In each simulation, the stochastic photoreceptor model generated 20 independent responses to the given 2-seconds-long (2,000 points) light pattern of a given intensity, following the published procedures (*5*). These 20 responses were used to estimate the model's rate of information transfer for that specific stimulus pattern (1/20) at that specific light level (1/15). So all together, we could have simulated 20 repeated photoreceptor outputs to 300 (20 x 15) different 2,000 points long stimulus patterns. But because the model's maximum information transfer rate estimates turned out to be relatively straightforward to determine for many light patterns (*cf.* **Box S2.3**), the total number of simulations never reached this limit. Nevertheless, being computationally expensive, the stochastic simulations took months to complete.

Crucially, in all the simulations, the photoreceptor model was exactly the same. Its stochastic bump production dynamics (waveform, latency and refractory distributions) were governed by light-adapted values with every single parameter fixed, and these parameter values were unchanged in each simulation. The supplement of the reference (*6*) lists these parameter values, which were collected from intracellular experiments or logically extrapolated to be biophysically realistic for light-adapted *Drosophila* photoreceptors. This supplement is downloadable from:

http://www.sciencedirect.com/science/article/pii/S0960982212006343

Figure S5 shows the simulated voltage responses (traces above) that carried the maximum information transfer rates for the 20 tested light patterns (traces below) and the corresponding intensity levels (as effective photons/s) that evoked them. The simulations match the overall size, appearance and dynamics of the real recordings astonishingly well (**Fig. 1B** and **fig. S1B**), indicating that the photoreceptor model, with its photomechanics optimizing light input intensity, samples and integrates light information much like its real-life counterparts. Notice that the optimal light intensity is the same for the different bandwidth (20, 50, 100, 200 and 500 Hz) stimuli within one BG. But for each BG (BG0, BG0.5, BG1 and BG1.5) this optimum is different. For example, for BG0, which results in bursty (high contrast) stimulation,

the optimal light intensity level is ~8 x 10^5 photons/s. Whereas for BG1.5 of low contrast Gaussian white-noise stimulation, this is 10-times lower (~8 x 10^4 photons/s).

We express these intensity levels in units of *effective* photons/s. This is because, theoretically, we have deduced the mean photon throughput that effectively fluxes into microvilli for a *Drosophila* photoreceptor to sample the best estimates of the given light stimuli. In other words, if the photomechanical screening mechanisms set the light input intensity for maximal information capture, as is our hypothesis, then these light-intensity values should also closely approximate the actual absolute photon flux changes that drive phototransduction in the real experiments (as recorded intracellularly from wild-type R1-R6 photoreceptors, which have the normal pupil mechanism and photomechanical rhabdomere contractility; **Figs 1** and **fig. S1**).

Because of the remarkable dynamic correspondence between the experiments (**Figs 1-2** and **figs S1-2**) and the theory (**figs S5-6**) over the whole tested encoding space, we now judge that this hypothesis must be largely true. Importantly, this realization opens up new ways to analyze photoreceptor function. For example, by making the general assumption that photon flux from our LED light source follows Poisson statistics, we could further estimate the lower information transfer rate bound for each tested light intensity pattern, and consequently the upper bound for the *Drosophila* photoreceptor's encoding efficiency (*e.g.* **Fig. 2D**, **figs S2D** and **6D**). More details about this assumption and the analysis are given at the end of this supplement (**Text S2**).

New insight into maximal visual encoding of different stimulus statistics
The reasons why and how the optimal light intensity input (that drives a photoreceptor's information transfer maximally) is different for bursts and Gaussian white-noise stimulation are summarized in **Image S2.3**. Here we assess both cases using data from the stochastic *Drosophila* R1-R6 photoreceptor model simulations, starting with light bursts.

Bursts (**Box S2.3A-D**). These light intensity time series characteristically contain periods of longer dark contrasts, intertwined with brief and bright high-contrast events, as shown for 100 Hz bandwidth stimulation (**Box S2.3A,** dark-yellow trace). Based on our previous analyses (*5*), longer dark contrasts help to recover more refractory microvilli than equally-bright stimuli without these features, improving neural information capture. This makes it more difficult for bursty stimuli to saturate the photoreceptor output. By increasing the stimulus intensity 8-fold, here from 1 x 10^5 to 8 x 10^5 effective photons/s, simply evoked larger macroscopic responses. These, thus, integrated more samples (bumps); as indicated by the larger (black) and smaller (blue) trace, respectively.

Because noise changes little in light-adapted photoreceptor output (*5-7, 45, 64*) (**fig. S3**), the larger responses to brighter bursts have higher and broader signal-to-noise ratio, $SNR_{output}(f)$ (**Box S2.3B**). This, in turn, results in higher information transfer rate estimates, R_{output} (**Box S2.3C**), following Shannon's equation (*25*):

$$R_{output} = \int_0^\infty \left(\log_2\left[SNR_{output}(f) + 1\right]\right)df \qquad (S2.1)$$

Note that with 1 kHz sampling rate used in every experiment, this estimation did not integrate information rate for frequencies from 0 to infinite, but from 2 to 500 Hz instead. However, the limited bandwidth would not considerably affect estimation results because: (i) high-frequency components have SNR << 1 and therefore contain mostly noise. (ii) Whereas even a high SNR_{output} contains little information in its low-frequency components, below 2 Hz.

Note also that we have previously shown the generality of Shannon's information theory for estimating information transfer rates of continuous (analogue) repetitive responses, irrespective of their statistical structure (*5, 44*). That is, for sufficient amount of data, Shannon's equation and triple extrapolation method, which is free of signal and noise additivity and Gaussian distribution assumptions, give comparable rate estimates. Thus, these estimates should evaluate the simulations' relative information rate differences truthfully; i.e. consistently with only small errors.

Box S2.3. Estimating optimal light intensity for 100 Hz high-contrast bursts and Gaussian white-noise (GWN) stimuli for a photoreceptor's maximal information transfer. We hypothesize that the role of photomechanical adaptations, which include the intracellular pupil and contracting rhabdomere (**Text S7**) of a photoreceptor, is to maximize information capture of microvilli by dynamically adjusting the light input falling upon them. The left side of the figure shows how encoding of light bursts (**A-D**) depends upon light intensity; the right side shows the same for GWN (**E-H**). (**A**) Owing to sufficient dark periods, a photoreceptor's sampling units (microvilli) have enough time to recover from their refractoriness even after they have responded to very bright bursts. This enables a photoreceptor to maintain a large pool of available microvilli to sum up high sample (bump) rate changes to any new incoming input, generating larger macroscopic responses to the brighter bursts (8 x 10^5 photons/s black trace) than to the less bright bursts (1 x 10^5 photons/s blue trace). (**B**) Macroscopic responses with larger sample rate changes (black trace, grey area) have higher and broader signal-to-noise ratios (*5*). (**C**) Correspondingly, as the sample sizes (bumps) are similar to both stimuli (*cf.* **fig. S3B**), the larger responses carry a higher information transfer rate (*5*), irrespective of the tested stimulus bandwidth. (**D**) Therefore, a photoreceptor's information transfer rate to bursty inputs increases with light intensity, until the sample rate changes eventually saturate at 8 x 10^5 photons/s; when most of 30,0000 microvilli become refractory (i.e. more microvilli are refractory than available to be light-activated). (**E**) A photoreceptor generates similar size responses to the brighter (8 x 10^5 photons/s black trace) and the less bright (1 x 10^5 photons/s blue trace) GWN inputs. But the response to the less bright input shows more high-frequency modulation. (**F**) Consequently, the response to the less bright input (blue area) has higher and broader signal-to-noise ratio than the response to the brighter input (black trace, grey area). (**G**) This is reflected also in the photoreceptor's information transfer rate, regardless of the GWN bandwidth. (**H**) Information transfer rate in macroscopic photoreceptor output to GWN stimulation saturates at 8-times less bright intensity levels than to bursts (**D**), reaching its maximum at 1 x 10^5 photons/s.

Markedly, a photoreceptor's performance is systematically better to the brighter bursts (black line) than to the less bright ones (blue line), irrespective of the stimulus bandwidth (**Box S2.3C**). Thus, for the brighter bursts, more microvilli are dynamically activated, generating larger sample rate changes. These bumps sum up larger (and more accentuated – see (*5*))

macroscopic responses, packing in more information than the corresponding responses to the less bright bursts.

However, because a *Drosophila* photoreceptor has a finite amount of microvilli, each of which - once activated by a photon's energy - stays briefly refractory, its sample rate changes and thus signaling performance first increases monotonically until about 6 x 10^5 photons/s, before gradually saturating, and eventually decreasing, with increasing burst brightness (**Box S2.3D**). The photoreceptor model's maximum information transfer rate estimate ($R_{max} = 631 \pm 31$ bits/s; marked by a square) for 100 Hz bright bursts is reached at the optimal stimulus intensity of 8 x 10^5 effective photons/s. In other words, this is the amount light the photomechanical adaptations, including the intracellular pupil mechanism and rhabdomere contractions (see **Text S7**), should let through in bright daylight for the fly to see bursty real-world events best. The corresponding performance estimate with the less bright bursts (1 x 10^5 effective photons/s) is 493 ± 12 bits/s (circle).

Gaussian white noise (GWN, **Box S2.3E**-**H**). Because GWN lacks long dark contrasts, refractory microvilli have fewer chances to recover (*5*). Consequently, photoreceptor output to GWN begins to show signs of saturation at lower light intensity levels. **Box S2.3E** shows responses to 100 Hz bandwidth GWN with the mean intensity of 1 x 10^5 (blue) or 8 x 10^5 (black) effective photons/s, respectively. Both responses are about the same size, but the one to the brighter stimulation has less high-frequency modulation. As more microvilli become refractory, smaller sample rate changes (modulation) map light changes into macroscopic voltage responses. (How refractoriness dynamically modulates bump counts and macroscopic response waveforms was analyzed in detail recently (*5*), and thus is not repeated here). Hence, the response to the brighter GWN (black/grey) has lower and narrower signal-to-noise ratio (**Box S2.3F**) over the frequency range than the responses to the 8-times less bright GWN (blue). Naturally, the same holds true for the photoreceptor's information transfer rate estimates (**Box S2.3G**); the less bright GWN gives consistently a better performance (blue), irrespective of the used stimulus bandwidth.

Again, the amount of microvilli and their refractoriness curb a photoreceptor's encoding (or signaling) performance. But to minimize their impact on encoding GWN, the photomechanical screening need to be more restrictive, letting in less light. With brightening 100 Hz GWN (**Box S2.3H**), the model's information transfer rate estimate first steeply increases until its peak ($R_{max} = 369 \pm 15$ bits/s; marked by a circle) at 1 x 10^5 photons/s, and then swiftly declines as progressively more microvilli become refractory and fewer samples are being produced. The corresponding transfer rate estimate for 8 x 10^5 photons/s GWN is 249 ± 17 bits/s (square). Notice, however, that although these results quantify the optimal light throughput for generating maximally informative responses to GWN, such performance is far from the estimated R1-R6 models' information capacity of 631 bits/s (*cf.* **Box S2.3D** and **fig. S6C**).

Simulations' maximal information transfer largely match those of recordings
We next compare the maximum information transfer rate estimates (squares) of the photoreceptor model simulations to those of corresponding *in vivo* recordings (circles) for all BG0 (bursts) and BG1 (GWN) stimuli (**Box S2.4A**). The simulated performance is very close to the measured mean performance for all the tested stimuli, typically falling within the standard deviation of the recordings' information transfer.

In further inspection, two interesting observations can be drawn from this data. First, for the GWN stimuli, irrespective of their bandwidth, the maximum information transfer rate estimates of the model (dotted line) are just a few bits/s (1-10%) higher than the corresponding mean estimates of the real recordings (continuous line). These small differences are probably caused by recording noise. Second, the simulations to bursty stimuli carry less information than the corresponding best recordings, and the recordings show variations in their information transfer.

Box S2.4. Information transfer rate estimates, R_{output}, of in vivo recordings and model simulations show similar encoding dynamics. (**A**) Comparison of corresponding information transfer rates of R1-R6 recordings and stochastic model simulations to light bursts and Gaussian white noise (GWN) stimuli of different bandwidths. The recorded and simulated information transfer estimates correspond closely over the whole tested encoding space (*cf.* **figures S2** and **S6**). (**B**) Their differences to light bursts help to identify extra information in the recordings, which likely comes from the lamina network (through gap-junctions (*29*) and feedback synapses (*30*)) to individual photoreceptors. The clear variability between different recordings from individual cells (continuous thin lines) indicates that some R1-R6s may receive up to 200-250 bits/s of information from the network, whereas others receive less (cyan background). Some recordings likely contained more instrumental/experimental noise (pink background), which could render their information transfer rates (in particular to low-frequency bursts) less than that of the model; some of this noise likely comes from low-frequency eye and photoreceptor movements (*cf.* **fig. S3**). Thick line and error bars give the average information transfer rate difference between the recordings and the model (~ 0-50 bits/s). The data implies that the extra network information to R1-R6s in vivo is mostly at high burst frequencies (100-500 Hz).

Each R1-R6 receives different amounts of information from the network
In **Box S2.4B**, the difference in information rate estimates between the corresponding recordings and simulations to the bursty stimuli is plotted for each complete recording series (thin lines) of individual cells. The thick line gives the mean difference to all these cells' performance. Most noticeably, some photoreceptor cells carry ~100-200 bits/s more information from the bursty stimuli, but many other cells also show information rates that surpass the model's performance (*cf.* **fig. S7**). Any information surplus (cyan background) presumably comes from the lamina network (*29, 30*); through gap-junctions and feedback synapses from the cells that sample information from the same small visual area (due to neural superposition (*65-67*)). The photoreceptor model lacks this network information.

Box S2.5. Synaptic connectivity between neurons within a lamina cartridge. Synapse numbers color-coded as indicated by the column on the right. Large monopolar cells, L1-L5; amacrine cell, Am; C2 and C3 are retinotopic centrifugal fibers from the next synaptic processing layer, medulla. Image from (28)

In fact, each of the six R1-R6s, which pool their inputs in the same lamina cartridge for feed-forward synaptic transmission, should show different information transfer rates. This is because the lamina connections are asymmetric (**Box S2.5**). Electron micrographs have shown that R1, R2, R3, R4, R5 and R6 make different amounts of feedback synapses with the lamina interneurons (28, 68). Most feedbacks are provided by neurons belonging to the L2/L4 circuits (28, 68). Whilst same-cartridge connections are selectively from L2 to R1 and R2 and from L4 to R5, all R1-R6s receive feedback signals from L4 of neighboring cartridges. There are further connections from Am to R1, R2, R4 and R5, and glia are also synaptically connected to the network (28, 68), but only R6 makes direct gap-junctions (69) with R7 or R8. These asymmetric functional connections (30) may largely explain the variability in photoreceptor output (**fig. S1B**) and information rates (**fig. S2C**).

Our recent work (29) further showed that during naturalistic stimulation R6 can receive up to ~200 bits/s of information from R8, as channeled through gap-junctions between these cells. Therefore, we infer here that the two recordings with the highest information transfer rates (~850 bits/s) were probably of R6-type, which directly receive extra information from its R8y and R7y neighbors (29, 69, 70) (**Box S2.6**). Conversely, the recordings, in which information rates are lower than those of the simulations (**Box S2.4B**, pink background), carried presumably more recording/experimental noise, with one potential source being minute retinal movements (see **Text S4, Text S6-9**).

Box S2.6. Because of axonal gap-junctions between R6 and R7/R8 photoreceptors in the lamina (69, 70), R1-R6s that have been genetically engineered to express UV-sensitive Rh3-rhodopsin ("UV-flies") can still respond to green light by different degrees (29). This flow of extra "color"- information can be readily identified in intracellular responses of different R1-R6 photoreceptors in the same "UV-fly" to very bright UV (385 nm) and green-yellow (505 nm) flashes. (**A**) First cell responded to UV but not to green. (**B**) Next cell (likely R6 in the same or neighboring neuro-

ommatidium) responded to both UV and green. This cell cannot be R7y/p, which are less green-sensitive, or R8y/p, which are less UV-sensitive. Inset highlights a hypothetical recording path, somewhere close to the retina/lamina border (red arrow), and gap-junctions (black arrows) between photoreceptor axons. Histaminergic L1 and L2 cells receive visual information from R1-R6 photoreceptors' output synapses in the same neuro-ommatidium. (**C**) Another cell responded to UV and weakly to green-yellow. Modified from (*29*).

Box S2.7. R1-R6 photoreceptors' response waveforms and frequency range of reliable encoding vary cell-to-cell, and this variation does not reflect recording quality. Intracellular responses to the same repeated stimuli recorded in vivo from two different R1-R6s (Cell #18 and Cell #12) in two wild-type *Drosophila*. (**A**) 20 Hz light bursts drive both cells vigorously, but the output of Cell #18 rises and decays faster. Both outputs have maximum signal-to-noise ratios (SNR_{max}) > 5,000, but because of its faster response dynamics Cell #18 encodes better high stimulus frequencies. (**B** and **C**) for 100 and 500 Hz light bursts, respectively, Cell #12's output has a higher SNR_{max} but again lags behind Cell #18's output. The corresponding information transfer rate estimates, R_{output}, for Cell #18 are 787 bits/s (20 Hz bursts), 850 bits/s (100 Hz) and 625 bits/s (500 Hz) and for Cell #12: 569 bits/s (20 Hz), 711 bits/s (100 Hz) and 503 bits/s (500 Hz). *Cf.* fig. S7.

Our intracellular recordings establish that during bright light stimulation, the voltage output of an individual photoreceptor is highly repeatable (*cf.* **Fig. 1**). Consequently, our recording system could be used to study variability among individual R1-R6 photoreceptors of the fly eye. We discovered that for the same stimuli the characteristic output waveforms and frequency distributions of one particular cell are typically different to those of another photoreceptor (**Box S2.7**), even when recorded from the neighboring cells in the very same eye (by the same microelectrode). Because the signal-to-noise ratios of the recordings were very high (**Fig. 2A** and **fig. S2A**), sometimes over 6,000, it was evident that the observed cell-to-cell variability had little to do with the quality of the recordings. Hence, in the *Drosophila* retina, R1-R6 photoreceptors show intercellular variability that is far greater than the observed intracellular variability.

Collectively, these results strongly suggest that every R1-R6 photoreceptor, which is pooled in one lamina cartridge under the developmental neural superposition principle (*65*) to transmit information about light changes in a small area of visual space to visual interneurons (L1-L3 and Am) (*28, 30, 68, 71*), has, in fact, its own unique output. Besides asymmetric connectivity within a neuro-ommatidium, some of the observed variations in R1-R6 photoreceptors response properties may also reflect different microelectrode recording locations

across the fly eyes. For example, *Drosophila* photoreceptors in the front of the eye might show slightly different responsiveness to those at the back, as already shown for unique localized polarization-sensitivity differences (*72*). Compound eyes of many insects exhibit structural adaptations that alter their lens sizes and shapes locally, such as bright or acute zones for increasing sensitivity or resolution, respectively (*73*). Furthermore, electrophysiological recordings in some fly species suggest that their photoreceptor output vary across the eyes and could be tuned to the spatial and temporal characteristics of the light environment (*74-76*).

Variable sampling matrix protects from aliasing, improving vision
With each R1-R6 photoreceptor having variable "network-tuned" (and possibly "location-tuned") encoding properties and output, and with each image pixel being sampled through variable size rhabdomeres (see **Text S5**, **Box S5.1**) and ommatidial lenses (the photoreceptors' receptive fields vary; see **Text S4**, **Box S4.6** and interommatidial angles change progressively from front to back (*32*)), the *Drosophila* eye should generate reliable neural estimates of the variable world. This is because a sampling matrix made out of variable pixels (neurommatidia), in which size and sensitivity show random-like constituents,

- *prevents aliasing of image information* (*40, 58, 77*) (**Box S2.8**).
- *mixes color information to the R1-R6 motion vision channel*, whitening its spectral sensitivity (*29*) (**Box S2.8E**), which is a prerequisite for an optimal motion detector (*78*).

Aliasing effects are reduced by sampling faster and/or with finer spatial resolution, and eliminated by sampling more than twice over the highest stimulus frequency (*79*). The Nyquist–Shannon sampling theorem establishes a sufficient condition for a sample rate that enables a discrete sequence of samples to capture all the information from a continuous-time signal of finite bandwidth. Specifically, it only applies to a class of mathematical functions having a Fourier transform that is zero outside of a finite region of frequencies. This condition, however, cannot be fully realized in sensory systems, which show finite spatiotemporal sampling resolution and evolved around $1/f$-stimulus (*80, 81*) distributions of the real-world objects and events. Because any physical transformation affects signal and noise equally (data processing theorem (*25, 79*)) and because real-world low-pass filters, such as a lens, cannot cut-off sharply at an exact point, but instead gradually eliminate frequency components and exhibit a fall-off or roll-off slope, aliasing effects would not be removed completely in an ordered sampling matrix. Therefore, to prevent phantom sensations of aliased signals fooling the brain and perception of physical reality, sampling matrixes of sensory systems must entail stochastic variations.

Box S2.8. Projected image of Sine(x^2+y^2) function is used to illustrate the effect of aliasing and how stochastic variability in the sampling matrix combats aliasing effectively. (**A**) Sine(x^2+y^2) is plotted with 0.1 resolution. (**B**) Under-sampling the same image by an ordered matrix leads to aliasing: ghost rings appear when the image (of the function) is sampled with 0.2 resolution. Aliasing is a critical problem, as the nervous system cannot differentiate the fake image rings from the original real image. (**C**) Sampling the image (A) with a random matrix may lose some of its fine resolution, due to broadband noise, but such sampling is anti-aliasing; sampling with random points at 0.2 resolution. (**D**) Color photoreceptor distributions across macaque (*82*) (red, green and blue cones; left) and *Drosophila* retina (*83*) (R7y and R7p receptors; right) show random-like sampling matrixes, suggesting that this sampling matrix sensitivity randomization would have an anti-aliasing role. **E**, Crucially, by integrating and redistributing R1-R6 outputs with additional gap-junctional inputs from randomized R7/R8 color channels (*29*) (**D** and **Box S2.6**) for each image pixel during synaptic transmission to LMCs, any broadband sampling noise should be much reduced and the R1-R6 (motion) channel's spectral range whitened (*29*). Note how LMC output peaks before the corresponding R1-R6 output. Scale bars: 10 mV / 20 ms. Sub-figure **E** modified from (*29*).

Whilst temporal and topological sampling matrix variations in retinae combat aliasing (*40, 58*), their trade-off is broad-band noise (*77*) (**Box S2.8C**). This noise, however, is much reduced (or nearly eliminated) by parallel sampling of the same information (*5, 40*). For example, noise reduction occurs naturally in the fly eye - both in time and in space. In every R1-R6 photoreceptor, 30,000 microvilli sample discrete information stochastically in time, generating virtually aliasing-free macroscopic responses of very high signal-to-noise (**Box S2.7**). Whereas, across the lamina neuro-ommatidia of variable connectivity and spectral sensitivity, neural superposition integrates local R1-R8 signals of overlapping information from each pixel (a small largely aligned area in the visual space) to improve the signal-to-noise ratio of the sampled images (see **Text S5**). Such images should provide the brain reliable and maximally informative estimates of the environment.

Estimating a R1-R6's encoding efficiency

A photoreceptor's encoding efficiency, η, is the ratio between the information rates of the voltage output, R_{output}, and the effective light input, R_{input}, that drove it:

$$\eta = \frac{R_{output}}{R_{input}} \tag{S2.2}$$

As we already had determined the maximum photoreceptor output information rates, R_{output} (**Eq. S2.1**; **Figs 2C**, **figs S2C** and **S6C**), only the corresponding information rates of the effective light stimuli, R_{input}, needed to be worked out. Because the output simulations' maximum information rates matched well the corresponding mean rates of the real recordings (**Box S2.4A**), we had extrapolated successfully each effective light intensity time series that drove the voltage response. Therefore, we could now estimate the rate of information transfer of the light input by making the following assumption:

- Photon emission from the light source (LED) follows Poisson statistics; this may or may not be true (see the discussion below)

Photons are thought to be emitted by the light source, such as the LEDs, at random, exhibiting detectable statistical fluctuations (shot noise). Such dynamics can be modelled by Poisson statistics (*5*). Therefore, as each light stimulus trace differs from any other, with their mean equaling their variance, we could estimate through simulations their average signals and noise, and signal-to-noise ratios, $SNR_{input}(f)$. The corresponding information transfer rates, R_{input}, could then be estimated by Shannon's equation (Eq. S2.1). For each tested stimulus pattern, this was done by using the same amount of simulated input data as with the output data (2,000 points x 20 repetitions) to control estimation bias. More details and examples about Poisson stimulus simulation procedures are given in (*5*).

For the simulated inputs and outputs, the data processing theorem (*25*) dictates that $R_{input} \geq R_{output}$; thus $\eta \leq 1$ ($\leq 100\%$). If not, then one or both estimates are biased or incorrect; information cannot be created out of nothing. However, for the efficiency estimates based on the real recordings, it is quite possible that $R_{output} > R_{input}$, and thus $\eta > 1$ ($> 100\%$), because R1-R6s receive extra information from the network (**Boxes S2.5** and **S2.6**) that is missing from the R_{input} estimates (*cf.* **Box S2.4B**).

We recognize that there are methodological limitations and unknowns, which may affect the accuracy and consistency of these estimates:
- Experimental and theoretical evidence suggests that photon output of some light sources might be sub-Poisson (*84*); meaning, not maximally random. If this were true for our LED, then our approach would slightly underestimate the information content of the light input, used in the experiments, and consequently overestimate *Drosophila* photoreceptors' encoding efficiency.
- Shannon's equation can bias information transfer rate estimates for any corresponding light input and photoreceptor output differently. This is because the signal and noise components of the input and output may deviate from the expected Gaussian by different amounts. Even though we used systematically the same amount of data for both estimates (20 x 2,000 data points), in the cases where light distribution is skewed (bursty stimuli) but the photoreceptor output is more Gaussian, it is possible that Shannon's equation would underestimate input but not (or less so) output information, causing us to overestimate efficiency.
- Small datasets limit analyses. In the past, we have compared information transfer rate estimates, as obtained by Shannon's equation to those estimated through the triple extrapolation method (*44*), which is directly derived from Shannon's information theory. For ergodic datasets of different distributions, and when appropriately applied, both methods provided similar estimates (*5, 44*). However, the triple extrapolation method works best with large datasets; preferably containing >30 responses to the same stimulus (*44*). In the current study, because of the practical limitations (to map a photoreceptor's whole encoding space within a reasonable time), all the selected recordings and simulations consisted responses to 20 stimulus repetitions. This data size was deemed insufficient for an accurate estimate comparison between the two methods and was not done here. In the analyses, to provide fair comparison between simulations and recordings in all tested conditions, all the data chunks (for the recordings, simulations and

stimuli) were exactly the same size (20 x 2,000 points) and they were processed systematically in the same way (apart from the two exceptions we discuss next). Therefore, the data-size bias should be under control and the results comparable within these limits.

- Implementation of Shannon's equation (Eq. S2.1) in digital computers typically requires windowing of the data chunks (for signal and noise) before calculating their power spectra though Fast Fourier Transfer (FFT). Windowing combats spectral leakage (smearing), but this affects especially low-frequency signals, in which information content is low. So this trade-off can be considered reasonable, and its effect on most performance estimates is marginal. But here as the input and output information transfer is calculated separately, windowing affects more 20 Hz GWN light input than its corresponding photoreceptor output. This is because windowing clips lower frequency power from 20 Hz GWN input, whereas in the simulated and real voltage responses much of this power is nonlinearly translated (through adaptation) to higher frequencies, including those over 20 Hz. The simulated light input, of course, carries no information on frequencies >20 Hz, but now it has also lost in windowing some of its low-frequency modulation, which the photoreceptors could translate into high-frequency voltage modulation (note, photomechanical phase enhancement can further contribute to this nonlinearity, see **Text S3**). For the two lowest intensity levels only, we judge that because of this methodological bias, the efficiency estimates for the 20 Hz GWN input-output datasets became unrealistic by a small margin of 10-40 bits/s, implying that R_{output} > R_{input}. Therefore, for these two stimulus dataset only, we applied box-car windowing (instead of the normal Blackman-Harris type), to retain its low frequency information content, and so to reduce this bias.

Because of all these possible error and bias sources, a *Drosophila* photoreceptor's encoding efficiency (η) estimates given in this publication must be considered as upper bounds. Nonetheless, for real photoreceptors, it is realistic to expect their maxima to approach 100%, and in some cells (likely in R6s) be beyond, for the tested low-frequency stimuli (20 Hz). This is because of the extra information from the network, which is missing from the simulated light stimulus estimates. Overall, these maximum values are slightly higher but consistent with our previous (conservative) estimates of 90-95%, in which the light input intensity to microvilli was inferred by comparing the wild-type photoreceptor performance to that of white-eye mutants, lacking the intracellular pupil. Therefore, we conclude (again conservatively) that the error margin of these new encoding efficiency estimates may reach ±5%.

Text S3: Similarities and differences in encoding bursts and other stimuli

Overview

This supplement explains why bursts encoding is both highly informative and reliable, proving additional insight to the results presented in **Figures 1-2** in the main paper.

Why do bursty responses carry more information than those to natural stimuli?

Figure S4 shows that R1-R6 photoreceptors can sample more information from maximally bursty light patterns than from highly dynamic naturalistic light intensity time series ($1/f$-stimuli). We can explain this higher performance by the bursty stimuli' proportionally more, and more evenly distributed, long dark contrasts. Such events enable refractory microvilli to recover efficiently from their previous light-activation so that large numbers of them are continuously available to sample ongoing light changes. This leads to larger sample rate changes and, thus, to a higher rate of information transfer (*cf.* (*5*)).

Moreover, fast high-contrast events survive the slower intracellular pupil mechanism (**Text S2**, *e.g.* **Box S2.2**) and photomechanical rhabdomere contractions (**Text S7**) well (**Fig. 4C**). And consequently, fewer photons are being filtered out (lost) from high-contrast bursts than from naturalistic stimulation (**fig. S4B**, pink trace) or Gaussian white-noise (grey), which adapt the photoreceptors more continuously to the given light background. Therefore, at the level of light source, bursty stimuli can have much lower power than the other two stimuli to drive photon-to-quantum bump sampling efficiently by the photoreceptor's 30,000 microvilli (see **fig. S2D**).

Relevance of bursty responses for seeing germane visual patterns

From an information theoretic point of view, the amount of visual information that is encoded by a photoreceptor can be similar for bursty (phasic) and Gaussian (tonic) signals (**Fig. 2C**; *cf.* 20 Hz 0.6-contrast bursts and 100 Hz 0.32-contrast GWN). However, when signal-detection-theoretic measures are applied, bursts outperform 'tonic' GWN signals in indicating visual "things"; i.e. the occurrence of perceptually relevant changes in light input. Bursts appear somewhat like all-or-none events (**Box S3.1**), having much higher local signal-to-noise ratios (in rising or decaying phases) in a photoreceptor's voltage responses (*30*). They tower over the background noise, making their detection much easier than for the tiny blips of the GWN signals. Accordingly, photoreceptors' voltage bursts support robust transmission of behaviorally relevant visual information and should further improve the reliability of synaptic transmission and perception (*30, 71*).

Box S3.1. Whilst having similar information transfer rates, the responses to bursty stimuli (left) show higher signal-to-noise ratio within brief (10-100 ms) perceptually relevant time windows than the responses to Gaussian white-noise (GWN, right). The data are from the same cell. **A**, 25 intracellularly recorded voltage responses to repeated bursty and GWN light stimuli. Individual responses (superimposed) are shown in light grey and their mean in black traces. Both of these responses series carry similar information contents (~350 bits/s), as estimated by Shannon's equation (see Eq. S2.3). **B**, The signal (average response) standard deviation (SD) in 10 ms windows to bursty stimulation vary much more than that to white-noise. The noise variability (blue traces) in the two set of responses (SD in 10 ms time windows) is similar. **C**, Signal-to-noise ratio is much greater in the responses to light bursts than to white-noise stimulation; it was calculated as the ratio between signal SD and noise SD, using 10 ms time resolution.

Text S4: Spatial resolution (visual acuity) of the fly eye (conventional measure)

Overview

This supplement describes in detail a new method to measure a *Drosophila* photoreceptor's receptive field, and provides important background information about the experimental and theoretical results presented in **Figures 3-5** in the main paper.

In this supplement:

- We test the hypothesis that in the *Drosophila* eye visual information is integrated laterally in dim conditions, and fed back synaptically to its photoreceptors, contributing to their spatial responsiveness.
- We measure dark- and moderately light-adapted wild-type R1-R6 photoreceptors' receptive fields by their acceptance angles, $\Delta\rho$, using intracellularly recorded voltage responses to light flashes, delivered from randomized positions of an orthogonal stimulation array.
- We compare these measurements to those of histamine-mutants (*34, 48, 85*) (hdc^{JK910}), in which first-order interneurones are blind (receiving no neurotransmitter from photoreceptors) and so incapable of feedback-modulating the photoreceptor output.
- We show that the average acceptance angles of dark-adapted wild-type photoreceptors are 10.9% broader than those of the mutant, while light-adapted cells show no such difference.
- We further quantify the effect of stimulus history on photoreceptors' receptive fields.
- We characterize slow spontaneous retinal movements in the *Drosophila* eye and show how this activity can influence intracellular photoreceptor recordings.
- Our results suggest that in dim conditions spatial information is pooled in the lamina and fed-back to wild-type photoreceptors. Such excitatory lateral synaptic modulation, which is missing in the mutant, increases spatial sensitivity, broadening the photoreceptors' receptive fields.

Optical limits of the fly compound eyes' visual acuity

Visual acuity is defined as the minimum angle that the eye can resolve. In the fly compound eye, if each ommatidium constitutes a sampling point in space, then the eye's maximal spatial resolution is set by the density of its ommatidial array (*86*). Suppose a regular pattern of black and white stripes is presented to a fly. The maximum spatial frequency that the fly can resolve, v_s, is achieved when one ommatidium points to a black stripe and its adjacent ommatidium points to the next white stripe (**Box S4.1A**). Thus, the interommatidial angle (*87*), angle, $\Delta\varphi$, is the key parameter in determining v_s. For the compound eyes with hexagonal layout, as is the case of most flies, the effective interommatidial angle, $\Delta\varphi_e$ (**Box S4.1B**), can be calculated by the following equation:

$$\Delta\varphi_{e=} \cos(30°)\,\Delta\varphi = \frac{\sqrt{3}}{2}\Delta\varphi \tag{S4.1}$$

Thus, the upper limit of the fly eye's visual acuity is given by:

$$v_s = \frac{1}{2(\Delta\varphi_e)} = \frac{1}{\sqrt{3}(\Delta\varphi)} \tag{S4.2}$$

Whether this limit is achieved or not depends upon the spatial performance of a photoreceptor (*88*). However, when estimating a photoreceptor's receptive field, which is quantified by its width at half-maximum, or acceptance angle (*89*), Δρ, one needs to consider several contributing factors.

Firstly, since the ommatidium lens and a photoreceptor's rhabdomere are very small, optical quality is strongly affected by light diffraction, of which airy pattern (the point-spread function) depends upon light wavelength, λ, lens diameter, *D*, rhabdomere diameter, *d*, and focal distance, *f* (**Box S4.1C**). Theoretically, the blurring functions at the ommatidium lens and rhabdomere tip are both Gaussian and therefore can be combined(*88*) to yield a simple approximation of Δρ:

$$\Delta\rho = \sqrt{\left(\frac{\lambda}{D}\right)^2 + \left(\frac{d}{f}\right)^2} \qquad\qquad (S4.3)$$

However, owing to the rather complex waveguide properties of small-diameter rhabdomeres, this formula is somewhat inaccurate. Van Hateren (*90*) and Stavenga (*39, 91*) found that along a fly photoreceptor's rhabdomere only a limited number of light patterns (modes) could be formed and that this number depends upon the incident angle of light, leading to a smaller actual Δρ than what Eq. S4.3 implies.

Another contributing factor is the spatial cross talk, in which a photon escapes the rhabdomere it first travels in and enters an adjacent rhabdomere (*92*). Such an effect is likely to happen when the cross-talk index of the ommatidia/rhabdomere structure is less than three (*93*). This was indeed reported for *Drosophila* (*32*), suggesting that its neural images might have lower resolution than theoretically calculated from the optics.

Lastly, the intracellular pupil mechanism further affects Δρ estimation. Inside each photoreceptor cell, there are tiny pigment granules that migrate toward its rhabdomere boundary upon light adaptation (see **Text S2**, **Box S2.1**). These pigments absorb and scatter light that travels inside the rhabdomere, reducing the light influx absorbable by its rhodopsin molecules (*94-96*). Consequently, the pupil mechanism shapes a photoreceptor's angular and spectral sensitivity (*63*). Moreover, in **Text S2**, we show by experiments and theory that it further helps to maximize a photoreceptor's information transfer; by optimizing the light intensity passing into the rhabdomere.

Box S4.1. Classic theories of compound eye optics. (**A**) It is assumed that the minimum angle a compound eye can resolve is its interommatidial angle, Δϕ. (**B**) The effective interommatidial angle of an eye with a hexagonal layout, Δϕ$_e$, is smaller than its actual interommatidial angle, Δϕ. Eq. S4.1 gives their geometrical relation. (**C**) Light diffraction at the ommatidial lens and the rhabdomere tip strongly affects the optical quality of the image pixel that a photoreceptor samples. D = ommatidial lens diameter; d = rhabdomere tip diameter; f = focal length; λ = wavelength. Redrawn from (*2*).

Rationale for investigating wild-type and *hdc* R1-R6s' receptive fields
The fly compound eyes are small and size-constrained, presumably to save energy and improve

survival (*97-99*). This puts their sensitivity/acuity trade-off under intense evolutional pressure (*86, 100-102*). While an increase in ommatidium size would improve photon capture, it would also result in fewer sampling points (image pixels) in the eye, lowering the resolution of its neural responses (neural images of the world). In dim conditions, where photon noise is relatively large compared to available information (signal), the task to enhance visual reliability and sensitivity becomes challenging. Optical mechanisms, including the widening of photoreceptor receptive fields by pupil opening (*63, 103-105*), can increase the amount of light collected in each ommatidium only to some extent. Yet under dim illumination, insect visual behaviors appear remarkably robust (*106-108*), suggesting that their eyes' neural mechanisms could successfully overcome the apparent shortfall in photon supply (*109*).

Sensitivity can be increased neurally at the cost of decreasing acuity by (**i**) increasing photoreceptors' voltage/light intensity gain (*6, 110, 111*), (**ii**) increasing their integration time (temporal summation) (*6, 112*) and (**iii**) spatially summing information, or reducing lateral inhibition from neural neighbors (*113-115*). Experimentally, spatial summation has been shown to occur in the directionally-selective motion-detecting (DSMD) neurons of the fly lobula plate (*116*), but it is possible that such signals might also reflect upstream processing in the earlier optic neuropiles (the lamina, medulla and lobula).

Although the fly photoreceptor biophysics for adapting temporal summation are well characterized (*5-7, 40, 45, 51, 64*) (see **Texts S1-3**), the neural substrate for spatial summation is less well understood. In the lamina, electrical couplings by gap-junctions were found only between the photoreceptor axons that share the same optical axis (*69, 70, 117, 118*) in neural superposition. Hence, these presynaptic connections probably cannot distribute spatial information. Nonetheless, postsynaptically, the evidence is more suggestive. Intracellular responses of the histaminergic interneurons (large monopolar cells, LMCs) to narrow (point source) and wide-field light stimuli match well the theoretical predictions of spatiotemporal summation (*46, 119, 120*). This notion was further advocated by the structural study in the nocturnal bee *Megalopta genalis* (*121*) lamina, which revealed extensive synaptic connections between adjacent cartridges. Finally, early functional studies of the housefly (*Musca domestica*) photoreceptors (*119*) indicated that quantum bumps, recorded to dim light at the behavioral threshold, contain additional small-amplitude events. These were judged not to be generated by the impaled cells but by single photon captures in their neighbors; with the receptive fields being wider than what were expected from the optics alone (*122*).

In this supplement, we test whether or how spatial information is integrated laterally and fed back synaptically to photoreceptors, contributing to their acceptance angles, $\Delta\rho$. The tight coupling between feed-forward and feedback pathways in the photoreceptor-lamina circuit is known to have crucial roles in maintaining robust adaptation and temporal coding efficiency (*30, 71, 123*). Theoretically, similar spatial information regulation should further improve fly vision. Specifically here, we take advantage of *Drosophila* genetics and compare R1-R6 photoreceptor outputs of wild-type and hdc^{JK910} mutant. Synaptic transmission from hdc^{JK910} photoreceptors is blocked, making their interneurons effectively blind (*35*). Therefore, feedback from the mutant LMCs (and possibly from amacrine cells (*30, 124*), Am, which also receive histaminergic input from photoreceptors) to R1-R6s cannot contain any lateral modulation, neither inhibitory nor excitatory.

We show that the dark-adapted wild-type photoreceptors' mean acceptance angles are 10.9% broader than in the mutant, while no significant differences are found between the light-adapted cells. We further show how stimulus history and retinal movements affect the receptive

fields in the *Drosophila* eye. Our results suggest that in dim conditions spatial information is pooled in the lamina and channeled back to R1-R6 photoreceptors in the form of excitatory synaptic modulation, which increases spatial responsiveness by broadening the cells' receptive fields.

Measurement and calculation of a R1-R6's receptive field
A photoreceptor's receptive field can be estimated electrophysiologically by measuring its intracellular response amplitudes, V_n, to a light flash intensity, I_n, at varying angular positions, α_n. From all these V_n, I_n, and α_n values generated by a complete scan, the receptive field width can be computed by three different methods as comparatively reviewed below.

Method 1. V_n is clamped to a constant value in a closed-loop system, which accordingly vary I_n for each tested light source positions (*125, 126*). Sensitivity at each position, S_n, is then defined by:

$$S_n = \frac{I_o}{I_n} \tag{S4.4}$$

where I_0 is the intensity required from a point source at the center of the receptive field. The definition of sensitivity can be equivalently expressed as the light source at an off-axis position. The off-axis light intensity needs to be $\frac{1}{S_n}$fold brighter than the axial one to stimulate responses of the same amplitude.

   After corresponding S_n was computed for every S_n, the sensitivity-angle relation is fitted by a Gaussian function. The width at the half-maximum of this Gaussian curve is called the *acceptance angle*. This is the conventional parameter, $\Delta\rho$, for quantifying the receptive field width.

Method 2, in which the same light flash intensity I_n is tested at many different angular positions, is the most widely used (*32, 92, 127-129*). Initially, the $V/\log(I)$ relation of the impaled photoreceptor is determined at the center of its receptive field by presenting logarithmically intensified flashes from a point-like light source (through scaled neutral density filters). V_n elicited by the light at each off-axis angle, I_n, is then substituted into the $V/\log(I)$ function to estimate, I_a, the light intensity that was effectively absorbed by the cell's photopigments. Angular sensitivity, S_n, is given by Eq. S4.5:

$$S_n = \frac{I_a}{I_n} \tag{S4.5}$$

Gaussian fitting and the acceptance angle calculation are performed as in Method 1.
   Method 2 is based upon the same principle as Method 1, which is to assess the light intensity necessary to elicit a criterion response (*130*). When the light-point with intensity I_0 is exactly at the optical axis of the cell, sensitivity is the highest with the response amplitude V_0. To evoke $V_n = V_0$ by a light source located at an angular position, α_n, it is required that the effective intensity I_a equals to I_0. Given the angular sensitivity formula:

$$S_n = \frac{I_a}{I_n} = \frac{I_o}{I_n} \tag{S4.6}$$

the necessary intensity I_n would be $\frac{1}{s_n}$fold brighter than I_o.

Though Method 2 does not require a closed-loop system and is, therefore, less experimentally challenging, its effective intensity, I_a, estimation has drawbacks. Fitting V/log(I) function to a small number of maximum amplitude values, which are adaptation-dependent, can introduce scaling errors. Whilst its underlying assumptions, that the voltage/(effective intensity) relation is static and independent of light source position, neglect possible dynamic and lateral interactions between neighboring cells.

Nonetheless, the outcomes of both methods are theoretically independent of photoreceptor biophysics and the test flash intensity. Hence, they enable electrophysiological receptive field measurements to be compared with those derived from optical, morphological and waveguide theories.

Method 3. Similar to Method 2, each tested photoreceptor is stimulated by the same light intensity flashes at different angular positions around its optical axes. Response amplitude V_n to a light flash coming from an angle α_n is then normalized to the maximum response evoked by the on-axis light source, V_0. The receptive field is determined by the Gaussian fitting of the relation between ratios V_n/V_0 and incident angle α_n. This may yield wider half-maximum widths ($\Delta\rho$ values) than the acceptance angles (*131, 132*) estimated by using Method 1 and Method 2.

We used Method 3 to estimate R1-R6 photoreceptors' receptive fields from intracellular recordings, despite its disadvantages; the results would depend on the flash intensity and would not be fully comparable to other approaches and the previous studies in *Drosophila*. Our main rationale was that this method characterizes "how well the flies see" most directly and reliably, without making any assumptions about lateral interactions between photoreceptors and LMC feedbacks. Moreover, the method's limitations should not compromise our objectives to compare the receptive fields in different genotypes and to report how these are affected by different light conditions and stimulation history. And importantly, these receptive field estimates could be directly used in further calculations to assess the same photoreceptors' theoretical acuity to detect moving objects, as shown in **Texts S6-8**. Experimentally, it was also unfeasible to expand our recording set-up either with a closed-loop system (as in Method 1) or with easily exchangeable neutral density filter sets for characterizing $V/log(I)$ function (as in Method 2).

25 LED light-point array and LED pads. A R1-R6 photoreceptor's receptive field was scanned by using an array of 25 light-points, mounted on a Cardan arm (**Box S4.2**). Each light-point (small light guide end) subtended an angle of 1.7° as seen by the fly, transmitting its light output from its specific LED (1/25). The system was controlled by 2 channels, both with voltage inputs ranging from 0 V to 10 V. Channel 0 was used to select the light-point while Channel 1 was used to linearly set its intensity.

Box S4.2. 25 light-point stimulus array. Each tested dark-adapted photoreceptor's receptive field (red Gaussian) was assessed by measuring its intracellular responses to successive flashes from 25 light-points. In light-adaptation experiments, two 39-LED pads, (on both sides of the vertical stimulus array) provided background illumination. The intact fly was fixed inside the conical holder, which was placed upon a close-looped Peltier-element system, providing accurate temperature control (at 19 ^{o}C). The rig was attached on a black anti-vibration table, inside a black-painted Faraday cage, to reduce noise and light scatter.

Box S4.3a shows a typical light-point's spectral density measured at the light-guide end. Each light-point had a narrow spectral **Peak1** at ~450 nm and a broader **Peak2** at ~570 nm, in which intensities and wavelengths are listed in **Box S4.4**. Based on their relatively small variations, all 25 light-points provided reasonably uniform light input, except the 4-times brighter No.22 (see below). This is because a fly photoreceptor's response amplitudes differ only marginally until light intensity changes several-fold, as defined by the sigmoidal V/log(I) relationship (*111*). Standard light flashes, containing ~2×10^6 photons/s at **Peak1** and ~3×10^6 photons/s at **Peak2**, were produced by setting Channel 1 to an input value of 2 V. Here, the estimated photon counts are given at the light source, not at the level of photoreceptor sampling. Moreover, in the experiments, to evoke subsaturating responses, we used a neutral density filter plate to cut the light-point intensity by 100-fold.

Box S4.3. Spectral properties of the light stimuli. (**A**) Typical spectral density of the light impulses delivered by the 25-point array. Note the spectra has two prominent peaks, named **Peak1** (~450 nm) and **Peak2** (~570nm). (**B**) Spectral density of a single LED on the two Lamina pads, which were used to provide ambient background illumination during light-adaptation experiments. These spectral intensities (photon counts) were measured by a spectrometer (Hamamatsu Mini C10082CAH, Japan).

Light-point	Peak1 wavelength (nm)	Peak1 intensity (10^6 photons/s)	Peak2 wavelength (nm)	Peak2 intensity (10^6 photons/s)
No.1	448	2.720	571	2.805
No.2	452	1.790	570	2.818

No.3	448	2.618	565	2.900
No.4	451	1.840	576	3.020
No.5	451	2.570	575	3.710
No.6	451	2.214	572	3.640
No.7	452	1.430	575	2.020
No.8	446	2.203	570	2.990
No.9	453	1.465	571	2.350
No.10	451	3.300	578	5.100
No.11	453	1.877	575	3.080
No.12	455	1.763	575	3.020
No.13	451	2.334	575	3.440
No.14	451	2.009	576	2.634
No.15	454	2.400	568	4.480
No.16	452	2.390	572	4.165
No.17	455	3.190	566	3.010
No.18	452	1.990	578	3.320
No.19	455	1.958	578	3.336
No.20	454	1.745	569	2.670
No.21	450	2.642	573	2.314
No.22	452	9.520	572	13.300
No.23	452	2.420	575	3.380
No.24	452	2.658	573	3.284
No.25	452	1.670	570	2.750

Box S4.4. Light flash peak wavelengths and intensities delivered by each of the 25 light-points in the stimulus array. The given light intensities were measured at the light source by Hamamatsu Mini C10082CAH spectrometer, before 100-fold neutral density filtering. Thus, these values are estimated to be $10^{2\text{-}3}$-times higher than the corresponding effective photon rates at the level of R1-R6 sampling (see **Text S1**, **Box S2.2**). Accordingly at the optical axis, the center LED (No.13), with the neutral density filter on it, evoked subsaturating (~20-35 mV) responses from *Drosophila* R1-R6s (**Box S4.6C-D**).

Light-point No.22 was 4-fold brighter than the others. However, no attempt was made to correct its intensity for three reasons. Firstly, it was located at the tested receptive fields' periphery, and thus had limited influence on the measurements. Secondly, because this "error" occurred stereotypically in every experiment, the brighter No.22 would not bias the comparative studies (see below). Lastly, having one brighter light-point was beneficial for other experiments, as will be shown in **Text S6**.

Two Lamina LED pads, each with 39 similar LEDs, provided ambient illumination to

moderately light-adapt the tested photoreceptors. The pads were located at the outer half and outside each tested cell's receptive field (**Box S4.2**). Thus, a large portion of their light projected onto the neighboring photoreceptors. But the pads also illuminated the whole recording chamber, revealing its spatial structure and possibly inducing spatial processing in the lamina network. Light from each of the pad's LEDs peaked at 460 nm (**Box S4.3B**), delivering estimated ~2 × 10^5 photons/s.

Pseudo-random receptive field scans. Before recording intracellular voltage responses from a R1-R6 photoreceptor, we located its receptive field center. This was done by flashing the light-point No.13 (at the center of the array) and moving the array (with the Cardan arm along its XY-axes) until the maximum response amplitude was elicited. The array was then locked at this position.

In the dark-adaptation experiments, the photoreceptor faced darkness (**Box S4.5A**) for 30-60 s before its receptive field was measured. In the light-adaptation experiments, preselected background illumination (using the LED pads: **Box S4.2**) was turned on 30-60 s before the corresponding receptive field measurement.

Box S4.5. Measuring a R1-R6 photoreceptor's receptive field with intracellular recordings. (**A**) Schematic Image of how the LED stimulus array - seen as 25 light-points (light-guide-ends in a row) was centered by a Cardan arm system in respect to the studied photoreceptor and the fly eye. (**B**) Channel 0 input was used to select the LED (light-point) to be turned on. (**C**) Channel 1 input defined light intensity of the selected LED. A standard light impulse (flash) was produced by a 2 V input, which lasted 10 ms. (**D**) A photoreceptor's intracellular voltage responses to a complete receptive field scan. These sub-saturating responses were recorded at 19 °C. Amplitude V_n of each flash response was the local maxima. V_0 is the amplitude of the response to a light flash at the center of the receptive field (on-axis).

A complete scan of a photoreceptor's receptive field comprised 25 subsaturating flashes from all the light-points, one after another in a pseudo-random order (**Box S4.5B**). Each flash lasted 10 ms, and was 490 ms apart from the next one (**Box S4.5C**). Although this inter-flash-

interval largely rescued the photoreceptor sensitivity, spatiotemporal adaptation might have still affected their responses. For instance, a flash near the center of the receptive field would light-adapt the cell more than one at the periphery, possibly causing the response to the next flash to be artificially smaller. Therefore, by randomizing the order of flash positions - with the Matlab command *randperm(25)* - we could reduce this kind of potential adaptation effects.

Channel 0 input was turned on only when Channel 1 was set to zero V; that is, in the resting period when all the light-points were off. Otherwise, the transitions of Channel 0 input values would generate running dot images. Each tested photoreceptor's responses to 2-5 repetitions of pseudorandom scans were averaged (**Box S4.5D**) before the acceptance angle (or half-maximum width), Δρ, of its receptive field was determined (**Box S4.6**).

Box S4.6. Estimating a dark-adapted *Drosophila* R1-R6's receptive field and its half-width. Flash response amplitudes V_n were initially normalized to V_0, the maximum response elicited by an on-axis light-point. A Gaussian curve was then fitted to these normalized values, yielding an estimate of the receptive field. Half-maximum width of this Gaussian function, Δρ, defined the tested photoreceptor's acceptance angle. The schematic fly eye inset clarifies how a single photoreceptor integrates light from the world spatially through its receptive field (green area), whilst being bounded by the ommatidial lens system. For a standard measurement, each tested photoreceptor's intracellular responses to 2-5 repetitions of pseudorandom scans (as shown in **Box S4.4**) were averaged.

Gaussian white-noise (GWN) stimuli. Light-point No.13 intensity was controlled by setting Channel 0 input to 5 V and modulating Channel 1 input with a white-noise time series, which had a mean value of 2.5 V and cut-off frequency of 200 Hz. With these settings, light-point No.13 emitted 2.5×10^6 photons/s at **Peak1** (451 nm) and 3.75×10^6 photons/s at **Peak2** (575 nm) on average (**Box S4.4**). But, as in other experiments, these intensities were reduced 100-fold by neutral density filtering

Receptive fields of dark-adapted photoreceptors

In every experiment, we first assessed the recorded photoreceptor's receptive field after dark-adaptation (**Box S4.7A**). Wild-type R1-R6s' mean acceptance angle, measured as their receptive fields' half-maximum width, Δρ, was 9.47 ± 0.36º (± SEM, n = 19 cells), ranging from 7.00º to 11.65º. Interestingly, *hdc^{JK910}* R1-R6's receptive fields were 10.9% narrower (p = 0.0397, two-tailed t-test). Their mean, minimum and maximum acceptance angles were 8.44 ± 0.32º (n = 18 cells), 6.18º and 11.50º, respectively.

Because each photoreceptor's flash response amplitudes, V_n, were directly used to estimate its receptive field (**Box S4.7B**), rather than being converted to angular sensitivities (see above), the obtained Δρ metric depended upon the cell's output/input characteristics and the test flash intensity. To ensure that wild-type and *hdc^{JK910}* photoreceptors' Δρ comparison was unbiased by variable on-axis light sensitivities, we also compared their maximum response amplitudes, V_0, and $V_0/Δρ$ relations.

The corresponding V_0 values were very similar (**Box S4.7B**) and mostly within 20-35 mV sub-saturated linear range of the photoreceptors' V/log(I) curves (*35*). Moreover, in both wild-type and the *hdc* mutant, the linear correlations between V_0 and $\Delta\rho$ reflected only a weak trend of more sensitive photoreceptors (larger V_0) having wider receptive fields (larger $\Delta\rho$) (**Boxes S4.7C-D**).

Together, these findings indicate that the narrower $\Delta\rho$ of dark-adapted hdc^{JK910} photoreceptors neither resulted from altered phototransduction nor was an artefact of this measurement method.

Box S4.7. Dark-adapted wild-type and hdc^{JK910} R1-R6s' acceptance angles differ. The 25 light-point array stimulation evoked spatially variable photoreceptor output, providing a receptive field estimate for each tested cell as in **Boxes S4.5** and **S4.6**. (**A**) Dark-adapted wild-type photoreceptors' receptive fields (above), shown as the mean of their Gaussian fits (black), were wider than those of hdc^{JK910} photoreceptors (red). Their receptive field sizes (below) were quantified by the corresponding half-maximum widths, giving the mean acceptance angles: $\Delta\rho_{\text{wild-type}}$ = 9.47 ± 0.36°; $\Delta\rho_{hdc}$ = 8.44 ± 0.32°; p = 0.0397, two-tailed t-test. (**B**) Wild-type and mutant photoreceptors' peak responses (above), evoked by a sub-saturating 10 ms light flash (grey bar) at the center of the receptive field, showed similar dynamics and amplitudes, V_0 (below). $V_{0\text{wild-type}}$ = 28.77 ± 1.19 mV; V_{0hdc} = 28.11 ± 1.03 mV; p = 0.67, two-tailed t-test. This indicates that hdc^{JK910} phototransduction is functionally intact and wild-type-like. *Cf*. (*35*). (**C**) Linear correlation between $\Delta\rho$ and V_0 of dark-adapted wild-type photoreceptors. Adjusted R-squared = 0.1043. (**D**) Linear correlation between $\Delta\rho$ and V_0 of dark-adapted hdc^{JK910} photoreceptors. Adjusted R-squared = 0.072. (**A-D**) $n_{\text{wild-type}}$ = 19; n_{hdc} = 18. (**A-B**) Mean ± SEM; two-tailed t-test.

Receptive fields of light-adapted photoreceptors

A photoreceptor's $\Delta\rho$ measured under light-adaptation (at specific ambient illumination; **Box S4.2**) should be smaller than during dark-adaption. There are four reasons for this difference:

- Light-adaptation steepens a photoreceptor's V/log(I) function (*110, 111, 133*). This reduces the difference between I_0 and I_{a50} (or the effective intensity that could evoke response amplitude V_2^0), which in turn leads to a smaller α_{50} (or the corresponding angular position of I_{a50}) and thus to a smaller $\Delta\rho$, as reported by the chosen method.
- Because the test flash intensity was kept unchanged, their contrast would be lower during light-adaptation than in the dark-adaption experiments, further reducing the I_o/I_{a50} ratio, α_{50} and $\Delta\rho$.
- Light adaptation activates screening pigment migration, narrowing the intracellular pupil (*cf.* **Text S2**, **Box S2.1**). The narrower pupil effectively reduces the amount of light from

off-axis angles that can be absorbed by rhodopsin-molecules in the rhabdomere (*63, 125, 128, 134*), reducing Δρ.

- Theoretical studies and some experimental data suggest that in dim conditions neural signal summation from neighboring cells may enhance sensitivity. But in bright conditions, lack of summation, increased lateral inhibition or both improve image resolution (*109, 113, 114, 135, 136*).

To quantify how moderate ambient light affects the fly eye's spatial responsiveness, we analyzed the receptive fields of six wild-type (n = 6) and eight hdc^{JK910} (n = 8) photoreceptors both at their dark- and light-adapted states.

At the dark-adapted state, the acceptance angles of wild-type and hdc^{JK910} R1-R6s' receptive fields, Δρ, were 9.65 ± 1.06° and 8.16 ± 0.62° (mean ± SEM), respectively (**Box S4.8A**). However, owing to the smaller test and control group sizes than in **Box S4.7A**, this average difference (15.44%), though similar, was now statistically insignificant (p = 0.258, two-tailed t-test).

At the light-adapted state, under the given ambient illumination (**Box S4.8B**), the corresponding Δρ values were 7.70 ± 0.52° for wild-type photoreceptors and 6.98 ± 0.46° for their hdc^{JK910} counterparts. Thus, light-adaptation significantly reduced Δρ values from dark-adaptation (p = 3.49 x 10^{-4}, paired two-tailed t-test). Switching from the dark- to light-adapted states, wild-type photoreceptors' receptive fields narrowed down by 18.44 ± 3.5% (**Box S4.8C**), slightly more than those of mutants, which changed by 13.68 ± 3.37%. Yet, none of these parameters differed significantly between the wild-type and mutant photoreceptors.

Box S4.8. Light-adaptation narrows wild-type and hdc-mutant receptive fields similarly. Wild-type and hdc^{JK910} photoreceptors, of which receptive fields were assessed in both the dark- and light-adapted states, were compared. (**A**) Their dark-adapted and (**B**) and light-adapted Δρ values were similar. Dark-adapted: Δρ$_{wild-type}$ = 9.65 ± 1.06°; Δρ$_{hdc}$ = 8.16 ± 0.62°; p = 0.258, two-tailed t-test. Light-adapted: Δρ$_{wild-type}$ = 7.7 ± 0.52°; Δρ$_{hdc}$ = 6.98 ± 0.46°; p = 0.323, two-tailed t-test. (**C**) Predictably, their receptive fields narrowed during light-adaptation (only wild-type shown). The relative changes between the two adaptation states between were statistically similar in wild-type and hdc^{JK910} photoreceptors. Relative changes, calculated as $C = \frac{\Delta\rho_{Dark} - \Delta\rho_{Light}}{\Delta\rho_{Dark}} \times 100\%$. **C**$_{wild-type}$ = 18.44 ± 3.5%; **C**$_{hdc}$ = 13.68 ± 3.37%, p = 0.347, two-tailed t-test. **A-C:** Mean ± SEM; n$_{wild-type}$ = 6; n$_{hdc}$ = 8 cells.

Stimulus history affects photoreceptors' receptive fields

Other than the ambient illumination, which light-adapts the photoreceptors, prolonged light stimulation periods could also desensitize them, resulting in comparable effects. To quantify specific stimulation history effects, we performed recordings from five wild-type (n = 5) and six

hdc^{JK910} photoreceptors (n = 6). Each cell's receptive field was scanned twice in the dark condition, with one minute of Gaussian white-noise (GWN) stimulation in between. The 2nd scan was started within 10 s from ceasing the white-noise light modulation.

We found that the stimulus history affected wild-type and mutant photoreceptors similarly (**Image S4.9**), narrowing their receptive field widths by ~6.5% on average, which was quantitatively ~35-50% of the moderate light-adaptation effects (*cf.* **Box S4.8C**). The differences between the two photoreceptor groups' Δρ values were virtually unchanged by the GWN stimulation and at the margin of statistical significance, having the respective p-values of 0.09 and 0.106 for the 1st and 2nd scans.

Box S4.9. Stimulus-history-dependent changes in wild-type and *hdc*-mutant receptive fields are similar. (**A**) Dark-adapted receptive fields' half-widths of the considered wild-type and *hdc^{JK910}* photoreceptors, measured before one-minute Gaussian white-noise (GWN) stimulation: Δρ_{wild-type} = 9.73 ± 0.45°; Δρ_{hdc} = 8.66 ± 0.32°; p = 0.09, two-tailed t-test. (**B**) Δρ of the same photoreceptors measured in dark condition within 10 s from ceasing the white-noise stimulation: Δρ_{wild-type} = 9.14 ± 0.52°; Δρ_{hdc} = 8.05 ± 0.22°; p = 0.106, two-tailed t-test. (**C**) Relative changes in Δρ of photoreceptors belonging to the two groups: **C**_{wild-type} = 6.2 ± 2.48%; **C**_{hdc} = 6.65 ± 3.01%, p = 0.909, t-test. (**A**-C) Mean ± SEM; two-tailed t-test; n_{wild-type} = 5; n_{hdc} = 6 cells.

R1-R6 acceptance angles are broader than the theoretical prediction

Based on the ommatidium dimensions, as extracted from histological images of fixed/non-living retinae, and the waveguide optic theory, Stavenga (*39*) calculated that the acceptance angles or dark-adapted *Drosophila* R1-R6 photoreceptors should be from 3.8 to 5.0° (as amended for 16.5 µm diameter ommatidium lens). Yet, our current (Δρ = 9.47°; **Boxes S4.7-9)** and earlier (*32*) measurements (Δρ = 8.23°; using ***Method 2*** above) clearly showed that their acceptance angles *in vivo* are in fact about twice as large. What kind of physical mechanism(s) could explain this discrepancy between the theory and measurements?

We briefly introduce here some key points of the new "microsaccadic sampling hypothesis", which is examined in detail in **Texts S7-8** (*e.g.* **Box. S8.4**).

- Living R1-R6 photoreceptors are not still but transiently contract to light (*8*) (**Movie S2**). We show in **Figure 4** (*cf.* **Texts S7-8**) that this causes considerable horizontal rhabdomere movements (up to 1.4 µm, peaking ~60-150 ms after a flash onset and returning back slower). As the lens system stays practically put, the rhabdomere tips shift away from the central axis, skewing the light input and narrowing the photoreceptors' receptive fields dynamically.
- Rhabdomere contractions also move their tips axially; transiently down the focal plane (**Movie S1**).
  - In a dark-adapted state, rhabdomere tips are elongated upwards (closer to the

ommatidium lens), and possibly out of focus, collecting light through the lens system over a wider space. Thus, R1-R6s' acceptance angles would be broader for light flashes spaced by normal (500 ms) intervals, which recover refractory microvilli, returning rhabdomeres to their old positions. But during a bright passing light stimulus, which progressively contracts R1-R6s, their acceptance angles dynamically narrow as the rhabdomeres draw a bit deeper in the retina.

   o At a moderate light-adaptation state (**Box S4.8**), the intracellular pupil mechanism has reduced light input and the rhabdomere lengths should occupy a position (or set-point), which allows further contractions to light increments (*cf.* hair cells in the inner ear (*137*)). However, here, R1-R6s' acceptance angles would still be broader for flashes with 500 ms intervals. Thus, the rhabdomeres would return to their pre-flash positions, which are closer to the lens than during a bright passing light stimulus.

• Photomechanical R1-R6 contractions are partially levered by the adherence junctions (*37*) from their rhabdomeres to the above cone cells and epithelial pigment cells, which form the inner wall of the ommatidium underneath the lens (**Fig. 4F**). Thus, as the photoreceptors contract to light, their adherence junctions appear to pull the pigments cells, moving and narrowing the aperture (and the light beam) in the front of their rhabdomere tips (**Movie S3**).

Slow *Drosophila* retina movements

Despite a *Drosophila*'s head and thorax being immobilized to the conical holder (**Box S4.5A**), its eyes could still move affecting the electrophysiological recordings (*94*). Retinal movements, caused by spontaneous intraocular muscle activity, have been described in larger flies (*19, 138-141*) and treated by different methods, including cooling, anesthesia and fixing their slightly pulled-out antennae (*125*).

Here we report possibly related but slower retinal movements in fixed *Drosophila* preparations *in vivo*. **Box S4.10A** shows an example, in which the optical axis of a R1-R6 photoreceptor moved between two consecutive receptive field scans. In the 1st scan, the receptive field center, which was localized by the largest flash response, corresponded to the light-point No.12. However, the 2nd receptive field scan indicated that the cell's optical axis pointed toward the light-point No.10. This displacement corresponds to an angular movement of ~3^{o}. We found that about 50% of photoreceptors, in which receptive fields were scanned more than once (8/18 wild-type and 8/16 mutant cells), displayed similar retinal movements in the range of 1-3.5^{o}. Moreover, these movements occurred in both front-to-back and back-to-front directions, validating that they were not equipment related artefacts; for example, not caused by gravitational drift in the 25 light-point stimulus array (**Box S4.2**).

Box S4.10. Retinal movements in the *Drosophila* eyes shift photoreceptors' receptive fields, modulating their light input and hence the transduced voltage output. (**A**) A R1-R6 photoreceptor's receptive field shifted between consecutive measurements to 25 light-point stimuli. The 1st scan showed that the receptive field center was closest to light-point No.12; while in the 2nd scan, the peak response was evoked by the light-point No.10. The difference between the optical axes, as indicated by the two scans, was ~3°. (**B**) Examples of R1-R6 photoreceptors' intracellular responses that show slow spontaneous voltage drifts, saccades, jumps or modulation (red arrows and boxes) during repeated light intensity time series stimulation from a fixed point source. Characteristically, these perturbations do not correlate with the light stimuli but are erratically superimposed on the responses' normal adapting trends. Because they occur in the time scale of seconds and show variable rhythmicity, they are likely caused by intrinsic eye muscle activity. Before the recordings, the light stimulus source (3° light-guide-end, as seen by the fly) was carefully positioned at the center of each tested photoreceptor's receptive field.

It has been suggested that recordings from damaged fly photoreceptors may result in (i) extraordinarily wide acceptance angles, (ii) diminishing sensitivity (*127*), or (iii) markedly asymmetrical receptive fields, attributed to artificial electrical coupling between neighboring cells (*126*). To ensure that high-quality Δρ measurements were presented in this study, we only considered data from photoreceptors in which intracellular responses were stable and repeatable, and their receptive fields reasonably symmetrical. However, we acknowledge that *Drosophila* eye movements can inadvertently affect the receptive field assessment accuracy.

The *slow* retinal movements and drifts that shift R1-R6 photoreceptors' receptive fields are likely driven by eye muscle (*20*) activity. These movements can modulate light input to photoreceptors, causing spontaneous dips and peaks in their output during continuous repetitive stimulation, as is sometimes seen during long-lasting intracellular recordings (**Box S4.10B**). Because this additional input modulation seems largely occur in the timescale of seconds, it should reduce mostly the low-frequency signal-to-noise ratio in the R1-R6 output. In this study, to obtain as good estimates as possible of the *Drosophila* photoreceptors' encoding capacity, we only used data from the very best (most stable) recording series. These recordings (**Figs 1-2; figs S1-2**) showed very little or no clear signs of such perturbations.

Furthermore, in **Texts S7-8**, we quantify how all *Drosophila* photoreceptors exhibit *fast* light-triggered photomechanical micro-saccades, peaking ~100 ms after the stimulus onset and lasting 0.2-3 s, depending upon the stimulus intensity. These microsaccades directly modulate light input from the moving objects and consequently photoreceptor output, and we show through simulations and recordings how they can improve the spatiotemporal resolution of neural images (**Figs 4** and **5**; **figs S8** and **S9**).

Conclusions

In this supplement, we described how receptive fields (spatial responsiveness) and acceptance angles ($\Delta\rho$) of wild-type and synaptically-blind hdc^{JK910} R1-R6 photoreceptors were estimated, using intracellular voltage responses to novel light-point array stimulation. Their characteristics in the dark- and moderately light-adapted states, and after prolonged light stimulation, were compared to test the hypothesis that spatial information summation in the lamina contributes to *Drosophila* photoreceptor function. We found that the dark-adapted wild-type R1-R6s show wider receptive fields. But in the other tested lighting conditions, the two photoreceptor groups' $\Delta\rho$-estimates adapted similarly.

Could the difference between the dark-adapted wild-type and hdc^{JK910} receptive field widths result from recording artefacts? It is well known that LMCs' hyperpolarizing intracellular responses to light pulses depolarize the surrounding extracellular space (*142*), seen as the on-transient in the electroretinogram (ERG) recordings (*143*). Theoretically, these signals could be picked up by recording microelectrodes, adding artificial components to the intracellularly measured V_n, and so making wild-type photoreceptors' receptive fields appear wider (*144*). However, we can essentially rule out this notion because the Gaussian functions, which quantified the actual recordings, were fitted to their real flank amplitudes rather than to zero (**Box S4.6**). Consequently, any extra DC component would not affect the resulting $\Delta\rho$-estimates. Furthermore, our past comparative study, which included different synaptic mutant flies, failed to find clear signs for ERG contamination in high-quality intracellular wild-type photoreceptor recordings (*30*). Whereas by lacking neurotransmitter histamine, hdc^{JK910} LMCs cannot respond to light (*34, 48, 85*), and thus hdc^{JK910} photoreceptor recordings cannot be ERG contaminated. Hence, we conclude that ERG signals could only have marginal contribution to our results at most.

In fact, the dark-adapted photoreceptors' $\Delta\rho$-estimates constituted the most reliable data in this supplement, with the largest number of samples ($n_{\text{wild-type}} = 19$, $n_{\text{hdc}} = 18$ cells) obtained through consistent recording protocols. In every experiment, the receptive field assessment in the dark-adapted state was strictly the first examination, following the standard stimulus centering procedure. This minimized any potential downgrade in the recording quality or variation in the stimulation history. Notably, our estimate of WT *Drosophila* R1-R6 photoreceptors' average receptive field half-width in the dark-adapted state, $\Delta\rho = 9.47 \pm 0.36°$ (n = 19 cells; see **Method 3**, above), is reasonably similar to the previous estimate of $8.23 \pm 0.54°$ (n = 11 cells) (*32*), which was obtained through a less stationary recording apparatus/method with more assumptions (see **Method 2**, above).

The dominant factors determining a fly photoreceptor's receptive field are optical, waveguide properties and, particularly for the chosen measurement method, the phototransduction characteristics (*2, 39, 88, 91*). For blowflies, it has been shown that the receptive field shape can be largely derived from the optical structure dimensions with the waveguide theory (*125*). Given the hdc^{JK910} mutants' seemingly normal ommatidial and rhabdomere optics, as seen *in vivo* (**Text S7**) and under electron microscopy (**Text S5**; R1-R6 rhabdomere diameters, *d*, were ~96% of the wild-type), and their wild-type like photoreceptor voltage/intensity relations to brief light pulses (*35*), it is reasonable to expect that their $\Delta\rho$-estimates would be close to wild-type. This should especially be true in light-adaptation, which is predicted to make photoreceptor output more independent of its neighbors (*46, 114, 145*). And indeed, we found $\Delta\rho$-estimates of the light-adapted wild-type and hdc^{JK910} photoreceptors alike.

But because of this conformity, the 10.9% difference between their dark-adapted acceptance angles requires an additional explanation. Although at the flanks this difference increases (**Box S4.7A**), it still may seem rather small when compared to the measured cell-to-cell variation within each genotype, with the respective maxima being 66% and 88% wider than the minima, and its statistical significance becomes less with fewer samples (**Box S4.8A**). Nevertheless, the finding is conceptually important as it supports an expansion in the classic spatial vision paradigm; from the optical constraints to spatial information summation in the network (*146*).

The dark-adapted wild-type photoreceptors' wider receptive fields are consistent with what we know about how synaptic inputs are channeled from lamina interneurons to R1-R6 axons (*28, 30, 68, 71, 124, 147, 148*). Thus, the corresponding narrower receptive fields of mutant photoreceptors seem most sensibly attributed to the missing excitatory feedback modulation from their interneurons (*30, 35, 71, 123*). As shown by intracellular recordings (*30, 35*), feedforward and feedback signals dynamically contribute to photoreceptor and interneuron outputs *in vivo*. When the probability of light saturation is low, the stronger synaptic transmission in both pathways helps to amplify their response amplitudes.

Therefore, taken together with the findings of Dubs et al. (*119*) and Dubs (*122*), these results suggest that under dim illumination, lateral excitation spreads synaptically within the lamina of the fly visual system. Spatial information summation is likely implemented by the first-order interneurons and fed back to photoreceptors through connections (*28, 68*) that utilize excitatory neurotransmitters (*30, 124*). This model is further supported by our anatomical observations (**Text S5** and **Text S7**), which imply that there are no major developmental defects in hdc^{JK910} retina and that their lens systems and rhabdomere sizes are broadly wild-type-like.

Text S5: R1-R6 rhabdomere sizes, neural superposition and hyperacuity schemes

Overview

This supplement shows how R1-R6 rhabdomere diameters vary systematically and consistently in *Drosophila* ommatidia and provides supporting background information for the results presented in **Figures 3-5** in the main paper and **Text S4**.

In this supplement:

- We measure wild-type and hdc^{JK910} (histamine-mutant (*34, 48, 85*)) R1-R6 photoreceptors' rhabdomere sizes from the electron micrographs of their retinae.
- We compare these measurements to their electrophysiologically measured receptive field estimates (**Text S4**).
- We show that the mean wild-type R1-R6 rhabdomere diameter, d_{R1-R6}, is only ~4.1% wider than in hdc^{JK910} mutant eyes. This difference may contribute in part to their 10.9% wider average acceptance angle ($\Delta\rho$) in a dark-adapted state (**Text S4**), but cannot fully explain it (*cf.* **Text S7**).
- We further show that in both phenotypes R1, R3 and R6 rhabdomeres are systematically larger than R2 and R4 rhabdomeres. As for the maximum difference, the mean R1 rhabdomere diameter ($d_{R1} = 1.8433 \pm 0.1294$ μm, mean ± SD) is about 18% wider than that of R4 ($d_{R4} = 1.5691 \pm 0.0915$ μm, n = 25, p = 2.3419 x 10^{-11}, 2-tailed t-test).
- These findings imply that each R1-R6 photoreceptor should have a different receptive field size (coinciding with their considerable $\Delta\rho$ variation seen in **Text S4**, **Box S4.7C**).
- Our results further suggest an asymmetric information integration model, in which the neural superposition of different-sized overlapping receptive fields (of the neighboring R1-R6s) has a potential to contribute in enhancing *Drosophila*'s visual acuity beyond the presumed optical limits of its compound eyes (as revealed by behavioral experiments in **Text S9**).

Readjusting the current theoretical viewpoint

Neural superposition eyes provide more samples from local light intensity changes for each image pixel, represented by large monopolar cell (L1 and L2) outputs. In the conventional viewpoint, each pixel's signal-to-noise improves by $\sqrt{6}$ because its L1 and L2 receive similar inputs from six "physiologically identical" R1-R6 photoreceptors, which sample information from the same small area in the visual space. Thus, the conventional assumptions and limits for neural superposition performance are:

- Each R1-R6 functions virtually identically, generating similar outputs to the same light input
- Each R1-R6 in superposition has an identical receptive field size and shape
- The receptive fields in superposition overlap perfectly
- Each image pixel represents sampling and processing within its inter-ommatidial angle
- Inter-ommatidial angle sets the visual resolution of the *Drosophila* eye

This supplement presents anatomical and theoretical evidence that these assumptions and limits are overly simplistic and suggests ways the real neural images could be sharpened beyond them to improve *Drosophila* vision.

Electron micrographs

Fixation. Flies were cold anaesthetized on ice and transferred to a drop of pre-fixative (modified Karnovsky's fixative (*69*): 2.5% glutaraldehyde, 2.5% paraformaldehyde in 0.1 M sodium cacodylate buffered to pH 7.3) on a transparent agar dissection dish. Dissection was performed using a shard of a razor blade (Feather S). Flies were restrained on their backs with insect pins through their lower abdomen and distal proboscis. Their heads were severed, probosces excised, and halved. The left half-heads were collected in fresh pre-fixative and kept for 2 h at room temperature under normal lighting conditions.

After pre-fixation, the half-heads were washed (2 x 15 min) in 0.1 M Cacodylate buffer, and then transferred to a 1 h post-fixative step, comprising Veronal Acetate buffer and 2% Osmium Tetroxide in the fridge (4 $^{\circ}$C). They were moved back to room temperature for a 9 min wash (1:1 Veronal Acetate and double-distilled H_2O mixture), and serially dehydrated in multi-well plates with subsequent 9 min washes in 50%, 70%, 80%, 90%, 95% and 2 x 100% ethanol.

Post-dehydration, the half-heads were transferred to small glass vials for infiltration. They were covered in Propylene Oxide (PPO) for 2 x 9 min, transferred into a 1:1 PPO:Epoxy resin mixture (Poly/Bed® 812) and left overnight. The following morning, the half-heads were placed in freshly made pure resin for 4 h, and placed in fresh resin for a further 72 h at 60 $^{\circ}$C in the oven. Fixation protocol was kindly provided by Professor Ian Meinertzhagen at Dalhousie University, Halifax, Nova Scotia.

Sectioning and staining. Embedded half-heads were first sectioned (at 0.5 µm thickness) using a glass knife, mounted in an ultramicrotome (Reichert-Jung Ultracut E, Germany). Samples were collected on glass slides, stained using Toluidine Blue and observed under a light microscope. This process was repeated and the cutting angle was continuously optimized until the correct orientation and sample depth was achieved; stopping when approximately 40 ommatidia were discernible. The block was then trimmed and shaped for ultra-thin sectioning. The trimming was necessary to reduce cutting pressure on the sample-block and resulting sections, thus helping to prevent "chattering" and compression artefacts. Ultra-thin sections (85 nm thickness) were cut using a diamond cutting knife (DiATOME Ultra 45°, USA), mounted and controlled using the ultramicrotome. The knife edge was first cleaned using a polystyrol rod to ensure integrity of the sample-blocks. The cutting angles were aligned and the automatic approach- and return-speeds set on the microtome. Sectioning was automatic and samples were collected in the knife water boat.

Sections were transferred to Formvar-coated mesh-grids and stained for imaging: 25 min in Uranyl Acetate; a double-distilled H_2O wash; 5 min in Reynolds' Lead Citrate (*149*); and a final double-distilled H_2O wash.

Box S5.1. R1-R6 photoreceptors' rhabdomere sizes differ consistently. (**A**) Electron micrographs of a characteristic wild-type (left) and *hdc^JK910^* (right) ommatidia. Each ommatidium contains the outer receptors, R1-R6, and the inner receptors, R7/R8, which can be identified by their rhabdomeres' relative positions. Here R8s are not visible because these lie directly below R7s. Markedly, both wild-type and *hdc^JK910^* R1-R6 photoreceptor rhabdomere sizes vary systematically. (**B**) The mean rhabdomere sizes measured from 25 ommatidia from 10 flies. *hdc^JK910^* R1-R6 rhabdomere cross-sectional areas are smaller than those of the wild-type cells, but show similar proportional variations. Error bars show SEMs.

Rhabdomere measurements. Transmission EM images for R1-R6 rhabdomere size comparisons were taken below the rhabdomere tips, as sectioned 25 μm down from the corneal surface of the ommatidium lens. 25 wild-type and *hdc^JK910^* ommatidia (n = 25) from 10 flies of each phenotype were used to estimate R1-R6 rhabdomere sizes. The images were processed with ImageJ software. Because the rhabdomere cross-sectional area is often better approximated by an ellipse than a circle (**Box S5.1A**), for greater accuracy, its circumference was fitted with an ellipse.

For an EM rhabdomere area, *A*, its mean diameter was then taken: $d = 2 \times \sqrt{\frac{A}{\pi}}$. Note that the obtained mean rhabdomere diameter estimates are somewhat smaller than the previous estimates, which measured the minimum and maximum rhabdomere diameters from edge-to-edge (*32, 150*). Here, instead, we standardized the measurement protocol to reduce *d*-estimation bias between the wild-type and *hdc^JK910^* electron micrographs to obtain straightforward statistical comparisons of their means. Nevertheless, both these and the previous(*150*) *d*-estimates indicated systematic R1-R6 rhabdomere size differences.

R1-R6 rhabdomere sizes differ
Drosophila R1-R6 photoreceptors' rhabdomere sizes vary systematically and consistently in each ommatidium (**Box S5.1B**). R1 and R6 rhabdomere cross-sectional areas are always the largest and R4 rhabdomeres the smallest (**Box S5.2**).

Significance t-test 2-tail	wild-type R1	wild-type R2	wild-type R3	wild-type R4	wild-type R5	wild-type R6
wild-type R1	N/A	5.3626×10^{-7}	1.3033×10^{-4}	3.6882×10^{-10}	1.5564×10^{-5}	0.8282
wild-type R2	5.3626×10^{-7}	N/A	0.1531	0.0371	0.3760	3.4906×10^{-7}
wild-type R3	1.3033×10^{-4}	0.1531	N/A	0.0011	0.5736	1.3187×10^{-4}
wild-type R4	3.6882×10^{-10}	0.0371	0.0011	N/A	0.0047	1.2193×10^{-10}
wild-type R5	1.5564×10^{-5}	0.3760	0.5736	0.0047	N/A	1.3479×10^{-5}
wild-type R6	0.8282	3.4906×10^{-7}	1.3187×10^{-4}	1.2193×10^{-10}	1.3479×10^{-5}	N/A

Box S5.2. Statistical comparison of wild-type Canton-S R1-R6 rhabdomere cross-sectional areas. The table gives the differences as p-values, calculated for 25 ommatidia of 10 flies at the same retinal depth. Red indicates statistically significant difference. R4 rhabdomeres are smaller than the other rhabdomeres, whereas R1 and R6 are always the largest.

Theoretically, a fly photoreceptor's receptive field size depends upon its rhabdomere diameter, d (Eq. S4.3). This relationship was recently supported experimentally by comparing the rhabdomere diameters and acceptance angle estimates, $\Delta\rho$, of *Drosophila* R1-R6 photoreceptors to those of killer fly (*Coenosia attenuata*), both of which eyes have rather similar ommatidial lens sizes (16–17 vs 14–20 µm) and focal lengths (21.36 vs 24.70 µm)(*32*), *cf.* Eq. S4.3. In ♀ *Drosophila*, maximum d was ~2 µm and in ♀ *Coenosia* ~1 µm, while *Drosophila*'s mean $\Delta\rho$-estimate was 8.23 ± 0.54^{o} and *Coenosia*'s 2.88 ± 0.07^{o} (*32*). Thus, the wider rhabdomere tip correlates strongly with the wider receptive field.

Accordingly, with each *Drosophila* ommatidium hosting R1-R6 rhabdomeres of distinct size differences (**Box S5.1B**, left; **Box S5.2**), the receptive fields of the neighboring R1-R6, which are pooled together in neural superposition, should differ in size and overlap broadly. These observations and analysis concur with the broad variability of the intracellularly measured wild-type R1-R6s' receptive field widths (**Text S4**: **Box S4.7A** and **C**).

However, although collectively the mean wild-type R1-R6 rhabdomere cross-sectional areas (**Boxes S5.3** and **S5.4**) are larger than their hdc^{JK910} counterparts (**Box S5.1B**), their corresponding diameter differences are small. The average wild-type R1-R6 rhabdomere diameter, (WT $d_{R1-R6} = 1.70 \pm 0.15$ µm; mean $\pm$ SD, n = 150 rhabdomeres; **fig. S7**) is only ~4.1% wider than in hdc^{JK910} mutant eyes (hdc^{JK910} $d_{R1-R6} = 1.64 \pm 0.10$ µm; n = 150 rhabdomeres), as extrapolated from the ellipsoid rhabdomere area fits. Thus, hdc^{JK910} photoreceptors' smaller average rhabdomere diameters can contribute to their ~10.9% narrower $\Delta\rho$ (**Text S4**: **Box S4.7A** and **D**), but cannot fully explain it (**Box S5.1B**, right).

Significance t-test 2-tail	hdc^{JK910} R1	hdc^{JK910} R2	hdc^{JK910} R3	hdc^{JK910} R4	hdc^{JK910} R5	hdc^{JK910} R6
hdc^{JK910} R1	N/A	4.7387×10^{-10}	0.2281	6.3122×10^{-10}	1.0125×10^{-10}	0.2335
hdc^{JK910} R2	4.7387×10^{-10}	N/A	7.9300×10^{-11}	0.7006	0.9897	2.0471×10^{-10}
hdc^{JK910} R3	0.2281	7.9300×10^{-11}	N/A	1.9466×10^{-10}	5.0272×10^{-12}	0.9921
hdc^{JK910} R4	6.3122×10^{-10}	0.7006	1.9466×10^{-10}	N/A	0.6830	4.3275×10^{-10}
hdc^{JK910} R5	1.0125×10^{-10}	0.9897	5.0272×10^{-12}	0.6830	N/A	1.6775×10^{-11}
hdc^{JK910} R6	0.2335	2.0471×10^{-10}	0.9921	4.3275×10^{-10}	1.6775×10^{-11}	N/A

Box S5.3. Statistical comparison of hdc^{JK910} R1-R6 rhabdomere cross-sectional areas. The table gives the differences as p-values, calculated for 25 ommatidia of 10 flies at the same retinal depth. Red indicates statistically significant difference. R2, R4 and R5 rhabdomeres are smaller than R1, R3 and R6 rhabdomeres.

Significance t-test 2-tail	wild-type R1	wild-type R2	wild-type R3	wild-type R4	wild-type R5	wild-type R6
hdc^{JK910} R1	0.0014	0.0012	0.1434	1.0091×10^{-7}	0.0302	0.0015
hdc^{JK910} R2	1.8211×10^{-11}	0.0078	1.5520×10^{-4}	0.6952	7.1690×10^{-4}	3.5405×10^{-12}
hdc^{JK910} R3	5.1489×10^{-5}	0.0073	0.4818	1.8478×10^{-7}	0.1415	3.8432×10^{-5}
hdc^{JK910} R4	1.7483×10^{-11}	0.0051	1.0904×10^{-4}	0.5015	4.9066×10^{-4}	3.7421×10^{-12}
hdc^{JK910} R5	9.2505×10^{-12}	0.0056	1.0272×10^{-4}	0.6710	4.8280×10^{-4}	1.5313×10^{-12}
hdc^{JK910} R6	5.6987×10^{-5}	0.0083	0.4903	2.8960×10^{-7}	0.1480	4.3539×10^{-5}

Box S5.4. Statistical comparison of wild-type and hdc^{JK910} R1-R6 rhabdomere cross-sectional areas. The table gives the differences as p-values, calculated for 25 ommatidia of 10 flies at the same retinal depth. Red indicates statistically significant difference. The wild-type and mutant R3 and R4 rhabdomeres are the same size (highlighted in bold); the other wild-type rhabdomeres are larger than their respective counterparts.

Theoretical models for spatial hyperacuity

In the *Drosophila* compound eye, R1-R6 photoreceptors' mean acceptance angle ($\Delta\rho \sim 9.5^{o}$) is on average about twice as wide (see **Text S4**) as its mean inter-ommatidial angle ($\Delta\varphi \sim 4.5^{o}$) (*32*). Such an eye design could potentially facilitate spatial hyperacuity through neural image processing. In the vein of previous suggestions for manmade optoelectrical systems (*151*) and retinae of other species (*152*), we consider three alternative hypothetical scenarios based upon neighboring R1-R6s' overlapping receptive fields (RFs).

The first case (**Box S5.5**, left) considers RF variations of the photoreceptors in neural superposition. An image of an object (here, a thin 1^{o} vertical bar, grey) stimulates simultaneously eight photoreceptors (R1-R8), which gather light information about the same area in the visual space by their overlapping RFs (for clarity only R1-R3' RFs are shown; orange Gaussians). However, because the rhabdomere diameters, *d*, vary considerably and consistently between them (*e.g.* R1 and R6 rhabdomere diameters areas are ~18% wider than those of R4s; **Box S5.1**),

so could also be their RF sizes (see **Text S4**, Eq. S4.3), causing overlaps. Six of these inputs (R1-R6) are pooled in the lamina. Consequently, even a small (1^o) displacement of the vertical bar stimulus (grey) could lead to variable but specific light intensity modulations in each of the six converging input channels. Their signals to LMCs (L1-L3), which in high signal-to-noise conditions transforms input modulations into phasic responses (*29, 30, 71, 153, 154*), could therefore be different before and after the bar displacement and possibly neurally detectable.

In the second case (**Box S5.5**, middle), six photoreceptor cells of the same type (say R2s) in the neighboring ommatidia gather light information from the neighboring small visual areas (~4.5^o apart, as separated by the interommatidial angle), but their receptive fields (~9.5^o half-widths) overlap; for clarity only three R2s are shown: orange, blue and red. Lateral connections (L4, Lawf and Am cells) between lamina neuro-ommmadia could then be used by LMCs to compare their outputs, enhancing the spatial resolution of each neural channel (see **Text S2**, **Box S2.5**).

Box S5.2. Theoretical ways to neurally improve the optical image resolution in the fly compound eyes. **Left**, neural superposition with overlapping photoreceptor receptive fields. **Middle**, same type photoreceptors (here R2s) in the neighboring ommatidia ($\Delta\phi$ ~4.5^o) collect light from neighboring visual areas, but their receptive fields (RFs) overlap, having twice as large acceptance angles, $\Delta\rho$ ~9.5^o. **Right**, neighboring photoreceptors in the same ommatidium collect light from neighboring visual areas with their RFs overlapping. By comparing the resulting variable photoreceptor outputs from a small visual object (0.5 ^{o} vertical grey bar), neural circuitry may resolve objects finer than the inter-ommatidial angle ($\Delta\phi$).

In the third case (**Box S5.5**, right), R1-R6 photoreceptor cells in the same ommatidium gather light information from the neighboring small visual areas, but their receptive fields overlap; again for clarity only three: orange (R1), blue (R2) and red (R3) are shown. Lateral connections (L4, Lawf and Am cells) between adjacent lamina neuro-ommatidia may enable LMCs to compare their outputs, enhancing the spatial resolution of each neural channel.

All these circuit models could theoretically contribute to motion vision hyperacuity, which occurs when a stationary *Drosophila* eye resolves object motion finer than its ~4.5^o inter-ommatidial angle (the average sampling point or photoreceptor spacing). And crucially, in **Text S10**, we use a flight simulator system to demonstrate and quantify *Drosophila* hyperacute optomotor behavior to sub-interommatidial stimuli ($1-4^o$). However in **Text S7-8**, we further provide decisive evidence that the spatiotemporal resolution of the early neural images is improved by synchronized and coherent photomechanical rhabdomere contractions (*8*), which move and *narrow* R1-R6s' receptive fields. Together with refractory information sampling, this

enables photoreceptors to encode space in time. Therefore, as the RFs *narrow* with moving stimuli, this reduces the overlap between the neighboring RFs and, consequently, affects the potential resolving power of the circuit models in **Box S5.5**.

We conclude that *at the photoreceptor level* the overlapping RFs provide neither necessary nor sufficient mechanistic explanations for the *Drosophila* spatiotemporal hyperacuity. However, *at the lamina interneuron level*, such connectivity arrangements may further enhance hyperacute vision. It is also possible that any kind of retinal image enhancement would be further coordinated centrally to match the visual needs of a locomoting fly. In walking blowflies, intraocular muscles in both the left and right eye seem to contract synchronously with increasing rates, causing vergence eye movements (*19, 138-141*). Correspondingly, in lobula plate motion sensitive cells, responses to moving visual stimuli increase with locomotion (*155-158*).

Possible dynamics arising from connectivity and variable rhabdomere sizes
Temporal output modulation through gap-junctions between R1-R6 and R7-R8 photoreceptor axons could further contribute, as a possible network mechanism, to the acuity improvements that we report in this publication (**Figs. 3** and **4**; see **Text S6**, **Box S6.7D**). *Drosophila* R1-R6s have larger rhabdomeres, each with ~30,000 microvilli, whereas those of R7s and R8s contain only ~15,000. Owing to superposition in each neuro-ommatidium, both of these photoreceptor classes integrate photons from the same small area in space, but for given light changes, the macroscopic R1-R6 output rises and decays faster than those of R7s and R8s. This is because R1-R6 rhabdomeres integrate twice as many samples from the same stimulus and their membranes have likely smaller time constant (*159*). Therefore, at each moment in time, R1-R6 and R7-R8 outputs carry a dynamic phase difference. If their responses were to antagonize each other through the gap-junctions between R6 and R7/R8 axons (*29, 69, 70*), similar to the crosstalk between *Calliphora* R7 and R8 outputs (*160*), phasic R1-R6 output components could be enhanced even further.

Text S6: Neural images of moving objects (R1-R6 recordings vs. classic predictions)

Overview
This supplement describes a new method to measure *Drosophila* photoreceptor output to moving dots, and shows how these responses provide much higher visual resolution and motion blur resistance than what is predicted by the classic theories, supporting the results in **Figures 3-5** in the main paper.

In this supplement:
- We measure how well dark- and moderately light-adapted wild-type *Drosophila* R1-R6 photoreceptors resolve bright dots (point-objects), which cross their receptive fields at different speeds.
- We compare these intracellular recordings to those of histamine-mutants (*34, 48, 85*) (*hdc^{JK910}*), in which first-order interneurones are blind (receiving no neurotransmitter from photoreceptors) and thus are incapable of feedback-modulating the photoreceptor output.
- We further record voltage responses of blowfly (*Calliphora vicina*) R1-R6 photoreceptors to moving point-objects, as an additional test of our experimental setup, stimulus paradigm and mathematical analyses, validating this method.
- We evaluate the wild-type and mutant recordings against their respective classic model simulations, in which each recorded receptive field is convolved by the same cell's impulse response.
- Our results indicate that both wild-type and *hdc^{JK910}* R1-R6s resolve moving dots approximately equally well, and significantly better than the corresponding classic model simulations.
- These findings demonstrate that the classic deterministic photoreceptor models (*2-4*) for resolving moving objects grossly underestimate the visual resolving power of real photoreceptors.
- Consequently, the classic theory overestimates the effects of motion blur on *Drosophila* vision during saccadic behaviors.

Retinal limitations and capacity to resolve moving objects
A single photoreceptor's voltage responses can give important insight into neural image processing behind a fly's ability to detect small objects, whether in high-speed chasing flights or against a cluttered background (*161, 162*). Theoretically, when a point-object moves uniformly across an array of photoreceptors, each cell would produce a similar response. But these responses would be displaced in time, t, which it takes for the object to travel between two adjacent cells' receptive field centers (*4*). Therefore, the response of the whole array, as a collective neural representation of the moving point-object, would be a travelling pattern with a mirrored waveform. Mathematically, this can be extrapolated from a single photoreceptor's response. We now apply this classic approach to simplify the questions about neural images of moving point-objects and compare its predictions to R1-R6s' spatiotemporal responses.

First, we consider the problem of neural latency compensation in motion perception. This is related to the flash-lag effects observed in humans (*163, 164*). Due to the inevitable phototransduction delays, every sighted animal should encounter this problem and *Drosophila* is not an exception. At 19 ºC, its photoreceptors' voltage responses rise ~10 ms from the light flash

and peak 15-30 ms later (*45*). Given that a fly's saccadic turning speed (*22*) in flight can exceed 1,000 $^\circ$/s, such delays need compensating by network computations. Otherwise, neural images of its surroundings could lag behind their actual positions by more than 25°, making fast and accurate visual behaviors seemingly infeasible.

Such compensations have been shown to occur early on in vertebrate eyes. In the tiger salamander and rabbit retina, ganglion cells' firing rates lag behind flashing but not moving bars (*165*). Whether similar processing happens in insect eyes is unknown. By analyzing *Calliphora*, wild-type *Drosophila* and *hdc*^{JK910} R1-R6 output to a point-object crossing their receptive fields, we find their time-to-peak values broadly similar to those evoked by light flashes. Therefore, these lag times are not, or at most weakly, compensated by the signal spread between photoreceptors or synaptic feedback.

Second, we examine whether R1-R6 output to point-object motion displays directional preference, as suggested by the asymmetric synaptic feedback to them (*28, 68*). While both L1 and L2 monopolar cells mediate a major neural pathway in the lamina, L2s show richer connectivity. Only L2 projects feedback to R1-R6 and have reciprocal connections with L4, which in turn connects to L4s of the neighboring neural cartridges (*166*), providing further feedback to photoreceptors (*28, 68*). Although data is limited about this circuitry's specific functional roles (*36*), its connectivity suggests that the L2 pathway has stronger (**i**) lateral interactions and (**ii**) influence on R1-R6 output than the L1 pathway. Interestingly, neurogenetic behavioral tests (*167*) indicated that at low pattern contrast conditions, the L1 and L2 pathways are specialized to mediate back-to-front and front-to-back motion perception, respectively. Because flies locomote forwards more often than backwards, front-to-back is the dominant direction of visual stimuli. Thus, processing of directional object motion might involve L2 and L4 feedback to R1-R6.

We find that whilst the photoreceptors' peak responses show no clear directional preference, their rise and delay time courses to front-to-back and back-to-front moving point-objects often differ significantly. Although these differences may in part be augmented by the asymmetric network feedback, we show in **Texts S7-8** that they actually originate from photomechanical R1-R8 contractions (*8*).

Third, we investigate whether network regulation affects R1-R6 cell's spatiotemporal acuity. By using the classic approaches (*2-4*), we estimate the theoretical blur effects and the eye's ability to resolve bright dots moving at certain speeds. Furthermore, we apply the Volterra series, a widely used "black-box" modeling method (*57, 168-171*), to simulate wild-type and *hdc*^{JK910} R1-R6 output to these point-objects (*172, 173*). Remarkably, the simulations make it clear that these models cannot predict the recordings accurately (see **Text S1** and the main article, **Figs 3** and **4G-I**).

Spatiotemporal resolution of the eye is thought to be determined by two components with special characteristics: the static spatial resolution of light input, as channeled through the optics (*4, 174*), and the temporal response dynamics of photoreceptors. These characteristics are further influenced by the photoreceptors' adaptation state and synaptic feedback. The classic approaches suggest that because wild-type R1-R6s' acceptance angles ($\Delta\rho$) are 10.9% wider than those of *hdc*^{JK910} but their response dynamics are similar (see **Text S4**), they should produce blurrier images (of wider spatial half-width, *S*). Moreover, as *hdc*^{JK910} R1-R6s lack synaptic feedback modulation (*35*), their predicted higher acuity should reflect differences in spatiotemporal photon sampling dynamics. Our recordings show, however, that both wild-type and *hdc*^{JK910} R1-R6s resolve moving dots at least twice as well what the classic theory predicts, and that any

resolvability difference between these cells largely disappears against a lit background. Thus, in dim conditions, lateral summation within the network may sensitize R1-R6 output by trading-off acuity, whereas in bright conditions more independent photoreceptor output sharpens neural images. Nonetheless, the classic theory cannot account for these dynamics, as it greatly overestimates the effect of motion blur on photoreceptor output.

We later demonstrate in **Texts S7-8** how and why the model simulations (of this supplement) differ from the corresponding recordings. Essentially, this is because the classic theoretical approaches do not incorporate two interlinked biophysical mechanisms that are critical for high acuity. (**i**) Rapid photomechanical photoreceptor contractions(*8*) accentuate light input dynamically by shifting (front-to-back) and narrowing the cell's receptive field as moving bright point-objects enter in its view. While (**ii**) stochastic refractory photon sampling by microvilli accentuates the temporal dynamics in R1-R6 output. These mechanisms work together to improve the acuity and resolvability of moving objects far beyond the predictions of the classic models.

Moving visual stimuli

The 25 light-point array and LEDs pads, which we used for creating images of moving objects and providing ambient illumination, respectively, are described in **Text S4**. In the *Drosophila* experiments, the 25 light-point array was placed 6.7 cm away from the fly, subtending an angle of 40.92^{o}. This gave each light-point (dot) 1.7^{o} size and minimum inter-dot-distance. In the *Calliphora* experiments, these parameters were 17 cm (distance) and 16.73^{o} (viewing angle).

Images of one moving point-object were produced by briefly turning each light-point on and off, one after another in an incremental (for front-to-back direction) or decremental order (for back-to-front direction). Accordingly, Channel 0 input was driven with increasing or decreasing "ramp" (**Box S6.3A**), while Channel 1 input was set to 2 V. The travelling time of an object, or duration of the "ramp", was between 50 ms and 2 s, resulting in object speeds within naturalistic range (*22, 175, 176*): from 20 to 818 ^{o}/s for *Drosophila* and 8 to 334 ^{o}/s for *Calliphora*.

The ability to resolve two moving point-objects of *Drosophila* photoreceptors were tested in dark-and light-adaptation conditions. In light-adaptation experiments, two 39-LED pads, on both sides of the 25 light-point array, provided background illumination (see **Text S4**, **Box S4.2**). The two dots in the 25 light-point array were separated by 6.8^{o} (four dark points in between) and moved together at different speeds; typically, 205, 409 or 818 ^{o}/s. Each stimulus was presented 8-10 times to the fly and the resulting photoreceptor responses were averaged before being analyzed.

Gaussian white-noise (GWN) stimuli

To evaluate how well the classic theory of fly compound eye optics/function (*3, 4, 73*) explains single *Drosophila* R1-R6 photoreceptors ability to resolve moving point-objects (dots), we needed to estimate each photoreceptor's linear impulse response (the 1st Volterra kernel) separately. The cell's voltage response to moving dot could then be predicted by convolving each recorded receptive field by the same cell's impulse response

Light-point No.13 intensity was controlled by setting Channel 0 input to 5 V and modulating Channel 1 input with a Gaussian white-noise (GWN) time series, which had the mean value of 2.5 V and cut-off frequency of 200 Hz. With these settings, light-point No.13 delivered 2.5×10^{6} photons/s at **Peak1** (451 nm) and 3.75×10^{6} photons/s at **Peak2** (575 nm) on

average (*cf.* **Text S4**: **Box S4.4**). Finally, these intensities were reduced 100-fold by neutral density filtering

Volterra series model of each tested *Drosophila* R1-R6
The principal assumptions of the Volterra series method are that the system has finite memory and is time-invariant (*177*). That is, (**i**) the relationship between output (photoreceptor voltage response) $y(t)$ and input (light stimuli) $u(t)$ is characterized by an unchanging impulse response and (**ii**) $y(t)$ depends only on current and past values of $u(t - \tau) \rightarrow u(t)$ with limited regression time, τ. The continuous form of this input/output relationship is described by the following equation:

$$y(t) = k_0 + \int_0^T k_1(\tau)u(t - \tau)d\tau + \int_0^T \int_0^T k_2 u(t - \tau_1)u(t - \tau_2)d\tau_1 d\tau_2 \qquad \text{(S6.1)}$$

where k_0, k_1 and k_2 are the zero-, first- and second-order time-invariant kernels, which define the system's impulse response. T is the finite system memory limit.

Note that the model order is not limited, as expressed only up to 2^{nd}-order in Eq. S6.1, but instead could be extended arbitrarily further. However, it has been shown that a light-adapted fly photoreceptor's response to GWN light intensity time series stimulation, which was used in these experiments, can be approximated well by the linear terms (*64, 178*).

Therefore, the estimation of system output was simplified to a linear convolution of input with zero- and first-order kernels. Each measurement of photoreceptor voltage response and light stimuli could be fitted into the discrete and simplified form of Eq. S6.1 as:

$$y(n) = k_0 + k_1(0)u(n) + k_1(1)u(n-1) + \ldots + k_1(T)u(n-T)$$
$$y(n-1) = k_0 + k_1(0)u(n-1) + k_1(1)u(n-2) + \ldots + k_1(T)u(n-1-T)$$
$$\vdots \qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad \text{(S6.2)}$$
$$y(n-N) = k_0 + k_1(0)u(n-N) + k_1(1)u(n-N-1) + \ldots + k_1(T)u(n-n-T)$$

The group of Eq. S6.2, which approximates N values of photoreceptor output, was then re-arranged into matrix form:

$$\begin{pmatrix} y(n) \\ y(n-1) \\ \vdots \\ y(n-N) \end{pmatrix} = \begin{pmatrix} 1 & u(n) & u(n-1) & \cdots & u(n-T) \\ 1 & u(n-1) & u(n-2) & \cdots & u(n-1-T) \\ & & \vdots & & \\ 1 & u(n-N) & u(n-N-1) & \cdots & u(n-N-T) \end{pmatrix} \times \begin{pmatrix} k_0 \\ k_1(0) \\ k_1(1) \\ \vdots \\ k_1(T) \end{pmatrix} \qquad \text{(S6.3)}$$

Equivalently, Eq.S6.3 could be symbolized as:

$$Y = P\theta \qquad \text{(S6.4)}$$

where vector Y contained a sequence of N output values, P was the regression matrix constructed from the lagged input values, and the column θ elements were the kernel values. The problem of determining a photoreceptor's Volterra series model was hence broken down to designing input stimuli $u(t)$ and measuring output values $y(t)$ to construct matrices P and Y of Eq. S6.4, and estimating θ.

$u(t)$ was a GWN series with 200 Hz bandwidth, which thus activated the whole frequency range of photoreceptor output. Initially, each tested photoreceptor was steady-state-adapted to

the chosen light background (the average brightness of the GWN stimuli) for 30-60 s (*cf.* **Text S4**, page 21). Input was then delivered from the light-point No.13, by setting Channel 0 to 5 V and modulating Channel 1 input by the GWN series around the mean value of 2.5 V. Each time series was 3-second-long and was repeated 8-10 times before the responses were averaged. The first 1.5 s of the recorded data was used to estimate the kernel values.

Photoreceptor output $y(t)$ was sampled at 10 kHz. It was then preprocessed by removing the mean value and trends, and down-sampled.

Once the matrices Y and P of Eq. 10 are constructed, there are several approaches to estimate θ with minimal error, such as the least squares regression by using Gram-Schmidt orthogonalisation (*179, 180*) or Meixner functions (*181*). Here, θ was approximated by the single value decomposition method (*182, 183*), in which the factorization of matrix P and the calculation of its Moore-Penrose pseudoinverse matrix, P^+, were carried out by the command *pinv(P)* in MATLAB. The linear least-squares estimation of θ, θ, was given by:

$$\theta = P^+ Y \tag{S6.5}$$

The computed kernels and the second half of GWN stimuli were then substituted to Eq. S6.2 to yield the model prediction of the photoreceptor response. The accuracy, or fitness, F, of the prediction was quantified by the complement of its mean squared error:

$$F = 1 - MSE = 1 - \frac{\overline{(y'-y)^2}}{\overline{(y-\bar{y})^2}} \tag{S6.6}$$

where y were the actual data measured from the photoreceptor voltage response and y' were the values simulated by the mathematical model.

Conventional simulation of intracellular responses to object motion

After determining and testing the Volterra model, we next approximated the light stimuli (input) delivered by the point-objects crossing a photoreceptor's receptive field. Since a photoreceptor's voltage response was assumed to be linearly correlated to light input, the response amplitudes to the subsaturating light flashes during the receptive field scans (see **Text S4**) were also considered linear measurements of the effective intensity from each light-point. Therefore, $u_m(t)$, created by one moving point-object, was modelled as 25 intensity steps, in which amplitudes were proportional to their corresponding flash responses (**Box S6.1**). The temporal width of each step was calculated according to the object speed. Similarly, $u'_m(t)$ of two moving point-objects was constructed from the superimposition of $u_m(t)$ and $u_m(t+\Delta)$, where Δ was calculated according to the point-objects' speed and their separation angle.

Box S6.1. Estimating the light input from a moving object. The effective light stimulus, which was delivered to a photoreceptor by a moving point-object, was estimated by a linear transformation of the cell's receptive field. The width of each intensity step was calculated according to the point-object speed.

Lastly, a photoreceptor's voltage response to point-object motion was simulated by substituting input $u_m(t)$ or $u'_m(t)$ and the kernels values to the zero- and first-order terms of Eq. S6.2.

Maximal responses to moving dots lag behind their actual positions

Neural images in the fly retina, lamina and medulla are generated by retinotopically mapping the surrounding light intensity distribution. This means that light coming from each point in space is sampled and processed by one neural cartridge (or neuro-ommatidia) (*68*). While a stationary object might be seen by several photoreceptors belonging to neighboring ommatidia due to their large acceptance angles (**Text S4**) and overlapping receptive fields (**Text S5**), the object position is almost certainly perceived on the photoreceptor's optical axis. This position, along its corresponding lamina/medulla cartridge below, produces the largest/fastest intracellular responses (**Box S6.2**; *cf.* **Text S4**) as it channels the maximum light influx into the rhabdomere (see **Text S2**).

Box S6.2. A R1-R6 photoreceptor's sensitivity is the highest at the center of its receptive field. (**A**) schematic showing how subsaturating isoluminant light pulses were delivered at different locations within each tested photoreceptor's receptive field. (**B**) The center stimulus (light-point No.13) typically evoked a response (orange trace) with the larger amplitude and faster rise-time than any stimulation at the flacks (light-points No.9 and No.10, respectively). (**C**) Average voltage responses of six photoreceptors to corresponding center and side stimulation. (**D**) Normalized responses make it clear that the responses to the center stimulus rise faster. (**E**) Time to the half-width response is significantly briefer (p = 4.84 x 10^{-5}) with the center stimulus. However, time-to-peak of the responses to center and side light-points shows more variability between individual cells (p = 0.125). Mean ± SD; two-tailed t-test; n$_{wild-type}$ = 6 cells.

Furthermore, it is customarily assumed that a moving point-object's position would be associated with the peak of its neural image. However, this does not necessarily mean that a R1-R6's response maximum would indicate the object position. Rather, it is more plausible - especially during high signal-to-noise ratio conditions (bright stimulation) - that the lamina circuitry (*cf.* **Text S2**, **Box S2.8E**) would be amplifying more the photoreceptor signal derivative. This is because the large monopolar cells (LMCs) are then more tuned to responding to the rate of light changes (*30, 46, 154, 178*).

In fact, neural latency might be compensated at the subsequent processing stages in the visual system, so that the peak of the travelling network response wave would closely follow the object's actual position, as is the case of the vertebrate ganglion cells (*165*). Theoretically, if such "correction" occurred maximally at the single neuron level, it would mark the coincidence

of two events. (**i**) The neuron's voltage response would peak as (**ii**) the object travels pass its receptive field center.

To examine whether, or to what degree, neural images of moving objects are compensated for latency at the first processing stage in the fly R1-R6 photoreceptors, we measured their intracellular voltage responses while presenting the fly with a bright dot moving at different speeds.

Box S6.3. Photoreceptor response maxima lag moving objects. (**A**) Channel 1 input was driven by an incremental ramp to create an image of a bright dot (point-object) moving from the No.1 light-point to the No.25 (front-to-back). Similar decremental ramps were used to produce back-to-front motion. (**B**) Intracellular responses of Calliphora photoreceptors to a moving point-object showed two response peaks: a large peak at t_2, which corresponded to the moment it travelled pass the cell's optical axis at t_1, and a smaller peak at t_4 caused by the exceptional brightness of light-point No.22, which was turn on at t_3. x was the object's travelling time. (**C**) An example of the linear correlation between t_2 and x. Below each data point is its corresponding stimulus (dot) velocity. (**D**) An example of the linear correlation between t_4 and x. Again, the corresponding dot velocities are shown.

Box S6.3B depicts a typical response waveform of a blowfly (*Calliphora vicina*) R1-R6 to a bright moving dot (the point-object in **Box S6.3A**). Let x be the time needed for it to travel through the 25 light-point array, t_1 be the moment when the dot pass the cell's optical axis, i.e. the corresponding light-point is turned on, and t_2 be when the intracellular response peaks. With varying x, and thus the dot speed, t_1, can be computed as:

$$t_1 = a \times x \tag{S6.7}$$

where the coefficient a is a constant. The aim was to align the 25 light-point array so that the point No.13 lies at the tested photoreceptor's receptive field center. Therefore in theory, a is approximately 0.48. However, the light-point No.13 might, in fact, be off-axis. For example, the cell's receptive field center could lie in between No.13 and No.12, causing inaccuracy in the calculation of a, t_1 and lag time b, which is given by:

$$b = t_2 - t_1 \tag{S6.8}$$

To overcome this ambiguity, we plotted t_2 against x, given that:

$$t_2 = t_1 + b = a \times x + b \qquad (S6.9)$$

Box S6.3C illustrates an example of the relationship between t_2 and x obtained from a *Calliphora* R1-R6. The two parameters fitted exceedingly well to a linear relationship (adjusted R-squared > 0.9999), in which coefficient a and lag time b were found as 0.486 and 14.62 ms, respectively. These data show that in this particular case, indeed light-point No.13 was close to the center of the cell's receptive field and that lag time b was virtually unchanged for different object speeds. The on-axial position of the light-point No.13 was later confirmed by 11 receptive field scans, all of which indicated that response elicited by a light flash from No.13 was the largest (data not shown).

The same photoreceptor was also stimulated by repeating a light flash, to which its voltage response peaked 15 ms later. The small difference between the lag time of motion response and the response time-to-peak to a flash does not readily imply network compensation of neural latency because of the essence of the two parameters. In case of the moving stimuli, photoreceptor was stimulated when the dot entered its receptive field, causing its intracellular voltage to start depolarizing before the dot reached the cell's optical axis. Conversely, the photoreceptor's response to light impulse was only elicited after the stimulus onset, making its time-to-peak slightly longer than the lag time.

In the classic flash-lag psychological experiment, where a flashing bar and another uniformly illuminated one travelled together, the former was perceived to be trailing (*184, 185*). Thus, it is probable that at some stage in the visual system, the voltage response peak (maximum) caused by moving object would display shorter delay than those elicited by increasing light intensity. In the present study, the light-point No.22 was 4-fold brighter than the others, as discussed in **Text S4**, and indirectly played the role of the flashing bar, causing a "local peak" in a photoreceptor's voltage response (**Box S6.3B**). Hence, to further examine neural latency of R1-R6 output, we next assessed the lag time, b', corresponding to this peak in the response. Given t_3 is the moment when No.22 was turned on, which can be calculated as:

$$t_3 = a' \times x \qquad (S6.10)$$

and t_4 is the time of the local response peak (**Box S6.3B**), lag time is defined as their difference:

$$b' = t_4 - t_3 \qquad (S6.11)$$

The relationship between t_4 and x could also be described by linear fitting with almost zero residue (**Box S6.3D**), yielding a' and b' values of 0.83 and 13.29 ms, respectively.

These data exemplify that a photoreceptor's response maxima, no matter whether caused by a point-object moving across its receptive field or by an unexpected increase in light intensity, show similar lag time characteristics. Both b and b' were independent of the object speed and comparable to the response time-to-peak, as induced by a comparable flash. These features were reproducible and general; observed in all 7 tested *Calliphora* photoreceptors, without exception (**Box S6.4**).

Animal	Flash response time-to-peak (ms)	Peaks corresponding to the receptive field center		Peaks corresponding to the light-point No.22	
		Lag-time b (ms)	Adj. R-Sqr	Lag-time b' (ms)	Adj. R-Sqr

Animal	Flash response time-to-peak (ms)	Front-to-back		Back-to-front	
		Lag-time b (ms)	Adj. R-Sqr	Lag-time b (ms)	Adj. R-Sqr
Calliphora	14.85 ± 0.78	14.6 ± 0.64	0.99985 ± 0.00012	13.9 ± 3.59	0.99985 ± 0.00017
		n = 7		n = 5	

Box S6.4. Response latency to dot motion analyses in *Calliphora* R1-R6s (Mean ± SD). Intracellular recordings were performed at 19 °C. The tested moving dot (point-object) velocities were: 334.6, 167.3, 111.53, 83.65, 66.92, 55.77, 47.8, 41.83, 37.18 and 33.46 °/sec.

Animal	Flash response time-to-peak (ms)	Front-to-back		Back-to-front	
		Lag-time b (ms)	Adj. R-Sqr	Lag-time b (ms)	Adj. R-Sqr
Wild-type *Drosophila*	23.81 ± 1.41	21.41 ± 4.5	0.99649 ± 0.0065	22.54 ± 4.15	0.99378 ± 0.007
		n = 12		n = 5	
hdc^{JK910}	24.4 ± 1.08	21.82 ± 1.36	0.9992 ± 0.0008	23.79 ± 5.72	0.9978 ± 0.003
		n = 3			

Box S6.5. Response latency to dot motion analyses in *Drosophila* wild-type and hdc^{JK910} R1-R6s (Mean ± SD). Intracellular recordings were performed at 19 °C. The tested point-object velocities were: 818.4, 409.2, 272.8, 204.6, 163.68, 136.4, 116.91, 102.3, 90.93 and 81.84 °/s. Note, these statistics are collected from individual recordings, not from paired data.

Drosophila photoreceptors' voltage responses did not clearly exhibit the "local peak" to the light-point No.22; possibly owing to their slower temporal dynamics. Nevertheless, the temporal position of the "global peak", measured by t_2 of wild-type (n = 12) and hdc^{JK910} (n = 3) photoreceptor outputs, consistently showed linear correlation to x and comparable lag-time/time-to-peak values (**Box S6.5**). The corresponding maxima in *Drosophila* data showed slightly larger lag time variations, and thus the linear fits were not as error-free as with the *Calliphora* data. Nonetheless, overall, the mathematical relation between their peak response lag time and the object speed appeared similar. The comparable wild-type and hdc^{JK910} photoreceptor output maxima to the tested point-object velocities, as recorded from their somata, suggests that their response dynamics mostly reflect equivalent phototransduction processes, with possibly only marginal influence from the lamina network.

Altogether, these data suggest that fly photoreceptors utilize the same mechanism (phototransduction machinery, see **Text S1** and **Text S2**) to sample intensity changes and object motions. Because the fly photoreceptors' peak responses lag behind the actual positions of the moving objects, the neural latency of moving objects is most likely compensated downstream by image processing within the interneuron networks, starting with the LMCs (*cf*. **Text S2**, **Box S2.8E**).

Response rise and decay to object motion show directional selectivity
As summarized in **Box S6.5**, *Drosophila* photoreceptors' maximum responses to a front-to-back or back-to-front moving bright dot did not exhibit clear signs of *latency compensation*, as indicated by their similar time-to-peak durations (estimated from the population means of individual unpaired recordings). Interestingly, in the paired recordings, however, the response rise and decay time-courses often showed considerable *latency modulation*.

Box S6.6 depicts intracellular responses to a moving dot, passing a photoreceptor's receptive field front-to-back and back-to-front at (**A**) 409, (**B**) 205, (**C**) 136 and (**D**) 102 °/s. Although their time-to-peak values appeared similar, the response rise and decay dynamics showed clear differences (**Box S6.6E-H**), which correlated with the dot speed and motion direction. We shall later show in **Text S7**, using high-speed video recordings of photoreceptor rhabdomeres, that their photomechanical contractions (*8*) occur in back-to-front direction. Light input modulation by these directional microsaccades can much explain the phasic differences in photoreceptor output to different directional point-object motions.

Box S6.6. Photoreceptor output's time-to-peak is insensitive to object motion direction but its waveforms rise and decay faster to back-to-front motion. Examples of intracellular voltage responses of the same dark-adapted *Drosophila* photoreceptor to a bright dot (point-object), which crosses its receptive field in back-to-front (green) or front-to-back (brown) directions (Mean ± SEM). The grey insets show a schematic of the fly compound eye structure, with the green Gaussian representing the photoreceptor's receptive field. (**A**-**D**) Voltage responses to 409, 205, 136 and 102 º/s object speeds, respectively, plotted over the whole duration of the corresponding stimuli. (**E**) Whilst the similarly timed respective response peaks showed no clear directional preference, the response waveforms, nevertheless, systematically rose and decayed earlier to back-to-front (green) than to front-to-back motion (brown), irrespective of the object speed. (**F**) The response rise-time (measured by time to half-maximal response), decreased with increasing object motion but it was always less to corresponding back-to-front (green) than front-to-back (brown) motion. This delay difference (white bars) was significant for all the tested object speeds and varied between 2 and 10 ms (mean ± SEM, $0.01 \le p \le 2.18 \times 10^{-14}$, $9 \le n \le 12$ trials, two-tailed t-test). (**G**) Because the response rise and decay times changed in unity, the resulting response half-widths to corresponding back-to-front and front-to-back object motion were largely similar. (**H**) Accordingly, the

response half-width decreased with increasing object motion, but showed only small (<4 ms) inconsistent differences (bars) between the corresponding back-to-front and front-to-back object speeds. Crucially, all these results are consistent with the fast light-induced photoreceptor contractions, which move their receptive fields in front-to-back direction, as observed directly in high-speed video recordings in vivo (see **Text S7**).

When a R1-R6 contracts to light input, its receptive field moves front-to-back (**Texts S7-8**). Thus, its responses to the back-to-front dot motion raise systematically slightly earlier (**Box S6.6E-F**). This rise-time lag reduces because the dot moves against the receptive field motion, whilst the rise-time lag increases during comparable front-to-back motion when the dot moves along the receptive field motion. In other words, a dot stayed a bit longer within a cell's receptive field during front-to-back motion than back-to-front motion. Such phasic differences were consistently observed in most recordings over the tested speed range. For example, 205 °/s back-to-front movement evoked narrower temporal response half-widths in 8/10 R1-R6s than the opposite movement. Similarly, 409 °/s back-to-front movement evoked narrower temporal response half-widths in 6/10 R1-R6s (in 2 cells, these were identical; and wider in 2).

In summary, a prominent feature of R1-R6s' voltage responses to opposing object motion directions is their similar time-to-peak values. This was found in all somatic recordings of *Calliphora* and *Drosophila* photoreceptors. Intriguingly, though, we further identified small (2-10 ms) but significant differences in the response rise and decay to front-to-back and back-to-front object motion. These phasic differences in photoreceptor output can be largely explained by each cell's directional photomechanical contractions, and we later show how these contribute to improving the fly's visual acuity (see **Texts S7-8**). It is plausible that these directional preferences would be further enhanced downstream at the level of network processing. Such scenarios are readily suggested by LMCs' phasic response properties to bright stimuli - responding most strongly to the rate of change in photoreceptor output (*29, 30, 71, 178*) - and by the richness of the lamina and medulla connectivity (*28, 29, 68*).

Classic theory greatly overestimates motion blur in R1-R6 output

We showed in **Text S4** that dark-adapted hdc^{JK910} photoreceptors have narrower receptive fields (acceptance angles) than their wild-type counterparts but broadly similar response dynamics (**Box S4.7**). Thus, the prediction is that hdc^{JK910} R1-R6s should produce slightly sharper neural images than their wild-type counterparts after dark-adaptation. Classic theoretical approaches have been used to predict how the spatial and temporal factors might jointly affect visual acuity (*3, 4*). Accordingly here, we first predict with them the motion blur effects on wild-type and hdc^{JK910} R1-R6 outputs. Later on, we test the ability of these cells to distinguish two dots moving together, separated by less than the cell's acceptance angles.

Since a fast moving bright dot can stimulate several photoreceptors virtually at the same time (**Box S6.7A**), theoretically, it should not be perceived as a single point but a streak, of which length is a function of object speed. This motion blur effect is classically quantified by the spatial half-width S of object's neural image. Because the spatial response in the retina has a similar waveform with the temporal response of a single photoreceptor (*3, 4*), S can be calculated as:

$$S = w \times T_h \qquad (S6.12)$$

where w is the object speed and T_h (*cf.* **Box S6.6G**) is the temporal half-width of a single photoreceptor response.

Box S6.7. Object motion blur according to the classic theory, applied to the neural images in the *Drosophila* eye as it was thought to affect them in the past. A, Hypothetical spatial pattern of an instantaneous voltage response at an ommatidial array produced by a moving point-object. Figure redrawn from (*4*). **B**, Spatial half-width of neural image of a moving point-object as a function of its speed in Dim and Bright conditions. For all the tested speeds, S_{hdc} were significantly smaller than $S_{wild-type}$ (p = 0.0036-0.049, t-test), except for 205 °/s in Bright condition, where the statistical test yielded p = 0.137. S was calculated using data at 19 °C. Mean ± SEM, $n_{wild-type}$ = 4-15, n_{hdc} = 3-16, two-tailed student test.

Box S6.7B illustrates the predicted relationship between the object speed and neural image resolution in wild-type and hdc^{JK910} *Drosophila*. These estimates imply that the spatial half-width of wild-type neural images should be 1-2° wider than that of the hdc^{JK910} during both the dim and bright conditions, reflecting wild-type R1-R6s' wider acceptance angles ($\Delta\rho$) (see **Text S4**, **Boxes S4.7A** and **S4.8B**). This prediction agrees with the previous theoretical works (*3, 4*), which used similar methods to indicate two distinct regions of image resolution. Thus, theoretically, at low object speeds, visual acuity should be mostly determined by a photoreceptor's spatial receptive field, but at high speeds, the motion-blur effect should increase rapidly, becoming the dominating factor. The corresponding trend differences (as separated by a thin dotted line) suggest that the point-object speed threshold dividing the two regions would be about 100-120 °/s.

Remarkably, however, we next demonstrate how these theoretical predictions greatly overestimate motion blur effects in R1-R6 output, and that these cells can, in fact, resolve image details finer than the half-width of their receptive fields, even at very high saccadic speeds.

In the second type of experiment, two bright dots, which were less than the half-width of a R1-R6's receptive field (6.8°) apart, crossed its receptive field at 409 °/s (**Box S6.8A**). Each tested photoreceptor's ability to distinguish the dots was assessed whether its response showed two clear peaks (**Box S6.8B**, solid line) or only one (dotted line). Quite unexpectedly, even at the low room-temperature of 19 °C, where phototransduction is slower than at the flies' preferred temperature (25 °C) (*45, 186*), 12/18 of wild-type R1-R6 photoreceptors and 15/16 hdc^{JK910} R1-R6s could clearly resolve the two dots. (Note that at 25 °C, every tested R1-R6 resolved them well; *cf*. **figs S9E-F**). Amongst the wild-type responses, the trailing peak was often larger than the leading one (**Box S6.8B**, solid line), whereas all but one hdc^{JK910} R1-R6 had the larger leading peak (**Box S6.8C**). This observation suggests that excitatory synaptic feedback modulation, which hdc^{JK910} photoreceptors lack, may enhance the second peak in the wild-type responses. Resolvability was further quantified by *D*-values:

$$D = \frac{d}{P}\%$$ (S6.13)

where P is the amplitude of the smaller peak and d is the depth of the response dip between the two peaks (**Box S6.8C**). In darkness, *D*-values measured from the mutant photoreceptors were, on average, more than double those of wild-type ($D_{wild-type} = 4.51 \pm 2.33\%$, $D_{hdc} = 10.5 \pm 5.23\%$), indicating that hdc^{JK910} R1-R6s resolve the two points more clearly than their wild-type

counterpart (**Box S6.8D**). Under light-adaptation (ambient illumination), while R1-R6s of both genotypes exhibited significant improvements in their image resolution ($D_{\text{wild-type}} = 9.81 \pm 3.7\%$, $D_{hdc} = 14.85 \pm 6.95\%$), the difference between the two groups decreased and was at the margin of statistical significance (**Box S6.8E**; p = 0.058, t-test). Taking into account only the cells in which D-values were measured in both dim and bright conditions, the enhancement of D-values to the ambient light change was quantified by their relative change:

$$R_C = \frac{D_{\text{Bright}} - D_{\text{Dim}}}{D_{\text{Dim}}} \% \tag{S6.14}$$

Box S6.8. Resolving two moving point-objects. (**A**) Two light-points (bright dots) moving together were presented in the tested photoreceptor's receptive field. The dots were 6.8° apart and travelled at 409 °/s in the front-to-back direction. For clarity, neutral density filters and the LEDs pads used for background illumination are not shown in this picture. (**B**) In Dim, 12 out of 18 wild-type photoreceptors could resolve them, showing the response waveform depicted by the continuous line, with the larger trailing peak. While 6 other could not distinguish the two objects; showing the waveform of by the dotted line. In Bright, 5 out of 6 examined wild-type photoreceptors resolved the two dots. (**C**) Response waveform of hdc^{JK910} photoreceptors exhibited two distinct peaks, with the leading one was the larger. In Dim, 15 out of 16 tested mutant photoreceptors could resolve the two objects and 14 of them displayed this waveform. In Bright, all of 8 recorded hdc^{JK910} photoreceptor responses resolved the objects. D-values were calculated from the amplitude of the smaller peak and the dip in between. (**D**) In Dim, D-values of hdc^{JK910} R1-R6 responses were significantly larger than their wild-type counterpart. $D_{\text{wild-type}} = 4.51 \pm 0.67\%$, $D_{hdc} = 10.5 \pm 1.35\%$, p = 0.00075, t-test, $n_{\text{wild-type}} = 12$, $n_{hdc} = 15$. (**E**) In Bright, the difference of D-values of the two photoreceptor groups was statistical insignificant. $D_{\text{wild-type}} = 9.81 \pm 1.65\%$, $D_{hdc} = 16.71 \pm 1.86\%$, p = 0.117, t-test, $n_{\text{wild-type}} = 5$, $n_{hdc} = 8$. (**F**) Changing from Dim to Bright condition, D-values of wild-type photoreceptors appeared to exhibit larger changes than those of mutant photoreceptors. However, the difference was not statistically significant due to the large cell-to-cell variation. R_C wild-type = 262 ± 126%, $R_{C\ hdc}$ = 31 ± 11%, p = 0.164, $n_{\text{wild-type}} = 4$, $n_{hdc} = 7$. (**D-F**) Mean ± SEM, two-tailed student test.

On average, wild-type D-values improved by 262%, ranging from 43% to 604%. These changes appeared to be markedly larger than those observed in hdc^{JK910} photoreceptors, which varied from 4% to 89% and averaged as 31%. Yet, the difference between the two groups was not statistically significant because of the large individual variations (**Box S6.8F**).

(In **Figure 3**, we further analyze the resolvability of those high-quality wild-type and hdc^{JK910} R1-R6s, from which we recorded the impulse response and receptive field measurements at the two adapting backgrounds - dim and bright, as well as voltage responses to both 205 and 409 ^{o}/s moving dots. Such data allowed us to compare the classic theory to the real recordings even more thoroughly.)

Together, these results show that the theoretical spatial half-width, *S*, grossly underestimates R1-R6 photoreceptors' image resolution. Recordings clarify that two bright dots that travel 409 ^{o}/s can, in fact, be resolved by a single photoreceptor, even when the dots are (6.8^{o}) less than the photoreceptor's acceptance angle ($\Delta\rho = 9.5^{o}$) apart. Therefore, a R1-R6 photoreceptor's real spatial half-width for the same high (saccadic) speed must be less than half of the theoretical estimate (~15^{o}; **Box S6.7B**). In other words, the classic theory overestimates the role of motion blur on *Drosophila* vision, as its R1-R6 photoreceptors resolve fast-moving dots beyond the predicted motion blur limit.

The recordings further indicate, consistent with hdc^{JK910} R1-R6s' marginally narrower acceptance angles (*cf.* **Text S4**, **Boxes S4.7-8**), that their spatiotemporal resolution is somewhat better than that of wild-type photoreceptors, both in dim and moderately bright conditions. However, when ambient light intensity was changed, the spatiotemporal resolutions of wild-type R1-R6s improved more. Here, possible contributing factors include:

- Slight (~4%) differences in the photoreceptors' rhabdomere diameters (see **Text S5**)
- Dynamic and homeostatic regulation of $[Ca^{2+}]_i$, membrane properties and synaptic feedback (*35*)
- Intracellular pupil (see **Text S2** and **Text S4**)
- Differences in photomechanical rhabdomere contractions (see **Texts S7-8**)
- Electrical coupling between the cells

Their potential roles are further discussed in **Texts S7-8**

Modeling R1-R6 output by the Volterra series method

Volterra kernels of each photoreceptor model were computed from the first half (1.5 s) of GWN data, before the other half of recorded light stimuli and voltage responses were used to validate the model. Because the output simulation accuracy depends upon input statistics and the model computation specifications, the system identification process was optimized by selecting suitable parameters.

Firstly, to test whether the selected 200 Hz input bandwidth was appropriate (**Box S6.9A**), we analyzed the resulting signal-to-noise (SNR) ratio of photoreceptor output (**Box S6.9B**). SNR decayed below 1 at around 66 Hz; at which point photoreceptor response contained more noise than signal. Thus, the GWN stimuli predictably activated a R1-R6 photoreceptor's whole frequency range.

Secondly, we assessed different sampling rates. According to the Nyquist-Shannon sampling theorem, a signal without frequencies higher than *B* Hz can be perfectly sampled and reconstructed (*25*) by sampling rate F_s of 2*B* Hz. Because the bandwidth of interest was 0-66 Hz, the data could be processed, in theory, at any sampling rate from 132 Hz to the recorded rate of 10 kHz, without compromising its information content.

We found that higher sampling rates yielded models, which predicted photoreceptor output with slightly higher accuracy. However, their kernels also exhibited larger fluctuations, and the kernels did not decay to zero over time, most likely due to high-frequency noise. For F_s = 1,000 Hz and higher, such fluctuations undermined the physiological meaning of the Volterra 1st-order kernel (**Box S6.9C**), which is the photoreceptor's impulse response (*187*). Thus, the

kernels computed from too richly-sampled data would be useful only for response prediction to this particular GWN stimulus.

On the other hand, while computations performed with lower F_s data would produce smoother kernels, a low sampling rate would also limit the model's other applications. For example, Volterra series models were used to simulate photoreceptor response to the image of moving objects created by the 25 light-point array. For an object moving at 409 $^{\circ}$/s, its travelling time across the array was 100 ms, or 4 ms per light-point. As the simulation required at least 2 data-points per light-point, F_s was chosen to be 500 Hz, at which rate reasonably smooth kernels could still be produced (**Box S6.9C**). Moreover, because the 1st-order kernel values decayed to zero at 50-60 ms, it was deemed that a 80 ms kernel length was sufficient for the computations.

Volterra series models, computed from data sampled at 500 Hz, could consistently predict response of *Drosophila* photoreceptors to GWN stimuli (for example, see **Box S6.9E**). On average, the model simulation accuracy, given by Eq. S6.6, was ~86% for both wild-type and *hdc*^{JK910} photoreceptors (**Box S6.9D**). These high F values confirmed that a linear Volterra series model could approximate light-adapted *Drosophila* photoreceptor output to the test stimulation appropriately.

Classic theory underestimates how well R1-R6s resolve fast moving dots
By approximating the light input directly from the receptive field measurements (**Box S6.1**) and the corresponding R1-R6 output by Volterra series (**Box S6.9C**), we could estimate each tested *Drosophila* photoreceptor's responses to moving bright dots. This was done by convolving the extrapolated light stimuli with the corresponding impulse responses.

The model predictions for a single moving stimulus were far less consistent than those for GWN stimuli. **Box S6.10A-B** show representative simulations with broadly acceptable and clearly unacceptable accuracies, respectively, together with the corresponding intracellular recordings.

From both the recordings and simulations, we further calculated the theoretical dot motion effects on the neural image resolution, or spatial half-width, *S* (**Box S6.10C**). As explained above (*cf.* **Box S6.7B**), the classic theory can only broadly suggest the relative differences between wild-type and mutant performances. Here, its application to the simulations further underestimated the spatial half-width predictions of wild-type recordings and over-estimated those of the *hdc* mutant.

Most critically, however, Volterra series models consistently failed to predict the fast phasic components of the recorded voltage responses, and thus their real resolvability, to two fast moving dots (**Box S6.10D**; *cf.* **Fig. 3**). The model simulations, and hence its underlying classic theory, always predicted lower resolvability than what we saw in the actual recordings. Further investigations (see **Texts S7-8**) revealed that this discrepancy reflected the missing biophysical mechanisms of the empirical black-box models (see **Text S1**). Specifically, the used photoreceptor models lacked: (**i**) the photomechanical rhabdomere movements, which shift and narrow the receptive fields, and (**ii**) the refractory sampling, which allows many microvilli (after the 1st dot) to recover from refractoriness (*5, 6, 40*) to respond to the 2nd dot. Moreover, by tuning and validating the Volterra kernels to GWN stimuli at relative steady-state, we inadvertently limited its exposure to (**iii**) excitatory dynamic synaptic feedback modulation (*30, 35*), which accentuates sudden changes in photoreceptor output. For these reasons, the responses of real R1-R6s, which naturally utilize the given mechanisms, showed systematically larger widths and 2nd-peaks, bestowing higher resolvability.

Box S6.10. Prediction accuracy of the Volterra series photoreceptor models to moving point-objects varies considerably. (**A**) Two examples of model simulations that were reasonably close to the actual intracellular recordings to the tested dot motion. (**B**) Two examples of simulations that clearly differed from the recordings. (**C**) Theoretical predictions of photoreceptor output spatial half-width, calculated from the recordings and simulations as a function of the point-object speed. Mean ± SEM, $n_{wild-type}$ = 9, n_{hdc} = 8. The theoretical spatial half-widths of the simulations differ from those of the recordings. *E.g.* the wild-type recordings (black) predicted consistently narrower S than the corresponding simulations (blue). The predicted resolvability of the resulting neural image, or spatial half-width (S), was consistently lower for the simulations than for the recordings. (**D**) Crucially, Volterra series models failed to predict how well the actual photoreceptor output resolves two close objects moving together very fast (shown for 409 and 818 ^{o}/s). For the 818 ^{o}/s prediction, we used here the fastest impulse response, recorded from another cell, but even so, the model still could not resolve the two dots. Thus, the actual spatial half-width of R1-R6s, limiting *Drosophila*'s resolving power at high image velocities, is about half of that estimated in (**C**). (Note, the dynamic biophysical mechanisms causing this difference – both in light input and photoreceptor output - are explained in **Text S8**, and illustrated in **Box S8.4A** and **D**). Recordings and simulations were at 19 ^{o}C.

We show in **Figures 4G-I**, **figure S8** and **Texts S7-8** that a new biophysically realistic microsaccadic sampling model, which allows for realistic refractory quantal phototransduction (*5*) and photomechanical rhabdomere contractions (*8*), yields (significantly better) theoretical predictions that closely approximate the real R1-R6 output. Importantly, in **Text S9**, we further show that R1-R6 output to two *dark* dots, moving at saccadic speeds, has the same relative resolution as their output to the corresponding two *bright* dots. Collectively, our results demonstrate that *Drosophila* photoreceptors resolve fast moving objects far better than what was believed previously.

Conclusions

In this supplement, we used intracellular recordings and classic theoretical approaches to study how fly photoreceptors encode moving bright dots. Model simulations about each tested R1-R6s' spatiotemporal responses were compared to the actual recordings to the same stimuli. We found that both wild-type and hdc^{JK910} photoreceptors resolved moving dots nearly equally well, and significantly better than the corresponding deterministic simulations. These findings demonstrate that the classic dynamic photoreceptor models (*2-4*), which lack knowledge about the underlying

phototransduction biophysics and photomechanics, grossly underestimate the spatiotemporal resolution of the real cells.

Animals counter self-motion blur effects by moving their eyes. This compensates for head and body movements by keeping the neural image position near stationary as long as possible (*31*). Interestingly here, a fly photoreceptor's response to two moving point-objects represents an opposite case where image motion, in fact, improves acuity. In the classic theory, to resolve two stationary objects, at least three photoreceptors are required so that the intensity difference in between can be detected. Because *Drosophila* photoreceptors' interommatidial angles (*2, 32*) vary from 3.4^o to 9.0^o and their average acceptance angle is ~9.5^o (**Text S4**), its eye should not resolve two point-objects 6.8^o apart. Nevertheless, responses of single photoreceptors, during even very fast (saccadic-speed) (*23*) movements, show large enough dips (in their temporal dynamics) to indicate that the objects are resolved neurally. Thus, an individual R1-R6 encodes space in time. In the classic theory instead, the outputs of several adjacent photoreceptors had to be processed together to distinguish two moving point-objects from one stationary object, in which brightness changes over time. This example highlights the inseparability of spatiotemporal information processing and visual acuity.

The unique advantages of the present study were the bespoke experimental equipment and stimulus paradigm. These allowed high-quality photoreceptor recordings with precisely controlled moving point-objects stimulation. Therefore, we could directly test and compare the theoretically predicted relationship between the neural image resolution and the object speed (*3, 4*) to the experimental data. However, the equipment also had limitations to be improved in future research. Wider object speed range is necessary, especially for testing insect eyes with fast responses. Owing to the long transient time, each light-point now required 2 ms switching period. Consequently, the minimum travelling time was 50 ms and the object speed limit in *Calliphora* experiments was 334 ^{o}/s, which is far slower than observed during the flies' saccadic flight behaviors (*175, 176*) (2,000-4,000 ^{o}/s). Whilst positioning the light-point array closer to the fly eye would increase object angular speed, it would compromise resolution as fewer light-points would then lie within a tested cell's receptive field.

In **Texts S7-9**, we show how both the enhanced resolvability of moving point-objects and the phasic modulation of their rising and decaying phases, as was shown here, emerge from the joint contributions of photomechanical rhabdomere contraction and its refractory information sampling.

Text S7: Photomechanical microsaccades move photoreceptors' receptive fields

Overview

This supplement describes a new powerful high-speed video recording method to measure photomechanical rhabdomere movements *in situ*, and provides important experimental and theoretical background information for the results presented in **Figures 4-5** in the main paper.

In this supplement:

- We utilize the optical cornea-neutralization technique (*49, 188*) with antidromic deep-red (740 or 785 nm peak) illumination to observe deep pseudopupils (photoreceptor rhabdomeres that align with the observer's viewing axis) in the *Drosophila* eye. We use an ultra-sensitive high-speed camera with a purpose-built microscope system to record fast rhabdomere movements across the compound eyes, while delivering blue-green stimuli (470 + 535 nm peaks) orthodromically into the eye.

- We show that light-activation moves rhabdomeres (both in dark- and light-adapted eyes) side-ways (horizontally). This movement starts after a 8-20 ms delay from the light stimulus onset, and reaches its peak in about 70-150 ms. Because these movements have fast onset and light intensity-dependency, which are similar to those of the R1-R6 photoreceptors' intracellular voltage responses to comparable stimuli, they must result from individual photoreceptors' photomechanical contractions; see (*8*).

- We show that *trp/trpl*-mutant photoreceptors, which have normal phototransduction reactions but lack the light-gated ion channels, also contract to light. Since these photoreceptors cannot produce electrical responses and thus communicate electrically or synaptically with other cells, including eye muscles, their <u>*contractility cannot be caused by eye muscle activity but must be intrinsic*</u>, supporting the earlier hypothesis (of phototransduction reactants interacting locally with the plasma membrane) (*8*).

- We show that light moves rhabdomeres fast in the back-to-front direction, while darkness returns them back to their original positions slower. Because the ommatidium lens inverts images, R1-R8 photoreceptors' receptive fields move in the opposite direction - front-to-back after light and back-to-front after darkness. Therefore, when front-to-back moving bright dots cross the eyes, the photoreceptors' receptive fields move along. But when bright dots cross the eyes in the back-to-front direction, the photoreceptors' receptive fields move against them (*cf.* **Text S6**).

- At the level of rhabdomere tips, the horizontal movements can be up to 1.4 μm, as measured occasionally in light-adapted eyes. Therefore, given the known optical dimensions, these photomechanical microsaccades can rapidly shift R1-R6 photoreceptors' receptive fields by ~5°. Remarkably, such a large image pixel displacement reaches the average interommatidial angle, $\Delta\varphi \sim 4.5\text{-}5°$, in the *Drosophila* eye (*cf.* **Text S4**; **Box S4.1**).

- We show that the light stimulus also contracts rhabdomeres axially (inwardly: 0.5-1.7 μm), down away from the lens. This transient increase in focal length should contribute in narrowing R1-R6's receptive fields dynamically. We further show that specific cone- and pigment-cells inside each ommatidium form an aperture, which is connected to the rhabdomere tips. During light stimulation, this aperture moves laterally with the rhabdomeres but only half as much. And since the ommatidium lens remains practically immobile, the light beam falling upon the rhabdomeres is shaped dynamically. These

observations mean that a R1-R6's receptive field must both move and narrow during dynamic light stimulation.

- We show that rhabdomeres of hdc^{JK910} histamine-mutant (*34, 48, 85*) R1-R6 photoreceptors, in which visual interneurones are blind (receive no neurotransmitter from photoreceptors), have broadly wild-type-like contraction dynamics, again refuting the role of eye muscle activity in the data. But interestingly, their light-sensitivity is about 10-fold reduced, similar to their voltage responses (*35*). In part, this may reflect hdc^{JK910} photoreceptors' smaller size. Given that hdc^{JK910} rhabdomere diameters are ~4% smaller than in wild-type (**Text S5**), their length should also be reduced in the same proportion. As the average wild-type R1-R6 is ~100 μm, hdc^{JK910} R1-R6s should be ~4 μm shorter. And indeed we find *in situ* that hdc^{JK910} rhabdomere tips are ~4 μm further away from the lens than the wild-type tips. In addition, the higher $[Ca^{2+}]_i$, caused by tonic excitatory synaptic feedback overload(*35*), may further affect their mobility, possibly retaining them in a slightly more contracted state.

Rapid adaptation caused by light-induced R1-R8 contractions

Atomic force microscopy (AFM) at the dissected *Drosophila* eyes' corneal surface(*8*) has shown up to 275 nm radial movements to brief light pulses, caused by transient *photomechanical* R1-R8 photoreceptor contractions. Such movements are too small and fast to see with the naked eye, and were initially considered: (i) only to participate in gating photoreceptor's transduction-channels, and (ii) possibly too small to affect fly vision in general. In this supplement, we use high-speed video microscopy to show that *in vivo* the underlying photomechanical rhabdomere (light sensor) movements are larger both laterally (horizontally: 0.3-1.4 μm) and axially (inwardly: 0.5-1.7 μm). Because these movements are also synchronous, ubiquitous, robust and reproducible, they influence how the fly eyes sample visual information about the world.

High-speed video recordings of light-induced rhabdomere movements

Dark-adapted dissociated photoreceptors rapidly contract to light(*8*) (*e.g.* **Movie S1**). It has been suggested that this contraction results from light-induced phosphatidylinositol 4,5-bisphosphate (PIP$_2$) cleaving, which modulates their rhabdomere membrane volume and so participates in gating the phototransduction-channels (*trp* and *trpl*) (*51*).

Here, we directly test the hypotheses that (i) the photomechanical photoreceptor contractions occur also in intact flies in normal stimulus conditions, and (ii) these movements serve the purpose of modulating light input to photoreceptors and thus photoreceptor output. We do this by recording high-speed video of how *Drosophila* photoreceptor rhabdomeres move to different light stimuli *in vivo*, and by analyzing and characterizing how these movements affect R1-R6s' receptive fields. Later on, in **Text S8**, we include their light input parameter changes in biophysically-realistic mathematical models to predict R1-R6 voltage output to moving visual stimuli.

Imaging setup for recording photomechanical rhabdomere contractions

We used the optical cornea-neutralization method to monitor how light stimuli evoke *Drosophila* photoreceptor rhabdomere movements. The imaging system was constructed upon an upright microscope (Olympus BX51), secured to a XY-micrometer stage on an anti-vibration table (MellesGriot, UK) (**Box S7.1**). To minimize light pollution in the recordings, the system was light-shielded inside a black Faraday cage with black lightproof curtains in the front, and the

experiments were performed in a dark room. For collecting and recording deep pseudopupil images, the system was equipped with a 40x water immersion objective (Zeiss C Achroplan NIR 40x/0.8 w, ∞/0.17, Germany) and an ultra-sensitive high-speed camera (Andor Zyla, UK), respectively.

Box S7.1. Microscope system for high-speed video recording of light-induced photoreceptor movements. (**A**) High-speed camera (Andor Zyla, UK) recorded images of deep pseudopupils in the eye of an intact living *Drosophila* under deep-red antidromic illumination (here 740 nm LED + 720 nm long-pass edge filter underneath the fly head). Each studied fly was immobilized inside a pipette tip. (**B**) A 10 ms blue-green light flash, delivered through the microscope system (orthodromically) into the left fly eye (above), was used to excite R1-R8 photoreceptors; the inset below shows R1-R7 rhabdomeres (blue) of one ommatidium just before the flash. (**C**) Light caused the rhabdomeres to twitch photomechanically in back-to-front direction (arrows) after 8-16 ms delay, with the photoreceptors being maximally displaced in ~100 ms from the stimulus onset. Invariably, this was seen as a sudden jump in the recorded rhabdomere position (red). (**D**) The difference in the rhabdomere position (displacement) before and after the flash, depended upon the light intensity, ranging between 0.3-1.4 μm; note a typical R1-R6 rhabdomere diameter is about 1.7 μm (**Text S5**). The frame subtraction (before and after the light flash) indicates that only the rhabdomeres that aligned directly with the blue/green light source moved (within the 7 central ommatidia; yellow area), while the rest of the eye remained immobile. Accordingly, the difference image shows little ommatidial walls, as these and other immobile eye structures became mostly subtracted away. In contrast, eye muscle activity, which is every so often seen with this preparation (see **Text S4**, **Box. S4.10**) occurs more gradually and moves all the eye structures together.

A *Drosophila* was gently fastened to an enlarged fine-end of a 1 ml pipette tip by puffing air from a 100 ml syringe at the large end until the fly head and ~1/5 of the thorax emerged outside (**Box S7.1A**). The head and thorax were carefully fixed (from the proboscis and cuticle) to the pipette wall in a preferred orientation by melted beeswax, without touching the eyes. The fly was then positioned with a remote-controlled XYZ-fine resolution micromanipulator (Sensapex, Finland) underneath the water immersion objective, using both visual inspection and live video stream on a computer monitor.

Antidromic illumination (through the fly head) revealed the deep pseudopupils of the fly eyes. It was provided with a high-power deep-red light source (740 nm LED with 720 nm high-pass edge-filter; or 785 nm LED with ± 10 nm bandpass filter), driven by a linear current LED driver (Cairn OptoLED, UK). Note, even very bright deep-red illumination, which is a

prerequisite for good signal-to-noise ratio high-speed video imaging, activates R1-R8 photoreceptors marginally. This is because their different rhodopsins' absorbance maxima are at much lower wavelengths (*29, 189*). The photoreceptors' near insensitivity to >720 nm red light was separately confirmed *in vivo* by ERG recordings (**Box S7.2**).

Orthodromic light stimulation (through the 40x objective into the eye), which evoked the photoreceptor contractions, was delivered by two high-power LEDs: 470 nm (blue) and 545 nm (green), each separately controlled by its own driver (Cairn OptoLED, UK). These peak wavelengths were selected to activate R1-R6s' rhodopsin (Rh1) and its meta-form near maximally, and so through joint stimulation to minimize desensitization by prolonged depolarizing after-potentials (PDA) (*42*). Simultaneous stimuli from the two LEDs were merged into one focused beam by a 495 nm dichroic mirror and low-pass-filtered at 590 nm. Pseudopupil signals of the observed fly eye (left or right) were split spectrally by another dichroic mirror (600 nm), and essentially only red image intensity information (≥600 nm) was picked up by the high-speed camera.

Box S7.2. Testing R1-R8 photoreceptors sensitivity to deep-red illumination by electroretinogram (ERG) recordings. (**A**) ERGs were recorded in intact *Drosophila*, placed inside a conical holder, to 1 s long very bright green (545 nm), red (740 nm) and deep-red (740 nm LED with 720 nm high-performance long-pass filter) pulses. A normal ERG contains a large but slow photoreceptor (DC) component and the faster on- and off-transients (*143*), which signal histaminergic transmission to lamina interneurons. (**B**) The very small ERG response (wine) to the deep-red pulse indicated very little R1-R8 photoreceptor activation. Mean ± SD shown, n = 6-7 wild-type flies.

Recording procedures

Light stimulus generation was performed by a custom-written Matlab (MathWorks, USA) program (Biosyst; M. Juusola, 1997-2015) (*7, 44*) with an interface package for National Instruments (USA) boards (MATDAQ; H. P. C. Robinson, 1997–2008). The length of the light stimuli (including the continuous red background and the blue/green stimulus patterns) was made to match the number of frames to be acquired by the camera, and the stimuli were externally triggered by Andor camera software (Solis). During the recordings, the frames were first buffered in the RAM in high-speed and then transferred on the computer's hard drive. Light stimulus intensity could be attenuated by a neutral density filter set (Thorn Labs, USA), covering a 5.3 log intensity unit range.

Key observations from unprocessed high-speed footage

High-speed video microscopy (**Box S7.1A**; e. g. **Movie S2**) from intact wild-type *Drosophila* eyes (n >> 100 flies) showed repeatedly and unequivocally that:

- Full-field light flashes evoked rapid local R1-R7 rhabdomere movements within those 7 ommatidia, which at the center of the imaged view, faced the blue/green stimulus source directly (**Box S7.3A**, orange area). Rhabdomeres in few other neighboring ommatidia also moved marginally (yellow area), but not obviously in other ommatidia. This meant that only the ommatidia that aligned with the blue/green stimulation absorbed the incident light, while those to one side reflected it. This local area, which showed photomechanical rhabdomere movements, closely matched *Drosophila*' normal pseudopupil (**Box S7.3B**).

- The rhabdomere movement was in the back-to-front direction (**Box S7.1B-D**), whilst in darkness, the rhabdomeres returned in front-to-back to their original positions more slowly. These dynamics and their directions appeared similar in both the left and right eye.

- Because light always moved the rhabdomeres back-to-front, the corresponding neural images of the left and the right eye must comprise left-right mirror symmetry. We show later in **Text S8** how this symmetry may allow *Drosophila* photoreceptors to encode orientation information during saccades or image rotation.

- The rhabdomere movement directions seemed homogeneous (at least in the 1st approximation) across each tested eye, appearing similar in its different regions: whether measured at its up, down, front or back ommatidia. Such "pixel interlocking" across the whole eye's visual field may help to preserve, or enhance, the neural images' spatial resolution of the world.

Box S7.3. Photomechanical rhabdomere movements were localized inside those 7 ommatidia that form the normal pseudopupil. (**A**) High-speed video recordings in the *Drosophila* eye showed clear rhabdomere displacement before (marked blue) and after (marked red) a blue/green flash only within 7 ommatidia (orange area). This was revealed by subtracting the corresponding frame contours. The rhabdomeres of these 7 ommatidia aligned directly with the blue/green stimulus, which was carefully centered above in the microscope port (*cf.* **Box. S7.1**). Marginal rhabdomere movements were further detected in 6 other neighboring ommatidia (yellow area). These results meant that only the rhabdomeres that faced the centered Orthodromic blue/green stimulus absorbed its light and contracted, while the rest of the eye reflected this stimulus and remained immobile. Note that this local rhabdomere activation pattern was restricted by the same eye design principle that causes the insect eye pseudopupil. (**B**) The *Drosophila* eye, in which photoreceptors were made to express green-fluorescence, displayed a green pseudopupil only form those 7 ommatidia that directly faced the observer (and the blue light source through the microscope lenses). This happened because these ommatidia (their rhabdomeres) both absorbed the incident blue light and their GFP-molecules released green light back to the observer's eye/camera, while the other ommatidia around reflected the blue light.

Image analysis

To accurately quantify the size and direction of the observed rhabdomere movements in time we devised specific image-analysis procedures. First, the stored images (raw data frames of each

recording) in the hard drive were exported as a tiff stack, in which pixel intensity range was set by the frames' minimum and maximum values, using the camera software (Andor SOLIS). ImageJ software was then used to convert the tiff-stack into a single tiff-file. Finally, custom-written Matlab programs (scripts) were used to process and analyze the images (**Box S7.4A**). These methods included:

i. Loading the image stack.

ii. Subtracting the median and mean from each frame and setting its negative values to (0. 0) to remove the dark noise background. This process was repeated for every frame.

iii. Calculating 2D cross-correlation between each frame and the reference frame.

iv. Selecting the cross-correlation values, which were ≥95% of the maximum (peak) value. This was repeated for every frame.

v. Calculating the weighted average position of the peak by using all the positions of the previous selection and using the cross correlation values as weights both in x- and y-direction. This was repeated for every frame.

vi. Subtracting the reference frame position from every frame.

Box S7.4. Cross-correlation image analysis to estimate photomechanical R1-R7 rhabdomere movements. (**A**) Analytical steps are shown for the reference frame at time zero (f_0), 2 ms before the 10 blue/green light stimulus pulse (red), and for the frame at the maximum rhabdomere displacement (f_{49}), 98 ms after (dark yellow). High-speed camera images of rhabdomeres were recorded using 750 nm red light. (i) Image stacks were uploaded, and (ii) the median of each frame was subtracted to remove its noise background. (iii) 2D cross-correlation was calculated for each frame, and (iv) the values within 5% of their peak value were selected. (v) The weighted mean peak positions gave each frame's x- and y-positions at its specific time point, and their distance, sqrt(x^2+y^2), the total rhabdomere displacement (in pixels) against the reference frame position. Notice that the 2D cross-correlation images have flipped x- and y-axis directions (up, U, appears down, D; front, F, appears back, B). (**B**) The resulting rhabdomere displacement distance and the corresponding x- and y-positions are plotted for each frame in time at 2 ms resolution (500 frames/s), against the reference frame position, $P_0(0, 0, 0)$. A comparable (inverted) atomic force microscopy data (cyan) closely matches the rise-time dynamic of the cross-correlation rhabdomere displacement estimate, validating our analytical approach. The analysis also implies that well dark-adapted photoreceptors may respond weakly to deep-red (740 nm) light onset (black trace 0-100 ms). Note R8 rhabdomere, which lies directly below R7, likely contracts too.

Quantifying rhabdomere travels and their receptive field shifts

In this study, the rhabdomere displacement measurements are given in microns (µm) and the resulting receptive field movements in degrees (°). **Box S7.5A** shows a whole image of a *Drosophila*'s left eye, as focused upon its rhabdomeres in the center and magnified by the microscope system to use the camera's full 2,048 x 2,048 pixel range. By placing a high-resolution µm-graticule on the same focal plane (**Box S7.5B**), we calibrated that the whole image is 303 x 303 µm. After converting the recorded rhabdomere displacements from pixels (**Box S7.4**) into microns, we then used the published parameters (*39*) about the *Drosophila* ommatidium optics (**Box S7.5C**) to translate these measurements into corresponding receptive field movements in degrees.

Drosophila ommatidium optical parameters were described by Stavenga (*39*) and Gonzalez-Bellido et al (*32*). Its biconvex facet lens focuses light to a rhabdomere (grey rectangle) tip. The outer and inner lens curvatures, $r_1 = -r_2$, are 11 µm, and its thickness, l_1, is 8 µm. Distance from lens to the rhabdomere, l_2, is 15 µm. Reflective indices, *n*, for the object space, lens and image space, respectively, are: $n_1 = 1$, $n_2 = 1.45$ and $n_3 = 1.34$.

We used standard ray transfer matrix analysis (*190*) to determine optical properties between the lens surface, P_1, and the rhabdomere tip, P_2. Both of these are represented as vectors

of their positions, y, and angles, θ: $P_1 = \begin{bmatrix} y_1 \\ \theta_1 \end{bmatrix}$ and $P_2 = \begin{bmatrix} y_2 \\ \theta_2 \end{bmatrix}$. Then, the optical system of the facet lens follows equation $P_2 = M\,P_1$, where

$M = \begin{bmatrix} 0.23 & 1.61*10^{-5} \\ -3.63*10^4 & 0.71 \end{bmatrix}$, which was calculated by using the ray transfer matrix analysis.

The transform matrix clarifies that the distance, y_2 (at P_2), mostly depends upon the angle θ_1 (of P_1). Thus, 1 µm movement gives $1*10^{-6}/1.61*10^{-5} = 0.0621$ (rad) angular change, which is 3.56 °/µm. This movement is an inverse of the visual field movement. Note that by using the highly comparable optical parameter values of Gonzalez-Bellido et al (*32*) (and considering the normal lens f-value variation across the *Drosophila* ommatidia), gives practically the same movement ratio (± 5% error).

Box S7.5. Calibrating the rhabdomere displacements in microns and their receptive field movements in degrees. (**A**) A whole image of a *Drosophila*'s left eye, the camera chip's full 2,048 x 2,048 pixel range. (**B**) a high-resolution graticule placed at the same focal plane as the image (a) gives the full image size of 303 x 303 µm. Thus 1 pixel ~ 0.1479 µm. (**C**) A schematic of the main optical components in a normal *Drosophila* ommatidium. Its optical properties indicate that a 1 µm rhabdomere displacement shifts its receptive field by 3.56°. The details about the used parameters and the basic calculations are given in the text.

Trp/trpl-mutants confirm the contractions' photomechanical origin

We then tested whether the rhabdomere contractions were generated by the photoreceptors themselves (photomechanically) or by eye muscle activity. This was done by recording in *trp/trpl* null-mutants, which express normal phototransduction reactants but lack completely their light-gated ion channels. Consequently, these photoreceptors did not generate electrical responses to light, and their eyes showed no ERG signal (**Box S7.6A**). Nonetheless, high-speed video recordings revealed that *trp/trpl*-mutant photoreceptors contracted photomechanically (**Box S7.6B**; *cf.* **Movie S1**). These observations are consistent with the hypothesis of the light-induced phosphatidylinositol 4,5-bisphosphate (PIP$_2$) cleaving from the microvillar photoreceptor plasma membrane causing the rhabdomere contractions (*8*).

When rhabdomeres move, the ommatidium lens system above stays still

Using the high-speed video microscopy, we next tested whether the *Drosophila* lens system or any other ommatidium structures moved during the rhabdomere movements (**Box S7.7**). In the experiments, a z-axis micromanipulator (Sensapex, Finland) was used to shift and reposition *Drosophila* in piezo-steps vertically. This allowed the focused image, as projected on the camera, to scan through each studied ommatidium, providing exact depth readings in µm. We then recorded any structural movements inside the ommatidia at different depths; from their corneal lens down to the narrow base, where the cone and pigment cells form an intersection between the crystalline cone and the rhabdomere tips (*37*).

Box S7.7. Rhabdomere movement is not due to eye muscle activity. (A) High-speed video recordings at different depths inside ommatidia before and after a bright light flash. (B) Cornea lens and the optical structures to the narrow base of the crystal cone remained virtually immobile. Below these, the cone cells showed movement that was half of that seen in the rhabdomeres. In the schematic, red dots and lines indicate adherens junctions that link the photoreceptors to the pigment and cone cells.

We found that when the rhabdomeres moved photomechanically (**Box S7.7B**, red trace; **Movie S3**) the corneal lens and the upper ommatidium structures were essentially immobile (grey), and normally remained so throughout the recordings. However, clear stimulus-induced movements were detected at the basal cone/pigment cell layer (orange; also inset) that connects to the rhabdomere tips with adherens junctions (*37*). Although it is less clear how much these structures reflected the rhabdomere motion underneath or were pulled by it, it is quite certain that they formed an aperture in the light path, which moved less than the light-sensors (rhabdomeres) below (orange vs. red). In **Text S8**, we analyze how such an interaction might dynamically narrow the R1-R6s' receptive fields to visual motion.

Overall, these results further verified that in normal stable recordings the used blue/green light flash was not evoking eye muscle activity, which would otherwise move the whole eye.

Light intensity-dependence of rhabdomere movements (*in vivo* dynamics)

Through a systematic and wide-ranging testing regime, we further discovered (**Boxes S7.8** and **S7.9**) that:

- Light-induced R1-R7 rhabdomere movements were robust and repeatable. **Box S7.8A** shows 10 trials (thin grey traces) and their mean (black) to a 10 ms bright flash measured from the same ommatidium. Between each flash, the eye was dark-adapted for 30 s. Characteristically, the rhabdomeres contracted to every light flash without a failure. Whilst these movements showed amplitude variations, their dynamic behavior was similar. Here, they reached their peak (mean = 0.806 µm, **Box S7.8B**) in about 140 ms and then decayed back to the baseline slower, mean τ_r ~190 ms (**Box S7.8C**).

- Dark-adapted R1-R7 rhabdomeres' maximum movement range (**Box S7.9A-I**) was considerably larger (0.3-1.2 µm) than the displacement range measured *ex vivo* by atomic force microscope (**Box S7.9B**, AFM$_{max}$ ≤0.275 µm) (*8*) on the corneal surface. This difference is hardly surprising. AFM measures axial (inward) cornea displacements, presumably resulting from a large number of simultaneous photoreceptor contractions underneath, whereas our high-speed video microscopy method measures orthogonal

(horizontal) rhabdomere movements locally at their source. Owing to the slight excitation caused by the bright 740 nm red-light background needed for *in vivo* imaging (**Box S7.4B**), the actual rhabdomere movements in full dark-adapted conditions could be even larger.

Box S7.8. Photomechanical rhabdomere motion is robust and repeatable. (**A**) Consecutive rhabdomere motions (grey thin traces) of the same ommatidium and their mean (black) to 10 stimulus repetitions (bright flash). The mean response recovery is fitted with an exponential, τ_r. (**B**) Each response maximum is shown against its time delay (time-to-peak) with the mean and SD. (**C**) Each response recovery time constant is plotted against its maximum with the mean and SD. Recording at t = 20 ^{o}C.

- The rhabdomere movement recordings (thin grey traces) from single ommatidia of individual flies vary more than the corneal AFM data (black traces). The sizable variations in their movement range and fine dynamics (such as minor oscillations) imply both considerable trial-to-trial (**Box S7.8**) and fly-to-fly variability (**Box S7.9**). Much of this is clearly physiological, as rhabdomere movement sizes and waveforms to specific stimuli were similar in one fly but often slightly different to those seen in another fly. However, because of the extreme sensitivity of our method (**Box S7.4B**, providing subpixel movement resolution), some of the variations reflected experimental noise. Such noise included microscopic mechanical vibrations in the recording system, minute spontaneous eye muscle activity (see **Text S4**, **Box S4.10**), and Poisson-noise, in which the image signal-to-noise ratio - as captured by the camera's CMOS sensor – reduced the more the faster the sampling. Appropriately, the average responses (red) to different intensity flashes were smoother, yet still remained much larger than in the AFM data.
- The average rise and decay time courses of the normalized rhabdomere movement recordings followed closely those of the normalized AMF recordings (**Box S7.9J**). Such dynamic conformity strongly suggests that both the methods capture accurately photomechanical R1-R8 photoreceptor contractions in their fast natural time resolution. But these observations also provided further evidence that *ex vivo* AFM data underestimate the actual magnitude of rhabdomere movements within ommatidia. In fact,

it seems possible that to maximize neural images' spatial resolution, the eye's architectural design dampens the lens system movement, while its sensors (rhabdomeres) contract. This would inadvertently impede the AFM signal (axial down movement), and any horizontal lens shift (**Box S7.7**), measured on the corneal surface.

Box S7.9. Comparing optically resolved wild-type rhabdomere movements to corresponding atomic force microscope (AFM) recordings from the corneal surface. To ease the comparisons, the AFM data is inverted. (**A**) Rhabdomere motion within individual ommatidia (grey thin traces) and their mean (red) evoked by the brightest 10 ms test flash are plotted against the largest AFM recording (black) to the brightest 5 ms test flash (data from Hardie and Franze) (*147*). The rhabdomere movement range is larger than what the AFM data suggests. (**B**) AFM recordings to a broad logarithmic light flash intensity range. (**C-I**) Rhabdomere movements vs. AFM recordings to light flashes of broadly comparable diminishing intensities. Notice that some individual rhabdomere movement recordings show minor oscillations that could be related to recording noise or physiological activity. (**J**) the mean and SD of normalized rhabdomere movements to the brightest test flash are compared with the normalized AFM recordings to three different test flash intensities. All these AFM recordings fall within the SD of the given rhabdomere recordings.

In vitro rhabdomere movements

As further controls, we measured photomechanical R1-R8 rhabdomere contractions of freshly dissociated ommatidia (*8*) to green (480 nm) light flashes using high-speed video recordings with infrared 850 nm background illumination (**Box S7.10**). The benefit of this *in vitro* method was that it provided a clear side-view of the tested wild-type and mutant rhabdomeres, enabling us to estimate their axial (longitudinal inward) contraction component; or how much the rhabdomere tip moved away from the ommatidium lens (**Movie S1**). *In vivo*, such fast lengthwise light-sensor movements should contribute to R1-R6 photoreceptors' transiently narrowing receptive fields (see **Text S8**).

We found that after dark-adaptation bright flashes could evoke 0.8-1.7 µm longitudinal rhabdomere contractions. These were characteristically accompanied by synchronous (about equally large) crosswise movement (or twist), which likely forms the basis of the sideways rhabdomere displacement; seen during the *in vivo* recordings (*e.g.* **Box S7.1D**).

Box S7.10. Photomechanical rhabdomere contractions in dissociated ommatidia. (**A**) Two frames of high-speed video footage of contracting R1-R8 photoreceptor rhabdomeres in vitro, evoked by a bright flash. The size-view imaging reveals the size and dynamics of their photomechanical lengthwise changes. (**B**) Characteristic maximum longitudinal rhabdomere contractions range from 0.8 to 1.7 µm. Notably, many rhabdomeres also twist during these contractions, providing additional crosswise movements. See **Movie S1**.

Estimating light intensity falling upon the rhabdomeres

Only in 2 out of 21 tested *Drosophila* eyes, the rhabdomeres moved unmistakably (twitched) to a very dim 10 ms blue/green LED flash, in which intensity was reduced ~200,000-fold by neutral density filters. Therefore, in these two positive occasions: (i) the resulting response must have been quantal with (ii) the 10 ms flash maximally containing ~1-3 absorbed photons. Furthermore, because this flash only succeeded in ~1/10 eyes, its average maximum intensity could only be ≤3/10 photons/10 ms, i.e. ≤0.03 photons/ms. This means that the brightest flash (logI(0)), which was not filtered, could maximally contain ≤6,000 photons/ms, or ≤6 million photons/s, making the used light intensity range natural and directly comparable to that used for the intracellular recordings (**Figs 1-3** in the main paper).

This reasoning is in line with the similar LED driver settings used in all the experiments, and the similar V/log(I) and (µm)/log(I) functions, which resulted from these experiments.

Photomechanical rhabdomere movements vs. R1-R6s' voltage responses

We next compared intracellular wild-type and hdc^{JK910} R1-R6 voltage outputs, evoked by 10 ms light flashes after brief dark-adaptation, to their characteristic rhabdomere movements (**Box S7.11A-D**). R1-R6 output and rhabdomere motion exhibited broadly comparable delays (or dead-time), but overall the rhabdomeres moved considerably slower than how their voltage was changing (*cf.* the thick black traces in **A** to **B**, and the thick red traces in **C** to **D** for similar flash intensities).The voltage responses peaked 25-30 ms from the light onset, and the rhabdomere movements 40-120 ms later. At this point, the photoreceptors had almost repolarized back to their dark resting potential (indicated by zero ordinate). Moreover, the recordings suggested that

for a given flash the rhabdomeres in *hdc^JK910* eyes typically moved a smaller distance and returned faster to their original positions than their wild-type counterparts.

To further characterize their light-dependent differences, we plotted the maximum wild-type and *hdc^JK910* rhabdomere movements of many flies against the corresponding flash intensities over the whole tested light range. The analysis revealed that:

- Both wild-type and *hdc^JK910* rhabdomere movements increased with flash intensity (**Box S7.11E-F**), following a characteristic sigmoidal displacement/logI-relationship.
- The average *hdc^JK910* rhabdomere movements were slightly smaller than those in wild-type eyes. In part, this could result from *hdc^JK910* R1-R6s being smaller than their wild-type counterparts, and embedded deeper inside the ommatidia. As their rhabdomere diameters are ~4% smaller than those of the wild-type (**Text S5**), their length should also be reduced proportionally; with the average wild-type R1-R6 being ~100 μm tall, *hdc^JK910* R1-R6s should be ~4 μm shorter. Accordingly, by using the z-axis micromanipulator (see above), we measured *in situ* (rest) that *hdc^JK910* rhabdomere tips were 3.5 μm further away from the lens than the wild-type tips (*hdc^JK910*: 27.4 ± 1.2 μm; wild-type: 23.9 ± 1.2 μm; n = 6 ommatidia, 6 flies).
- *hdc^JK910* rhabdomere contractions further implied reduced sensitivity (**Box S7.11G**), seen as a 10-fold right-shift in their displacement/logI-curve in respect to the wild-type data.
- Interestingly, this sensitivity difference resembles that seen between the wild-type and *hdc^JK910* ERGs (**Box S7.11H**). We have recently provided compelling evidence that the missing histaminergic (inhibitory) neurotransmission from *hdc^JK910* photoreceptors to interneurons (LMCs and amacrine cells) causes a tonic excitatory synaptic feedback to R1-R6s, depolarizing them ~5 mV above the normal wild-type dark resting potential (*35*). Thus, *hdc^JK910* R1-R6s should experience a tonic Ca^{2+} influx and be permanently in a more "light-adapted" state. Here, our results suggested that Ca^{2+} overload may desensitize the biophysical machinery that moves the rhabdomeres, reducing its dynamic range.

Box S7.11. Comparing dark-adapted wild-type and hdc^{JK910} R1-R6 photoreceptors' photomechanical rhabdomere contractions to their corresponding electrophysiological responses at 19°C. (**A**) Intracellular voltage responses of dark-adapted wild-type R1-R6s to a 10 ms subsaturating light pulse, delivered at the center of their receptive field. (**B**) Characteristic non-averaged photomechanical rhabdomere movements (displacement in time) of a typical wild-type fly, as quantified by the cross-correlation analysis (*cf.* **Box S7.4**) and image calibration (*cf.* **Box S7.5**), to very dim (log(-3.5)), moderate (log(-2)) and very bright (log(0)) 10 ms light flashes. (**C**) Intracellular responses of dark-adapted hdc^{JK910} R1-R6s to a similar stimulus (as in a). (**D**) hdc^{JK910} rhabdomere movements to very dim, moderate and very bright 10 ms flashes are characteristically slightly smaller than the corresponding wild-type recordings (**B**). (**E**) The wild-type and (**F**) hdc^{JK910} rhabdomere displacements increase with logarithmic light intensity. (**G**) Mean wild-type rhabdomere movement was larger than that of hdc^{JK910} over the tested intensity range, with the hdc^{JK910} photoreceptors' apparent right-shift indicating a 10-fold reduced sensitivity. (**H**) This right-shift is broadly similar to these photoreceptors' V/LogI characteristics, measured from their ERG slow components (*35*).

Light-adapted rhabdomere motion reflects rhodopsin/meta-rhodopsin balance

Given the slightly reduced back-to-front rhabdomere mobility of hdc^{JK910} photoreceptors, we next asked whether prolonged light-adaptation itself would reduce wild-type photoreceptors' rhabdomere movement. To study this question, we examined how the rhabdomeres in seven individual flies responded to different light impulses at different adaptation states.

First, to obtain the baseline responses in each fly eye, we recorded their rhabdomere movements to a bright and a very bright full-field green-blue flash after 30 s of dark-adaption (**Box S8.10A-B**, respectively). As before, we found that the brighter the light flash, the larger and the faster their rhabdomere movements were on average.

Interestingly, however, when the flies were adapted to a moderate or bright blue (470 nm) light field for 30 s, which converts most (if not all) rhodopsin Rh1 to its active meta-form (causing PDA, or prolonged depolarizing afterpotential), the flashes now evoked only weak or no rhabdomere movement (**Box S8.9C-D**). Such reduced mobility somewhat resembled that in

some hdc^{JK910} rhabdomeres (**Box S7.11D**). These observations can be explained, at least in part, with the basic molecular model proposed by Hardie and Franze (*8*) for rhabdomere contraction. Here, meta-rhodopsin would continuously activate G-protein and in turn phospholipase C (PLC). PLC would then cleave most PIP_2 off the microvillar membrane, causing a tonic photoreceptor contraction, which facilitates light-gated channel openings and thus increases Ca^{2+} influx. But because a 10 ms bright green (545 nm) light pulse would convert only some fraction of meta-rhodopsin back to its non-activated form, this effect would be small proportionally, and could only partially rescue the rhabdomere contractibility.

Box S7.12. Wild-type photomechanical rhabdomere movement dynamics at different light-adaptation states. Characteristic non-averaged photomechanical rhabdomere movements of 7 wild-type flies and their means (thick traces) to bright (logI(-1); left) or to very bright (logI(0); right) light flashes. The recordings were performed in relative darkness (>720 nm) or when the eyes were adapted for 30 s to different green (545 nm) or blue (470 nm) light intensity levels; from weak (logI(-3)) to bright (logI(-1)). (**A-B**) In dark-adaptation, consistent with the results in **Box S7.11**, the brighter the flash, the larger and faster the evoked mean rhabdomere movements. (**C-D**) Prolonged blue light exposure converted virtually all Rh1-rhodopsins to their active meta-form, causing a PDA (prolonged depolarizing afterpotential). PDA increases the photoreceptors' intracellular calcium load, cleaving PIP_2 from the plasma-membrane and so keeping them in a contracted state. A very bright green flash rapidly converts some of the meta-Rh1 back to Rh1, enabling small and brief rhabdomere movements, which resemble those seen with hdc^{JK910} flies (*cf.* **Box S7.11**). (**E-H**) Adaptation to different green-light intensity levels does not abolish rhabdomere movements to light increments. These movements can, in fact, be larger than in the same cell's dark-adapted state, as was seen in Fly4 and Fly6 recordings. Notice that a 1.2-1.4 µm rhabdomere displacements means ~4-5° shifts in a R1-R6 photoreceptor's receptive field (*cf.* **Box S7.5**), which can be more than the average *Drosophila* interommatidial angle.

Rhabdomere contractions resensitise refractory sampling units

On the other hand, under more natural green light-adaptation (**Box S7.12E-H**), the eyes' normal back-to-front rhabdomere contractibility to light increments was retained, and sometimes even increased, in respect to their dark-adapted baseline responses (*cf.* **Box S7.12A-B**). Consequently in every fly eye (7/7), the rhabdomeres moved more when green-adapted than when moderately blue light-adapted (**Box S7.12C**), as quantified by their flash-induced maximum displacements (**Box S7.12I**).

These results indicate that in normal spectrally-broad natural environments, light increments (positive contrasts) will evoke fast evasive rhabdomere movements, steering them away from pointing directly to a bright light source. This novel photomechanical adaptation should together with the slower screening pigment migration (intracellular pupil mechanism, **Text S2**) help to recover (resensitise) a rhabdomere's refractory sampling units (30,000 microvilli). Thus, the rhabdomere movements likely participate in optimizing photon sampling for maximum information capture (see **Text S1-2**). This further means that a rhabdomere's state of contraction is constantly being reset to the ongoing light input, proving the capacity to respond to the next stimulus increment. Therefore, although being slower, in many sense, *Drosophila* photoreceptors' mechanical adaptation resembles the adaptive resensitization of the inner ear hair-cells (*137, 191*).

Text S8: Microsaccadic sampling hypothesis for resolving fast-moving objects

Overview

This supplement describes a new "microsaccadic sampling"-hypothesis that predicts a *Drosophila* photoreceptor's voltage responses to moving dots, and provides important background information about the experimental and theoretical results presented in **Figures 3-5** in the main paper.

In this supplement:

- We use our biophysical *Drosophila* R1-R6 model (**Text S1**), with different degrees of photomechanical rhabdomere movements modulating light input (**Text S7**), to simulate photoreceptor voltage output to moving bright dots that cross its receptive field in different directions, speeds and inter-distances.
- By comparing the model simulations to intracellular recordings, we reveal the likely biomechanics that allow *Drosophila* photoreceptors to resolve adjacent dots at high saccadic velocities (**Text S6**).
- We show that when a rhabdomere contracts away from the ommatidium lens's focal point, its receptive field must move and narrow dynamically. Together these processes actively reshape both the light input and photoreceptor output to separate and sharpen neighboring visual objects in time, improving their resolvability.
- Crucially, with such photomechanical light input modulation, the model photoreceptor output closely approximates that of the real R1-R6s, as recorded to two moving bright dots crossing their receptive fields at different speeds in **Text S6**.
- Hence, with refractory photon sampling and photomechanical rhabdomere movements, we can correctly predict and convincingly explain visual acuity of *Drosophila* photoreceptors to moving objects.

Modeling a R1-R6's receptive field dynamics to moving dots

Based on the combined results in **Text S1-7**, we develop a new "microsaccadic sampling"-hypothesis, which predicts how photomechanical rhabdomere contractions (microsaccades) move and narrow *Drosophila* R1-R6 photoreceptors' receptive fields (RFs) to resolve fast-moving objects. We present extensive analytical and experimental evidence to show how these mechanisms operate with the photoreceptors' refractory information sampling to reduce light-adaptation and to increase the spatiotemporal resolution of their voltage responses, improving visual acuity.

Using the results in **Text S1-7**, we can now work out the biomechanics, which allow a R1-R6 photoreceptor to resolve two close bright dots crossing its receptive field at saccadic speeds. We do this systematically by comparing the output of our biophysical model (*5, 6, 40*) (**Text S1**), in which input is modulated by different degrees of rhabdomere photomechanics (**Box S8.1A-C**), to the corresponding recorded real R1-R6 output (**Box. S8.1D**). Specifically, we consider three input modulation models:

- A.   *Stationary rhabdomere model (receptive field is fixed)*
- B.   *Photomechanical rhabdomere model (receptive field moves)*
- C.   *Photomechanical rhabdomere model (receptive field moves & narrows)*

In the following, to make these different models (*a-c*) directly comparable, we first present the

findings for two bright dots, which cross the receptive field at 205 $^\circ$/s (**Box S8.1**), before generalizing the results for a vast range of stimuli and giving more examples. Note that these simulations and recordings were performed at 19-20 $^\circ$C (room temperature). Later on in this supplement, we show how at the *Drosophila*'s preferred temperature range (24-25 $^\circ$C) (*186*) these dynamics are naturally faster (*45*) and improve visual acuity further.

A. Stationary rhabdomere model (receptive field is fixed). This approach is broadly analogous to the classic theory (*cf.* **Text S4** and **Text S6**). It was implemented in four steps (**Box S8.1A**):

(i)   Two bright dots, which were 6.8° apart, crossed a R1-R6's receptive field (RF; $\Delta\rho$ = 8.1 $^\circ$) front-to-back at the saccadic speed of 205 $^\circ$/s.

(ii) Concurrently, the ommatidium lens focused their light onto a rhabdomere tip.

(iii) The resulting dynamic light input at the rhabdomere tip was a convolution of the two dot intensities with the cell's receptive field over time (*3, 4*).

(iv) The light input (photons/s) drove the photon sampling and refractory quantum bump (QB) production of 30,000 microvilli (*5, 6, 40*), which formed the rhabdomere. The resulting macroscopic photoreceptor output (*51*) (mV) dynamically integrated the QBs.

Box S8.1. Microsaccadic sampling-hypothesis predicts R1-R6 output dynamics to fast moving bright dots. We compare stochastic sampling model predictions of increasingly realistic light input modulation (**A**-**C**) to actual recordings (**D**). In each case, the light stimulus is two fast moving bright dots (i), which cross a R1-R6 cell's receptive field (RF, half-width = 8.1°) front-to-back at 205 $^\circ$/s. The immobile ommatidium lens focusses their light onto a rhabdomere tip (ii). But because of diffraction, their images blur (airy disk). Interestingly, however, in *Drosophila*, the airy disk half-width (~1.7° or ~0.6 µm) is smaller than the average R1-R6 rhabdomere diameter (1.7 µm; **Text S5**, **Box S5.1**). The dynamic light input falling upon the rhabdomere (iii) is then estimated by convolving the two dot intensities with the cell's receptive field over time (i). This light input is sampled stochastically by 30,000 microvilli, in which quantum bumps integrate the dynamic voltage response (iv). (**A**) In the simulations, where the rhabdomere is stationary (ii) during the dots flyby (corresponding to the classic theory (*3, 4*)), their light fuses together (iii). This is because the simulated photoreceptor's receptive field half-width (8.1°) is wider than the distance between the dots (6.8°). Accordingly, the photoreceptor output

(iv) cannot distinguish the dots. (**B**) Next, we include the light-induced fast back-to-front rhabdomere movement in the model (ii; as measured in the high-speed video recordings in **Text S7**, **Box S7.1**), but keep the cell's receptive field shape the same. The resulting receptive field movement makes the light input rise and decay slightly later than in the previous case (iii; *cf.* the green and dotted grey (**A**) traces), but still cannot separate the two dots. The resulting photoreceptor output (iv) shows a single peak, which is slightly broader than the output from the immobile cell (**A**). (**C**) In the full model, as the contracting rhabdomere jumps away from the focal point (ii), the light input both moves with the cell's receptive field and transiently narrows it from 8.1^o to 4.0^o. These processes differentiate the light input (from the dots) into two separate intensity spikes over time (iii), which the photoreceptor output can clearly separate in two distinctive peaks (iv). (**D**) *In vivo* intracellular photoreceptor recordings from a wild-type R1-R6 to the two moving dot stimulus show comparable dynamics to the full model's prediction. In the simulation, the same cell's RF (i) was moved and narrowed dynamically (ii-iii) according to our hypothesis (**C**). This close correspondence, which occurred even without the additional response accentuation from the synaptic feedbacks (see **Text S6**), implies that *Drosophila* photoreceptors' RFs must move and narrow with their photomechanical rhabdomere contractions. The optical/structural mechanism behind the RF narrowing needs further examination. We propose four viable hypotheses for it in this supplement.

Characteristically in this approach, input modulation reduced to a single peak as the RF convolution fused the light from the two dots together (iii). This made them irresolvable to the discrete photon sampling (iv). Specifically here, light input could not separate the two dots because the given receptive field half-width (8.1^o) was wider than the distance between the dots (6.8^o). The resulting model output, thus, failed to capture the resolvability of the real R1-R6 output (**Box S8.1D**, iv).

Markedly, the model prediction (**Box S8.1A**, iv) much resembled those of the deterministic Volterra-models, in which similarly broad RFs were used in the calculations (see **Text S6**).

B. Photomechanical rhabdomere model (receptive field moves). In this case (**Box S8.1B**), we kept the cell's receptive field shape the same (i; $\Delta\rho = 8.1^o$) but included the light-induced fast horizontal (back-to-front) rhabdomere movement in the model (ii). The given model parameters (**Box S8.2**) were fixed to closely approximate the experimentally observed rhabdomere movement dynamics (see **Text S7**). Here, the following deductions were made:

- Each photoreceptor contracted independently to light. Although this movement is linked to the synchronous contractions of its neighbors inside the same ommatidium, such cooperativity makes no difference to the model.
- Rhabdomere movement started 8 ms after the first dot reached the outer rim of a photoreceptor's receptive field, called the trigger zone. This delay matched both the delay in R1-R6 output (**Text S7**: **Box S7.5A**) and the apparent dead-time in AFM data (*8*) to a bright light flash (**Text S7**: **Box S7.4J**). The trigger zone was 14.6^o from the photoreceptor receptive field center, matching the typical spatial threshold where dark-adapted wild-type R1-R6s responded faintly to subsaturating peripheral light flashes (**Text S4**: **Box S4.7C**).
- Maximum horizontal rhabdomere movement was set to be 0.58 µm, corresponding to a 1.6^o shift in its receptive field. This value is close to the measured average of the maximum light-induced rhabdomere movements in wild-type fly ommatidia (**Text S7**: **Box S7.7E**).
- Rhabdomere movements had two phases. In the 1st phase, a rhabdomere moved 1.6 ^{o} degrees in back-to-front direction for 100 ms, reaching its maximum displacement. This

caused a receptive field to shift in the opposite, α (front-to-back) direction. Importantly, the 1st phase could not be disturbed. In the 2nd phase: the rhabdomere slowly returned to the original position in 500 ms. The 2nd phase could be disturbed. Both the phases followed linear motion.

Box S8.2. Parameters for modeling a R1-R6's receptive field (RF) dynamics caused by its rhabdomere contraction							
Trigger zone: **trig**	Startin g RF half-width: $\Delta\rho_{start}$	Delay before rhabdomere motion: **lag**	Rhabdomere motion: (time-to-peak) **Phase 1**	Resulting parallel RF shift: RF_{shift}	RF shift direction: **α** (front-to-back)	Rhabdomere motion: **Phase 2**	Ending RF half-width: $\Delta\rho_{end}$
14.6 °	8.1°	8 ms	100 ms	1.6°	0	500 ms	4.0°

As in the first case, the moving receptive field model (**Box S8.1B**) was implemented for the same stimulus in four steps (i-iv):

(i) Two bright dots, 6.8° apart, crossed a photoreceptor's receptive field (RF; $\Delta\rho = 8.1°$) front-to-back at the saccadic speed of 205 °/s.

(ii) The ommatidium lens focused their light onto a rhabdomere tip. But here, after 8 ms delay, the rhabdomere started to move back-to-front as it contracted photomechanically.

(iii) The resulting dynamic light input was, therefore, a convolution of the two dot intensities and the cell's receptive field, which moved at different speeds in the same direction (front-to-back).

(iv) The light input drove the photon sampling and refractory quantum bump (QB) production of 30,000 microvilli, while the resulting macroscopic photoreceptor output summed up the QBs.

We found that the resulting receptive field movement caused (iii) the light input rise and decay slightly later than when the rhabdomere was immobile (*cf.* green and dotted grey traces). However, because the two dots were close and crossed the cell's receptive field fast, the given co-directional receptive field motion failed to separate the light from them. Thus, (iv) the resulting photoreceptor output showed a single peak, which was slightly broader than the output from the immobile model.

C. Photomechanical rhabdomere model (receptive field moves & narrows). There is a large disparity between the measured (**Text S4**: $\Delta\rho = 7.00 – 11.65°$) and the optical waveguide theory derived (*39*) ($\Delta\rho = 3.5 - 5.3°$) acceptance angles of dark-adapted R1-R6s. Even Snyner's simple formula (*88*) (**Text S4**: Eq. S4.3, page 17), which overestimates (*39, 90*) $\Delta\rho$ from the measured *Drosophila* ommatidium optical dimensions, gives a theoretical upper bound ($\Delta\rho <5°$) that is smaller than the smallest intracellularly measured values.

$$\Delta\rho = \left(\sqrt{\left(\frac{0.545}{16}\right)^2 + \left(\frac{1.7}{21.36}\right)^2} \right) \times \frac{180}{\pi} = 4.9601°$$

$$\Delta\rho = \left(\sqrt{\left(\frac{0.545}{17}\right)^2 + \left(\frac{1.7}{21.36}\right)^2} \right) \times \frac{180}{\pi} = 4.9161°$$

These upper bounds for green light (545 nm) were obtained for the smallest and largest ommatidium lens (16-17 µm) with average rhabdomere diameter EM measurements (1.7 µm) (**Text S5**), using the experimentally estimated focal length (21.36 µm) (*32*).

Moreover, interestingly, Götz estimated from *Drosophila* optomotor behavior, using the early flight simulator system (*38*), that in bright illumination R1-R6 $\Delta\rho$ would be 3.5^o.

To resolve the paradox between the conflicting experimental and theoretical $\Delta\rho$-estimates, which in the past were based upon histological measurements of fixed/stained (dead/immobile) retinal structures, we hypothesized that the photomechanical rhabdomere contractions not only move a photoreceptor's RF but also dynamically narrow it (**Box S8.1C**). What is more, we reasoned that the RF narrowing should depend upon stimulus history; the cell's ongoing light exposure. Therefore, our specific prediction was that when moving bright dot stimuli entered a cell's receptive field, the resulting dynamic input modulation would transiently sharpen R1-R6 output, improving its temporal resolution.

Again, the feasibility of the hypothesis was assessed by analyzing and comparing the resulting biophysical model output to real R1-R6 recordings. The model was implemented in four steps (**Box S8.1C**):

(i)   Two bright dots, 6.8^o apart, crossed a photoreceptor's receptive field (RF; $\Delta\rho = 8.1^o$) front-to-back at the saccadic speed of 205 ^{o}/s.

(ii) The ommatidium lens focused their light onto a rhabdomere tip. After 8 ms delay, the rhabdomere started to move back-to-front as it contracted photomechanically. And now, with this movement, its acceptance angle, $\Delta\rho$, also narrowed transiently, from 8.1^o to 4.0^o (**Box S8.2**). In the model, the further away the rhabdomere moved from its starting position at the focal plane, the more its receptive field narrowed (or skewed).

(iii) The resulting dynamic light input was, therefore, a convolution of the two dot intensities and the cell's receptive field, which narrowed and moved at different speeds in the same direction (front-to-back).

(iv) The light input drove the microvillar photon sampling and refractory quantum bump (QB) production, which were summed up over the whole rhabdomere to a macroscopic photoreceptor voltage output.

Importantly, the predicted photoreceptor output showed now two distinct peaks, indicating that the two dots (iv) were resolved neurally. Moreover, the simulations closely resembled the recordings to the similar stimulus (*cf.* **Box S8.1C** to **D**).

In another test (**Box S8.1D**), we estimated the same R1-R6's light input (iii) from its RF (i; through programmed look-up table operations, see **Box S6.1**) and used this to predict its output (iv; blue dotted trace). We discovered that the simulated output was indeed similar to the cell's actual recorded output to the same stimulus (black trace), with the timing of their two peaks matching closely. This close dynamic correspondence between the simulations and recordings was robust and reproducible in different tested stimulus conditions (*cf.* **Box S8.3C-D**), meaning that the given model structure likely incorporated the basic biophysical mechanisms that R1-R6 photoreceptors use in encoding moving stimuli.

Box S8.3. The hypothesis predicts R1-R6 output to double-fast moving point-objects. All the parameters of the simulations are fixed to the experimentally measured values, similar to Box S8.1. The stimulus is the same two bright dots, but this time, they cross the cell's receptive field (RF) double-fast, 409°/s (i). (A) Again, if the rhabdomere remains stationary during their flyby (corresponding to the classic theory, ii), their light fuses (iii), and the photoreceptor output cannot distinguish the dots (iv). (B) By including the light-induced back-to-front rhabdomere movement (ii) but with the same receptive field shape, the light input broadens slightly (iii), but cannot separate the two dots. Consequently, the photoreceptor output (iv) shows a slightly narrower single peak than in the previous case (A). (C) If, however, the rhabdomere contraction (away from the lens' focal point, ii) moves the receptive field and actively narrows it (from 8.8° to 4.9°), the light input from the dots is transformed into two intensity spikes (iii), which R1-R6 output separates into two peaks (iv). (D) Corresponding intracellular R1-R6 recordings show comparable dynamics to the full model c. These and other simulations and recordings, which all show good correspondence, establish that photoreceptors' receptive fields must move and narrow dynamically during light stimulation.

In summary, for the biophysical model to match up the real R1-R6 cells in resolving two bright moving dots, its receptive field must move and transiently narrow from its original size (**Text S4**: $\Delta\rho_{start}$ = 7.00 - 11.65°) to the size predicted by the optical waveguide theory ($\Delta\rho_{end}$ = 3.5 - 5.3°). Moreover, within this dynamic $\Delta\rho$ change range, the resulting model outputs become realistic and robust. Using this modeling approach, we could appropriately predict the real photoreceptors' voltage responses to the different moving dot stimuli (**Text S6**), irrespective of the tested dot speed, direction (front-to-back or back-to-front) and inter-dot distance. For example, by replacing the mean measured rhabdomere displacements with some of the larger values (**Text S7**), the simulations resolved moving dots similarly well to the recordings even at very high stimulus speeds 400-800 °/s.

From the neural coding point of view, this broad agreement between our "microsaccadic sampling"-hypothesis and the experiments makes it almost certain that photomechanical rhabdomere contractions (**Text T S7**) move and narrow R1-R6 photoreceptors' RFs to enhance visual acuity. But from the viewpoint of reducing light-adaptation, these processes seem like by-

products of a simple evasive action, which steers the rhabdomere away from pointing directly to the light source (to recover more refractory microvilli; see **Text S2**). Nevertheless, while elementary optics makes it clear why a horizontal rhabdomere motion must move a R1-R6's receptive field in the opposite way, it is harder to see what physical mechanisms could narrow it. We next consider four potential processes within the ommatidium lens system that could just do this.

What could cause the receptive field narrowing during moving light stimuli?
Four hypothetical mechanisms, together or separately, could explain the required RF narrowing (**Box S8.4**):

1. When a rhabdomere moves back-to-front (**Box S8.4A**), it moves away from the center axis, which remains fixed because the ommatidium lens system does not move (see **Text S7**, **Box S7.7**). Therefore, as the rhabdomere tip moves horizontally (**Movie S2**), the light input point-spread function (airy disc) should fall only partly upon it. This may clip or skew the rhabdomere's RF (acceptance angle, Δρ), narrowing it.

2. Besides moving rhabdomeres horizontally (**Text S7**), photomechanical photoreceptor contractions also move them 0.5-1.7 μm inwards (**Movie S1**; **Box S8.4B**). In a dark-adapted state, the rhabdomeres are elongated towards the lens with their tips possibly not being at the focal point. Hence, in this position, the rhabdomeres should collect light from broader angles (**Text S4**, from the brief test pulses: **Boxes S4.5-6**), and partially recover (re-elongate) before the next pulse comes. But during more continuous light stimulus, their contraction pulls their tips inwards, towards the possible focal point of the lens, which could narrow Δρ towards its theoretical values (3.5-5.3°).

3. Rhabdomere tips are linked by adherence junctions to the cone cells (above them) and pigment cells (at their upper corners) (*37*) (**Box S8.4C**). When the rhabdomeres contract these connections likely pull the pigment cells above, generating a dynamic aperture (**Video 3**), which moves and possibly tightens, to narrow Δρ (see **Text S7**, **Box S7.7**).

Box S8.4. How ommatidium dimensions change during light-induced photoreceptor contractions. The three images show the same transverse section of three *Drosophila* ommatidia, with their rhabdomeres appearing as six darker curvy strips. The rhabdomeres contract (cyan arrow) to light (yellow arrow, as focused by the lens), moving both (**A**) horizontally (back-to-front) and (**B**) vertically (down arrow), see **Text S7**. Because the rhabdomeres are connected to the surrounding structures by adherence junctions (*37*) (red boxes highlight the transition areas), their contraction induces (**C**) moving and possibly narrowing of the aperture (green horizontal arrow) formed by the cone and pigment cells, which are directly above the rhabdomeres (lighter blue areas). Ultimately, the curvature of the lens and focal distance might also change slightly. Image modified from (*32*).

4. In dark-adaptation, the waveguide crosstalk between neighboring rhabdomeres could broaden their receptive fields. But light-induced horizontal rhabdomere movement may eliminate the crosstalk between the neighbors, narrowing $\Delta\rho$ towards the theoretical values (3.5-5.3$^\text{o}$).

Specific predictions of our new hypothesis and their experimental validations
The microsaccadic sampling-hypothesis, as implemented by our biophysically realistic photoreceptor model (**Text S1**) with combined photomechanical rhabdomere dynamics, makes important predictions about the coding benefits of moving and narrowing R1-R6 receptive fields that can be tested experimentally.

The 1$^\text{st}$ prediction is that, for a given stimulus or saccadic velocity, R1-R6 output to a back-to-front moving bright dot should appear before the output to a front-to-back moving dot. This is because the back-to-front moving dot should enter and exit a contracting photoreceptor's front-back moving receptive field earlier; whereas the dot moving in the opposite direction should stay marginally longer inside its receptive field. **Box S8.5** compares the theoretical predictions (output simulations) of the same three models as earlier (**A**, **B** and **C**; their details are above) and corresponding exemplary intracellular recordings (**d**) for these two stimuli.

Both the full model simulations (**c**) and many recordings (**d**) indicate that this prediction is indeed what happens for a R1-R6 with a symmetrical RF (**i**). The responses (red traces) to back-to-front moving dots rise and decay faster than the responses (blue traces) to front-to-back moving dots. Similar response dynamics of another intracellular recording series from another R1-R6 are highlighted in **Text S6** (**Box S6.6**). Notice, however, that these results are explicitly true for symmetrical receptive fields. If, on the other hand, a R1-R6's RF was asymmetrical - say, profoundly skewed towards the front of the eye, then its response to the front-to-back moving dot might, in fact, rise earlier, or there could be little difference between the responses. Thus, it is the mathematical relationship between the scale of RF asymmetricity and the scale of the rhabdomere back-to-front movement, which ultimately sets whether a front-to-back or back-to-front stimulus would win. Notice also that dynamic photomechanical rhabdomere movements and eye muscle activity make RF recordings difficult to perform, and consequently, experimental inaccuracies and limitations can influence the results. Thus, some of the natural variations in the recordings may result from imprecise stimulation control. For example, imperfect positioning of a 25 light-point stimulus array - either off-center of a cell's receptive field or if not aligned perfectly parallel in respect to the eye's back-to-front axis - could bias a recording.

Box S8.5. The new hypothesis predicts and the recordings show that R1-R6 output rises and decays earlier to a back-to-front moving bright dot than to a front-to-back moving dot of equal velocity. (A) A classic model with (i) a symmetrical receptive field and (ii) an immobile rhabdomere leads to identical (iii) light inputs and virtually identical (iv) photoreceptor outputs (minute differences result from stochastic photon sampling), respectively, for the dots moving in the opposite directions. (B) A model with (ii) a moving rhabdomere generates both dynamic (iii) light input and (iv) photoreceptor output, which rise and decay earlier for the back-to-front moving dot (red traces) than the front-to-back moving dot (blue traces). (C) Our full model with rhabdomere contraction dynamics that move and narrow its receptive field. This makes the light input and photoreceptor output rise and decay faster than in the other two models, with the back-to-front waveforms leading the front-to-back counterparts. (D) The intracellular responses of a R1-R6 photoreceptor to the given two dot stimuli, as recorded *in vivo*, show similar dynamics to the full model (**c**), with its back-to-front signals (red) leading the front-to-back signals (blue).

Interestingly, for both opposing object directions, the narrowing of a R1-R6's receptive field (**Box S8.5C-D**) makes its voltage responses briefer than what would be the case without this process (**A-B**). Therefore, the resulting faster temporal photoreceptor output dynamics combat the effects of motion blur, supporting the theoretical and experimental results in **Text S6** (**Boxes S6.7** and **S6.10**).

The 2nd prediction is that, for high (saccadic) speeds, R1-R6s resolve two front-to-back moving bright dots better than when these move back-to-front (**Box S8.6**). Thus, the normal back-to-front rhabdomere movement should improve the fly eye's spatiotemporal resolution during fast forward locomotion or object motion. This is indeed what we saw in the full model simulations (**Box S8.6D**) and in some stable experimental recordings (**Box S8.6E**). Because of the back-to-front rhabdomere movement, which was inverted by the ommatidium lens, the light input (**C**) for two front-to-back moving dots were separated further apart as intensity spikes (blue trace) than that for the opposite motion (red). Consequently, the resolvability in the resulting R1-R6 output (**D**) also became greater for front-to-back moving stimuli. This of course further meant

that in comparison to the case of the immobile rhabdomere with the same narrow acceptance angle ($\Delta\rho = 4^{o}$), the neural resolvability of the back-to-front moving rhabdomere, which kept the same stimulus longer within its receptive field, would be still better.

Box S8.6. Fast rhabdomere movements are predicted to improve the resolvability of fast front-to-back moving objects the most. (**A**) Two bright dots, 6.8^{o} apart, cross a photoreceptor's receptive field (RF; $\Delta\rho_{start} = 8.1^{o}$) either in back-to-front (red, left) or front-to-back (blue, middle) at 409 ^{o}/s (i). (**B**) The new photoreceptor model translated the light-induced back-to-front rhabdomere motion into concurrent RF narrowing ($\Delta\rho_{end} = 4.0^{o}$) and front-to-back movement (as reversed by the ommatidium lens). (**C**) Consequently, the light input from the dots was transformed into two intensity spikes. These spikes were further apart in time for the front-to-back moving dots and for the opposing stimuli. (**D**) The two peaks in in the corresponding model-predicted R1-R6 output (blue) for front-to-back moving dots (highlighted by arrows) indicated that the dots were neurally detectable. In contrast, the predicted R1-R6 output for back-to-front moving dots (red) failed to separate these dots at 19^{o}C. (**E**) An example of intracellular recordings from one R1-R6 photoreceptor to the same two stimuli at 25^{o}C. This cell's voltage responses also resolved the front-to-back moving dots better than their back-to-front moving counterparts. These and other comparable simulations and recordings suggest that microsaccadic rhabdomere movements improve the neural resolution (and representation) of fast moving visual objects.

Photoreceptors resolve moving object up to high body-saccadic speeds

We next used the complete photoreceptor model (above) to estimate how well a typical R1-R6 photoreceptor can resolve two bright dots, which are less than the average acceptance angle ($\Delta\rho$) apart, moving together at increasingly fast (saccadic) velocities. This time, however, the simulations were performed at the flies' preferred temperature (*186*) of 25 ^{o}C, rather than at 19^{o}C (as in the previous data). We have shown earlier that warming fastens R1-R6s' phototransduction dynamics and refractory microvilli recovery (*5, 6, 45*). Because the resulting increase in their sample (quantum bump) rate changes improves information transfer rate (*5, 6, 45*), we expect here that their output to moving dots should also show improved resolvability. We later compare these estimates to the measured head/body-saccade speeds of freely locomoting *Drosophila* (*22, 23*).

      Box S8.7 shows the simulated light input (**c**) and photoreceptor output (**D**) to the dots (6.8^{o} apart) crossing a R1-R6 photoreceptor's RF ($\Delta\rho_{start} = 8.1^{o}$, $\Delta\rho_{end} = 4.0^{o}$) in front-to-back direction at 205 (left), 409, 600, 700, 800 and $1,000^{o}$/s (right). We found that the slower the dots moved (**A**), the better the predicted photoreceptor output distinguished them as two separate

events (peaks) in time (**D**). Remarkably, the output resolved the dots at speeds until ~600 ^{o}/s (cyan background). As a neural threshold for representing sub-RF details, this image speed is indeed very high. It means that *Drosophila* should lose little neural image detail during its saccadic body rotations during walking; the measured body rotation speed range is about 200-800 ^{o}/s (*23*).

At the higher speeds, the two response peaks fused into one. Notice that because the predicted light input (as modulated by the rhabdomere contraction) resolved the dots even at 1,000 ^{o}/s (**Box S8.7C**), the resolution limit in the photoreceptor output (**D**) resulted from its intrinsic signal integration time limit; for the given (experimentally measured) quantum bump size, latency and refractoriness distributions (*5-7*). Notice also how the maximum amplitude and half-width of the voltage responses was reduced more the faster dots crossed its receptive field. Thus, the rhabdomere then simply captured and integrated fewer photons in a given time unit.

Box S8.7. The hypothesis predicts that at the preferred temperature (25 ^{o}C) *Drosophila* R1-R6 photoreceptors can distinguish two bright dots 6.8 ^{o} apart travelling together at ~600 ^{o}/s. (A) The starting and ending receptive field size (RF;$\Delta\rho_{start}$ = 8.1^o; $\Delta\rho_{end}$ = 4.0^o) in all the simulation was the same. (**B**) in every case, the rhabdomere contracted, moving and narrowing the RF. (**C**) This translated light input into two clear sharp peaks. (**D**) Photoreceptor output separated the dots as two peaks at velocities until above 600 ^{o}/s (left, cyan background), indicating that they were neurally resolvable. At 700 ^{o}/s or higher speeds (right), the dots were not resolved as the outputs had only a single peak. Mean ± SD shown, n = 6 repeated stochastic simulations to each stimulus.

Prediction that R1-R6s encode hyperacute images in space-time

Given that R1-R6 output shows unexpectedly high acuity even at very fast saccadic velocities (**Box S8.7**), we asked how well these cells could in fact resolve slower moving point-objects. Could a normal R1-R6 encode image details, which were less than the average interommatidial angle apart? That is, could *Drosophila* actually see the world in finer resolution than their compound eyes maximum sensor (or pixel) spacing, which is the limit predicted by the classic optical theory (*2*)?

We first tested this hypothesis theoretically by using the full "microsaccadic sampling"-model (**Fig. 5**, main article). In these simulations, two bright dots were now either 1^o, 2^o, 3^o or 4^o apart. Thus the dot spacing was less than the *Drosophila* compound eye' average interommatidial angle ($\Delta\varphi \sim 4.5\text{-}5^o$). The dots were then moved across a R1-R6 photoreceptor's RF ($\Delta\rho_{start} = 8.1^o$; $\Delta\rho_{end} = 4.0^o$) at different speeds, ranging from 5 (slow gaze fixation) to 400 ^{o}/s (very fast body saccade).

Crucially, the model predicted that at suitable velocities, a typical R1-R6 photoreceptor would resolve the dots in hyperacute details (**Fig. 5**). When the dots (or the *Drosophila* eye) moved at 10 ^{o}/s, a single photoreceptor could theoretically capture image details at 1^o resolution. However, if their movement was slower (2.5 ^{o}/s), the fast photoreceptor adaptation fused the dots together, making them neurally unresolvable. On the other hand, if the dots were 2^o apart, they could be neurally resolved using 5 to 20 ^{o}/s speeds, and when 3^o apart, they could be seen over a broad speed range; from 5 to 100 ^{o}/s. Better yet, the dots that were 4^o apart could be resolved even during fast saccadic body-rotations (200-300 ^{o}/s).

Therefore, according to the "microsaccadic sampling"-hypothesis, *Drosophila* should have hyperacute vision, and through its own behavior, be able to adjust the resolution of its neural images. These results suggest that if a fly needs to see high image details it should scan the world slowly (which probably happens during normal gaze fixation). Yet the fixation speed cannot be too slow; otherwise, the photoreceptor adaptation removes the fine details. And when a fly locomotes or spins around faster, in a saccadic or bursty fashion, its photoreceptors' information capture goes up (**Figs 1-2**), while surprisingly little spatial fine details about the surrounding scene would be lost.

These theoretical predictions were broadly confirmed by intracellular recordings (**Fig. 5**; **fig. S9**), whilst flight simulator experiments verified that *Drosophila* indeed have hyperacute vision (**Fig. 6**).

Refractory sampling improves hyperacute motion vision
To quantify how refractory photon sampling contributes to the sharpening of the macroscopic responses during moving hyperacute 2-dot stimuli (brightness as in **Box S8.1**), we further compared the outputs of two different photoreceptor models for the same stimuli (**Box S8.8**). For both cases, the resulting dynamic light input – reflecting the narrowing and moving receptive field, as caused by photomechanical rhabdomere contraction - was the same, but the models' photon sampling differed. The test model had 30,000 stochastically operating refractory microvilli and the control was a comparable mock model, which converted every incoming photon into a quantum bump.

Firstly, we found that the refractory sampling consistently improved the response resolution beyond that of the control for bright hyperacute dots (inter-dot-distance <4.5^o) at slow velocities (≤ 20 ^{o}/s) (**Box S8.8A-B**). Thus, refractoriness enhances neural image resolution during slow self-motion or when high-resolution objects move slowly. Its effects were particularly well seen in the differing rising phases of the normalized light-induced current (LIC)

responses of the two models. With the slow moving stimuli, the rising responses of the refractory sampling model (black traces) always led those of the non-refractory model (red).

Additionally, in other trials, we found that when the dots were more than the average acceptance angle apart ($\Delta\rho_{start} \geq 9.5^{o}$) but moved across a R1-R6 photoreceptor's receptive field at very fast saccadic speeds ($\geq 400^{o}$/s), the refractoriness often enhanced neural image resolution beyond that of the controls (**Box S8.8C**). This observation is consistent with our previous finding that stochastic refractoriness of light-activated microvilli exerts a memory of past events in bump integration (*5*). This memory accentuates certain stimulus features relative to others so that a R1-R6 samples information from different inter-dot-distance/speed-combinations differently. Ultimately, it could well be that the real R1-R6 photoreceptors' refractory photon sampling statistics are adapted (through their visual lifestyle) to the statistics of moving high-resolution natural images. Of course, here, the used 2-dot stimuli and the models, which were isolated from the lamina network feedbacks, were too simple to fully explore such statics and the intricacies of hyperacute *Drosophila* vision.

Box S8.8. **Refractory sampling enhances neural resolution for different aspects of hyperacute images**. (**A**) Examples of simulated macroscopic light-induced current responses (normalized LICs) of two different photoreceptor models. Both models have 30,000 microvilli. In the 1st model (red traces), every photon causes a quantum bump; hence, its photon sampling has no refractoriness. In the 2nd model (black), photon sampling is refractory. Both models are stimulated with the same moving two bright dots, with their actual light inputs being first modulated by photomechanical rhabdomere contractions (following the microsaccadic sampling hypothesis). Based on the tests with different velocities and inter-dot-distances, refractoriness consistently causes a phase lead in LIC responses. (**B**) Refractoriness improves response resolution for hyperacute stimuli (inter-dot-distance <4.5^{o}) at slow velocities ($\leq 20^{o}$/s). The resolvability, D, of the recordings and simulations, was determined by Raleigh criterion (*cf.* **Fig. 3C**). (**C**) An example of how refractoriness can enhance response resolution for larger stimulus separations at high saccadic velocities. The difference from the lower peak to the trough is larger in the black trace (refractory photon sampling) than in the red trace (complete photon sampling).

Horizontal vs. vertical motion hyperacuity

The two most important biophysical factors of *Drosophila* photoreceptors, which lead to motion hyperacuity - whereupon space is encoded in time, are their sufficiently narrow receptive fields ($\Delta\rho_{end} < 5^{o}$) and refractory photon sampling (quantum bump dynamics). Therefore, theoretically,

as R1-R6s' receptive fields should narrow when an object crosses them, irrespective of its motion direction; *Drosophila* is expected to have hyperacute vision for both horizontal and vertical motion.

However, as we considered in **Box S8.5**, R1-R6 photoreceptors' neural resolvability should be the best for front-to-back moving objects. In this case, due to their back-to-front sweeping rhabdomeres, R1-R6s' receptive field can broaden slightly in horizontal direction. This dynamic may in part contribute to the curious observation that L2 monopolar cell terminals' receptive fields (in the medulla) are anisotropic, elongated in horizontal (yaw) direction and narrower in vertical (pitch) direction, as measured by calcium-imaging experiments (*33*).

Mirror symmetric contractions may improve rotatory motion detection

As we showed in **Text S7**, light increments evoke mirror symmetric back-to-front rhabdomere movements in the left and right fly eye. Interestingly, during fast saccadic body rotations, this phenomenon could surprisingly help a fly's visual orientation (**Box S8.9**). The microsaccadic sampling-hypothesis predicts that image rotation causes a phasic difference in photoreceptor outputs between the left and the right eye. Because this difference depends upon the rotation speed, it could be used for signaling changes in the fly's heading direction or to improve visual navigation.

Box S8.9. During a left saccade, signals from the left-eye lead the signals from the right-eye. Because incoming light, in both the left and right eyes, contracts their rhabdomeres in the back-to-front direction, *Drosophila* photoreceptors can transmit information about the field rotation or the fly's orientation changes to the brain. Specifically, the output from the eye towards which the fly turns phase leads, and thus neurally signals rotation direction. Here we illustrate the underlying mechanism by using a brief 205 °/s left saccade (in a world of 6.8° black-and-white gratings), as an example. (**A**) During the left saccade, the flow field facing the left eye moves back-to-front (red arrow) across its photoreceptors' receptive fields (i). The image of the moving flow field is inverted by the ommatidium lens, and so moves front-to-back (blue arrow) while being sampled by the rhabdomere (ii). Light contracts the rhabdomeres in back-to-front direction (red arrow). With the projected image

and the photon-sampler (rhabdomere) moving against each other, the light input from two bright bars (iii) becomes narrowed and accelerated. Accordingly, photoreceptor output shows two prominent peaks, in which distance to each other is compacted in time (iv). (**B**) During the left saccade, the flow field facing the right eye moves front-to-back (blue arrow) across its photoreceptors' receptive fields (i). Again, the ommatidium lens inverts the image, which now moves back-to-front (red arrow) over the rhabdomere (ii), which contracts in the same direction (red arrow). With the projected image and the rhabdomere moving together, the light input from two bright bars (iii) excites the photoreceptor longer, in which output shows two prominent peaks elongated in time (iv). (**C**) The light input to the left eye photoreceptors (red trace) rise and decay faster than the corresponding light input (blue trace) to the right eye photoreceptors. (**D**) R1-R6 output in the left eye (red trace) is faster and briefer than the photoreceptor output (blue trace) in the right eye. (**E**) Consequently, by correlating the outputs from the left and right eye photoreceptors in time, the fly brain can obtain information about the directional changes in the visual space, in respect to its head orientation.

Text S9: Microsaccadic rhabdomere movements and R1-R6 information capture

Overview
This supplement describes how photomechanical rhabdomere movements affect *Drosophila* R1-R6 photoreceptors' information capture, compares microsaccadic information sampling of dark and bright objects and provides useful background information about the experimental and theoretical results presented in **Figures 1-5** in the main paper.

In this supplement:

- We explain why and how rhabdomere movement noise influences R1-R6 output mostly at low frequencies, causing relatively little information loss.
- We test and compare how microsaccadic sampling affects encoding of bursty bright or *dark* image contrasts, using intracellular recordings.
- We further examine how well R1-R6s encode two *dark* moving point-objects (dots), and compare these recordings with those to corresponding bright moving dots.
- The results confirm that *Drosophila* R1-R6 photoreceptors resolve both bright and dark moving hyperacute patterns (< interommatidial angle, $\Delta\varphi \sim 4.5$-5^o), and can respond to bright or dark point-objects, which are less than their acceptance angles ($\Delta\rho \sim 9.5^o$) apart, even at high saccadic velocities. Thus, microsaccadic sampling hypothesis provides a robust functional explanation for *Drosophila*'s hyperacute vision, shown by the behavioral experiments in **Text S10**.
- The results support the idea that *Drosophila*'s optimal viewing strategy would involve fixating on dark features, which recover refractory microvilli, and then shifting gaze to bright features, to maximize information capture in time.

Microsaccades accentuate high-frequency resolution but generate low-frequency noise
Photomechanical rhabdomere contractions (microsaccades; **Text S7**) can maximally shift the center of a R1-R6 photoreceptor's receptive field by $\sim 5^o$, and through this self-induced light input modulation (**Fig. 4**) cause variations (noise) in its voltage output. Such "rhabdomere movement noise" is inevitable if the photoreceptor signal is classified and estimated as the average of the repeated responses, just as we did in the performance calculations (**Fig. 2** and **fig. S2**).

The condition itself bears resemblance to taking snapshots of a stationary scene from different positions and averaging these. The mean image shows an obvious smear, even if the positions were only a fraction of a photoreceptor's receptive field ("pixel") apart. However, during repeated light stimulation, the rhabdomere movements adapt rapidly (**Fig. 4E** and **fig. S3E**), with this noise affecting less the subsequent performance estimates. Thus, when quantifying the photoreceptor performance to repeated light intensity time series stimulation (*e.g.* **Fig. 2**), we removed the first 3-10 responses, in which these movements had the largest effect. In the recordings, this noise would then be rather constant across the collected responses.

Rhabdomere movement noise is missing from the simulated photoreceptor output (**figs S5-6**). Therefore, given that the stochastic photoreceptor model's transduction noise is adapted to the mean light stimulus intensity (**Text S2**), similarly to that of the recordings (**figs S3A-B**), we could isolate it as the difference between the recorded and simulated R1-R6 output (**figs S3C-D**). The analysis suggests that rhabdomere movement noise affects mostly low-frequency R1-R6

output, reducing its signal-to-noise ratio, and importantly, it effectively matches the rhabdomere jitter in high-speed video footage (**Text S7**; **fig. S3E-F**).

The rhabdomere contractions deviate them from directly facing the light source, reducing photon influx especially during bright stimulation (**Figs 4C-D**). Such evasive action, however, has surprisingly little detrimental effect on the photoreceptors' information transfer. This is because bright stimulation ($>10^6$ photons/s) contains too much light to be transduced by 30,000 microvilli into quantum bumps, and R1-R6s actively screen off excess photons to maximize information in their voltage output. In **Text S2**, we showed that R1-R6s' photomechanical adaptations (the contractions and intracellular pupil mechanism) are jointly optimized with refractory sampling (to modulate quantum efficiency (*5, 6*)) for maximal information intake at different stimulus conditions. Moreover, owing to bump adaptation and microvilli refractoriness, which accentuate light fast changes in macroscopic voltage output (*5, 6, 40*) (*e.g.* (*5*): Figs 9-10, improving high-frequency resolution), slower signals in return become compressed. Importantly, this low-frequency response range (<~10Hz), where also rhabdomere movement noise mostly resides, carries relatively little information (*5*) (about the behaviorally more relevant faster changes in the world).

Are saccades and fixations optimized to microsaccadic sampling?
Recordings and simulations (**Figs. 1-2** and **figs. S1-2** and **S4-6**) showed unequivocally that R1-R6s information capture is maximized for high-frequency saccade-like bursts with dark fixation intervals. This suggests that the optimal daytime viewing strategy would be to fixate on dark features in the visual scenes, as this recovers refractory microvilli, and then rapidly move gaze to over bright features, as this increases quantum bump (sample) rate changes and thus information capture in time. And indeed, in behavioral experiments, *Drosophila* readily fixates on and track dark objects, such as vertical bars (*158, 192, 193*). But because the fly eye photoreceptors sample a continuous panoramic view of the world, many of them - at any one time - would unavoidably face bright contrasts, which reduce their sensitivity even when their photomechanical adaptations (**Text S2**) operate maximally. We therefore also tested by intracellular R1-R6 recordings how encoding of dynamic bright or dark contrast changes may differ.

Box S9.1A shows examples of consecutive responses recorded from a R1-R6 to repeated high-frequency saccadic bright or dark contrast bursts (with 100 Hz cut-off). Expectedly, following the stochastic adaptive visual information sampling theory (**Texts S1-2**), the responses to bright (positive) contrasts after dark "fixation" periods (background, BG) were significantly larger than those to dark (negative) contrasts after bright "fixation" periods. Notice, however, that although in terms of absolute light intensity (*I*) changes (or peak-to-peak amplitude modulation) the two stimuli were equal, the negative bursts had smaller absolute contrast values ($c = \frac{\Delta I}{I}$) than the positive ones. This is because the darkening bursts reached their absolute contrast maximum (darkness, -1) only occasionally, whereas the corresponding brightening bursts reached higher absolute contrasts (>1; owing to their lower mean light intensity).

Box S9.1. Optimal saccadic viewing strategy for maximizing R1-R6 photoreceptors' information capture requires fixating on darker image features. (**A**) Intracellular responses of a R1-R6 to repeated saccadic bright (left) and dark (right) bursts. (**B**) Output to bright saccadic bursts was always more vigorous when mixed with dark fixation periods (BG, above) than with bright periods (middle), irrespective of the stimulus frequency distribution. When the same R1-R6s were fixated (light-adapted) on a bright background (below), they still responded well to dark saccadic bursts (of the inverse waveforms), but the amplitude range of these responses was less than to bright bursts (above, dark BG). (**C**) Corresponding R1-R6 output signal-to-noise ratios were the highest for the bright-saccadic-burst/dark-fixation-period stimuli (black traces). Signal-to-noise ratios were lower but alike for the brighter saccades with bright fixation periods (grey) and the dark saccades with bright fixation periods (light grey). (**D**) Information transfer rates confirmed the global maximum for 100 Hz bright saccade-like bursts with dark fixations (black bars; *cf.* **Fig. 2C**). Whilst bright fixation periods reduced information (with more microvilli becoming refractory), the bit rates for bright (grey) and dark (light grey) saccadic modulation, of equal but opposite contrasts, were similar.

To counter this bias, we used three sets of stimuli to examine individual R1-R6s' response dynamics to both positive and negative bursts of equal or different contrast distributions (**Box S9.1B**). First (top row), the photoreceptors were stimulated with positive contrast bursts (peak-to-peak modulation = 1 intensity units), which contained high-frequency saccade-like events on a dark background, having different cut-off frequencies (from 20 to 500 Hz, as in **Fig. 1**). In the second set (middle), the same stimuli were superimposed on a bright background (1 unit). Finally (bottom), the stimulus modulation was inverted (to negative contrast bursts) and superimposed on the same bright background (1 unit). Thus, for the 2nd and 3rd sets, the stimulus contrasts were equal but opposite.

We found (again in tone with the theory) that whilst the responses were always the largest to positive contrast bursts on a dark background, the corresponding responses to the positive and negative contrast bursts on bright background, although smaller, were about the same size (**Box S9.1B**). Since we further know that the larger responses contain more quantum bumps (**Fig. 2**), with the average bumps light-adapting to about the same size (*e.g.* **fig. S3**), the responses' signal-to-noise ratios (**Box S9.1C**) and information transfer rates (**Box S9.1D**) were predictable. R1-R6s' signaling performance was the greatest to the larger positive contrast bursts on a dark background (black), and more than halved to the smaller corresponding positive (grey) and negative (light grey) contrast bursts over the test bandwidths. Notably, the responses to the opposite but equal positive and negative contrast bursts carried effectively equal information contents, underscoring the importance of contrast invariance at the primary visual encoding stage (*6, 40, 194*).

Therefore, given the fast speed of adaptation (microvilli refractoriness and dynamic quantum bump size modulation) and its photomechanical counterbalancing (**Text S2** and **Text S7**), the sensitivity of neighboring photoreceptors across the eyes can differ greatly at any one moment, depending upon whether they face dark or bright contrasts. This realization also implies that when a fly moves its gaze in saccades, the dark and bright spatial contrast differences in the world should be automatically translated into large temporal contrast changes between the neighboring retinotopic image pixels (neuro-ommatidia). Enhancement of local differences and similarities in neural images by spatiotemporal synaptic (*30, 33*) and gap-junctional (*29*) co-processing (including network adaptation (*71, 123*)) across the first optic ganglia, the lamina and medulla, should further improve object detection and fly vision.

Microsaccadic sampling of bright or dark moving point-objects

Natural scenes are rich with dark features: shadows, object boundaries, surfaces of lesser reflectance etc., which have shaped visual circuit functions, perception and behaviors (*5, 36, 59, 195-197*). Consequently, a fly's self-motion generates both dark and bright moving features travelling across its eyes. We have already shown that R1-R6s can resolve fast-moving and hyperacute bright dots (**Figs. 3-5** and **fig. S9**). But how well can these cells resolve dark moving point-objects?

We studied this question with the 25 light-point stimulus array (explained in **Text S4** and **Text S6**). As before, the stimulus array was first carefully placed at the studied R1-R6's receptive field center, but this time, all the light-points were switched on, and we generated (by software Biosyst) two travelling dark points of specified speeds and interdistances. Because, during these experiments, the cells were light-adapted (depolarized) by the lit stimulus array, the two dark dots moving on it evoked hyperpolarizing responses.

Box S9.2A shows R1-R6s' characteristic responses to two dark (black traces) and bright (red) dots of specific speeds (102, 205 and 409 ^{o}/s) and interdistances (3.4, 5.1 and 6.8^{o} apart), recorded from the same cells. In all these cases, the hyperpolarizing responses resolved the two dots, generating two troughs separated by a peak, but these responses were considerably smaller than those to the corresponding bright dots. However, when normalized, the photoreceptors' relative neural resolvability of the dark dots matched that of the bright dots (**Box S9.2B**).

Thus, in concordance with the behavioral experiments (**Fig. 6**), these and other intracellular recordings established that *Drosophila* R1-R6 photoreceptors see both bright and dark moving hyperacute patterns (< interommatidial angle, $\Delta\varphi \sim 4.5$-5^{o}), and can resolve point-objects, which are less than their acceptance angles ($\Delta\rho \sim 9.5^{o}$) apart, even at high saccadic velocities.

Box S9.2. Comparing R1-R6s' characteristic intracellular responses to hyperacute or saccadic bright or dark moving dots. (**A**) Mean (thick traces) and individual responses (thin) of three individual R1-R6s to two bright (red, above) or dark (black, below) dots (of different inter-dot-distances and speeds), crossing the cells' receptive fields in front-to-back direction. Responses to the hyperacute dots (3.4^{o} inter-dot-distance) are shown left; responses to the saccadic speed stimuli in the middle (205 ^{o}/s) and right (409 ^{o}/s). In all cases, these outputs resolved the two dots, but predictably the responses were always smaller to the dark dots at the bright background than to the bright dots at the dark background (*cf.* **Fig. 1**; as more microvilli should remain refractory when adapting to bright background). (**B**) The normalized and sign-inverted R1-R6 outputs to dark dots were similar to their normalized outputs to bright dots, with both showing equally good Raleigh-resolvability (*cf.* **Fig. 3C**). Naturally, as the *Drosophila* eye samples light information from each point in space by 8 photoreceptors (due to neural superposition) and balances this estimate with those of the neighboring lamina cartridges, its perception/resolvability of hyperacute and saccadic image motion is improved further.

Microsaccadic sampling hypothesis and efficient coding

Our results indicate that R1-R6 photoreceptors' information transfer adapts to the context of stimulus statistics; with refractory microvilli, fast quantum bump adaptation (*6, 7, 15*) and photomechanical microsaccades maximizing encoding of phasic information from high-frequency saccadic bursts. Remarkably, the extraction of phasic stimulus features, which characterize object boundaries and line elements in visual scenes, already starts during sampling and integration of visual information in the microvilli, at the first visual processing stage. The darker periods in stimuli relieve the effects of microvilli refractoriness, enabling greater sensitivity: more and slightly larger samples (quantum bumps) can be generated transiently to the next light change (**Fig. 1, figs S1** and **S5**). This increases the signal-to-noise ratio of the integrated macroscopic response, especially in its phasic (fast rising/decaying) components (**Text S3**, **Box S3.1**). However, unlike later information processing in the network, during which presynaptic inputs are often translated to postsynaptic spike-bursts of high sparseness for specific features, the earliest neural code of photoreceptors must consider all stimulus features together. It adapts to allocate information in high-contrast bursts into continuous Gaussian broadband voltage signals (**Figs 2A-B**, **figs S2A-B** and **S6A-B**), utilizing the output range optimally.

In the viewpoint of efficient coding, stochastic refractory sampling, fast quantum bump adaptation and photomechanical contractions benefit vision in three important ways:

- They exchange redundant information in mean voltage to more useful information in relative modulation, enabling photoreceptors to encode reliable estimates of the world within their limited output ranges, despite strongly and quickly changing intensities.
- They lower the metabolic cost of information with fewer bumps integrating a lower membrane potential, consuming less ATP (*5*).
- And when linked to bursty saccadic head/body movements, they increase high-frequency information capture from the world and reduce motion blur.

These results further imply that saccadic behaviors enable the fly eye to convey to the fly brain a far more efficient and accurate neural image of the variable world than what was believed before. Thus, saccades not only contribute to gaze-stabilization (*17, 198*), which historically is

considered to be their major function (*31*), but they also "burstify" light input for efficient and accurate sampling.

Text S10: *Drosophila* behavior confirms hyperacute vision

Overview

This supplement describes the used optomotor behavioral paradigm in a classic *Drosophila* flight simulator system to study visual resolution, and provides important background information about the experimental and theoretical results presented in **Figure 6** in the main paper.

In this supplement:
- We test in open-loop conditions whether wild-type *Drosophila* can generate yaw-torque (optomotor responses) to hyperacute vertical black-and-white stripe-scenes that rotate clockwise and counterclockwise at 45 ^{o}/s.
- Our results clearly indicate that wild-type *Drosophila* have hyperacute vision (seeing finer image details than the average interommatidial angle, $\Delta\varphi \sim 4.5$, of their compound eyes), with the measured behavioral responses closely following the prediction of our new "microsaccadic sampling"-hypothesis (see **Text S8**).

Testing *Drosophila* visual behavior in flight simulator system

Wild-type "Berlin" *Drosophila* were raised on standard food medium at 25°C and 60% relative humidity with a 12-h light and 12-h dark cycle, with light on at 8 a.m. In the experiments, we used 2- to 3-day-old female flies. Under cold-anesthesia (lasting < 3 min), a small copper-wire hook was fixed with a droplet of UV-light sensitive glue (Loctite) between each fly's head and thorax. After preparation, flies were left to familiarize themselves with their hooks overnight in single vials, which provided them water and sucrose.

A custom-built, computer controlled flight simulator system (*29*) was used to study *Drosophila*'s optomotor behavior. A tethered *Drosophila* was connected to the torque-meter (*50*) by a small clamp holding the copper-wire hook, which fixed the fly's head in a rigid position and orientation, but allowed stationary flight (*38, 199*). The torque meter transduced yaw torque into electrical voltage.

A fly, tethered from the torque meter, was lowered by a mechanical micromanipulator in the center of a white featureless plastic hollow cylinder (a diffuser). Inside it, we placed high-resolution visual patterns (bars, stripe patterns, etc.), which were laser-printed on a transparent film, forming a 360^o panorama around the fly's long axis. The panorama could be rotated around its vertical axis by a servomotor. Outside, the diffuse cylinder faced a surrounding ring-shaped light-tube (special full-band: 350-900 nm) that provided uniform illumination on the panorama. The light intensity during the panoramic motion stimulation, although bright, was always less (0.5-1.5 log-intensity units) than the direct stimuli used in the intracellular recordings (*cf.* **Fig. 1**).

Open-loop experiments

Inside the flight simulator, a flying fly saw a continuous (360^o) stripe-scene (black-and-white bars) of predetermined spectral and spatial resolution, which was free of motion artefacts, flashing or aliasing. After one second of viewing the still scene, it was spun to right (clockwise) by a linear stepping motor for two seconds, stopped for two seconds, before rotating to left (counterclockwise) for two seconds, and stopped again for a second. This eight-second stimulus was repeated 10-25 times and each trial, together with the fly's coincident yaw torque responses, was sampled at 1 kHz and stored in a PC's hard-drive for later analysis, using custom-written software (Biosyst) (*7*). Presumably to stabilize gaze, flies tend to follow the scene rotations,

generating yaw torque responses (optomotor responses to right or left), the strength of which is believed to reflect the strength of their motion perception (*29, 38, 199*). The fixed stimulus parameters for moving stripe scenes, as shown in the figures, were: azimuth ±360°; elevation ±45°; velocity, 45°/s; contrast, 1.0, as seen by the fly. **Figure 6A** show the averages (*n* = 9 flies) of the mean optomotor responses (*n* = 22-35 trials for each fly) for each tested fly population (**fig. S10**).

　　We tested optomotor responses of wild-type flies to moving black-and-white stripe-scenes (spectral full-width: 380-900 nm) of three different spatial resolutions (wavelength: 1.16°, 2.88° and 14.40°), as shown in **Figure 6**. To verify that air flow, or some hidden features in the stimulus panorama, was not affecting optomotor responses, we used the white diffuser cylinder alone, which showed no clear contrast to human eye, as the control stimulus (**fig. S10**). These control field rotation experiments were repeated using the same flies. We found by that the white control stimulus did not evoke torque responses.

Quantifying optomotor behavior. The optomotor responses of individual flies to the same repeated field rotations vary in strength and repeatability (**fig. S10A**), but their visual performance to different spatial resolution stripe scenes is clearly different. These differences can be quantified by measuring the mean torque response of a single fly to stimulus repetitions and by averaging the mean responses of the many flies of the same stripe scene resolution (**fig. S10B**; here 9 flies). This reduces noise and non-systematic (arbitrary) trends of single experiments, revealing the underlying response strength and optomotor behavior characteristics. These population responses are shown in **Figure 6** for a straightforward comparison.

　　In open-loop experiments, a *Drosophila*'s torque response returns gradually to baseline after the optomotor stimulus stops, but this can take many seconds (varying with individual flies). Accordingly, in our experiments, which contain only brief 2 second-long inter-stimulus-intervals, the torque responses typically show recover only fractionally (10-70%) during these still periods toward the baseline. Therefore, for comparing the optomotor behavior different stripe scene resolutions, we used the maximum range (or peak-to-peak) of the torque response, evoked by the combined leftward and rightward field rotation stimulus. The maximum range and variability in the torque responses to the same optomotor stimulus are shown with controls in **figures S10C** and **10D**, respectively.

　　Markedly, the optomotor responses to hyperacute stripe-scenes were not caused by aliasing. This is because perceptual aliasing (such as the wagon-wheel effect or Moiré patterns), if induced by the rotating hyperacute scenes, would have been perceived as slowed down image rotation, eventually reversing to the opposite direction (the reverse rotation effect). And thus, if the tested flies had seen such motion patterns, they would have consequently followed them slower and rotated against the real scene rotation direction. Such optomotor behavior was never observed in our experiments.

Why did the previous behavioral studies not find hyperacute vision?
In 1976, Buhner probed *Drosophila*'s visual acuity by stimulating the upper frontal part in one of its eyes with small local moving grating patterns (covering about 50 ommatidia) while a fly walked on track-ball (*200*). Notably, the aim of his study was not to find the finest resolution what a *Drosophila* can resolve but instead to deduce the likely columnar organization of its directionally sensitive elementary motion detectors from a fixed fly's tendency to follow moving stimulus patterns. Thus, this was also an open-loop paradigm, but the used microscope-mediated

local grating stimulation was very different from the global hyperacute panoramic visual scenes of our study. Specifically, we note that in Buchner's study:

- Visual acuity was not tested below the interommatidial angle ($\Delta\varphi \sim 4.5$); with the overall results deduced by eliminating the presumed boundary elements and contrast attenuation from the data.
- The used mean stimulus light intensity (luminance; 16 cd/m^2) was low. Therefore, the resulting image grating at the level of individual photoreceptors would have been dim and spatiotemporal signal-to-noise ratio of light input and photoreceptor output low. Based on our intracellular data (*6, 7*), this dim light intensity would have made it practically impossible for R1-R6 photoreceptors to resolve very fine (or hyperacute) visual patterns.
- The sensitivity and the time resolution of the used trackball system (*200*) seem significantly less than in our bespoke torque meter (*201*), requiring extensive data averaging. This would have made it far more difficult for the trackball system to resolve the weaker (small amplitude) behavioral responses to fine spatial contrast changes (**Fig. 6A**).

More recently, because of the historical belief that interommatidial angle limits a fly's visual acuity, many experimentalists have started using coarse LED-matrixes, typically with 4.5-5$^\text{o}$ maximum resolution, to probe visual learning and optomotor responses. As our study here shows, these kinds of visual stimuli are very different from the panoramic high-resolution printed scenes with thin continuous lines and symbols and thus are expected evoke quite different neural responses.

Supplementary Figures S1-S10

Fig. S1. R1-R6 output varies more cell-to-cell than trial-to-trial (*cf.* Fig. 1) but show consistent stimulus-dependent dynamics over the whole encoding range. (**A**) The mean voltage response and SD of 15 R1-R6 cells to the same repeated 20 Hz bandwidth bursts. Photoreceptor output adapts within ~2 s to the stimulation. (**B**) Population means (thick) and 15 mean voltage responses of individual photoreceptors (thin traces) to 20 different stimuli; each with specific bandwidth (columns: from 20 Hz, red to 500 Hz, blue) and mean contrast (rows). Stimulation changes from Gaussian white-noise (GWN; bottom) to bursts (top) with the light background: from BG0 (dark) to BG1.5 (very bright). Left top: the traces from (**A**). The yellow box indicates the responses with the highest entropy and information content. Vertical dotted rectangle (orange square) and horizontal rectangle (black circle): responses for contrast and bandwidth analyses in **fig. S2A**. All recordings were done at 25 °C. Compare this data to **Figure 1**.

Fig. S2. Signaling performance vary cell-to-cell but adapts methodically to given stimulus statistics. (**A**) Response signal-to-noise ratios (SNR) to 20 (red), 100 (yellow) and 500 Hz (blue) bandwidth (left) saccadic bursts, and to 50 Hz bandwidth stimuli of different contrasts (right); color scheme as in **Fig. 1**. SNR increases with contrast (right), reaching (in some cells) ~6,000 maximum for 20 Hz bursts (left, red). All R1-R6s showed the broadest frequency range for 100 Hz bursts (yellow). (**B**) Highly skewed bursts drove mostly Gaussian responses (exception: 20 Hz, red), with 100 Hz bursts evoking the broadest amplitude range (yellow). (**C**) Information transfer of all cells peaked for 100 Hz stimuli, irrespective of the tested contrast (or BG; left), having global maxima (infomax) between 600-850 bits/s (yellow box). (**D**) Mean encoding efficiency (R_{output}/R_{input}) reached >100% for 20 Hz bursts, with its extra information coming from the neighboring cells. For determining R_{input} see **fig. S5**. Encoding efficiency fell with increasing stimulus bandwidth, but less with contrast. Note: encoding efficiency for bursts (η^{burst}; black trace, left) was lower than for GWNs (η^{GWN}; grey and light grey traces). Because photomechanical adaptations let optimally 8-times brighter intensity modulation through for

high-contrast bursts (8 x 10^5 photons/s) than for GWN (1 x 10^5 photons/s), their higher input information is higher; $R_{input}^{burst} \gg R_{input}^{GWN}$. Thus, whilst $R_{output}^{burst} > R_{output}^{GWN}$, $\eta^{burst} < \eta^{GWN}$ (see **Supplementary Text 2**).

Fig. S3. Light-adapted R1-R6 noise is similar for all the test stimuli, with its high-frequencies reflecting the mean quantum bump shape and its low-frequencies the rhabdomere jitter. (**A**) Photoreceptor noise (from **Fig. 1**) remained largely constant for all the test stimuli; extracted from the responses (output) to stimulus repetition (input); see Methods. (**B**) Corresponding photoreceptor noise of model (*5, 6*) simulations was also broadly constant but lacked the recordings' low-frequency noise (**A**). (**C**) The mean simulated noise power ascends with membrane impedance, which here was larger than that in the recordings (**A**). Yet, the high-frequency parts of the real and simulated noise (>60 Hz), indicating the corresponding average light-adapted quantum bump waveform (*5-7, 24, 45*), sloped similarly. (**D**) Overlaying these exposed the low-frequency noise difference (<60 Hz). Our results (**Fig. 4**) predicted that this difference was a by-product of photomechanical rhabdomere and eye muscle movements, which the simulations lacked. (**E**) Mean rhabdomere movement responses (± SD, grey) in five different flies to the same repeated 20 Hz high-contrast bursts. These were smaller than those to 1 s flashing (**Fig. 4E, Supplementary Text 7**). (**F**) Average variability of the recording series (mean – individual response) shown as rhabdomere movement noise power spectra. (**G**) Mean rhabdomere movement noise (black trace; from **F**) matched its prediction (grey; from **D**).

Therefore, the recordings' extra noise resulted from variable rhabdomere contractions; jittering light input to R1-R6s. Crucially, this noise is minute; for 20 Hz saccadic bursts, ~1/6,000 of R1-R6 signal power (**Fig 2A**).

Fig. S4. Strong responses to naturalistic stimulation (NS) carry only about half the information of the strongest responses to bursts. (**A**) The mean (signal; purple) and voltage responses (pink) of a R1-R6 photoreceptor to naturalistic light intensity time series. (**B**) At the light source, NS (purple), which is dominated by low-frequency transitions between darker and brighter events, had higher power than GWN (grey) or bursty high-contrast stimuli (yellow), but its mean contrast (0.58) is between the other two. (**C**) Signal-to-noise ratio of responses to NS has a similar low-frequency maximum to responses to bursty 100 Hz stimuli, but lower values at high-frequencies, similar to GWN-driven responses (grey). Both of these signaling performance estimates are from the same R1-R6 (a). (**D**) Information transfer rate in photoreceptor output directly depends upon the mean stimulus contrast. Photoreceptors encode more information during naturalistic stimulation than during GWN stimulation (*cf.* (*5*)). But encoding can further double during high-contrast bursts, which utilize better the refractory sampling dynamics of 30,000 microvilli, generating the largest sampling rate changes (*cf.* (*5*)). Significance by two-tailed t-test. For more explanation, see **Supplementary Text 3**.

Fig. S5. Model output is realistic over the whole encoding range. (**A-B**) Simulated responses of a stochastic *Drosophila* photoreceptor model to the tested light stimuli show similar response dynamics to the corresponding real recordings (*cf*. **Fig. 1** and **fig. S1**). The model has 30,000 microvilli (sampling units) that convert absorbed photons to quantum bumps (samples). Simulations at each background (BG) were set for the mean light level (effective photons/s) that generated responses with the maximum information transfer (**Supplementary Text 2**). These effective light levels should correspond to the optimal photomechanical screening throughput (by intracellular pupil mechanism and photomechanical rhabdomere contractions), which minimize saturation effects (refractory microvilli) on a *Drosophila* photoreceptor; see **figs S2** and **S6**. Notice that the model had no free parameters - it was the same in all simulations and had not been fitted to data. Thus, these macroscopic voltage responses emerged naturally as a by-product of refractory information sampling by 30,000 microvilli. Yellow box: maximum information responses. Vertical dotted rectangle (orange square) and horizontal rectangle (black circle): responses for contrast and bandwidth analyses in **fig. S6A**.

Fig. S6. Model encodes light information realistically. Encoding capacity of a stochastic photoreceptor model peaks to high-contrast bursts with 100 Hz cut-off, much resembling that of the real recordings (*cf*. **Fig. 2**). (**A**) Inserts show simulations for the bursty input patterns of similar high contrast values (left) that drove its responses (outputs) with maximum information transfer rates. Output signal-to-noise ratios peaked for 20 Hz bursts (red), but was the broadest for 100 Hz bursts (yellow). Signal-to-noise ratio rose with stimulus contrast (right). (**B**) The corresponding probability density functions show that 100 Hz bursts evoked responses with the broadest Gaussian amplitude distribution (yellow). Only responses to low-frequency bursts (20-50 Hz) deviated from Gaussian (skewed). (**C**) Information transfer of the model output reached its global maximum (infomax) of 632.7 ± 19.8 bits/s (yellow) for 100 Hz (left) bursts (right). Corresponding information transfer for Gaussian white-noise stimuli was significantly lower. (**D**) Encoding efficiency peaked for low-frequency stimuli (left), decaying gradually with increasing contrast. For details see **Supplementary Text 2**.

Fig. S7. Each R1-R6 has a different diameter rhabdomere and network connections, and thus should extract different amounts of information from the same stimulus. (**A**) Electron micrograph of an ommatidium, showing R1-R7 rhabdomeres with characteristic cross-sectional area differences. (**B**) R1 and R6 rhabdomeres are always the largest and R4 the smallest (statistics: **Supplementary Text 5**). (**C**) R6 can receive ~200 bits/s of network information through axonal gap-junctions from R7/R8 (*29*) in the lamina about local light changes - due to their neural superposition. Gap-junctions between R1-R6 axons and synapses (*28, 30*) in the lamina redistribute information (**Supplementary Text 2**). (**D**) R1-R6s' response waveforms and frequency range varied cell-to-cell; as evidenced by the recording system's low noise and the cells' high signal-to-noise ratios (~1,000). Here, Cell #17 encoded 100 Hz bursts reliably until ~140 Hz, but Cell #11 only until ~114 Hz. (**E**) Maximum information (for 100 Hz bursts) of 18 R1-R6s, grouped in ascending order and used for typifying the cells. Because R6s' rhabdomeres are large (**B**) and their axons communicate with R7/R8 (**C**), the cells with the distinctive highest infomax were likely this type (blue). Conversely, R3, R4 and R2 rhabdomeres are smaller and their axons furthest away from R7/R8, and thus they should have lower infomaxes. Notably, our photoreceptor model (*6*) (grey), which lacked network information, had a similar infomax. The mean infomax of the recordings was 73 bits/s higher than the simulation infomax.

Fig. S8. Microsaccadic sampling hypothesis predicts realistic voltage output to two dots crossing a R1-R6's receptive field in saccadic speeds. (**A**) Receptive fields of four R1-R6 photoreceptors, measured with 25 light point stimulator (see **Supplementary Text 4**). (**B**) Microsaccadic sampling hypothesis predicts that because the rhabdomeres move photomechanically, the photoreceptors' receptive fields move in the opposite direction and narrow transiently (acceptance angles, $\Delta\rho$, change from 8.2-9.5^o to 3.5-4.5^o). (**C**) The resulting light input for each tested photoreceptor was predicted from its measured receptive field (**A**) by the microsaccadic sampling hypothesis. (**D-E**) These inputs then drove our biophysically realistic R1-R6 model, predicting the photoreceptor voltage output, which was compared to the corresponding real recordings. The simulated R1-R6 output closely resembled the recorded R1-R6 output of the same cells to saccadic two dot stimuli.

Fig. S9. Encoding space in time - intracellular R1-R6 recordings to two bright dots crossing the receptive field show how their responses convey hyperacute spatial information in time. 25 light-point-array positioned at a R1-R6's receptive field center, generating two bright front-to-back moving dots. (**A**) Characteristic responses of a R1-R6 at 25 °C to the dots, 1.7, 3.4 or 6.8° apart, travelling 102 °/s in front-to-back direction. (**B**) Individual outputs resolved the dots, which were less than the interommatidial angle ($\Delta\varphi = 4.5°$) apart (yellow box); resolvability given by the Raleigh criterion (**Fig. 3C**). Microsaccadic sampling model (**Fig. 5**) predicted a comparable resolvability threshold (dotted line). (**C**) At lower dot velocities (20-50 °/s), corresponding to normal gaze-fixation speeds in close-loop flight simulator experiments (**Supplementary Text 10**), each R1-R6 responded to the tested hyperacute dot separations (1.7° and 3.4°) even stronger. Notice the small staircase-like steps in voltage responses. These represent light from 25 individual light-guide-ends being turned on/off in sequence to generate the moving dots, crossing the receptive field slowly (see **Supplementary Text 6**). (**D**) At 20 °/s velocity, neural resolvability to the tested hyperacute dot separations was between 10-20%. (**E**) R1-R6 output to the two dots, having the same separations as above, but now moving at fast saccadic velocity (409 °/s). (**F**) Although, at such a high speed, R1-R6 output could not resolve the hyperacute dot separations (1.7° and 3.4°) consistently, the dots were nevertheless clearly detected when at 5.1° apart (cyan bar), which is about $\Delta\varphi$.

Fig. S10. Optomotor behavior in a flight simulator system demonstrates that *Drosophila* see hyperacute visual patterns. (**A**) 22-35 individual optomotor responses (thin grey traces) and their mean (think trace) of the same tethered fly to 45^{o}/s clockwise and counterclockwise panoramic field rotations (light grey sections), having a full wavelength of 1.16^{o} (left, black) or 14.40^{o} (center, blue), or no-bar (right, orange) stimuli. (**B**) Every tested fly (n = 9) responded to hyperacute ($<\Delta\varphi \sim 4.5^{o}$) bar stimuli ($1,16^{o}$ and 2.88^{o}, red; having 0.58^{o} and 1.44^{o} inter-bar-distances, respectively) consistently. (**C**) Their maximum (peak-to-peak) responses to the hyperacute stimuli and the no-bar and coarse-bar controls. (**D**) The relative response strength to the hyperacute stimuli varied between 30-80% of the maximum responses to the coarse-bar stimulus.

Movie S1: *Drosophila* R1-R8 photoreceptors in dissociated ommatidia contract photomechanically to light flashes.

Wild-type and *trp/trpl*-mutant R1-R8 photoreceptors contract photomechanically to light flashes. The panels show: top, a sideview of *ex vivo* wild-type R1-R8 photoreceptors of a single dissociated ommatidium contracting to 1 ms bright light flash; bottom left, R1-R8 of a *trp/trpl* null-mutant, which express normal phototransduction reactants but lack completely their light-gated ion channels, contracting to a similar flash. Notably, *trp/trpl* photoreceptors cannot generate electrical responses to light, with their eyes showing no ERG signal (**Text S7**). Nonetheless, *trp/trpl*-mutant photoreceptors contract photomechanically (but require ~5 min dark-adaptation between flashes to restore their contractility). These observations are consistent with the hypothesis of the light-induced phosphatidylinositol 4,5-bisphosphate (PIP$_2$) cleaving from the microvillar photoreceptor plasma membrane causing the rhabdomere contractions (*8*). Video playback slowed down and down-sampled to reveal the contractions, which otherwise would be too fast to see with a naked eye. Each video clip is repeated three times with a running timer giving the time course of the contractions.

Movie S2: *Drosophila* R1-R8 photoreceptors contract photomechanically to light flashes *in vivo*, moving back-to-front inside each observed ommatidium.

We utilised the optical cornea-neutralisation technique with antidromic deep-red (740 or 785 nm peak) illumination to observe deep pseudopupils (photoreceptor rhabdomeres that align with the observer's viewing axis) in the *Drosophila* eye. High-speed video captures fast rhabdomere movements to bright orthodromic blue-green flashes (470 + 535 nm peaks). The panels show: left, R1-R7 photoreceptor rhabdomere tips moving rapidly back-to-front and returning slower to each 10 ms flash, delivered repeatedly every second; right, the cross-correlated horizontal (blue) and vertical (red) components as the time series of this movement. Grey vertical lines indicate each flash. The rhabdomere movement is caused by the photomechanical photoreceptor contractions (not by muscle activity). These *in vivo* movements are large, here 1.7 μm from dark-adapted rest-state; causing up to 5 degree transient shift in the R1-R6 photoreceptors receptive fields (**Text S7**). Note average diameter of R1-R6 rhabdomeres is 1.7 μm (**Text S5**). The high-speed video rate was 500 frames/s. Video playback slowed down and down-sampled to reveal the contractions, which otherwise would be too fast to see with a naked eye.

Movie S3: While photoreceptors contract, the ommatidium lens above is virtually immobile but a cone-cell aperture, connected to the rhabdomere tips by adherens junctions, moves half as much as the rhabdomeres.

We used a z-axis micromanipulator to shift and reposition *Drosophila* in piezo-steps vertically underneath the microscope. This allowed the focussed image, as projected on the camera, to scan through each studied ommatidium, providing exact depth readings in μm. We then recorded any structural movements inside the ommatidia to light flashes at different depths; from the corneal lens down to the narrow base, where the cone and pigment cells form an intersection between the crystalline cone and the rhabdomere. The left panels show: up, ommatidium lens; middle, basal cone/pigment cell layer; down, R1-R7 photoreceptor rhabdomeres tips during and after flash stimulation. The right panels show the cross-correlation time series of these high-speed videos: up, the corneal lens and the upper ommatidium structures were essentially immobile), and normally remained so throughout the recordings; Middle, cone cells that connect to the rhabdomere tips with adherens junctions (*37*) showed clear light-induced movements; down, R1-

R7 rhabdomeres moved half as much as the cone cells above. The high-speed video rate was 500 frames/s. Video playback slowed down and down sampled to reveal the contractions, which otherwise would be too fast to see with a naked eye.

References

1. H. B. Barlow, The Size of Ommatidia in Apposition Eyes. *J Exp Biol* **29**, 667 (1952).
2. M. F. Land, Visual acuity in insects. *Annu Rev Entomol* **42**, 147 (1997).
3. M. Juusola, A. S. French, Visual acuity for moving objects in first- and second-order neurons of the fly compound eye. *J Neurophysiol* **77**, 1487 (Mar, 1997).
4. M. V. Srinivasan, G. D. Bernard, Effect of Motion on Visual-Acuity of Compound Eye - Theoretical-Analysis. *Vision Res* **15**, 515 (1975).
5. Z. Song, M. Juusola, Refractory sampling links efficiency and costs of sensory encoding to stimulus statistics. *J Neurosci* **34**, 7216 (May 21, 2014).
6. Z. Song *et al.*, Stochastic, Adaptive Sampling of Information by Microvilli in Fly Photoreceptors. *Curr Biol* **22**, 1371 (Aug 7, 2012).
7. M. Juusola, R. C. Hardie, Light adaptation in *Drosophila* photoreceptors: I. Response dynamics and signaling efficiency at 25 degrees C. *J Gen Physiol* **117**, 3 (Jan, 2001).
8. R. C. Hardie, K. Franze, Photomechanical Responses in *Drosophila* Photoreceptors. *Science* **338**, 260 (Oct 12, 2012).
9. R. W. Ditchburn, B. L. Ginsborg, Vision with a Stabilized Retinal Image. *Nature* **170**, 36 (1952).
10. L. A. Riggs, F. Ratliff, The Effects of Counteracting the Normal Movements of the Eye. *J Opt Soc Am* **42**, 872 (1952).
11. E. Ahissar, A. Arieli, Figuring space by time. *Neuron* **32**, 185 (Oct 25, 2001).
12. S. Viollet, Vibrating makes for better seeing: from the fly's micro-eye movements to hyperacute visual sensors. *Frontiers in Bioengineering and Biotechnology* **2**, 1 (2014).
13. K. Donner, S. Hemilä, Modelling the effect of microsaccades on retinal responses to stationary contrast patterns. *Vision Res* **47**, 1166 (Apr, 2007).
14. R. Juston, L. Kerhuel, N. Franceschini, S. Viollet, Hyperacute edge and bar detection in a bioinspired optical position sensing device. *IEEE/ASME Transactions on Mechatronics* **19**, 1025 (2014).
15. S. R. Henderson, H. Reuss, R. C. Hardie, Single photon responses in *Drosophila* photoreceptors and their regulation by Ca2+. *J Physiol-London* **524**, 179 (Apr 1, 2000).
16. G. Blaj, J. H. van Hateren, Saccadic head and thorax movements in freely walking blowflies. *J Comp Physiol A* **190**, 861 (Nov, 2004).
17. M. F. Land, Head Movement of Flies during Visually Guided Flight. *Nature* **243**, 299 (1973).
18. C. Schilstra, J. H. van Hateren, Stabilizing gaze in flying blowflies. *Nature* **395**, 654 (Oct 15, 1998).
19. N. Franceschini, R. Chagneux, paper presented at the Gottingen Neurobiology Report, Stuttgart, 1997.
20. R. Hengstenberg, Eye Muscle System of Housefly Musca-Domestica .1. Analysis of Clock Spikes and Their Source. *Kybernetik* **9**, 56 (1971).
21. S. Martinez-Conde, J. Otero-Millan, S. L. Macknik, The impact of microsaccades on vision: towards a unified theory of saccadic function. *Nat Rev Neurosci* **14**, 83 (Feb, 2013).
22. S. N. Fry, R. Sayaman, M. H. Dickinson, The aerodynamics of free-flight maneuvers in *Drosophila*. *Science* **300**, 495 (Apr 18, 2003).
23. B. R. H. Geurten, P. Jahde, K. Corthals, M. C. Gopfert, Saccadic body turns in walking *Drosophila*. *Front Behav Neurosci* **8**, (Oct 22, 2014).

24. F. Wong, B. W. Knight, F. A. Dodge, Adapting Bump Model for Ventral Photoreceptors of Limulus. *J Gen Physiol* **79**, 1089 (1982).

25. C. E. Shannon, A Mathematical Theory of Communication. *At&T Tech J* **27**, 379 (1948).

26. R. R. D. van Steveninck, S. B. Laughlin, The rate of information transfer at graded-potential synapses. *Nature* **379**, 642 (Feb 15, 1996).

27. J. Howard, B. Blakeslee, S. B. Laughlin, The Intracellular Pupil Mechanism and Photoreceptor Signal - Noise Ratios in the Fly Lucilia-Cuprina. *Proc R Soc Ser B-Bio* **231**, 415 (Sep 22, 1987).

28. M. Rivera-Alba *et al.*, Wiring Economy and Volume Exclusion Determine Neuronal Placement in the *Drosophila* Brain. *Curr Biol* **21**, 2000 (Dec 6, 2011).

29. T. J. Wardill *et al.*, Multiple Spectral Inputs Improve Motion Discrimination in the *Drosophila* Visual System. *Science* **336**, 925 (May 18, 2012).

30. L. Zheng *et al.*, Feedback network controls photoreceptor output at the layer of first visual synapses in *Drosophila*. *J Gen Physiol* **127**, 495 (May, 2006).

31. M. F. Land, Motion and vision: why animals move their eyes. *J Comp Physiol A* **185**, 341 (Oct, 1999).

32. P. T. Gonzalez-Bellido, T. J. Wardill, M. Juusola, Compound eyes and retinal information processing in miniature dipteran species match their specific ecological demands. *P Natl Acad Sci USA* **108**, 4224 (Mar 8, 2011).

33. L. Freifeld, D. A. Clark, M. J. Schnitzer, M. A. Horowitz, T. R. Clandinin, GABAergic Lateral Interactions Tune the Early Stages of Visual Processing in *Drosophila*. *Neuron* **78**, 1075 (Jun 19, 2013).

34. M. G. Burg, P. V. Sarthy, G. Koliantz, W. L. Pak, Genetic and Molecular-Identification of a *Drosophila* Histidine-Decarboxylase Gene Required in Photoreceptor Transmitter Synthesis. *Embo J* **12**, 911 (Mar, 1993).

35. A. Dau *et al.*, Evidence for Dynamic Network Regulation of *Drosophila* Photoreceptor Function from Mutants Lacking the Neurotransmitter Histamine. *Frontiers in Neural Circuits* **10**, (2016).

36. M. Joesch, B. Schnell, S. V. Raghu, D. F. Reiff, A. Borst, ON and OFF pathways in *Drosophila* motion vision. *Nature* **468**, 300 (Nov 11, 2010).

37. U. Tepass, K. P. Harris, Adherens junctions in *Drosophila* retinal morphogenesis. *Trends Cell Biol* **17**, 26 (Jan, 2007).

38. K. G. Götz, Optomotory Investigation of the Visual-System of Some Eye Mutations of the *Drosophila* Fruit-Fly. *Kybernetik* **2**, 77 (1964).

39. D. G. Stavenga, Angular and spectral sensitivity of fly photoreceptors. II. Dependence on facet lens F-number and rhabdomere type in *Drosophila*. *Journal of Comparative Physiology a-Neuroethology Sensory Neural and Behavioral Physiology* **189**, 189 (Mar, 2003).

40. M. Juusola, Z. Song, R. C. Hardie, in *Encyclopedia of Computational Neuroscience,* D. Jaeger, R. Jung, Eds. (Springer, New York, 2015), pp. 2359-2376.

41. M. Juusola, A. Dau, L. Zheng, D. Rien, Electrophysiological method for recording intracellular voltage responses of *Drosophila* photoreceptors and interneurons to light stimuli in vivo. *Journal of Visualized Experiments*, e54142 (6/19/2016, 2016).

42. B. Minke, The History of the Prolonged Depolarizing Afterpotential (PDA) and Its Role in Genetic Dissection of *Drosophila* Phototransduction. *J Neurogenet* **26**, 106 (Jun, 2012).

43. J. H. van Hateren, Processing of natural time series of intensities by the visual system of the blowfly. *Vision Res* **37**, 3407 (Dec, 1997).

44. M. Juusola, G. G. de Polavieja, The rate of information transfer of naturalistic stimulation by graded potentials. *J Gen Physiol* **122**, 191 (Aug, 2003).

45. M. Juusola, R. C. Hardie, Light adaptation in Drosphila photoreceptors: II. Rising temperature increases the bandwidth of reliable signaling. *J Gen Physiol* **117**, 27 (Jan, 2001).

46. J. H. van Hateren, Theoretical Predictions of Spatiotemporal Receptive-Fields of Fly Lmcs, and Experimental Validation. *J Comp Physiol A* **171**, 157 (Sep, 1992).

47. J. H. van Hateren, H. P. Snippe, Information theoretical evaluation of parametric models of gain control in blowfly photoreceptor cells. *Vision Res* **41**, 1851 (Jun, 2001).

48. J. Melzig *et al.*, Genetic depletion of histamine from the nervous system of *Drosophila* eliminates specific visual and mechanosensory behavior. *J Comp Physiol A* **179**, 763 (Dec, 1996).

49. N. Franceschini, K. Kirschfeld, Phenomena of Pseudopupil in the Compound Eye of *Drosophila*. *Kybernetik* **9**, 159 (1971).

50. S. M. Tang, A. Guo, Choice behavior of *Drosophila* facing contradictory visual cues. *Science* **294**, 1543 (Nov 16, 2001).

51. R. C. Hardie, M. Juusola, Phototransduction in *Drosophila*. *Current opinion in neurobiology* **34C**, 37 (Jan 29, 2015).

52. Z. Song, Y. Zhou, M. Juusola, Random Photon Absorption Model Elucidates How Early Gain Control in Fly Photoreceptors Arises from Quantal Sampling. *Frontiers in computational neuroscience* **10**, 61 (2016).

53. J. E. Niven *et al.*, The contribution of Shaker K+ channels to the information capacity of *Drosophila* photoreceptors. *Nature* **421**, 630 (Feb 6, 2003).

54. M. Vähäsöyrinki, J. E. Niven, R. C. Hardie, M. Weckström, M. Juusola, Robustness of neural coding in *Drosophila* photoreceptors in the absence of slow delayed rectifier K+ channels. *J Neurosci* **26**, 2652 (Mar 8, 2006).

55. A. S. French *et al.*, The Dynamic Nonlinear Behavior of Fly Photoreceptors Evoked by a Wide-Range of Light Intensities. *Biophys J* **65**, 832 (Aug, 1993).

56. U. Friederich, D. Coca, S. Billings, M. Juusola, Data Modelling for Analysis of Adaptive Changes in Fly Photoreceptors. *Lect Notes Comput Sc* **5863**, 34 (2009).

57. M. Juusola, M. Weckstrom, R. O. Uusitalo, M. J. Korenberg, A. S. French, Nonlinear models of the first synapse in the light-adapted fly retina. *J Neurophysiol* **74**, 2538 (Dec, 1995).

58. J. I. Yellott, Spectral-Analysis of Spatial Sampling by Photoreceptors - Topological Disorder Prevents Aliasing. *Vision Res* **22**, 1205 (1982).

59. H. Barlow, B., in *Sensory communication,* W. A. Rosenblith, Ed. (MIT, Cambridge, 1961).

60. C. A. Hofstee, D. G. Stavenga, Calcium homeostasis in photoreceptor cells of *Drosophila* mutants inaC and trp studied with the pupil mechanism. *Visual Neurosci* **13**, 257 (Mar-Apr, 1996).

61. N. Franceschini, K. Kirschfeld, Automatic-Control of Light Flux in Compound Eye of Diptera - Spectral, Statical, and Dynamical Properties of Mechanism. *Biol Cybern* **21**, 181 (1976).

62.   Y. M. Elyada, J. Haag, A. Borst, Different receptive fields in axons and dendrites underlie robust coding in motion-sensitive neurons. *Nat Neurosci* **12**, 327 (Mar, 2009).

63.   D. G. Stavenga, Angular and spectral sensitivity of fly photoreceptors. III. Dependence on the pupil mechanism in the blowfly Calliphora. *Journal of Comparative Physiology a-Neuroethology Sensory Neural and Behavioral Physiology* **190**, 115 (Feb, 2004).

64.   M. Juusola, E. Kouvalainen, M. Jarvilehto, M. Weckstrom, Contrast Gain, Signal-to-Noise Ratio, and Linearity in Light-Adapted Blowfly Photoreceptors. *J Gen Physiol* **104**, 593 (Sep, 1994).

65.   E. Agi *et al.*, The Evolution and Development of Neural Superposition. *J Neurogenet* **28**, 216 (Sep-Dec, 2014).

66.   P. Vigier, The reception of light stimulant in the compound eyes of insects, particularly the Muscidae. *Cr Hebd Acad Sci* **145**, 633 (1907).

67.   P. Vigier, On the photoreceptive endings in the compound eyes of Muscids. *Cr Hebd Acad Sci* **145**, 532 (1907).

68.   I. A. Meinertzhagen, S. D. Oneil, Synaptic Organization of Columnar Elements in the Lamina of the Wild-Type in *Drosophila melanogaster*. *J Comp Neurol* **305**, 232 (Mar 8, 1991).

69.   S. R. Shaw, A. Frohlich, I. A. Meinertzhagen, Direct Connections between the R7/8 and R1-6 Photoreceptor Subsystems in the Dipteran Visual-System. *Cell Tissue Res* **257**, 295 (Aug, 1989).

70.   S. R. Shaw, Early Visual Processing in Insects. *J Exp Biol* **112**, 225 (1984).

71.   L. Zheng *et al.*, Network Adaptation Improves Temporal Representation of Naturalistic Stimuli in *Drosophila* Eye: I Dynamics. *Plos One* **4**, (Jan 30, 2009).

72.   M. F. Wernet *et al.*, Genetic Dissection Reveals Two Separate Retinal Substrates for Polarization Vision in *Drosophila*. *Curr Biol* **22**, 12 (Jan 10, 2012).

73.   M. F. Land, in *Adaptive Mechanisms in the Ecology of Vision,* S. Archer, M. B. Djamgoz, E. Loew, J. C. Partridge, Eds. (Kluwer Academic Publishers, London, 1998), pp. 51-72.

74.   B. G. Burton, B. W. Tatler, S. B. Laughlin, Variations in photoreceptor response dynamics across the fly retina. *J Neurophysiol* **86**, 950 (Aug, 2001).

75.   R. C. Hardie, in *Progress in Sensory Physiology,* D. Ottoson, Ed. (Springer, New York, (1985)), vol. 5, pp. 1–79.

76.   S. B. Laughlin, M. Weckstrom, Fast and Slow Photoreceptors - a Comparative-Study of the Functional Diversity of Coding and Conductances in the Diptera. *J Comp Physiol A* **172**, 593 (May, 1993).

77.   M. A. Z. Dippe, E. H. Wold, Antialiasing through stochastic sampling. *Computer Graphics* **19**, 69 (1985).

78.   M. V. Srinivasan, Shouldnt Directional Movement Detection Necessarily Be Color-Blind. *Vision Res* **25**, 997 (1985).

79.   T. Cover, J. Thomas, *Elements of Information Theory*. (John Wiley & Sons, Inc., New York, 1991).

80.   D. J. Field, Relations between the Statistics of Natural Images and the Response Properties of Cortical-Cells. *J Opt Soc Am A* **4**, 2379 (Dec, 1987).

81.   J. H. van Hateren, Processing of natural time series of intensities in the early visual system of the blowfly. *Perception* **26**, 6 (1997).

82.   G. D. Field *et al.*, Functional connectivity in the retina at the resolution of photoreceptors. *Nature* **467**, 673 (Oct 7, 2010).

83. D. Vasiliauskas *et al.*, Feedback from rhodopsin controls rhodopsin exclusion in *Drosophila* photoreceptors. *Nature* **479**, 108 (Nov 3, 2011).

84. M. C. Teich, B. E. A. Saleh, J. Perina, Role of Primary Excitation Statistics in the Generation of Antibunched and Sub-Poisson Light. *J Opt Soc Am B* **1**, 366 (1984).

85. J. Melzig, M. Burg, M. Gruhn, W. L. Pak, E. Buchner, Selective histamine uptake rescues photo- and mechanoreceptor function of histidine decarboxylase-deficient *Drosophila* mutant. *J Neurosci* **18**, 7160 (Sep 15, 1998).

86. A. W. Snyder, D. G. Stavenga, S. B. Laughlin, Spatial Information Capacity of Compound Eyes. *J Comp Physiol* **116**, 183 (1977).

87. A. W. Snyder, W. H. Miller, Photoreceptor Diameter and Spacing for Highest Resolving Power. *J Opt Soc Am* **67**, 696 (1977).

88. A. W. Snyder, Acuity of Compound Eyes - Physical Limitations and Design. *J Comp Physiol* **116**, 161 (1977).

89. E. J. Warrant, P. D. Mcintyre, Arthropod Eye Design and the Physical Limits to Spatial Resolving Power. *Prog Neurobiol* **40**, 413 (Apr, 1993).

90. J. H. van Hateren, Waveguide Theory Applied to Optically Measured Angular Sensitivities of Fly Photoreceptors. *J Comp Physiol* **154**, 761 (1984).

91. D. G. Stavenga, Angular and spectral sensitivity of fly photoreceptors. I. Integrated facet lens and rhabdomere optics. *Journal of Comparative Physiology a-Neuroethology Sensory Neural and Behavioral Physiology* **189**, 1 (Jan, 2003).

92. G. A. Horridge, K. Mimura, R. C. Hardie, Fly Photoreceptors .3. Angular Sensitivity as a Function of Wavelength and Limits of Resolution. *Proc R Soc Ser B-Bio* **194**, 151 (1976).

93. W. Wijngaard, D. G. Stavenga, Optical Crosstalk between Fly Rhabdomeres. *Biol Cybern* **18**, 61 (1975).

94. K. Kirschfeld, N. Franceschini, A Mechanism for Directing Flow of Light in Rhabdomeres of Complex Eyes in Musca. *Kybernetik* **6**, 13 (1969).

95. C. B. Boschek, Fine Structure of Peripheral Retina and Lamina Ganglionaris of Fly, Musca-Domestica. *Z Zellforsch Mik Ana* **118**, 369 (1971).

96. J. G. H. Roebroek, D. G. Stavenga, Insect Pupil Mechanisms .4. Spectral Characteristics and Light-Intensity Dependence in the Blowfly, Calliphora-Erythrocephala. *J Comp Physiol A* **166**, 537 (1990).

97. S. B. Laughlin, R. R. D. van Steveninck, J. C. Anderson, The metabolic cost of neural information. *Nat Neurosci* **1**, 36 (May, 1998).

98. M. F. Land, *Compound eye structure: Matching eye to environment. Adaptive Mechanisms in the Ecology of Vision.*, (Springer Netherlands, 1999).

99. J. E. Niven, J. C. Anderson, S. B. Laughlin, Fly photoreceptors demonstrate energy-information trade-offs in neural coding. *Plos Biol* **5**, 828 (Apr, 2007).

100. D. E. Nilsson, in *Facets of vision,* D. G. Stavenga, Hardie, R. C. , Ed. (Springer, Berlin, 1989).

101. S. B. Laughlin, The Role of Sensory Adaptation in the Retina. *J Exp Biol* **146**, 39 (Sep, 1989).

102. E. Warrant, P. Mcintyre, in *Nonlinear vision*. (1992), pp. 391-421.

103. D. S. Williams, Rhabdom Size and Photoreceptor Membrane Turnover in a Muscoid Fly. *Cell Tissue Res* **226**, 629 (1982).

104. S. B. Laughlin, Retinal Information Capacity and the Function of the Pupil. *Ophthal Physl Opt* **12**, 161 (Apr, 1992).

105. D. E. Nilsson, A. I. Ro, Did Neural Pooling for Night-Vision Lead to the Evolution of Neural Superposition Eyes. *J Comp Physiol A* **175**, 289 (Sep, 1994).

106. B. Pick, E. Buchner, Visual Movement Detection under Light-Adaptation and Dark-Adaptation in the Fly, Musca-Domestica. *J Comp Physiol* **134**, 45 (1979).

107. E. Warrant, T. Porombka, W. H. Kirchner, Neural image enhancement allows honeybees to see at night. *P Roy Soc B-Biol Sci* **263**, 1521 (Nov 22, 1996).

108. A. Honkanen, J. Takalo, K. Heimonen, M. Vahasoyrinki, M. Weckstrom, Cockroach optomotor responses below single photon level. *J Exp Biol* **217**, 4262 (Dec, 2014).

109. E. J. Warrant, Seeing better at night: life style, eye design and the optimum strategy of spatial and temporal summation. *Vision Res* **39**, 1611 (May, 1999).

110. S. B. Laughlin, R. C. Hardie, Common Strategies for Light Adaptation in the Peripheral Visual Systems of Fly and Dragonfly. *J Comp Physiol* **128**, 319 (1978).

111. T. Matic, S. B. Laughlin, Changes in the Intensity-Response Function of an Insects Photoreceptors Due to Light Adaptation. *J Comp Physiol* **145**, 169 (1981).

112. P. Skorupski, L. Chittka, Differences in Photoreceptor Processing Speed for Chromatic and Achromatic Vision in the Bumblebee, Bombus terrestris. *J Neurosci* **30**, 3896 (Mar 17, 2010).

113. M. V. Srinivasan, S. B. Laughlin, A. Dubs, Predictive Coding - a Fresh View of Inhibition in the Retina. *Proc R Soc Ser B-Bio* **216**, 427 (1982).

114. J. H. van Hateren, A Theory of Maximizing Sensory Information. *Biol Cybern* **68**, 23 (Nov, 1992).

115. J. H. van Hateren, Spatiotemporal Contrast Sensitivity of Early Vision. *Vision Res* **33**, 257 (Jan, 1993).

116. M. V. Srinivasan, D. R. Dvorak, Spatial Processing of Visual Information in the Movement-Detecting Pathway of the Fly - Characteristics and Functional-Significance. *J Comp Physiol* **140**, 1 (1980).

117. W. A. Ribi, Gap Junctions Coupling Photoreceptor Axons in the 1st Optic Ganglion of the Fly. *Cell Tissue Res* **195**, 299 (1978).

118. J. H. van Hateren, Electrical Coupling of Neuro-Ommatidial Photoreceptor Cells in the Blowfly. *J Comp Physiol A* **158**, 795 (1986).

119. A. Dubs, S. B. Laughlin, M. V. Srinivasan, Single Photon Signals in Fly Photoreceptors and 1st Order Interneurones at Behavioral Threshold. *J Physiol-London* **317**, 317 (1981).

120. J. H. van Hateren, Real and Optimal Neural Images in Early Vision. *Nature* **360**, 68 (Nov 5, 1992).

121. B. Greiner, W. A. Ribi, E. J. Warrant, A neural network to improve dim-light vision? Dendritic fields of first-order interneurons in the nocturnal bee Megalopta genalis. *Cell Tissue Res* **322**, 313 (Nov, 2005).

122. A. Dubs, The Spatial Integration of Signals in the Retina and Lamina of the Fly Compound Eye under Different Conditions of Luminance. *J Comp Physiol* **146**, 321 (1982).

123. A. Nikolaev *et al.*, Network Adaptation Improves Temporal Representation of Naturalistic Stimuli in *Drosophila* Eye: II Mechanisms. *Plos One* **4**, (Jan 30, 2009).

124. W. Hu, T. T. Wang, X. Wang, J. H. Han, I-h Channels Control Feedback Regulation from Amacrine Cells to Photoreceptors. *Plos Biol* **13**, (Apr, 2015).

125. J. G. J. Smakman, J. H. Vanhateren, D. G. Stavenga, Angular Sensitivity of Blowfly Photoreceptors - Intracellular Measurements and Wave-Optical Predictions. *J Comp Physiol* **155**, 239 (1984).

126. J. G. J. Smakman, D. G. Stavenga, Angular Sensitivity of Blowfly Photoreceptors - Broadening by Artificial Electrical Coupling. *J Comp Physiol A* **160**, 501 (1987).

127. M. Wilson, Angular Sensitivity of Light and Dark Adapted Locust Retinula Cells. *J Comp Physiol* **97**, 323 (1975).

128. R. C. Hardie, Electro-Physiological Analysis of Fly Retina .1. Comparative Properties of R1-6 and R7 and 8. *J Comp Physiol* **129**, 19 (1979).

129. K. Mimura, Receptive-Field Patterns in Photoreceptors of the Fly. *J Comp Physiol* **141**, 349 (1981).

130. E. Warrant, D. E. Nilsson, *Invertebrate vision*. (Cambridge University Press, Cambridge, 2006).

131. Y. Washizu, D. Burkhardt, P. Streck, Visual Field of Single Retinula Cells and Interommatidial Inclination in the Compound Eye of the Blowfly Calliphora-Erythrocephala. *Z Vergl Physiol* **48**, 413 (1964).

132. D. Burkhardt, Vision of Insects. *J Comp Physiol* **120**, 33 (1977).

133. E. Eguchi, T. Horikoshi, Comparison of Stimulus-Response (V-Log-I) Functions in 5 Types of Lepidopteran Compound Eyes (46 Species). *J Comp Physiol* **154**, 3 (1984).

134. D. G. Stavenga, Visual acuity of fly photoreceptors in natural conditions - dependence on UV sensitizing pigment and light-controlling pupil. *J Exp Biol* **207**, 1703 (Apr, 2004).

135. J. H. van Hateren, 3 Modes of Spatiotemporal Preprocessing by Eyes. *J Comp Physiol A* **172**, 583 (May, 1993).

136. A. Klaus, E. J. Warrant, Optimum spatiotemporal receptive fields for vision in dim light. *J Vision* **9**, (2009).

137. J. Howard, W. M. Roberts, A. J. Hudspeth, Mechanoelectrical Transduction by Hair-Cells. *Annu Rev Biophys Bio* **17**, 99 (1988).

138. N. Franceschini, in *Biocybernetics of Vision: Integrative mechanisms and CognitiveProcesses,* C. Taddei-Ferretti, Ed. (World Scientific, Singapore, 1998), pp. 341-361.

139. N. Franceschini, R. Chagneux, in *Gottingen Neurobiology Report* N. Elsner, H. Breer, Eds. (Georg Thieme Verlag, Stuttgart, 1994).

140. N. Franceschini, R. Chagneux, K. Kirschfeld, paper presented at the Gottingen Neurobiology Report, Stuttgart, 1995.

141. N. Franceschini, R. Chagneux, K. Kirschfeld, A. Mucke, paper presented at the Gottingen Neurobiology Report, Stuttgart, 1991.

142. S. R. Shaw, Retinal Resistance Barriers and Electrical Lateral Inhibition. *Nature* **255**, 480 (1975).

143. M. Heisenberg, Separation of Receptor and Lamina Potentials in Electroretinogram of Normal and Mutant *Drosophila*. *J Exp Biol* **55**, 85 (1971).

144. R. C. Hardie, N. Franceschini, W. Ribi, K. Kirschfeld, Distribution and Properties of Sex-Specific Photoreceptors in the Fly Musca-Domestica. *J Comp Physiol* **145**, 139 (1981).

145. J. J. Atick, Could Information-Theory Provide an Ecological Theory of Sensory Processing. *Network-Comp Neural* **3**, 213 (May, 1992).

146. A. L. Stockl, D. C. O'Carroll, E. J. Warrant, Neural Summation in the Hawkmoth Visual System Extends the Limits of Vision in Dim Light. *Curr Biol* **26**, 821 (Mar 21, 2016).

147. A. N. Abou Tayoun *et al.*, The *Drosophila* SK Channel (dSK) Contributes to Photoreceptor Performance by Mediating Sensitivity Control at the First Visual Network. *J Neurosci* **31**, 13897 (Sep 28, 2011).

148. I. Sinakevitch, N. J. Strausfeld, Chemical neuroanatomy of the fly's movement detection pathway. *J Comp Neurol* **468**, 6 (Jan 1, 2004).

149. E. S. Reynolds, Use of Lead Citrate at High Ph as an Electron-Opaque Stain in Electron Microscopy. *J Cell Biol* **17**, 208 (1963).

150. P. T. Gonzalez-Bellido, the University of Sheffield (2009).

151. G. P. Luke, C. H. G. Wright, S. F. Barrett, A Multiaperture Bioinspired Sensor With Hyperacuity. *Ieee Sens J* **12**, 308 (Feb, 2012).

152. D. B. Zurek, X. J. Nelson, Hyperacute motion detection by the lateral eyes of jumping spiders. *Vision Res* **66**, 26 (Aug 1, 2012).

153. M. Järvilehto, F. Zettler, Electrophysiological-Histological Studies on Some Functional Properties of Visual Cells and Second-Order Neurons of an Insect Retina. *Z Zellforsch Mik Ana* **136**, 291 (1973).

154. F. Zettler, M. Järvilehto, Lateral Inhibition in an Insect Eye. *Z Vergl Physiol* **76**, 233 (1972).

155. M. E. Chiappe, J. D. Seelig, M. B. Reiser, V. Jayaraman, Walking Modulates Speed Sensitivity in *Drosophila* Motion Vision. *Curr Biol* **20**, 1470 (Aug 24, 2010).

156. J. Haag, A. Wertz, A. Borst, Central gating of fly optomotor response. *P Natl Acad Sci USA* **107**, 20104 (Nov 16, 2010).

157. G. Maimon, A. D. Straw, M. H. Dickinson, Active flight increases the gain of visual motion processing in *Drosophila*. *Nat Neurosci* **13**, 393 (Mar, 2010).

158. S. M. Tang, M. Juusola, Intrinsic Activity in the Fly Brain Gates Visual Information during Behavioral Choices. *Plos One* **5**, (Dec 30, 2010).

159. J. C. Anderson, S. B. Laughlin, Photoreceptor performance and the co-ordination of achromatic and chromatic inputs in the fly visual system. *Vision Res* **40**, 13 (2000).

160. R. C. Hardie, Properties of Photoreceptor-R7 and Photoreceptor-R8 in Dorsal Marginal Ommatidia in the Compound Eyes of Musca and Calliphora. *J Comp Physiol* **154**, 157 (1984).

161. B. G. Burton, S. B. Laughlin, Neural images of pursuit targets in the photoreceptor arrays of male and female houseflies Musca domestica. *J Exp Biol* **206**, 3963 (Nov, 2003).

162. R. S. A. Brinkworth, E. L. Mah, J. P. Gray, D. C. O'Carroll, Photoreceptor processing improves salience facilitating small target detection in cluttered scenes. *J Vision* **8**, (2008).

163. B. Krekelberg, M. Lappe, Neuronal latencies and the position of moving objects. *Trends Neurosci* **24**, 335 (Jun, 2001).

164. R. Nijhawan, Neural delays, visual motion and the flash-lag effect. *Trends Cogn Sci* **6**, 387 (Sep, 2002).

165. M. J. Berry, I. H. Brivanlou, T. A. Jordan, M. Meister, Anticipation of moving stimuli by the retina. *Nature* **398**, 334 (Mar 25, 1999).

166. V. Braitenberg, P. Debbage, Regular Net of Reciprocal Synapses in Visual System of Fly, Musca-Domestica. *J Comp Physiol* **90**, 25 (1974).

167. J. Rister *et al.*, Dissection of the peripheral motion channel in the visual system of *Drosophila* melanogaster. *Neuron* **56**, 155 (Oct 4, 2007).

168. P. Z. Marmarelis, G. D. Mccann, Development and Application of White-Noise Modeling Techniques for Studies of Insect Visual Nervous-System. *Kybernetik* **12**, 74 (1973).

169. H. Eckert, L. G. Bishop, Nonlinear Dynamic Transfer Characteristics of Cells in Peripheral Visual Pathway of Flies .1. Retinula Cells. *Biol Cybern* **17**, 1 (1975).

170. R. Gemperlein, G. D. Mccann, Study of Response Properties of Retinula Cells of Flies Using Nonlinear Identification Theory. *Biol Cybern* **19**, 147 (1975).

171. M. J. Korenberg, M. Juusola, A. S. French, Two methods for calculating the responses of photoreceptors to moving objects. *Ann Biomed Eng* **26**, 308 (Mar-Apr, 1998).

172. M. Juusola, J. E. Niven, A. S. French, Shaker K+ channels contribute early nonlinear amplification to the light response in *Drosophila* photoreceptors. *J Neurophysiol* **90**, 2014 (Sep, 2003).

173. J. E. Niven, M. Vahasoyrinki, M. Juusola, A. S. French, Interactions between light-induced currents, voltage-gated currents, and input signal properties in *Drosophila* photoreceptors. *J Neurophysiol* **91**, 2696 (Jun, 2004).

174. E. P. Hornstein, D. C. O'Carroll, J. C. Anderson, S. B. Laughlin, Sexual dimorphism matches photoreceptor performance to behavioural requirements. *P Roy Soc B-Biol Sci* **267**, 2111 (Oct 22, 2000).

175. C. Schilstra, J. H. Van Hateren, Blowfly flight and optic flow I. Thorax kinematics and flight dynamics. *J Exp Biol* **202**, 1481 (Jun, 1999).

176. J. H. van Hateren, C. Schilstra, Blowfly flight and optic flow II. Head movements during flight. *J Exp Biol* **202**, 1491 (Jun, 1999).

177. M. Schetzen, *The Volterra and Wiener theories of nonlinear systems.*, (1980).

178. M. Juusola, R. O. Uusitalo, M. Weckstrom, Transfer of Graded Potentials at the Photoreceptor Interneuron Synapse. *J Gen Physiol* **105**, 117 (Jan, 1995).

179. M. J. Korenberg, S. B. Bruder, P. J. Mcilroy, Exact Orthogonal Kernel Estimation from Finite Data Records - Extending Wiener Identification of Nonlinear-Systems. *Ann Biomed Eng* **16**, 201 (1988).

180. M. J. Korenberg, L. D. Paarmann, Applications of Fast Orthogonal Search - Time-Series Analysis and Resolution of Signals in Noise. *Ann Biomed Eng* **17**, 219 (1989).

181. M. H. Asyali, M. Juusola, Use of Meixner functions in estimation of Volterra kernels of nonlinear systems with delay. *Ieee T Bio-Med Eng* **52**, 229 (Feb, 2005).

182. G. H. Golub, C. Reinsch, Singular Value Decomposition and Least Squares Solutions. *Numer Math* **14**, 403 (1970).

183. C. L. Lawson, R. J. Hanson, *Solving least squares problems.* (SIAM, 1974).

184. R. Nijhawan, Motion Extrapolation in Catching. *Nature* **370**, 256 (Jul 28, 1994).

185. E. Brenner, J. B. J. Smeets, Motion extrapolation is not responsible for the flash-lag effect. *Vision Res* **40**, 1645 (2000).

186. O. Sayeed, S. Benzer, Behavioral genetics of thermosensation and hygrosensation in *Drosophila*. *P Natl Acad Sci USA* **93**, 6079 (Jun 11, 1996).

187. J. D. Victor, in *Nonlinear vision: Determination of neural receptive fields, function, and networks*. (CRC, 1992), pp. 1-37.

188. N. Franceschini, K. Kirschfeld, Optical Study in-Vivo of Photoreceptor Elements in Compound Eye of *Drosophila*. *Kybernetik* **8**, 1 (1971).

189. S. G. Britt, R. Feiler, K. Kirschfeld, C. S. Zuker, Spectral Tuning of Rhodopsin and Metarhodopsin in-Vivo. *Neuron* **11**, 29 (Jul, 1993).

190. G. Laufer, *Introduction to optics and lasers in engineering*. (Cambridge University Press 1996).
191. D. P. Corey *et al.*, TRPA1 is a candidate for the mechanosensitive transduction channel of vertebrate hair cells. *Nature* **432**, 723 (Dec 9, 2004).
192. A. Bahl, G. Ammer, T. Schilling, A. Borst, Object tracking in motion-blind flies. *Nat Neurosci* **16**, 730 (Jun, 2013).
193. K. G. Götz, Visual guidance in *Drosophila*. *Basic Life Science* **16**, 391 (1980).
194. M. Juusola, Linear and Nonlinear Contrast Coding in Light-Adapted Blowfly Photoreceptors. *J Comp Physiol A* **172**, 511 (May, 1993).
195. C. I. Yeh, D. J. Xing, R. M. Shapley, "Black" Responses Dominate Macaque Primary Visual Cortex V1. *J Neurosci* **29**, 11753 (Sep 23, 2009).
196. J. Kremkow *et al.*, Neuronal nonlinearity explains greater visual spatial resolution for darks than lights. *Proc Natl Acad Sci U S A*, (Feb 10, 2014).
197. C. P. Ratliff, B. G. Borghuis, Y. H. Kao, P. Sterling, V. Balasubramanian, Retina is structured to process an excess of darkness in natural scenes. *P Natl Acad Sci USA* **107**, 17368 (Oct 5, 2010).
198. J. L. Fox, M. A. Frye, Figure-ground discrimination behavior in *Drosophila*. II. Visual influences on head movement behavior. *J Exp Biol* **217**, 570 (Feb 15, 2014).
199. M. Heisenberg, E. Buchner, Role of Retinula Cell-Types in Visual Behavior of *Drosophila melanogaster*. *J Comp Physiol* **117**, 127 (1977).
200. E. Buchner, Elementary Movement Detectors in an Insect Visual-System. *Biol Cybern* **24**, 85 (1976).
201. S. M. Tang, R. Wolf, S. P. Xu, M. Heisenberg, Visual pattern recognition in *Drosophila* is invariant for retinal position. *Science* **305**, 1020 (Aug 13, 2004).