Transposable element exaptation is the primary source of novelty in the primate gene regulatory landscape

Marco Trizzino[1,#,*], YoSon Park[1,*], Marcia Holsbach-Beltrame[1], Katherine Aracena[1], Katelyn Mika[2], Minal Caliskan[1], George H. Perry[3], Vincent J. Lynch[2] and Christopher D. Brown[1,#]

1. Department of Genetics, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, 19104
2. Department of Human Genetics, University of Chicago, Chicago, IL, 60637
3. Departments of Anthropology and Biology, Pennsylvania State University, University Park, PA, 16802

* Co-first author
# Correspondence: chrbro@upenn.edu (C.D.B.), marco.trizzino83@gmail.com (M.T.)

Summary

Gene regulation plays a critical role in the evolution of phenotypic diversity. We investigated the evolution of liver promoters and enhancers in six primate species. We performed ChIP-seq for two histone modifications and RNA-seq to profile cis-regulatory element (CRE) activity and gene expression. The primate regulatory landscape is largely conserved across the lineage. Conserved CRE function is associated with sequence conservation, proximity to coding genes, cell type specificity of CRE function, and transcription factor binding. Newly evolved CREs are enriched in immune response and neurodevelopmental functions, while conserved CREs bind master regulators. Transposable elements (TEs) are the primary source of novelty in primate gene regulation. Newly evolved CREs are enriched in young TEs that affect gene expression. However, only 17% of conserved CREs overlap a TE, suggesting that target gene expression is under strong selection. Finally, we identified specific genomic features driving the functional recruitment of newly inserted TEs.

M.T. Current affiliation: Gene Expression and Regulation Program, The Wistar Institute, Philadelphia, PA, 19104

Introduction

The evolution of cis-regulatory elements (CREs) plays an important role in phenotypic and behavioral evolution (King and Wilson, 1975; Rockman et al., 2005; Loisel et al., 2006; Pollard et al., 2006; Prabhakar et al., 2008; Warner et al., 2009; Babbitt et al., 2010; Cain et al., 2011; Marnetto et al., 2014; Zhou et al., 2014; Villar et al., 2015). Many aspects of CRE evolution in mammals have been characterized (Schmidt et al., 2010; Cain et al., 2011; Zhou et al., 2014; Prescott et al., 2015; Reilly et al., 2015; Villar et al., 2015; Emera et al., 2016), and suggest a role for transposable elements (TEs) in the evolution of gene regulation (McClintock, 1950, 1984; Britten and Davidson, 1969; Davidson and Britten, 1979; Jordan et al., 2003; Bejerano et al., 2006; Wang et al., 2007; Bourque et al., 2008; Sasaki et al., 2008; Markljung et al., 2009; Kunarso et al., 2010; Lynch et al., 2011, 2015; Chuong et al., 2013, 2016; Schmidt et al., 2012; Xie et al., 2013; del Rosario et al., 2014; Sundaram et al., 2014; Du et al., 2016; Rayan et al., 2016). However, validating the functional contribution of TEs in the mammalian gene regulation remains a challenge. Lynch et al. (2011, 2015) demonstrated that the recruitment of novel regulatory networks in the uterus was likely mediated by ancient mammalian TEs. However, Emera and colleagues (2016) suggested that neocortical enhancers do not exhibit strong evidence of transposon exaptation.

Many important questions remain unanswered: to what extent are poised and active regulatory elements functionally conserved? Are specific genomic features predictive of CRE conservation? To what extent have TEs driven the evolution of gene regulation? And finally, what determines which TE insertions are recruited as functional CREs? Establishing answers to these questions is critical for understanding how the evolution of regulatory elements contributes to the conservation and divergence of gene expression and complex traits.

With a goal of answering these questions, we collected liver samples from six primate species. Core liver functions are largely conserved across primate species. However, different environmental exposures, diets, and lifestyles likely directed the adaptation of liver functions, and associated regulatory evolution, making this tissue an optimal model in which explore the conservation and divergence of the gene regulation.

To characterize primate liver cis-regulatory evolution, we performed chromatin immunoprecipitation followed by sequencing (ChIP-seq) for Histone H3 Lysine 27

acetylation (H3K27ac), which marks active enhancers and promoters, and Histone H3 Lysine 4 mono-methylation (H3K4me1), which marks poised regulatory elements on liver tissues from six primate species, including at least one species from each major primate clade to maximize evolutionary diversity within primates (Perelman et al., 2012). We generated whole transcriptome sequencing (RNA-seq) data from the same specimens to quantify gene expression variation across species. We estimated the degree of evolutionary conservation of regulatory activity and gene expression levels across the entire lineage, and characterized the genomic features associated with evolutionary conservation of gene regulation, to understand why some enhancers and promoters are conserved across species whereas others are subject to rapid turnover.

The activity of the majority of human CREs is conserved across the entire primate lineage, and the differences in gene expression and regulation reflect the phylogenetic distance between species. Conservation of cis-regulatory activity is associated with nucleotide sequence conservation, gene function, gene distance, cell-specificity of CRE function, and transcription factor binding site (TFBS) density. Strikingly, human- and ape-specific enhancers and promoters are significantly enriched for evolutionarily young TEs. In particular, the majority of human- and ape-specific CREs are derived from SINE-VNTR-*Alus* (SVAs) and Long-Terminal-Repeats (LTRs), respectively. On the other hand, only a minor fraction of evolutionarily conserved CREs are derived from TEs, indicating that purifying selection on the associated genes likely preserve these CREs from being disrupted by TE insertions, thus conserving the expression of the associated genes. In addition, we characterized SVAs that evolved into regulatory elements, and estimated potential impacts of these SVAs on gene regulation across the lineage. We validated the regulatory activity of several TE families, and conclude with a new model describing specific genomic features that strengthen the potential adaptation of TEs into functional regulatory elements (exaptation; Brosius and Gould; 1992; de Souza et al., 2013).

Results

Data generation, quality assessment, and validation
We generated a total of 757 million RNA-seq reads and 1.70 billion ChIP-seq reads (H3K27ac, H3K4me1, and input) from *post mortem* livers of three or four individuals per

species of mouse lemur (*Microcebus murinus*), bushbaby (*Otolemur garnettii*), marmoset (*Callithrix jacchus*), rhesus macaque (*Macaca mulatta*), chimpanzee (*Pan troglodytes*), and human (*Homo sapiens*) (Fig. 1; Table S1). The six species were selected to include at least one species from each of the major primate clades, thus maximizing phylogenetic diversity within primates. The samples were all from young adults and, with the exception of the bushbaby, included both males and females. In total, 18 RNA-seq and 14 ChIP-seq samples remained post-QC and were used for analyses (Table S1). On average, we sequenced 42.1 million RNA-seq reads and 40.6 million ChIP-seq reads per sample (Table S1). We applied stringent quality control (QC) measures to assess library construction, sequencing, and peak-calling methods. Read mappability after the filtering steps was consistent across species and assays (77% in ChIP-seq, 78% in RNA-seq; Table S1), suggesting that differences in genome assembly quality do not introduce large biases.

ChIP-seq data were used to map the distribution of active (H3K27ac) and poised (H3K4me1) CREs in the six primate genomes (Fig. 1). We mapped ChIP-seq reads to their respective reference genomes with BWA-mem (Li, 2013) and identified regions of significant H3K27ac and H3K4me1 enrichment in the human liver, treating all human individuals as replicates in the peak calling procedure with MACS2 (Zhang et al., 2008). 85.6% of the regions marked by H3K4me1 overlapped regions marked by H3K27ac. A total of 84,253 human peaks remained after merging overlapping peaks from the two histone markers. Next we identified regions orthologous to the human consensus peaks from the genomes of non-human primates using the Ensembl multiple sequence alignment (MSA) database. We catalogued 47,673 total human CREs with orthologs in all six species: 40,527 enhancers (distance from transcription start site (TSS) > 1 kb) and 7,146 promoters (distance from TSS ≤ 1 kb).

Figure 1 - **Experimental design and analytical pipeline**. (A) Sampling included three to four specimens from six species representing all of the major primate clades. (B) ChIP-seq and RNA-seq profiles were produced from the liver samples. Differential histone modification and gene expression analyses were performed on the orthologous CREs and genes in each species, respectively. Outputs shown in the diagram were used for downstream analyses and validations.

Several lines of evidence indicate that the regions of histone modification we have identified represent active CREs. First, 99.0% and 73.3% of ENCODE HepG2 H3K27ac and H3K4me1 regions (ENCODE Project Consortium, 2012), respectively, overlapped with one or more human peaks. Similarly, 63.2% of the 43,011 permissive enhancers that were predicted by the FANTOM5 Consortium based on enhancer RNA expression (Andersson et al., 2014) overlapped with one or more of human peaks identified in our study. Further, the promoters of 98.1% of genes expressed in the liver overlapped a region of histone modification (Fig. S1). Finally we compared our human H3K27ac data to a recent study focused on liver CREs in mammals (Villar et al., 2015) and demonstrated that peaks bearing signatures of robust and broad regulatory activities are largely reproducible across studies, despite variation attributable to different study designs (Fig. S2).

ChIP-seq experiments can identify regions of the genome bound by histones and other proteins that characterize regulatory elements, however, this does not guarantee

these regions are functional regulatory elements (Pickrell et al., 2011; Cusanovic et al., 2014; Jain et al., 2015). Therefore, we used a novel parallelized reporter assay (Melnikov et al., 2012; Patwardhan et al., 2012; Sharon et al., 2012; see methods) to validate the regulatory function of predicted human liver CREs. Specifically, we tested the regulatory activity of 1 kb fragments from 122 putative regulatory elements in HepG2 cells, including both evolutionarily conserved and recently evolved CREs (see below). Among 122 tested elements, 79 drove significant reporter gene expression levels (42/53 enhancers [79.2%] and 36/69 promoters [52.2%]; Table S6), suggesting that the majority of the CREs predicted in our study based on the enrichment of active histone modification states are likely functional regulatory elements in the human liver.

The majority of human CREs are functionally conserved across primates

After extracting ChIP-seq read counts for the six species from 47,673 orthologous regions, we assessed evidence of differential histone modification between species with DESeq2 (Love et al., 2014), using the ChIP-seq input data as a covariate. We compared ChIP-seq read counts in the 47,673 regions by means of all possible human-centric species × species and group × group pairwise comparisons (see methods). This approach provides a quantitative assessment of histone modification profiles across species, while avoiding issues arising from many potential experimental variables that may confound peak calling (Waszak et al., 2015). A specific analysis of human and marmoset, the latter being the species with the smallest number of peaks called in this study (Supplemental File S1), strongly supported the validity and robustness of our approach (Fig. S3).

The majority of the 47,673 human CREs (63.8%) did not show significant differential histone modifications in any of the tested pairwise comparisons (FDR < 10%; Fig. 2). This suggests that these regulatory regions are consistently active across the primate lineage and thus represent evolutionarily conserved primate CREs. Although the absence of differential histone modifications in a pairwise comparison between two species is not a direct proof of CRE conservation, we demonstrated that the selected FDR threshold does not affect downstream conclusions (Table S7). As an additional control, we performed a chimpanzee-centric analysis for the regions orthologous to chimpanzee consensus peaks (hereafter, chimpanzee CREs), and demonstrated that 62.5% of these regions were not differentially histone modified in any of the pairwise comparisons. This observation is consistent with 63.8% conserved CREs identified in the human-centric analysis, indicating

that the differential histone modification analysis is robust and species-specific bias is unlikely.

Promoters are significantly more conserved than enhancers (68.9% and 62.8%, respectively; Fisher's exact test *p* < 2.2×10^{-16}; Fig. 2), as observed previously (Villar et al., 2015). On the other hand, 36.2% of orthologous CREs exhibited differential histone modification state across species (Fig. 2). We detected 57 human-specific CREs (0.13%; Fig. 3) and 544 ape-specific CREs (1.42%; Fig. 3). Together, our differential histone state analysis results are broadly supported by several studies that have consistently suggested a high degree of regulatory element conservation between closely related species in metazoans (Cotney et al., 2013; Boyle et al., 2014; Prescott et al., 2015; Reilly et al., 2015; Emera et al., 2016).

Ape-specific CRE
chr15: 73972589-73974747

Conserved CRE
chr13: 42270792-42273375

All CREs (H3K27ac+H3K4me1)
N = 47,673

Conserved (63.79 %)

Recently evolved (36.21%)

- Human specific (0.13 %)
- Ape specific (1.42%)
- Other recently evolved (34.66%)

evolutionarily conserved. (A) Plots show examples of ...nd conserved CREs. (B) Fractions of conserved and recently ...breakdown of enhancers and promoters.

...ssessed the extent to which primate-conserved CREs identified in this study are also evolutionarily conserved across a broader range of mammals. In particular, we compared our conserved H3K27ac CREs with the H3K27ac profile of the opossum, the species with the earliest divergence from humans (>180 million years) in the Villar et al.

(2015) dataset. 2,854 primate-conserved promoters and 9,456 primate-conserved enhancers have orthologous regions in the opossum genome. Among these, 71.3% of the promoters and 19.1% of the enhancers had significant H3K27ac enrichment in both species, supporting that most of the primate conserved promoters show conserved activity in all of the mammalian clade, whereas only a fraction of the primate conserved enhancers is also conserved across mammals. Further, the two studies come to consistent estimates of the fraction of differentially active CREs per million years: 0.06–0.12% in primates and 0.07% in mammals.

The conservation of the nucleotide sequence is associated with conservation of regulatory activity

Previous studies have suggested that the sequence conservation is associated with conservation of regulatory activity, especially in absence of comparative functional assays (Brown et al., 2007; Cooper and Brown, 2008; Pollard et al., 2010; Gittelman et al., 2015; Holloway et al., 2015; Villar et al., 2015; Yang et al., 2015; Dong et al., 2016; Lewis at al., 2016). For each human-centric species × species comparison, we estimated: i) the fraction of differentially modified CREs; ii) the fraction of differentially expressed genes from a set of 10,243 genes with six way orthologs (Table S3); and iii) the per-nucleotide pairwise sequence divergence for each species with respect to humans for each of the 47,673 unique orthologous CREs

Differential histone state fractions ranged from 0.77% in the human × chimpanzee, to 21.9% in the human × mouse lemur comparisons (Fig. 3a). Similarly, differential gene expression ranged from 5.93% in the human × chimpanzee to 16.0% in the human × mouse lemur comparisons (Fig. 3b). Both differential histone modification and differential gene expression fractions reflected phylogenetic distance between humans and other tested species. Differentially expressed genes were significantly more likely to be associated with a differentially modified CRE than expected by chance (9.90%; Fisher's exact test $p < 2.2 \times 10^{-16}$).

Sequence conservation was significantly correlated with regulatory activity (human × chimpanzee, logistic regression $p = 5.7 \times 10^{-16}$; human × rhesus macaque, $p = 6.7 \times 10^{-8}$; human × marmoset, $p = 4.0 \times 10^{-9}$; human × bushbaby, $p < 2.2 \times 10^{-16}$; human × mouse lemur, $p < 2.2 \times 10^{-16}$; Fig. 3c). 24,691 CREs overlapped 94,578 placental mammal phastCons elements (i.e. regions of the genomes with consistent nucleotide sequence conservation across species; Siepel et al., 2005). The fraction of evolutionary conserved

231 CREs overlap

232 (Fisher's exact

233 conserved nuc

...ng genes

...nes (Fisher's exact test p = 0.5×10⁻) or lincRNAq (Fisher's exact test p |

). For closely related species, regulatory activity was conserved, regardless

stance to the nearest TSS (human × chimpanzee, logistic regression p = 0.261;

numan × rhesus macaque, p = 0.336; Fig. 4b). However, for more distantly related species

pairs, the evolutionary conservation of the CRE activity was significantly lower in regions

more distant from TSSs (human × marmoset, logistic regression p = 6.0×10^{-15}; human ×
bushbaby, p = 2.1×10^{-5}, human × mouse lemur, p = 2.8×10^{-5}; Fig. 4b). Intronic enhancers
were significantly more conserved than intergenic enhancers (63.0% and 55.2%
respectively; Fisher's exact test p < 2.2×10^{-16}). These data demonstrate increased selective
pressure to maintain regulatory activity in the vicinity of protein coding genes.

Multiple genomic features indicative of broad regulatory element activity were
significantly associated with the conservation of regulatory activity across the primate
phylogeny. Promoters and enhancers overlapping regions of chromatin accessibility in
many cell types (ENCODE Project Consortium, 2012) were significantly more functionally
conserved than those that are functional in only a small number of tissues (logistic
regression p < 2.2×10^{-16}; Fig. 4c). Similarly, CREs that overlapped many TFBS, as
identified by ENCODE ChIP-seq in HepG2 cells, were significantly more evolutionarily
conserved than those with fewer binding sites (logistic regression p < 2.2×10^{-16}; Fig. 4d).
Further, liver enhancers that generate consistent enhancer RNA transcription were
significantly more evolutionarily conserved than the untranscribed enhancers (Fig. S3;
Fisher's exact test p = 0.04613), validating previous findings (Andersson et al., 2014).

Finally, we used GOrilla (Eden et al., 2007; Eden et al., 2009) to identify biological
processes Gene Ontology terms that are enriched in genes found within 10-kb of
evolutionarily conserved CREs, using as background all of the genes found within 10 kb
from any of the 47,673 orthologous CREs. We found an enrichment for housekeeping
functions involved in the regulation of cellular, transcriptional and developmental processes
(Table S4). These findings support previous observations that conserved CREs are
proximal to housekeeping genes (FANTOM5 Consortium, 2014; Villar et al., 2015).

Specific transcription factor motifs are associated with regulatory conservation and
turnover
We used the MEME Suite (Bailey et al., 2009) to identify sequence motifs enriched in
human-specific, ape-specific, and evolutionarily conserved liver CREs. Human-specific
CREs were enriched with motifs for TFs associated with immune response and
hematopoietic maintenance (Fig. 4e; Supplemental File S2), such as PRDM1 (BLIMP1)
which is induced upon viral infection and represses beta-interferon (β-IFN) gene
expression. The rapid evolution of immune response genes and TFs is supported by many
studies in vertebrates and in *Drosophila melanogaster*, demonstrating that while the central
machinery of immune responses is strongly conserved, several components of the

extended molecular networks can evolve rapidly or diversify as a consequence of
evolutionary competition between hosts and pathogens (Jansa et al., 2003; Vallender 2004;
Sackton et al., 2007; Obbard et al., 2009; Schadt et al., 2009; Grueber et al., 2014; Lazzaro
and Schneider 2014; Salazar-Jaramillo et al., 2014; Zak et al., 2014; Sironi et al., 2015;
Wertheim, 2015; Chuong et al., 2016). Similarly, recently evolved promoters and enhancers
in primates are enriched in functions associated to neuronal proliferation, migration and
cortical map organization (Boyd et al., 2015; Reilly et al., 2015; Emera et al., 2016). In
contrast, ape-specific CREs were instead enriched with motifs representing binding sites for
TFs involved in liver function but, remarkably, also in brain and neural system proliferation
and development (Fig. 4e; Supplemental File S2).

Evolutionarily conserved CREs were enriched in TFBSs for master regulators and
homeobox genes that establish cell-type identity in liver cells (Fig. 4e; Supplemental File
S2). Among these master regulators, HNF4A is essential for the differentiation of human
hepatic progenitor cells by establishing the expression of the network of transcription
factors that controls the onset of hepatocyte cell fate (DeLaForest et al., 2011). Likewise,
CEBPA is required for the liver cell specification and gene function, and the associated
TFBSs are highly conserved across mammals (Ballester et al., 2014). Both CEBPA and
HNF4A have conserved cis-regulatory activity and significantly higher numbers of shared
TF binding events than expected by chance alone across distant vertebrates (Schmidt et
al., 2010). These results demonstrate that evolution shapes the regulatory landscape by
preserving the regulatory activity in essential metabolic and developmental pathways, while
permitting incessant renovation of specific networks that are under strong selective
pressures.

Figure 4 - **Genomic features associated to CRE conservation**. (A) Fractions of conserved and recently evolved CREs associated to protein coding genes, lincRNAs and pseudogenes. (B) Distribution of CRE conservation along the genome. (C) Correlation between CRE conservation and cell-type specificity based on ENCODE data. (D) Correlation between CRE conservation and number of ENCODE HepG2 TFBSs present of each CRE. (E) Examples of i) enriched motifs associated to immune response (BLIMP1, IZKF1, STAT3) in human-specific CREs; ii) enriched motifs associated to language and neural development (FOXA2, GSC, TBR1) in ape-specific CREs; iii) enriched motifs associated to master regulators (CEPBA, GATA2, HNF4A) in conserved CREs.

Exaptation of TEs into functional CREs is pervasive in the primate genomes

Previous studies have found that TEs can contribute to the origin of CREs. To quantify the contribution of TEs in the liver gene expression regulation in primates, we annotated each liver CRE based on overlap with RepeatMasker elements (Smit, Hubley & Green, 2013-2015; http://www.repeatmasker.org). We found that 28.7% of human liver CREs overlapped an annotated TE, most of which were SINEs (59.8% of the total) and LINEs (21.2%), although LTRs (9.3%) and DNA transposons (8.4%) were also abundant. 27 TE families were significantly enriched within the set of 47,673 orthologous CREs (FDR < 1%), nearly all of which were SINE-VNTR-*Alu*s (SVAs), LTRs, and *Alu* (Figure 5), suggesting these TEs contributed to the regulatory landscape in the primate liver (Jordan et al., 2003; Schmidt et al., 2012; Sundaram et al., 2014; Lynch et al., 2015).

The majority (75.0%) of the enriched TE families were relatively young, and specific to humans (SVA-F), Hominidae (SVA-B, SVA-C, and SVA-D), Hominoidea (the LTR12 subfamily), Simiiformes (LTRs), or primates (Alu elements), whereas the remaining 25.0% were Eutherian-specific or older (Fig. 5, Fig. S7, and Table S5). We therefore investigated whether these recent TE insertions altered the expression patterns of nearby genes in primates. Specifically, we focused on the enriched TE families younger than the Strepsirrhini/Haplorrhini divergence, thus not present in mouse lemur and bushbaby. We found that 22.6% of the CREs overlapping SVAs, and 12.5% of the CREs overlapping LTRs were differentially expressed between the Strepsirrhini (human, chimpanzee, rhesus macaque and marmoset) and the Haplorrhini (mouse lemur and bushbaby). Both of these fractions were significantly higher than expected by chance (binomial test $p < 2.2 \times 10^{-16}$). Together, these findings indicate that TEs have played a key role in shaping primate gene regulation, introducing novel gene expression patterns as a consequence of their recruitment as functional CREs.

The vast majority of newly evolved CREs are derived from TE insertions

84.2% of ape-specific CREs and 94.7% of human-specific CREs overlap at least one TE (Fig. 5; Fig. S7). In contrast, only 17.0% of evolutionarily conserved CREs contain an annotated TE (Fisher's exact test $p < 2.2 \times 10^{-16}$). However, we hypothesize that this may be an underestimate due to the inability to recognize ancient TE insertions accumulating mutations over time. LTRs and SVAs were the most frequently exapted TEs in newly evolved CREs (LTR = 40.1% of the exapted TEs in ape-specific CREs; SVA = 75.3% of the exapted TEs in the human-specific CREs; Fig. 5), despite being among the least common

Class

☐ LTR	☐ DNA	☐ SVA
☐ LINE	☐ SINE	

Lineage Specificity

☐ Metazoa	☐ Amniota/Euteleostomi?	☐ Amniota/Theria	☐ Eutheria+Metatheria	☐ Boreotheria	☐ Primates/Indep	☐ Catarrhini
☐ Vertebrata	☐ Amniota	☐ Theria	☐ Eutheria/Primates	☐ Euarchontoglires	☐ Primates	☐ Hominoidea
☐ Euteleostomi	☐ Mammalia	☐ Eutheria/Theria?	☐ Eutheria	☐ Euarchontoglires/Primates?	☐ Simiiformes	☐ Hominidae
☐ Tetrapoda	☐ Theria/Mammalia	☐ Eutheria/Theria	☐ Boreotheria/Eutheria?	☐ Primates/Glires	☐ Haplorrhini	☐ Human

Figure 5 - **Newly evolved CREs are enriched in TEs**. (A) Proportion of CREs that overlap TEs in the different primate lineages. (B) Number of enriched TE families within CREs in the different primate lineages. (C) Most enriched TE families in primates.

The contribution of LTRs in primate gene regulation has been characterized in previous studies (Wang et al., 2007; Cohen et al., 2009; Sundaram et al., 2014; Janoušek et al., 2016). In our data, a remarkable example of an ape-specific CRE derived from LTR insertion is an enhancer at the gene *GRIN3A*. This gene is involved in physiological and pathological processes in the central nervous system and has been associated with several complex human diseases, including schizophrenia (Takata et al., 2013). Our differential histone modification analysis identified an ape-specific ChIP-seq peak overlapping a 1-Kb long ape-specific insertion (present also in orangutan and gorilla, but not in other primates; GRCh38 chr9:101,723,127-101,724,197). This insertion, located 13 Kb from the TSS of *GRIN3A*, is entirely derived from an LTR-12C. The insertion drove strong enhancer activity upon transfection into HepG2 cells (Wilcoxon's rank sum test $p = 0.00017$; Fig. S5), suggesting that the transposable element was recruited as functional enhancer in the *GRIN3A* locus.

SVAs are instead a hominid-specific family of composite retrotransposons that are
active in humans (Hancks and Kazazian, 2010), with more than 3,500 annotated copies.
Given that nearly all human-specific liver CREs were derived from SVA insertions in our
analysis, we further investigated the genomic features of SVA insertions that lead to
exaptation (Table S5). 49.5% of human SVAs overlapped regions of significant histone
modification, and 97.8% of those were enhancers. These exapted SVAs are significantly
more likely to be associated with protein coding genes than the non-exapted SVAs (Fisher's
exact test p = 0.017) and are significantly closer to the TSS of the associated gene (mean
of 52.9 kb versus 64.1 kb; Wilcoxon rank-sum test $p < 2.2 \times 10^{-16}$). Exapted and non-exapted
SVAs lie within open chromatin regions in approximately the same number of cell types
(3.44 and 3.94 respectively; logistic regression p = 0.827) and host on average a
comparable density of TFBSs (3.28 and 1.95; logistic regression p = 0.679). However,
exapted SVAs have a significantly higher number of TFBSs in the neighboring regions (8.12
versus 5.86 in +/- 10kb; logistic regression $p = 1.91 \times 10^{-5}$). Taken together, these data
suggest a model where an SVA has a higher probability of becoming a CRE if it inserts in
TFBS-dense regions near protein coding gene promoters. (Fig. 6).

Among the exapted SVAs, our data predicted an intronic CRE for the gene *JARID2.*
This gene is an accessory component of Polycomb Repressive Complex-2 (PRC2), recruits
PRC2 to chromatin, and is involved in liver, brain, neural tube development, and embryonic
stem cell differentiation (Kaneko et al., 2014). Our differential histone state analysis
revealed a human-specific ChIP-seq peak overlapping a human-specific 1.9 kb-long
insertion, entirely derived from an SVA-F retrotransposon. JARID2 is significantly
downregulated in humans compared to all the other primates (Benjamini–Hochberg p =
0.019). Exapted SVAs-Fs exhibit significant enrichment for binding sites of known
transcriptional repressors such as PAX-5, FEV, and SREBF1 (Maurer et al., 2003; Fazio et
al., 2008; Lecomte et al., 2010). Indeed, the JARID2 SVA-F insertion leads to significantly
decreased expression in HepG2 reporter assays (Wilcoxon's rank sum test p = 0.00275;
Fig. S6), supporting the role of this SVA-F as a transcriptional repressor.
402

Figure 6 - **Model for exaptation of SVAs into human functional CREs**. A given SVA has a higher probability of being recruited as a functional cis-regulatory element if it is found within 50 kb from a protein coding gene, in a genomic region already enriched in TF binding motifs in the surrounding 10 kb up- and downstream.

Broad regulatory activity of TE insertions in the primate liver

Our findings strongly suggest that the majority of novel CREs in primates is derived from TE insertions. To validate the predicted regulatory activity of recent TE insertions, we tested the cis-regulatory activity of 69 TE subfamilies, covering all of the main classes and families of TEs (Table S6). TEs from these families overlap 3,897 of our predicted CREs. We synthesized the mammalian consensus sequence (see experimental procedures) for 69 different TE families, cloned them into a luciferase reporter vector with a minimal promoter, and transfected them into HepG2 cells to perform dual luciferase reporter assays. We found that luciferase expression for 66 of the 69 (95.6%) tested TE families was significantly different from the negative control (Fig. 7a; Wilcoxon's rank sum test p-values in Table S6). Strikingly, only 17 (25.7%) of these, mostly LTRs and DNA transposons, produced activity

significantly higher than the negative control (Fig. 7a), whereas the remaining 49 (74.3%), mostly LINEs, repressed transcription. Consistent with results from the JARID2 locus presented above, SVA-Fs were confirmed to function as transcriptional repressors.

Figure 7 - **Regulatory ability of TE families found in poised and active regulatory elements in HepG2 cells.** (A) The p-value, class, and lineage specificity for the 69 TE families tested in HepG2 cells for regulatory ability, the empty vector control (Basic[minP]), and the positive control (TAP2_C) are all shown above the 6 luciferase assay replicates conducted and the average regulatory ability found across the replicates. Red indicates luciferase expression higher than the empty vector control; blue indicates luciferase expression less than Basic[minP]. (B) CTCF, FOX, USF2, GABP, and HNF4a binding motifs were found to be enriched in the sequences of the 66 TE families that drive expression significantly different from background. The top row shows the enriched sequence found while the bottom shows the Jaspar binding motifs recognized for each transcription factor.

These findings support that LTRs, among the most enriched TEs in our peak set, and the most common exapted TEs in apes, are likely co-opted as active enhancer elements. The putative repressor activity of LINEs is consistent with their underrepresentation in the human ChIP-seq peak set. However, we hypothesize that, at least for some of the TE families, such observed repressing activity levels could be the

byproduct of secondary biological mechanisms leading the cell to recognize these elements
as newly inserted TEs, and therefore silencing them via epigenetic mechanisms. However,
further tests will be needed to support this hypothesis.

The consensus sequences for the 66 TEs that drove reporter expression significantly
different from background were analyzed with MEME to identify enriched motifs. Motifs for
known master regulators of liver cell identity, including FOX, USF2, GABP, and HNF4A
(Wallerman et al., 2009), were significantly enriched within the sequences of the 66 TE
families with significant regulatory activity (Fig. 7b). In summary, most TE families function
as CREs in the primate liver, either as strong enhancers or as repressors. TEs are actively
recruited into the regulatory landscape, further supporting our findings on the pervasive
involvement of TEs in the primate gene regulation.

Discussion

The primate regulatory landscape is evolutionarily conserved
Only a small fraction (<1.50%) of the CREs were differentially active between humans and
chimpanzees. This suggests that even modest changes in gene regulation produce
observable phenotypic differentiation, and confirm that cis-regulatory evolution plays a
central role in primate diversification (Davidson 2001, 2006; Wray, 2007; Ho et al., 2009;
Tsankov et al., 2010; Smith et al., 2013; Martin et al., 2012; Coolon et al., 2014; Martin and
Reed, 2014; Guo et al., 2015; Lynch et al., 2015; Villar et al., 2015; Adachi et al., 2016;
Landeen et al., 2016; Lesch et al., 2016; Zhang and Reed, 2016). Our approach for the
comparison of CREs across species, based on the analysis of differential histone
modification state in orthologous regions, demonstrated that cis-regulatory divergence
across species may be overestimated when based on the peak overlap status as a binary
variable.

Specific genomic features are associated with CRE conservation
Evolutionarily conserved promoters and enhancers have conserved nucleotide sequence,
are close to protein coding genes, are functional in many cell types, and harbor many
TFBSs. Many regulatory pathways, specifically those involved in the regulation of liver
function and housekeeping functions, are strongly conserved across primates, while other
pathways, such as immune response, are less constrained and evolve more rapidly. This

observation is consistent with the expected arms-race in evolution between host and pathogens.

Newly evolved CREs are derived from TE exaptation

Based on our findings, exaptation of transposable elements into functional promoters and enhancers is a pervasive phenomenon in primate genomes. LTRs and SVAs are the most frequently exapted in humans and other apes, despite not being among the most common transposable elements in the genome. Primate liver CREs are enriched in young TEs. These young TEs, after being recruited into the primate regulatory network, introduced novel gene expression in the associated species. To our knowledge, this is the first study demonstrating how specific genomic features are associated to the recruitment of TEs as functional elements in the primate regulatory landscape. In contrast, only a minor fraction of the evolutionarily conserved CREs overlap an annotated TE. This suggests the action of strong selection against the disruption of these regulatory elements, in order to maintain stable gene expression. Further, these data suggest that the core regulatory network that establishes liver cell-type identity is very conserved, whereas adaptive evolution occurs on the periphery of the network, where TEs have the most impact on gene regulatory evolution.

Data availability

All non-human raw sequence data have been deposited in the Sequence Read Archive under following BioProject IDs: PRJNA349047 for RNA-seq and PRJNA349046 for ChIP-seq data.

Supplemental information

Supplemental can be found with this article online at **xxxxxxxx**

Author contributions

MT, MC and CDB conceived the project. MT, YP, GHP and CDB designed the taxon sampling and experiments. MT performed ChIP-seq and RNA-seq experiments. MT and MHB performed the parallelized reporter assays. MHB and KA produced luciferase assay data on *GRIN3A* and *JARID2*. KM and VJL designed and performed the luciferase assays on the 69 TE families. MT, VJL and KM performed the TE enrichment analysis. YP designed computational pipelines for the detection of orthologous regions in the Ensembl

MSA alignment and all related analyses. MT, YP and CDB analyzed the data and wrote the

paper. All authors read and approved the manuscript.

Acknowledgements

We thank Texas Biomedical Research Institute, Duke University Lemur Center and the

Hospital of University of Pennsylvania for providing the samples. We thank Jonathan Schug

and other members of UPenn NGS Core Facility. We are grateful to Sarah Tishkoff, Yoseph

Barash, Benjamin Voight, Yana Kamberov, Erin Fry, and Barbara Engelhardt for providing

insightful comments on the preliminary versions of the results and of the manuscript.

References

References

Adachi, N., Robinson, M., Goolsbee, A., and Shubin, N.H. (2016). Regulatory evolution of
Tbx5 and the origin of paired appendages. Proceedings of the National Academy of
Sciences.

Andersson, R., Gebhard, C., Miguel-Escalada, I., Hoof, I., Bornholdt, J., Boyd, M., Chen, Y.,
Zhao, X., Schmidl, C., Suzuki, T., et al. (2014). An atlas of active enhancers across human
cell types and tissues. Nature *507*, 455–461.

Austenaa, L.M.I., Barozzi, I., Simonatto, M., Masella, S., Della Chiara, G., Ghisletti, S.,
Curina, A., de Wit, E., Bouwman, B.A.M., de Pretis, S., et al. Transcription of Mammalian
cis-Regulatory Elements Is Restrained by Actively Enforced Early Termination. Molecular
Cell *60*, 460–474.

Babbitt, C.C., Fedrigo, O., Pfefferle, A.D., Boyle, A.P., Horvath, J.E., Furey, T.S., and Wray,
G.A. (2010). Both Noncoding and Protein-Coding RNAs Contribute to Gene Expression
Evolution in the Primate Brain. Genome Biology and Evolution *2*, 67–79.

Bailey, T.L. (2011). DREME: motif discovery in transcription factor ChIP-seq data.
Bioinformatics *27*, 1653–1659.

Bailey, T.L., Boden, M., Buske, F.A., Frith, M., Grant, C.E., Clementi, L., Ren, J., Li, W.W.,
and Noble, W.S. (2009). MEME SUITE: tools for motif discovery and searching. Nucleic
Acids Res *37*, W202-208.

Ballester, B., Medina-Rivera, A., Schmidt, D., Gonzàlez-Porta, M., Carlucci, M., Chen, X.,
Chessman, K., Faure, A.J., Funnell, A.P., Goncalves, A., et al. (2014). Multi-species, multi-
transcription factor binding highlights conserved control of tissue-specific biological
pathways. eLife *3*, e02626.

Barnabas, S., Hai, T., and Andrisani, O.M. (1997). The Hepatitis B Virus X Protein
Enhances the DNA Binding Potential and Transcription Efficacy of bZip Transcription
Factors. Journal of Biological Chemistry *272*, 20684–20690.

Bejerano, G., Lowe, C.B., Ahituv, N., King, B., Siepel, A., Salama, S.R., Rubin, E.M., James
Kent, W., and Haussler, D. (2006). A distal enhancer and an ultraconserved exon are
derived from a novel retroposon. Nature *441*, 87–90.

Bourque, G., Leong, B., Vega, V.B., Chen, X., Lee, Y.L., Srinivasan, K.G., Chew, J.-L.,
Ruan, Y., Wei, C.-L., Ng, H.H., et al. (2008). Evolution of the mammalian transcription factor
binding repertoire via transposable elements. Genome Research.

Boyd, J.L., Skove, S.L., Rouanet, J.P., Pilaz, L.-J., Bepler, T., Gordân, R., Wray, G.A., and
Silver, D.L. Human-Chimpanzee Differences in a FZD8 Enhancer Alter Cell-Cycle
Dynamics in the Developing Neocortex. Current Biology *25*, 772–779.

Boyle, A.P., Araya, C.L., Brdlik, C., Cayting, P., Cheng, C., Cheng, Y., Gardner, K., Hillier,
L.W., Janette, J., Jiang, L., et al. (2014). Comparative analysis of regulatory information and
circuits across distant species. Nature *512*, 453–456.

Brawand, D., Soumillon, M., Necsulea, A., Julien, P., Csardi, G., Harrigan, P., Weier, M.,
Liechti, A., Aximu-Petri, A., Kircher, M., et al. (2011). The evolution of gene expression
levels in mammalian organs. Nature *478*, 343–348.

Breschi, A., Djebali, S., Gillis, J., Pervouchine, D.D., Dobin, A., Davis, C.A., Gingeras, T.R.,
and Guigó, R. (2016). Gene-specific patterns of expression variation across organs and
species. Genome Biology *17*, 1–13.

Britten, R.J., and Davidson, E.H. (1969). Gene Regulation for Higher Cells: A Theory.
Science *165*, 349–357.

Brosius, J., Gould, S.J. (1992). On "genomenclature": a comprehensive (and respectful)
taxonomy for pseudogenes and other "junk DNA.". Proceedings of the National Academy of
Sciences USA *89*, 10706–10710.

Brown, C.D., Johnson, D.S., and Sidow, A. (2007). Functional Architecture and Evolution of
Transcriptional Elements That Drive Gene Coexpression. Science *317*, 1557.

Cain, C.E., Blekhman, R., Marioni, J.C., and Gilad, Y. (2011). Gene Expression Differences
Among Primates Are Associated With Changes in a Histone Epigenetic Modification.
Genetics *187*, 1225–1234.

Chuong, E.B., Rumi, M.A.K., Soares, M.J., and Baker, J.C. (2013). Endogenous
retroviruses function as species-specific enhancer elements in the placenta. Nat Genet *45*,
325–329.

Chuong, E.B., Elde, N.C., and Feschotte, C. (2016). Regulatory evolution of innate
immunity through co-option of endogenous retroviruses. Science *351*, 1083–1087.

Cohen, C.J., Lock, W.M., and Mager, D.L. (2009). Endogenous retroviral LTRs as promoters for human genes: A critical assessment. Gene *448*, 105–114.

Coolon, J.D., McManus, C.J., Stevenson, K.R., Graveley, B.R., and Wittkopp, P.J. (2014). Tempo and mode of regulatory evolution in Drosophila. Genome Research.

Cooper, G.M., and Brown, C.D. (2008). Qualifying the relationship between sequence conservation and molecular function. Genome Research *18*, 201–205.

Core, L.J., Martins, A.L., Danko, C.G., Waters, C.T., Siepel, A., and Lis, J.T. (2014). Analysis of nascent RNA identifies a unified architecture of initiation regions at mammalian promoters and enhancers. Nat Genet *46*, 1311–1320.

Cotney, J., Leng, J., Yin, J., Reilly, S.K., DeMare, L.E., Emera, D., Ayoub, A.E., Rakic, P., and Noonan, J.P. The Evolution of Lineage-Specific Regulatory Activities in the Human Embryonic Limb. Cell *154*, 185–196.

Creyghton, M.P., Cheng, A.W., Welstead, G.G., Kooistra, T., Carey, B.W., Steine, E.J., Hanna, J., Lodato, M.A., Frampton, G.M., Sharp, P.A., et al. (2010). Histone H3K27ac separates active from poised enhancers and predicts developmental state. Proceedings of the National Academy of Sciences *107*, 21931–21936.

Cusanovich, D.A., Pavlovic, B., Pritchard, J.K., and Gilad, Y. (2014). The Functional Consequences of Variation in Transcription Factor Binding. PLoS Genet *10*, e1004226.

Davidson, E., and Britten, R. (1979). Regulation of gene expression: possible role of repetitive sequences. Science *204*, 1052–1059.

Degner, J.F., Pai, A.A., Pique-Regi, R., Veyrieras, J.B., Gaffney, D.J., et al. (2012). DNaseI sensitivity QTLs are a major determinant of human expression variation. Nature, *482*, 390–394.

DeLaForest, A., Nagaoka, M., Si-Tayeb, K., Noto, F.K., Konopka, G., Battle, M.A., and Duncan, S.A. (2011). HNF4A is essential for specification of hepatic progenitors from human pluripotent stem cells. Development *138*, 4143.

De Souza, F.S.J., Franchini, L.F., and Rubinstein, M. (2013). Exaptation of Transposable Elements into Novel Cis-Regulatory Elements: Is the Evidence Always Strong? Molecular Biology and Evolution *30*(6), 1239-1251.

Dong, X., Wang, X., Zhang, F., and Tian, W. (2016). Genome-Wide Identification of
Regulatory Sequences Undergoing Accelerated Evolution in the Human Genome. Mol Biol
Evol *33*, 2565–2575.

Du, J., Leung, A., Trac, C., Lee, M., Parks, B.W., Lusis, A.J., Natarajan, R., and Schones,
D.E. (2016). Chromatin variation associated with liver metabolism is mediated by
transposable elements. Epigenetics Chromatin *9*, 28.

Eden, E., Lipson, D., Yogev, S., and Yakhini, Z. (2007). Discovering Motifs in Ranked Lists
of DNA Sequences. PLoS Comput Biol *3*, e39.

Eden, E., Navon, R., Steinfeld, I., Lipson, D., and Yakhini, Z. (2009). GOrilla: a tool for
discovery and visualization of enriched GO terms in ranked gene lists. BMC Bioinformatics
10, 1–7.

Emera, D., Yin, J., Reilly, S.K., Gockley, J., and Noonan, J.P. (2016). Origin and evolution
of developmental enhancers in the mammalian neocortex. Proceedings of the National
Academy of Sciences *113*, E2617–E2626.

ENCODE Project Consortium (2012). An integrated encyclopedia of DNA elements in the
human genome. Nature *489*, 57–74.

Fazio, G., Palmi, C., Rolink, A., Biondi, A., Cazzaniga, G. (2008). PAX5/TEL acts as a
transcriptional repressor causing down-modulation of CD19, enhances migration to
CXCL12, and confers survival advantage in pre-BI cells. Cancer Research *68*(1), 181–189.

FANTOM Consortium and the RIKEN PMI and CLST (DGT) (2014). A promoter-level
mammalian expression atlas. Nature *507*, 462–470.

Gittelman, R.M., Hun, E., Ay, F., Madeoy, J., Pennacchio, L., Noble, W.S., Hawkins, R.D.,
and Akey, J.M. (2015). Comprehensive identification and analysis of human accelerated
regulatory DNA. Genome Res *25*, 1245–1255.

Gross, L. (2005). Selection on a Neural Gene Regulator Sheds Light on Human Evolution.
PLoS Biol *3*, e417.

Grueber, C.E., Wallis, G.P., and Jamieson, I.G. (2014). Episodic Positive Selection in the
Evolution of Avian Toll-Like Receptor Innate Immunity Genes. PLoS ONE *9*, e89632.

Guo, C., Ludvik, A.E., Arlotto, M.E., Hayes, M.G., Armstrong, L.L., Scholtens, D.M., Brown,
C.D., Newgard, C.B., Becker, T.C., Layden, B.T., et al. (2015). Coordinated regulatory

variation associated with gestational hyperglycaemia regulates expression of the novel
hexokinase HKDC1. Nat Commun *6*.

Hah, N., Murakami, S., Nagari, A., Danko, C.G., and Kraus, W.L. (2013). Enhancer
transcripts mark active estrogen receptor binding sites. Genome Research *23*, 1210–1223.

Hancks, D.C., and Kazazian Jr., H.H. (2010). SVA retrotransposons: Evolution and genetic
instability. Seminars in Cancer Biology *20*, 234–245.

Ho, M.C.W., Johnsen, H., Goetz, S.E., Schiller, B.J., Bae, E., Tran, D.A., Shur, A.S., Allen,
J.M., Rau, C., Bender, W., et al. (2009). Functional Evolution of cis-Regulatory Modules at
a Homeotic Gene in Drosophila. PLoS Genet *5*, e1000709.

Holloway, A.K., Bruneau, B.G., Sukonnik, T., Rubenstein, J.L., and Pollard, K.S. (2016).
Accelerated Evolution of Enhancer Hotspots in the Mammal Ancestor. Molecular Biology
and Evolution *33*, 1008–1018.

Innocenti, F., Cooper, G.M., Stanaway, I.B., Gamazon, E.R., Smith, J.D., Mirkov, S.,
Ramirez, J., Liu, W., Lin, Y.S., Moloney, C., et al. (2011). Identification, Replication, and
Functional Fine-Mapping of Expression Quantitative Trait Loci in Primary Human Liver
Tissue. PLoS Genet *7*, e1002078.

Jain, D., Baldi, S., Zabel, A., Straub, T., and Becker, P.B. (2015). Active promoters give rise
to false positive "Phantom Peaks" in ChIP-seq experiments. Nucleic Acids Res *43*, 6959–
6968.

Jansa, A.S., Lundrigan, L.B., and Tucker, K.P. (2003). Tests for Positive Selection on
Immune and Reproductive Genes in Closely Related Species of the Murine Genus Mus.
Journal of Molecular Evolution *56*, 294–307.

Jordan, I.K., Rogozin, I.B., Glazko, G.V., and Koonin, E.V. (2003). Origin of a substantial
fraction of human regulatory sequences from transposable elements. Trends in Genetics
19, 68–72.

Kaneko, S., Bonasio, R., Saldaña-Meyer, R., Yoshida, T., Son, J., Nishino, K., Umezawa,
A., and Reinberg, D. Interactions between JARID2 and Noncoding RNAs Regulate PRC2
Recruitment to Chromatin. Molecular Cell *53*, 290–300.

Kapusta, A., Kronenberg, Z., Lynch, V.J., Zhuo, X., Ramsay, L., Bourque, G., Yandell, M.,
and Feschotte, C. (2013). Transposable Elements Are Major Contributors to the Origin,

Diversification, and Regulation of Vertebrate Long Noncoding RNAs. PLoS Genet *9*,
e1003470.

King, M., and Wilson, A. (1975). Evolution at two levels in humans and chimpanzees.
Science *188*, 107–116.

Kunarso, G., Chia, N.-Y., Jeyakani, J., Hwang, C., Lu, X., Chan, Y.-S., Ng, H.-H., and
Bourque, G. (2010). Transposable elements have rewired the core regulatory network of
human embryonic stem cells. Nat Genet *42*, 631–634.

Landeen, E.L., Muirhead, C.A., Wright, L., Meiklejohn, C.D., and Presgraves, D.C. (2016).
Sex Chromosome-wide Transcriptional Suppression and Compensatory Cis-Regulatory
Evolution Mediate Gene Expression in the Drosophila Male Germline. PLoS Biol *14*,
e1002499.

Lazzaro, B.P., and Schneider, D.S. (2014). The Genetics of Immunity. G3:
Genes|Genomes|Genetics *4*, 943–945.

Lecomte, V., Maugnier, E., Euthine, V., Durand, C., Freyssenet, D., Nemoz, G., Rome, S.,
Vidal, H. and Lefai, E. (2010). A new role for sterol regulatory element binding protein 1
transcription factors in the regulation of muscle mass and muscle cell differentiation.
Molecular and Cell Biology *30*(5), 1182–1189.

Lesch, B.J., Silber, S.J., McCarrey, J.R., and Page, D.C. (2016). Parallel evolution of male
germline epigenetic poising and somatic development in animals. Nat Genet *48*, 888–894.

Lewis, J.J., van der Burg, K.R.L., Mazo-Vargas, A., and Reed, R.D. ChIP-Seq-Annotated
Heliconius erato Genome Highlights Patterns of cis-Regulatory Evolution in Lepidoptera.
Cell Reports *16*, 2855–2863.

Li, H. (2013) Aligning sequence reads, clone sequences and assembly contigs with BWA-
MEM. arXiv:1303.3997v1

Li, W., Notani, D., and Rosenfeld, M.G. (2016). Enhancers as non-coding RNA transcription
units: recent insights and future perspectives. Nat Rev Genet *17*, 207–223.

Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general purpose
program for assigning sequence reads to genomic features. Bioinformatics *30*, 923–930.

Loisel, D.A., Rockman, M.V., Wray, G.A., Altmann, J., and Alberts, S.C. (2006). Ancient
polymorphism and functional variation in the primate MHC-DQA1 *5′* cis-regulatory region.
Proceedings of the National Academy of Sciences *103*, 16331–16336.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biology *15*, 1–21.

Lynch, V.J., Leclerc, R.D., May, G., and Wagner, G.P. (2011). Transposon-mediated
rewiring of gene regulatory networks contributed to the evolution of pregnancy in mammals.
Nat Genet *43*, 1154–1159.

Lynch, V.J., Nnamani, M.C., Kapusta, A., Brayer, K., Plaza, S.L., Mazur, E.C., Emera, D.,
Sheikh, S.Z., Grützner, F., Bauersachs, S., et al. (2015) Ancient Transposable Elements
Transformed the Uterine Regulatory Landscape and Transcriptome during the Evolution of
Mammalian Pregnancy. Cell Reports *10*, 551–561.

Markljung, E., Jiang, L., Jaffe, J.D., Mikkelsen, T.S., Wallerman, O., Larhammar, M., Zhang,
X., Wang, L., Saenz-Vash, V., Gnirke, A., et al. (2009). ZBED6, a Novel Transcription
Factor Derived from a Domesticated DNA Transposon Regulates IGF2 Expression and
Muscle Growth. PLoS Biol *7*, e1000256.

Marnetto, D., Molineris, I., Grassi, E., and Provero, P. (2014). Genome-wide Identification
and Characterization of Fixed Human-Specific Regulatory Regions. American Journal of
Human Genetics *95*, 39–48.

Martin, A., and Reed, R.D. (2014). Wnt signaling underlies evolution and development of
the butterfly wing pattern symmetry systems. Developmental Biology *395*, 367–378.

Martin, A., Papa, R., Nadeau, N.J., Hill, R.I., Counterman, B.A., Halder, G., Jiggins, C.D.,
Kronforst, M.R., Long, A.D., McMillan, W.O., et al. (2012). Diversification of complex
butterfly wing patterns by repeated regulatory evolution of a Wnt ligand. Proceedings of the
National Academy of Sciences *109*, 12632–12637.

Maurano, M.T., Humbert, R., Rynes, E., Thurman, R.E., Haugen, E., et al. (2012).
Systematic Localization of Common Disease-Associated Variation in Regulatory DNA.
Science, *337*, 1190–1195.

Maurer, P., Tsas, F., Coutte, L., Callens, N., Brenner, C., Van Lint, C., de Launoit, Y., Baert,
JL (2003). FEV acts as a transcriptional repressor through its DNA-binding ETS domain
and alanine-rich domain. Oncogene *22*(21), 3319–3329.

McClintock, B. (1950). The Origin and Behavior of Mutable Loci in Maize. Proceedings of
the National Academy of Sciences of the United States of America *36*, 344–355.

McClintock, B. (1984). The significance of responses of the genome to challenge. Science
226, 792–801.

Melnikov, A., Murugan, A., Zhang, X., Tesileanu, T., Wang, L., Rogov, P., Feizi, S., Gnirke,
A., Callan, C.G., Kinney, J.B., et al. (2012). Systematic dissection and optimization of
inducible enhancers in human cells using a massively parallel reporter assay. Nat Biotech
30, 271–277.

Musunuru, K., Strong, A., Frank-Kamenetsky, M., Lee, N.E., Ahfeldt, T., Sachs, K.V., Li, X.,
Li, H., Kuperwasser, N., Ruda, V.M., et al. (2010). From noncoding variant to phenotype via
SORT1 at the 1p13 cholesterol locus. Nature, *466* (7307), 714–719.

Patwardhan, R.P., Hiatt, J.B., Witten, D.M., Kim, M.J., Smith, R.P., May, D., Lee, C.,
Andrie, J.M., Lee, S.-I., Cooper, G.M., et al. (2012). Massively parallel functional dissection
of mammalian enhancers in vivo. Nat Biotech *30*, 265–270.

Obbard, D.J., Welch, J.J., Kim, K.-W., and Jiggins, F.M. (2009). Quantifying Adaptive
Evolution in the Drosophila Immune System. PLoS Genet *5*, e1000698.

Ori, A., Atzmony, D., Haviv, I., and Shaul, Y. (1994). An NF1 Motif Plays a Central Role in
Hepatitis B Virus Enhancer. Virology *204*, 600–608.

Perelman, P., Johnson, W.E., Roos, C., Seuánez, H.N., Horvath, J.E., Moreira, M.A.M.,
Kessing, B., Pontius, J., Roelke, M., Rumpler, Y., et al. (2011). A Molecular Phylogeny of
Living Primates. PLoS Genet *7*, e1001342.

Pickrell, J.K., Gaffney, D.J., Gilad, Y., and Pritchard, J.K. (2011). False positive peaks in
ChIP-seq and other sequencing-based functional assays caused by unannotated high copy
number regions. Bioinformatics *27*, 2144–2146.

Pollard, K.S., Salama, S.R., King, B., Kern, A.D., Dreszer, T., Katzman, S., Siepel, A.,
Pedersen, J.S., Bejerano, G., Baertsch, R., et al. (2006). Forces Shaping the Fastest
Evolving Regions in the Human Genome. PLoS Genet *2*, e168.

Pollard, K.S., Hubisz, M.J., Rosenbloom, K.R., and Siepel, A. (2010). Detection of
nonneutral substitution rates on mammalian phylogenies. Genome Research *20*, 110–121.

Prabhakar, S., Visel, A., Akiyama, J.A., Shoukry, M., Lewis, K.D., Holt, A., Plajzer-Frick, I.,
Morrison, H., FitzPatrick, D.R., Afzal, V., et al. (2008). Human-Specific Gain of Function in a
Developmental Enhancer. Science *321*, 1346–1350.

Prescott, S.L., Srinivasan, R., Marchetto, M.C., Grishina, I., Narvaiza, I., Selleri, L., Gage,
F.H., Swigut, T., and Wysocka, J. Enhancer Divergence and cis-Regulatory Evolution in the
Human and chimpanzee Neural Crest. Cell *163*, 68–83.

Rayan, N.A., Del Rosario, R.C.H., and Prabhakar, S. (2016). Massive contribution of
transposable elements to mammalian regulatory sequences. Semin Cell Dev Biol *57*, 51–
56.

Reilly, S.K., Yin, J., Ayoub, A.E., Emera, D., Leng, J., Cotney, J., Sarro, R., Rakic, P., and
Noonan, J.P. (2015). Evolutionary changes in promoter and enhancer activity during human
corticogenesis. Science *347*, 1155–1159.

del Rosario, R.C.H., Rayan, N.A., and Prabhakar, S. (2014). Noncoding origins of
anthropoid traits and a new null model of transposon functionalization. Genome Research
24, 1469–1484.

Sackton, T.B., Lazzaro, B.P., Schlenke, T.A., Evans, J.D., Hultmark, D., and Clark, A.G.
(2007). Dynamic evolution of the innate immune system in Drosophila. Nat Genet *39*,
1461–1468.

Salazar-Jaramillo, L., Paspati, A., van de Zande, L., Vermeulen, C.J., Schwander, T., and
Wertheim, B. (2014). Evolution of a cellular immune response in Drosophila: a phenotypic
and genomic comparative analysis. Genome Biol Evol *6*, 273–289.

Sasaki, T., Nishihara, H., Hirakawa, M., Fujimura, K., Tanaka, M., Kokubo, N., Kimura-
Yoshida, C., Matsuo, I., Sumiyama, K., Saitou, N., et al. (2008). Possible involvement of
SINEs in mammalian-specific brain formation. Proceedings of the National Academy of
Sciences *105*, 4220–4225.

Schadt, E.E. (2009). Molecular networks as sensors and drivers of common human
diseases. Nature *461*, 218–223.

Schmidt, D., Wilson, M.D., Ballester, B., Schwalie, P.C., Brown, G.D., Marshall, A., Kutter,
C., Watt, S., Martinez-Jimenez, C.P., Mackay, S., et al. (2010). Five-Vertebrate ChIP-seq
Reveals the Evolutionary Dynamics of Transcription Factor Binding. Science *328*, 1036.

Schmidt, D., Schwalie, P.C., Wilson, M.D., Ballester, B., Gonçalves, Â., Kutter, C., Brown,
G.D., Marshall, A., Flicek, P., and Odom, D.T. (2012). Waves of Retrotransposon
Expansion Remodel Genome Organization and CTCF Binding in Multiple Mammalian
Lineages. Cell *148*, 335–348.

Sharon, E., Kalma, Y., Sharp, A., Raveh-Sadka, T., Levo, M., Zeevi, D., Keren, L., Yakhini,
Z., Weinberger, A., and Segal, E. (2012). Inferring gene regulatory logic from high-
throughput measurements of thousands of systematically designed promoters. Nat Biotech
30, 521–530.

Siepel, A., Bejerano, G., Pedersen, J.S., Hinrichs, A.S., Hou, M., Rosenbloom, K., Clawson,
H., Spieth, J., Hillier, L.W., Richards, S., et al. (2005). Evolutionarily conserved elements in
vertebrate, insect, worm, and yeast genomes. Genome Research *15*, 1034–1050.

Sironi, M., Cagliani, R., Forni, D., and Clerici, M. (2015). Evolutionary insights into host-
pathogen interactions from mammalian sequence data. Nat Rev Genet *16*, 224–236.

Smedley, D., Haider, S., Durinck, S., Pandini, L., Provero, P., Allen, J., Arnaiz, O., Awedh,
M.H., Baldock, R., Barbiera, G., et al. (2015). The BioMart community portal: an innovative
alternative to large, centralized data repositories. Nucleic Acids Research *43*, W589–W598.

Sundaram, V., Cheng, Y., Ma, Z., Li, D., Xing, X., Edge, P., Snyder, M.P., and Wang, T.
(2014). Widespread contribution of transposable elements to the innovation of gene
regulatory networks. Genome Res *24*, 1963–1976.

Takata, A., Iwayama, Y., Fukuo, Y., Ikeda, M., Okochi, T., Maekawa, M., Toyota, T.,
Yamada, K., Hattori, E., Ohnishi, T., et al. A Population-Specific Uncommon Variant in
GRIN3A Associated with Schizophrenia. Biological Psychiatry *73*, 532–539.

Tsankov, A.M., Thompson, D.A., Socha, A., Regev, A., and Rando, O.J. (2010). The Role
of Nucleosome Positioning in the Evolution of Gene Regulation. PLoS Biol *8*, e1000414.

Vallender, E.J., and Lahn, B.T. (2004). Positive selection on the human genome. Human
Molecular Genetics *13*, R245–R254.

Villar, D., Berthelot, C., Aldridge, S., Rayner, T.F., Lukk, M., Pignatelli, M., Park, T.J.,
Deaville, R., Erichsen, J.T., Jasinska, A.J., et al. Enhancer Evolution across 20 Mammalian
Species. Cell *160*, 554–566.

Wallerman, O., Motallebipour, M., Enroth, A., Patra, K., Bysani, M.S.R., Komorowski, J.,
and Wadelius, C. (2009). Molecular interactions between HNF4a, FOXA2 and GABP
identified at regulatory DNA elements through ChIP-sequencing. Nucleid Acids Research
37, 7498–7508.

Wang, T., Zeng, J., Lowe, C.B., Sellers, R.G., Salama, S.R., Yang, M., Burgess, S.M.,
Brachmann, R.K., and Haussler, D. (2007). Species-specific endogenous retroviruses

shape the transcriptional network of the human tumor suppressor protein p53. Proceedings
of the National Academy of Sciences *104*, 18613–18618.

Warner, L.R., Babbitt, C.C., Primus, A.E., Severson, T.F., Haygood, R., and Wray, G.A.
(2009). Functional consequences of genetic variation in primates on tyrosine hydroxylase
(TH) expression in vitro. Brain Research *1288*, 1–8.

Wertheim, B. (2015). Genomic basis of evolutionary change: evolving immunity. Frontiers in
Genetics *6*, 222.

Wray, G.A. (2007). The evolutionary significance of cis-regulatory mutations. Nat Rev
Genet *8*, 206–216.

Xie, M., Hong, C., Zhang, B., Lowdon, R.F., Xing, X., Li, D., Zhou, X., Lee, H.J., Maire, C.L.,
Ligon, K.L., et al. (2013). DNA hypomethylation within specific transposable element
families associates with tissue-specific enhancer landscape. Nat Genet *45*, 836–841.

Yang, S., Oksenberg, N., Takayama, S., Heo, S.-J., Poliakov, A., Ahituv, N., Dubchak, I.,
and Boffelli, D. (2015). Functionally conserved enhancers with divergent sequences in
distant vertebrates. BMC Genomics *16*, 1–13.

Yates, A., Beal, K., Keenan, S., McLaren, W., Pignatelli, M., Ritchie, G.R.S., Ruffier, M.,
Taylor, K., Vullo, A., and Flicek, P. (2015). The Ensembl REST API: Ensembl Data for Any
Language. Bioinformatics *31*, 143–145.

Young, R.S., Kumar, Y., Bickmore, W.A., and Taylor, M.S. (2016). Bidirectional
transcription marks accessible chromatin and is not specific to enhancers. bioRxiv.

Zak, D.E., Tam, V.C., and Aderem, A. (2014). Systems-Level Analysis of Innate Immunity.
Annu. Rev. Immunol. *32*, 547–577.

Zhang, L., and Reed, R.D. (2016). Genome editing in butterflies reveals that spalt promotes
and Distal-less represses eyespot colour patterns. Nat Commun *7*.

Zhang, Y., Liu, T., Meyer, C.A., Eeckhoute, J., Johnson, D.S., Bernstein, B.E., Nusbaum,
C., Myers, R.M., Brown, M., Li, W., et al. (2008). Model-based Analysis of ChIP-Seq
(MACS). Genome Biology *9*, 1–9.

Zhou, X., Cain, C.E., Myrthil, M., Lewellen, N., Michelini, K., Davenport, E.R., Stephens, M.,
Pritchard, J.K., and Gilad, Y. (2014). Epigenetic modifications are associated with inter-
species gene expression variation in primates. Genome Biology *15*, 1–19.

Materials and Methods

Tissue sampling

We obtained liver tissue samples for three to four individuals belonging to each of the studied primate species (three bushbabies, four chimpanzees, three humans, three marmosets, three mouse lemurs, and three rhesus macaques; Table S1). Samples for chimpanzees, marmoset and rhesus macaque were obtained from Texas Biomedical Research Institute (San Antonio, TX); bushbaby and mouse lemur livers were obtained from the Lemur Center of Duke University (Durham, NC). Tissue samples were collected and flash-frozen immediately. With the exception of the bushbaby, samples for all of the species included both males and females (Table S1). Age was comparable across species (young adults) and all individuals died of causes unrelated to liver disease.

RNA-seq sample processing

We processed samples from all species in random batches of four in order to minimize batch effects. For each sample, 25 mg of frozen liver tissue was used to extract total RNA and genomic DNA, using QIAGEN AllPrep DNA/RNA/miRNA Universal Kit. Quality of total RNA was assessed computing the RNA Integrity Number (RIN) using Agilent Bioanalyzer. All RNA samples had a RIN > 8. We used 4µg aliquots of total RNA to produce barcoded RNA sequencing libraries using the Illumina TruSeq Stranded mRNA kit. The quality of generated libraries was assessed using Agilent Bioanalyzer High Sensitivity DNA Kit and Kapa metrics. Libraries were pooled in two different pools based on barcode compatibility, and each pool was sequenced in two Illumina HiSeq2500 lanes, producing on an average of 42.1 million single end (SE) 100-bp reads per sample.

ChIP-seq sample processing

We processed samples in six randomly assigned groups in order to minimize batch effects. For each sample, we cut 90 mg of frozen liver tissue into 1 mm³ pieces, washed the cut tissue samples with cold phosphate-buffered saline (PBS), and fixed with 1% formaldehyde for 5 minutes at room temperature. We prepared nuclei of each washed sample using the Covaris truChIP Tissue Chromatin Shearing Kit. Chromatin was then sheared for 16 minutes using a Covaris S220 Focused-ultrasonicator. We quantified shearing efficiency and chromatin concentration using Agilent Bioanalyzer High Sensitivity DNA Kit.

From each specimen, we kept aside a 0.5 μg aliquot of sheared chromatin to be
used as input. We used two 5 μg aliquots of chromatin per sample to perform
immunoprecipitation (IP) with antibodies directed at H3K27ac (ab4729) and H3K4me1
(ab8895) respectively. We performed each IP using 5 μg of antibody with an overnight
incubation at 4°C as specified by the Magna ChIP A/G Chromatin Immunoprecipitation Kit
protocol. After elution and protein-DNA crosslink reversal, we extracted DNA using Zymo
Research ChIP DNA Clean & Concentrator kit, and quantified extracted DNA using Agilent
High Sensitivity kit and Qubit 2.0.

We used 5 to 15 ng of input and immunoprecipitated DNA to generate sequencing
libraries using the NEBNext Ultra ChIPseq library kit, following protocols specified by the
manufacturer. We assessed the quality of each constructed library using Agilent
Bioanalyzer High Sensitivity DNA Kit and Kapa metrics. Libraries were multiplexed, pooled
and sequenced on a total of 16 Illumina HiSeq2500 lanes, producing on an average of 40.6
million SE 100-bp reads per sample.

Sequence QC: ChIP-seq and RNA-seq
We performed the standard quality control (QC) measures on both ChIP-seq and RNA-seq
fastq files using FastQC v0.11.3 (Andrews, 2010). We then trimmed sequencing adapters
and low quality base calls using TrimGalore! v0.4.1 with the following parameters: -
stringency   5   -length   50   -q   20
(http://www.bioinformatics.babraham.ac.uk/projects/trim_galore/).

RNA-seq alignment and gene expression quantification
We aligned all sequences that passed QC to the reference genomes from the Ensembl
database (bushbaby: otoGar3; chimp: CHIMP2.1.4; humans: GRCh38; rhesus macaque:
Mmul1; marmoset: C_jacchus3.2.1; mouse lemur: micMur1) using STAR v2.5 (Dobin et al.,
2013) in 2-pass mode with the following parameters: --quantMode TranscriptomeSAM --
outFilterMultimapNmax 10 --outFilterMismatchNmax 10 --outFilterMismatchNoverLmax 0.3
--alignIntronMin 21 --alignIntronMax 0 --alignMatesGapMax 0 --alignSJoverhangMin 5 --
runThreadN 12 --twopassMode Basic --twopass1readsN 60000000 --sjdbOverhang 100.
We filtered bam files based on alignment quality (q = 10) and sorted using Samtools
v0.1.19 sort function (Li, 2009). We used the latest annotations for each species obtained
from Ensembl to build reference indexes for the STAR alignment:
Homo_sapiens.GRCh38.82.chr.gtf;   Pan_troglodytes.CHIMP2.1.4.82.chr.gtf;

Macaca_mulatta.MMUL_1.82.chr.gtf;   Callithrix_jacchus.C_jacchus3.2.1.82.chr.gtf;
Otolemur_garnettii.OtoGar3.82.gtf; Microcebus_murinus.micMur1.82.gtf (Aken et al., 2016).
We used FeatureCounts (Liao et al., 2014) to count reads mapping to each protein coding
gene/lincRNA/pseudogene, according to Ensembl annotations for the six studied species.
Read counts were then normalized based on feature (gene) length.
925
926 **Differential gene expression analysis**
927 We analyzed differential gene expression levels for each species using read counts,
928 normalized by feature length with the DESeq2 software (Love et al., 2014), and with the
929 following model:

930 *design = ~condition*

931
where condition indicates the species or the group of species (e.g. apes).
We used a set of 10,243 genes annotated as orthologs in the six species according
to Ensembl (BioMart v. 0.9; Smedley et al., 2015; Table S3) and used 5% False Discovery
Rate (FDR) as our multiple-testing-corrected significance threshold. The overall analysis
included three main comparisons of our interest: 1) pairwise comparisons between human
and each of the other five species; 2) comparisons for human-specific differential
expression of orthologous genes (human versus other five primates grouped together); 3)
comparisons for ape-specific differential expression of orthologous genes (human +
chimpanzee versus other four primates).
4) comparisons for Catarrhini-specific differential expression (human+chimpanzee+rhesus
macaque versus other primates). Finally comparison between Strepsirrhini (human,
chimpanzee, rhesus macaque and marmoset) and Haplorrhini (mouse lemur and
bushbaby) was computed.
945
ChIP-seq QC and alignment
We applied standard QC measures to ChIP-seq data as described above for RNA-seq data
processing. We aligned the sequences that passed QC to the reference genomes from the
Ensembl database (bushbaby: otoGar3; chimpanzee: CHIMP2.1.4; humans: GRCh38;
rhesus macaque: Mmul1; marmoset: C_jacchus3.2.1; mouse lemur: micMur1) using
Burrows Wheeler Alignment tool (BWA), with the MEM algorithm (Li, 2013). We sorted the
filtered bam files using Samtools v0.1.19 (Li, 2009).
953

ChIP-seq peak calling and QC

We called peaks for each individual using MACS2 (Zhang et al., 2008), at 1% FDR, and with parameters recommended for histone modifications (Liu, 2014): --m 30 40 --ext size 147 -B. Input was used as a control in all differential histone modification analyses. We performed QC on peaks called for each specimen using metrics recommended by ENCODE (Landt et al., 2012): Fraction of Reads in Peaks (FRiP), Normalize Strand Correlation coefficient (NSC) and Relative Strand Correlation coefficient (RSC) and ENCODE quality score.

In order to compute FRiP, we used Bedtools (v2.25.0; Quinlan and Hall, 2010) to intersect bed files containing all coordinates of called peaks (narrowPeak output of MACS2) with the original sorted bam file of the specific ChiP-seq sample. Then, we used a publicly available perl script to count the reads mapping in the intersection regions(https://github.com/mel-astar/mel-ngs/blob/master/mel-chipseq/chipseq-metrics/getCnt.pl). As recommended by the ENCODE consortium, we selected a threshold of 1% as acceptable FRiP values. We computed the two strand correlation metrics (NSC, RSC) using Phantompeakqualtools (Landt et al., 2012). For H3K27ac, NSC ≥ 1.05 and RSC ≥ 0.8 were used as threshold for retaining samples. For H3K4me1, that tends to produce broader peaks, we used NSC ≥ 1.05 and RSC ≥ 0.5 (Table S1).

Samples that did not pass at least two of the three main QC metrics (FRiP, NSC, RSC) were excluded for any downstream analysis. We then called human consensus peaks for H3K27ac and H3K4me1 using MACS2 and the above described parameters with the 1% FDR threshold. All human samples passing QC were considered as replicates of each other for the consensus peak calling. These human consensus H3K27ac and H3K4me1 profiles were used to perform all of the below described human-centric downstream analyses. Peaks called for the other species were only used for the above mentioned QC purposes but were not utilized for any of the downstream analyses, with the exception of the chimpanzee consensus peaks (see below). In order to assess how our data compares to known liver related regulatory regions, we overlapped our set of human consensus peaks to the set of H3K27ac and H3K4me1 peaks generated for HepG2 cells from the ENCODE consortium and to the set of permissive enhancers generated by the FANTOM5 consortium.

Comparison to previous findings using human liver ChIP-seq data

We compared our human H3K27ac ChIP-seq data, with a set of published human liver
H3K27ac peaks (Villar et al., 2015). We used the window function of Bedtools to quantify
the number of H3K27ac peaks in the replication dataset that either overlap with, or are
found within 1 kb up- and downstream from each of our human H3K27ac peaks. With this
approach, we quantified and assessed overlaps between discovery and replication
datasets.

Next, we characterized sets of replicated and unreplicated peaks between the two
datasets. Using the procedures previously described for the comparison of conserved and
recently evolved CREs, we annotated several genomic features for both replicated and
unreplicated peaks. Specifically, we included information regarding: 1) the average distance
from the closest TSS; 2) the class of the associated gene (e.g., protein coding, lincRNA); 3)
the number of cell types with an overlapping ENCODE DHS site and; 4) the number of
ENCODE TFBS overlapping the peak. Moreover, a logistic regression on the q-values of
replicated and not replicated peaks was performed.

Parallelized reporter assay
We obtained a list of 334 putative 1-kb long CREs overlapping liver eQTLs from Brown and
collaborators (Brown et al., 2013). This data included both enhancers (distance from TSS >
1Kb) and promoters (distance from TSS < 1kb). 122 CREs out of these 334 CREs
overlapped our human ChIP-seq peaks (53 enhancers and 69 promoters; Table S6). Within
each of the loci defined by the investigated liver eQTLs, we predicted a 1-kb CRE. These
predicted CREs were amplified in individual PCRs performed on 120 pooled Yoruban
HapMap DNA samples. PCR products from each reaction therefore represent a complex
mixture of haplotypes. We inserted barcodes (hereafter, tags) consisting of a 160-bp oligo,
including a randomized 20-bp unique barcode for each construct, into luciferase reporter
vectors (pGL4.23 and pGL4.10), immediately downstream of the luciferase gene, after
linearizing the vector with the XbaI restriction enzyme.

We pooled and cloned DNAs from each putative CRE into uniquely barcoded
luciferase reporter vectors (pGL4.23 were used for enhancers and pGL4.10 for promoters),
using the Gibson Assembly Kit (New England BioLabs). The CREs were specifically
inserted upstream of the luciferase gene, after linearizing the vector with the restriction
enzymes KpnI and XhoI. We then transfected the complex pool of CRE reporters into
HepG2 cells in two replicates. 24 hours after transfection, we extracted total RNA, purified
poly-A RNA, and produced cDNA that was used to amplify the tag, with the QIAGEN One

Step RT-PCR Kit with primers that included Illumina adapters for sequencing. Tag libraries were pooled and sequenced on a single Illumina HiSeq2500 lane, producing single end (SE) 50-bp reads. We amplified the tags from the vector before the transfection and sequenced them in the same pool with the tag-RNA libraries as a control for tag read counts.

In parallel, reporter tags were unambiguously associated with each specific CRE by sequence based sub-assembly. Briefly, we cut the luciferase gene from the vector by inverse PCR and then re-ligated the vector using the T4 Polynucleotide Kinase (PNK) + T4 ligase kit from NEB. In this way CREs and tags were flanking each other and CRE-tag complexes. The CRE-tag complexes were then PCR amplified using a reverse primer that included Illumina adapter for sequencing. Next, the CRE-tag PCR product was digested for 5 minutes at 55°C using Nexetera Tn5 Transposase (TDE1) in order to produce fragments of variable length (from ca. 150 bp to the entire length of the construct). When cutting the fragments, TDE1 also inserts an Illumina compatible adapter in proximity of the cutting site. We performed a PCR to enrich the libraries using the TDE1 inserted adapter as forward primer and the previously included Illumina adapter as reverse primer.

We pooled the two libraries (one for pGL4.10 and one for pGL4.23 constructs) and sequenced them on an Illumina MiSeq, producing paired-end (PE) reads (250 + 50 bp). After performing QC with FastQC v0.11.3, we aligned the sub-assembly sequences to the human genome (GRCh37/hg19) using BWA mem and the bam files were sorted and indexed with Samtools v0.1.19. Finally, we produced a matrix listing all of the CRE-tag associations. Tags associated with more than one CRE were discarded and not used for further analyses. After attributing each tag to its uniquely associated CRE, we used sequence based tag counts (HiSeq reads), normalized by sequencing depth, to quantify the gene expression level driven by each CRE, and therefore its functionality as enhancer/promoter.

For each CRE, we used a count-based generalized linear model to quantify differential expression between RNA (after transfection) and DNA (before transfection), assuming a Poisson error function:

model= count~condition

where condition indicates that the read count comes either from RNA (replicates 1 and 2) or DNA-control.

055   In presence of a significant p-value, the model indicates a significant difference
056   between the expression of the tags in the RNA samples compared to their DNA control.
057   The effect size estimate was then used to infer whether the RNA samples were
058   upregulated, hence showing significantly higher level of expression of the tags compared to
059   their DNA controls, and therefore indicating that the CRE is a functional regulatory element.
060
Detection of orthologous regions for human peaks in each primate
We mapped orthologous sequences using all identified consensus ChIP-seq peak regions
in both H3K27ac and H3K4me1 experiments. We used the Ensembl multiple sequence
alignment (MSA) reference database with the following specifications: 39 Eutherian
mammals; method_link_type: "EPO_LOW_COVERAGE"; species_set_name: "mammals"
(Herrero et al., 2015). For orthologous sequence analysis, 500 bp up- and downstream
regions were considered to be a part of the identified consensus peaks in all six species.
We queried all regions directly from the reference database using the REST API (Yates et
al., 2015).
All orthologous sequences retained gaps generated by MSA. In cases of incomplete
chromosome assembly (e.g. mouse lemur), composite sequence representations
containing parts of multiple scaffolds are used as a reference as provided by the Ensembl
database. As a result, we independently queried each peak region as well as regions
covering each peak +/- 500 bp. All orthologous sequences pulled from the references for
downstream analysis contained only directly aligned sequences. All regions with no
orthologous regions represented in the MSA reference were excluded from further
analyses. All query results in .json format and extracted sequences formatted for the MSA
alignment as well as genomic position information are provided in the repository mentioned
in the final section.
080
Correlation between human and marmoset read counts within orthologous regions
We assessed human and marmoset (i.e. the species with the smallest number of peaks
called; Supplemental File S1) normalized read counts at the 47,673 orthologous CREs,
after splitting them into two groups: 1) regions with overlapping peaks present in both
marmoset and human, and 2) regions with a peak present only in human. Spearman's
correlation (ρ) between human and marmoset normalized read depths was then computed
for each the two groups.
088

Differential histone modification analysis

Using the above described procedure, for both H3K27ac and H3K4me1, we produced a single matrix including the human peaks having an ortholog in each of the studied species and the associated read count for the specific histone mark and for the input in all of the six species. The normalized read counts were used for differential ChIP-seq analysis with DESeq2, performing an interaction analysis using the Wald statistic between the histone marks read counts and their associated input values:

$$design = \sim assay + condition + assay{:}condition$$

where the assay indicates either histone marks data or input data, and condition indicates instead the species or the group of species (e.g. apes, Catarrhini, Haplorrhini).

Differential histone mark analysis included the same species × species and group × group comparisons described for RNA-seq. We used 10% FDR as our multiple testing corrected significance threshold. Further, different FDRs (5%, 10%, 20%, 30%, 40%) were tested to assess the robustness of our approach.

We analyzed differential histone modifications for the two marks independently. However, in order to quantify the fraction of differentially marked cis-regulatory elements (CREs) in all of the above described pairwise comparisons, we used bedtools to identify the CREs predicted by H3K27ac (i.e. H3K27ac peaks) that would overlap those predicted by H3K4me1 for at least a 25% of their length. The list of unique CREs was used to estimate the fraction of differentially bound CREs for each of the above mentioned human-centric pairwise comparisons. We defined CREs as evolutionarily conserved if they did not show significant differential histone mark in any of the above mentioned pairwise comparisons. Otherwise, the CRE were defined as recently evolved.

Sequence conservation versus functional conservation analysis

We estimated per-nucleotide pairwise divergence for all five species in comparison to humans using MSA aligned sequences of orthologous regions for consensus peaks +/- 500 bps. All gaps in human were excluded from analysis. Regions not included in the set of six-way orthologous CREs were also pruned. Finally, we removed outliers - with respect to the distribution of the genetic distances in the given pairwise comparison - using the R package *outliers* (Komsta, 2006). We intersected the set of 47,673 orthologous liver CREs with the UCSC phastConsElements30wayPlacental track (Siepel et al., 2005), to assess whether

genomic regions characterized by conserved nucleotide sequence (i.e. phastCons elements) are significantly more associated to CREs detected as evolutionarily conserved in primates.

Analysis of features associated with evolutionary conservation of CREs

The following features were associated to each of predicted human CREs: nearest gene, distance from TSS, functional categories of genes, CRE category and histone mark (Table S2). Any human CREs without orthologous regions in other five species have been excluded from our analyses. To assess the correlation between the degree of conservation of a CRE and the number of cell types where the CRE is functional, we obtained publicly available data for DNase hypersensitivity sites (DHS) for over 200 cell types (ENCODE Project Consortium, 2012). For each CRE overlapping one or more DHS regions, we annotated the number of cell types where the specific CRE is putatively functional.

We estimated the correlation between the degree of conservation of CREs and the number of TFBSs by comparing our human consensus peaks with previously published HepG2 TF-binding profiles (ENCODE Project Consortium, 2012). Further, we quantified the proportion of putative primate CREs overlapping known transcribed enhancers (eRNAs) by using 43,011 known permissive transcribed enhancers (FANTOM5 Consortium, 2014). Similarly, for liver-specificity of human CREs analysis, we used coordinates of published liver eQTLs (Innocenti et al., 2011). We then selected all genes within 10 kb distance from evolutionarily conserved CREs for gene set enrichment analysis using GOrilla software (Eden et al., 2007; Eden et al., 2009). All genes found within 10 kb of any of the 47,673 orthologous CREs are used as a background for the enrichment test.

Known and *de novo* motif analysis

Genomic coordinates of orthologous regions were used to extract target sequences from the Ensembl references without MSA alignment gaps. All regions containing consensus peaks identified as human- and apes-specific and primate-conserved were used for the motif discovery and enrichment analysis. MEME-chip was used for known motif discovery and enrichment analysis using the Jaspar database (Bailey et al., 2009). We used DREME (Bailey, 2011). De novo motif identification was performed with AME (McLeay et al., 2010). Jaspar and Hocomoco (v10) databases were used as references to estimate similarities to known motifs. All motif discovery and enrichment analysis used default settings and parameters provided by the developers except for the maximum *de novo* motif discovery

threshold (changed from 1 to 1000 for maximum threshold). Shuffled input sequences were

used to estimate the background distribution of motifs.

Overlap of transposable elements (TEs) with primate CREs

We used Bedtools to overlap the RepeatMasker track for GRCh38 to the set of unique

human CREs that would overlap a TE for at least 25% of their length. TE enrichment

analysis was performed using the *TEAnalysis* pipeline with TE-analysis_Shuffle_bed v. 2.0,

setting 1000 replicates (https://github.com/4ureliek/TEanalysis; Kapusta et al., 2013). To

test the regulatory effect of enriched young TEs on primate gene expression, we performed

a differential gene expression analysis between Strepsirrhini (human, chimp, rhesus

macaque, marmoset) and Haplorrhini (bushbaby, mouse lemur) and quantified the number

of CREs associated to differential expressed genes that overlapped a TE younger than the

Haplorrhini-Strepsirrhini divergence.

Analysis of SINE-VNTR-*Alu*s (SVA) TEs enriched in human CREs

We intersected all known SVAs annotated in the human genome with all human consensus

peaks from our study. We used the same 25% overlap threshold as described above and

considered all human consensus peaks regardless of presence of orthologous regions in

the other species. The two SVA lists (overlapping and not overlapping the CREs,

respectively) were annotated for: the average distance from the closest TSS, functional

categories of associated gene, the average number of cell types with available DHS data,

the average number of TFBS overlapping the SVAs, and finally the average number of

TFBS within 10 kb up- and downstream of the SVAs. We used AME (McLeay et al., 2010)

to look for motifs enriched in the exapted SVAs, using the not exapted SVAs as a control.

Luciferase reporter assay validation of *GRIN3A* and *JARID2*

To test for species- or clade-specific regulatory activity, we compared activity of two

predicted functional CREs with the empty pGL4.23 vector as a negative control. For

GRIN3A we PCR amplified the CRE (Table S6), and cloned the fragment into pGL4.23

using the NEB Gibson Assembly Kit. The *JARID2* CRE was synthesized by GenScript and

cloned into the same pGL4.23 vector. Cells were grown in DMEM high glucose (Gibco

#11965084) supplemented with 10% fetal bovine serum (FBS) (GE Healthcare Life

Sciences #SH3091003) containing antibiotic and antimycotic (Gibco #15240062) in a

humidified incubator with 5% CO_2 at 37°C. HepG2 cells were seeded in 48-well CellBIND

surface plates (Costar #3338) with 1.5 x 10⁵ cells per well 24 h prior to transfection.

Transfection complexes were formed using 800 ng of each construct with 1 µL of TransIT-

LT1 transfection reagent (Mirus #MIR2304) and Opti-MEM (Gibso #31985070) in a total

volume of 27 µL, incubated for 20 min and then added to cells. After transfection, cells were

incubated for 24 h and were lysed in passive lysis buffer. To read firefly luciferase activity,

100 µL of LARII were added to 20 µL of cell lysate (from the dual-luciferase reporter assay

system from Promega #E1910). We read Luminescence for 2 seconds per well on a 96-well

compatible plate luminometer (ThermoFisher Luminoskan Ascent). The constructs were

tested using three vector preparations in three to four technical transfection replicates (9 to

12 measurements for for each construct). We normalized for transfection replicates effect

using a linear model:

lm(log10(luciferase) ~ replicate + element).

Validation of the gene regulatory functionality of TE families

HepG2 cells were cultured in DMEM + GlutaMAX (Gibco) supplemented with 10% Fetal

Bovine Serum (Gibco) and Normocin (InvivoGen). Transposable element constructs were

built by synthesizing (GenScript) the Dfam (Hubley et al., 2016) consensus sequence for

each element and cloning into the pGL3 Basic vector (Promega) with an added minimal

promoter (pGL3 Basic[minP]). pGL3 BASIC[minP] with no insert was used as the negative

expression control. pRL null (Promega) was the renilla control for transfection efficiency.

TAP2 cloned into the pGL3 Basic[minP] was the positive control. Confluent HepG2 cells in

opaque 96 well plates in 90ml of Opti-MEM (Gibco) were transfected according to the

Lipofectamine p3000 protocol (Invitrogen) with 100 ng of the luciferase containing plasmid,

1 ng of pRL null, 0.3 ml of Lipofectamine 3000, and 0.2ml of p3000 reagent in 10 ml of Opti-

MEM per well. The cells incubated in the transfection mixture for 24h hours then the media

was then changed to the regular FBS containing media for an additional 24 hours. Dual

Luciferase Reporter Assays (Promega) were started by incubating the cells for 15 mins in

20 ml of 1x passive lysis buffer. Luciferase and renilla expression were then measured

using the Glomax multi+ detection system (Promega). Luciferase expression values of the

transposable elements and TAP2 were standardized by the renilla expression values and

background expression values as determined by pGL3-Basic expression. Enriched motifs

were found by analyzing the Dfam (Hubley et al., 2016) consensus sequences of the TEs

found to have a regulatory ability significantly different from the pGL3 Basic[minP] empty

vector using the MEME Suite. TomTom (Gupta et al., 2007) was used to match binding site

motifs in the Jaspar database to the enriched motifs found in our data.

Additional notes on analyses used throughout the project

All statistical analyses (DESeq2 analysis, Fisher's exact tests, logistic regressions,

Spearman's correlations, Wilcoxon tests, General Linear Models, binomial test, and

quantiles calculations) were performed using R v3.3.1. Figures were made with the

package ggplot2 (Wickham, 2009) in R v3.3.1. Bedtools v2.25.0 (Quinlan et al., 2010) was

used for overlap and closest feature/window analyses. All relevant scripts and pipelines are

available online (https://github.com/ypar/cre_evo_primates.git).

Supplementary references

237
Aken, B.L., Ayling, S., Barrell, D., Clarke, L., Curwen, V., Fairley, S., Fernandez Banet, J.,
Billis, K., García Girón, C., Hourlier, T., et al. (2016). The Ensembl gene annotation system.
Database 2016.

Andrews S., (2010). FastQC: a quality control tool for high throughput sequence data.
Available online at: http://www.bioinformatics.babraham.ac.uk/projects/fastqc

Bailey, T.L. (2011). DREME: motif discovery in transcription factor ChIP-seq data.
Bioinformatics *27*, 1653–1659.

Bailey, T.L., Boden, M., Buske, F.A., Frith, M., Grant, C.E., Clementi, L., Ren, J., Li, W.W.,
and Noble, W.S. (2009). MEME SUITE: tools for motif discovery and searching. Nucleic
Acids Research *37*, W202-208.

Brown, C.D., Mangravite, L., and Engelhardt B. (2013) Integrative Modeling of eQTLs and
Cis-Regulatory Elements Suggests Mechanisms Underlying Cell Type Specificity of eQTLs.
Plos Genetics http://dx.doi.org/10.1371/journal.pgen.1003649

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P.,
Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner.
Bioinformatics *29*(1), 15–21.

Eden, E., Lipson, D., Yogev, S., and Yakhini, Z. (2007). Discovering Motifs in Ranked Lists
of DNA Sequences. PLoS Comput Biol *3*, e39.

Eden, E., Navon, R., Steinfeld, I., Lipson, D., and Yakhini, Z. (2009). GOrilla: a tool for
discovery and visualization of enriched GO terms in ranked gene lists. BMC Bioinformatics
10, 1–7.

FANTOM Consortium and the RIKEN PMI and CLST (DGT) (2014). A promoter-level
mammalian expression atlas. Nature *507*, 462–470.

Herrero, J., Muffato, M., Beal, K., Fitzgerald, S., Gordon, L., Pignatelli, M., Vilella, A.J.,
Searle, S.M.J., Amode, R., Brent, S., et al. (2016). Ensembl comparative genomics
resources. Database 2016.

Hinrichs, A.S., Karolchik, D., Baertsch, R., Barber, G.P., Bejerano, G., Clawson, H.,
Diekhans, M., Furey, T.S., Harte, R.A., Hsu, F., et al. (2006). The UCSC Genome Browser
Database: update 2006. Nucleic Acids Research *34*, D590–D598.

Hubley, R., Finn, R.D., Clements, J., Eddy, S.R., Jones, T.A., Bao, W., Smit, A.F.A., and
Wheeler, T.J. (2016). The Dfam database of repetitive DNA families. Nucleid Acids
Research *44*, D81-D89.

Gupta, S, Stamatoyannopoulos, J.A., Bailey, T.L., and Noble, W.S. (2007). Quantifying
similarity between motifs. Genome Biology *8*, R24.

Innocenti, F., Cooper, G.M., Stanaway, I.B., Gamazon, E.R., Smith, J.D., Mirkov, S.,
Ramirez, J., Liu, W., Lin, Y.S., Moloney, C., et al. (2011). Identification, Replication, and
Functional Fine-Mapping of Expression Quantitative Trait Loci in Primary Human Liver
Tissue. PLoS Genet *7*, e1002078.

Kapusta, A., Kronenberg, Z., Lynch, V.J., Zhuo, X., Ramsay, L., Bourque, G., Yandell, M.,
and Feschotte, C. (2013). Transposable Elements Are Major Contributors to the Origin,
Diversification, and Regulation of Vertebrate Long Noncoding RNAs. PLoS Genet *9*,
e1003470.

Komsta, L. (2006). Processing data for outliers. R News, 6(2), 10-13.

Landt, S.G., Marinov, G.K., Kundaje, A., Kheradpour, P., Pauli, F., Batzoglou, S., Bernstein,
B.E., Bickel, P., Brown, J.B., Cayting, P., et al. 2012. ChIP-seq guidelines and practices of
the ENCODE and modENCODE consortia. Genome Research *22*(9): 1813–1831.

Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general purpose
program for assigning sequence reads to genomic features. Bioinformatics *30*, 923–930.

Liu, T. (2014). https://github.com/taoliu/MACS/wiki/Call-differential-binding-events.

Li, H. (2013) Aligning sequence reads, clone sequences and assembly contigs with BWA-
MEM. arXiv:1303.3997v1

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biology *15*, 1–21.

McLeay, R.C., and Bailey, T.L. (2010). Motif Enrichment Analysis: a unified framework and
an evaluation on ChIP data. BMC Bioinformatics *11*, 1–11.

Melnikov, A., Murugan, A., Zhang, X., Tesileanu, T., Wang, L., Rogov, P., Feizi, S., Gnirke,
A., Callan, C.G., Kinney, J.B., et al. (2012). Systematic dissection and optimization of
inducible enhancers in human cells using a massively parallel reporter assay. Nat Biotech
30, 271–277.

Patwardhan, R.P., Hiatt, J.B., Witten, D.M., Kim, M.J., Smith, R.P., May, D., Lee, C.,
Andrie, J.M., Lee, S.-I., Cooper, G.M., et al. (2012). Massively parallel functional dissection
of mammalian enhancers in vivo. Nat Biotech *30*, 265–270.

Quinlan, A.R., and Hall, I.M. (2010). BEDTools: a flexible suite of utilities for comparing
genomic features. Bioinformatics *26*, 841–842.

R Core Team (2016). R: A language and environment for statistical computing. R.
Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Sharon, E., Kalma, Y., Sharp, A., Raveh-Sadka, T., Levo, M., Zeevi, D., Keren, L., Yakhini,
Z., Weinberger, A., and Segal, E. (2012). Inferring gene regulatory logic from high-
throughput measurements of thousands of systematically designed promoters. Nat Biotech
30, 521–530.

Siepel, A., Bejerano, G., Pedersen, J.S., Hinrichs, A.S., Hou, M., Rosenbloom, K., Clawson,
H., Spieth, J., Hillier, L.W., Richards, S., et al. (2005). Evolutionarily conserved elements in
vertebrate, insect, worm, and yeast genomes. Genome Research *15*, 1034–1050.

Smedley, D., Haider, S., Durinck, S., Pandini, L., Provero, P., Allen, J., Arnaiz, O., Awedh,
M.H., Baldock, R., Barbiera, G., et al. (2015). The BioMart community portal: an innovative
alternative to large, centralized data repositories. Nucleic Acids Research *43*, W589–W598.

Villar, D., Berthelot, C., Aldridge, S., Rayner, T.F., Lukk, M., Pignatelli, M., Park, T.J.,
Deaville, R., Erichsen, J.T., Jasinska, A.J., et al. Enhancer Evolution across 20 Mammalian
Species. Cell *160*, 554–566.

Wickham., H. (2009). ggplot2: Elegant graphics for data analysis. Springer-Verlag New
York, 2009.

Yates, A., Beal, K.,Keenan, S., McLaren, W., Pignatelli, M., Ritchie, G.R.S., Ruffler, M.,
Tylor, K., Vullo, A. and Flicek, P. (2015). The Ensembl REST API: Ensembl Data for any
language. Bioinformatics *31*(1): 143–145.
322

H3K4me1
4.78%

H3K27ac & H3K4me1
54.15%

H3K27ac
41.07%

326 histone modified Venn diagram showing the distribution marks on the
326 promoters of genes that are expressed in the primate liver.

	REPLICATED	NOT REPLICATED
Distance from TSS (Kb)	29.6	90
Number of ENCODE cell types	17.7	9.3
Number of ENCODE HepG2 TFBS	10.7	3.3

Figure S2: Peaks bearing signatures of robust and broad regulatory activities are largely reproducible across studies. We compared our human H3K27ac data to a recent study focused on liver CREs in mammals (Villar et al., 2015). Overall, less than 40% of the peaks identified by Villar et al. (2015) overlapped one of our H3K27ac regions. Moreover, only 19% of the ENCODE HepG2 H3K27ac peaks were replicated in the Villar et al. (2015) dataset. We thus investigated possible features associated with ChIP-seq peak reproducibility.

(A) Peak discovery significance (q-value) is significantly correlated with cross-dataset reproducibility (logistic regression $p < 2.2\times10^{-16}$). (B) Replicated peaks are significantly more likely than non-replicated peaks to be associated with protein coding genes rather than with lincRNAs or pseudogenes (Fisher's exact test $p < 2.2\times10^{-16}$). (C) Replicated peaks are: 1) systematically closer to the nearest TSS (29.6 kb for replicated peaks, 90.8 kb for not replicated; Wilcoxon rank-sum test $p < 2.2\times10^{-16}$); 2) overlap chromatin accessible regions in significantly higher numbers of ENCODE cell types (an average of 17.7 cell types for replicated peaks and 9.3 cell types for unreplicated peaks; logistic regression $p < 2.2\times10^{-16}$); 3) contain a significantly higher number of transcription factor binding sites (TFBSs) per peak region as identified by ENCODE in HepG2 cells (10.7 TFBSs in the replicated peaks, 3.6 in the unreplicated; logistic regression $p < 2.2\times10^{-16}$).

350
351
Figure S3: Correlation of the normalized ChIP-seq read depths between human and
marmoset. Human consensus peaks with orthologous in all the six species were split into
two groups for this analysis: 1) regions with overlapping peaks present in both marmoset
and human, and 2) regions with a peak present only in human. While human and marmoset
normalized read counts were more highly correlated with each other in group 1
(Spearman's ρ = 0.67; p < 2.2×10^{-16}), we found a nearly as strong correlation in group 2
(Spearman's ρ = 0.57; p < 2.2×10^{-16}). These findings are consistent with the results of our
differential histone modification state analyses, which demonstrated that only a small
fraction of the 47,673 orthologous CREs (5.97%, FDR < 10%) are differentially modified,
despite the fact that we had a much smaller total number of peak calls in the marmoset
samples.
363

Figure S4: Enhancers with signature of transcription are conserved. Fraction of conserved and recently evolved CREs with signature of transcription (eRNA) based on FANTOM5 Consortium data. Fraction of conserved and recently evolved CREs with no signature of transcription (eRNA) based on FANTOM5 Consortium data, Fraction of conserved and recently evolved CREs overlapping and not overlapping liver eQTLs. Specifically, predicted human CREs overlapped 500 eQTLs detected in a recent study on human liver (Innocenti et al., 2011). We tested whether enhancers and promoters overlapping liver eQTLs would lean toward being more conserved or more labile than a random liver CRE, and we found that neither of these two conditions are satisfied (Fisher's Exact Test $p = 0.05076$), as we show that liver eQTLs behave as "average" liver regulatory elements.

Fig S5: **Functional analysis on *GRIN3A locus***. ChIP-seq read depth distributions and luciferase assays reporter activity for the CRE associated to *GRIN3A*.

JARID2 repressor ChIP-seq

SVA-F (**Hominid specific TE**)

Chr 6: 15,414,808-15,416,756

381
382
383 **Fig S6**: **Functional analysis on *JARID2 locus.*** ChIP-seq read depth distributions and
384 luciferase assays reporter activity for the CRE associated to *JARID2*.
385

Figure S7: **Lineage specificity of enriched TEs.** Word-cloud representing the lineage specificity of the enriched TE families.