Widespread patterns of sexually dimorphic

gene expression in an avian hypothalamic–pituitary–gonadal (HPG) axis

Matthew D. MacManes[1*], Suzanne H. Austin[2], Andrew Lang[1], April Booth[2], Victoria Farrar[2],

Rebecca M. Calisi[2]

[1] Department of Molecular, Cellular, and Biomedical Sciences. University of New Hampshire,

Durham NH 03824

[2] Department of Neurobiology, Physiology, and Behavior. University of California, Davis. Davis

CA. 95616

[*] Corresponding author: MacManes@gmail.com, @MacManes

Keywords: reproductive axis, rock dove, pigeon, *Columba livia*, gene expression, RNAseq,

reproduction, sexual behavior

ABSTRACT

The hypothalamic-pituitary-gonadal (HPG) axis is a key biological system required for

reproduction and associated sexual behaviors to occur. In the avian reproductive model of the

rock dove (*Columba livia*), we characterized the transcript community of each tissue of the HPG

axis in both sexes, thereby significantly expanding our mechanistic insight into HPG activity. We

report greater sex-biased differential expression in the pituitary as compared to the

hypothalamus, with multiple genes more highly expressed in the male pituitary being related to

secretory function, and multiple genes more highly expressed in the female pituitary being

related to reproduction, growth, and development. We report tissue-specific and sex-biased

expression in genes commonly investigated when studying reproduction, highlighting the need

for sex parity in future studies. In addition, we uncover new targets of investigation in both

sexes, which could potentially change our understanding of HPG function.

INTRODUCTION

The hypothalamic-pituitary-gonadal (HPG) axis is a system comprised of endocrine

glands whose function is vital to the regulation of reproduction and associated behaviors (Fig.

1). In all vertebrates studied, from humans to Agnatha, the jawless fishes, the HPG axis is

present and its function is generally conserved [1]. For reproduction to occur, the hypothalamus

must produce and secrete gonadotropin-releasing hormone (GnRH), which causes the pituitary

gland to secrete gonadotropins, luteinizing hormone (LH) and follicle stimulating hormone (FSH)

[2]. LH and FSH travel through the bloodstream and act upon receptors in the gonads (i.e. testes

or ovaries), stimulating gametogenesis and the secretion of sex steroids, such as testosterone

and estradiol. These sex steroids then bind with receptors within the HPG axis to create a

feedback system that facilitates reproduction and sexual behaviors. The HPG axis has been

lauded for providing a foundation to guide reproductive endocrinology investigations, but it has

also been criticized as an overly simplistic depiction of the mechanisms mediating reproduction.

Now, burgeoning genomic sequencing technologies are permitting a deeper and more complete

understanding of the mechanistic drivers of reproduction. They are allowing us to understand

how differential gene expression between the sexes could underpin sexually dimorphic

reproductive behavior, and how the full complement of genes expressed in the HPG axis could

work in concert to define physiology and behavior [3].

Previous understanding of HPG function has come about by measuring the amount of

circulating hormones in blood, conducting immunochemistry to visualize protein presence or

absence, conducting *in situ hybridization* to visualize a target gene, and the use of endocrine

and receptor agonists and antagonists and their resulting physiological and behavioral effects.

Due to recent technological advances, researchers are beginning to identify correlative and

causative links between gene activity and phenotype [4–10]. Concerning the genomics of

reproduction, some studies have generated whole-organism or tissue-specific sequence data of

which the HPG axis was a subset [11,12]. Even fewer studies have reported observing differential

gene expression of the HPG axis as a global system [13]. Here, we focus specifically on patterns

of tissue and sexually dimorphic gene expression in the HPG axis of the rock dove (*Columba*

livia) and discuss how they might relate to male and female reproductive strategies. Doves have

been historically used to study reproductive behavior [14,15] and now are proving to be a valuable

model for genomics research [16–18].

We characterize the first sex-specific avian HPG axis transcriptome using RNA extracted

from whole hypothalamus, pituitary, and gonads of sexually mature male and female rock doves

(*Columba livia*). We constructed an annotated *de novo* transcriptome assembly for these

tissues, which then enabled us to quantify the abundance of all mRNA transcripts per tissue.

We elected to perform a *de novo* assembly, rather than use any other alternative approach

(*e.g.,* mapping reads to the chicken or rock dove genome), after preliminary mapping

experiments suggested that genes important to the behavioral phenotypes in question were not

well represented in existing genomic resources. By incorporating candidate gene and *ab initio*

approaches across all tissue types and between sexes, we were able to both characterize

general patterns of gene expression and identify specific patterns of sex-biased expression. It is

our intention that these data illuminate sex differences in gene presence and abundance

throughout the HPG axis when birds are at a basal state (i.e. reproductively mature but sampled

when not actively breeding). By doing this, we create a resource for the scientific community to

devise further studies of how gene expression patterns change over different reproductive

stages and in response to physical and social events. As the HPG axis is fairly well-conserved

across vertebrates, these data may be useful for formulating potential therapeutic strategies for

abnormal HPG axis function.

103 Figure 1. Depiction of the hypothalamus, pituitary, and gonads of male and female rock doves.

104 Illustration by Natalia Duque.

109 <u>**Results**</u>

Sequence Read Data & Code Availability

In total, 20 hypothalamus (11 male, 9 female), 23 pituitary (14 male, 9 female), 13 testes, and

10 ovary samples from 24 birds were sequenced. Each sample was sequenced with between

2.3 million and 24.5 million read pairs. All read data are available using the project ID

PRJEB16136. Code used for the analysis of these data are available at https://git.io/vPA09.

Transcriptome Assembly and Evaluation

The Rock Dove transcriptome assembly consists of 88,011 transcripts (available here

https://goo.gl/PYrzas, dryad on acceptance). This assembly, along with the annotations

contained in gff3 format, is available at https://goo.gl/DyZ8pw (dryad on acceptance). The

transcriptome contains 25,696 with at least 1 hit to the Pfam database, 46,854 hits to

OrthoDB[19], 51,522 hits to the Uniref90 database, 3,108 hits to the transporter database [20], and

452 hits to the Rfam database [21]. These 88,011 assembled transcripts map to 15,102 unique

genes in the *Gallus* genome. The evaluation using BUSCO and TransRate are presented in

Table 1 and 2 respectively.

Table 1: Quality control statistics for the assembled transcriptome: BUSCO metrics include
statistics regarding the number of universal vertebrate single copy orthologs found in the
assembly. 85.9% of the Avian BUSCOs were identified as full length (complete) sequences,
while 4.3% were found to be fragmented. 27.5% were found in greater than 1 copy in the Rock
Dove transcriptome.
131

	BUSCO		
	Complete	Duplicated	Fragmented
Rock Dove 1.0.3	85.9%	27.5%	4.3%

133   Table 2: Quality control statistics for the assembled transcriptome: TransRate metrics are
134   derived from mapping RNAseq reads to the assembly, with higher scores indicating a higher
135   quality assembly. A score of 0.41 ranks this assembly higher than the majority of other
136   published transcriptomes, with 90% of reads mapping, and only 2% of bases uncovered (no
137   read support) and 0% contigs low-covered (mean per-base read coverage of < 10).
138

	TRANSRATE					
	Score	Number of Contigs	Assembly Size	Percent Mapping	Percent Bases Uncovered	Percent Contigs Lowcovered
Rock Dove 1.0.3	0.41	88011	146.8Mb	90	2	0

Sequence Read Mapping and Estimation of Gene Expression.
Raw sequencing reads corresponding to individual samples of hypothalami, pituitary glands,
and gonads were mapped to the Rock Dove reference HPG transcriptome using Salmon, which
resulted in between 70% and 80% read mapping. These mapping data were imported into R
and summarized into gene-level counts using tximport, after which, edgeR was used to
generate normalized estimates of gene expression. Patterns of transcript-expression overlap
are presented in Figure 1. Of the 15,102 genes expressed in the HPG, 7,529 are expressed in
all tissues (median CPM > 10 - Figure 1). A total of 746 transcripts are expressed in the ovary,

hypothalamus, and pituitary, 732 are expressed uniquely in the testes, 529 uniquely in ovary,

418 uniquely in the hypothalamus, and 230 uniquely in the pituitary. Dyads (genes expressed

uniquely in pairs of tissues) follow an expected pattern, with the number of genes expressed

uniquely in the gonads (testes and ovary) being more than the number of genes expressed

uniquely in the pituitary and hypothalamus, followed by the pituitary and ovary, hypothalamus

and ovary, pituitary and testes, and hypothalamus and testes. Understanding tissue-specific

patterns of expression can provide a glimpse into the HPG axis as an interconnected functional

system.

A gene ontology-based pathway analysis[27] of the transcripts unique to each tissue provides

some insights into the specific biological processes important to each tissue. Of the genes that

are uniquely expressed in the testes, the three most highly represented pathways include

Inflammation mediated by chemokine and cytokine signaling pathway (Panther pathway number

P00031), CCKR signaling map (P06959) and Nicotinic acetylcholine receptor signaling pathway

(P00044). The top three pathways for genes uniquely expressed in the ovary are related to

Gonadotropin-releasing hormone receptor pathway (P06664), Wnt signaling pathway (P00057),

and angiogenesis (P00005). Genes uniquely expressed in the hypothalamus cluster around the

pathway terms oligodendrocyte differentiation, regulation of glial cell differentiation, positive

regulation of synapse assembly, negative regulation of neurogenesis and modulation of

synaptic transmission. The pituitary contains uniquely expressed genes related to the Wnt

signaling pathway (P00057), the Cadherin signaling pathway (P00012), and the gonadotropin-

releasing hormone receptor pathway (P06664).

Figure 1. Description of transcript presence overlap in the HPG axis. Bars are annotated by

connected dots which indicate a given number of transcripts expressed in those tissues. For

example, 746 transcripts are expressed in the pituitary, hypothalamus, and ovary, but these

genes are not expressed in the testes. Bars annotated by single blue dots indicate the number

of transcripts expressed uniquely in that tissue. For example, 732 transcripts are expressed only

in the testes.

Evaluation of Candidate Gene Expression.

Using the assembled transcriptome, we characterized expression in males and females across

the HPG axis (*e.g.,* Figure 2*)*. For the candidate genes evaluated, which include gonadotropin

releasing hormone (*GnRH-1*) and its receptor (*GnRH-1*-R), LH receptor (*LH-R*), FSH receptor

(*FSH-R*), androgen receptor (*AR*), estrogen receptor alpha and beta (*ESR alpha*, *ESR beta*). In

addition, we examined gonadotropin inhibitory hormone (*GnIH*), due to its inhibitory effect on the

HPG axis and reproductive behavior [22], vasoactive intestinal peptide (*VIP*), due to its role in

stimulating the release of the pituitary hormone, prolactin [23], and its receptor (*VIP-R*), prolactin

(PRL) due to its facilitation of parental care behaviors, including crop milk production [24], and its

receptor (*PRL-R*), arginine vasopressin-like receptor 1A and 1B (*AVPR1A, AVPR1B*),

mesotocin (*MT*) and its receptor (*MTR*), due to their role in social bonding behaviors [25,26],

progesterone receptor (*PGR*) due to its role in reproductive behavior [27], CYP19, which encodes

the aromatase enzyme (*ARO*), which is responsible for the conversion of testosterone into

estradiol [28].

Using a generalized linear model and least-squares means for post-hoc tests of significance

(P<0.05), we uncovered greater expression of *GnRH-1-R*, *AR*, and *PGR* in the female pituitary

as compared to the male pituitary (z-ratio 2.324, p-value =0.0201; z ratio = -3.842, p-value =

0.0001; z ratio = 5.559, p-value = <.0001, respectively). We also found greater expression of

PRL in the female pituitary (z ratio = 2.912, p-value = 0.0036) as compared to the male pituitary,

but *PRL* was more highly expressed in the male hypothalamus as compared to the female

hypothalamus (z ratio = -2.433, p-value = 0.0150). The *PRL* receptor was also more highly

expressed in the male hypothalamus and pituitary gland as compared to corresponding female

tissues (z ratio -2.674, p-value = 0.0075 and z ratio -2.564, p-value = 0.0104, respectively).

Finally, we discovered greater expression of *AVPR1A* in the male pituitary as compared to the

female pituitary (z ratio = -2.416, p-value = 0.0157).

Figure 2A-F. Box and whisker plots illustrating the differential expression between the sexes of

six candidate genes. The Y-axis indicates levels of gene expression as presented by taking the

log of transcripts per million plus one (TPM+1) while tissue and sex are presented on the X-axis.

The bottom and top of the box indicates the first and third quartiles. The band inside the box is

the median. Whiskers indicate the 95% confidence intervals of the data. ** denotes statistically

significant differences in expression.

Table 3: Mean gene expression (normalized counts per million (CPM)) +/- standard deviation)

for a list of candidate genes known to be important mediators of reproductive biology.

	Hypo		Pit		Gonads	
Gene	Female	Male	Female	Male	Female	Male
GnRH-I	116.1 +/- 75.9	142.3 +/- 121.6	0.3 +/- 0.5	1.3 +/- 2.3	0.6 +/- 0.7	1.2 +/- 1.1
GnRH-I-R	2.34 +/ 2.67	2.85 +/- 2.75	245.6 +/- 212.4	112.8 +/- 83.76	14.0 +/- 6.7	1.38 +/- 1.09
LH-R	19.0 +/- 11.3	23.03 +/- 12.2	0.9 +/- 0.95	0.38 +/- 0.67	264.45 +/- 77.54	42.59 +/- 39.1
FSH-R	4.02 +/- 4.42	5.13 +/- 2.62	9.4 +/- 3.75	11.5 +/- 5.62	151.59 +/- 80.2	181.0 +/- 51.56
AR	37.7 +/- 13.4	27.8 +/- 13.1	226.6 +/- 121.4	111.1 +/- 72.7	204.1 +/- 62.5	80.3 +/- 23.1
ERa	49.2 +/- 14.0	57.2 +/- 18.6	724.5 +/- 329.3	836 +/- 283.4	166.2 +/- 152.5	64.6 +/- 32.6
ERb	41.9 +/- 18.7	36.1 +/- 14.5	53.8 +/- 16.5	58.6 +/- 27.9	173.4 +/- 90.5	7.4 +/- 3.9
GnIH	287.8 +/- 148.7	383.2 +/- 510.0	4.0 +/- 3.8	2.9 +/- 3.1	0.1 +/- 0.2	0.7 +/- 0.7
VIP	1.3e+02 +/- 60.1	1.3e+02 +/- 98.3	1.19 +/- 1.37	.46 +- .77	1.3e+01 +/- 15.6	0 +/- 0
VIP-R	70.78 +/- 18.5	69.2 +/- 24.5	47.3 +/- 141.5	376.35 +/- 134.64	42.3 +/- 17.3	18.8 +/- 7.9
PRL	389.9 +/- 170.4	766.0 +/- 436.7	295840.2 +/- 320026.2	136607.5 +/- 121634.2	250.9 +/- 95.2	397.9 +/- 45.1
PRL-R	463.0 +/- 141.7	803.4 +/- 558.4	540.4 +/- 143.2	790.85 +/- 204	338.5 +/- 131.8	550.66 +/- 100.7
AVPR1A	117.3 +/- 58.8	98.3 +/- 34.4	202.2 +/- 111.3	362.2 +/- 181.6	19.0 +/- 13.5	3.2 +/- 2.3
AVPR1B	5.4 +/- 5.4	5.4 +/- 3.1	1538.3 +/- 503.5	1900.1 +/- 709.1	3.8 +/- 1.3	2.3 +/- 1.5
MT	264.9 +/- 82.1	206.9 +/- 101.1	173 +/- 39.9	172.78 +/- 60.9	182.11 +/- 265.4	11.1 +/- 2.98
MTR	<1	<1	<1	<1	<1	<1
PGR	50.6 +/- 24.6	51.4 +/- 20.8	874.4 +/- 294.9	216.7 +/- 82.6	1004.6 +/- 1193	126.7 +/- 14.4
CYP19	212.6 +/- 137.5	325.8 +/- 200.2	26.4 +/- 10.3	49.5 +/- 20.2	786.6 +/- 446.2	1.7 +/- 1.3

Global Evaluation of Gene Expression

Analysis of global patterns of gene expression, aimed at uncovering previously unknown sex-

specific differences in gene expression in the hypothalamus and pituitary by examining the

entire transcriptome, was conducted using edgeR. After controlling for over 15,000

comparisons, we normalized the count data using the TMM method [29], which is done by finding

a set of scaling factors for the library sizes that minimize the log-fold changes between the

samples for most genes. This analysis revealed sex-specific differences in gene expression,

particularly in the pituitary, where 218 genes were more highly expressed in males, while 153

genes were more highly expressed in females (Figure 3). The five most differentially expressed

genes are Zinc Finger AN1-Type Containing 5 (*ZFAND5*), Betacellulin (*BTC*) which appears to

be responsive to LH [30], Cbp/P300 Interacting Transactivator With Glu/Asp Rich Carboxy-

Terminal Domain 4 (*CITED4*), Growth Regulation By Estrogen In Breast Cancer 1 (*GREB1*)

which is regulated by estrogen [31], and Kinesin Family Member 24 (*KIF24*). All had false

discovery rates (FDR) < 7e-09. The full table of differentially expressed genes is available at

https://git.io/vXJWp, which includes several tantalizing novel candidates, including Ecto-NOX

Disulfide-Thiol Exchanger 1 (*ENOX1*; Fig. 4). In contrast to the pituitary, the hypothalamus

exhibited only a single differentially expressed gene. That gene was potassium voltage-gated

channel subfamily Q member 1 (*KCNQ1*, FDR =0.007). KCNQ1, which we found to be more

highly expressed in females, is known to be imprinted (expressed in a parent-of-origin-specific

manner) in mammals [32].

A number of expression patterns in the pituitary were identified by gene ontology analysis,

which uses a controlled vocabulary to describe gene function and relationships between these

terms. First, gene ontology terms that describe genes differentially expressed in the female

pituitary (Table 4) are clustered around terms related to the binding of lipid-based hormones,

immune function, growth, and development. Gene ontology terms from genes differentially

expressed in male pituitary (Table 5) are clustered around terms related to muscles and

movement.

275   Figure 3. MA plot for the *ab initio* comparison between genes differentially expressed in the

276   male and female pituitary. Red dots indicate statistically different genes. Those with positive

277   values on the log fold change (logFC) Y axis are more highly expressed in the male pituitary,

278   while those with negative values are more highly expressed in the female pituitary. The x-axis is

279   a measure of gene expression, with higher numbers indicating higher levels of gene expression.

Figure 4. Box and whisker plots illustrating sex-biased expression of two genes, Betacellulin

(*BTC*) and Ecto-NOX Disulfide-Thiol Exchanger 1 (*ENOX1*), identified using our global analysis.

The Y-axis indicates levels of gene expression as presented by taking the log of transcripts per

million plus one (TPM+1) while tissue and sex are presented on the X-axis. The bottom and top

of the box indicates the first and third quartiles. The band inside the box is the median. Whiskers

indicate the 95% confidence intervals of the data. ** denotes statistically significant differences

in expression.

Table 4: **Top 20 Gene Ontology (GO) Terms** describing differentially expressed genes in the

female pituitary gland. These terms describe the putative function of genes using a controlled

vocabulary. Abbreviations: Ontology (Ont), Molecular Function (MF), Biological Processes

(BP). N signifies the number of genes in the entire dataset that are linked to the specific

ontology term. F+ indicates the number of differentially expressed genes significantly more

highly expressed in the female pituitary, while M+ indicates the number of expressed genes

significantly more highly expressed in the male pituitary. Results are limited to terms where the

specific term contains >4 genes. In all cases, p-values are corrected for multiple hypothesis

tests using the Bonferroni correction.

GO number	Term	Ont	N	F+	M+	P-value
GO:0008289	lipid binding	MF	50	6	1	1.92E-05
GO:0048469	cell maturation	BP	13	3	0	4.29E-04
GO:0021700	developmental maturation	BP	18	3	0	1.18E-03
GO:0043178	alcohol binding	MF	7	2	0	2.93E-03
GO:0033293	monocarboxylic acid binding	MF	8	2	0	3.88E-03
GO:0014066	regulation of phosphatidylinositol 3-kinase signaling	BP	8	2	0	3.88E-03
GO:0006869	lipid transport	BP	27	3	2	3.92E-03
GO:0007548	sex differentiation	BP	27	3	0	3.92E-03
GO:0010876	lipid localization	BP	29	3	2	4.82E-03
GO:0048017	inositol lipid-mediated signaling	BP	9	2	0	4.95E-03
GO:0014065	phosphatidylinositol 3-kinase signaling	BP	9	2	0	4.95E-03
GO:0048015	phosphatidylinositol-mediated signaling	BP	9	2	0	4.95E-03
GO:0005496	steroid binding	MF	9	2	0	4.95E-03
GO:0005319	lipid transporter activity	MF	10	2	1	6.14E-03
GO:0043627	response to estrogen	BP	10	2	0	6.14E-03
GO:0004867	serine-type endopeptidase inhibitor activity	MF	12	2	0	8.86E-03
GO:0042698	ovulation cycle	BP	13	2	0	1.04E-02
GO:0022602	ovulation cycle process	BP	13	2	0	1.04E-02
GO:0005102	receptor binding	MF	114	5	0	1.07E-02
GO:0046545	development of primary female sexual characteristics	BP	14	2	0	1.20E-02

Table 5: **Gene Ontology (GO) Terms** describing genes differentially expressed genes in the

male pituitary gland. These terms describe the putative function of genes using a controlled

vocabulary. Abbreviations: Ont= Ontology (Ont), MF (Molecular Function (MF), BP (Biological

Processes (BP). N signifies= the number of genes in the entire dataset that are linked to the

specific ontology term. F+ indicates the number of differentially expressed genes significantly

more highly expressed in the female pituitary, while M+ indicates the number of expressed

genes significantly more highly expressed in the male pituitary. Results are limited to terms

where the specific term contains >4 genes. In all cases, p-values are corrected for multiple

hypothesis tests using the Bonferroni correction.

GO number	Term	Ont	N	F+	M+	P-value
GO:0006936	muscle contraction	BP	21	0	4	3.90E-04
GO:0006941	striated muscle contraction	BP	10	0	3	5.53E-04
GO:0003012	muscle system process	BP	26	0	4	9.15E-04
GO:0050879	multicellular organismal movement	BP	4	0	2	1.77E-03
GO:0050881	musculoskeletal movement	BP	4	0	2	1.77E-03
GO:0003009	skeletal muscle contraction	BP	4	0	2	1.77E-03
GO:0008092	cytoskeletal protein binding	MF	101	1	6	7.20E-03
GO:0003779	actin binding	MF	53	0	4	1.28E-02

Discussion

The HPG axis is a system comprised of endocrine tissues whose function is vital to the

regulation of reproduction and associated behavior. Here, we report patterns of gene expression

amongst tissues of the HPG axis as well as patterns of sex-biased gene expression within them.

We describe patterns of tissue-specific expression and shared incidence of expression between

tissues, providing an important picture of the functional-connectivity of these tissues. We also

describe sex-biased patterns of gene expression for a set of candidate genes, currently known

to play important roles in reproduction and associated behaviors. Lastly, we use an *ab initio*

approach to identify all differentially expressed transcripts between male and female pituitary

and hypothalamus. Our findings provide a vital resource for researchers examining the

molecular basis of reproductive behavior and the mechanisms underlying fundamental

physiological and behavioral differences in males and females.

The Columba livia HPG transcriptome. We present an HPG transcriptome that is substantially

complete, with fewer than 10% of universal avian orthologs missing (Table 1). This number

suggests that all, or nearly all, transcripts present in the avian HPG have been successfully

reconstructed, and the missing avian orthologs may be expressed in tissues other than those

studied here. The transcriptome is structurally sound, as is evident from its TransRate score

of .41, with 90% of reads mapping concordantly to the assembled reference (Table 2).

Evaluation of Candidate Gene Expression. Decades of previous research describing the

HPG axis and its role in reproductive biology have elucidated much of the molecular machinery

underlying phenotypes [33]. Based on this previous work, we targeted well-known substrates

involved in the facilitation and mediation of reproductive processes and their associated

behaviors for investigation (Table 3). Other genes not highlighted here can be found at

https://git.io/vXJW.

We discovered previously unrecognized sex-specific differences in several of these candidate

genes. Arginine vasopressin-like receptor 1A (*AVPR1A*) is known to be implicated in

reproductive behaviors such as pair-bonding and parental care [34–37]. *AVPR1A* is most highly

expressed in the pituitary, followed by the ovary, the hypothalamus, and lastly the testes. We

found statistically significant differences in expression of this gene in the pituitary, with

expression of this gene to be 1.8x higher in males as compared to females. Although a small

number of studies have investigated sex-specific differences in gene expression [38–40], including

one in which expression in males was linked to pair-bond formation [41], it is not well known if

AVPR1A is differentially expressed by sex. As fields of behavior and genomics further integrate,

links between sex-specific roles in pair-bonding and reproductive behavior in the context of

corresponding differences in gene expression will transform the way we understand how

behavior is regulated.

We also investigated other candidate genes involved in reproduction and associated behaviors.

Prolactin (*PRL*) is an important promoter of parental care, promoting lactation [42] and related to

appetite and weight gain [43]. We found *PRL* to be more highly expressed in the male

hypothalamus as compared to the female hypothalamus. The receptor for prolactin (*PRL-R*)

was also more highly expressed in the male hypothalamus and pituitary as compared to

females. In contrast, *PRL* was more highly expressed in the female pituitary as compared to the

male pituitary. Though this species exhibits bi-parental care, including the production of crop

milk by both males and females, neither sex was caring for eggs or chicks at the time of

collection to help control for potential reproductive stage confounds. However, the interesting

sexually-biased differences we note in inter-tissue gene expression inspires the question as to

whether *PRL* is regulated in a sex-specific manner to produce a sex-specific result? Or, despite

the difference in location of expression, does *PRL* activation lead to sexually monomorphic

prolactin-mediated reproductive processes and behaviors? In other words, the differences

observed in underlying genetic expression between the sexes may converge to the same

behavioral endpoint, permitting certain factors in one sex to offset the effects in the other,

making the sexes more similar [44]. Post our discovery that *PRL* expression differs relative to sex
and tissue of the HPG axis, the next step would be to investigate how the action of *PRL* on each
of these tissues manifests in males versus females.
381
In addition, we found *GnRH-1-R*, *AR*, and *PRG* to be more highly expressed in the female
pituitary as compared to the male pituitary. GnRH-1, produced in the hypothalamus, is a major
mediator of gonadotropin release in the pituitary. These gonadotropins signal the gonads to
produce androgens, which feedback onto androgen receptors along the HPG axis. We found
that androgen receptors are more highly expressed in females as compared to males. Females
tend to have lower circulation of androgens than males, so could this be a way to increase
sensitivity to androgen signals in females? Finally, progesterone receptor is associated with the
mediation of female breeding cycles [45], but it can also serve as an important precursor to the
creation of multiple hormones in males and females, including testosterone and estradiol. We
found that the androgen receptor is more highly expressed in the female pituitary relative to the
male pituitary. While expression in both sexes has been linked to reproductive development [46–
48], to our knowledge, sex differences in expression have never been reported. To better
elucidate the phenomenon of these sexually-dimorphic genetic expression phenotypes, further
studies of functional genomics are required. Indeed, study of the most well-characterized
candidate genes underlying reproductive behavior could reveal important and previously
unrecognized sex-differences in expression. As the field of behavioral genomics advances,
important questions to ask will be, are these differences biologically relevant, and if so, how do
differences in expression lead to differences in behavior and reproductive output? Future work
focused on these relationships will significantly shape our understanding of the molecular
mechanisms underlying the observed patterns.
402
Global Evaluation of Gene Expression. Avian reproductive behavior has been shown to be
heavily influenced by the endocrine function of the hypothalamus, pituitary, and gonads.
Although sex-biased differences in vertebrate reproductive behavior have been well-noted, data
on sex-specific patterns of gene expression that could be influencing these behaviors are
lacking. This may largely be due to the novelty of various genomic technologies and the
expense of collecting such data. However, it is becoming increasingly feasible and affordable to
conduct such studies, making possible the potential for a more integrative understanding of
reproductive behavior [3]. Thus, in addition to investigating sexually dimorphic candidate gene
expression in the HPG axis, we used an *ab initio* RNAseq approach to explore differences in *all*

sexually dimorphic HPG gene expression in each tissue. The newly discovered sex-biased

genetic differences that we report have the potential to offer a more in-depth picture of sexual

dimorphism at the molecular level.

To conduct a global analysis of gene expression, we compared levels of expression in the

tissues of the HPG axis shared by both sexes, the pituitary glands and hypothalami. Unlike our

previous analyses where we targeted genes of interest, here we analyzed all genes expressed,

controlling for over 15,000 multiple comparisons. We uncovered 361 sex-based differentially-

expressed genes in the pituitary: 218 were more highly expressed in males, while 153 were

more highly expressed in females. A table detailing differences in expression of all expressed

genes, including those statistically and non-statistically differentially expressed, can be

accessed at https://git.io/vXJWp. In contrast, only a single gene, the potassium voltage-gated

channel subfamily Q member 1 (*KCNQ1*), was statistically differentially expressed in the

hypothalamus (full table available here https://git.io/vXJW5). This profound disparity in patterns

of differential expression between hypothalamic and pituitary tissues begs a question of function

- how do the different tissues of the HPG axis contribute to physiological and behavioral

differences between males and females? Is the pituitary more involved in the maintenance of

sex-specific physiology and behavior than the hypothalamus? Studies comparing gene

expression in males and females are rare - most often studies have combined the sexes (but

see [38–40,49–54]). However, Nishida and colleagues [39] found similar patterns of gene expression in

mice, reporting that 43 genes were more highly expressed in the female pituitary while only 3

were more highly expressed in the hypothalamus as compared to males. Could this pattern of

sex-biased expression be conserved across vertebrates, and if so, can we conclude that the

pituitary, more so than the hypothalamus, plays a greater role in the maintenance of sex-specific

reproductive physiology and behavior?

Another hypothesis as to why we may observe less differential expression at the level of the

hypothalamus is because of the high heterogeneity of the region, which could result in the

dilution of observable differences in gene expression existing in more discrete locations. On the

other hand, high heterogeneity of the hypothalamus might only mask results of differential

expression if individual nuclei express antiparallel sex-specific changes, resulting in an

averaging of gene expression signals. Given this particular scenario is likely to be exceptionally

rare, it is possible that nuclei heterogeneity of the hypothalamus does not contribute significantly

to the observed patterns, though future investigations may yield more telling data if specific

nuclei are examined within the hypothalamus.

Gene ontology (GO) analysis of the differential expression in the pituitary revealed interesting
patterns of enrichment, or patterns of gene expression related to a common biological function
or process. GO terms enriched for genes expressed more highly in males were often related to
motor or muscle function, with specific terms such as muscle contraction being more highly
expressed in males as compared to females. The differentially expressed genes linked to these
GO terms include *Myosin* and other components of the myosin actin skeleton. Previous
research has identified a potential role for myosin in the pituitary, including in its primary function
of secretion [55–57]. For example, GnRH-induced secretion of LH can be altered by manipulating
the action of myosin in the anterior pituitary [56]. In females, a surge in LH will trigger ovulation.
We were unable to control for the specific ovulatory stage of the females sampled, though by
chance we can assume that most were not sampled at the exact point they were experiencing
their LH surge. However, males were sexually mature and actively paired with female partners
at the time of sampling. In males, LH stimulates Leydig cell production of testosterone. Thus,
one explanation for having GO term enrichment for genes more highly expressed in males
under this label of motor control is that the male pituitary could be more sensitive to the GnRH
signal to stimulate LH release due to myosin-related actions. The biological relevance of sex-
specific differential expression of *Myosin* remains unknown and deserves future study.

Gene ontology terms enriched for genes expressed more highly in the female pituitary were
often related to aspects of female growth and development. This may be due to their varying
stages of follicular development. Females, unlike males, must grow and maintain follicles and
the oviduct, which is energetically costly [59]. In a passerine species, females increased their
resting metabolic rate by 22% during egg laying, of which 18% was attributed to maintenance of
the oviduct [59]. While birds in our study were not actively nesting, we could not control for
reproductive stage of the gonads. When we extracted female gonads, we found that they were
experiencing different stages of follicular development. Some birds had regressed follicles, but
many had a follicular hierarchy. The presence of follicular hierarchy could explain the presence
of gene ontologies related to ovulation. Similarly, lipid binding and transport may be associated
with follicular development because females are laying down lipid-rich yolk. On the other hand,
males never had regressed testes. The active state of the male gonads is typical of pigeons
who breed year-round, except for a brief photorefractory period in the North temperate fall
where they may regress [60]. Gonads of females are more plastic and vary with the presence of

suitable mates and breeding stage [60].

Using the global analysis approach, we were able to identify sex-biased differentially expressed
genes that have not previously been targeted for investigation of HPG function. Such findings
that are yielded from this discovery-based approach offer promising targets for future study,
inspiring new lines of investigation [3]. One particularly promising target for understanding sex
differences in reproductive biology is Betacellulin (*BTC*), which is highly expressed in the female
pituitary as compared to the male pituitary (4.4 FC - Figure 4A). *BTC* is an epidermal-like growth
hormone known to regulate female reproduction [61] via mediation of LH [30] and regulation of
progesterone receptors [62]. In addition to being differentially expressed in the pituitary, it is also
highly expressed in the female ovary, but not the hypothalamus. It is lowly expressed, but
present in all male tissues, which begs the question, what role could *BTC* have in male
reproductive physiology, if any?

The Ecto-NOX Disulfide-Thiol Exchanger 1 (*ENOX1*) gene, an electron transport gene, is more
highly expressed in the male pituitary relative to the female pituitary (1.2x FC - Figure 4B). This
gene, known to be involved in circadian rhythms [63], has also been associated with weight gain
[64], litter size in the pig [65], and eggshell thickness in a genome-wide association study (GWAS) of
chickens [66]. What role this gene plays in the pituitary, and specifically in the male pituitary,
where it is differentially expressed, is completely unclear. Further work aimed at understanding
this and the other differentially expressed genes can be the key to gaining the most complete
understanding yet of how the HPG axis functions.

<u>Summary</u>
Researchers have described many important characteristics of the molecular basis for
reproductive behaviors, including the characterization of multiple genes expressed in the HPG
axis that are critical to reproduction. Our sex-specific, global analyses have uncovered hundreds
of sexually-biased genes. These genes can provide novel clues to assist behavioral biologists
and functional genomicists in more fully understanding the mechanisms underlying sexual
dimorphism in reproductive behaviors. Indeed, future work can now use this information to
better understand causal relationships and functions of genes expressed differently in males
and females.

In conclusion, we reveal patterns of tissue specific and sexually dimorphic gene expression in

the HPG axis. We report sex-biased expression in genes commonly investigated when studying

reproduction. In addition, we offer up promising new targets of investigation that could lead to a

better understanding of HPG function in both sexes. Our results highlight the need for sex parity

in transcriptomic studies, providing new lines of investigation of the mechanisms of reproductive

function.

Methods

Animal Collection Methods

Birds were housed at the University of California, Davis, in large, outdoor aviaries (5'x4'x7'), with

8 sexually reproductive adult pairs per aviary. All birds were three years old and had been

housed in their respective aviaries for one year at the time of collection. All birds were also

successful breeders, having paired naturally and produced multiple clutches throughout the year

thus far. However, to control for reproductive stage as much as was possible, all birds used in

this study were sampled at a time when they were naturally without eggs or chicks, creating a

type of "baseline" sampling point. All sampling occurred during late fall of 2015 and winter of

2016 between 10:00-12:00 (PST) to avoid potential seasonal and circadian rhythm confounds.

While birds were exposed to natural light, we also augmented their aviaries with artificial lights

set to 14L:10D photoperiod in order to maintain reproductive condition. All methods were carried

out in accordance with relevant guidelines and regulations, and all experimental protocols were

approved by the University of California, Davis, IACUC (#18895). Fourteen males and ten

females were sacrificed within 3 min of entering their aviary. They were anesthetized using

isoflurane prior to decapitation. Brains, pituitaries, and gonads were extracted and immediately

placed on dry ice and transferred within the hour to a -80°C freezer until further processing.

Frozen brains were coronally sectioned on a cryostat (Leica CM 1860) at 100µM and the

hypothalamus was isolated using surgical punches. We used Karten and Hodos' stereotaxic

atlas [67] of the brain of the pigeon to locate the hypothalamus and collect it in its entirety. In brief,

we, collected hypothalamic tissue beginning at the point of bifurcation of the tractus

septomesencephalicus and ending after the cerebellum was well apparent. Hypothalamic,

pituitary, and gonadal tissue were preserved at -80°C in RNALater and shipped from the

University of California, Davis, to the University of New Hampshire for further processing.

Library Preparation and Sequencing

Tissues frozen in RNALater were thawed on ice in an RNAse-free work environment. Total RNA

was extracted using a standard Trizol extraction protocol (Thermo Fisher Scientific, Waltham,

MA). The quality of the resultant extracted total RNA was characterized using the Tapestation

2200 instrument (Agilent, Santa Clara, CA), after which Illumina sequence libraries were

prepared using the TruSeq RNA Stranded LT Kit (Illumina). The Tapestation was, again, used

to determine the quality and concentration of these libraries. Each library was diluted to 2nM

with sterile ddH$_2$O, and pooled in a multiplexed library sample. The multiplexed library sample

was then sent to the New York Genome Center for 125 or 150 base pair paired-end sequencing

on a HiSeq 2500 platform.

Sequence Quality Control and Assembly

Sequence read data were downloaded and quality checked using FastQC version 0.11.5 [68]. A

de novo transcriptome for the HPG was assembled following the Oyster River Protocol for

Transcriptome Assembly [69]. In brief, 59 million paired end reads from two individuals (1 male

and 1 female) were error corrected using RCorrector version 1.0.2 [70]. Adapters, as well as

bases with a Phred score <2 were trimmed using Skewer 0.2.2 [71], and assembly was carried out

using Trinity version 2.2.0 [72], Binpacker version 1.0 [73], and Shannon version 0.0.2 [74], after which

the resultant transcriptomes were merged using the software package transfuse version 0.5.0

(https://github.com/cboursnell/transfuse). Lowly expressed transcripts, defined as those with an

abundance of less than 0.5 transcript-per-million (TPM<0.5), were filtered out of the dataset.

The resultant assembly was annotated using the software package dammit

(https://github.com/camillescott/dammit), and evaluated using BUSCO 2.0 beta 3 [75] and

TransRate version 1.0.1 [76].

Mapping and Differential Gene Expression All samples were used (20 hypothalamus (11

male, 9 female), 23 pituitary (14 male, 9 female), 13 testes, and 10 ovary samples from 24

birds.) Raw reads were mapped to the annotated transcriptome after a quasi-mapping index

was prepared using Salmon 0.7.0 [77]. Rock dove transcripts were mapped to genes from the

Gallus gallus genome version 5, using BLAST [78]. All data were then imported into the R

statistical package (version 3.3.0) [79] using tximport [80] for gene level evaluation of gene

expression, which was calculated using edgeR (version 3.1.4) [81] following TMM normalization

and correction for multiple hypothesis tests by setting the false discovery rate (FDR) to 1%.

Gene Expression Evaluation

Although excellent genomic resources exist for the Rock Dove [17], the current Rock Dove

genome available lacks functional annotation (*e.g.,* gene ontology terms, information about

function and protein networks). Thus, we elected to establish orthologous relationships between

transcript sequences we generated in Rock Dove with those from the *Gallus gallus* genome,

version 5.

Patterns of gene expression were evaluated using two distinct methods. First, we selected *a*

priori genes of interest based on their known involvement in reproduction and associated

behaviors (Table 3). Differences in gene expression were evaluated between these genes in the

hypothalamic, pituitary, and gonadal tissues and between both sexes using a generalized linear

model framework (expression ~ sex * tissue) with post-hoc significance for all pairwise

combinations of factors tested using the Bioconductor package lsmeans (https://cran.r-

project.org/package=lsmeans). Second, a more global investigation of differential expression

was conducted to characterize the general presence and expression of all genes per tissue and

how they might differ between the sexes. A comparison of differential expression between

tissues was not carried out secondary to results from preliminary analysis suggesting that most

genes are expressed differently, thus, making meaningful comparisons uninformative. Instead,

to characterize patterns of expression in each tissue, we generated a transcript

presence/absence matrix (1=present, 0=absent) using median CPM (counts-per-million,

generated in edgeR) > 10 as the metric. Overlaps were visualized using the R package UpSet

[82]. Gene ontology enrichment analysis was carried out using the Kolmogorov-Smirnov test [83] for

significance in the R package, topGO [84].

Acknowledgements:

This work is supported by NSF IOS 1455960 to RMC and MM. We thank Jesse Krause,
606   Jonathan Peréz, and members of the Calisi Lab for their help in maintaining the rock dove
607   aviaries and aiding in tissue collection. We also thank Stacia Sower, Samuel Díaz-Muñoz,
608   and the University of California, Davis, Environmental Endocrinology Group (EEG),
609   comprised of the labs of Rebecca Calisi, Tom Hahn, Marilyn Ramenofsky, Karen Ryan, and
610   John Wingfield, for their comments on this project and manuscript. We are grateful to
611   Natalia Duque for her illustration (Fig. 1).

613   **Author Contribution Statements**

614   Matthew MacManes and Andrew Lang constructed sequencing libraries, did bioinformatics

615   analyses, and wrote the paper. Suzanne H. Austin, April Booth, Victoria Farrar and Rebecca

616   Calisi conducted live animal experiments and wrote the paper.

Additional Information

Read data is available at PRJEB16136

Competing Financial Interests

The authors declare no competing financial interests.

REFERENCES

1. Sower, S. A., Freamat, M. & Kavanaugh, S. I. The origins of the vertebrate hypothalamic–pituitary–gonadal (HPG) and hypothalamic–pituitary–thyroid (HPT) endocrine systems: New insights from lampreys. *Gen. Comp. Endocrinol.* **161,** 20–29 (2009/3).

2. Plachetzki, D. C., Tsai, P.-S., Kavanaugh, S. I. & Sower, S. A. Ancient origins of metazoan gonadotropin-releasing hormone and their receptors revealed by phylogenomic analyses. *Gen. Comp. Endocrinol.* **234,** 10–19 (2016).

3. Calisi, R. M. & MacManes, M. D. RNAseq-ing a more integrative understanding of animal behavior. *Current Opinion in Behavioral Sciences* **6,** 65–68 (2015).

4. Lake, J. I., Lange, H. S., O'Brien, S., Sanford, S. E. & Maney, D. L. Activity of the hypothalamic-pituitary-gonadal axis differs between behavioral phenotypes in female white-throated sparrows (Zonotrichia albicollis). *Gen. Comp. Endocrinol.* **156,** 426–433 (2008).

5. Dhillo, W. S. *et al.* Kisspeptin-54 stimulates the hypothalamic-pituitary gonadal axis in human males. *J. Clin. Endocrinol. Metab.* **90,** 6609–6615 (2005).

6. Saldanha, C. J., Walters, B. J. & Fraley, G. S. Neurons that co-localize aromatase- and kisspeptin-like immunoreactivity may regulate the HPG axis of the Mallard drake (Anas platyrhynchos). *Gen. Comp. Endocrinol.* **166,** 606–613 (2010).

7. Wang, R.-L., Bencic, D., Lazorchak, J., Villeneuve, D. & Ankley, G. T. Transcriptional

regulatory dynamics of the hypothalamic-pituitary-gonadal axis and its peripheral pathways

as impacted by the 3-beta HSD inhibitor trilostane in zebrafish (Danio rerio). *Ecotoxicol.*

Environ. Saf. **74,** 1461–1470 (2011).

8. Zhang, X. *et al.* Real-time PCR array to study effects of chemicals on the Hypothalamic-

Pituitary-Gonadal axis of the Japanese medaka. *Aquat. Toxicol.* **88,** 173–182 (2008).

9. Qin, F. *et al.* Gene expression profiling of key genes in hypothalamus-pituitary-gonad axis

of rare minnow Gobiocypris rarus in response to EE2. *Gene* **552,** 8–17 (2014).

10. Han, H. *et al.* Fluoride exposure changed the structure and the expressions of reproductive

related genes in the hypothalamus-pituitary-testicular axis of male mice. *Chemosphere* **135,**

297–303 (2015).

11. Peterson, M. P. *et al.* De novo transcriptome sequencing in a songbird, the dark-eyed junco

(Junco hyemalis): genomic tools for an ecological model system. *BMC Genomics* **13,** 305

(2012).

12. Cánovas, A. *et al.* Multi-tissue omics analyses reveal molecular regulatory networks for

puberty in composite beef cattle. *PLoS One* **9,** e102551 (2014).

13. Xu, J. *et al.* Defining global gene expression changes of the hypothalamic-pituitary-gonadal

axis in female sGnRH-antisense transgenic common carp (Cyprinus carpio). *PLoS One* **6,**

e21057 (2011).

14. Lehrman, D. S. The Physiological Basis of Parental Feeding Behavior in the Ring Dove

(Streptopelia Risoria). *Behaviour* **7,** 241–285 (1955).

15. Ball, G. F. & Silver, R. Timing of incubation bouts by ring doves (Streptopelia risoria). *J.*

Comp. Psychol. **97,** 213–225 (1983).

16. Gillespie, M. J. *et al.* Transcriptome analysis of pigeon milk production � role of

cornification and triglyceride synthesis genes. *BMC Genomics* **14,** 1–1 (2013).

17. Shapiro, M. D. *et al.* Genomic Diversity and Evolution of the Head Crest in the Rock

Pigeon. *Science* **339,** 1063–1067 (2013).

18. Domyan, E. T. & Shapiro, M. D. Pigeonetics takes flight: Evolution, development, and

genetics of intraspecific variation. *Dev. Biol.* (2016). doi:10.1016/j.ydbio.2016.11.008

19. Waterhouse, R. M., Tegenfeldt, F., Li, J., Zdobnov, E. M. & Kriventseva, E. V. OrthoDB: a

hierarchical catalog of animal, fungal and bacterial orthologs. *Nucleic Acids Res.* **41,** D358–

65 (2013).

20. Saier, M. H., Jr, Tran, C. V. & Barabote, R. D. TCDB: the Transporter Classification

Database for membrane transport protein analyses and information. *Nucleic Acids Res.* **34,**

D181–6 (2006).

21. Griffiths-Jones, S. *et al.* Rfam: annotating non-coding RNAs in complete genomes. *Nucleic*

Acids Res. **33,** D121–4 (2005).

22. Calisi, R. M. An integrative overview of the role of gonadotropin-inhibitory hormone in

behavior: Applying Tinbergen's four questions. *Gen. Comp. Endocrinol.* **203,** 95–105

(2014).

23. Cloues, R., Ramos, C. & Silver, R. Vasoactive intestinal polypeptide-like immunoreactivity

during reproduction in doves: influence of experience and number of offspring. *Horm.*

Behav. **24,** 215–231 (1990).

24. Silver, R. Prolactin and parenting in the pigeon family. *J. Exp. Zool.* **232,** 617–625 (1984).

25. Ondrasek, N. R. Emerging Frontiers in Social Neuroendocrinology and the Study of

Nonapeptides. *Ethology* **122,** 443–455 (2016).

26. Kelly, A. M. & Goodson, J. L. Hypothalamic oxytocin and vasopressin neurons exert sex-

specific effects on pair bonding, gregariousness, and aggression in finches. *Proc. Natl.*

Acad. Sci. U. S. A. **111,** 6069–6074 (2014).

27. Liley, N. R. The role of estrogen and progesterone in the regulation of reproductive

behaviour in female ring doves (Streptopelia risoria) under long vs. short photoperiods.

Can. J. Zool. **54,** 1409–1422 (1976).

28. Balthazart, J. & Ball, G. *Brain Aromatase, Estrogens, and Behavior*. (OUP USA, 2012).

29. McCarthy, D. J., Chen, Y. & Smyth, G. K. Differential expression analysis of multifactor

RNA-Seq experiments with respect to biological variation. *Nucleic Acids Res.* **40,** 4288–

4297 (2012).

30. Park, J.-Y. *et al.* EGF-like growth factors as mediators of LH action in the ovulatory follicle.

Science **303,** 682–684 (2004).

31. Ghosh, M. G., Thompson, D. A. & Weigel, R. J. PDZK1 and GREB1 are estrogen-regulated

genes expressed in hormone-responsive breast cancer. *Cancer Res.* **60,** 6367–6375

(2000).

32. Lee, M. P., Hu, R. J., Johnson, L. A. & Feinberg, A. P. Human KVLQT1 gene shows tissue-

specific imprinting and encompasses Beckwith-Wiedemann syndrome chromosomal

rearrangements. *Nat. Genet.* **15,** 181–185 (1997).

33. Nelson & Kriegsfield. *Introduction to Behavioral Endocrinology*. (Sinauer, 2017).

34. Castelli, F. R., Kelley, R. A., Keane, B. & Solomon, N. G. Female prairie voles show social

and sexual preferences for males with longer avpr1a microsatellite alleles. *Anim. Behav.*

82, 1–10 (2011).

35. Staes, N. *et al.* Chimpanzee sociability is associated with vasopressin (Avpr1a) but not

oxytocin receptor gene (OXTR) variation. *Horm. Behav.* **75,** 84–90 (2015).

36. Walum, H. *et al.* Genetic variation in the vasopressin receptor 1a gene (AVPR1A)

associates with pair-bonding behavior in humans. *Proceedings of the National Academy of*

Sciences **105,** 14153–14156 (2008).

37. Turner, L. M. *et al.* Monogamy evolves through multiple mechanisms: evidence from V1aR

in deer mice. *Mol. Biol. Evol.* **27,** 1269–1278 (2010).

38. Trabzuni, D. *et al.* Widespread sex differences in gene expression and splicing in the adult

human brain. *Nat. Commun.* **4,** 2771 (2013).

39. Nishida, Y., Yoshioka, M. & St-Amand, J. Sexually dimorphic gene expression in the

hypothalamus, pituitary gland, and cortex. *Genomics* **85,** 679–687 (2005).

40. Shan, L. *et al.* Transcriptome Profiling Identifies Differentially Expressed Genes in Postnatal

Developing Pituitary Gland of Miniature Pig. *DNA Res.* (2013). doi:10.1093/dnares/dst051

41. Lim, M. M. *et al.* Enhanced partner preference in a promiscuous species by manipulating

the expression of a single gene. *Nature* **429,** 754–757 (2004).

42. Angelier, F. & Chastel, O. Stress, prolactin and parental investment in birds: a review. *Gen.*

Comp. Endocrinol. **163,** 142–148 (2009).

43. Buntin, J. D., Hnasko, R. M. & Zuzick, P. H. Role of the ventromedial hypothalamus in

prolactin-induced hyperphagia in ring doves. *Physiol. Behav.* **66,** 255–261 (1999).

44. McCarthy, M. M., Arnold, A. P., Ball, G. F., Blaustein, J. D. & De Vries, G. J. Sex

differences in the brain: the not so inconvenient truth. *J. Neurosci.* **32,** 2241–2247 (2012).

45. Askew, J. A., Georgiou, G. C., Sharp, P. J. & Lea, R. W. Localization of progesterone

receptor in brain and pituitary of the ring dove: influence of breeding cycle and estrogen.

Horm. Behav. **32,** 105–113 (1997).

46. Walters, K. A. *et al.* Targeted loss of androgen receptor signaling in murine granulosa cells

of preantral and antral follicles causes female subfertility. *Biol. Reprod.* **87,** 151 (2012).

47. Wacker, D. W., Wingfield, J. C., Davis, J. E. & Meddle, S. L. Seasonal changes in

aromatase and androgen receptor, but not estrogen receptor mRNA expression in the brain

of the free-living male song sparrow, Melospiza melodia morphna. *J. Comp. Neurol.* **518,**

3819–3835 (2010).

48. Chang, C., Lee, S. O., Wang, R.-S., Yeh, S. & Chang, T.-M. Androgen receptor (AR)

physiological roles in male and female reproductive systems: lessons learned from AR-

knockout mice lacking AR in selective cells. *Biol. Reprod.* **89,** 21 (2013).

49. Hodes, G. E. *et al.* Sex Differences in Nucleus Accumbens Transcriptome Profiles

Associated with Susceptibility versus Resilience to Subchronic Variable Stress. *J. Neurosci.*

35, 16362–16376 (2015).

50. Balakrishnan, C. N., Lin, Y.-C., London, S. E. & Clayton, D. F. RNA-seq transcriptome

analysis of male and female zebra finch cell lines. *Genomics* **100,** 363–369 (2012).

51. Zhang, Z. *et al.* Transcriptome Analysis of Female and Male Xiphophorus maculatus Jp 163

A. *PLoS One* **6,** e18379 (2011).

52. Sun, L., Wang, C., Huang, L., Wu, M. & Zuo, Z. Transcriptome analysis of male and female

Sebastiscus marmoratus. *PLoS One* **7,** e50676 (2012).

53. Jansen, R. *et al.* Sex differences in the human peripheral blood transcriptome. *BMC*

Genomics **15,** 33 (2014).

54. Weickert, C. S. *et al.* Transcriptome analysis of male–female differences in prefrontal

cortical development. *Mol. Psychiatry* **14,** 558–561 (2009).

55. Lloyd, R. V. *et al.* Myosin XVA expression in the pituitary and in other neuroendocrine

tissues and tumors. *Am. J. Pathol.* **159,** 1375–1382 (2001).

56. Rao, K. *et al.* Wortmannin-Sensitive and -Insensitive Steps in Calcium-Controlled

Exocytosis in Pituitary Gonadotrophs: Evidence That Myosin Light Chain Kinase Mediates

Calcium-Dependent and Wortmannin-Sensitive Gonadotropin Secretion. *Endocrinology*

138, 1440–1449 (1997).

57. Ostlund, R. E., Jr, Leung, J. T. & Kipnis, D. M. Myosins of secretory tissues. *J. Cell Biol.* **77,**

827–836 (1978).

58. Lovell-mansbridge, C. & Birkhead, T. R. Do female pigeons trade pair copulations for

protection? *Anim. Behav.* **56,** 235–241 (1998).

59. Vézina, F., Salvante, K. G. & Williams, T. D. The metabolic cost of avian egg formation:

possible impact of yolk precursor production? *J. Exp. Biol.* **206,** 4443–4451 (2003).

60. Murton, R. K. & Westwood, N. J. *Avian breeding cycles*. (Oxford University Press, USA,

1977).

61. Gratao, A. A., Dahlhoff, M., Sinowatz, F., Wolf, E. & Schneider, M. R. Betacellulin

overexpression in the mouse ovary leads to MAPK3/MAPK1 hyperactivation and reduces

litter size by impairing fertilization. *Biol. Reprod.* **78,** 43–52 (2008).

62. Shimada, M., Hernandez-Gonzalez, I., Gonzalez-Robayna, I. & Richards, J. S. Paracrine

and Autocrine Regulation of Epidermal Growth Factor-Like Factors in Cumulus Oocyte

Complexes and Granulosa Cells: Key Roles for Prostaglandin Synthase 2 and

Progesterone Receptor. *Mol. Endocrinol.* **20,** 1352–1365 (2006).

63. Dick, S. S., Ryuzoji, A., Morré, D. M. & Morré, D. J. Identification of the constitutive

ultradian oscillator of the circadian clock (ENOX1) in Saccharomyces cerevisiae. (2013).

64. Serão, N. V. *et al.* Single nucleotide polymorphisms and haplotypes associated with feed

efficiency in beef cattle. *BMC Genet.* **14,** 94 (2013).

65. Sell-Kubiak, E. *et al.* Genome-wide association study reveals novel loci for litter size and its

variability in a Large White pig population. *BMC Genomics* **16,** 1049 (2015).

66. Liu, W. *et al.* A Genome-Wide SNP Scan Reveals Novel Loci for Egg Production and

Quality Traits in White Leghorn and Brown-Egg Dwarf Layers. *PLoS One* **6,** e28600 (2011).

67. Karten, H. J. & Hodos, W. *A stereotaxic atlas of the brain of the pigeon: Columba livia*. **696,**

(Johns Hopkins Press, 1967).

68. Andrews, S. R. *FastQC*. (2016).

69. MacManes, M. D. Establishing evidenced-based best practice for the de novo assembly

and evaluation of transcriptomes from non-model organisms. *biorxiv.org* 1–23 (2015).

70. Song, L. & Florea, L. Rcorrector: efficient and accurate error correction for Illumina RNA-

seq reads. *Gigascience* **4,** 48 (2015).

71. Jiang, H., Lei, R., Ding, S.-W. & Zhu, S. Skewer: a fast and accurate adapter trimmer for

next-generation sequencing paired-end reads. *BMC Bioinformatics* **15,** 182 (2014).

72. Haas, B. J. *et al.* De novo transcript sequence reconstruction from RNA-seq using the

Trinity platform for reference generation and analysis. *Nat. Protoc.* **8,** 1494–1512 (2013).

73. Liu, J. *et al.* BinPacker: Packing-Based De Novo Transcriptome Assembly from RNA-seq

Data. *PLoS Comput. Biol.* **12,** e1004772 (2016).

74. Kannan, S., Hui, J., Mazooji, K., Pachter, L. & Tse, D. *Shannon: An Information-Optimal de*

Novo RNA-Seq Assembler. (Cold Spring Harbor Labs Journals, 2016).

75. Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V. & Zdobnov, E. M.

BUSCO: assessing genome assembly and annotation completeness with single-copy

orthologs. *Bioinformatics* **31,** 3210–3212 (2015).

76. Smith-Unna, R. D., Boursnell, C., Patro, R., Hibberd, J. M. & Kelly, S. TransRate: reference

free quality assessment of de-novo transcriptome assemblies. *bioRxiv* 1–25 (2015).

77. Patro, R., Duggal, G. & Kingsford, C. Accurate, fast, and model-aware transcript expression

quantification with Salmon. *biorxiv.org* 1–35 (2015).

78. Camacho, C. *et al.* BLAST+: architecture and applications. *BMC Bioinformatics* **10,** 421

(2009).

79. RStudio Team. RStudio: Integrated Development for R. (2015). Available at:

http://www.rstudio.com/. (Accessed: 1st September 2016)

80. Soneson, C., Love, M. I. & Robinson, M. D. Differential analyses for RNA-seq: transcript-

level estimates improve gene-level inferences. *F1000Res.* **4,** 1521–1519 (2015).

81. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for

differential expression analysis of digital gene expression data. *Bioinformatics* **26,** 139–140

(2010).

82. Lex, A., Gehlenborg, N., Strobelt, H., Vuillemot, R. & Pfister, H. UpSet: Visualization of

Intersecting Sets. *IEEE Trans. Vis. Comput. Graph.* **20,** 1983–1992 (2014).

83. Young, I. T. Proof without prejudice: use of the Kolmogorov-Smirnov test for the analysis of

histograms from flow systems and other sources. *J. Histochem. Cytochem.* **25,** 935–941

(1977).

84. Alexa, A. & Rahnenfuhrer, J. topGO: enrichment analysis for gene ontology. *R package*

version **2,** (2010).

81. Zhang Z, Wang Y, Wang S, Liu J, Warren W, Mitreva M, Walter RB (2011) Transcriptome

Analysis of Female and Male Xiphophorus maculatus Jp 163 A. PLoS One 6:e18379

Available at:

http://journals.plos.org/plosone/article/asset?id=10.1371/journal.pone.0018379.PDF.